A research monograph

The Geometry and Arithmetic of Iterated Theta Lifts

A reconstruction and study of the shortened Alpöge–Claude proof of Dirichlet GRH

Proof-gap closure edition III

Working mathematical manuscript – November 2026

Scope. This seventh edition builds upon the 849-page second proof-closure edition. It attacks the remaining precise proof interfaces rather than adding general background: each new chapter gives a written derivation or a checked literature bridge and records its residual hypotheses. Not every specialized comparison has yet been discharged. The final ledger explicitly identifies what is still required for a truly independent conventional proof.

Attribution. The original mathematical argument is attributed by the supplied artifact to Levent Alpöge with Claude. The shortened Lean development is the primary source of this exposition. No claim of independent kernel verification, new authorship of the GRH proof, or completion of every remaining paper-proof obligation is made. The shortened Lean code on which this exposition is based is credited to Luke Sernau; see his repository, https://github.com/lsernau/grh-lean.

AI disclaimer. The text of this document was written by AI, ChatGPT-6.2 Sol, and has not been checked by a human.

Preface: the purpose of this monograph

The compressed formal proof of the generalized Riemann hypothesis for Dirichlet L-functions presents a very unusual task for an expository work. There is a clearly identifiable central argument, but the mathematics needed to implement its quantitative estimates comes from several distinct areas. A reader who knows the analytic theory of L-functions will not necessarily be familiar with the integral model of a unitary theta lift. A specialist in p-adic Hodge theory may have little reason to know the exact Bessel normalization in a trace calculation. And a specialist in automorphic forms may be familiar with the norm formulas of the theta correspondence, but not with the torsion or crystalline comparison estimates that are needed uniformly over all coefficient primes. The difficulty is not merely that there are many auxiliary lemmas; the lemmas belong to different mathematical languages and are linked in numerical ways that must survive translation.

The 49-page first exposition supplied a guide to the mathematical architecture. The 162-page specialist version then expanded many of the central estimates, organized the proofs of the analytical endgame, and gave the interfaces to the remaining arithmetic statements. The 333-page third exposition supplied systematic technical companion chapters. The fourth, 550-page exposition organized the complete argument as a detailed reference monograph, with explicit outstanding specialized inputs. The fifth, 711-page edition undertook a focused proof-closure pass. The sixth, 849-page edition supplied source-matched derivations of further analytic, arithmetic, and local interfaces. The present seventh edition develops the weighted filling and frequency calculus, actual Bessel levels and majorants, singular-coefficient induction and period exchange, carrier analysis, and the arithmetic passage from integral trace functions to the Ribet and denominator operators. It preserves that logical spine but extends it to include proofs of more of the hypotheses of the earlier companion chapters. In particular we reconstruct the elementary analytic inputs, rather than invoking uniform prime distribution, and develop the normalized global data, the exact finite and local computations, and the specialized geometric and arithmetic reductions of the main argument. The order is chosen to make the relationship between the conceptual proof and its technical realization visible.

For whom the book is written. We assume a graduate-level acquaintance with algebra, number fields, linear algebra and complex analysis. A reader with experience in automorphic forms, representation theory, arithmetic geometry or Galois representations should be able to follow the chapters of their own specialty and to understand the role of the others. In a largely self-contained mathematical monograph it is reasonable to take major classical theorems, such as global reciprocity, classical Hermitian mass formulas, and standard foundations of p-adic Hodge theory, from the literature rather than re-establish the entire theory. The distinction is that specialized strengthenings used here, especially exact integral or quantitative variants, must be separately justified. Where this has not been possible, the exact missing assertion is

identified rather than silently treated as a textbook theorem.

What constitutes a proof here. The label *formal-source input*, where not discharged in a later chapter of this edition, marks an assertion taken from a specified theorem in the Lean repository whose conventional proof, including its precise hypotheses and constants, is not completely reconstructed in this manuscript. Such an input is not a new conjecture or an additional axiom of the formal development. It is an outstanding obligation for a proof that a mathematician could verify without using that development as a certificate. In the passages presented as ordinary proofs, the argument is given from the stated hypotheses and inputs. The geometric constructions and some of the most technical integral and uniform estimates are still mathematical work requiring further checking. The final chapter lists them by theme. The aim is to make genuine progress in conventional exposition without disguising a gap by calling a result "standard".

Reading strategy. Read the introductory chapter, the analytic reduction and the final arithmetic assembly first. These explain why three specific estimates suffice and where the critical exponent arises. Read the parameter and common arithmetic-model chapters next. The upper-bound, spanning and lower-bound parts can then largely be studied separately. Each includes both the main chain of deductions and complementary chapters which derive supporting lemmas and specialize classical theorems. Earlier statements are sometimes named using their formal declaration to preserve concordance even when individual components are subsequently proved. Consult the final dependency audit and the internal cross-references to distinguish the complete parts of a proof from those whose exact specialized hypotheses still need to be established.

Part	Main mathematical question
I. Analytic reduction and models	How can a one-sided smoothed prime sum exclude a zero, and what are the fields, weights, levels and integral spaces that will produce it?
II. Upper bounds	Why do derivatives of CM theta functions contain the valuations needed to count prime powers, and how can lifts of integral forms be made short without a prohibitively large index?
III. Spanning	How can the short forms be made to span almost all of a large automorphic space, with an explicit bound on the number of translates?
IV. Lower bounds	Why must an almost-basis have large total volume, and how do exchange identities, special values, and congruences control the integral costs?
V. Assembly and audit	How do these estimates cancel, yield a square-root saving, and imply GRH? Which mathematical dependencies remain to be independently justified?

A conceptual point. Products of positive integers that encode primes can give large archimedean values; factorials are the simplest example. The proof at issue is much more elaborate, but one can recognize a familiar product-formula pattern. A value whose finite valuations are large cannot simultaneously be too small at every infinite place, unless it vanishes. Here the relevant object is not a single nonzero value but a family of automorphic vectors, its determinant, and an auxiliary family of twists measuring different finite places. The main term of the upper estimate measures the cost of derivatives after subtracting guaranteed divisibilities. The lower estimate recovers exactly that cost with the opposite sign via volume ratios. The canonical cancellation between them, not an absolute estimate of a high-degree

automorphic L-function, gives the saving. This is the mathematical thread to keep in mind across the many local constructions.

Conventions. The prime d defines $K = \mathbb{Q}(\sqrt{-d})$, and $\mathcal{D}$ denotes a chosen admissible datum. The integer m is the number of theta lifts, r the initial row rank, $k = r + 1 - m$ the initial column rank, and $t = 5r$ the rank increment on the row chain. The symbol $n = \dim W$ is the dimension of the row form space, not a Hermitian rank. We reserve d_{ind} for a logarithmic index cost. The scale b is real and large, $B = fa(\log b)^3$ is the basic error allowance, $L = B \log b$, and $H = mk\mathcal{U}$ is the common main term. The normalized prime error R_M differs from the arithmetic remainder $\mathcal{R}$. A glossary at the end of the volume collects these and other important symbols, with references to the chapters where they enter. We use the compact edition's numerical constants unless otherwise specified. For source references, a module path beginning with `New/` or `Lib/` is relative to `grh-compact/src/` of the supplied repository.

Contents

Preface: the purpose of this monograph **i**

I The analytic reduction and arithmetic models **1**

1 Introduction and mathematical status **2**
1.1 The architecture of the proof 2
1.2 Primary material, conventions, and the remaining challenge 4

2 The analytic problem: a one-sided bound for weighted prime sums **5**
2.1 The particular smoothing used in the proof 5
2.2 Why the prime sum contains information about zeros 7
2.3 An elementary form of Landau's one-sided principle 8
2.4 Quadratic twists avoiding a prescribed half-strip 9
2.5 The one-sided zero-free criterion 12
2.6 A good second scale in a long window 13
2.7 From the target bound to GRH 14

3 Classical analytic foundations without a prime number theorem **16**
3.1 Finite characters, Gauss sums, and the Fourier bound 16
3.2 The quadratic symbol and the choice of the family 18
3.3 Holomorphy and nonvanishing on the line at one 19
3.4 Reciprocal primes in a fixed residue class 21
3.5 The theta transform and the full functional equations 22
3.6 Analytic details of the bad-twist estimate 24
3.7 A uniform disk at one for powers of a fixed modulus 25
3.8 Prime existence in long logarithmic windows 27
3.9 Final conditional theorem and quantifier audit 29

4 The admissible data and coefficient spaces **32**
4.1 The CM extension and the different 32
4.2 The scales, and the reason for so many parameters 33
4.3 Admissibility and construction of the data 34
4.4 Derived weights, level and integral structures 36
4.5 Two kinds of automorphic forms 36

4.6 Precise weights and levels of the source and target . . . 37
4.7 Integral gauges and the precise uniform index . . . 39
4.8 The theta chains and their relation . . . 40
4.9 Norms, squared covolume and index . . . 40
4.10 The common leading term and the prime remainder . . . 41

5 Adelic coefficient models and integral linear algebra 43
5.1 Hermitian spaces, places and adelic lattices . . . 43
5.2 Why the extra level removes stabilizer denominators . . . 45
5.3 Normalization of class sums and change of level . . . 46
5.4 A useful elementary form of weak approximation . . . 47
5.5 Algebraic modular forms and the finite-dimensional model . . . 48
5.6 The integral Weyl model . . . 48
5.7 Fischer pairing and the highest-weight hook constant . . . 50
5.8 Examples of the coefficient normalizations . . . 51
5.9 The class-sum metric and the real matrix-coefficient model . . . 51
5.10 Transported lattices and gauges . . . 52
5.11 Finite-place Hecke operators on explicit lattices . . . 54
5.12 Class correspondences and adjoints . . . 55
5.13 Where this elementary model stops . . . 56

6 The splitting character, integral unitary models and the definite mass 57
6.1 Notation and the local norm theorem . . . 57
6.2 A character extension with exactly the required sign . . . 58
6.3 The self-dual lattice and half of the different . . . 61
6.4 Reduction of finite ideals and their adelic generators . . . 62
6.5 Completions of the coefficient primes and the choice of a side . . . 63
6.6 Unimodular lattices over split and unramified places . . . 64
6.7 Existence and equivalence of the framed global spaces . . . 65
6.8 Counting the local finite groups and their congruence quotients . . . 66
6.9 An exact norm-one class-sum identity . . . 67
6.10 The classical mass formula in the present normalization . . . 68
6.11 Useful consequences and the separation of normalizations . . . 69

7 Heisenberg operators and the finite vacuum multiplier 71
7.1 The saturating lattice at a finite place . . . 71
7.2 The multiplier and its transformation laws . . . 72
7.3 The finite vacuum and the Heisenberg cocycle . . . 73
7.4 The lattice model and the absolute vacuum coefficient . . . 74
7.5 The Gauss-sum phase and the finite splitting . . . 75
7.6 The real harmonic model and its basic normalization . . . 77
7.7 The rational theta functional in a polarized model . . . 78

8 Foundational theorem audit and universal coefficient comparison 80
8.1 The universal integral Schur basis . . . 80
8.2 The norm subgroup and the normalization of the splitting character . . . 83
8.3 Classical CM and the discriminant normalization . . . 84
8.4 The exact classical finite local condition . . . 86
8.5 The integral characteristic-polynomial interface in Ribet's construction . . . 88
8.6 How to read the input labels in the preceding chapters . . . 89

9 The normalized real oscillator, global theta functional and mixed model 91
9.1 The real vacuum identity . . . 91
9.2 The real phase from a Gaussian determinant . . . 92
9.3 Adelic duality and summable vacuum correlations . . . 94
9.4 Existence, uniqueness and normalization of theta . . . 95
9.5 The normalized hyperbolic Schrödinger model . . . 97
9.6 The zero-term character on the actual parabolic . . . 98
9.7 The mixed-model expansion and equality with the source zero term . . . 100
9.8 Rational invariance in the exact nonpolar representation . . . 101
9.9 Application to a unitary dual pair . . . 102
9.10 Closure ledger for this chapter . . . 103

10 Qualitative theta growth and convergence in the marked lattices 104
10.1 Lattice height and the compactness criterion . . . 104
10.2 A bounded archimedean step that improves the height . . . 105
10.3 Moderate growth of the theta kernel . . . 106
10.4 Small real shifts in the Fock norm . . . 107
10.5 The finite annihilation test . . . 108
10.6 Decay on a full-rank split support . . . 108
10.7 Convergence and transposition of theta periods . . . 110
10.8 Status of the growth interfaces . . . 110

11 The convergent-range Siegel–Weil formula in the integrable model 112
11.1 The classical theorem and the range used here . . . 112
11.2 Identifying the splitting of the Weil representation . . . 113
11.3 The evaluation at a rational Lagrangian . . . 114
11.4 The maximum of a coherent wave vacuum . . . 114
11.5 A common integrable majorant for coherent translates . . . 115
11.6 The graph term in the doubled datum . . . 116

II Upper bounds: CM jets, theta chains and Hecke repair 118

12 The CM differential, theta normalization and integral frame 119
12.1 An elliptic curve normalized at a point of order two . . . 119
12.2 Theta sections and their integral size . . . 121

12.3 Isogenies, their multipliers, and Tate's inequality . . . 122
12.4 The integral formal parameter and the Gauss principle . . . 123
12.5 Logarithms, norm-pair limits, and the formal height . . . 125
12.6 A unit-adjusted Chowla–Selberg formula . . . 127
12.7 The finite idele and the period scale . . . 129
12.8 Constructing arithmetic tight frames of prescribed length . . . 130
12.9 Frobenius on the frame matrix . . . 133
12.10 Theta-group orthogonality and the finite realization . . . 134
12.11 From an arithmetic frame to a theta section . . . 136
12.12 Integral vacuum and the remaining reciprocity law . . . 137
12.13 What the arithmetic normalization proves . . . 139

13 Local CM theta analysis and Schur jets **140**
13.1 The arithmetic local scales . . . 140
13.2 The CM period and the differential coordinate . . . 141
13.3 Integral and logarithmic theta identities . . . 142
13.4 The regularized CM parameter at nonsplit primes . . . 143
13.5 The exceptional exponential factor . . . 144
13.6 Tight frames and theta vectors . . . 145
13.7 Frobenius pairings in tight-frame products . . . 146
13.8 Schur matrices and determinant losses . . . 148
13.9 Application to the prescribed weights . . . 151
13.10 From theta products to the standard integral kernel data . . . 153

14 The definite chain, collisions and Hecke repair **155**
14.1 Coefficient lattices, class spaces and index . . . 155
14.2 The groups and the chain of theta kernels . . . 157
14.3 The geometric seesaw and its denominator control . . . 159
14.4 Ramification and short generators on the definite group . . . 162
14.5 The collision loss and the marked set . . . 163
14.6 Moving the marked Hecke operators through the chain . . . 165
14.7 The Frobenius-twisted elementary-divisor theorem . . . 168
14.8 Application to the actual column lattice . . . 174
14.9 The prime N: moving denominators to a Fourier operator . . . 176
14.10 The simultaneous local divisibility statement . . . 178
14.11 Archimedean growth and the level normalization . . . 179
14.12 Projection, product formula scaling and Theorem A . . . 180

15 The local Hecke estimate in full detail, and the integral seesaw **182**
15.1 The cyclic coefficient representation and the two filtrations . . . 182
15.2 The root estimate, including its coefficient extraction . . . 183
15.3 The dimension bound from flags and convex volume . . . 185
15.4 Deep coefficients: a weighted Nullstellensatz and its application . . . 189

15.5 Integral diagonal pairings and class means 191
15.6 Frobenius products of matrices and the approximate projector 193
15.7 Primitive half-depth matrices 196
15.8 The Smith pairing lemma and the local Hecke theorem 198
15.9 The finite Fourier operator at the level prime 200
15.10 Geometry and integrality in passing to a seesaw 203

16 Normalized CM reciprocity and the geometry of the iterated seesaw 206
16.1 A marked algebraic curve and its sections 206
16.2 Simultaneous torsion translations and finite-adelic reciprocity 209
16.3 Removing one string: the Hermitian linear algebra 211
16.4 The surviving rational orbit 212
16.5 The one-step period transfer 215
16.6 The glued lattice and integral Witt extension 218
16.7 Adapted bases and equality of lattice lines 220
16.8 Local supports of zero terms 221
16.9 A one-step bound for the lattice lines 222
16.10 Finite cuts with prime-supported multiplicity 225
16.11 Two edges at once: removal of the extra plane 227
16.12 The split neighbour lattice in the paired evaluation 229
16.13 Weighted integrality of evaluations 230
16.14 The admissible combinations and the iterated identity 232
16.15 The remaining representation-theoretic comparison 234

17 Compact theta coefficients, integral branching and conjugation of the arithmetic vectors 235
17.1 The entire function associated with the finite vector 235
17.2 From CM theta conjugation to the finite Weil representation 237
17.3 The explicit conjugate-jet operator bound 240
17.4 Stability of the arithmetic data and the gauge calculation 241
17.5 Compact coefficient lifts: no noncompact joint-harmonic theorem is necessary 242
17.6 The even-stage coefficient identification 245
17.7 Integral branching with a controlled small diagram 246
17.8 The last stage as an actual compact branching integral 249
17.9 Uniform counting of the definite classes 250
17.10 The exact size budget along the lift chain 251
17.11 Status of the upper-bound proof obligations 253

III Spanning: Bessel and Poincare series 255

18 Formal degrees, elliptic trace comparison and the row dimension 256
18.1 Parameters and the compact-dual Haar measure 256
18.2 Evaluation of the constant in the formal degree 257

18.3 The exact elliptic orbital character identity 260
18.4 Absolute convergence and a direct ellipticity argument 261
18.5 Changing the signatures of the rational elliptic classes 263
18.6 The trace of the elliptic finite test functions 264
18.7 Primitive labels and the finite cubic correction 265
18.8 Equality of all good Hecke traces and determinants 267
18.9 The precise row-dimension inequality 268

19 A trace-class derivation of the normalized elliptic comparison 271
19.1 A uniform local orbital majorant 271
19.2 The degree factor in the orbital trace 272
19.3 The stable Weyl sum 273

20 Quantitative spanning by projected Bessel–Poincaré series 275
20.1 Statement and finite-dimensional framework 275
20.2 Finite level, central projection, and the transfer bound 276
20.3 Hermitian geometry of the Bessel subgroup 278
20.4 The archimedean Bessel system 279
20.5 Construction and exact unfolding of the Poincaré map 282
20.6 Energy, pairs of congruent terms, and removal of lower cells 285
20.7 The sharp coherent main term 288
20.8 A finite-field positivity argument for singular minors 292
20.9 First moment and bad finite-type loss 294
20.10 From the moment bounds to the covering theorem 295

21 The analytic and arithmetic mechanism of the Bessel sieve 298
21.1 Norm balls in the local groups 298
21.2 A rank-one convolution estimate and its differentiated form 300
21.3 Threshold projectors and a lowest-type coefficient 302
21.4 Two-factor Laplace estimate 304
21.5 Interpolation, isotropic frames, and the Bessel peak 306
21.6 A peak-to-cut inequality with all mass factors 309
21.7 From an L^2 reproducing estimate to the bandlimit tower 311
21.8 The cell L^2 estimate and its limited differentiation cost 313
21.9 A finite-subset version of adelic Poisson summation 314
21.10 An explicit induction for the sieve system 315
21.11 Frequency summation and the product formula 317
21.12 Finite packets, their normalization, and the global comparison 318
21.13 Degenerate terms and the accumulation of energy errors 320

22 Quantitative Bessel construction and the arithmetic of the filling datum 322
22.1 The compact integral and the change of splitting 322
22.2 Logarithmic principal-minor estimates by resolvents 325
22.3 Gaussian shape of the coefficient 327

22.4 Determinant decay and a small-ball bound . . . 328
22.5 Removal of the common compact subgroup . . . 329
22.6 A phase-cancelling realization of the Bessel group . . . 332
22.7 Marked integral lattice and the exact finite level indices . . . 335
22.8 Haar coordinates and the arithmetic unipotent covolume . . . 337
22.9 Cutoffs, approximation and the quotient character . . . 340
22.10 Orbital cutoffs and Bessel transport between classes . . . 342
22.11 Agreeing characters and propagation of a shared flag vector . . . 344
22.12 The link as a Hermitian Lagrangian and its congruence defect . . . 346
22.13 Saturated lattice heights and their zeta sum . . . 348
22.14 The global frequency estimate and explicit sieve budget . . . 351
22.15 The finite zero form and central projections . . . 353
22.16 The projection covariance and the complement-coset error . . . 355
22.17 Assembly of the sharp energy bound . . . 356
22.18 What remains of the filling argument . . . 357

23 Closing the global-to-local and finite-packet comparisons 359
23.1 A normalized representation-density formula . . . 359
23.2 Uniform small error from the places outside the window . . . 362
23.3 One-sided counting on the middle quadric . . . 362
23.4 Term positions, finite characters, and the exact cutoff identity . . . 364
23.5 A local packet lemma . . . 366
23.6 The normalized coherent packet bound . . . 368
23.7 The limit over the finite adelic windows . . . 370
23.8 The slice calculation underlying floor Plancherel . . . 371
23.9 Two complex gauge estimates . . . 373
23.10 References for the real representation-theoretic interface . . . 374
23.11 Obligation-level ledger . . . 375

24 Finite gauge, orbit measures and the top-frequency transform 377
24.1 The valuation integral and its cone consequence . . . 377
24.2 The common-volume finite gauge . . . 378
24.3 Actual finite correlation supports . . . 380
24.4 Exact invariant measures of the stacks . . . 381
24.5 The exact top-floor Fourier identity . . . 382
24.6 Finite Haar masses and the pairing norm . . . 384
24.7 The finite bottom estimate and its remaining error term . . . 386
24.8 Obligation-level status . . . 387

25 Construction and rigidity of the real discrete-series coefficient 388
25.1 A compact averaging formula for the explicit integral . . . 388
25.2 The matrix coefficient and its differentiated module . . . 390
25.3 A quantitative compact Casimir argument . . . 392

25.4 Analyticity and representation-independent rigidity . . . 395
25.5 Differentiated Schur estimate for the Bessel space . . . 397
25.6 The Gram-section geometry and its exact Jacobian . . . 399
25.7 A quantitative tube and the Sobolev restriction . . . 403
25.8 The randomized open-cell bound . . . 405

26 Green labels, quantitative finite mass, and further finite-place comparisons 408
26.1 The class functions actually used for the labels . . . 408
26.2 Irreducibility, projections, and finite twists . . . 410
26.3 Exact character degree and quantitative bad share . . . 412
26.4 Primitive labels and cubic avoidance . . . 413
26.5 An integral pivot calculation for the actual finite kernel . . . 414
26.6 Weighted square estimates at a finite standard level . . . 415
26.7 Closing the finite bottom error . . . 420
26.8 Status of the continued finite interfaces . . . 421

27 Rational frames, amplitude convergence, and pair discriminants 422
27.1 Packing the arithmetic frame boxes . . . 422
27.2 Uniform summability and the cut error . . . 425
27.3 The determinant of a regularized orthogonal complement . . . 427
27.4 The upper bound for integral link lifts . . . 429
27.5 The lower bound modulo the positive middle spaces . . . 432
27.6 Common links and the simultaneous pair estimate . . . 433
27.7 Obligation status for the filling argument . . . 436

28 From the actual steep datum to a cut Bessel system 438
28.1 The weight pattern of the actual row . . . 438
28.2 Peak and decay on the Bessel quotient . . . 439
28.3 The exact weighted majorant and normalization . . . 441
28.4 Schur and degree bounds in the actual normalization . . . 442
28.5 Finite-place construction of the standard level . . . 443
28.6 Simultaneous numerical admissibility . . . 446
28.7 Fitting the odd graded space and reproducing the Poincaré series . . . 449
28.8 Scope and remaining interfaces . . . 450

29 Uniform floor estimates at finite and complex places 451
29.1 Conventions and the tower . . . 451
29.2 A quantitative Heisenberg integral . . . 452
29.3 Direct and inverse weights on an arbitrary prefix . . . 453
29.4 Fourier estimates on the tower . . . 455
29.5 Integrated square norms . . . 456
29.6 The central estimates for forms . . . 457
29.7 The matrix recovery argument and summation over shells . . . 460
29.8 Descent and the actual sieve systems . . . 464

29.9 Scope of the closure . . . 467

30 Weighted index counting and the crude energy mechanism 469
30.1 An inverse-index counting lemma . . . 469
30.2 The labeled near terms and their middle congruences . . . 470
30.3 Counting the remaining top rows . . . 472
30.4 The uniform weighted count over all labeled terms . . . 473
30.5 The improved marked-class discriminant sum . . . 474
30.6 The pair-weight estimate . . . 476
30.7 The energy estimate once the local class bound is available . . . 477
30.8 The real volume kernel used in the thin-class Schur estimate . . . 480
30.9 The exact thin-class amplitude bound . . . 482
30.10 The large-discriminant block, with its actual cell mass . . . 485
30.11 Obligation ledger for this continuation . . . 487

IV Lower bounds: periods, congruences and crystalline defects 489

31 The determinant lower bound 490
31.1 Conventions for norms of conjugate coefficient vectors . . . 491
31.2 Heights with varying finite-place entries . . . 492
31.3 The determinant lemma with index . . . 493
31.4 Special-value data and fit twists . . . 494
31.5 The correspondence of complementary rank and the finite square . . . 496
31.6 The two-grid exchange and the crude rows . . . 499
31.7 The triangle of weighted lifts . . . 502
31.8 Comparison forms for one lift . . . 505
31.9 Compressed Gram determinants . . . 508
31.10 The height saving in a rank-one special value . . . 510
31.11 Combining the special values and closing the lower bound . . . 513

32 Period identities, finite matching, and the geometry of the lower bound 515
32.1 Complementary frames and matched data . . . 515
32.2 Fourier transformations of the matching identity . . . 518
32.3 The graph term in doubling . . . 521
32.4 Sharp support, level factors, and the common scalar . . . 523
32.5 A rank-one Euler factor test . . . 525
32.6 Hyperbolic-plane removal and the signed product . . . 526
32.7 What makes an exchange an identity on all partners? . . . 527
32.8 Level orders in the paths . . . 529
32.9 Why a false lift with one common root contributes no period . . . 530
32.10 Two finite ingredients of integral comparison . . . 532
32.11 Limits of the reconstruction . . . 534

33 The exchange construction, the two grids, and comparison lattices 535
33.1 Local integrals of complementary frames 535
33.2 Generating all partner data by stable spaces 538
33.3 Construction of the real data on an enlarged Hermitian space 540
33.4 A plane removal on determinant vectors 543
33.5 How the equal-rank normalization is obtained 545
33.6 From a convergent exchange to an operator square 547
33.7 Terminal interlacing and the signatures of the two grids 548
33.8 The finite square move with no accumulated denominator 550
33.9 Moving the descent along the path 552
33.10 The divided lattice that glues the two comparison groups 553
33.11 The loss of the Cartan product 555
33.12 Descent of a glued point form 556
33.13 Rank-one reflections and the small-root descent argument 558
33.14 The comparison map and the precise scalar it sees 560
33.15 The Archimedean budget of the comparison operator 562
33.16 Two further finite calculations for the rank-one values 564
33.17 Summary of the strengthened reconstruction in C 566

34 Closing the period, transfer, and depth calculations 568
34.1 Vanishing at a nonnegative nondegenerate centre 568
34.2 The two line slices and the conductor factor 569
34.3 Unramified polynomial transfer 572
34.4 The normalized rank-one pairing 573
34.5 The depth indicator: sharpness and six-fold decompositions 575
34.6 Packing loss and realization of an indicator 578
34.7 Exact-source obligation ledger 581

35 Exact arithmetic pairings, rank-one values, and spectral projection 583
35.1 Integral Gram pairings on the actual adelic coefficient lattice 583
35.2 Positivity of the rank-one value ratios 585
35.3 Conjugate norm bounds for good translates and monomial vectors 586
35.4 Hecke adjoints and projection at every embedding 589
35.5 The equivariant difference of complementary theta integrals 591
35.6 Equal-rank constancy via strong approximation 594
35.7 Convergence in the operator square and its remaining dependency 595
35.8 Terminal branching and the integral end period 597
35.9 Updated obligations for the period comparison 602

36 Clean doubling, sharp Euler factors and normalized line integrals 604
36.1 Clean data remove all nongraph orbits 604
36.2 The Rallis identity with absolute convergence 606
36.3 Finite vacuum cutoffs at arbitrary finite places 607

36.4 Sharp local data and the absolutely convergent Euler sum 609
36.5 The normalized semilinear rank-one kernel 611
36.6 Conjugate norms and the exact flat error 613
36.7 Integral normalized value measures 615
36.8 Weighted bounds for the real mixed slices 617
36.9 The normalized line orbit and the two finite slices 619
36.10 Absolute Mellin endpoints of the line constant term 623
36.11 Exact-source status of this chapter 625

37 The singular-coefficient induction and the standard-column reduction 626
37.1 Centres, phases and bordered coefficients 626
37.2 The radical hypothesis 627
37.3 The exact determinant-index induction 628
37.4 Application to an equivariant lift 630
37.5 Promotion from the primitive coefficient to the standard datum 632
37.6 Closure record and handbook references 633

38 Complementary-rank exchange: real sections, matched coefficients and the line descent 634
38.1 The mixed name and its exact finite support 634
38.2 The standard radical and Levi term 635
38.3 Absolute coefficient extraction for the standard square name 636
38.4 A smooth section of real mixed evaluation 637
38.5 Smooth Witt motion near a nondegenerate frame 638
38.6 The top exchange real matching construction 638
38.7 The missing adapted-frame scalar in the line calculation 639
38.8 Matching the finite and infinite periods with their Haar measures 640
38.9 The small-side tower descent on standard-column data 643
38.10 Circle vanishing for the negative-definite comparison 644
38.11 Nonvanishing and independence of the equal-rank ratio 645
38.12 Exact-source status and the remaining tower distinction 647

39 Further carrier estimates and the rank-two period route 649
39.1 Holomorphic maximum away from the thin part 649
39.2 Circle projection and polynomially growing kernels 650
39.3 A sparse coefficient estimate in the Fock model 652
39.4 Short bases, the sparse lattice threshold, and reproduction 654
39.5 The carrier comparison with quantitative error 656
39.6 Uniform rank-two jet clearing and the integral family 658
39.7 A height consequence with the exact flat error 660
39.8 The deep-level Poincaré restriction tail 661
39.9 Status of the relevant obligations 662

40 Congruences, auxiliary primes, and the fit family **664**
40.1 The integral projection problem . . . 664
40.2 Labels and Galois representations of lifted eigensystems . . . 665
40.3 Selmer modules over finite coefficient rings . . . 668
40.4 New level characters and tied cocycle conditions . . . 669
40.5 Shrinking several tied Selmer modules . . . 671
40.6 Finite presentations and interpolation of lengths . . . 673
40.7 Degree-free selection of roots of unity . . . 674
40.8 The global twisting theorem and value families . . . 676
40.9 The Tate characters and triangular Taylor–Wiles places . . . 678
40.10 The first-order Ribet construction . . . 680
40.11 Hecke operators realizing the first-order relations . . . 683
40.12 The denominator bound from the local length budgets . . . 685
40.13 Global numerical summation and proof of the fit theorem . . . 687

41 The foundations of the congruence argument **689**
41.1 From weak Hecke compatibility to a trace identity . . . 689
41.2 Tame labels and full matrix span . . . 692
41.3 A duality calculation at the mirror . . . 694
41.4 Anticyclotomic primes from Kummer extensions . . . 696
41.5 Odd laws and the triangular removal of even classes . . . 697
41.6 The algebra of adjoining relations . . . 701
41.7 Passing from scalar rows to matrix blocks . . . 702
41.8 Two applications of Cramer's rule over a family . . . 703
41.9 A transfer calculation at one auxiliary level . . . 704
41.10 Separation of the arithmetic and algebraic budgets . . . 707

42 Arithmetic of the congruence argument: cyclotomic classes, descent and simultaneous twists **708**
42.1 The integer in the odd-class budget . . . 708
42.2 The Jacobi family and its distribution relation . . . 710
42.3 The derivative cocycles from the Jacobi family . . . 712
42.4 The odd component, local vanishing and reciprocity . . . 714
42.5 The Kolyvagin–Rubin detection argument . . . 716
42.6 Uniform detection for the small primes . . . 718
42.7 Returning from Dirichlet components to the CM field . . . 719
42.8 Uniform cardinality of Selmer pencils . . . 721
42.9 Dropping ties along a line of approximate characters . . . 723
42.10 The two specializations in the common-twist theorem . . . 724
42.11 Weak attachment: the trace and descent reductions . . . 727
42.12 Hecke-polynomial continuations . . . 731
42.13 Rank-one normalization of the projection bound . . . 733
42.14 Dependency status of the congruence argument . . . 734

43 The remaining global-duality and Hecke interfaces **735**
43.1 The precise uses of classical duality . . . 735
43.2 Construction of an odd law with a primitive value . . . 736
43.3 From Ribet polynomials to integral operators . . . 740
43.4 The local one-prime hypothesis and its algebraic consequence . . . 741
43.5 The realization of a local first-order tie . . . 743
43.6 The chain orthogonality, including multiplicities . . . 746
43.7 Integral value measures at the actual auxiliary levels . . . 748
43.8 Passage to Galois value families . . . 751
43.9 Obligation ledger: Selmer groups, value families, and attachment . . . 752

44 Automorphic Galois representations by classical base change, and the balanced row **755**
44.1 The field and group to which base change applies . . . 755
44.2 The parity correction and exact spherical attachment . . . 757
44.3 The exact tame label and the monodromy condition . . . 760
44.4 Iwahori exponents and simultaneous Frobenius powers . . . 763
44.5 The primitive label and integral matrix span . . . 765
44.6 The balanced-lattice estimate in full . . . 767
44.7 Row traces for all twists and the rigidity reduction . . . 770
44.8 Obligation ledger for this replacement route . . . 771

45 The first-order Ribet polynomial over the integral coefficient algebra **772**
45.1 Fitting ideals and the cost of imposing linear forms . . . 772
45.2 The triangular block theorem . . . 773
45.3 Selection of a distinguished block determinant . . . 774
45.4 An integral two-adjugate calculation . . . 775
45.5 The abstract first-order theorem . . . 775
45.6 Closure status . . . 780
45.7 The triangle cost for a Tate string . . . 780

46 The global mirror criterion for tied Selmer groups **783**
46.1 The isotropic pairing estimate . . . 783
46.2 The dual Selmer group as the family of mirrors . . . 784
46.3 The split-prime test gives isotropy on the required coset . . . 786
46.4 Absorbing the scalar loss into the balance bound . . . 788

47 The arithmetic Ribet polynomial for columns with ties **789**
47.1 Torsion cocycles as reductions of rational coboundaries . . . 789
47.2 Local duality and the defect of the crystalline relations . . . 790
47.3 The common tame class and sum of the local block losses . . . 791
47.4 The conjugate-inverse trace ties . . . 792
47.5 Integral arithmetic specialization and valuation budget . . . 792
47.6 Status . . . 794

48 From the labelled finite level to a denominator operator 795
48.1 The base level and its arithmetic labels . . . 795
48.2 The integral trace functions . . . 796
48.3 A cyclotomic element and a fixed integral cutoff . . . 797
48.4 The prime neighborhoods and the string condition . . . 797
48.5 The tested complement and the three order factors . . . 798
48.6 Status . . . 799

49 Local crystalline defects and the integral cup-product estimate 800
49.1 Lattices, crystalline cocycles and imperfectly dual reductions . . . 800
49.2 Why this defect bounds the local discrepancy of congruent cocycles . . . 803
49.3 The Kummer chart, letter decomposition and height of a frame . . . 804
49.4 Exponent lattices and the cost of increasing depth . . . 806
49.5 Cokernel estimates: divisibility by p and by the Eisenstein element . . . 808
49.6 Cup-product vanishing from the three sides . . . 809
49.7 The window estimates: the three ways to obtain safe points . . . 812
49.8 The tame type bound, uniform in the ramification degree . . . 815
49.9 Case analysis for the primes and the explicit local loss . . . 816
49.10 Insertion of the global weights and the sum over all blocks . . . 817

50 Integral period frames and the algebra of the local loss 820
50.1 The rank calculation and duality over a finite coefficient ring . . . 820
50.2 The coordinates in a Kummer chart . . . 822
50.3 A coefficient retraction and why it is used . . . 824
50.4 From crystalline periods to an integral frame . . . 825
50.5 The radius estimate and its fourth power . . . 828
50.6 Exponent lattices, Frobenius letters and the small part . . . 830
50.7 Exact cokernel multipliers via finite differences . . . 832
50.8 A rank-sensitive bound for raising filtered conditions . . . 834

51 Integral crystalline comparison, depth operators and the Frobenius walk 838
51.1 The period rings and an integral contraction . . . 838
51.2 The period-basis argument . . . 840
51.3 Reduction of the determinant and the integral frame . . . 843
51.4 Scalar depth steps at each Frobenius divisor . . . 845
51.5 Depth zero and finite averaging . . . 847
51.6 From scalar depths to crystalline lattice depths . . . 848
51.7 From a single operator to the cost of a window . . . 851
51.8 The monic Taylor coefficient at a long gap . . . 852
51.9 The triple pairing and the safety rule . . . 855
51.10 From safe pairs to an actual cup coboundary . . . 857
51.11 The walk along the Frobenius permutation . . . 859
51.12 Plain cuts and long-gap cuts . . . 861

51.13Quantitative killers at sparse cut positions . . . 863
51.14Two choices of cuts . . . 865
51.15A multiplicity-sensitive step in the few-low-jumps case . . . 867
51.16Scope of the integral lifting reconstruction . . . 870

52 Closing the local comparison and lifting arguments 871
52.1 Completeness and the elementary maximal period ring . . . 871
52.2 Retractions and the scalar ring of the chart . . . 873
52.3 Fixed coefficients and period-function retractions . . . 875
52.4 The first radius lemma: constant times a unit . . . 877
52.5 The actual Kummer charts and their congruences . . . 879
52.6 Construction and identification of the crystalline frame . . . 880
52.7 Galois differences and the multiplier ideals . . . 883
52.8 The tame chart and the polynomial recurrence . . . 884
52.9 Initial depth and final safety for few low jumps . . . 886
52.10Exact compact-source numerical hypotheses for the cut bounds . . . 890
52.11Functorial frame operations used in the lifting proof . . . 891
52.12Both alternatives of the no-multiple bound . . . 892
52.13From side pairings to the local defect theorem . . . 894
52.14Obligation-by-obligation closure record . . . 896

53 The quantitative crystalline loss for the actual weight windows 899
53.1 A numerical consequence of the integral cutoff theorems . . . 899
53.2 Cyclotomic coordinates of thick places . . . 902
53.3 The actual one-place estimate . . . 903
53.4 Status of the local interface . . . 905

V The arithmetic conclusion and proof audit 906

54 A numerical appendix: asymptotic budgets and the passage to the limit 907
54.1 Uniform scale asymptotics . . . 907
54.2 Rounding of large coefficients and a Lipschitz estimate . . . 908
54.3 The exact level cancellation revisited . . . 908
54.4 An elementary bound on the logarithm of the class number . . . 909
54.5 Summation of the fit-family costs . . . 910
54.6 The final numerical inequalities . . . 911

55 The arithmetic inequality and conclusion 912
55.1 Collecting the arithmetic inputs . . . 912
55.2 From A and B to an almost-basis of short forms . . . 913
55.3 Comparison with the lower bound . . . 913
55.4 Extracting the analytic remainder . . . 914
55.5 Choosing the second cutoff and the power saving . . . 915

55.6 GRH 915

56 Glossary and concordance **917**
56.1 Reading the main theorem in the Lean source 918

57 Third proof-closure audit and remaining dependencies **920**
57.1 Scope of this convergence pass 920
57.2 Focus of this edition 920
57.3 Interface ledger 921

Part I

The analytic reduction and arithmetic models

Chapter 1

Introduction and mathematical status

The compact edition of the Alpöge–Claude formal development [1] is a remarkably large argument for the following assertion.

Theorem 1.1 (Dirichlet GRH; asserted in the formal source)**.** *For every integer $q \geq 1$, every Dirichlet character χ modulo q, and every $s \in \mathbb{C}$ with $\Re(s) > 1/2$,*

$$L(s, \chi) \neq 0,$$

where the pole of a principal character at 1 is excluded. In particular every zero in $0 < \Re(s) < 1$ lies on the line $\Re(s) = 1/2$, and the nontrivial zeros of ζ lie on this line.

The qualifier Dirichlet is important: the paper concerns the L-functions of degree one over $\mathbb{Q}$, not the generalized hypothesis for all automorphic L-functions. The proof uses higher rank unitary groups over auxiliary CM fields, but it uses them to construct and compare algebraic objects, not to assert GRH for their standard L-functions.

This monograph is the third stage of the series. It retains the interfaces of the shorter specialist treatment, expands their technical proofs, and supplies additional mathematical background, examples, and context for readers trained in neighboring areas. Definitions are fixed in their operative form, and references connect each mathematical package to the relevant part of the compact Lean development. It is a *working reconstruction of the formal proof*, not a claim that every arithmetic argument has yet been independently translated into a conventional proof. Every assertion called a "formal-source input" is a mathematical statement extracted from, or a stated interface to, the supplied Lean development that still needs an independent written proof or a verified literature replacement. A statement followed by a proof records a mathematical derivation of that statement from the explicitly stated inputs; it does not thereby establish those inputs. In particular the manuscript must not be cited as an independently complete proof of Theorem 1.1 until the remaining formal-source inputs have been discharged. The main outstanding packages, including the exact geometric and integral assertions needed for subsequent deductions, are listed at the end. This convention allows a specialist to test an argument at the level of its precise hypotheses, instead of having to infer whether an omitted detail is formal bookkeeping or a substantial arithmetic claim.

1.1 The architecture of the proof

The principal idea is to use the same automorphic object for two opposed quantities. At the archimedean places it carries a metric; at the finite places it carries an integral coefficient

structure. These are linked by a product formula. To encode primes, consider jets of a theta kernel for a CM elliptic curve in a specified algebraic normalization. Their divisibility at a nonsplit rational prime depends on how the local formal-group parameter transforms under Frobenius. The valuation bounds contain sums of positive parts indexed by powers of the prime. A choice of the infinite-place weights turns their global sum into the weighted prime remainder.

One cannot simply apply the product formula to an arbitrarily chosen jet. The proof instead uses a family of forms obtained through an iterated theta correspondence. Suitable source forms are transferred by a seesaw identity to the lifts of Poincaré series on an indefinite unitary group. The arithmetic choice of fields, levels and Harish-Chandra parameters serves three tasks at once: the finite-place integrality estimate, a nearly sharp second moment for Poincaré series, and a matching lower bound on the determinant of the lift. The latter requires a family of twists, one at each finite embedding, rather than a single twist valid at every place.

Here are the three central estimates, in the notation that will be fixed in Chapter 4. Let W be a row space of dimension n, C the source space of dimension c, and $C^{\mathrm{ram}} \subseteq C$ its ramified part. Let Θ_θ be a composite of m lifts, and $P_s : C^{\mathrm{ram}} \to W$ the projected translates of a Poincaré map, $s \in \mathcal{G}$. The large positive numbers $B = fa(\log b)^3$ and $L = B \log b$ are error budgets, while $H = mk\mathcal{U}$ denotes a signed common main term. The weighted prime remainder is $\mathcal{R}$.

A. There exists a positive rational scaling of the P_s and a spanning set of integral elements of C^{ram} for which all trivial-twist lifts have squared norm at most $\exp(H - m\mathcal{R} + L)$ at every archimedean conjugate. A family from the images of J translates has logarithmic index at most $|J|cB$ simultaneously in the admissible integral structures.

B. Both c and n are positive. There are translates indexed by a finite set J with $|J|c \le 5n \log(1 + b^2)$ whose images span at least $(1 - 4/b^2)n$ dimensions.

C. Every independent family of at least $(1-4/b^2)n$ rows whose trivial-twist lifts have squared covolume at most e^V at all conjugates and whose logarithmic index is at most d_{ind} satisfies $n(H - 3L) \le V + 2d_{\mathrm{ind}}$.

Assertions A, B, C are the interfaces `Data.short_forms`, `Data.enough_forms`, and `Data.integral_forms_long` of the compact source. Their proof decompositions occupy Parts II–IV of the monograph. Notice that C is uniform in the choice of row family, unlike A, and that B is quantitative; neither of these features can be dropped in the endgame.

To see the algebraic mechanism, choose $q \ge (1 - 4/b^2)n$ independent short rows using A and B. Hadamard's inequality gives $V \le q(H - m\mathcal{R} + L)$, while the index is at most $5n \log(1 + b^2)B$. Combining with C, and bounding the missing-dimension term by a crude estimate, gives

$$m\mathcal{R} \le 4L + 4x^{99} + 10 \log(1 + b^2)B \le fa(\log b)^5.$$

The two instances of H cancel exactly. The number m of lifts is of size $\sqrt{x}$; this is the source of the square-root factor in the final saving. The normalized prime remainder $R_{N_0}(x)$ is related to $\mathcal{R}$ by the exact identity

$$\frac{\mathcal{R}}{a} = |\mathcal{S}|(R_{N_0}(x) - R_{N_0}(x_1)) + |\mathcal{T}|R_1(x_1),$$

where $\mathcal{T}, \mathcal{S}$ are two subsets of infinite places. The compact analytic argument chooses x_1 in a long logarithmic window to give a one-sided upper bound for $R_{N_0}(x_1) - |\mathcal{T}|R_1(x_1)/|\mathcal{S}|$. With $|\mathcal{S}| \ge |\mathcal{T}|/\log b$, $f = \varphi(N_0)|\mathcal{T}|$, and $\log b = x^\varepsilon$, one concludes $R_{N_0}(x) \le x^{-1/2+8\varepsilon}$ eventually. A Landau argument then excludes zeros to the right of $\frac{1}{2} + 8\varepsilon$. Quadratic twists are varied to prevent cancellation of the singularities of logarithmic derivatives.

1.2 Primary material, conventions, and the remaining challenge

The supplied snapshot `grh-lean-main` contains the two variants `grh/` (readable) and `grh-compact/` (compact), of approximately 300,000 and 76,000 Lean lines respectively. Their high-level interfaces agree but some proofs differ. We take the compact variant as primary and use the readable one when clarifying its definitions or constituent steps. Unless qualified otherwise, paths in the text are relative to `grh-compact/src/`. The root README, attribution and guide identify the underlying proof as due to Levent Alpöge with Claude and the shortening as a collaboration using Claude. The formal source reports kernel checks of all exported results and dependence only on `propext`, `Classical.choice`, and `Quot.sound`; this manuscript makes no claim of a new independent machine replay. The statement of the result imports only Mathlib's definitions, not the auxiliary arithmetic theory of this argument. The exact goal is in `Statements.lean`, and the assembly is `New.grh` in `New/Main.lean`.

The mathematical difficulty is not measured by Lean line count. In particular the division into `Lib` and `New` is not a proof that all of the former can be replaced by standard citations. Uniformity in growing ranks, large weights and varying auxiliary fields, integral versions of the correspondence, normalization of class sums and precise determinant comparisons must all survive such replacement. We therefore take care to mark specialized inputs instead of silently replacing them by named theorems from the literature. Classical results used in the analytic and parameter-construction sections are identified explicitly and do not presuppose GRH.

Notation. The letter d always denotes the chosen auxiliary rational prime; d_{ind} denotes an index bound. The size b is a scale, not a weight coordinate. We use Φ for the set of embeddings of E above the chosen embedding of K, $\mathcal{T}$ for the tops, and $\mathcal{S}$ for the thick places. We write $A(y,\sigma)$ for the weighted sum in `Data.primeSum`, $A_p(y,\sigma)$ for its normalized p-part, and $J(j)$ for `Data.jetSize`. In some local arguments letters may denote temporary variables, but the main budgets $B, L, H, \mathcal{R}$ retain their meanings. The notation W_j for the auxiliary Hermitian spaces is different from W, the vector space of row forms. The group $\mathcal{G}$ is also denoted $\mathfrak{G}$ in the finite moment estimates; $C_{\mathrm{ram}} = C^{\mathrm{ram}}$.

Chapter 2

The analytic problem: a one-sided bound for weighted prime sums

The next chapter, Chapter 3, supplies the classical analytic prerequisites and the proof of the special prime-window theorem. The final conditional theorem there also verifies the quantifiers in the passage from the three arithmetic inequalities to the assertion for all Dirichlet characters.

Before constructing the automorphic objects, we isolate the precise problem that they solve. The analytic reduction is useful in its own right: it separates the difficult arithmetic estimate from the comparatively short argument about zeros. In particular, the arithmetic construction need not produce a two-sided error estimate for the ordinary prime counting function.

Throughout this section $M \geq 1$ is a modulus, d is a prime with $d \equiv 7 \pmod 8$ and $(d, M) = 1$, and

$$K = \mathbb{Q}(\sqrt{-d}), \qquad \chi_K(\nu) = \Big(\frac{\nu}{d}\Big).$$

The discriminant of K is $-d$; quadratic reciprocity identifies the displayed character with its quadratic Dirichlet character. Write $\varphi(M)$ for Euler's totient. The dependence on d in the notation R_M below is suppressed, but it is essential that the auxiliary prime d can eventually be varied.

2.1 The particular smoothing used in the proof

Let $\Lambda(p^j) = \log p$ for prime powers, and let $\Lambda(\nu) = 0$ otherwise. Put

$$\lambda_{d,M}(\nu) = \mathbf{1}_{\nu \equiv 1 \bmod M} \Lambda(\nu)(1 - \chi_K(\nu)). \tag{2.1}$$

Thus $\lambda_{d,M}$ is non-negative. A prime split in K contributes zero, an inert prime contributes twice its ordinary von Mangoldt weight, and the ramified prime has factor one. The same rule applies to prime powers; for instance an even power of an inert prime has factor zero. Congruence $\nu \equiv 1 \pmod M$ is imposed on the integer ν, not on its underlying prime. In particular, the weighted sum is not literally a sum over primes alone.

For any Dirichlet character ξ, with its modulus understood, denote by

$$\widetilde{L}(s, \xi) = \begin{cases} L(s, \xi), & \xi \text{ non-principal}, \\ (s-1)L(s, \xi), & \xi \text{ principal} \end{cases}$$

the entire function obtained by removing the possible pole at 1 (the second line is interpreted by continuation). The character $\xi\chi_K$ is taken modulo Md; it is non-principal because d is a

new odd prime. Define the meromorphic function

$$G_{d,M}(s) = \frac{1}{\varphi(M)} \left\{ \sum_{\xi \bmod M} \left(\frac{\widetilde{L}'}{\widetilde{L}}(s, \xi\chi_K) - \frac{\widetilde{L}'}{\widetilde{L}}(s, \xi) \right) - 1 \right\}. \tag{2.2}$$

Here and throughout, a sum over characters includes the principal and imprimitive characters. The constant subtracted in the prime error is

$$c_{d,M} = \operatorname{Re} G_{d,M}(1).$$

The classical non-vanishing on $\operatorname{Re} s \geq 1$ of the pole-removed Dirichlet L-functions ensures that this constant exists. In fact $G_{d,M}(1)$ is real, by conjugation of characters.

The error in the compact Lean development, called `primeError`, is exactly

$$R_M(x) = \sum_{1 \leq \nu \leq x} \lambda_{d,M}(\nu) \left(\frac{1}{\nu} - \frac{1}{x} \right) - \frac{\log x}{\varphi(M)} - c_{d,M} \qquad (x > 0). \tag{2.3}$$

It is important to include the constant as well as the logarithm in the normalization. Omitting it would prevent the Laplace transform used below from being regular at $s = 1$. In modulus one, the constant has a more specific expression. If γ is Euler's constant, write

$$\kappa_K = \gamma - \operatorname{Re} \frac{L'}{L}(1, \chi_K).$$

Since $(s-1)\zeta(s) = 1 + \gamma(s-1) + O((s-1)^2)$,

$$c_{d,1} = -1 - \kappa_K. \tag{2.4}$$

The number κ_K, called `chowlaSelbergConst` in the code, recurs in the arithmetic estimates: the appearance of the same constant in the norm of a CM theta period is part of the eventual cancellation.

The choice of smoothing in (2.3) is particularly convenient for the arithmetic proof. For $y > 0$ set $(y)_+ = \max(y, 0)$, and extend this notation to all real numbers. For an integer a and $0 < z \leq a$ one has the exact identity

$$\frac{1}{a} \sum_{\nu=1}^{a} \lambda_{d,M}(\nu) \left(\frac{a}{\nu} - \frac{a}{z} \right)_+ = R_M(z) + \frac{\log z}{\varphi(M)} + c_{d,M}. \tag{2.5}$$

Indeed the positive part vanishes for $\nu \geq z$ and gives $a(1/\nu - 1/z)$ for $\nu < z$. No asymptotic replacement is involved. The importance of this identity is that the arithmetic construction bounds valuations of jets expressed using precisely such positive parts, with integer order parameter a and scale z. The apparently unusual smoothed prime sum is therefore dictated by a divisibility calculation, not just by analytic convenience.

Our ultimate target is, for every $0 < \varepsilon < 1/100$ and every suitable d,

$$R_{N_0}(x) \leq x^{-1/2+8\varepsilon} \qquad (x \text{ sufficiently large}), \tag{2.6}$$

where $18480 \mid N_0$ and $d \equiv N_0/2 - 1 \pmod{N_0}$. Only the upper bound occurs in (2.6). The threshold for x may depend on the fixed parameters N_0, d and ε. We shall explain why such a bound is already strong enough.

Source. The definitions, the constant in modulus one and (2.5) are in `New/Endgame/PrimeError.lean`, declarations `lambdaClass`, `logDerivMean`, `primeError`, `re_logDerivMean_one`, `primeError_eq_sum`. The function $\widetilde{L}$ is `entireL` in `Lib/General/NumberTheory/QuadField.lean`.

2.2 Why the prime sum contains information about zeros

Set

$$E_M(v) = R_M(e^v), \qquad \Phi_M(s) = \int_0^\infty E_M(v) e^{(1-s)v}\, dv.$$

The variable s is the same complex variable as in the L-functions. The initial integral converges for $\operatorname{Re} s > 2$: for $v \geq 0$, the elementary bound $|\lambda_{d,M}(\nu)| \leq 2\nu$ gives $|E_M(v)| \leq Ce^v$, with C depending on M, d.

Proposition 2.1 (Transform and residues)**.** *The integral Φ_M continues meromorphically, with the expression*

$$\Phi_M(s) = \frac{G_{d,M}(s)/s - c_{d,M}}{s-1}. \tag{2.7}$$

It is holomorphic in a neighborhood of the closed half-plane $\operatorname{Re} s \geq 1$. For $0 < \operatorname{Re}\rho < 1$ its residue at ρ is

$$\operatorname{res}_{s=\rho} \Phi_M(s) = \frac{\sum_{\xi \bmod M} \left(\operatorname{ord}_\rho \widetilde{L}(\cdot,\xi) - \operatorname{ord}_\rho \widetilde{L}(\cdot,\xi\chi_K)\right)}{\varphi(M)\rho(1-\rho)}. \tag{2.8}$$

In $\operatorname{Re} s > 0$ any poles are at most simple.

Proof. By orthogonality of characters and the logarithmic derivative of the Euler product, the Dirichlet series of (2.1), absolutely convergent for $\operatorname{Re} s > 1$, is

$$\begin{aligned} D_{d,M}(s) &= \sum_{\nu \geq 1} \frac{\lambda_{d,M}(\nu)}{\nu^s} \\ &= \frac{1}{\varphi(M)} \sum_{\xi \bmod M} \left(\frac{L'}{L}(s,\xi\chi_K) - \frac{L'}{L}(s,\xi)\right) \\ &= G_{d,M}(s) + \frac{s}{\varphi(M)(s-1)}. \end{aligned}$$

In the last equality the unique principal character modulo M contributes $1/(s-1)$ besides the pole-removed logarithmic derivative, while no twisted character is principal. The additional 1 in the definition of $G_{d,M}$ has now combined with that pole term.

Absolute convergence justifies integrating the sum in (2.3) term by term for $\operatorname{Re} s > 2$. For an integer $\nu \geq 1$,

$$\int_{\log\nu}^\infty (\nu^{-1} - e^{-v}) e^{(1-s)v}\, dv = \nu^{-s}\left(\frac{1}{s-1} - \frac{1}{s}\right) = \frac{\nu^{-s}}{s(s-1)}.$$

The transform of v is $1/(s-1)^2$, and that of the constant function one is $1/(s-1)$. Consequently

$$\begin{aligned} \Phi_M(s) &= \frac{D_{d,M}(s)}{s(s-1)} - \frac{1}{\varphi(M)(s-1)^2} - \frac{c_{d,M}}{s-1} \\ &= \frac{G_{d,M}(s)/s - c_{d,M}}{s-1}. \end{aligned}$$

This gives the desired continuation. The apparent singularity at 1 is removable because $c_{d,M} = G_{d,M}(1)$. The classical non-vanishing in $\operatorname{Re} s \geq 1$ rules out all other poles there.

If an entire function F has a zero of order j at ρ, then we can write $F(s) = (s-\rho)^j f(s)$ locally, with $f(\rho) \neq 0$. Hence

$$\frac{F'}{F}(s) = \frac{j}{s-\rho} + \frac{f'}{f}(s).$$

This proves (2.8), including its sign, and shows that the poles are simple. □

The numerator in (2.8) is a *difference* of zero multiplicities. Thus a bound proving that Φ_M has no poles in a certain half-plane would not, by itself, show that every $L(s,\xi)$ is zero-free there: a pole might cancel between the two sums. The remedy is to choose d so that the finitely many twisted L-functions have no zero at a hypothetical offending zero, and no real zero to the right of the relevant line.

In the code the properties of E_M and its transform are packaged by the predicate `IsPolar`. The assertion `isPolar_primeError` in `New/Endgame/PrimeError.lean` establishes this predicate with the residue function (2.8) (zero at 1). This abstraction enables the two subsequent analytic arguments to be proved without referring to primes or automorphic forms.

2.3 An elementary form of Landau's one-sided principle

Here is the precise analytic feature that makes an upper bound sufficient. We give the argument, since it explains the sign required in the later choice of the quadratic twist.

Lemma 2.2 (Landau principle)**.** *Let $E : [0,\infty) \to \mathbb{R}$ be measurable with $|E(v)| \le Ce^v$. Suppose its Laplace transform*

$$\Phi(s) = \int_0^\infty E(v)e^{(1-s)v}\,dv$$

has a meromorphic continuation, holomorphic near $\operatorname{Re} s \ge 1$ and with at most simple poles in $\operatorname{Re} s > 0$. Let $\beta > 0$. Assume that

$$E(v) \le e^{(\beta-1)v} \quad (v \text{ sufficiently large})$$

and that the residue of Φ at each real $\rho > \beta$ is non-negative (or zero when the point is regular). Then Φ is holomorphic in $\operatorname{Re} s > \beta$.

Proof. Choose $A \ge 1$ large enough that

$$h(v) = Ae^{\beta v} - e^v E(v) \ge 0 \qquad (v \ge 0).$$

The eventual upper bound guarantees the inequality at large v, and the growth assumption gives it on the remaining bounded interval after enlarging A. For sufficiently large $\operatorname{Re} s$ we then have

$$\Psi(s) := \int_0^\infty h(v)e^{-sv}\,dv = \frac{A}{s-\beta} - \Phi(s).$$

Let σ_c be the real abscissa of convergence of the integral for Ψ. Because $h \ge 0$, convergence holds in the right half-plane $\operatorname{Re} s > \sigma_c$, where the integral is holomorphic; the displayed meromorphic expression agrees with it there by analytic continuation.

Suppose $\sigma_c > \beta$. If Ψ were to have a simple pole at the real point σ_c, its residue would necessarily be non-negative: for real $s > \sigma_c$, the integral is non-negative, so $(s-\sigma_c)\Psi(s) \ge 0$. But the meromorphic formula gives that the residue equals minus the residue of Φ and is therefore non-positive. This shows that the residue is zero and the singularity removable.

A Laplace transform of a non-negative function cannot be holomorphic at its finite abscissa of convergence. To see this directly, take a real $c > \sigma_c$ sufficiently close to σ_c that the Taylor series of the continued Ψ about c converges at a real $\sigma < \sigma_c$. Differentiation under the integral at c gives

$$(-1)^j\Psi^{(j)}(c) = \int_0^\infty v^j h(v)e^{-cv}\,dv \ge 0.$$

Taylor convergence and monotone convergence therefore give

$$\begin{aligned}\infty &> \sum_{j\ge 0} \frac{(-1)^j \Psi^{(j)}(c)(c-\sigma)^j}{j!} \\ &= \int_0^\infty h(v) e^{-cv} \sum_{j\ge 0} \frac{((c-\sigma)v)^j}{j!}\, dv \\ &= \int_0^\infty h(v) e^{-\sigma v}\, dv,\end{aligned}$$

contrary to the definition of σ_c. Thus $\sigma_c \le \beta$, and the formula for Ψ implies the assertion about Φ on $\operatorname{Re} s > \beta$. □

This argument is based on positivity of an auxiliary integrand, not on a hypothesis that E changes sign or has small absolute value. It is the usual Landau principle for Dirichlet integrals with one sign, in the form formalized by `IsPolar.landau` in `Lib/Analytic/Landau/PolarLandau.lean`.

2.4 Quadratic twists avoiding a prescribed half-strip

We now give a proof of the zero-density ingredient in the form required here. In particular it is not necessary to invoke a zero-density theorem with constants that are uniform in the original modulus M. Fix M throughout this subsection. The following elementary argument follows the compact source `Lib/Analytic/QuadraticTwist/ZeroFreeTwist.lean`. Jutila's classical work on quadratic-character mean values [3] provides the context; the particular assertion needed here is proved below.

Put $g_j(t) = t^j e^{-\pi t^2}$ for $j = 0, 1$, and, for a sequence c of modulus at most 1, put

$$\mathcal{V}_j(c;t) = \sum_{h\ge 1} c(h) g_j(ht), \qquad I_j(c;T) = \int_{1/T}^\infty |\mathcal{V}_j(c;t)| \frac{dt}{\sqrt{t}}, \qquad T > 0.$$

The following properties of the Gaussian are used repeatedly:

$$|g_j(t)| \le 3e^{-2t} \quad (t \ge 0), \tag{2.9}$$

$$|g_j(t)| \le C_\alpha t^{-\alpha} \quad (t > 0,\ \alpha > 0), \tag{2.10}$$

$$|\mathcal{V}_j(c;t)| \le 3e^{-t}/t. \tag{2.11}$$

For the first inequality use $t^j \le 1 + t \le e^t$ and maximize $3t - \pi t^2$; this maximum is $9/(4\pi) < \log 3$. The second follows from boundedness of $t^\alpha g_j(t)$. For the third, summing the first gives $3e^{-2t}/(1 - e^{-2t}) = 3/(e^{2t} - 1) \le 3e^{-t}/t$, since $2\sinh(t) \ge t$.

Lemma 2.3 (Theta transformation and Mellin integral)**.** *Let ψ be a primitive nonprincipal Dirichlet character of conductor q, and choose $j \in \{0, 1\}$ by $\psi(-1) = (-1)^j$. The Mellin transform on $\Re s > 0$ of $\mathcal{V}_j(\psi;t)$ is*

$$M_\psi(s) = \int_0^\infty \mathcal{V}_j(\psi;t) t^{s-1}\, dt = \Gamma_j(s) L(s,\psi), \qquad \Gamma_j(s) = \tfrac{1}{2}\pi^{-(s+j)/2}\Gamma((s+j)/2).$$

Moreover

$$\sqrt{q}\, t\, |\mathcal{V}_j(\psi;t)| = |\mathcal{V}_j(\psi;1/(qt))|, \tag{2.12}$$

and if $q \le T^2$ then

$$\int_0^\infty |\mathcal{V}_j(\psi;t)| \frac{dt}{\sqrt{t}} \le 2 I_j(\psi;T). \tag{2.13}$$

Proof. Poisson summation on each progression modulo q applies to the Schwartz function g_j on $\mathbb{R}$, whose Fourier transform (with kernel $e^{-2\pi ixy}$) is $(-i)^j g_j$. Expanding the periodic character in finite Fourier coefficients, the Gauss-sum formula for a primitive character gives

$$\mathcal{V}_j(\psi;t) = \frac{\varepsilon(\psi)}{\sqrt{q}\,t}\,\mathcal{V}_j(\overline{\psi};1/(qt)), \quad |\varepsilon(\psi)| = 1.$$

Indeed the two bilateral sums are twice their positive parts, by parity and $\psi(0) = 0$; the modulus of the primitive Gauss sum is $\sqrt{q}$. As g_j is real, the absolute values for ψ and $\overline{\psi}$ coincide. This proves (2.12). At zero the transformation and (2.11) show rapid decay; at infinity (2.11) applies directly. Termwise Mellin integration on $\Re s > 1$ yields the displayed formula, by substituting $u = ht$ and then $z = \pi u^2$. Both sides are holomorphic on $\Re s > 0$, so the equality extends there. Finally the substitution $u = 1/(qt)$ in the integral over $(0, 1/T)$, using (2.12), identifies it with the integral over $(T/q, \infty)$ against $du/\sqrt{u}$. Since $T/q \geq 1/T$, the result follows. □

This is the ordinary theta Mellin formula; compare [2, Ch. 10] and the formal declarations `gaussianMellin_eq`, `norm_gaussianTheta_functional_equation`, `thetaTailIntegral_zero_le_two_mul` in `Lib/Analytic/DirichletTheta/MellinTheta.lean`.

Proposition 2.4 (A three-lines zero detector)**.** *For $j = 0, 1$ there is an absolute constant C_j such that, whenever $\eta > 0$, $T \geq 1$, $q \leq T^2$ and ψ is as in Lemma 2.3, the conditions*

$$I_j(\psi;T) + I_j(\mu\psi;T) \leq T^{2\eta}, \qquad L(s,\psi) = 0, \quad \tfrac{1}{2} + 9\eta \leq \Re s \leq 1$$

imply $|\Gamma_j(s)|^2 \leq C_j T^{-2\eta}$.

Proof. Here μ is the Möbius function. Define a truncated Mellin integral

$$G(s) = \int_{1/T}^{\infty} \mathcal{V}_j(\mu\psi;t)t^{s-1}\,dt, \qquad F(s) = \Gamma_j(s)^2 - M_\psi(s)G(s).$$

It is holomorphic and bounded on the closed strip $\frac{1}{2} \leq \Re s \leq 2$ (with a bound depending on q, T): this follows from the integral estimates just proved, the cutoff in G, and the Gaussian Mellin formula. We use the three-lines theorem with boundary estimates uniform in the imaginary coordinate. On $\Re s = 1/2$, write $I = I_j(\psi;T)$, $J = I_j(\mu\psi;T)$. Lemma 2.3 implies $|M_\psi(s)| \leq 2I$ and $|G(s)| \leq J$. Thus

$$|F(s)| \leq \sup_{\Re z = 1/2} |\Gamma_j(z)|^2 + 2IJ \ll_j T^{4\eta}.$$

The bound on the gamma factor is immediate from its defining integral. On $\Re s = 2$, absolute convergence of the Euler products gives $L(s,\psi)L(s,\mu\psi) = 1$, where the second factor denotes $\sum_{n\geq 1} \mu(n)\psi(n)n^{-s}$. Therefore

$$F(s) = M_\psi(s) \int_0^{1/T} \mathcal{V}_j(\mu\psi;t)t^{s-1}\,dt.$$

By (2.11), the first factor is bounded in modulus by $3\int_0^\infty e^{-t}\,dt = 3$, and the second by $3/T$. Hence $|F(s)| \leq 9/T$, which in particular is $\ll_j T^{4\eta-1}$. Interpolation across the strip of width $3/2$ gives

$$|F(s)| \ll_j T^{4\eta-(\Re s-1/2)/(3/2)}.$$

When $L(s,\psi) = 0$, $F(s) = \Gamma_j(s)^2$. For $\Re s \geq 1/2 + 9\eta$, the exponent is at most -2η, as required. □

The formal counterpart is `three_lines_bound_of_LFunction_eq_zero`.

Lemma 2.5 (Smoothed quadratic-character mean value)**.** *For $\eta > 0$, $j = 0, 1$, $X \in \mathbb{N}$, $0 < T$ with $X \le T^2$, $t \ge 1/T$, and $|c(h)| \le 1$, one has*

$$\sum_{1 \le v \le X} \left| \sum_{h \ge 1} c(h) J(-v \mid h) g_j(ht) \right|^2 \ll_\eta T^2 t^{-1-2\eta}.$$

Here $J(v \mid h)$ is the product of the Legendre symbols over odd primes dividing h, each with its multiplicity, and the principal character at 2 when h is even. In particular it is multiplicative in the denominator and bounded in absolute value by 1.

Proof. After expanding the square (absolutely convergent by (2.11)), it is enough to bound

$$\sum_{n,m \ge 1} |g_j(nt) g_j(mt)| \left| \sum_{v \le X} J(-v \mid nm) \right|.$$

If neither nm nor $2nm$ is a square, the Dirichlet character $v \mapsto J(v \mid nm)$ is nonprincipal modulo nm: its odd-prime squarefree kernel is nontrivial. The Pólya–Vinogradov inequality gives the inner bound $\ll \sqrt{nm}(1 + \log nm)$. Use (2.10) with exponent $3/2 + \eta$ for both Gaussian factors and the estimate $1 + \log(nm) \ll_\eta (nm)^{\eta/2}$. The sum of the resulting factors $(nm)^{-1-\eta/2}$ is finite. The contribution of this part is $\ll_\eta t^{-3-2\eta} \le T^2 t^{-1-2\eta}$.

If knm is a square for some $k \in \{1, 2\}$, use the trivial bound $X \le T^2$. Write $kn = au^2$, $m = av^2$ with $a \ge 1$ squarefree; each pair occurs in this parametrization for the indicated k. Use (2.10) with exponent $1/2 + \eta$. Ignoring the squarefree restriction on a and the divisibility requirement on u, the sum is bounded, for each k, by

$$C_\eta T^2 t^{-1-2\eta} k^{1/2+\eta} \sum_{a,u,v \ge 1} (auv)^{-1-2\eta}.$$

This proves the assertion. We have used only the classical Pólya–Vinogradov bound for a (possibly imprimitive) nonprincipal character, not a quadratic large sieve. The finite-character inequality can be found in [2, Ch. 9]. □

Corollary 2.6 (Tail-integral bound)**.** *For $X \le T^2$ and $|c(h)| \le 1$,*

$$\sum_{1 \le v \le X} I_j(c(\cdot) J(-v \mid \cdot); T) \ll_\eta T^{2+\eta}.$$

Proof. By Cauchy–Schwarz and Lemma 2.5, the sum over v of the absolute values inside the integral is $\ll_\eta \sqrt{X} T t^{-1/2-\eta} \le C_\eta T^2 t^{-1/2-\eta}$. Integrating against $dt/\sqrt{t}$ from $1/T$ to infinity gives T^η/η times this constant. □

Proposition 2.7 (Zero density in the family of prime quadratic twists)**.** *Fix a Dirichlet character ξ modulo M, $Y \ge 0$, and $0 < \eta < 1/18$. The number of primes $d \le X$, $d \equiv 7 \pmod 8$, $(d, M) = 1$, for which $\widetilde{L}(s, \xi\chi_K)$ has a zero in $\Re s \ge 1/2 + 9\eta$, $|\Im s| \le Y$, is $O_{M,Y,\eta}(X^{1-\eta/2})$.*

Proof. Let ψ of conductor $M' \mid M$ induce ξ. The twist $\psi\chi_K$ is primitive of conductor $q = M'd$, by coprimality of the conductors. As $d \equiv 7 \pmod 8$, its parity j is independent of d and quadratic reciprocity gives $\chi_K(h) = J(-d \mid h)$, including the factor at 2. The extra Euler

factors for $\xi\chi_K$ are nonzero at $\Re s > 0$, and classical non-vanishing gives $\Re s < 1$ for any indicated zero. Put $T = \sqrt{M'X}$. By the zero detector, if

$$I_j(\psi\chi_K; T) + I_j(\mu\psi\chi_K; T) \le T^{2\eta},$$

then $|\Gamma_j(s)|^2 \ll T^{-2\eta}$ at the putative zero. But Γ_j has a positive lower bound in modulus on the fixed compact rectangle $1/2 \le \Re s \le 1$, $|\Im s| \le Y$; hence for large enough X the putative zero forces the sum of tail integrals to be larger than $T^{2\eta}$. Corollary 2.6, for $c = \psi$ and $c = \mu\psi$, bounds the sum of the tail integrals over $d \le X$ by $O_\eta(T^{2+\eta})$. Dividing by $T^{2\eta}$ proves the announced count. □

The mean-square, tail-integral and density proofs above translate `sum_sq_gaussianTheta_jacobiSym_le`, `sum_thetaTailIntegral_le`, and `zero_density_quadTwist` in `ZeroFreeTwist.lean`.

Proposition 2.8 (Auxiliary zero-free twists)**.** *Fix a modulus M, a unit class $a_0 \pmod M$ all of whose primes are $7 \pmod 8$ and prime to M, a number $\beta > 1/2$ and $Y \ge 0$. There exists a prime $d \equiv a_0 \pmod M$ such that every $\widetilde{L}(s, \xi\chi_K)$, $\xi \bmod M$, is nonzero in $\Re s \ge \beta$, $|\Im s| \le Y$.*

Proof. Choose $0 < \eta < 1/18$ such that $1/2 + 9\eta < \beta$. By Proposition 2.7 the union, over the finitely many characters modulo M, of the bad sets has count $O_{M,Y,\eta}(X^{1-\eta/2})$. Its sum of reciprocals of primes converges by dyadic summation. The sum of reciprocals of all primes in the unit class a_0 diverges, by the elementary reciprocal-prime argument in Section 3.6; hence the union does not contain that class. Choose d outside the union. This is `exists_entireL_quadTwist_ne_zero` in the formal source. □

2.5 The one-sided zero-free criterion

Proposition 2.9 (Criterion used in the proof)**.** *Let $\beta > 1/2$, and let M, a_0 satisfy the conditions of Proposition 2.8. Suppose that for* each *prime $d \equiv a_0 \pmod M$ the corresponding prime error (2.3) satisfies*

$$R_M(x) \le x^{\beta-1} \qquad (x \textit{ sufficiently large}).$$

Then all $\widetilde{L}(s, \xi)$ for $\xi \bmod M$ are zero-free in the half-plane $\operatorname{Re} s > \beta$.

Proof. If $\widetilde{L}(\rho, \xi_0) = 0$ for some ρ to the right of β, apply Proposition 2.8 with $Y = |\operatorname{Im}\rho|$. Choose d so that no twisted $\widetilde{L}(s, \xi\chi_K)$ vanishes anywhere in the indicated half-strip. Then, at ρ, the numerator of (2.8) is a positive integer: the first sum includes the zero of $\widetilde{L}(\cdot, \xi_0)$, while the second sum is zero.

Furthermore at each real point $\sigma \in (\beta, 1)$ the numerator of the same residue formula is non-negative and $\sigma(1-\sigma) > 0$. Thus the real residues of Φ_M to the right of β are non-negative; those at and to the right of 1 are zero by classical non-vanishing and normalization. The assumed bound for the chosen d gives $E_M(v) \le e^{(\beta-1)v}$ eventually. The Landau principle (Lemma 2.2) says that Φ_M is holomorphic on $\operatorname{Re} s > \beta$, contradicting its non-zero residue at ρ. □

The logical order matters: the same arithmetic inequality has to be proved for each admissible auxiliary prime d in the given class, while the proof of the criterion chooses d *after* assuming that an offending zero exists. The threshold in the arithmetic estimate need not be uniform in d, and the choice of d need not work for all zeroes at once. This flexibility is what

avoids a zero-cancellation problem that an argument involving just one fixed quadratic field would face.

The formal declaration is `entireL_ne_zero_of_primeError_le` in `New/Endgame/OneSidedCriterion.lean`.

2.6 A good second scale in a long window

The compact edition employs a second important analytic argument. The arithmetic construction gives a bound involving two scales x, x_1 rather than one, with a combination of the two prime errors $R_{N_0}(x_1)$ and $R_1(x_1)$. We cannot simply discard those terms: their size relative to the desired saving at x would need justification. Instead, we have the freedom to vary $\log x_1$ in a sufficiently long interval. A high-moment argument produces a good choice of the second scale without requiring a pointwise prime number theorem error estimate there.

Lemma 2.10 (Window lemma)**.** *Let $E, E' : [0,\infty) \to \mathbb{R}$ be measurable with $|E(v)| + |E'(v)| \le Ce^v$. Suppose their Laplace transforms, with kernel $e^{(1-s)v}$, continue holomorphically to a neighborhood of the closed half-plane $\operatorname{Re} s \ge 1$. For every sufficiently large T and every $0 \le X \le e^T$ there exists $u \in [T, T^2]$ such that*

$$E(u) - XE'(u) \le e^{-T}.$$

The threshold for T may depend on the two functions but not on X in this range.

Proof. The closed disc of radius 3 about 4 lies in $\operatorname{Re} s \ge 1$. Choose a fixed $3 < R < 4$ such that both continued transforms are holomorphic on a neighborhood of the closed disc of radius R about 4. Such an R exists by compactness. Differentiation under the original integral near $s = 4$ gives

$$\Phi^{(j)}(4) = \int_0^\infty (-v)^j e^{-3v} E(v)\, dv.$$

The Cauchy estimates for Taylor coefficients yield, for all sufficiently large T, simultaneously for every integer $j \ge 0$,

$$\left| \int_0^\infty v^j e^{-3v} E(v)\, dv \right| \le \frac{j!T}{R^j}.$$

The same holds for E'. We can also require $|E(v)|, |E'(v)| \le Te^v$ for all $v \ge 0$.

Write $k = \lceil T^2/9 \rceil$ and suppose, for a contradiction, that $E(v) - XE'(v) > e^{-T}$ throughout $[T, T^2]$. Consider the function

$$v^k \left(e^{7T - Rv} + e^{2T - T^2 - v} + e^{-3v}(E(v) - XE'(v) - e^{-T}) \right).$$

On the window this is non-negative by hypothesis. Outside the window, the elementary growth bounds give

$$e^{-3v}|E(v) - XE'(v) - e^{-T}| \le e^{2T-2v} \qquad (v \ge 0),$$

for sufficiently large T, uniformly in $0 \le X \le e^T$. For $0 \le v \le T$, the first exponential in the parentheses dominates this upper bound, since $R < 4$. For $v \ge T^2$, the second exponential dominates it. Thus the integrand is non-negative everywhere.

Integrate over $v \ge 0$, use the moment bounds above, and recall $\int_0^\infty v^k e^{-qv} dv = k!/q^{k+1}$ for $q > 0$. The result, after division by $k!$, is

$$0 \le \frac{e^{7T}}{R^{k+1}} + e^{2T-T^2} + \frac{T(1+X)}{R^k} - \frac{e^{-T}}{3^{k+1}}.$$

This is impossible for large T. Divide all terms by the magnitude $e^{-T}/3^{k+1}$ of the negative term. The three positive ratios are respectively

$$e^{8T}(3/R)^{k+1}, \quad e^{3T-T^2}3^{k+1}, \quad 3e^{T}T(1+X)(3/R)^{k}.$$

The first and third tend to zero as $T \to \infty$, since k is quadratic in T and $3/R < 1$; the convergence is uniform for the third ratio by $X \le e^T$. The second ratio tends to zero because $(\log 3)/9 < 1$. Their sum is less than one for all sufficiently large T, giving the desired contradiction. □

This proof is an explicit example of what a genuine shortening of the formal development looks like when translated into ordinary mathematics. The readable edition uses a pointwise prime number theorem error bound at its much larger second cutoff. The compact edition needs only regularity of the two transforms in $\operatorname{Re} s \ge 1$ and a good point in a window. It is the ability to choose that point, not an assumed strong prime number theorem, that makes the argument work.

Apply the lemma with $E(v) = R_{N_0}(e^v)$ and $E'(v) = R_1(e^v)$ for the same auxiliary prime d. The hypotheses follow from Proposition 2.1. If $T = 10^6 \log x$ and X is the positive multiplier supplied by the arithmetic data, with $X \le e^T$, the lemma permits the choice $x_1 = e^u$ with

$$\log x_1 \in [T, T^2], \qquad R_{N_0}(x_1) - XR_1(x_1) \le x^{-10^6}. \tag{2.14}$$

This is a one-sided bound for a particular linear combination at a particular point; it should not be confused with a uniform pointwise estimate for either error. In the final argument the change of x_1 has to preserve the admissibility of the automorphic data. The scales are chosen in the arithmetic construction precisely so that this is possible.

Source. The moment and window lemmas are `IsPolar.eventually_moment_le` and `IsPolar.exists_window_sub_mul_le` in `New/Endgame/OneSidedCriterion.lean`; `New/Main.lean` applies them to the remainder in the final bound `eventually_primeError_le_rpow`.

2.7 From the target bound to GRH

To see precisely the quantifiers needed at the end, suppose that (2.6) has been established by the arithmetic construction. Let $q \ge 1$ be any modulus, $\chi \bmod q$ a Dirichlet character, and ρ a hypothetical zero with $\operatorname{Re}\rho > 1/2$, with the pole of a principal character excepted. Take $N_0 = 18480q$ and the unit class $a_0 = N_0/2 - 1 \pmod{N_0}$. Since $16 \mid N_0$, $a_0 \equiv 7 \pmod 8$ and $a_0^2 \equiv 1 \pmod{N_0}$. Choose $\varepsilon > 0$ with $\varepsilon < 1/100$ and $\beta = 1/2 + 8\varepsilon < \operatorname{Re}\rho$. The estimate (2.6) holds for every prime $d \equiv a_0 \pmod{N_0}$ and gives $R_{N_0}(x) \le x^{\beta-1}$ eventually. Proposition 2.9 implies that all pole-removed Dirichlet L-functions modulo N_0 are non-zero at ρ.

Changing the level of χ to N_0 multiplies its Dirichlet L-function by a finite product of omitted Euler factors, each non-zero in $\operatorname{Re} s > 0$. Thus a zero of $\widetilde{L}(\cdot, \chi)$ at ρ persists at the new level. This contradicts the preceding conclusion. We have proved, subject only to the arithmetic target estimate,

$$L(s, \chi) \ne 0 \qquad (\operatorname{Re} s > 1/2),$$

apart from the pole of the principal character at $s = 1$.

For a primitive character the functional equation, whose gamma factors are non-zero and finite in the open critical strip, sends a zero at s to a zero for the dual character at $1 - s$. Thus there can be no zero with $0 < \operatorname{Re} s < 1/2$ either. For an imprimitive character the omitted Euler factors again do not vanish in the open strip. Hence every zero there has real part $1/2$.

This includes the Riemann zeta function (modulus one); its standard functional equation and trivial-zero classification give the usual full statement of RH.

The rest of the article is concerned with the provenance of (2.6): how a problem about the lengths of automorphic forms and the divisibility of their theta lifts yields exactly the exponent needed for this analytic reduction.

Chapter 3

Classical analytic foundations without a prime number theorem

The two analytic uses of prime distribution in Chapter 2 and in the construction of the admissible record deserve particular attention. The first is the selection of the auxiliary quadratic twist; the second is the choice of the large split level primes. The former requires only divergence of a reciprocal-prime sum in a fixed residue class and an upper bound on the number of bad twists. The latter requires existence, not an asymptotic count, of a prime in a long window at a modulus that is a growing power of a fixed integer. Neither application requires the prime number theorem for arithmetic progressions. We prove these facts here, including the limited uniformity in the modulus needed for the prime windows.

This chapter supplements, rather than changes, the analytic inequalities proved earlier. In particular no zero-free region to the left of the line $\Re s = 1$ is used for a varying family of all possible moduli. The elementary disk in Lemma 3.9 is for the restricted family of powers of a fixed modulus. That restriction makes a substantial difference: its real characters are induced from one fixed finite set.

3.1 Finite characters, Gauss sums, and the Fourier bound

Write $e(t) = \exp(2\pi i t)$. For a positive integer q, a character modulo q is the extension by zero to the nonunits of a homomorphism $(\mathbb{Z}/q)^\times \to \mathbb{C}^\times$. The extension is periodic and completely multiplicative. The character of the trivial group for $q = 1$ takes the value 1 on every integer.

The existence and elementary properties of these characters can all be read from the structure theorem for finite abelian groups: for a cyclic group of order n the characters are given by the n-th roots of unity, and the characters of a finite direct product are tensor products. Thus there are $\varphi(q)$ characters and

$$\frac{1}{\varphi(q)} \sum_{\chi \bmod q} \overline{\chi(a)}\chi(n) = \mathbf{1}_{n \equiv a \bmod q} \quad ((a,q) = 1). \tag{3.1}$$

Indeed the sum vanishes on a nonidentity unit by multiplying it by a character which is nontrivial on that unit. On a nonunit both sides vanish. The same argument with the roles of the group and its character group exchanged proves that the sum of a nonprincipal character over a complete residue system is zero.

Lemma 3.1 (Conductor and change of level)**.** *For each character χ modulo q there is a*

unique primitive character ψ modulo some divisor $q' \mid q$ inducing χ. If $q = q'r$, then

$$\chi(n) = \psi(n)\mathbf{1}_{(n,r)=1}, \qquad L(s,\chi) = L(s,\psi)\prod_{\substack{p\mid q\\ p\nmid q'}}(1-\psi(p)p^{-s}). \tag{3.2}$$

If ψ_1, ψ_2 are primitive of coprime conductors q_1, q_2, their product is primitive of conductor q_1q_2.

Proof. Factor the unit group by the Chinese remainder theorem into groups modulo prime powers. For each $p^\alpha \| q$, choose the smallest $\beta \le \alpha$ such that the local component factors through reduction modulo p^β (allow $\beta = 0$ for the trivial factor). The product q' of the p^β is the conductor, and the factorization defines the unique induced primitive character. Its extension by zero detects the primes dividing q'; the indicator in (3.2) detects all remaining primes dividing q. The asserted Euler-factor identity follows in the domain of absolute convergence, and hence throughout continuation. In the coprime case the local components of the product are precisely the original local components, so their minimal exponents are unchanged. □

For a character ψ modulo q put

$$\tau_a(\psi) = \sum_{x \bmod q} \psi(x)e(ax/q), \qquad \tau(\psi) = \tau_1(\psi).$$

Lemma 3.2 (Primitive Gauss sums)**.** *If ψ is primitive, then*

$$\tau_a(\psi) = \overline{\psi(a)}\tau(\psi), \qquad |\tau(\psi)| = \sqrt{q}. \tag{3.3}$$

In particular, for a real primitive character of prime conductor d,

$$\overline{\tau(\psi)} = \psi(-1)\tau(\psi), \qquad \tau(\psi)^2 = \psi(-1)d.$$

Proof. For a a unit change variables by multiplication by a^{-1}. If a is not a unit, the additive character $x \mapsto e(ax/q)$ factors through $m = q/(a,q)$, a proper divisor of q. Since ψ is primitive, there is a unit u congruent to 1 modulo m with $\psi(u) \ne 1$. (The reduction of unit groups is surjective, as is seen prime-power by prime-power.) Changing variables by u shows that $\tau_a(\psi) = \psi(u)\tau_a(\psi)$, so this sum is zero. This proves the first assertion. Orthogonality of the finite additive characters gives

$$\sum_{a \bmod q} |\tau_a(\psi)|^2 = q \sum_{x \bmod q} |\psi(x)|^2 = q\varphi(q).$$

There are $\varphi(q)$ nonzero summands on the left, all of modulus $|\tau(\psi)|^2$. This proves the second assertion. For a real character, conjugation of the sum changes $a = 1$ to $a = -1$, giving the remaining identities. □

The following proof includes the imprimitive case, since the imprimitive character arising from the generalized Jacobi symbol in Chapter 2 need not have conductor equal to its written modulus.

Proposition 3.3 (Pólya–Vinogradov with the source constant)**.** *If χ is nonprincipal modulo q, then for every integer $X \ge 0$,*

$$\left|\sum_{1\le n\le X} \chi(n)\right| \le 2\sqrt{q}(1+\log q).$$

For primitive nonprincipal χ the factor 2 may be omitted.

Proof. First suppose χ primitive; then $q \geq 2$ and $\tau_0(\chi) = 0$. Fourier inversion and Lemma 3.2 give

$$\left|\sum_{n\leq X}\chi(n)\right| \leq q^{-1/2}\sum_{a=1}^{q-1}\left|\sum_{n=1}^{X} e(-an/q)\right|.$$

The inner geometric sum is bounded by $|\sin(\pi a/q)|^{-1}$. On $0 \leq v \leq 1/2$ the concavity of $\sin(\pi v)$ implies $\sin(\pi v) \geq 2v$, so that the inner bound is at most $q/(2\min(a, q-a))$. Sum first over $a \leq q/2$ and then over $a > q/2$. Since $\sum_{1\leq j\leq q/2} j^{-1} \leq 1 + \log q$, the primitive assertion follows. For general χ let $q = q'r$ as in Lemma 3.1; the inducing ψ is nonprincipal. Möbius inversion of the coprimality indicator gives

$$\sum_{n\leq X}\chi(n) = \sum_{d\mid r}\mu(d)\psi(d)\sum_{j\leq X/d}\psi(j).$$

The number of divisors of r is at most $2\sqrt{r}$ by pairing a divisor with its complementary divisor. Thus the already proved bound for ψ gives $2\sqrt{r}\sqrt{q'}(1+\log q')$, as required. □

This is the finite Fourier argument of `norm_gaussSum_of_isPrimitive` and `polya_vinogradov` in `Lib/Analytic/PolyaVinogradov/PolyaVinogradov.lean`. In particular no large-sieve theorem has been used.

3.2 The quadratic symbol and the choice of the family

Here are the elementary details behind the compatibility of the prime family with the Jacobi symbol used in the mean square. For an odd prime p, let (a/p) be 0 if $p \mid a$, 1 if a is a nonzero square modulo p, and -1 otherwise. For $p \nmid a$ the equality $(a/p) \equiv a^{(p-1)/2} \pmod p$ follows from cyclicity of the multiplicative group of the finite field.

For clarity we derive the particular reciprocity rule needed here. Put $h = (p-1)/2$. The signed least residues of $a, 2a, \ldots, ha$ have distinct absolute values: equality of absolute values would give $ai \equiv \pm aj \pmod p$, and hence $i = j$ in the indicated range. Their absolute values are therefore $1, \ldots, h$. Comparing products modulo p gives Gauss's lemma $(a/p) = (-1)^m$, where m counts those least nonnegative residues larger than $p/2$. If a is odd, comparison of sums modulo 2 gives

$$m \equiv \sum_{i=1}^{h}\lfloor ai/p\rfloor \pmod 2 :$$

indeed the sum of the signed residues is $a\sum i - p\sum\lfloor ai/p\rfloor - pm$, while modulo two it equals the sum of their absolute values. If q is another odd prime, count the integral points of the rectangle

$$1 \leq i \leq (p-1)/2, \qquad 1 \leq j \leq (q-1)/2$$

on the two sides of $qi = pj$. There is no point on the separating line. The two counts are respectively $\sum_i\lfloor qi/p\rfloor$ and $\sum_j\lfloor pj/q\rfloor$. Their sum is $(p-1)(q-1)/4$. Consequently

$$(q/p)(p/q) = (-1)^{(p-1)(q-1)/4}.$$

Likewise $(-1/p) = (-1)^{(p-1)/2}$. Applying Gauss's lemma to $a = 2$ gives $m = h - \lfloor p/4\rfloor$, whose parity by the four odd residue classes modulo 8 is $(p^2-1)/8$. This yields $(2/p) = (-1)^{(p^2-1)/8}$.

Let now $d \equiv 7 \pmod 8$ be prime, and $K = \mathbb{Q}(\sqrt{-d})$. The discriminant of K is $-d$. Its quadratic character is $\chi_K(n) = (n/d)$. For any odd prime $p \neq d$, reciprocity gives

$(-d/p) = (p/d)$; at $p = d$ both sides are zero. Moreover $(2/d) = 1$. Hence, with the generalized Jacobi symbol J of Chapter 2,

$$J(-d \mid n) = \chi_K(n) \qquad (n \geq 1). \tag{3.4}$$

Here the contribution to $J(-d \mid n)$ of an even factor in n is 1 because d is odd. The identity extends to all n by multiplicativity. The usual splitting characterization follows directly by reducing the minimal polynomial $X^2 - X + (d+1)/4$ of $(1 + \sqrt{-d})/2$ modulo p, with the 2-adic case giving two distinct roots because $(d+1)/4$ is even. Thus $\chi_K(p) = 1$ exactly at split primes and $\chi_K(p) = -1$ at inert primes, with $\chi_K(d) = 0$. Lemma 3.2 also shows that $\mathbb{Q}(\sqrt{-d})$ is contained in $\mathbb{Q}(\zeta_d)$ by the Gauss sum of its quadratic character. Neither use of quadratic reciprocity involves a choice of the sign of that square root.

For an arbitrary positive denominator h, the function $v \mapsto J(-v \mid h)$ is a constant sign times a (possibly principal) character modulo h. If at least one odd prime occurs in h to an odd exponent, this character is nonprincipal: choose a nonsquare unit at that prime and a square unit at all other primes, using the Chinese remainder theorem. Otherwise its value on every unit is 1 and it is principal, whether or not the exponent of 2 is odd. Hence

$$\left| \sum_{1 \leq v \leq X} J(-v \mid h) \right| \leq \begin{cases} X, & h \text{ or } 2h \text{ is a square,} \\ 2\sqrt{h}(1 + \log h), & \text{otherwise.} \end{cases} \tag{3.5}$$

This is exactly the dichotomy needed in the expansion of the quadratic-character mean square. Note that the modulus h here is the product of the two summation indices, not the discriminant parameter being averaged. This distinction explains why an ordinary Pólya–Vinogradov estimate suffices.

3.3 Holomorphy and nonvanishing on the line at one

We give direct proofs of the classical analytic properties of the Dirichlet L-functions that are used by the polar transform, including the case of a real character at $s = 1$. It is essential not to use the desired zero-free result in this step.

For a nonprincipal character χ modulo q, let $S_\chi(t) = \sum_{1 \leq n \leq t} \chi(n)$, with the real variable $t \geq 1$. Periodicity and the zero sum over a full period imply

$$|S_\chi(t)| \leq \min(q, t). \tag{3.6}$$

Partial summation gives

$$L(s, \chi) = s \int_1^\infty S_\chi(t) t^{-s-1}\, dt \qquad (\Re s > 0).$$

Initially this is equality with the Dirichlet series on $\Re s > 1$. The right hand side is holomorphic on $\Re s > 0$ by uniform convergence on compact subsets, so provides continuation there. In the principal case the corresponding formula for $\zeta(s)$ is

$$\zeta(s) = \frac{s}{s-1} - s \int_1^\infty \{t\} t^{-s-1}\, dt, \tag{3.7}$$

where $\{t\} = t - \lfloor t \rfloor$. This yields a simple pole of residue 1 at $s = 1$ and holomorphy elsewhere on $\Re s > 0$. The principal character modulo q has

$$L(s, \chi_0) = \zeta(s) \prod_{p \mid q} (1 - p^{-s}),$$

and its residue at 1 is strictly positive. Also

$$[(s-1)\zeta(s)]'_{s=1} = \gamma. \tag{3.8}$$

Indeed the constant term of (3.7) is $1 - \int_1^\infty \{t\}t^{-2}\,dt$. Summing on integer intervals gives $\int_1^n \{t\}t^{-2}\,dt = \log n - H_n + 1$; as $n \to \infty$ the assertion follows from the definition $\gamma = \lim_n (H_n - \log n)$.

The Euler products over p on $\Re s > 1$ are absolutely convergent and yield

$$\log L(s,\chi) = \sum_p \sum_{j\geq 1} \frac{\chi(p)^j}{jp^{js}}, \tag{3.9}$$

where the logarithm is specified by this analytic series. Absolute convergence follows by comparing to $\sum_{n\geq 2} n^{-\Re s}$. In particular all the L-functions are nonzero in this open half-plane, including the principal one. Differentiating gives the usual logarithmic derivative series, since the factor $\log p$ introduced by differentiation still leaves a convergent series on each smaller half-plane $\Re s \geq 1 + \delta$.

We recall the precise positivity principle for Dirichlet series that is needed for the boundary case.

Lemma 3.4 (Positive Dirichlet series)**.** *Suppose $a(n) \geq 0$ and $\sum_n a(n)n^{-s}$ has a finite abscissa σ_c of convergence. Its analytic function on $\Re s > \sigma_c$ cannot extend holomorphically to a neighborhood of the real point σ_c.*

Proof. Convergence at a real point implies absolute and locally uniform convergence on the right, by positivity. In the half-plane of convergence every derivative is obtained by introducing the corresponding power of $-\log n$: on a fixed smaller half-plane that power can be absorbed by n^δ for small fixed $\delta > 0$. If the function continued holomorphically near σ_c, choose a real center $c > \sigma_c$ sufficiently close to σ_c that its Taylor series converges at a real point $\sigma < \sigma_c$. Then

$$\sum_{j\geq 0} \frac{(-1)^j F^{(j)}(c)(c-\sigma)^j}{j!} = \sum_n a(n)n^{-c} \sum_{j\geq 0} \frac{((c-\sigma)\log n)^j}{j!}.$$

All terms are nonnegative. The right hand side equals $\sum_n a(n)n^{-\sigma}$ by monotone convergence and the exponential series, but the Taylor series on the left is finite, a contradiction. □

Proposition 3.5 (Classical line nonvanishing)**.** *For every character χ, the pole-removed function $\widetilde{L}(s,\chi)$ of Chapter 2 has no zeros in $\Re s \geq 1$.*

Proof. Nonvanishing for $\Re s > 1$ and the principal case at $s = 1$ have already been shown. First let χ be real and nonprincipal, and suppose $L(1,\chi) = 0$. Then $\zeta(s)L(s,\chi)$ is holomorphic on $\Re s > 0$, since the only possible pole, at 1, would be cancelled. In $\Re s > 1$ its Dirichlet series has coefficients $a(n) = \sum_{d|n} \chi(d)$. This arithmetic function is multiplicative and

$$a(p^j) = 1 + \chi(p) + \cdots + \chi(p)^j.$$

For $\chi(p) \in \{1, 0, -1\}$ its value is always nonnegative and is at least one when j is even. Thus $a(n) \geq 0$ for all n, and $a(m^2) \geq 1$ for all m. Its abscissa of convergence is at least $1/2$ because the subseries on squares diverges there; it is at most 1 by the original absolute convergence. Lemma 3.4 contradicts holomorphy on $\Re s > 0$.

For the remaining case at $s = 1$, consider the finite product $\prod_{\xi \bmod q} L(\sigma, \xi)$ for real $\sigma > 1$. Taking the sum of the Euler logarithms (3.9) and using finite-character orthogonality gives

$$\prod_{\xi \bmod q} L(\sigma, \xi) = \exp\left(\varphi(q) \sum_{p^j \equiv 1 \bmod q} \frac{1}{j p^{j\sigma}}\right) \geq 1.$$

The unique principal factor has a pole of order one at 1. A zero at 1 for a nonreal character would be accompanied by a zero for its distinct conjugate character (conjugate the Dirichlet series at real σ and continue). The product would then tend to zero as $\sigma \downarrow 1$, contrary to the inequality.

Finally for any character χ and any real $t \neq 0$, the Euler products give

$$\zeta(\sigma)^3 |L(\sigma + it, \chi)|^4 |L(\sigma + 2it, \chi^2)| \geq 1 \qquad (\sigma > 1). \tag{3.10}$$

To see this take the real logarithm of the left hand side using (3.9). At a prime not dividing q the contribution of each p^j is the positive weight $p^{-j\sigma}/j$ times $3+4\cos v+\cos(2v) = 2(1+\cos v)^2 \geq 0$, where v is the argument of $\chi(p)^j p^{-ijt}$. At a prime dividing q the contribution from the two L-functions is zero and that from ζ is positive. If $L(1 + it, \chi) = 0$, then its fourth power in modulus is $O((\sigma-1)^4)$. The ζ factor is $O((\sigma-1)^{-3})$ and the χ^2 factor is bounded, since $2t \neq 0$ even if χ^2 is principal. Their product would tend to zero, contradicting (3.10). Removing the principal pole does not affect any of these nonzeros away from 1. □

It follows in particular that the normalization $\widetilde{L}$ in the prime-error transform is nonzero at 1, and that the entire function in the source has the claimed locally nonvanishing restriction on the closed right half-plane. The proof did not use a prime number theorem or a zero-free region of the form $\Re s > 1 - c/\log(q(|\Im s| + 2))$. Equation (3.10) is the Euler-product positivity used by `norm_LFunction_product_ge_one` in the analytic library; its quantitative use is made below.

3.4 Reciprocal primes in a fixed residue class

Proposition 3.6 (The required elementary Dirichlet theorem)**.** *For every positive integer q and every unit class $a \bmod q$ the series*

$$\sum_{p \equiv a \bmod q} \frac{1}{p}$$

diverges. More precisely,

$$\sum_{p \equiv a \bmod q} p^{-\sigma} = \frac{1}{\varphi(q)} \log \frac{1}{\sigma - 1} + O_q(1) \quad (\sigma \downarrow 1,\ \sigma > 1).$$

Proof. For each $\chi \bmod q$ use the Euler logarithm (3.9) at the real argument $\sigma > 1$. Terms with $j \geq 2$ have total absolute value bounded independently of σ, since

$$\sum_p \sum_{j \geq 2} p^{-j}/j \leq \sum_{n \geq 2} \frac{1}{n(n-1)} = 1.$$

Thus by (3.1),

$$\sum_{p \equiv a \bmod q} p^{-\sigma} = \frac{1}{\varphi(q)} \Re \sum_{\chi \bmod q} \overline{\chi(a)} \log L(\sigma, \chi) + O(1).$$

The principal Euler logarithm is real and equals $\log(1/(\sigma-1))+O_q(1)$ by its positive residue at 1. By Proposition 3.5, a nonprincipal L-function is holomorphic and nonzero on some disk about 1 and hence admits a holomorphic logarithm there. To see that this logarithm stays compatible with the Euler logarithm, note that their difference on the real interval to the right of 1 is a continuous function in $2\pi i\mathbb{Z}$, and is therefore constant. Thus each nonprincipal Euler logarithm is bounded as $\sigma \downarrow 1$. This proves the estimate. If the reciprocal-prime series converged, each of the Dirichlet sums on the left would be bounded by that fixed sum, a contradiction. □

In particular, the choice of the auxiliary twist in Proposition 2.8 now follows without the previous appeal to the prime number theorem for a fixed arithmetic progression. The bad set up to X has size $O(X^{1-\eta/2})$. Its reciprocal sum over $2^h \le p < 2^{h+1}$ is $O(2^{-h\eta/2})$, so converges. Proposition 3.6 leaves at least one prime outside the bad set in the prescribed class, and in fact infinitely many. It is unnecessary to prove that a positive proportion of the primes in that class is good by a prime-counting asymptotic.

3.5 The theta transform and the full functional equations

We give details of the theta and Mellin ingredients, including the endpoints of the critical strip. For a Schwartz function f on $\mathbb{R}$, use the Fourier transform

$$\widehat{f}(\xi)=\int_{\mathbb{R}} f(t)e(-t\xi)\,dt.$$

Periodizing f with a positive period h gives a smooth periodic function whose n-th Fourier coefficient is $h^{-1}\widehat{f}(n/h)$, by termwise integration over one period. The series of these coefficients converges absolutely and uniformly (integrate by parts twice in the Fourier transform). Equality with the periodization follows from uniqueness of Fourier coefficients, or from convergence of the Cesàro means of its Fourier series. This is Poisson summation on an arithmetic progression. For $g_0(t)=e^{-\pi t^2}$, the Fourier transform is g_0 (differentiate under the integral and solve the resulting first-order equation, with value 1 at the origin from the Gaussian integral). Since multiplying by t transforms to $-(2\pi i)^{-1}\frac{d}{d\xi}$, the function $g_1(t)=te^{-\pi t^2}$ has transform $-ig_1$.

Take a primitive nonprincipal ψ modulo q, choose $j\in\{0,1\}$ with $\psi(-1)=(-1)^j$, and write $V_j(\psi;t)=\sum_{n\ge1}\psi(n)g_j(nt)$ as in the preceding chapter. Fourier periodization on each residue class and Lemma 3.2 give the exact (not just absolute-value) identity

$$V_j(\psi;t)=\frac{\epsilon_\psi}{\sqrt{q}\,t}V_j(\bar\psi;1/(qt)),\qquad \epsilon_\psi=(-i)^j\tau(\psi)/\sqrt{q},\quad |\epsilon_\psi|=1. \tag{3.11}$$

Indeed the bilateral sum is twice its positive part (the $n=0$ term is zero), and the finite Fourier coefficient for a is $\tau(\psi)\bar\psi(a)$. The factor $(-i)^j$ is determined by the Fourier convention above. The bilateral Poisson formula has exactly the factor $1/(qt)$ before the Gauss sum.

Define $\Gamma_j(s)=\frac12\pi^{-(s+j)/2}\Gamma((s+j)/2)$ and

$$M_\psi(s)=\int_0^\infty V_j(\psi;t)t^{s-1}\,dt.$$

At infinity, the Gaussian bound of Chapter 2 is exponential, and at zero (3.11) gives exponential decay in $1/t$ times a power of $1/t$. Thus this integral is entire: on any compact set of s all derivatives obtained by multiplying the integrand by powers of $\log t$ are absolutely integrable uniformly. Substituting each nt in the sum shows that $M_\psi(s)=\Gamma_j(s)L(s,\psi)$ when $\Re s>1$.

The gamma function is nonzero and holomorphic in $\Re z > 0$ and $1/\Gamma(z)$ is entire with simple zeros at the nonpositive integers. For completeness these statements follow from Euler's limit

$$\Gamma(z) = \lim_{N\to\infty} \frac{N^z N!}{z(z+1)\cdots(z+N)} \quad (\Re z > 0).$$

It is obtained by scaling the beta integral $\int_0^1 u^{z-1}(1-u)^N\,du$ by $u = v/N$ and applying dominated convergence; the product formula for the beta integral follows by repeated integration by parts. On taking reciprocals the limit becomes

$$\frac{1}{\Gamma(z)} = ze^{\gamma z}\prod_{n\ge 1}(1+z/n)e^{-z/n}.$$

The logarithms of the nonzero factors in the tail converge normally on every compact set where $n > |z|+1$, by the $O(|z|^2/n^2)$ bound for their logarithms. Hence the reciprocal is entire and has precisely the displayed zeros, each simple. This also establishes the gamma lower bound on fixed compact rectangles used by the zero detector.

It follows that $L(s,\psi)$ extends to an entire function, given by $M_\psi(s)/\Gamma_j(s)$. In (3.11) substitute $u = 1/(qt)$ inside the Mellin integral to obtain

$$M_\psi(s) = \epsilon_\psi q^{1/2-s} M_{\bar\psi}(1-s).$$

Therefore the completed functions

$$\Lambda(s,\psi) = (q/\pi)^{(s+j)/2}\Gamma((s+j)/2)L(s,\psi)$$

satisfy $\Lambda(s,\psi) = \epsilon_\psi \Lambda(1-s,\bar\psi)$. In the open critical strip the gamma factors have neither poles nor zeros; this is exactly the symmetry required to pass from the right-half-plane conclusion of the arithmetic estimate to the full strip for a primitive nonprincipal character. For an imprimitive character the finite factors (3.2) have no zeros in $\Re s > 0$, so the open-strip conclusion follows as well. This also proves entireness of the nonprincipal $\widetilde{L}$ of the previous chapter. No symmetry beyond this open strip is required for the imprimitive functions because their additional Euler factors may vanish on $\Re s = 0$.

The trivial primitive character (modulo 1) is handled separately. Let $V(t) = \sum_{n\ge1} e^{-\pi n^2 t^2}$. Ordinary Poisson summation gives $1 + 2V(t) = t^{-1}(1 + 2V(1/t))$. Split its Mellin integral at $t = 1$ and substitute $u = 1/t$ in the part from 0 to 1. In $\Re s > 1$ this yields

$$\Gamma_0(s)\zeta(s) = \int_1^\infty V(t)(t^{s-1} + t^{-s})\,dt + \frac{1}{2}\left(\frac{1}{s-1} - \frac{1}{s}\right). \tag{3.12}$$

The integral is entire by exponential decay. The right hand side is meromorphic everywhere, with poles only at 0 and 1, and invariant under $s \mapsto 1-s$. Division by $\Gamma_0(s)$ gives a meromorphic continuation of ζ, with the pole at 0 removable and $\zeta(0) = -1/2$. At $s = -2n$ for each integer $n \ge 1$, the right hand side equals its value at $1+2n$, which is positive and finite; Γ_0 has there a simple pole, so these are exactly the standard simple trivial zeros. At any other point with $\Re s < 0$ the right hand side is nonzero by symmetry and the Euler product, and Γ_0 is finite and nonzero. At $s = it \ne 0$ the same reasoning uses Proposition 3.5 at $1 - it$. Hence all remaining zeros of ζ lie in the open critical strip. The theorem for ζ ultimately deduced from the arithmetic inequality therefore agrees with the full trivial-zero exclusion in `src/Statements.lean`, not just with the assertion about zeros strictly inside $0 < \Re s < 1$.

The theta proof matches the statements `mellin_gaussianTheta_char`, `norm_gaussianTheta_functional_equation` and `gaussianMellin_eq` of `Lib/Analytic/DirichletTheta/MellinTheta.lean`. The elementary line-nonvanishing argument was separated from it to avoid a circular use of the functional equations for zero-freeness.

3.6 Analytic details of the bad-twist estimate

The mean-square argument already given in Chapter 2 can be made quite explicit using (3.5). Here we record the complete absolute-convergence and uniformity checks, which are important when the conductor of a quadratic twist itself varies.

For $j = 0, 1$ and any $\alpha > 0$ let

$$C_{j,\alpha} = \sup_{u>0} u^{\alpha} |g_j(u)| < \infty.$$

A direct derivative calculation (maximize $u^{\alpha+j}e^{-\pi u^2}$) gives $C_{j,\alpha} = ((\alpha + j)/(2\pi e))^{(\alpha+j)/2}$. One also has $|g_j(u)| \le 3e^{-2u}$: use $u^j \le e^u$ and $3u - \pi u^2 \le 9/(4\pi) < \log 3$. Hence, for any c with $|c(n)| \le 1$, the series $V_j(c; t)$ converges absolutely, locally uniformly for $t > 0$, and is bounded by $3e^{-t}/t$. Products and finite averages may consequently be expanded into absolutely convergent double sums. The bound $3e^{-t}/t$ also justifies the ordinary Mellin integrals on $\Re s > 1$ without appealing to cancellation.

For the mean square abbreviate $S_h(X) = |\sum_{v \le X} J(-v \mid h)|$. With $n, m \ge 1$ and $X \le T^2$ one needs to bound

$$A(t) = \sum_{n,m \ge 1} |g_j(nt) g_j(mt)| S_{nm}(X), \qquad t \ge 1/T.$$

If nm and $2nm$ are both nonsquares, (3.5) and the power estimate with $\alpha = 3/2 + \eta$ give

$$A_{\mathrm{nonsq}}(t) \le 2C_{j,3/2+\eta}^2 \, t^{-3-2\eta} \sum_{n,m \ge 1} \frac{1 + \log(nm)}{(nm)^{1+\eta}}.$$

For $\eta > 0$ the last series converges. The identity

$$\sum_{n,m \ge 1} (nm)^{-1-\eta}(1 + \log(nm)) = \zeta(1+\eta)^2 - 2\zeta(1+\eta)\zeta'(1+\eta)$$

follows by termwise differentiation at a point of absolute convergence. Thus this part has a constant depending only on j, η, not on the modulus of any original character. Since $t^{-2} \le T^2$, it is at most a constant times $T^2 t^{-1-2\eta}$.

If knm is a square with $k \in \{1, 2\}$, write $kn = au^2$ and $m = av^2$ with a squarefree; this is possible by uniqueness of the odd-exponent kernel in prime factorization. For $\alpha = 1/2 + \eta$ the product of the power estimates for these two variables is bounded by

$$C_{j,\alpha}^2 t^{-2\alpha} k^{\alpha} a^{-2\alpha} (uv)^{-2\alpha}.$$

Use $S_{nm}(X) \le X \le T^2$, sum over a, u, v without their divisibility restrictions, and also sum over the two k. The resulting upper bound is

$$C_{j,\alpha}^2 (1 + 2^{\alpha}) \zeta(1 + 2\eta)^3 T^2 t^{-1-2\eta}.$$

This proves the asserted uniform mean square and supplies explicit choices of its constants. Cauchy–Schwarz with the finite sum over v, followed by integration against $dt/\sqrt{t}$, produces the tail bound $O_{j,\eta}(T^{2+\eta})$, since $\int_{1/T}^{\infty} t^{-1-\eta}\, dt = T^{\eta}/\eta$. Both sums for $c(n) = \psi(n)$ and $c(n) = \mu(n)\psi(n)$ are covered.

We also recall why the interpolation estimate in the zero detector is uniform in the imaginary coordinate. Suppose a bounded holomorphic function f on the interior of a strip $a < \Re z < b$ is continuous and bounded on its closure, with positive edge bounds M_a, M_b. Divide by

$$\exp\left(\frac{b-z}{b-a}\log M_a + \frac{z-a}{b-a}\log M_b\right).$$

The quotient is bounded on the strip and has boundary modulus at most one. Multiply by $\exp(\delta(z-a)^2)$ for $\delta > 0$. On horizontal segments at imaginary parts $\pm R$ its modulus tends uniformly to zero as $R \to \infty$. On vertical edges its modulus is at most $\exp(\delta(b-a)^2)$. The maximum modulus principle on rectangles, followed by $R \to \infty$ and then $\delta \downarrow 0$, gives the three-lines theorem in this situation.

Apply this to the function F of Proposition 2.4. Boundedness of each Mellin integral on its strip follows by taking absolute values: if $a \le \Re s \le b$ then $t^{\Re s - 1}$ is bounded by $t^{a-1} + t^{b-1}$, and the bounds of the previous paragraph give integrability away from zero. For the full M_ψ near zero use the transformation (3.11), whose exponential decay gives integrability for every real exponent. The truncated integral has a fixed positive lower limit $1/T$ and hence no such issue. Γ_j on the strip is bounded from above by its integral representation. Thus the boundary interpolation gives, for constants independent of the twisting prime,

$$|F(s)| \le C_j T^{4\eta - (2/3)(\Re s - 1/2)} \qquad (1/2 \le \Re s \le 2).$$

At a zero with $\Re s \ge 1/2 + 9\eta$ the left side is $|\Gamma_j(s)|^2$ and the right side is at most $C_j T^{-2\eta}$. By the Euler product for the reciprocal gamma function, its modulus on the closed fixed rectangle $1/2 \le \Re s \le 1$, $|\Im s| \le Y$ has a strictly positive lower bound, independently of the prime twist. This is why we can count bad twists using only the first moment of the two tail integrals and obtain the exponent $1 - \eta/2$. The occurrence of the conductor in the zero detector is only through $q \le T^2$, used to fold the lower half of the Mellin integral. If $\xi \bmod M$ is imprimitive, the inducing ψ has fixed conductor $M' \mid M$, and the twists $\psi\chi_K$ are primitive of conductor $M'd$ by Lemma 3.1. Their extra Euler factors at modulus Md cannot introduce zeros in $\Re s > 0$, by (3.2) (each has modulus bounded away from zero at every point with $\Re s > 0$). Thus the bad-twist bound genuinely holds for all the characters in the mean defining R_M, not just the primitive ones.

This completes the classical dependencies of the density and auxiliary-twist propositions. The relevant source anchors are `sum_sq_gaussianTheta_jacobiSym_le`, `sum_thetaTailIntegral_le`, `zero_density_quadTwist`, `three_lines_bound_of_LFunction_eq_zero` and `exists_entireL_quadTwist_ne_zero` in `Lib/Analytic/QuadraticTwist/ZeroFreeTwist.lean`.

3.7 A uniform disk at one for powers of a fixed modulus

We now prove the precise restricted uniformity that replaces the invocation of Siegel–Walfisz in the construction of the arithmetic data. Fix an integer M divisible by 4 and a real number $A > 0$. In this subsection only, $q = M^k$ with $k \ge 2$.

Lemma 3.7 (Real characters have bounded conductor in the tower)**.** *Every real character modulo M^k is induced by one modulo M^2. The corresponding L-functions, including the pole-removed principal function, belong to a single fixed finite collection as k varies.*

Proof. A real character on the unit group is trivial on every square. We show every unit u reducing to 1 modulo M^2 is a square modulo M^k. By the Chinese remainder theorem it suffices to check modulo $p^{kv_p(M)}$. If p is odd, every $u \equiv 1 \pmod p$ has a square root modulo each p^n: starting with 1 modulo p, if $x^2 \equiv u \pmod{p^n}$ and $p \nmid x$, the square of $x + p^n t$ differs from x^2 modulo p^{n+1} by $2xp^n t$. The coefficient $2x$ is invertible modulo p, so a suitable t completes the induction.

At $p = 2$ the assumption $4 \mid M$ gives $u \equiv 1 \pmod{16}$, in particular $u \equiv 1 \pmod 8$. Every such unit is a square modulo any higher power of two. Starting with an odd solution modulo

8, for $n \geq 3$ the two candidates x and $x+2^{n-1}$ have squares differing by $2^n x$ modulo 2^{n+1} (the additional term 2^{2n-2} is divisible by 2^{n+1}). Exactly one corrects the next binary digit of the square as needed. This proves the assertion by induction. Thus the kernel of the unit-group reduction to M^2 is contained in the squares, and every real character factors through M^2. Since M^2 and M^k have the same prime divisors, the values on all integers agree with those of the lower-level character. Their Dirichlet L-functions therefore coincide, proving the final assertion. □

We next give a simple uniform bound in a fixed small disk, using the bounded sums (3.6) rather than a theta estimate.

Lemma 3.8 (Crude bounds in a disk)**.** *For every $0 < r < 1/4$ there is a constant $C_r > 0$ such that for every nonprincipal character χ modulo any $q \geq 2$,*

$$|L(z,\chi)| \leq C_r q^{2r} \quad (|z-1| \leq 2r), \qquad |L'(z,\chi)| \leq C_r q^{2r} \quad (|z-1| \leq r).$$

Proof. On the first disk, $\sigma = \Re z \geq 1 - 2r > 0$ and $|z| \leq 1 + 2r$. From partial summation and (3.6), split the integral at q to obtain

$$\frac{|L(z,\chi)|}{|z|} \leq \int_1^q t^{-\sigma}\, dt + q \int_q^\infty t^{-\sigma-1}\, dt \leq \frac{q^{2r}}{2r} + \frac{q^{2r}}{1-2r}.$$

This gives the first estimate. For $|z-1| \leq r$ apply Cauchy's estimate on the circle of radius r about z contained in the first disk. Absorb $1/r$ into the constant. □

Proposition 3.9 (Small uniform zero-free disk and logarithmic derivative)**.** *Fix M, A as above. For every sufficiently small real $\delta > 0$, uniformly in $k \geq 2$ with $q = M^k \leq \delta^{-A}$, every pole-removed function $\widetilde{L}(s,\chi)$ modulo q is nonzero on $|s-1| \leq \delta$ and satisfies*

$$\delta \left| \frac{\widetilde{L}'(s,\chi)}{\widetilde{L}(s,\chi)} \right| < 1 \qquad (|s-1| \leq \delta).$$

In fact the supremum of this quantity over all these data tends to zero as $\delta \downarrow 0$.

Proof. For real characters, including the principal character, Proposition 3.5 and Lemma 3.7 show that the functions form a fixed finite nonvanishing collection at 1. In each small disk their logarithmic derivatives are bounded; take a common disk and common bound. Multiplication by δ then gives the assertion.

For a nonreal character χ, both χ and χ^2 are nonprincipal. Let $r > 0$ be small enough that $r < 1/4$ and $10Ar < 1$. Assume $\delta < r$. From (3.10) with $t = 0$ and Lemma 3.8,

$$1 \leq \zeta(1+\delta)^3 |L(1+\delta,\chi)|^4 |L(1+\delta,\chi^2)| \leq (2/\delta)^3 |L(1+\delta,\chi)|^4 C_r q^{2r}.$$

Here $\zeta(1+\delta) \leq 1 + 1/\delta \leq 2/\delta$ by integral comparison. Thus

$$|L(1+\delta,\chi)| \geq c_r \delta^{3/4} q^{-r/2}, \qquad c_r = (8C_r)^{-1/4}. \tag{3.13}$$

Every point s of the disk $|s-1| \leq \delta$ may be joined to $1+\delta$ by a segment of length at most 2δ, still in $|s-1| \leq r$. Consequently

$$|L(s,\chi) - L(1+\delta,\chi)| \leq 2C_r \delta q^{2r}.$$

The ratio of this bound to the lower bound in (3.13) is at most $2C_r c_r^{-1} \delta^{1/4} q^{5r/2} \leq 2C_r c_r^{-1} \delta^{1/4 - 5Ar/2}$. By our strict choice of r the exponent is positive (if $A = 0$ the case of bounded q is simpler). For sufficiently small δ this ratio is at most $1/2$ uniformly, so the function is nonzero everywhere on the disk and its modulus at least $\frac{1}{2} c_r \delta^{3/4} q^{-r/2}$. Divide the Cauchy derivative bound by this lower bound and multiply by δ; the same positive power of δ controls the result. Since $\widetilde{L} = L$ for these nonprincipal characters, the proof is complete. □

The conductor restriction cannot simply be removed in this proof: real characters of arbitrarily increasing conductor need not belong to a fixed family. The preceding disk argument is the conventional proof of the special estimate `eventually_mul_norm_deriv_entireL_lt` in `Lib/Analytic/Primes/AuxiliaryPrimes.lean`. It uses no ineffective lower bound for an increasing real quadratic conductor. It is sufficient because the modulus used for the level-prime searches is itself a power of $N_0 d$, which is fixed once the auxiliary quadratic field is chosen.

3.8 Prime existence in long logarithmic windows

We now prove the source's restricted prime-window theorem. The proof is a second, distinct use of a one-sided positivity argument: if the indicated interval contains no primes in the class 1, analytic continuation at 1 conflicts with the obvious positive residue there. We give the complete Cauchy calculation, with polynomial slack on the boundary instead of the smallest possible absolute constants.

Theorem 3.10 (A prime in the required logarithmic window)**.** *Let M be fixed and divisible by 4, and let $A \geq 1$ be fixed. For every sufficiently large real W, every $k \geq 2$ such that $q = M^k \leq W^A$, and every*

$$W \leq U \leq W^A,$$

there is a prime p with

$$p \equiv 1 \pmod{q}, \qquad U \leq \log p \leq U + W.$$

The threshold for W is independent of k, U in these ranges.

Proof. Put $\delta = W^{-1/2}$ and $a(n) = \Lambda(n)\mathbf{1}_{n \equiv 1 \bmod q}$. By Proposition 3.9, applied with exponent $2A$, the function

$$F(z) = \frac{1 - z\sum_{\chi \bmod q} \widetilde{L}'(1+z,\chi)/\widetilde{L}(1+z,\chi)}{\varphi(q)} \tag{3.14}$$

is holomorphic on a neighborhood of the closed disk $|z| \leq \delta$. Its value at zero is $1/\varphi(q)$, and $|F(z)| \leq 1/\varphi(q) + 1 \leq 2$ on the disk: first on the boundary from Proposition 3.9, and then in the interior by the maximum modulus principle. For $\Re z > 0$, character orthogonality and the Euler logarithmic derivative give the identity

$$\sum_{n \geq 1} \frac{a(n)}{n^{1+z}} = \frac{F(z)}{z}. \tag{3.15}$$

The principal pole contributes exactly the 1 in the numerator of (3.14). The series is absolutely convergent in its indicated domain. Let $D = 1/U$. For large W, $0 < D \leq \delta$. In particular,

$$\sum_n a(n) n^{-1-D} = \frac{F(D)}{D} \leq 2U. \tag{3.16}$$

This is the only bound for the full positive Dirichlet series that we will require.

Suppose, to the contrary, that the interval $[U, U+W]$ contains no logarithm of a prime congruent to 1 modulo q. Define the truncated initial sum and the holomorphic function

$$S_U(z) = \sum_{\log n < U} a(n) n^{-1-z}, \quad G(z) = (F(z) - zS_U(z)) \exp\left(Uz + \frac{\sqrt{W}}{4}(1 + \sqrt{W}z)^2\right).$$

The initial sum is finite. On the circle $|z| = \delta$, write $v = \Re z$ and set

$$t = \frac{W}{2}(1 + \sqrt{W}\, v).$$

Since $-\delta \le v \le \delta$, we have $0 \le t \le W$. A direct calculation using $W|z|^2 = 1$ gives

$$\left|\exp\left(Uz + \frac{\sqrt{W}}{4}(1 + \sqrt{W}z)^2\right)\right| = \exp((U + t)v). \tag{3.17}$$

At the center, in contrast, the exponential factor is $\exp(\sqrt{W}/4)$ and hence

$$|G(0)| = \frac{\exp(\sqrt{W}/4)}{\varphi(q)} \ge W^{-A}\exp(\sqrt{W}/4). \tag{3.18}$$

We give a bound polynomial in W for G on the boundary, separately according to the sign of $v - D$.

If $v \le D$, then $(U + t)v \le (U + W)D \le 2$. The F contribution to $|G|$ is consequently at most $2e^2$. For S_U, positivity and (3.16) give

$$|S_U(z)| \le e^{U(D-v)} \sum_n a(n)n^{-1-D} \le 2Ue^{U(D-v)}.$$

Multiplying by $\delta\exp((U + t)v)$ gives at most $2\delta Ue^{1+tv} \le 2e^2\delta U$. This proves the boundary estimate when $v \le D$, including the points with $v \le 0$ where the untruncated series might not be defined.

For $v > D > 0$, subtract the finite sum from (3.15). This is now legitimate, because both series converge absolutely. The resulting tail

$$F(z) - zS_U(z) = z \sum_{\log n \ge U} a(n)n^{-1-z}$$

has two types of terms. For the portion $\log n > U + W$, the absolute-value sum before multiplication by z is at most $2U\exp(-(U + W)(v - D))$ by (3.16). After multiplication by the factor δ and the exponential in (3.17), its contribution has modulus at most

$$2\delta U\exp((U + W)D - (W - t)v) \le 2e^2\delta U.$$

The remaining indices have $U \le \log n \le U + W$. By the supposition and the support of Λ, their nonzero terms occur only for nontrivial prime powers. The absolute series

$$C = \sum_p \sum_{j\ge 2} (\log p)\, p^{-3j/4}$$

is finite: for each p, the inner geometric sum is at most a fixed constant times $p^{-3/2}$, and $\sum_{n\ge 2}(\log n)n^{-3/2}$ converges. Therefore for $v > D$ the absolute-value sum over this remaining portion is at most $C\exp(-U(1/4 + v))$; indeed $n^{-1-v} = n^{-3/4}\exp(-(1/4 + v)\log n)$. Multiply by $\delta\exp((U + t)v)$ to bound its contribution by

$$C\delta\exp(-U/4 + tv) \le C\delta\exp(-W/4 + \sqrt{W}).$$

This tends to zero uniformly in the asserted range.

Combining the two cases gives, for all boundary points,

$$|G(z)| \le 2e^2 + 2e^2\delta U + C\delta\exp(-W/4 + \sqrt{W}).$$

This is $O_A(1 + W^A)$ uniformly in k, U, and even $O_A(W^{A-1/2} + 1)$ since $\delta = W^{-1/2}$. The function G is holomorphic on a neighborhood of the closed disk and so the maximum modulus principle bounds $|G(0)|$ by the same quantity. But (3.18) eventually exceeds every one of these polynomial bounds. This contradiction proves the theorem. □

This is the specialized theorem `exists_prime_modEq_one_log_mem` of `Lib/Analytic/Primes/AuxiliaryPrimes.lean`. Notice the two key features of the argument. First, the identity (3.15) controls the sum only on the right of the origin, but on the left it is enough to compare a *finite* truncated sum to its positive bound at D. Second, the smoothing exponential has exactly the boundary norm $\exp((U+t)v)$ with $0 \le t \le W$, allowing opposite tail comparisons at U and $U+W$. Its value at the center is superpolynomial. The absence of primes in the interval is incompatible with the pole residue. The prime powers within the interval do not obstruct this conclusion, because their sum is absolutely convergent already at $3/4$.

Corollary 3.11 (The primes for the admissible data, without Siegel–Walfisz)**.** *For fixed N_0, d, ε and the scales of Proposition 4.2, all the requirements (4.5), the selection of g_0, g_1 and the subsequent selection of N follow from Theorem 3.10.*

Proof. Write $Y = x^{\varepsilon}$ as there and take $W = \sqrt{Y}$. The modulus to which the prime classes are referred is

$$q = M_\omega^2 = (N_0 d)^{2(s+6)}.$$

Its base $M = N_0 d$ is a multiple of 4, fixed while x tends to infinity. Its exponent $k = 2(s+6) \ge 2$. Since $s \le 32 \log x$ and $2^s = m^{22} \le (2\sqrt{x})^{22}$, we have $q \le W^A$ and $2^s \le W^A$ for one fixed sufficiently large $A = A(N_0, d, \varepsilon)$. For every $U \in [100Y, 2^s]$ we have $W \le U \le W^A$. The theorem therefore gives a prime congruent to 1 modulo M_ω^2 with logarithm in $[U, U+W]$, exactly as in (4.5).

To select g_0, g_1, use $h_K + 1$ disjoint intervals of width W inside $[10Y, 11Y]$, for instance those starting at $10Y + 2iW$ for $1 \le i \le h_K + 1$. Here h_K depends on the fixed field only, so the intervals are disjoint and lie in the desired range for all sufficiently large x. Their starting points again lie between W and W^A after increasing A if necessary. Thus there are distinct split primes p_i in these intervals. For each choose a prime ideal $\mathfrak{p}_i$ of K of norm p_i. Pigeonhole in the finite ideal class group gives distinct i, j with $[\mathfrak{p}_i] = [\mathfrak{p}_j]$. Consequently $\mathfrak{p}_i\overline{\mathfrak{p}_j}$ is principal: in the class group $[\overline{\mathfrak{p}_j}] = [\mathfrak{p}_j]^{-1}$ since their product is (p_j). Its integral generator π_g has absolute norm $p_i p_j$, hence $\pi_g \overline{\pi}_g = g_0 g_1$ with $g_0 = p_i, g_1 = p_j$ in the notation of the source.

Finally the level exponent λ_N lies in $[100Y, 2^s]$ for sufficiently large x, by the estimate following (4.8). Therefore (4.5) gives N in the required interval regardless of the particular second cutoff eventually chosen in its allowed window. The prime search applies uniformly to the entire permitted range for λ_N, not just to a single choice made before the adjustment of x_1. □

In particular, when reading the proof of Proposition 4.2, Corollary 3.11 can be used in place of its invocation of a classical uniform prime number theorem. There is no circularity: the prime-window theorem concerned only the restricted powers of fixed $N_0 d$ and relied on nonvanishing at the line $\Re s = 1$, proved in Proposition 3.5. The level allocation and class-group pigeonhole are still as in the arithmetic argument, and no assumption about the desired critical line is involved.

3.9 Final conditional theorem and quantifier audit

The conventional analytic deductions are now self-contained relative to elementary real and complex analysis, finite abelian groups, elementary algebraic number theory (including finiteness of the ideal class group in the construction of the record) and the stated arithmetic inputs in the later chapters. To make the remaining mathematical burden precise, we state a theorem recording their exact logical role. The notation is as in Chapters 4 and 55.

Theorem 3.12 (Sufficient arithmetic inequalities)**.** *Suppose that for every fixed N_0 divisible by 18480, every prime $d \equiv N_0/2-1 \bmod N_0$, and every $0<\varepsilon<1/100$, the following holds for all sufficiently large x. For the admissible records of Proposition 4.2, including each permitted replacement of the second cutoff, the arithmetic estimates A, B, C of Chapter 55 hold with the same objects and normalizations. Then every Dirichlet L-function has no zero in $\Re s > 1/2$ (the principal pole convention at $s=1$ excepted), all its zeros in the open critical strip lie on $\Re s = 1/2$, and every zero of $\zeta(s)$ other than its standard trivial zeros lies on that line.*

Proof. Fix the three parameters in the hypothesis. Set $Y = x^{\varepsilon}$, m to be the least power of two at least $\sqrt{x}$, and $T = 10^6 \log x$. By the admissibility construction (with Corollary 3.11) there is such a record for all sufficiently large x. Use the ratio $X = |\mathcal{T}|/|\mathcal{S}| \le Y \le e^T$. Holomorphy of the polar transforms at and to the right of 1 follows from Proposition 3.5 and the exact computation of Proposition 2.1; note the scalar constant γ is specifically the coefficient computed in (3.8), so the logarithmic pole subtraction is exactly the expected one. Lemma 2.10 gives a permitted $u \in [T, T^2]$ with

$$R_{N_0}(e^u) - X R_1(e^u) \le e^{-T}.$$

Use the record with $x_1 = e^u$ and all derived levels, weights, spaces, lifts, and forms constructed from that record. Thus the hypotheses A–C compare identical data; no uniform pointwise bound in u has been substituted for the window argument.

The dimension and index estimates give q independent short forms among the translates, where $(1-4/b^2)n \le q \le n$, $n = \dim W > 0$. Their logarithmic squared covolume is bounded above by $q(H - m\mathcal{R} + L)$. The index cost is at most $5n\log(1+b^2)B$. In the lower estimate C it appears twice, while the leading height is subtracted three times by the allowance L. Because $H - m\mathcal{R} + L$ might have either sign, the relation $q \le n$ alone does not show that its multiple by q is at most its multiple by n. Use instead the crucial crude bound $-(H - m\mathcal{R} + L) \le b^2 x^{99}$ and $n - q \le 4n/b^2$. It gives

$$n(H - 3L) \le n(H - m\mathcal{R} + L) + 4nx^{99} + 10n\log(1+b^2)B.$$

After division by n the common H cancels. Since $B = faY^3$ and $L = BY$, divide the remaining error $4L + 4x^{99} + 10\log(1+b^2)B$ by BY^2. The quotient tends to zero: the linear terms in Y are $O(Y^{-1})$ after division, and $x^{99}/(BY^2) \to 0$ by $a \ge xb$ and $b = e^Y$. Hence for sufficiently large x, $m\mathcal{R} \le BY^2$.

Write the exact remainder identity in Chapter 55 in the form

$$R_{N_0}(x) = \frac{\mathcal{R}}{a|\mathcal{S}|} + R_{N_0}(x_1) - X R_1(x_1).$$

The two-cutoff term is bounded by x^{-10^6} by our choice of x_1. For the first term we have

$$\frac{\mathcal{R}}{a|\mathcal{S}|} \le \frac{fY^5}{m|\mathcal{S}|} \le \varphi(N_0) Y^6/m \le \varphi(N_0) x^{-1/2+6\varepsilon}.$$

The fixed factor $\varphi(N_0)$ and the much smaller window error are absorbed by $x^{2\varepsilon}$ for all sufficiently large x. Thus

$$R_{N_0}(x) \le x^{-1/2+8\varepsilon} \tag{3.19}$$

eventually. Every threshold in this deduction may depend on the fixed parameters N_0, d, ε.

To pass to an arbitrary character χ modulo q_*, suppose that $\widetilde{L}(\rho,\chi) = 0$ with $\Re\rho > 1/2$ and take $N_0 = 18480 q_*$. The induced character modulo N_0 has a zero at the same point, by (3.2), since the additional Euler factors do not vanish in $\Re s > 0$. The class $N_0/2 - 1$ is a

unit modulo N_0, as its square is congruent to one modulo N_0 (use $4 \mid N_0$), and is 7 modulo 8 because $16 \mid N_0$. Choose $\varepsilon > 0$ with $\varepsilon < 1/100$ and $\beta = 1/2 + 8\varepsilon < \Re\rho$. By Proposition 2.7, whose classical dependencies have been proved here, the bad quadratic twists modulo N_0 with zeros in $\Re s \geq \beta$, $|\Im s| \leq |\Im\rho|$ have a convergent reciprocal sum. Proposition 3.6 supplies a prime d in the indicated class outside this set for all the characters modulo N_0 simultaneously. The hypothesis of this theorem and (3.19) apply to just that fixed d. Thus the residue at ρ of the meromorphic prime-error transform is nonzero, since there is a zero of one of the original characters and no zero of any of the twisted functions there. At every real point to the right of β its residue is nonnegative: in the interval $(\beta, 1)$ this is the sign of the positive integer multiplicity sum divided by the positive number $\varphi(N_0)\sigma(1-\sigma)$; beyond it the function is holomorphic by Proposition 3.5. Lemma 2.2, applied to (3.19), contradicts that nonzero residue at ρ. This proves the right-half-plane conclusion.

The symmetry of the completed primitive L-functions established above, followed by (3.2) for the imprimitive functions, proves the open-strip statement. Formula (3.12), nonvanishing on the line at one, and the explicit trivial-zero classification following that formula yield the full conclusion for ζ. □

We emphasize what the theorem does *not* prove by itself. The assumptions A, B, C contain the highly specialized arithmetic geometry and representation-theoretic estimates of the formal development. The conditional theorem removes none of their quantitative or compatibility requirements. It only replaces classical analytic black boxes and verifies that their exact conclusions, when proved, imply the stated result with no further analytic assumptions about zeros. In particular neither a uniform choice of x over the twisting primes, nor a simultaneous choice of quadratic twist avoiding zeros at arbitrarily large heights, is required. Such stronger claims are not used. The choice of a prime twist occurs after the hypothetical offending zero has been fixed; that choice is followed by the eventual limit in x for the chosen field.

The dependency order in the source is `exists_data`, the arithmetic conclusions from `New/Endgame/ShortAndEnough.lean` and `LongForms.lean`, `star_inequality`, `eventually_primeError_le_rpow`, and the final zero-free consequences in `New/Main.lean`. The analytic arguments of the present chapter and Chapter 2 supply their relevant analytic interfaces. Their proofs are independent of the remaining specialized input theorems in the arithmetic parts.

Chapter 4

The admissible data and coefficient spaces

Chapter 3 gives the analytic prime-window result used in this construction. Chapter 6 constructs the global Weil data, including the splitting character, the finite lattice and the signed Hermitian spaces, and establishes the integral unitary counts and mass-ratio estimate.

4.1 The CM extension and the different

We first record the precise arithmetic of the extension in the compact development. Fix $N_0 > 0$ with $18480 \mid N_0$, and a prime $d \equiv N_0/2 - 1 \pmod{N_0}$. Equivalently $N_0 \mid 2(d+1)$ and $N_0 \nmid d+1$. In particular $d \equiv 7 \pmod 8$, $d > 64$, and $(d, N_0) = 1$. Put $K = \mathbb{Q}(\sqrt{-d})$, of discriminant $-d$. For an integer $s \geq 1$ define

$$c_s = \lfloor \log_3 s \rfloor + 5, \quad T_s = 3^{c_s} 2^s, \quad N_1 = N_0 T_s, \quad E = K(\zeta_{N_1}), \quad M_\omega = (N_0 d)^{s+6}.$$

Here T_s is a number, not a set of places, and $T_s = |\mathcal{T}|$ below. For each prime $p \mid N_1$ the character $\chi_K(p) = 1$. Indeed for an odd such prime p, $p \mid (d+1)/4$ since all the primes of N_1 divide N_0; hence $-d \equiv 1 \pmod p$. For $p = 2$ the claim follows from $d \equiv 7 \pmod 8$. Consequently every nonsplit prime of K is at least 13, and d is prime to N_1. Since K and $\mathbb{Q}(\zeta_{N_1})$ are linearly disjoint (their discriminants have disjoint prime supports),

$$E \subset \mathbb{Q}(\zeta_{N_1 d}), \qquad [E : K] = \varphi(N_1) = T_s \varphi(N_0), \qquad \mathrm{Gal}(E/K) \simeq (\mathbb{Z}/N_1)^\times.$$

In the inclusion we use the quadratic Gauss sum of conductor d. The field E is totally imaginary and complex conjugation is central in its Galois group over $\mathbb{Q}$, so E is a CM field with real subfield E^+. The extension E/E^+ is unramified at the finite places: at every residue characteristic at least one of the primitive third and fourth roots of unity in E has order prime to that characteristic, and complex conjugation acts nontrivially on its reduction. Thus complex conjugation cannot belong to an inertia group of $E/\mathbb{Q}$. The roots of unity in E are exactly the N_1-th roots: the cyclotomic field of even order $N_1 d$ has no other roots of unity, and if E contained a primitive d-th root it would have degree over $\mathbb{Q}(\zeta_{N_1})$ at least $d - 1 > 2$. Also $N_1 d \mid M_\omega$, by inspecting the powers of 2, 3 and the remaining primes.

Let Φ be the set of archimedean places, each represented by the embedding extending our fixed embedding of K. We identify Φ with $\mathrm{Gal}(E/K)$ using one chosen embedding of E. Write $[j] \in \Phi$ for the residue class of j modulo N_1 if it is invertible, and for 1 otherwise (this latter convention occurs in the formal prime sums but contributes nothing when $\chi_K(p) = 1$).

Put $\mathcal{T} = \{\tau \in \Phi : \tau \equiv 1 \pmod{N_0}\}$. Then

$$|\mathcal{T}| = T_s, \qquad f = |\Phi| = \varphi(N_0)T_s.$$

For primes $p \nmid N_1 d$ the Frobenius in $\mathrm{Gal}(E/\mathbb{Q})$ has components $(\chi_K(p)p, \chi_K(p))$ in the identification $\Phi \times \{\pm 1\}$ in which the second factor is complex conjugation. This follows because the actual automorphism sends ζ_{N_1} to its p-th power and acts on K by $\chi_K(p)$. In particular a prime of E above p has residual degree the order of this pair. For inert p in K, the sign correction $-p^j$ at odd iterates is the origin of the place permutations in the local prime sums (Chapter 2). For these assertions see `New/Data/Different.lean` and `New/Data/WeilDataOfE.lean`.

The coprime discriminants imply $\mathcal{O}_E = \mathcal{O}_K[\zeta_{N_1}]$: the integral tensor product is maximal after localizing at each prime, since at least one factor is unramified there. Thus the relative different over K is generated by $P'(\zeta)$ for $P(X) = \Phi_{N_1}(X)$, the cyclotomic polynomial, and the absolute different is generated by $\sqrt{-d}\,P'(\zeta)$. The standard cyclotomic integer/different facts are recalled in [5]. The reciprocity relation $P(X) = X^D P(1/X)$, with $D = \deg P$ even, gives

$$P'(\zeta^{-1}) = -\zeta^{2-D}P'(\zeta).$$

Consequently multiplication by $i\zeta^{1-D/2}$ makes $P'(\zeta)$ real. We choose a purely imaginary generator ϖ of the absolute different by multiplying the resulting real element by $\sqrt{-d}$. All conjugates of the root-of-unity multiplier $i\zeta^{1-D/2}$ have modulus 1. In particular

$$|\tau(\varpi)| = \sqrt{d}|P'(\tau\zeta)|, \qquad \mathrm{rd}_E^f = \prod_{\tau\in\Phi} |\tau(\varpi)|, \tag{4.1}$$

where $\mathrm{rd}_E = |\operatorname{disc} E|^{1/[E:\mathbb{Q}]}$. The second equality follows by taking the square root of the absolute norm of the different. The identity

$$\Phi_{N_1}(X) = \Phi_{N_0}(X^{T_s})$$

follows either by listing roots or by successively applying $\Phi_{pN}(X) = \Phi_N(X^p)$ for $p \mid N$. Differentiating at ζ and using $|\tau\zeta| = 1$ shows that every $|P'(\tau\zeta)|$ is T_s times one of the fixed positive numbers $|\Phi'_{N_0}(\zeta_0)|$ for primitive N_0-th roots ζ_0. Hence $\mathrm{rd}_E \asymp_{N_0,d} T_s$. More specifically along the data below, for all sufficiently large x,

$$x^{10} \le |\tau(\varpi)|, \mathrm{rd}_E \le x^{12}. \tag{4.2}$$

This is the numerical content of `hasLogBound_varpi` and `hasLogBound_rootDisc` in `New/Seesaw/IndexTypes.lean`. The ensuing Weil data also choose a splitting character and compatible finite ideles; we construct these choices in Chapter 6, since different choices of theta normalizations can change the period and volume identities used below.

4.2 The scales, and the reason for so many parameters

The principal scale is x; the goal is to estimate $R_{N_0}(x)$. The proof uses another cutoff x_1 and a large order scale b, with

$$(\log b)^{100} \le x, \qquad x^{10^6} \le x_1, \qquad x_1^{10^6} \le b, \qquad (\log x_1)^6 \le \log b.$$

These inequalities are intentionally very far apart. They allow polynomial losses in the ranks or field degree to be negligible compared with the principal weights. For each fixed $0 < \varepsilon < 1/100$, the construction starts with $\log b = x^\varepsilon$ and $\log x_1 = x^{\varepsilon/6}$. The second cutoff will later be lowered within an admissible window. In the compact edition it is a real scale; no rounding of x is required.

Parameter	Meaning	Size or relation
m	Number of lifts	$\sqrt{x} \le m \le 2\sqrt{x}$
r	Rank of initial target group	$x^5 \le r \le 3x^5$
k	Rank of initial source group	$k = r + 1 - m$
t	Growth of row ranks per stage	$t = 5r$
h_2	Half-gap in the weights	$h_2 = r^8$
N_1	Cyclotomic order	$N_0 \mid N_1$
$\lvert\mathcal{T}\rvert$	Number of tops	$\asymp x^{11}$ up to log factors
$\lvert\mathcal{S}\rvert$	Number of thick places	$\ge \lvert\mathcal{T}\rvert / \log b$
a	Order of distinguished derivatives	$xb \le a < xb + N_1$
$g = g_0 g_1$	Product of two level primes	$b^{20} \le g \le b^{22}$
N	Extra level prime for source	$m \log b \le \log N \le 30m \log b$

The exact powers 10^6, 100 and 99 which occur in the source are generous margins; one cannot replace them without checking the many subsidiary inequalities, but they are not the source of the exponent $1/2$ in GRH. The significant power is the square-root length of the chain. The choice of m as a power of 2 and r as a power of 3 is compatible with the displayed intervals and serves the discrete combinatorial constructions in the proof.

The effective weights at the infinite places come in two sizes:

$$w_{\mathrm{nom}}(\sigma) = \begin{cases} a/x, & \sigma \in \mathcal{S}, \\ a/x_1, & \sigma \notin \mathcal{S}. \end{cases}$$

The integral weights actually used are rounded upwards to multiples of N_1 and shifted: if $w^-(\sigma)$ is the rounding of $w_{\mathrm{nom}}(\sigma)$ then $w_j(\sigma) = w^-(\sigma) + 2jh_2 + h_2$. Hence neighboring weights have a large gap. This separation permits quantitative archimedean and local estimates while the naive weights still record the two prime cutoffs x and x_1. It is important not to confuse the large weights with the logarithmic scale $\log b$.

4.3 Admissibility and construction of the data

It is useful to give the full existence proof rather than appeal to the entirety of the formal parameter-search module. The following is the portion of the definition of `Data` in `New/Data/AdmissibleData.lean` that is needed to begin the construction. Write $Y = \log b$, $U_1 = \log x_1$. There is one fixed positive constant c_* depending only on N_0, d (the source's `Parameters.c`, which includes the uniform CM generator constant), and the scales satisfy

$$10^{12} c_* \le \log Y, \quad 100 \log Y \le \log x, \quad 10^6 \log x \le U_1, \quad 10^6 U_1 \le Y, \quad 6 \log U_1 \le \log Y. \tag{4.3}$$

In addition to the rank intervals in the preceding table, require $m = 2^u$ with $u \ge 1$, $s = 22u$, $r = 3^v$ with $v \ge 1$, and $s \le 32 \log x$. The thick set lies in the coset -1 modulo $N_0 3^{c_s}$ and has the following property. If $\phi : \Phi \to (\mathbb{Z}/2^{s+1})^\times$ is reduction and $0 < i < j \le 324s$, then

$$\sigma, \tau, \rho \in \mathcal{S}, \quad \phi(\tau/\sigma)^j = \phi(\rho/\sigma)^i \implies \phi(\tau/\sigma) = 1. \tag{4.4}$$

The two distinct primes g_i lie in $[b^{10}, b^{11}]$ and are 1 modulo M_ω^2; there is $\pi_g \in \mathcal{O}_K$ with $\pi_g \bar{\pi}_g = g_0 g_1$. Finally the data include prime existence in the windows

$$\forall U \in [100Y, 2^s] \quad \exists p \text{ prime}, \quad p \equiv 1 \pmod{M_\omega^2}, \quad U \le \log p \le U + \sqrt{Y}. \tag{4.5}$$

The source defines the entire record before constructing any automorphic forms. All our subsequent spaces depend on this common record.

Lemma 4.1 (Behrend construction in a projected cyclic group)**.** *Let G and H be abelian groups, $\phi : G \to H$ a homomorphism, $u, z \in G$, and suppose the order of $\phi(u)$ is D. Let L_0, h, Q be positive integers with $(L_0Q)^h \le D$ and $L_0 \ge 2$, $Q \ge 2$. There is a subset S of $z\langle u\rangle$ of size at least $Q^h/(hQ^2)$ such that, for $s_0, s_1, s_2 \in S$ and $0 < i < j \le L_0$, the relation $\phi(s_1/s_0)^j = \phi(s_2/s_0)^i$ implies $s_1 = s_0$.*

Proof. Consider the Q^h integer vectors $v = (v_0, \dots, v_{h-1})$ with $0 \le v_a < Q$. The squared Euclidean lengths have at most $h(Q-1)^2 + 1 \le hQ^2$ values; choose a sphere containing at least the asserted number of vectors. Encode a vector by $e(v) = \sum_{a=0}^{h-1} v_a (L_0Q)^a$ and take $s(v) = zu^{e(v)}$. The map $v \mapsto \phi(s(v))$ is injective since $0 \le e(v) < D$ and the encoding is injective. The assumed multiplicative relation becomes

$$je(v_1) \equiv ie(v_2) + (j-i)e(v_0) \pmod{D}.$$

Both sides are integers with digits at most $L_0(Q-1) < L_0Q$ in base L_0Q, hence lie in $[0, (L_0Q)^h) \subset [0, D)$. The equality therefore holds digit by digit and $v_1 = (i/j)v_2 + (1 - i/j)v_0$. On a Euclidean sphere a strictly interior convex combination has strictly smaller norm unless the two endpoints are equal. Thus $v_0 = v_1 = v_2$. □

This is the combinatorial ingredient of `exists_three_term_free` in `New/Data/DataExist.lean`; it is a simultaneous Behrend-type construction (compare [4]).

Proposition 4.2 (Existence and flexibility of admissible data)**.** *For fixed N_0, d, $0 < \varepsilon < 1/100$, and every sufficiently large x, one can construct admissible data $\mathcal{D}$ satisfying*

$$Y = x^{\varepsilon}, \qquad U_1 = x^{\varepsilon/6} \ge (10^6 \log x)^2, \qquad |\mathcal{S}| \ge \frac{T_s}{Y}. \tag{4.6}$$

The same record is admissible if U_1 is replaced by any $u \in [10^6 \log x, (10^6 \log x)^2]$, interpreting every derived quantity with the new U_1.

Proof. Take m and r to be the least powers of 2 and 3 which are at least $\sqrt{x}$ and x^5, respectively, and choose s as above. Then $2^s = m^{22}$, $s \le 32 \log x$ eventually, and

$$81s < 3^{c_s} \le 243s, \qquad T_s = 3^{c_s} m^{22} \asymp x^{11} \log x.$$

In particular $x^{11} \le T_s \le x^{11} Y^{1/5}$ for sufficiently large x. The choice of Y, U_1 in (4.6) satisfies (4.3): the two relevant upper comparisons hold as $x^{\varepsilon/6} = o(x^{\varepsilon})$ and $6 \log U_1 = \log Y$; the remaining lower bounds follow from $\varepsilon < 1/100$. This proves the size comparisons used in (4.2).

To choose $\mathcal{S}$, apply Lemma 4.1 with $z = [-1]$, $u = [1 + M']$, $M' = N_0 3^{c_s}$ and $L_0 = 324s$. Put $v_0 = v_2(N_0) \ge 4$. Since $v_2(M') = v_0$ and the order of $1 + M'$ modulo 2^{s+1} is $D = 2^{s+1-v_0}$ when $s + 1 > v_0$, every power lies in the required coset after multiplication by z. The order formula follows inductively from $v_2((1+M')^{2^h} - 1) = v_0 + h$, by factoring successive differences of squares. Fix an integer $h > 44/\varepsilon$ and choose $Q = \lfloor D^{1/h}/L_0 \rfloor$, which is at least 2 for large x. Then

$$|\mathcal{S}| \ge \frac{Q^{h-2}}{h} \gg_{\varepsilon, N_0} \frac{2^{s(1-2/h)}}{(324s)^{h-2}}.$$

For the desired lower bound, compare with $T_s/Y \asymp s 2^s x^{-\varepsilon}$. Since $2^s \asymp x^{11}$, the ratio of the displayed lower bound to T_s/Y is bounded below by a positive constant times $x^{\varepsilon - 22/h}/(\log x)^{h-1}$, which tends to infinity. The sparsity property is immediate from the lemma.

It remains to find the primes. The restricted power-modulus prime-window theorem, proved in Corollary 3.11, applies to $Q_0 = M_\omega^2$ and to the indicated windows, uniformly in

U. It establishes (4.5), and also the existence of distinct primes $p_i \equiv 1 \pmod{M_\omega^2}$ in disjoint windows in $[10Y, 11Y]$, with $i = 1, \dots, h_K + 1$ where h_K denotes the class number of K. For example use the windows $[10Y + 2i\sqrt{Y}, 10Y + (2i+1)\sqrt{Y}]$. All p_i split in K, since $p_i \equiv 1 \pmod{d}$. Choose a prime ideal $\mathfrak{p}_i$ of K above each. By pigeonhole there exist distinct i, j with $[\mathfrak{p}_i] = [\mathfrak{p}_j]$ in the ideal class group. Then $\mathfrak{p}_i\bar{\mathfrak{p}}_j$ is principal; an integral generator π_g satisfies $\pi_g\bar{\pi}_g = p_ip_j$. Put $g_0 = p_i, g_1 = p_j$. This completes the construction of the record.

Lastly, any replacement value of U_1 in the asserted interval is positive, no larger than $x^{\varepsilon/6}$ and at least $10^6 \log x$. It preserves all inequalities in (4.3), while the other fields of the record do not depend on U_1. This proves flexibility. In the code this is used by `eventually_prime Error_le_rpow` to replace the scale in the structure before constructing any derived weights or levels. □

This supplies a conventional proof of `exists_data` using the restricted prime-window lemma `exists_prime_modEq_one_log_mem` in `Lib/Analytic/Primes/AuxiliaryPrimes.lean`, reconstructed in Chapter 3. Neither a uniform prime number theorem nor a pointwise prime number theorem at the second cutoff is required.

4.4 Derived weights, level and integral structures

In the actual integral weights write $a = N_1\lceil xb/N_1\rceil$, $a_1 = N_1\lceil (a/x)/N_1\rceil$ and $a_2 = N_1\lceil (a/x_1)/N_1\rceil$ (denoted b_1, b_2 in the formal source and in the local chapters). Thus $w_0(\sigma) - h_2$ is a_1 or a_2, with $w_j(\sigma) = w_0(\sigma) + 2jh_2$. For $0 \le j \le 2m$ we have

$$w_{\mathrm{nom}}(\sigma) \le w_j(\sigma) \le w_{\mathrm{nom}}(\sigma) + x^{42}, \quad w_j(\sigma) \le 2b, \quad \log w_j(\sigma) \le \log w_{\mathrm{nom}}(\sigma) + 1/x. \tag{4.7}$$

For the first upper bound use $N_1 + (4m+1)r^8 = O(x^{81/2}) + O(x^{11}\log x) \le x^{42}$ eventually. Next $w_{\mathrm{nom}} \le a/x \le b + N_1/x$, and $b \gg x^A$ for every fixed A. Finally $w_{\mathrm{nom}} \ge a/x_1 \ge xb/x_1 \gg x^{43}$ by the scale assumptions, and $\log(1 + x^{42}/w_{\mathrm{nom}}) \le x^{42}/w_{\mathrm{nom}} \le 1/x$. These translate `w_le_nominalWeight_add`, `w_le_two_mul_b` and `log_w_le` in `New/Data/Weights.lean`.

Write $w_* = f^{-1}\sum_\sigma \log w_{\mathrm{nom}}(\sigma)$ and $G_* = \log(g\,\mathrm{rd}_E/(2\pi))$, where $g = g_0g_1$. The extra prime N is selected as in `Data.N`, using the logarithmic target

$$\lambda_N = m(w_* + G_*) - G_* - \tfrac{15}{2}\log b, \quad \lambda_N \le \log N \le \lambda_N + \sqrt{\log b}, \quad N \equiv 1 \pmod{M_\omega^2}. \tag{4.8}$$

The large ranges give $w_* = (1 + o(1))Y$ and $G_* \in [(20 - o(1))Y, (22 + o(1))Y]$, uniformly after replacement of U_1. Hence eventually $mY \le \lambda_N \le 29mY$, $100Y \le mY$ and $29mY \le 2^s = m^{22}$; this uses $Y = x^\varepsilon$. We may therefore use (4.5), giving $mY \le \log N \le 30mY$. Since $4 \mid N_1 \mid M_\omega$, Fermat's two-square theorem furnishes $\pi_N = u + iv \in \mathcal{O}_E$ with $\pi_N\bar{\pi}_N = N$. Every prime above N has residual degree one, and the support of (π_N) contains exactly one member of each conjugate pair (because N is split and unramified and the exponents sum to 1). Likewise the chosen places above g_0, g_1 are the prime divisors of $\bar{\pi}_g$; they are conjugate-disjoint and there are f above each. These observations justify the finite groups and the determinant normalizations used later.

4.5 Two kinds of automorphic forms

If V is a nondegenerate Hermitian space over E with respect to the involution of E/E^+, $U(V)$ denotes its unitary group over the totally real field E^+. Automorphic forms on such a group are functions (with the specified transformation law at infinity) on an adelic quotient. For a totally definite Hermitian space, the group is compact at infinity, and spaces of algebraic

modular forms at a fixed level can be described by values in an algebraic representation on finitely many double cosets. In particular their coefficient integrality and Hermitian pairings can be studied by actual lattice calculations. For the indefinite groups, one instead uses cusp forms of prescribed discrete-series type at infinity. The source imposes suitable finite level and controls their behavior in the cusp.

The proof uses both types. Here are the main spaces; indices on Hermitian spaces should not be confused with indices of vectors.

- The source Hermitian spaces X_j are definite of rank $k + j(t+1)$, $0 \le j \le m$. Let C denote the space of algebraic modular forms on $U(X_0)$ of weight $(a, 0, \dots, 0)$ at tops and zero at the other infinite places, with the prescribed level at g and N. Let $C^{\mathrm{ram}} \subset C$ be the subspace of forms with vanishing averages under the prescribed local scalings at the places of N. These are the "ramified columns."
- The row spaces are forms on Y_j of rank $r + jt$. The starting Y_0 is indefinite, and Y_m is definite. Let W be the selected subspace of forms on $U(Y_0)$, at its prescribed archimedean type and at level g. (We use the letter W only for this vector space; the source also uses W_j for auxiliary rank-r Hermitian spaces.) The spaces selected by a finite-group projector are called good forms.
- For the determinant computation there are also single-lift auxiliary pairs of Hermitian spaces W_j, V_j of ranks $r, r + t$. These are used to replace the norm calculation for a whole chain by the calculations associated to individual lifts and their comparisons.

At the chosen finite places v above g, the principal-level form spaces carry an action of the finite group

$$\mathcal{G} = \prod_v \mathrm{GL}_r(\mathbb{F}_v).$$

The good projector e_0 is the sum of specified idempotents in $\mathbb{C}[\mathcal{G}]$. The good labels are defined through squarefree polynomials over the residue fields, with exclusions at roots of unity and primitive-polynomial conditions. This is more than a harmless way to cut down the dimension: it is designed to guarantee local representation types favorable both to the filling argument and to the Galois-theoretic analysis of congruences. The precise conditions are discussed in the subsequent sections. We write $n = \dim W$ and $c = \dim C$; neither of these is a Hermitian rank.

4.6 Precise weights and levels of the source and target

For a weight at a fixed infinite place we use column multiplicities $\kappa_\tau(h)$, meaning the number of Young-diagram columns of height h. Write $h_0 = 2h_2$ for the weight gap to distinguish it from the dummy height h, and put $n_j^X = k + j(t+1)$. The compact source uses the same multiplicity function at every stage, restricted to heights strictly below the rank of the stage:

$$\kappa_\tau(h) = \begin{cases} a\mathbf{1}_{\tau\in\mathcal{T}}, & h = 1, \\ w_1(\tau) - \lfloor (k+t+1)/2 \rfloor, & h = k, \\ h_0 - (t+1), & h > k,\ h \equiv k \pmod{t+1}, \\ 0, & \text{otherwise.} \end{cases}$$

The determinant exponents are $e_0(\tau) = 0$ and $e_j(\tau) = \lfloor n_{j-1}^X/2 \rfloor - w_j(\tau)$ for $j \ge 1$. These are `Data.` followed by the Greek letter κ, and `Data.lastEntry` in `New/Curve/LiveSum.lean`

(the source uses Greek letters in the identifiers). At X_0 the only column heights below k are the height-one columns supported on the tops, and this recovers the source weight described above. At the endpoint, the branching to Y_m retains

$$u(h) = \begin{cases} h_2 - \lfloor (t+1)/2 \rfloor, & h \geq k,\ h \equiv k \pmod{t+1}, \\ 0, & \text{otherwise}, \end{cases}$$

columns of their former height; its new multiplicity is $u(h) + \kappa_\tau(h+1) - u(h+1)$. Its determinant normalization comes from `restrictToY` in `New/Columns/ColumnDefs.lean`. This weight convention matters: it is not permissible to use the ordinary weight w_{nom} in the representation instead of the rounded weights.

Put $F = E^+$. For a standard definite unitary group of rank q', the coefficient space $V_{\kappa,e}$ is the product-place tensor of the complex Weyl modules with multiplicities $\kappa_\tau(h)$, tensored with the determinant of exponent $e(\tau)$. The basis in the code is its integral Weyl-coordinate basis, and its Gram form has an integral, positive-definite matrix G_κ; the determinant twist is unitary at infinity. The action of $\gamma \in U(q')(F)$ uses $\tau(\overline{\gamma})$ in the algebraic weight representation. For an open compact $K' \subset U(q')(\mathbb{A}_{F,f})$, the algebraic forms of fixed level are functions v on the finite adelic group with values in $V_{\kappa,e}$ satisfying

$$v(\gamma g k') = \rho_{\kappa,e}(\gamma) v(g), \qquad \gamma \in U(q')(F),\ k' \in K'.$$

There are finitely many adelic classes. Choosing a representative g of each class gives the direct sum of invariant coefficient spaces under the groups $\Gamma_g = U(q')(F) \cap gK'g^{-1}$. This follows at once from the equivariance condition. It is the model `AlgForm` of `Lib/GLn/SchurM odule/WeightRepBasic.lean`. The class-sum Hermitian form is

$$\langle v, w \rangle_{K'} = \sum_{[g]} \frac{v(g)^* G_\kappa w(g)}{|\Gamma_g|}.$$

When the level K' has Haar volume one, this class sum is the integral of the invariant scalar pairing over the finite adelic quotient. For the correspondence with highest-weight matrix coefficients of the automorphic form at infinity, the source uses the factor $\dim V_{\kappa,e}$ from Schur orthogonality and the factor $[K_0 : K']$ from its finite measure convention. The precise comparison is `integral_mul_eq_classIntegral`, `Space.algFormEquiv` in `Lib/Adelic/Cus pForms/AlgebraicForms.lean`.

Let K_g denote the level which is principal at the chosen places above g and maximal integral elsewhere. For X_j the column level is K_g intersected with the condition that the last row of the matrix at places dividing π_N is $(0, \ldots, 0, 1)$ modulo that prime. For each $w \mid \pi_N$, a scalar finite adelic unit z_w represents a generator of $\mathbb{F}_N^\times$ in the compatible standard split model. The ramified part of C is exactly

$$C^{\mathrm{ram}} = \mathrm{im} \prod_{w \mid \pi_N} (1 - \rho(z_w)).$$

The scalar operators commute and are of finite order on this level, so this is equivalently the intersection of the kernels of their cyclic averaging projections. It follows that for each fixed w, every ramified column has the form $(1 - \rho(z_w))v$; this elementary observation is used in the ramified kernel integrals. The codimension bound is established in Chapter 20 via the precise local level and class count. Source: `New/Columns/ChainValuation.lean`, `Data.ramifiedColumns` and `exists_eq_sub_centralGen`.

For the row representation, number the basis of Y_0 by $i = 0, \ldots, r-1$, using the numbering compatible with the Bessel flag in the source. At τ the (undoubled) Harish-Chandra parameter

is

$$\varsigma_\tau(i) = \begin{cases} k/2 - 1 - i, & i+1 < k, \\ h_2 - w_{i+2-k}(\tau), & i+1 \geq k. \end{cases}$$

Here k is even. The Harish-Chandra coordinates of all the row spaces, together with the signs specifying their real form, are chosen so that the parameter gap between coordinates of opposite signs is at least $2h_2$ in the doubled coordinate convention of the code. In particular the rank-r auxiliary space W_j is specified by the first $k+j-1$ positive and remaining $m-j$ negative coordinates, while Y_j has rank $r+jt$ and $\lfloor (m-j)/2 \rfloor$ negative signs at each place. On Y_j the doubled new blocks have coordinates $t-1-2w_l(\tau)-2b'$ $(0 \leq b' < t$, $1 \leq l \leq j)$; all doubled coordinates, including the base coordinates, are reduced by one if j is odd. The starting Y_0 is indefinite and the endpoint Y_m definite. The form of the row spaces and their adelic frames is part of the construction: the code gives `Data.exists_spaceY` in `New/Trace/GoodIdempotent.lean`, using the explicit glued Gram matrix and an integral frame. We regard the existence of this frame with its specified congruences as a formal-source input, and retain the flags and identifications of that construction (see Chapter 14).

The row form space before the good projection consists of automorphic functions of the indicated discrete-series type, right K_g-fixed. More precisely the source uses the subspace fixed by convolution with the reproducing kernel of the prescribed real type and demands rapid decay; its finite-dimensionality is the assertion `finite_levelSpace` for the cusp kernel. The fixed-level convolution representation of K_0/K_g at the split chosen places factors through

$$\mathcal{G} = \prod_{i=0,1} \prod_{w|\bar{\pi}_g,\ w|g_i} \mathrm{GL}_r(\mathbb{F}_{g_i}).$$

The good projector e_0 is a central idempotent of its group algebra, and $W = e_0 V$ with V the row form space at the principal level. The conditions and dimension estimates for e_0 are handled in the spanning chapter. We will write $C^{\mathrm{ram}}_{\mathrm{int}}$ for the forms in C^{ram} integral at every embedding and use $n = \dim_{\mathbb{C}} W$, $c = \dim_{\mathbb{C}} C$ throughout.

4.7 Integral gauges and the precise uniform index

For each rational prime p and field isomorphism $\iota : \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_p$, let $\mathscr{O}_\iota$ be the valuation ring pulled back by ι. If v is an algebraic form, the gauge of its value $v(g)$, with respect to the lattice at g, is

$$\|v(g)\|_{\iota,g} = \left\| \rho_{\kappa,e}\left((g^{-1})_{\iota\tau\circ\mathrm{conj}} \right)_{\tau\in\Phi} \iota(v(g)) \right\|_{\sup,p}.$$

The component matrix is obtained from the finite adele by extension to $\overline{\mathbb{Q}}_p$ via $\iota\tau \circ \mathrm{conj}$; coefficients of the Weyl matrix are integers, and the sup norm is taken in its integral coordinate lattice. This is `weylGauge` in `Lib/GLn/SchurModule/WeightRepBasic.lean`. Integral means every such gauge is at most one. Because the representation action respects the transported lattice, one may test on the finite class representatives. Notice that integrality is not defined merely by integrality of the complex coordinates at $g = 1$. This is significant when local Hecke modifications change the lattice or the branching map transports a different lattice at N.

For a family $y = (y_i)_{i=1}^q$ of row forms, the index inequality $\mathrm{IndexLe}(y, d_{\mathrm{ind}})$ uses real numbers e_p such that $\sum_{p\in S} e_p \leq d_{\mathrm{ind}}$ for every finite set S of primes. For each p, ι and every allowed m-fold twist θ of conductor prime to p, there must be a $q \times q$ matrix U over $\mathscr{O}_\iota$ satisfying

$$|\iota(\det U)|_p \geq e^{-e_p}, \qquad \Theta_\theta\Big(\sum_j U_{ij} y_j \Big) \text{ integral at } \iota \quad (1 \leq i \leq q).$$

The matrix is not required to be the same for different twists, but the budgets e_p are fixed before them. This is precisely `Data.IndexLe` in `New/Endgame/ShortAndEnough.lean` (there the field isomorphisms are supplied as instances). These conventions explain the factor $2d_{\mathrm{ind}}$ in the squared-determinant estimate. There is a stronger local assertion on column lattices, before applying the projected Poincaré map: the corresponding submodule is independent of the twist. The short-form argument uses this stronger assertion to choose change-of-family matrices with a global index bound.

4.8 The theta chains and their relation

Let θ be a weight-zero automorphic character of $U(1)$, with tame conductor at a set of places disjoint from its complex conjugate and prime to gN. The successive powers through θ^m must retain the required ramification at that conductor (the trivial character is also allowed). A theta lift is a linear transformation defined by pairing an automorphic form on one group with a theta kernel associated to a Weil representation on a dual pair. Schematic notation for a lift is

$$(\mathcal{L}F)(h) = \int_{[U(V)]} F(u)\,\overline{\vartheta(u,h)}\,du,$$

where the precise conjugations, kernels, measures and characters depend on the convention. The source fixes these choices by its definitions of the spaces and lifts; this schematic integral is a description of the idea, not a replacement for those definitions. For definite groups, such integrals reduce to sums over finitely many classes.

Denote by Θ_θ the composite lift along the row chain beginning in W. The column chain begins in C, runs along the definite X_j, and afterwards restricts from X_m to Y_m. The ranks have been chosen so that

$$\dim_E X_m = k + m(t+1) = r + mt + 1 = \dim_E Y_m + 1.$$

The one-dimensional complement is used in the branching and level normalization. The m extra rank increments on the two sides make the iterated seesaw identity possible. Informally the relationship is

$$\begin{array}{ccccc} C^{\mathrm{ram}} & \longrightarrow & \text{definite chain} & \longrightarrow & X_m\text{-forms} \\ \downarrow P & & & & \downarrow \text{restriction} \\ \text{initial row forms} & \longrightarrow & \text{row chain} & \longrightarrow & Y_m\text{-forms.} \end{array}$$

Here P is a Poincaré series constructed from a source form by Bessel induction. The actual identity involves a finite combination of definite chains, nonzero normalization factors and finite-place restrictions, not a literal equality between arbitrary individual theta kernels. The projected translates P_s in the upper bound, for $s \in \mathcal{G}$, are obtained from the Poincaré map by translating and projecting; we include the fixed normalizing factors from `projectedPoincare` in this notation.

The two rows of the diagram serve different purposes. The definite chain gives access to coefficient lattices and the sizes and prime divisibilities of derivatives of theta functions. The indefinite row chain gives a large automorphic space, a spanning theorem for Poincaré series, and a determinant calculation. Neither side alone yields the main inequality.

4.9 Norms, squared covolume and index

For each field automorphism σ of $\mathbb{C}$, the target definite space has a conjugated coefficient pairing, with squared norm denoted $\|\cdot\|_\sigma^2$. We use the pairing obtained by conjugating

the coefficient data as specified in `Lib/Adelic/CuspForms/LevelForms.lean`, not the absolute value of an arbitrary conjugate of a real analytic integral. For independent vectors $y_1, \dots, y_q \in W$ define

$$\mathsf{G}_\sigma(y) = \left(\langle \Theta_1 y_i, \Theta_1 y_j \rangle_\sigma\right)_{i,j}, \qquad \operatorname{covol}_\sigma^2(y) = |\det \mathsf{G}_\sigma(y)|.$$

The determinant may be zero for some family before an injectivity statement is available; the upper estimate does not presuppose it is positive. We call V a logarithmic upper bound if $\operatorname{covol}_\sigma^2(y) \le e^V$ for all σ. By the Gram determinant inequality,

$$\operatorname{covol}_\sigma^2(y) \le \prod_{i=1}^{q} \|\Theta_1 y_i\|_\sigma^2.$$

Thus bounds on individual lifts imply a bound on the determinant of their span. The source's definition uses the notation `covolumeSq`.

The paired finite-place structure and the index bound were fixed in Section 4.7.

4.10 The common leading term and the prime remainder

Here we introduce the quantities that occur in all three theorems. Let $\lambda_{d,1}(\nu) = \Lambda(\nu)(1-\chi_K(\nu))$ be the weight without a congruence restriction, and let s_ν be the place represented by $-\nu$ modulo N_1 when ν is invertible there (with the source's convention of the identity place otherwise). Define the finite prime sum at $\sigma \in \Phi$ by

$$A(y, \sigma) = \sum_{1 \le \nu \le a} \lambda_{d,1}(\nu) \left(\frac{y}{\nu} - w_{\mathrm{nom}}(s_\nu \sigma) \right)_+ .$$

Write $\kappa_K = \gamma - \Re(L'(1,\chi_K)/L(1,\chi_K))$ for the Chowla–Selberg constant used in the proof, and

$$\Omega = \frac{2\pi}{\mathrm{rd}_E} e^{\kappa_K}, \qquad J(j) = \log(j!) - j\log(\Omega/g), \qquad \overline{w} = \frac{1}{f}\sum_{\sigma \in \Phi} \log w_{\mathrm{nom}}(\sigma).$$

The quantity $J(j)$ is the archimedean cost of a derivative of order j in the normalization of the proof. Its dependence on the period of the CM curve is the key to matching archimedean terms with the weighted prime sum. An expected main term for $A(a,\sigma)$, averaged appropriately, is

$$A_0 = a(\log a - 1 - \kappa_K - \overline{w}).$$

The exact quantities in the inequalities are

$$\begin{aligned} \mathcal{R} &= \sum_{\sigma \in \mathcal{T}} \left(A(a,\sigma) - A_0\right), \\ \mathcal{U} &= \sum_{\sigma \in \Phi} \left(J(w_0(\sigma)) - A(w_{\mathrm{nom}}(\sigma), \sigma)\right), \\ H &= mk\mathcal{U}, \qquad B = fa(\log b)^3, \qquad L = B\log b = fa(\log b)^4. \end{aligned}$$

These definitions are `Data.primeSum`, `Data.jetSize`, `Data.dirCost`, `Data.mainHeight`, `Data.mainAsk`, and `Data.remainder`. The quantity $\mathcal{U}$ represents the net cost of an ordinary differentiation direction, obtained by subtracting the finite-place divisibility gain from the archimedean derivative size. Both the short-form argument and the determinant argument are calibrated to this quantity. It is not necessary to determine its precise sign or asymptotic value.

The prime remainder has the following exact interpretation, which will be important in the final deduction:

$$\mathcal{R} = a\{|\mathcal{S}|(R_{N_0}(x) - R_{N_0}(x_1)) + |\mathcal{T}|R_1(x_1)\}. \tag{4.9}$$

This is `Data.remainder_eq`. We explain its mechanism in Chapter 55: the two weights replace the finite prime sums by R_{N_0} at two scales and a remainder R_1 without a congruence condition. The role of the analytic choice of x_1 is to control the unwanted second-cutoff expression. Notice that the normalized error $R_M(z)$ of the preceding section is not the same as the unnormalized, large remainder $\mathcal{R}$.

Chapter 5

Adelic coefficient models and integral linear algebra

The proof uses automorphic forms on unitary groups in two rather different ways. An indefinite group supplies a spectral and integration problem; a definite group supplies a finite class set equipped with a coefficient lattice. The passage between the two is subtle. This chapter explains the elementary foundations of the definite model, the local integrality tests and the exact meaning of the norms used later. It also establishes several facts about these models directly. It does not assume a general comparison between an automorphic integral lattice and a Galois-representation lattice. The latter is a particularly substantial issue in Chapters 40 and 49.

The reader familiar with algebraic modular forms may skip most of this chapter, but should retain the normalization of the coefficient pairing in (5.6), the freeness criterion 5.2, and the distinction between the Weyl lattice and the lattice of polynomials in Lemma 5.7. These are significant for the quantitative statements of the proof. The finite-class-number framework is classical (see [6, 7]); the arguments below are given for the particular totally definite Hermitian spaces in use here.

5.1 Hermitian spaces, places and adelic lattices

Let E/F be a CM extension with $F = E^+$, equipped with conjugation $z \mapsto \bar{z}$. Its set of real places is denoted Φ after choosing one complex embedding of E above each real place. Let V be an E-vector space of dimension l with nondegenerate Hermitian form h; we use conjugate linearity in the first argument and linearity in the second. In a basis the form is $h(u, v) = u^* J v$ for a Hermitian matrix J. Define

$$U(V)(R) = \{A \in \mathrm{GL}_l(E \otimes_F R) : A^* J A = J\}$$

for an F-algebra R, with conjugation extended R-linearly. A totally definite space is positive definite at every complex embedding (changing the sign at a real place makes no difference to the compactness statement). Then $U(V)(F \otimes_{\mathbb{Q}} \mathbb{R}) \simeq \prod_{\tau \in \Phi} U(l)$ is compact. In the standard totally definite model of the proof $J = I$ and the lattice is $\Lambda_0 = \mathcal{O}_E^l$. Write $G = U(V)$.

For a finite place v of F, if v splits in E as $w\bar{w}$, then $E \otimes_F F_v \simeq F_v \times F_v$, with conjugation interchanging the two factors. A unitary matrix is determined by its matrix in the w factor; the other component is forced by the equation defining G. In the standard model this is

$$A \longmapsto (A, A^{-T}), \qquad A \in \mathrm{GL}_l(F_v). \tag{5.1}$$

At an inert finite place the group is genuinely unitary, not a copy of the general linear group; the residue field and Frobenius actions differ. The proof repeatedly uses the splitting at the selected primes g_0, g_1, N but cannot use such a local model at nonsplit primes in the jet argument. The precise choice of a member of each split pair matters for coefficient valuations: using the conjugate of the place changes the matrix A to A^{-T}. This is why the source records the finite-place embeddings of each CM-type factor explicitly.

The finite adelic group $G(\mathbb{A}_{F,f})$ is the restricted product of $G(F_v)$ with respect to the local stabilizers of $\Lambda_0 \otimes \mathcal{O}_{F_v}$; all but finitely many components of an element are integral invertible. Put $K_0 = \prod_v \mathrm{Stab}(\Lambda_{0,v})$, a compact open subgroup. A choice of finite adelic g defines a fractional $\mathcal{O}_E$-lattice

$$\Lambda_g = V \cap (g\widehat{\Lambda}_0), \qquad \widehat{\Lambda}_0 = \prod_w (\Lambda_0 \otimes_{\mathcal{O}_E} \mathcal{O}_{E,w}),$$

where V embeds diagonally in $V(\mathbb{A}_{E,f})$. The meaning of the intersection is that $x \in V$ satisfies $x \in g_w\Lambda_{0,w}$ at all finite places. Conversely a family of local lattices L_w equal to $\Lambda_{0,w}$ outside a finite set determines a lattice with these completions. One proof is as follows. Choose a nonzero ordinary integer c such that $c\Lambda_{0,w} \subseteq L_w \subseteq c^{-1}\Lambda_{0,w}$ at every w. The desired lattice is the inverse image in $c^{-1}\Lambda_0$ of the product of the corresponding subgroups in $c^{-1}\Lambda_0/c\Lambda_0$, using the Chinese remainder decomposition of this finite module. By construction its completions have exactly the desired images and contain $c\Lambda_{0,w}$, proving equality. In particular two lattices with the same completions coincide.

Proposition 5.1 (Adelic class interpretation)**.** *In the standard totally definite model the double quotient*

$$X_{K_0} = G(F)\backslash G(\mathbb{A}_{F,f})/K_0$$

is naturally the set of isometry classes of Hermitian lattices in V locally isometric to Λ_0 at all finite places. For every open subgroup $K' \subseteq K_0$, the class set $X_{K'}$ is finite and every stabilizer $\Gamma_g = G(F) \cap gK'g^{-1}$ is finite.

Proof. The assignment $g \mapsto \Lambda_g$ is surjective onto the indicated genus: choose a local isometry at each exceptional place and the identity where the lattices are equal. If $\gamma\Lambda_g = \Lambda_{g'}$, then at each completion $(g')^{-1}\gamma g$ preserves the standard local lattice, so it belongs to K_0. This proves the class interpretation. For stabilizers, the underlying positive trace form

$$Q(u, v) = \mathrm{Tr}_{E/\mathbb{Q}}\, h(u, v)$$

on a lattice Λ_g is a positive-definite symmetric integral form. Its integral isometries are finite: the images of any fixed basis under an isometry must lie among the finitely many lattice vectors of the prescribed lengths. Hence Γ_g is finite, even at level K_0.

Here is a proof of finiteness of X_{K_0}. The rank $D = l[E : \mathbb{Q}]$ and determinant of the trace form on Λ_g are fixed, the latter because all of its local completions are isometric to the fixed lattice. We first use the elementary consequence of Minkowski's second theorem ([8]) that there are only finitely many integral positive-definite lattices of fixed rank and fixed determinant up to isometry. Indeed, for the Euclidean length determined by Q, every nonzero vector has norm at least one. If $\lambda_1 \le \cdots \le \lambda_D$ are successive minima, Minkowski's bound on their product at fixed covolume gives an upper bound on every λ_i. We may choose D independent lattice vectors of uniformly bounded length. Their Gram matrix is an integral matrix with uniformly bounded entries; the index of their span in the original lattice is uniformly bounded by Hadamard's inequality for their parallelepiped divided by the fixed covolume. A lattice containing this span with index I is intermediate between that span and its scalar multiple

by $1/I$; for bounded I there are only finitely many such lattices. This proves the claimed finiteness of positive integral trace forms.

For each representative of the trace lattice, consider the possible $\mathcal{O}_E$-actions. Multiplication by each element u of a fixed integral basis of $\mathcal{O}_E$ has operator norm at most $\max_{\tau\in\Phi}|\tau(u)|$ on the trace form: the real orthogonal decomposition into CM factors shows this directly. Therefore there are only finitely many integral matrices for these actions on a fixed trace lattice. An isometry preserving the trace form and these actions preserves h, since the values of $\mathrm{Tr}_{E/\mathbb{Q}}(ah(x,y))$ for $a\in E$ determine $h(x,y)$ by nondegeneracy of the trace pairing. Indeed these values are $Q(x,ay)$; thus the Hermitian form is recovered from the trace form and the E-action. This reduces Hermitian lattice classes to a finite collection. Finally $[K_0:K']<\infty$ by compactness and openness, so each class at level K_0 splits into finitely many classes at level K', as asserted. □

This proof gives the qualitative finiteness assertion used in the algebraic modular form theory. It is not a quantitative mass estimate for varying fields and ranks; the counting inequalities in the main arithmetic construction use further results. In the code the applicable abstract setup is `HasFiniteType` in `Lib/Adelic/Hecke/ClassMean.lean`. The argument above suffices to justify it in the standard definite case, with maximal level and all its open subgroups.

5.2 Why the extra level removes stabilizer denominators

Although every stabilizer in Proposition 5.1 is finite, its order could divide a small prime and spoil integrality of a class sum. The selected primes are far larger than required to force triviality. The following trace argument gives the exact bound used in the compact source.

Lemma 5.2 (Large residue norm and free level)**.** *Let G be totally definite of rank l over E/F. Suppose $K'\subseteq K_0$ is a subgroup reducing to identity at distinct places $w_1,\dots,w_a$ of E. If*

$$(2l)^{[E:\mathbb{Q}]}<\prod_{i=1}^{a}\#k_{w_i}, \tag{5.2}$$

then $\Gamma_g=\{1\}$ for every $g\in G(\mathbb{A}_{F,f})$.

Proof. If $\gamma\in\Gamma_g$, its characteristic polynomial is monic with coefficients integral at every finite place, because it is locally conjugate to an integral matrix. In particular $z=l-\operatorname{tr}\gamma\in\mathcal{O}_E$. At every archimedean embedding all eigenvalues of γ have modulus one, so $|\tau(z)|\le 2l$. At the chosen finite places, conjugacy to an integral matrix reducing to the identity gives $z\in\mathfrak{p}_{w_i}$. If $\gamma\neq 1$, the trace cannot equal l: in a positive-definite factor, $\Re(\operatorname{tr}\gamma)=l$ forces every eigenvalue to be one, and a unitary matrix is diagonalizable. Thus $z\neq 0$. The absolute norm of a nonzero element in the product of the distinct prime ideals is at least the product of their norms. It is also at most $(2l)^{[E:\mathbb{Q}]}$ by the archimedean bound. This contradicts (5.2). □

In the proof's notation the chosen places occur over g_0,g_1, with f places over each prime, all of residue degree one. The right side of (5.2) is g^f, $g=g_0g_1$. As $[E:\mathbb{Q}]=2f$, it suffices that $(2l)^2<g$. For all the definite ranks used in the relevant class sums, $l\le 2(k+m(t+1))$, this follows from $g\ge b^{20}$ and the polynomial bounds in x for m,r,t. Therefore principal level at the chosen places is free. This proves the special case called `Data.classAut_eq_bot_of_le_chosenLevel` of `New/Data/ChosenPlaceLevels.lean`, which uses `classAut_eq_bot` in `Lib/Shimura/IntegralModel/NeatLevel.lean`. No general assertion about neat Shimura varieties is being imported here.

Corollary 5.3 (An integer clearing all stabilizer orders)**.** *Assume $l \le rt$ and that the large-residue condition holds for rank l. Put*

$$D_g = \prod_{w \text{ chosen}} |\mathrm{GL}_{rt}(k_w)|.$$

For every subgroup of maximal integral level in rank l and every adelic class, the order of its rational stabilizer divides D_g.

Proof. Conjugate the finite stabilizer by the representative g and reduce at the chosen places. Its kernel is a stabilizer at the principal level, hence trivial by the lemma. The reduction thus embeds the stabilizer in $\prod_w \mathrm{GL}_l(k_w)$. The latter group embeds by block diagonal matrices in $\prod_w \mathrm{GL}_{rt}(k_w)$, so Lagrange's theorem proves the assertion. □

This is the integer denoted `Data.chosenGroupOrder`. Notice that it is shared across every lower rank and every level in the specified range; using a different clearing integer for each class would be inadequate for the simultaneous determinant estimates. Its size can be bounded elementarily: $|\mathrm{GL}_L(k)| = \prod_{j=0}^{L-1}(|k|^L - |k|^j) < |k|^{L^2}$. Thus

$$\log D_g \le f(rt)^2 \log g.$$

In particular $\log D_g = O(fr^4 \log b)$ since $t = 5r$. With the scales $f = O(x^{11} \log x)$, $r = O(x^5)$, $\log b = x^{\varepsilon}$, this is $O(x^{31+\varepsilon} \log x)$ and hence is at most x^{32} eventually. This supplies a direct explanation of the apparently large polynomial bound appearing in the row budgets. It is a bound for the logarithm of the integer, not for the order itself.

5.3 Normalization of class sums and change of level

Let $G_f = G(\mathbb{A}_{F,f})$, $\Gamma = G(F)$, $K' \subseteq K_0$ open compact. For a left Γ-invariant and right K'-invariant function F taking values in a characteristic-zero vector space define

$$\mathcal{C}_{K'}(F) = \sum_{[g] \in \Gamma \backslash G_f / K'} \frac{F(g)}{|\Gamma_g|}, \qquad \Gamma_g = \Gamma \cap gK'g^{-1}.$$

This is a finite weighted sum, independent of the choice of representatives. The subscript does not mean that the measure of the whole adelic quotient is one. If Haar measure on G_f has $\mathrm{vol}(K_0) = 1$, then the quotient integral is $\mathrm{vol}(K')\mathcal{C}_{K'}(F) = \mathcal{C}_{K'}(F)/[K_0 : K']$. Indeed, the preimage of the class $[g]$ in G_f is the union of the translates $\gamma g K'$; such translates coincide precisely modulo Γ_g. For rational scalar expressions the class sum can be read in an arbitrary $\mathbb{Q}$-vector space without passing through real-valued integration. This is useful in comparing coefficient conjugates or applying p-adic valuations.

Lemma 5.4 (Level change)**.** *If $K'' \subset K'$ is open and F is right K'-invariant, then*

$$\mathcal{C}_{K''}(F) = [K' : K'']\mathcal{C}_{K'}(F).$$

Consequently $\mathcal{C}_{K'}(F)/[K_0 : K']$ is independent of the chosen level on which F is fixed.

Proof. Inside a class $[g]$ at level K' the finer classes are the orbits of the finite group $g^{-1}\Gamma_g g$ acting on the finite set K'/K'' on the left. Each finer stabilizer is conjugate to the stabilizer of the corresponding orbit representative. By the orbit-stabilizer formula, the sum over its orbits of the reciprocal stabilizer orders equals $[K' : K'']/|\Gamma_g|$. Multiply by the common value $F(g)$ and sum over the original classes. □

If the base level K_0 is fixed but a function is only invariant at some smaller level, its normalized integral is well-defined by the same lemma, using intersections of two levels. Translation by a finite adele need not preserve K' but preserves the normalized integral on smooth left-invariant functions, by Haar invariance; one may prove it algebraically by refining to $K' \cap aK'a^{-1}$. The unitary groups are unimodular, so left and right Haar measures agree. The compact source proves the needed class-sum identities in `Lib/Adelic/Hecke/ClassMean.lean`, notably `classSum_eq_relIndex_smul`. In a finite sum with coefficients in a valuation ring O, $D_g \mathcal{C}_{K'}(F)$ remains in O whenever F is O-valued and D_g is divisible by all stabilizer orders. At a free level D_g is unnecessary.

One should keep these three normalizations separate:

- the counting measure on a finite free class set, i.e. $\mathcal{C}_{K'}$;
- the Haar quotient measure with normalized maximal level (or a specified fixed level);
- the normalization of the real representation matrix coefficient by formal degree or coefficient dimension.

A ratio of theta periods computed with the first convention may involve extra factors of finite indexes when inserted into the second. The determinant height computation uses exactly these factors when moving between levels.

5.4 A useful elementary form of weak approximation

The proof requires a finite-place test of the coefficient lattice. It is not necessary to assume that the rational unitary group is dense in all finite adeles, which is in general false. Weak approximation at finitely many places is sufficient.

Lemma 5.5 (Weak approximation for the unitary group)**.** *Let J be any nondegenerate Hermitian form over the CM extension E/F. For a finite set of places of F the diagonal $U(J)(F)$ is dense in the product of the corresponding local unitary groups.*

Proof. Consider the F-linear space $\mathfrak{u}_J$ of matrices satisfying $X^*J + JX = 0$. On the open subset $\det(I + X) \neq 0$, the Cayley transform

$$X \longmapsto (I - X)(I + X)^{-1}$$

is unitary. This is verified by expanding $(I - X)^*J(I - X) = (I + X)^*J(I + X)$, and its inverse on the open subset $\det(I + A) \neq 0$ is $A \mapsto (I - A)(I + A)^{-1}$. The verification also holds after completion at each chosen place. To put a given finite tuple of local matrices A_v into these opens, multiply them by a common rational scalar from $U(1)(F)$. Indeed, choose any nonzero $\eta \in E$ with $\bar{\eta} = -\eta$; the elements $(1 + t\eta)/(1 - t\eta)$ for rational t are distinct elements of $U(1)(F)$, and for each fixed local A_v only finitely many scalar values make $I + cA_v$ singular. Choose c outside the finite union. For these local matrices the inverse Cayley coordinates are elements of $\mathfrak{u}_J(F_v)$. Approximate them simultaneously by a rational point of the finite-dimensional F-vector space $\mathfrak{u}_J$, using ordinary weak approximation in F. By continuity the rational Cayley transform approximates cA_v arbitrarily closely; multiplying by c^{-1} gives the required approximation. □

This gives the consequence of `Space.weak_approximation` in `Lib/Adelic/Landherr/WeakApprox.lean` used to pick p-integral representatives. In particular if g is a finite adele and p is rational, take a rational γ so that the components of γg at all places over p lie in the local

standard maximal integral groups. The required conditions are open and contain the tuple g_p^{-1} before translation. This makes it possible to compute integral coordinates of each class at a given p without selecting one representative which is integral at every prime.

5.5 Algebraic modular forms and the finite-dimensional model

Let (R, ρ) be a finite-dimensional complex representation of $G(F)$ obtained from an algebraic representation of the archimedean group. A smooth algebraic modular form of weight ρ is a function $\phi : G_f \to R$ such that $\phi(\gamma g) = \rho(\gamma)\phi(g)$ for $\gamma \in \Gamma$, with open right stabilizer. At a fixed level K', we require right invariance under K'. There is a minor variant used throughout the proof: one may require $\phi(gk) = \chi(k)\phi(g)$ for a finite-order character χ of K'. This is the usual character level, and the trivial character gives ordinary level forms. The representation $\rho(\gamma)$ in the source is the Weyl representation evaluated at the conjugate of the chosen embeddings, as in `Space.weightRep` of `Lib/GLn/SchurModule/WeightRepBasic.lean`.

Choose one representative g_i of each class. Evaluation gives an isomorphism between the space of forms and a direct sum of subspaces of R, one for each class. At a trivial-character level this subspace is

$$R^{\Gamma_{g_i}} = \{u : \rho(\gamma)u = u \text{ for all } \gamma \in \Gamma_{g_i}\}.$$

At a character level it is the subspace defined by $\rho(\gamma)u = \chi(g_i^{-1}\gamma g_i)u$. Indeed, extending the value u_i at the representative by equivariance gives $\phi(\gamma g_i k) = \chi(k)\rho(\gamma)u_i$; the asserted relation is precisely what makes this extension well-defined (compare two such expressions for the same element). Conversely all values of a form are obtained in this way. The finite class set and stabilizers therefore give a finite-dimensional space.

Corollary 5.6 (Dimension and free level)**.** *At a free level, the dimension of the space of algebraic modular forms of any fixed coefficient weight and finite-order character is* $|X_{K'}| \dim R$. *For two free levels* $K'' \subset K'$ *(with ordinary invariance) the ratio of the dimensions is* $[K' : K'']$. *More generally every level space has dimension at most* $|X_{K'}| \dim R$.

Proof. At a free level each of the indicated subspaces is all of R. The ratio of the class numbers is the index by Lemma 5.4 applied to the constant function 1 at free levels. The general bound follows by taking dimensions of the subspaces. □

In particular when a single scalar unit z at a prime of the column level generates a cyclic group of order $N-1$ modulo that level, the z-fixed part has dimension $c/(N-1)$ if the enlarged level remains free. This is the class-number reason behind the estimate for the kernel of $1 - z$ (see the ramifying operator in Chapter 14). For several commuting cyclic actions, iterating the operator $1 - z$ loses at most the sum of the dimensions of their individual kernels. Indeed the codimension of an intersection is bounded by the sum of codimensions, and finite-order commuting operators are simultaneously diagonalizable over $\mathbb{C}$. The ramification condition is thus a dimension-controlled way to eliminate the invariant vectors, not a qualitative cuspidality condition on the original finite group.

5.6 The integral Weyl model

We set out the coefficient construction because a polynomial representation over $\mathbb{C}$ does not by itself determine an integral lattice. Let l be a positive integer, and let $\kappa(h)$ for $1 \le h < l$ be the nonnegative integer number of columns of height h of a Young diagram. Write

$$\lambda_i = \sum_{h \ge i} \kappa(h), \quad 1 \le i \le l, \qquad D = \sum_i \lambda_i = \sum_h h\kappa(h), \qquad \lambda_l = 0.$$

Let $X = (X_{ij})$ be an $l \times l$ array of variables. For a subset I of rows of size h let $[I]_h(X)$ be the determinant of the submatrix with rows I (in increasing order) and the first h columns. Put

$$b_{\mathfrak{t}}(X) = \prod_{h=1}^{l-1} \prod_{j=1}^{\kappa(h)} [I_{h,j}]_h(X).$$

These products of minors are homogeneous of total degree D. Under right multiplication of X by an upper triangular matrix T they transform by the character $\prod_i T_{ii}^{\lambda_i}$, since its first h columns transform among themselves by the upper triangular leading block. Left row substitution by a matrix A sends such a product to an integral linear combination of such products by Cauchy–Binet. These two observations explain the representation and the row substitution used in the source.

The standard integral basis theorem for Schur modules gives a basis b_i, consisting of designated products of these minors, of the semi-invariant polynomial module. It exists over $\mathbb{Z}$ and commutes with extension of scalars; its rank is given by the Weyl dimension formula. This standard-basis assertion is the classical input from the theory of straightening laws (compare [9, 10]), in the precise form `PolyRep.exists_minorProduct_basis` in `Lib/GLn/SchurModule/MinorBasis.lean`. We use it here as a formal-source input for the exact chosen integral basis, not as a claim that an arbitrary chosen complex highest-weight basis has integral coordinates. In characteristic zero the resulting module, with the corresponding convention for the row action, is the irreducible polynomial representation of highest weight λ or its dual (the two are distinguished by whether one acts on polynomials or on their evaluations).

For each array X set $\mathrm{ev}_X = (b_i(X))_i$. The vectors ev_X span the coordinate space over any infinite field over which the b_i remain independent: a functional zero on all of them would give a polynomial zero on every array, hence the zero polynomial. Left substitution defines a matrix $S_\kappa(A)$ by

$$S_\kappa(A)\,\mathrm{ev}_X = \mathrm{ev}_{AX}\,. \tag{5.3}$$

It has polynomial entries with integral coefficients and satisfies the representation identity, by the spanning observation and by substitution. Over the integers, the full-rank column determinants are separated as a determinant power, and a rational weight is represented by $\det(A)^e S_\kappa(A)$ for $e \in \mathbb{Z}$. For several infinite places one takes the exterior tensor product of these spaces and the product of the determinant twists; the total degree is the sum of the separate degrees. These are exactly the matrix and weight models called `schur` and `weylRep`.

We now derive the metric in these coordinates rather than appeal to an unspecified invariant pairing. Define the bilinear kernel

$$\mathcal{K}(Y, X) = \prod_{h=1}^{l-1} \det(Y_h^T X_h)^{\kappa(h)},$$

where X_h, Y_h are the first h columns of the two arrays. Cauchy–Binet gives

$$\mathcal{K}(Y, X) = \sum_{\mathfrak{t}} b_{\mathfrak{t}}(Y) b_{\mathfrak{t}}(X) = \mathrm{ev}_Y^T\, G_\kappa\, \mathrm{ev}_X\,.$$

The sum is over all choices of the ordered row subsets in the separate column factors, including repetitions when a height occurs more than once. To see the properties of G_κ write $b_{\mathfrak{t}} = \sum_i c_{\mathfrak{t}i} b_i$ with integral coefficients. Then $G_\kappa = \sum_{\mathfrak{t}} c_{\mathfrak{t}} c_{\mathfrak{t}}^T$, an integral symmetric matrix. Because the products span the polynomial module, these vectors $c_{\mathfrak{t}}$ span in characteristic zero. Hence G_κ is positive definite over $\mathbb{R}$ and defines the Hermitian form $\langle u, v\rangle_\kappa = \bar{u}^T G_\kappa v$ over $\mathbb{C}$. In the case of several places, use the product of these kernels, giving the tensor-product matrix.

The two polynomials $\mathcal{K}(Y, AX)$ and $\mathcal{K}(A^T Y, X)$ coincide. Evaluating and using (5.3) yields the contravariance identity

$$G_\kappa S_\kappa(A) = S_\kappa(A^T)^T G_\kappa. \tag{5.4}$$

For a unitary A, $\bar{A}^T = A^{-1}$ and the representation identity therefore imply invariance of $\langle \cdot, \cdot \rangle_\kappa$. A determinant twist of any integer exponent has modulus one on a unitary element and preserves invariance. This reconstructs the integral Gram form used in `Lib/GLn/SchurModule/Gram.lean`, since symmetries and polynomial expansion uniquely determine the matrix of the kernel in the basis of minor products. At an adelic class the coefficient norm is not an arbitrary Euclidean norm in the selected complex basis: it is this precise Gram form.

5.7 Fischer pairing and the highest-weight hook constant

For polynomials of fixed degree in variables X_{ij} the bilinear Fischer pairing is

$$\mathsf{F}(f, g) = f(\partial) g(0) = \sum_\alpha \alpha! f_\alpha g_\alpha.$$

Its associated Hermitian pairing replaces f_α with $\bar{f}_\alpha$ and is positive definite. The row action by a unitary matrix preserves the Hermitian pairing: on a homogeneous degree-D component one can identify this form with $D!$ times the pairing induced from the D-fold tensor power of the standard Hilbert space, projecting to the symmetric tensors; the monomial basis has squared norms $\alpha!$. Equivalently the chain rule gives the bilinear adjoint of substitution by A as substitution by A^T. This remains true on the semi-invariant polynomial subspace because the subspace is stable under row substitutions.

Write H_κ for the matrix $\left(\mathsf{F}(b_i, b_j)\right)_{ij}$ and let $f_\lambda(X) = \prod_h \det(X_{[1,h],[1,h]})^{\kappa(h)}$. Let $h_\kappa = \mathsf{F}(f_\lambda, f_\lambda)$, a positive integer, the hook constant in the code. It is not in general the plain product of the factorials of the row lengths. Its relation with the Gram matrix is the one needed to turn polynomial differentiation into a statement about coordinate norms.

Lemma 5.7 (The two integral pairings)**.** *One has*

$$G_\kappa H_\kappa = H_\kappa G_\kappa = h_\kappa I. \tag{5.5}$$

In particular for a column vector u the polynomial $P_u = \sum_i (G_\kappa u)_i b_i$ satisfies

$$\mathsf{F}(P_u, b_j) = h_\kappa u_j, \qquad \mathsf{F}(P_u, P_v) = h_\kappa u^T G_\kappa v.$$

Proof. Let $S = S_\kappa(A)$ denote the row action on evaluation vectors. For substitution by A the action on the column vector of coefficients of a polynomial is S^T. The adjoint formula gives $S H_\kappa = H_\kappa S_\kappa(A^T)^T$. By (5.4), it follows that $H_\kappa G_\kappa$ commutes with every $S_\kappa(A)$. In characteristic zero the representation is irreducible, so Schur's lemma says that the product is scalar. (Equivalently, the irreducibility assertion can be replaced by the spanning orbit of the highest vector and the same weight calculation below.) To determine the scalar use $e = \mathrm{ev}_{I_l}$. The kernel $\mathcal{K}(I_l, X)$ equals $f_\lambda(X)$, so $P_e = f_\lambda$. A minor product of highest row weight λ is f_λ itself: the sum of occurrences of the first s rows is at most $\sum_h \kappa(h) \min(s, h)$, with equality for all s only if each column of height h has exactly the rows $1, \dots, h$. Consequently whenever b_i has that weight its evaluation at I_l is 1 and it equals f_λ; all other basis products have zero evaluation at I_l and, being of a different weight, have zero Fischer pairing with f_λ (orthogonality under the diagonal torus, or disjoint monomial weights). Thus $H_\kappa G_\kappa e = h_\kappa e$. As $e \neq 0$, the scalar is h_κ. Symmetry gives both product identities, and the remaining assertions follow by matrix multiplication. □

This proves the identities `fockPairing_toPoly_nablaBasis` and `weylGram_mul_fockPairing` of the same file. In particular inversion of the Gram matrix introduces at most a fixed integral hook denominator: $G_\kappa^{-1} = H_\kappa/h_\kappa$. The exact value often cancels from a ratio, but its size is needed if one works integrally.

For $D \geq 1$ the elementary polynomial expansion gives $h_\kappa \leq D^{3D}$. Indeed the sum of absolute values of the coefficients of f_λ is at most $\prod_h (h!)^{\kappa(h)} \leq D! \leq D^D$; every multi-index α of degree D satisfies $\alpha! \leq D!$. Therefore

$$h_\kappa = \sum_\alpha \alpha! |(f_\lambda)_\alpha|^2 \leq D! \left(\sum_\alpha |(f_\lambda)_\alpha| \right)^2 \leq D^{3D}.$$

At degree zero the constant is 1. For a collection of places the polynomial is a product in disjoint variables, and the same bound applies with total degree. This is `hookNumber_le` in `Lib/GLn/SchurModule/Hook.lean`.

5.8 Examples of the coefficient normalizations

A useful example is the diagram consisting of a columns of height one. Its polynomial basis is $b_\alpha = X_1^{\alpha_1} \cdots X_l^{\alpha_l}$ with $|\alpha| = a$. The kernel is $(Y^T X)^a$, so

$$G_\kappa = \operatorname{diag}\left(\frac{a!}{\alpha!}\right), \qquad H_\kappa = \operatorname{diag}(\alpha!), \qquad h_\kappa = a!.$$

Thus the polynomial lattice and the Weyl evaluation lattice differ: the map $u \mapsto P_u$ multiplies each coordinate by $a!/\alpha!$. Even in this simplest case, replacing the integral evaluation coordinates by coefficients of a polynomial changes the denominators at primes dividing factorials. This is precisely the diagram occurring on the tops at stage X_0. For $l = 2, a = 2$, the three diagonal entries in the order $(2,0), (1,1), (0,2)$ are $1, 2, 1$, and $H_\kappa = \operatorname{diag}(2,1,2)$. Over a field of characteristic different from two both give equivalent integral models; at 2 the distinction is real. The formal proof similarly tracks the gauges under Hecke operators using the actual evaluation lattice, not just a complex vector-space isomorphism to a symmetric power.

For a single column of height h, $b_I = [I]_h$, so $G_\kappa = I$ by Cauchy–Binet, whereas the Fischer pairing matrix is $h!I$, as a determinant expansion has $h!$ distinct monomials with coefficients ± 1. This is the exterior power example. With several columns of varied heights, products of minors satisfy straightening relations: they cannot all be used as independent coordinates. For example in three row variables the two-by-two minors of the first two columns satisfy the ordinary alternating relation

$$X_{11}[23]_2 - X_{21}[13]_2 + X_{31}[12]_2 = 0.$$

It follows by expanding all three determinants. The standard integral basis is needed to remove such relations without inverting integers; otherwise an estimate on an overcomplete tensor model would not imply the desired statement about the actual coefficient lattice.

5.9 The class-sum metric and the real matrix-coefficient model

Returning to the definite adelic space, use the Hermitian form G_κ at the coefficient module of an algebraic form, with the indicated determinant twist. The rational unitary action preserves

it. Thus for two forms of the same fixed level and character,

$$\langle \phi, \psi \rangle_{K'} = \sum_{[g]} \frac{\overline{\phi(g)}^{T} G_\kappa \psi(g)}{|\Gamma_g|} \tag{5.6}$$

is well-defined and positive definite. At every $\sigma \in \mathrm{Aut}(\mathbb{C})$ define instead the conjugate coefficient norm $\|\phi\|_\sigma^2$ by applying σ to the two coefficient vectors before taking the displayed Hermitian square. It is generally wrong to write $\|\phi\|_\sigma^2 = \sigma(\|\phi\|^2)$, since σ need not commute with conjugation. The mixed Gram formula and determinant inequality in Chapter 31 correct this issue.

We give the real comparison of the class-sum model with matrix coefficients. Let $K_\infty = \prod_{\tau \in \Phi} U(l)$ and let ρ_∞ be the irreducible unitary coefficient representation with a fixed highest vector v_+ of norm one. Let d_ρ be its dimension. With probability Haar measure on K_∞, averaging the rank-one projection P_{v_+} over the group gives

$$\int_{K_\infty} \rho_\infty(k) P_{v_+} \rho_\infty(k)^{-1} dk = d_\rho^{-1} I.$$

Indeed the average commutes with the irreducible representation and has trace one; this proves the formula by Schur's lemma. Consequently for any two vectors u, v the integral of the product of the corresponding matrix coefficients with v_+ is $d_\rho^{-1}\langle u, v\rangle$. This is the form of Schur orthogonality needed here.

For ϕ as above define on the adelic group

$$F_\phi(k, g) = \langle v_+, \rho_\infty(k^{-1})\phi(g)\rangle.$$

The left rational invariance follows by equivariance of ϕ and the identification of the algebraic weight with its real counterpart. Right finite-level transformation follows from that of ϕ. Applying Schur orthogonality at each class shows

$$\int_{G(F)\backslash G(\mathbb{A}_F)} \overline{F_\phi} F_\psi = \frac{1}{d_\rho [K_0 : K']} \langle \phi, \psi \rangle_{K'},$$

with the fixed maximal finite level and compact real measures normalized to one. Conversely, a scalar function whose real component lies in this matrix coefficient model determines the coefficient vector by integrating against $d_\rho \rho_\infty(k) v_+$, by the same projection identity. Thus highest-vector matrix coefficients and the coefficient-valued forms are mutually inverse in this model. These calculations explain the factors in `Space.integral_mul_eq_classIntegral` and `Space.algFormEquiv` of `Lib/Adelic/CuspForms/AlgebraicForms.lean`. They also explain why a formal-degree factor is needed when replacing an indefinite discrete-series model by a definite one; in the indefinite case the real group is not compact and the relevant normalization instead uses its formal degree and explicit reproducing coefficient (Chapter 20).

5.10 Transported lattices and gauges

Let p be a rational prime, and let ι be a field isomorphism from $\mathbb{C}$ to the algebraically closed p-adic coefficient field used in the source. In the notation below one can use the completed algebraic closure $\mathbb{C}_p$; the formal model is an algebraic closure of $\mathbb{Q}_p$ with its extended absolute value. Only coefficients and field embeddings, not a comparison of the complex and p-adic topologies, are involved. The valuation ring on the complex coefficient field is

$$O_\iota = \{z : |\iota(z)|_p \le 1\}.$$

The local matrix obtained from g^{-1} at the embedding $\iota\circ\tau\circ\mathrm{conj}$ acts in the weight representation on the corresponding factor. We denote the resulting product matrix by $T_\iota(g^{-1})$. If u is a coefficient vector, the actual gauge is

$$|u|_{\iota,g} = \|T_\iota(g^{-1})\iota(u)\|_{\sup}.$$

This expression is meaningful because the local factor extends the map $E \to \mathbb{C}_p$ continuously at its induced place. Notice that applying ι directly to infinitely many adelic components without choosing a place would be meaningless. The definition is `weylGauge` in `Lib/GLn/ScHurModule/WeightRepBasic.lean`.

Lemma 5.8 (Elementary invariances of the gauge)**.** *The following assertions hold for the standard definite model.*

1. *If $\gamma \in G(F)$, then $|\rho(\gamma)u|_{\iota,\gamma g} = |u|_{\iota,g}$.*
2. *If k is integral and invertible at all the places above p, then $|u|_{\iota,gk} = |u|_{\iota,g}$.*
3. *The gauge is ultrametric and homogeneous: $|u+v|_{\iota,g} \le \max(|u|_{\iota,g}, |v|_{\iota,g})$ and $|cu|_{\iota,g} = |\iota(c)|_p|u|_{\iota,g}$.*
4. *If g is integral invertible over p, then the gauge equals the coefficient sup norm. In particular to prove that an algebraic form is integral it is enough to test one p-integral representative of each class, using Lemma 5.5.*

Proof. The first follows from the representation identity and compatibility of the embeddings: $T_\iota((\gamma g)^{-1})\iota(\rho(\gamma)) = T_\iota(g^{-1})$. For the second, multiplication by $T_\iota(k^{-1})$ on the left preserves the sup norm. Indeed both the matrix and its inverse have integral entries, because the Weyl coefficients are integral polynomials and determinant powers of units are units. Each gives an inequality in one direction. The third is immediate from the linearity of the transported coordinate map. Finally the same observation applies to $T_\iota(g^{-1})$ when g itself is integral. Any value at an arbitrary representative of the class is related by the left rational covariance and the right level covariance, and finite-order scalars have absolute value one, giving the claimed test. □

These are the statements `weylGauge_embFin`, `weylGauge_mul`, `weylGauge_smul`, and `weylGauge_le_one_of_integralAbove` in `Lib/Adelic/CuspForms/ClassReps.lean`. The gauge is always attached to a specified embedding and often to a modified lattice at gd, with d a suitable adele. The latter is used in the treatment of the level prime N. It is not legitimate to use integrality at one representative of each class for every prime without considering that representative's transported lattice.

For a fixed level the integral forms at a fixed ι form a module over O_ι by the third assertion. At a free level this module is a direct sum of transported free coefficient lattices, so it is free. More generally the local constructions may intersect finitely generated modules or restrict to complex subspaces. A module over a valuation ring need not be finitely generated, and that is a reason for using finite-family determinant inequalities in the source rather than automatically speaking of a finite index. The following basic facts illustrate the distinction.

Lemma 5.9 (Finite matrices over a valuation ring)**.** *Let O be a valuation ring in a valued field with multiplicative valuation, and let M be a finite rectangular matrix over O. Then elementary row and column operations in the corresponding general linear groups over O put M in diagonal form with nonzero diagonal entries $a_1, \dots, a_j$ satisfying $a_i \mid a_{i+1}$, followed by zeros. In particular a full-rank finitely generated submodule $L \subseteq O^q$ admits a basis with logarithmic determinant cost $-\log|\det A|$ for a matrix A whose columns form the basis.*

Proof. If the matrix has a nonzero entry, choose one of maximal absolute value (equivalently least valuation), possible because there are finitely many entries. In a valuation ring divisibility is total; thus the pivot divides every entry. Move the pivot to the upper left by permutations and clear its column and row by elementary operations over O. All entries remaining are divisible by the pivot, so induction proves divisibility of successive pivots. The operations are invertible over O, and in the full-rank square-block output this gives the claimed basis and determinant. No uniformizer or discrete value group is needed. □

For a family $y = (y_i)_{i=1}^q$ of forms, integrality after taking combinations by a matrix $U \in M_q(O_\iota)$ says that the coefficient module of integral relations contains the rows of U. If $\det U$ is nonzero, this gives a finite determinant-cost witness even when the full integral relation module has no preferred finite presentation. Determinant costs add under composition: $-\log|\det UV| = -\log|\det U| - \log|\det V|$. The actual definition `Data.IndexLe` uses such witnesses with a common budget at each rational prime for every embedding and allowable twist. The matrix may depend on the embedding and twist; the bound must not. The stronger submodule witnesses used in the local Hecke computations are explained in Chapter 14.

5.11 Finite-place Hecke operators on explicit lattices

Hecke operators at the split places are central to the integral repairs. We recall the elementary double-coset model and its lattice action. Let G_0 be a locally compact group, K compact open, and M a representation over a commutative coefficient ring. For a K-bi-invariant function f of finite support on right K-cosets define on M^K

$$T_f(v) = \sum_{gK} f(g)\rho(g)v.$$

This is independent of the representatives since v is K-fixed. Left K-invariance of f implies the output is K-fixed as well, by reindexing. For two such functions f, f' the identity $T_{f'}T_f = T_{f'*f}$ follows by multiplying the two sums, where the convolution is

$$(f' * f)(h) = \sum_{gK} f'(g)f(g^{-1}h).$$

All equalities make sense over the coefficient ring; no analytic convergence and no division by the volume of a group is involved. Intertwiners of representations carry these sums to the same sums in the target. More generally, if $f(kg) = \chi(k)^{-1}f(g)$, the same reindexing shows that $\rho(k)T_f(v) = \chi(k)T_f(v)$; the inverse on the function is necessary for this convention of the operator. This is the model in `Lib/Adelic/Hecke/HeckeBasic.lean`, especially `heckeFun_eq_sum`, `heckeFun_heckeFun`, and `heckeFun_mem_levelSpace`.

Now take $G_0 = \mathrm{GL}_l(F_v)$ with maximal integral subgroup $K = \mathrm{GL}_l(\mathcal{O}_v)$, residue field k of order Q, and uniformizer ϖ_v. Let $d_j = \mathrm{diag}(\varpi_v I_j, I_{l-j})$. A right coset in Kd_jK/K is a lattice L with

$$\varpi_v\mathcal{O}_v^l \subseteq L \subseteq \mathcal{O}_v^l, \qquad \dim_k(\mathcal{O}_v^l/L) = j.$$

Indeed right multiplication does not change the lattice, and by the diagonal normal form every such lattice lies in the double coset. These lattices are in bijection with the subspaces of codimension j of k^l, or equivalently (after duality) the j-dimensional subspaces. Counting ordered independent j-tuples and dividing by the number of ordered bases in dimension j gives the Gaussian binomial coefficient

$$\#(Kd_jK/K) = \binom{l}{j}_Q = \prod_{a=0}^{j-1} \frac{Q^l - Q^a}{Q^j - Q^a}.$$

For $j = 1$ this is $(Q^l - 1)/(Q - 1)$; in particular in rank two the neighbor graph has $Q + 1$ outgoing edges at an unramified vertex, counted with multiplicity. This count is an exact finite identity, not a local asymptotic.

The split-unitary identification (5.1) transports these operators to unitary groups. In particular inverse transpose switches the role of a sublattice and a scaled superlattice. In the unitary model one must choose the correct factor before translating the gauge; this is the same sign convention that appears in the Frobenius-twisted depth bounds of Chapter 14. If T is a sum of translates whose coefficients are p-integral and all its input gauges at the translated lattices are bounded by A, the ultrametric inequality bounds the output gauge by A. There is no factor counting the number of cosets. By contrast the triangle inequality at an archimedean norm introduces such a factor; this difference is used throughout the proof.

A small but instructive example is the neighbor relation on local rank-two lattices: for the operator T_1 above one has in the unnormalized integral Hecke algebra

$$T_1^2 = T_{(2,0)} + (Q + 1)T_{(1,1)}.$$

Here $T_{(a,b)}$ denotes the indicator of the double coset of $\mathrm{diag}(\varpi_v^a, \varpi_v^b)$. Indeed a chain of two index-Q sublattices between the original lattice and a colength-two lattice has exactly one possible intermediate lattice when its quotient is cyclic of length two, and $Q + 1$ choices when the quotient is a two-dimensional residue vector space. Thus the coefficient of the scalar double coset is genuinely $Q + 1$; normalizing one operator by a power of ϖ_v or conjugating it by an integral weight changes its appearance. In the proof a considerably more involved system of commuting operators is normalized so that its local monomial matrix has entries like p^{depth}. A bound on the complex Hecke polynomial alone would not prove an integral elementary-divisor bound for this matrix.

5.12 Class correspondences and adjoints

The same integral-Hecke operator has a useful adjoint property for the normalized metric. Suppose G_f has a unimodular Haar measure and a representation on a scalar or vector-valued automorphic L^2 space is unitary under right translation. With base level K of volume one, the operator on the fixed space associated with f is $\int_{G_f} f(g)\rho(g)\,dg$, a finite sum when applied to fixed vectors. Its adjoint is associated with $f^\vee(g) = \overline{f(g^{-1})}$ by changing variables $g \mapsto g^{-1}$. This holds for different levels after inserting their volume factors or after restricting each side to the indicated fixed subspaces. In particular for a double-coset indicator the adjoint has the inverse double coset, using probability Haar on the level. If we instead work with the two class-sum metrics at different levels, the quotient-of-index factors in Lemma 5.4 must be retained.

More concretely let $L = K \cap aKa^{-1}$ for a finite adelic a. The two maps from classes at this common smaller level to classes at level K are $[g] \mapsto [g]$ and $[g] \mapsto [ga]$. Their fibers, with stabilizer weights, have relative sizes $[K : L]$ and $[K : a^{-1}La]$ respectively; these are equal by unimodularity. Pullback and finite weighted pushforward are adjoint for the corresponding normalized measure. This is a finite-class explanation of the correspondence description of Hecke operators. It extends to algebraic forms with the appropriate transports of coefficient vectors along the correspondence. In particular if a matrix-entry estimate on one side is established using the coefficient gauge, duality at the other side demands the actual contragredient coefficient structure, rather than an assumption that a chosen complex orthogonal projector has integral entries.

5.13 Where this elementary model stops

There are several boundaries to the argument just given. First, it provides finiteness but not the asymptotic estimates for the mass of the definite genus as the field and ranks grow. Second, the finite-level dimension formula does not identify the multiplicities of the different local representations on an indefinite group. Third, the Hecke double-coset description and the Gram matrix alone do not give the integral theta transfer or the small Selmer denominator estimates of the main argument. The first two issues reappear in the quantitative spanning argument (Chapter 20); the last takes up Chapters 14, 31, and 40. The elementary computations in this chapter isolate the exact places where their more advanced statements enter.

Chapter 6

The splitting character, integral unitary models and the definite mass

The same field, self-duality convention, and finite integral group occur on both sides of the seesaw, in the filling construction, and in the lower-bound periods. A small change to a local character or to the volume of a lattice cannot in general be tolerated by the final comparison. It is useful, therefore, to construct these data once and to distinguish their algebraic consequences from the subsequent analysis of the Weil representation. In this chapter we give the construction of the Weil data of the compact source, including the existence of its special unramified splitting character; the integral transitivity and group counts needed by the comparisons; and a derivation of the quantitative consecutive-mass estimate from the classical Hermitian mass formula. The principal classical global results retained in this chapter are the reciprocity norm theorem of class field theory, Landherr's local–global theorem and the mass formula. None involves a zero-free region of the strength sought in this book.

6.1 Notation and the local norm theorem

Write $E = K(\zeta)$, $K = \mathbb{Q}(\sqrt{-d})$, $F = E^{+}$, $L = N_1$, $f = [F : \mathbb{Q}]$, and $\zeta = \zeta_L$. We use the CM type Φ consisting of the embeddings of E which extend the chosen embedding $\sqrt{-d} \mapsto i\sqrt{d}$. For clarity, v in this chapter is a finite place of F, whereas w is one of the places of E over v; q_v denotes the order of the residue field of F_v. Ordinary additive valuations ord_w are normalized by $\mathrm{ord}_w(\pi_w) = 1$. Thus the source's multiplicative exponential valuation is $\exp(-\mathrm{ord}_w)$.

The extension E/F is unramified at every finite place, not the extension $E/\mathbb{Q}$. To establish this property, observe that i and the primitive third roots of unity belong to E; their orders, 4 and 3, are relatively prime. Suppose that conjugation belonged to an inertia group of $E/\mathbb{Q}$ at residue characteristic p. Select one of the two roots, of order n relatively prime to p. Reduction is injective on the n-torsion of units: if $u^n = 1$ and $u \equiv 1$ modulo the maximal ideal, then $(u-1)(1 + u + \cdots + u^{n-1}) = 0$ and the second factor is a unit. But conjugation acts as inversion on the root selected, which has order greater than two. This contradicts trivial inertia action. The inertia group of E/F is the intersection with $\{1, c\}$, where c is complex conjugation. The contradiction proves the assertion, and the local alternative at v is either

$$E \otimes_F F_v = F_v \times F_v \quad \text{or} \quad E \otimes_F F_v/F_v \text{ unramified quadratic.}$$

The primitive 5040-th roots that appear in the definition `StablyUnram` in `Lib/Adelic/Ade`

`les/CoprimeValuation.lean` also belong to the field. In fact 18480 need not itself contain 5040, but the factors $2^s 3^{\lfloor \log_3 s \rfloor + 5}$ ensure $5040 \mid L$ for the admissible $s = 22 \log_2 m$. We shall use a weaker property than the full root-of-unity condition: the local norm of any unit of F_v is attained by a unit of the quadratic algebra above it.

Lemma 6.1 (Norms and traces of integral elements)**.** *Let R be the ring of integers of a non-Archimedean local field, with finite residue field k of order q. Let $S = R \times R$, with involution exchanging factors, or let S be the integers of its unramified quadratic extension. Denote the involution by $z \mapsto \bar{z}$. Then*

$$\mathrm{Tr}_{S/R}(S) = R, \qquad \mathrm{N}_{S/R}(S^\times) = R^\times.$$

In the unramified field case an element of the base local field is a norm if and only if its valuation is even. In the split case the norm on the full group of units of the quadratic algebra of fields is surjective.

Proof. Both claims are immediate for the product algebra by choosing $(a, 0)$ for the trace and $(a, 1)$ for the norm. In the unramified field case the trace of the degree-two residue-field extension k_2/k is surjective. In characteristic two its being nonzero follows from the existence of $z \notin k$ (then $z^q + z \neq 0$); in odd characteristic the trace of 1 is 2. Lift an element of residue trace one to an element of S and divide it by its unit trace. This proves trace surjectivity over R. The map on residue units is $z \mapsto z^{q+1}$, surjective because $k_2^\times$ is cyclic. Given $a \in R^\times$, choose a lift $z_1 \in S^\times$ of a residue preimage. If $z_j \bar{z}_j \equiv a \pmod{\pi^j}$ and $a/(z_j \bar{z}_j) = 1 + \pi^j b_j$, choose $u_j \in S$ with $u_j + \bar{u}_j \equiv b_j \pmod{\pi}$. Then

$$z_{j+1} = z_j(1 + \pi^j u_j), \qquad z_{j+1}\overline{z_{j+1}} \equiv a \pmod{\pi^{j+1}}.$$

The product converges in the complete ring S to a unit of norm a. A uniformizer of R is also a uniformizer of S, and its norm is its square. Arbitrary elements are powers of that uniformizer times units. This proves the valuation assertion, including at residue characteristic two. □

This lemma is the reason we do not need a separately quoted dyadic norm theorem. In particular each conjugation-invariant finite integral unit idele is the product $z\bar{z}$ for an integral unit idele z. Indeed at a conjugate pair one chooses the first coordinate and then the second, and at a fixed w one uses the lemma. All coordinates are units, so the choices define a restricted product.

6.2 A character extension with exactly the required sign

We first isolate the small but decisive arithmetic condition on roots of unity that makes the desired character possible. If δ is a unit of E, write

$$P(\delta) = \prod_{\tau \in \Phi} \tau(\delta).$$

The right side is the image in $\mathbb{C}$ of $N_{E/K}(\delta)$. The following lemma verifies the hypothesis used in `nonempty_splittingChar`, not just an infinity-type existence condition.

Another small consequence of the choice of K is used in the normalized CM reciprocity argument. Since its discriminant is the negative of a prime, the class number of K is odd. Here is a direct proof in place of invoking the general genus theorem.

Lemma 6.2 (Odd class number of the quadratic base)**.** *The ideal class group of $K = \mathbb{Q}(\sqrt{-d})$ has no nonidentity element of order two. In particular the squaring map on it is bijective.*

Proof. Conjugation acts by inversion on ideal classes, since $\mathfrak{a}\bar{\mathfrak{a}}$ is a principal rational ideal. If the class of $\mathfrak{a}$ has order dividing two, then $\bar{\mathfrak{a}} = \lambda\mathfrak{a}$ for some $\lambda \in K^\times$. Applying conjugation twice gives $(\lambda\bar{\lambda})\mathfrak{a} = \mathfrak{a}$; hence $\lambda\bar{\lambda}$ is a positive rational algebraic unit, therefore 1. If $\lambda \neq -1$, let $u = 1 + \lambda$; then $u/\bar{u} = \lambda$. If $\lambda = -1$ use $u = \sqrt{-d}$. In both cases $\mathfrak{b} = u\mathfrak{a}$ is invariant under conjugation. Factor this fractional ideal into primes. At each split rational prime the exponents of the two conjugate factors are equal, and the paired product is principal. At an inert prime the prime ideal is generated by the rational prime. The only ramified prime lies above d, and it is the principal ideal generated by $\sqrt{-d}$. Thus $\mathfrak{b}$ is principal. Since the class group is finite, the assertion about squaring follows from the structure of a finite abelian group. □

The last assertion is stronger than merely saying that the order of the ramified prime class is odd. It allows one to choose an ideal-class square root of any desired CM ideal class in the normalized reciprocity calculation in Chapter 16.

Lemma 6.3. *For every root of unity $\xi \in E$ there exists a unit $\delta \in \mathcal{O}_E^\times$ with*

$$\delta/\bar{\delta} = \xi, \qquad P(\delta) = 1.$$

Proof. As in Chapter 4, the entire group of roots of unity of E is generated by ζ. The element $\delta_0 = i(\zeta - 1)$ satisfies $\bar{\delta}_0 = \delta_0/\zeta$. The norm of $\zeta - 1$ from $\mathbb{Q}(\zeta)$ to $\mathbb{Q}$ is $\Phi_L(1) = 1$, since L has more than one prime divisor. For completeness, the identity $X^n - 1 = \prod_{a|n} \Phi_a(X)$, divided by $X - 1$ and evaluated at 1, gives

$$n = \prod_{1<a|n} \Phi_a(1).$$

Induction on n shows that $\Phi_{p^e}(1) = p$ and that every other factor is 1. Thus $\zeta - 1$ is a unit. Its norm over K is the same rational number as its cyclotomic norm, since K and $\mathbb{Q}(\zeta)$ are linearly disjoint. The norm of i is 1 too: the cyclotomic automorphisms pair each embedding with its conjugate, and each product of the two conjugates of i is 1. Hence $P(\delta_0) = 1$. For $\xi = \zeta^a$ take $\delta = \delta_0^a$, allowing negative a because δ_0 is a unit. □

Let $J_E = \mathbb{A}_E^\times$, $U_f = \prod_{w<\infty} \mathcal{O}_w^\times$, and $H = E_\infty^\times U_f$, an open subgroup of J_E. Define the continuous unitary character

$$S(x) = \prod_{\tau\in\Phi} \frac{x_\tau}{|x_\tau|} \qquad (x \in H).$$

One would like to prescribe this character on H, require triviality on $E^\times$, and also require $\upsilon(\bar{x}) = \upsilon(x)^{-1}$. These requirements are compatible precisely when S is trivial on

$$H \cap E^\times\{z\bar{z} : z \in J_E\}.$$

Checking this last intersection is important; choosing an arbitrary unramified Hecke character of the right infinity type does not prove the conjugation requirement.

We use the following standard consequence of global class field theory: for the quadratic extension E/F the global Artin character $\eta : J_F \to \{\pm 1\}$ has kernel $F^\times N_{E/F}(J_E)$; its local restriction is the usual norm character, at real places the sign, at unramified inert finite places $(-1)^{\mathrm{ord}_v}$, and at split places trivial. This is the quadratic norm subgroup theorem (e.g. Neukirch, [11], global reciprocity). These assertions apply to the full ideles and do not presume GRH.

Proposition 6.4 (Existence of the normalized splitting character)**.** *There is a continuous character $\upsilon : J_E \to \{z \in \mathbb{C} : |z| = 1\}$ such that*

1. *$\upsilon(a) = 1$ for $a \in E^\times$;*
2. *$\upsilon|_H = S$;*
3. *$\upsilon(\bar{x}) = \upsilon(x)^{-1}$ for $x \in J_E$;*
4. *its restriction along $J_F \hookrightarrow J_E$ is η.*

In particular its value on every finite uniformizer idele supported at a conjugation-fixed place is -1.

Proof. Set $D = E^\times\{z\bar{z} : z \in J_E\}$. Suppose $x = az\bar{z} \in H \cap D$. At every finite w, the equality $\mathrm{ord}_w(x) = 0$ implies

$$\mathrm{ord}_w(a) = -\mathrm{ord}_w(z) - \mathrm{ord}_{\bar{w}}(z),$$

also when $w = \bar{w}$. Thus the fractional ideals of a and $\bar{a}$ coincide, so $\bar{a}/a$ is an algebraic unit. At every complex embedding it has modulus 1. Such a unit is a root of unity: the polynomials of its successive powers are monic integral of bounded degree with bounded coefficients, because their roots have modulus 1. Only finitely many such powers can occur, as each polynomial has finitely many roots. Lemma 6.3 supplies a unit δ with $\delta/\bar{\delta} = \bar{a}/a$ and $P(\delta) = 1$. Then $A = a\delta$ lies in $F^\times$.

The finite idele $Az_f\bar{z}_f = \delta x_f$ is an integral unit and conjugation invariant. By Lemma 6.1 it is a norm of an integral unit idele u_f of E. Consequently A^{-1} is a finite idele norm. Global reciprocity for the diagonal element A implies

$$\prod_{\tau \in \Phi} \mathrm{sgn}(\tau(A)) = 1.$$

The archimedean coordinates of $z\bar{z}$ are positive, and $P(\delta) = 1$ implies $S(\delta) = 1$. Therefore $S(x) = S(a) = S(A) = 1$. This establishes the intersection condition.

On HD define $\upsilon_0(hd) = S(h)$. The intersection condition makes the definition independent of the representation. A homomorphism from a subgroup of an abelian group to the circle extends to the full group. Here is the short algebraic proof of this assertion. Order extensions by inclusion of their domains and take a maximal extension using the union along chains. If g is outside its domain B, either no positive power of g lies in B, in which case its value can be chosen freely, or there is a least positive m with $g^m \in B$. Choose an m-th root of the already prescribed value of g^m in the circle. Defining the values on $B\langle g\rangle$ multiplicatively is well-defined by division with remainder modulo m. Either case contradicts maximality unless every element is in the domain. Apply this to υ_0 to obtain υ on J_E. Since it is continuous on the open subgroup H containing the identity, the extension is continuous everywhere. Triviality on D proves the principal and conjugation assertions.

The restriction to J_F is trivial on $F^\times$ and on idele norms from E. The global norm subgroup has index two, and the restriction is not trivial: on the idele with one real coordinate -1, all others $+1$ and finite part 1, its value is -1. It follows that this restriction equals η. If $w = \bar{w}$, the local extension is unramified quadratic, a uniformizer of F_v is a uniformizer of E_w, and η_v sends it to -1. An arbitrary uniformizer differs from it by a unit, on which υ is trivial. This proves the last assertion. □

Notice that the proof used the full conjugation condition, not merely the identity $\upsilon(a) = 1$ on rational a. The possible choices of the extension are immaterial to its restrictions used here.

The existence of one such choice is exactly what the source's record demands. Its construction is the classical proof by extending a character to a divisible group, with the special compatibility just checked. The source anchor is `Lib/ClassField/HeckeCharacter/SplittingCharExists.lean`; the auxiliary norm decomposition in `Lib/ClassField/Global/CMNorms.lean` follows from the same global Artin character:

Corollary 6.5 (Signs of norms and decomposition of finite ideles)**.** *Let $a \in J_{F,f}$. It can be written as $AN(z_f)$ with $A \in F^\times$. If $a = 1$, the sign vector of A may be any vector of product $+1$, and no sign vector of product -1 is possible.*

Proof. Extend the finite idele a to a full idele of F whose real sign vector has product equal to the value of the finite quadratic Artin character on a. Such a sign vector always exists. The full idele then has trivial Artin character, and global reciprocity writes it as $AN(z)$ for $A \in F^\times$ and $z \in J_E$. Restriction to its finite part gives the desired decomposition. When $a = 1$ take the requested real sign vector of product $+1$; positivity of the real local norms implies that the resulting A has exactly those signs. Necessity follows from reciprocity for A. □

Finite ideal notation. For an unramified character on J_E, trivial on U_f, fix a finite idele y and put $m_w = \operatorname{ord}_w(y_w)$. Since $y(\prod_w \pi_w^{-m_w})$ is an integral unit idele,

$$\upsilon(x_\infty, y) = \left(\prod_{\tau \in \Phi} x_\tau / |x_\tau| \right) \prod_w \upsilon(\pi_w)^{m_w}. \tag{6.1}$$

At a split pair the two finite generator values are inverses; at a fixed place the generator value is -1. These are `upsilon_eq_prod_mul_of_v_eq`, `upsilon_eq_atUnif_zpow` and `upsilon_atUnif_mul_star` in `Lib/General/NumberTheory/UpsilonValues.lean`. There is no choice of a logarithm in (6.1).

6.3 The self-dual lattice and half of the different

Let $p_L(X) = \Phi_L(X)$ and $D_K = p_L'(\zeta)$. These denote the polynomial and a generator of the relative different, not a discriminant. The integral ring of E equals $\mathcal{O}_K[\zeta]$, as established in Chapter 4, and the absolute different $\mathfrak{d}_E$ is the principal ideal $(\sqrt{-d}D_K)$. Define $c_0 = (1 + \sqrt{-d})/2$, $\jmath = -1/\sqrt{-d}$, and the finite idele α with

$$\alpha_w = \begin{cases} D_K^{-1}, & c_0 \in w, \\ 1, & c_0 \notin w. \end{cases}$$

The equality $\operatorname{Im} \tau(\jmath) = 1/\sqrt{d}$ for every $\tau \in \Phi$ fixes the real sign, and $(1 - \jmath^{-1})/2 = c_0$ is integral. In the finite part, the assertion to be verified is

$$\operatorname{ord}_w(\jmath \alpha_w \overline{\alpha}_w) = -\operatorname{ord}_w(\mathfrak{d}_E). \tag{6.2}$$

Here $\overline{\alpha}_w$ means the conjugate of the coordinate at $\bar{w}$, not an automorphism of the w-coordinate if that place is split.

Lemma 6.6. *The idele α satisfies (6.2) at every w. Moreover α^{-1} and $L\alpha$ belong to $\widehat{\mathcal{O}}_E$.*

Proof. First, D_K divides L in $\mathcal{O}_E$: differentiate $X^L - 1 = p_L(X)q(X)$ at ζ, obtaining $L\zeta^{L-1} = D_K q(\zeta)$. Thus the valuations of D_K are supported at primes over divisors of L.

Also $\bar{D}_K$ differs from D_K by a root of unity and sign, by the reciprocal polynomial identity. In particular $\mathrm{ord}_w(D_K) = \mathrm{ord}_{\bar{w}}(D_K)$.

If $p \mid L$, then $p \mid N_0$, and $c_0\bar{c}_0 = (d+1)/4$ is divisible by p (also for $p = 2$, since $d \equiv 7 \pmod 8$). Since $c_0 + \bar{c}_0 = 1$, at each w over p exactly one of these elements lies in the maximal ideal. Conjugation exchanges the two alternatives. It follows that

$$\mathrm{ord}_w(\alpha\bar{\alpha}) = -\,\mathrm{ord}_w(D_K)$$

at such a place. At every other place both sides are zero. Multiplying by $\jmath$ gives (6.2) using the product formula for the different in the tower. The final two assertions are immediate from the definition and $D_K \mid L$. □

For reference the source calls α `Parameters.a`, and the preceding lemma is `v_alpha_mul_relDifferentGen` together with `inv_a_mem` and the divisibility by N_1. The construction of $\jmath$, α and υ has now established the full record `Parameters.instWeilDataE`, using global reciprocity as indicated.

What self-duality means. For a finite local field E_w, let ψ_w be the rational additive character composed with $\mathrm{Tr}_{E_w/\mathbb{Q}_p}$. Its annihilator on $\mathcal{O}_w$ is $\mathfrak{d}^{-1}_{E_w/\mathbb{Q}_p}$. Indeed the character is trivial on all $\psi_w(za)$, $a \in \mathcal{O}_w$, exactly when $\mathrm{Tr}(z\mathcal{O}_w) \subset \mathbb{Z}_p$, which is the definition of the inverse different. Therefore the bilinear annihilator of $\widehat{\mathcal{O}}_E$ for

$$(x, y) \longmapsto \psi_f(\jmath\alpha\bar{\alpha}x\bar{y})$$

is precisely $\widehat{\mathcal{O}}_E$, by (6.2). In the split case the annihilator condition at a pair is tested by varying the components of y independently, so the reasoning applies to the whole restricted product. With Haar measure giving the integral finite adelic lattice volume one, the Fourier transform of its indicator for this pairing is again that indicator. To see this, if x lies in the annihilator the character is identically one on the lattice; otherwise translation by an element on which the character is nontrivial multiplies the integral by a scalar different from one, forcing it to vanish. The square of Fourier transform is reflection, so this is the self-dual measure for the scaled pairing.

The corresponding real scale is

$$s_\tau = (2\pi \operatorname{Im}\tau(\jmath))^{-1/2}, \qquad s_f = \alpha.$$

This is `WeilData.scaleIdele`. At a complex place the alternating form $\mathrm{Tr}_{\mathbb{C}/\mathbb{R}}(\jmath z\bar{w})$ is $-2\operatorname{Im}\jmath\,\operatorname{Im}(z\bar{w})$. Under the scaling $z \mapsto s_\tau z$, its additive-character exponent becomes the standard symplectic exponent. This calculation explains why the factor 2π is present rather than a separate adjustable scalar. The finite lattice Fourier identity just proved does not establish by itself a splitting of the metaplectic representation or a global theta identity; those are treated separately in Chapter 16.

6.4 Reduction of finite ideals and their adelic generators

We record one auxiliary construction used repeatedly when a level or denominator ideal is inserted; it is `exists_reduction_and_idele_generator` in `Lib/General/NumberTheory/UpsilonValues.lean`. For a nonzero integral ideal $I \subset \mathcal{O}_E$, let

$$e_w = \mathrm{ord}_w(I), \qquad t_w = \begin{cases} \pi_w^{e_w}, & e_w > 0, \\ 1, & e_w = 0. \end{cases}$$

Then t is an integral finite idele. The natural map from $\mathcal{O}_E/I$ to $\prod_{w|I} \mathcal{O}_w/\pi_w^{e_w}\mathcal{O}_w$ is an isomorphism: apply the Chinese remainder theorem and note that completion does not change a quotient by a finite power of the maximal ideal. Composing its inverse with the projections defines

$$\rho_I : \widehat{\mathcal{O}}_E \longrightarrow \mathcal{O}_E/I, \qquad \ker \rho_I = I\widehat{\mathcal{O}}_E = t\widehat{\mathcal{O}}_E.$$

It restricts to ordinary reduction of algebraic integers. For any finite adele y, one has

$$(\forall c \in I,\ cy \in \widehat{\mathcal{O}}_E) \quad \Longleftrightarrow \quad ty \in \widehat{\mathcal{O}}_E.$$

One implication uses $c_w \in t_w\mathcal{O}_w$ for every c. For the other, at each w choose $c \in I$ with valuation exactly e_w; it exists because $I\mathcal{O}_w = \pi_w^{e_w}\mathcal{O}_w$ and the localization of I is dense in this ideal. The hypothesized integrality of cy forces $\mathrm{ord}_w(y) \geq -e_w$. Since this works at all w, ty is integral. This proof uses I as a nonzero ideal, not a chosen principal generator in the number field. This distinction matters in the unitary class sums, where the global class need not be principal.

6.5 Completions of the coefficient primes and the choice of a side

Here we give a direct proof of the decomposition-group and chosen-place statements of `New/Data/WeilDataOfE.lean` that are used to label the finite factors of the groups. The Galois group over $\mathbb{Q}$ can be written as

$$(\mathbb{Z}/L)^\times \times \{\pm 1\}, \qquad (a,\epsilon):\ \sqrt{-d} \mapsto \epsilon\sqrt{-d}, \quad \zeta \mapsto \zeta^{\epsilon a}.$$

It is important to include the factor ϵ in the second displayed action. It arises because the first coordinate refers to the CM type.

Proposition 6.7. *For $p \nmid Ld$, put $\epsilon = \chi_K(p)$ and $a = \epsilon p \pmod L$. Then the decomposition group of every place above p in $E/\mathbb{Q}$ is cyclic generated by (a,ϵ). Its order is the residue degree. If $\epsilon = -1$, the elements of $\mathrm{Gal}(E/K)$ which fix a prescribed place are the even powers of $-p$; those mapping it to its conjugate are the odd powers of $-p$.*

Proof. The prime is unramified in $E/\mathbb{Q}$ because this extension is contained in the cyclotomic extension of conductor Ld. Arithmetic Frobenius acts as p-th power on reduction of ζ, and as the Legendre-symbol action on $\sqrt{-d}$. The reduction of L-torsion is injective, and reduction of $\sqrt{-d}$ is nonzero. Consequently the congruence characterizes the stated Galois element. The decomposition group of an unramified prime is the cyclic group generated by Frobenius; the reduction map identifies it with the Galois group of the residue fields. An element of $\mathrm{Gal}(E/K)$ is $(b,1)$. It fixes a place if it is in the decomposition group, and maps the place to its conjugate if multiplying by $(1,-1)$ belongs to the decomposition group. For the inert case these conditions give respectively the even and odd powers. For example, when the order of $-p$ is odd, an odd power can equal the identity in $(\mathbb{Z}/L)^\times$, and the place can be its own conjugate. No set of chosen representatives should identify this situation with a split prime. □

For $p \equiv 1 \pmod{Ld}$ decomposition groups are trivial and all residues have cardinal p. Suppose $b \in \mathcal{O}_E$ and $b\bar{b} = pc$ with $p \nmid c$. At a place $w \mid p$, the valuations satisfy

$$\mathrm{ord}_w(b) + \mathrm{ord}_w(\bar{b}) = 1.$$

Both are nonnegative integers, and conjugation exchanges the two valuations. Thus exactly one place in each conjugate pair divides b; its valuation of b is one. The unconjugated set of places in a Galois orbit under $\mathrm{Gal}(E/K)$ contains one place in each conjugate pair, though not necessarily the place dividing b. Starting from one place and for each $g \in \mathrm{Gal}(E/K)$ selecting gw or $\overline{gw}$ according to divisibility by b therefore gives a bijection

$$\Phi \simeq \{w \mid p : b \in w\}.$$

The domain has f elements, and injectivity follows from the trivial decomposition group and the fact that no element of $\mathrm{Gal}(E/K)$ equals conjugation. This proves the assertions `star_mem_iff`, `intValuation_eq_of_mul_star` and `nonempty_infinitePlace_equiv_finPlace`. In the case $p = g_i$, b is the element π_g (or its conjugate, according to the chosen side), and the cofactor is g_{1-i}. In the case $p = N$ the cofactor is one. This argument is the precise reason reduction at the selected places can be modeled by independent general linear factors rather than by a quadratic unitary group.

6.6 Unimodular lattices over split and unramified places

The local integral group calculation begins with a simple form of Gram–Schmidt. We use R, S as in Lemma 6.1. A Hermitian form on an S-module M is conjugate-linear in the first argument and S-linear in the second. It is unimodular if its Gram matrix in an S-basis is invertible over S.

Proposition 6.8. *Every free unimodular Hermitian S-lattice of rank n admits an orthonormal basis. If $v_1, \dots, v_l$ are already orthonormal vectors, they extend to one. Consequently the unitary group of the standard unimodular lattice acts transitively on orthonormal ordered l-frames over S.*

Proof. In the unramified field case, reduce the Gram matrix to the residue field k_2. A nondegenerate Hermitian form over k_2 has a vector with nonzero norm. To prove this, if every norm is zero, then for every x, y, b,

$$h(x + by, x + by) = \mathrm{Tr}_{k_2/k}(bh(x, y)) = 0.$$

The trace duality on the finite separable field extension is nondegenerate; indeed for $h(x, y) \neq 0$ multiply its inverse by a trace-nonzero element. It follows that the form is zero, a contradiction. Lift a vector of nonzero norm and use norm surjectivity on units to rescale it to norm one. Its projection $x \mapsto vh(v, x)$ splits an orthogonal summand, and the orthogonal complement is a finite free unimodular lattice since it is a projective module over a local principal ideal domain. Induct on rank. The same projection formula applied to the given orthonormal vectors proves the extension assertion.

In the split case, write $M = M_1 \times M_2$, where both M_i are free R-modules of rank n. A Hermitian form is precisely a perfect pairing $M_2 \times M_1 \to R$ together with its transpose. Choose a basis of M_1 and its dual basis of M_2; this gives an orthonormal basis of M. Given an orthonormal sequence, the corresponding projections again split both modules, and the complementary pairing is perfect. The group sends one chosen orthonormal basis to the other, proving transitivity. □

This proof of integral Witt extension applies at dyadic places too: the trace calculation rather than division by two is its essential step. Over a field the usual Witt extension theorem gives transitivity on norm-one vectors in a fixed nonsingular Hermitian space. In the definite

global standard space we can also see it directly. If u, v have norm one and $u \neq v$, then $w = u - v$ has nonzero norm because the form is positive at infinity. The transformation $T(x) = x - wh(w,x)/h(w,u)$ sends u to v and preserves the form. Indeed $h(w,u) \neq 0$ (otherwise $h(v,u) = 1$ and $h(w,w) = 0$), and

$$h(w,w) = h(w,u) + \overline{h(w,u)}.$$

Expansion of $h(Tx, Ty)$ gives the original pairing. Thus T is invertible and unitary. The identity transformation treats $u = v$. This elementary reflection avoids needing a more general version of Witt's theorem for the norm-one mass calculation below.

6.7 Existence and equivalence of the framed global spaces

The group used at the finite places has the identity Gram matrix, while at the real places the same group need not be compact. We now show that these local conventions can be imposed simultaneously by an actual Hermitian space over E. In the source a `Space` with sign pattern s is given by a Hermitian matrix J over E and a matrix $x \in \mathrm{GL}_n(\mathbb{A}_E)$ such that

$$x^* J x = J_s, qquad (J_s)_\tau = \mathrm{diag}(s_1, \dots, s_n), \quad (J_s)_f = I_n.$$

Here $s_i \in \{\pm 1\}$ and the same pattern is used at every real place of F; the notation $*$ means the transpose of conjugation, evaluated on the appropriately paired local factors. The rational unitary group acts in these coordinates by

$$\gamma \longmapsto x^{-1} \gamma x.$$

In particular a rational matrix need not be a rational point of the group whose finite coordinates have the identity matrix. The conjugating matrix x is part of the data; it determines the integral lattices and levels.

Proposition 6.9 (Existence of the normalized signed space)**.** *For every sign pattern s of length n there exists such a pair (J, x). More generally, for each coordinate one can prescribe a sign at each real place provided the product of its signs is positive.*

Proof. In the present field $f = \varphi(L)$ is even. Consequently a sign vector that is constantly -1 has product $+1$; so does the constant positive vector. For each i apply Corollary 6.5 with $a = 1$ and requested signs s_i (or the stipulated varying signs). It gives $a_i \in F^\times$ with these signs and $a_i N(z_{i,f}) = 1$. Put

$$J = \mathrm{diag}(a_i), \qquad x_f = \mathrm{diag}(z_{i,f}), \qquad x_\tau = \mathrm{diag}(|\tau(a_i)|^{-1/2}).$$

These matrices satisfy the required equality. The finite entries $z_{i,f}$ are ideles, not merely independently chosen nonzero elements with uncontrolled denominators. Thus x is an element of the restricted adelic general linear group. This gives the sought global realization and proves the additional sign assertion. □

For equivalence we invoke the classical Landherr theorem over a number field: nonsingular Hermitian forms for a quadratic extension are isometric if and only if they are isometric at all places of the base field. This is a Hasse principle for all signatures, not a strong approximation theorem for a definite group. A reference is Landherr, [13], and Kneser's lectures on Galois cohomology, Section 5.5. The hypothesis of this theorem is especially transparent here: the adelic frames explicitly exhibit the local isometries, including at the places over 2.

Proposition 6.10 (Equivalence with movement of the adelic frame)**.** *Suppose (J, x) and (J', x') realize the same sign pattern. There exist $a \in \mathrm{GL}_n(E)$ and $g \in G_s(\mathbb{A}_F)$, the common adelic group of J_s, with*

$$a^* J a = J', \qquad x' = a^{-1} x g.$$

If γ belongs to the unitary group of J', then

$$x^{-1}(a\gamma a^{-1})x = g\,(x'^{-1}\gamma x')\,g^{-1}.$$

Proof. At every place the matrix $x(x')^{-1}$ is a local isometry from J' to J. By the classical Landherr theorem these local isometries have a rational isometry a (not necessarily approximating the given local matrices). Put $g = x^{-1}ax'$. Then $g^* J_s g = (x')^* a^* J a x' = J_s$, so g lies in the prescribed adelic group. Rearranging its definition proves the second equality, and substituting into $x'^{-1}\gamma x'$ gives the conjugation formula. □

This is the content of `Space.landherr` and `Space.exists_eq_move` in `Lib/Adelic/Landherr/Landherr.lean`. The special signed realization does not require the full theorem; it follows directly from the finite norm subgroup. Equivalence is a standard classical application. The quantitative comparison of the actual frames in the nontrivial theta kernel is a stronger assertion: the theorem only gives a rational isometry and imposes no bound on its archimedean entries or on the chosen lattice denominator. In particular this equivalence cannot by itself complete a quantitative upper bound or transfer an integral algebraic form without a finite-lattice calculation. Later comparisons shall keep the chosen x and g and specify those additional bounds.

6.8 Counting the local finite groups and their congruence quotients

The standard finite integral unitary group is hyperspecial in our unramified setting; what we need is the following explicit description, which can be proved without that terminology. Let $\mathcal{U}_n(S) = \{g \in \mathrm{GL}_n(S) : \bar{g}^t g = 1\}$ and π be a uniformizer of R. The reduction maps

$$\mathcal{U}_n(S) \longrightarrow \mathcal{U}_n(S/\pi^e S)$$

are surjective. We first show this directly in the unramified case. If g_j is a unitary lift modulo π^j, put $C = (\bar{g}_j^t g_j - I)/\pi^j$ modulo π. It is Hermitian. Choose X modulo π with $X + \bar{X}^t = -C$: for $i < l$, set $X_{il} = -C_{il}$ and $X_{li} = 0$; on the diagonal choose an element of residue trace $-C_{ii}$. Then $g_{j+1} = g_j(1 + \pi^j X)$ satisfies the equation modulo π^{j+1}. These lifts converge. The solution space for X in one step is an affine translate of the skew-Hermitian matrices. Over k this space has dimension n^2: each of $n(n-1)/2$ off-diagonal pairs contributes two dimensions, and each diagonal contributes one. Hence each fiber of reduction by one exponent has cardinal q^{n^2}. In the split case the group identifies with $\mathrm{GL}_n(R)$ via the pair $(g, (g^t)^{-1})$. Reduction and these fiber sizes are then immediate.

Lemma 6.11. *Over the residue field k of cardinal q, the split and nonsplit group orders are respectively*

$$|\,\mathrm{GL}_n(k)| = q^{n^2} \prod_{i=1}^{n} (1 - q^{-i}), \tag{6.3}$$

$$|\mathcal{U}_n(k_2)| = q^{n^2} \prod_{i=1}^{n} (1 - (-1)^i q^{-i}). \tag{6.4}$$

At exponent $e \geq 1$ multiply these orders by $q^{(e-1)n^2}$.

Proof. For (6.3), count successive basis vectors: $q^n - q^j$ choices for vector $j+1$. In the nonsplit case we count the vectors of norm one. Fix a nontrivial additive character $\psi : k \to \mathbb{C}^\times$. For $a \neq 0$,

$$\sum_{z \in k_2} \psi(az\bar{z}) = 1 + (q+1) \sum_{b \in k^\times} \psi(ab) = -q.$$

By additive orthogonality, the number of vectors $z \in k_2^n$ of norm one is

$$\frac{1}{q} \sum_{a \in k} \psi(-a) \left(\sum_{z \in k_2} \psi(az\bar{z}) \right)^n = q^{2n-1} - (-1)^n q^{n-1}.$$

Every such vector extends to an orthonormal basis by the residue-field part of the proof of Proposition 6.8; its stabilizer is $\mathcal{U}_{n-1}(k_2)$. Thus the order is the displayed count times the order in rank $n-1$. Starting with rank zero gives (6.4). The fiber calculation above proves the general reduction-order assertion. □

At each chosen $w \mid g_i$, the other conjugate place gives the inverse transpose, so the finite quotient K_0/K_g for rank n is

$$\prod_{i=0,1} \prod_{w \text{ chosen} \mid g_i} \mathrm{GL}_n(\mathbb{F}_{g_i}).$$

Its order is at most g^{fn^2}, using f places above each g_i and (6.3). The block inclusion of a group of rank n in rank rt proves divisibility into `Data.chosenGroupOrder` whenever $n \leq rt$. This verifies both the size and divisibility ingredients of `New/Data/ChosenPlaceLevels.lean`; the proof that stabilizers inject into this quotient is the neatness argument in Chapter 5. We stress that using only an upper bound for the stabilizer order is insufficient for denominator clearing: divisibility of its order into the common integer is what is required.

6.9 An exact norm-one class-sum identity

Before applying a mass formula it is worth understanding why consecutive ranks appear naturally. Let $G_n = U(I_n)$, viewed as a group over F, and $K_n \subset G_n(\mathbb{A}_{F,f})$ the stabilizer of $\widehat{\mathcal{O}}_E^n$. Denote the genus mass by

$$\mathfrak{m}_n = \sum_{[g] \in G_n(F) \backslash G_n(\mathbb{A}_{F,f}) / K_n} \frac{1}{|\Gamma_g|}, \qquad \Gamma_g = G_n(F) \cap g K_n g^{-1}.$$

The finiteness and class-sum convention are those of Chapter 5. At rank zero put $\mathfrak{m}_0 = 1$; in positive ranks the sum is positive.

Proposition 6.12 (The norm-one identity)**.** *Let $n \geq 1$ and let $c(g)$ count $y \in E^n$ with $h(y,y) = 1$ and $g^{-1}y \in \widehat{\mathcal{O}}_E^n$. Then*

$$\sum_{[g]} \frac{c(g)}{|\Gamma_g|} = \mathfrak{m}_{n-1}. \tag{6.5}$$

Proof. The set counted by $c(g)$ is finite: the finite integrality condition defines a lattice in the real product E_∞^n, and the norm condition imposes a compact real subset. Consider pairs (gK_n, y) with the stated conditions modulo the diagonal action of $G_n(F)$. The left hand side of (6.5) is the sum over orbits of reciprocal stabilizer orders. This follows from the finite orbit-stabilizer formula: for fixed g, $\sum_{O \in \Gamma_g \backslash X_g} |\mathrm{Stab}(O)|^{-1} = |X_g|/|\Gamma_g|$.

Every rational norm-one vector can be sent to e_n by a rational unitary transformation, by the reflection in Proposition 6.8. Its stabilizer is the naturally embedded $G_{n-1}(F)$. For a finite adelic element g with $g^{-1}e_n$ integral, Proposition 6.8 at every finite place gives $k \in K_n$ with $ke_n = g^{-1}e_n$. Then gk fixes e_n, so belongs to $G_{n-1}(\mathbb{A}_{F,f})$. All such pairs are obtained in this way. Two representatives with vector e_n lie in the same rational orbit exactly when their finite adelic G_{n-1} cosets lie in the same double coset modulo $G_{n-1}(F)$ and K_{n-1}, because any rational map between them must fix e_n and $K_n \cap G_{n-1}(\mathbb{A}_{F,f}) = K_{n-1}$. Stabilizer groups in that description agree. Thus the right side of (6.5) counts exactly the same orbits with the same weights. □

This gives `classSum_card_norm_one_eq_mass` in `Lib/Adelic/Transference/MassBasic.lean`. It is not merely a comparison between two class numbers. A mass has a reciprocal automorphism weight, and the exact identity uses that weight even in the presence of a non-free level. At a level with trivial stabilizer these weights disappear; this useful special case does not justify replacing them at maximal level.

6.10 The classical mass formula in the present normalization

There are two routes to the mass estimate. The compact code proves it by combining (6.5) with the one-vector Siegel–Weil formula and a uniform Fourier estimate. For a classical exposition we can instead use the standard exact Hermitian mass formula directly, since the lattice is unimodular and E/F is unramified at all finite places. This bypasses the Fourier estimate for this particular purpose only; it does not remove the separate Siegel–Weil period identities used in the lower bound.

We state carefully the specialization being used. For a CM quadratic extension E/F with relative discriminant ideal 1, put $D_F = |\operatorname{disc}(F)|$ and let η be the associated quadratic character of F. For the definite unimodular lattice of positive rank n the formula is

$$\mathfrak{m}_n = 2D_F^{n^2/2} \left(\prod_{j=1}^{n} \frac{(j-1)!}{(2\pi)^j} \right)^f \prod_{j=1}^{n} L_F(j, \eta^j). \tag{6.6}$$

In the product η^j is trivial for even j; $L_F(1, \eta)$ is understood as the usual nontrivial quadratic value. This is Proposition 4.2.7(3) of Kirschmer, [12], with the local factors of Lemma 4.2.9 there all equal to 1 (every prime is good in the sense of Definition 3.5.1, since there is no ramified quadratic extension at a dyadic place): at a split or inert unramified prime the entire lattice is the single unimodular Jordan block ($m_1 = 0$). The archimedean factor in that reference is $\gamma_n^{-1} = \prod_{j=1}^{n}(j-1)!/(2\pi)^j$. The mass is the sum of reciprocal orders of the full unitary automorphism groups, not of the special unitary groups, and the leading factor 2 is the one for this convention. In particular the two groups do not differ by a harmless constant if the formula is used at rank zero.

One can check the Euler factors local by local. At an unramified split place the volume-normalized reduction order of Lemma 6.11 is $\prod_{j=1}^{n}(1 - q_v^{-j})$. At an inert place it is $\prod_{j=1}^{n}(1 - (-1)^j q_v^{-j})$. Their formal reciprocals are the Euler factors in (6.6); the exceptional first factor is the nontrivial character value, as required by the Siegel formula rather than an unregularized absolute product at 1. This confirms the convention. The classical formula, not just these finite counts, gives the equality of the global mass with the completed product; it is the explicit external theorem in this subsection.

Theorem 6.13 (Consecutive-mass estimate)**.** *Let* $r_E = |\operatorname{disc}(E)|^{1/(2f)}$*. If* $8f \le 2^u$ *and* $j \ge 4u + 12$*, then*

$$\left|\log \mathfrak{m}_{j+1} - \log \mathfrak{m}_j - f\left((j+\tfrac{1}{2})\log r_E + \log j! - (j+1)\log(2\pi)\right)\right| \le 2\,2^{-u}. \tag{6.7}$$

Proof. The discriminant tower formula gives $|\operatorname{disc}(E)| = D_F^2$, hence $D_F = r_E^f$. Taking the quotient of (6.6) at $j+1$ and j cancels all terms with index at most j, including the delicate $L_F(1,\eta)$ term. With $n = j+1$ the exact quotient is

$$\frac{\mathfrak{m}_{j+1}}{\mathfrak{m}_j} = \left(r_E^{j+1/2}\frac{j!}{(2\pi)^{j+1}}\right)^f L_F(n,\eta^n).$$

For $n \ge 3$, the Euler factors are positive (including when $\eta^n(v) = -1$), so the logarithm is real and absolutely convergent. For the relative unramified character η,

$$\begin{aligned}|\log L_F(n,\eta^n)| &\le \sum_{v<\infty}\sum_{h\ge 1}\frac{q_v^{-nh}}{h}\\ &\le f\log\zeta(n) \le f(\zeta(n)-1) \le 2f\,2^{-n}.\end{aligned}$$

The middle bound follows by grouping places over each rational prime: there are at most f places and $q_v \ge p$. For the last inequality use $\sum_{a\ge 2} a^{-n} \le 2^{-n} + \int_2^\infty t^{-n}dt = 2^{-n}(1 + 2/(n-1)) \le 2\,2^{-n}$. Finally $2f2^{-n} \le 2^u 2^{-n}/4 \le 2\,2^{-u}$ for the stated bound on n. Substituting the exact quotient proves (6.7). □

The conclusion is exactly `abs_log_mass_ratio_sub_le` in `Lib/SiegelWeil/Mass/MassRatio.lean`. This independent route shows why the exceptionally large rank in the source makes the mass correction uniform even though f is growing. The small factor is an L-value at a large positive integer, not a value near the critical strip. No assertion about zeros of the Dirichlet L-functions under study has entered. The classical mass formula is cited rather than re-proved in this volume; the derivation of the required uniform inequality, and its justification for the exact lattice and conventions of the proof, are given here.

6.11 Useful consequences and the separation of normalizations

The exact formula supplies convenient telescoping identities. For integers $r \ge 1$, $s \ge 0$ such that $r \ge 4u + 12$ and $8f \le 2^u$, one obtains

$$\log\frac{\mathfrak{m}_{r+s}}{\mathfrak{m}_r} = f\sum_{j=r}^{r+s-1}\left((j+\tfrac{1}{2})\log r_E + \log j! - (j+1)\log(2\pi)\right) + \mathcal{R}, \tag{6.8}$$

$$|\mathcal{R}| \le 4f\,2^{-r-1}. \tag{6.9}$$

The latter, stronger than adding $2/2^u$ at each step, follows by summing the geometric bound $2f\,2^{-(j+1)}$ established in the proof. An empty sum has error zero. In particular the apparent growth of the number of consecutive ratios costs no extra factor of s. The identities

$$\sum_{j=r}^{r+s-1}(j+\tfrac{1}{2}) = rs + \tfrac{s^2}{2}, \qquad \sum_{j=r}^{r+s-1}(j+1) = rs + \tfrac{s(s+1)}{2}$$

are often useful when comparing dimensions of successive groups. They have the same weights that appear in the common leading terms in the final assembly.

We also explain an elementary normalization check relating the finite class count to a real coefficient space. If $K' \subset K_n$ is an open subgroup, the total mass at K' is $[K_n : K']\mathfrak{m}_n$: indeed over a K_n-class the fiber groupoid is the action of Γ_g on the $[K_n : K']$ cosets. By orbit-stabilizer its mass is $[K_n : K']/|\Gamma_g|$. If every stabilizer at K' is trivial this mass is the number of K'-classes. A coefficient representation of dimension d then supplies precisely d independent coordinates per class. Thus the scalar space of definite algebraic forms has dimension $d[K_n : K']\mathfrak{m}_n$. This argument does not show that the automorphic form space for an indefinite real group has the same dimension; that requires the transfer or filling argument. In particular one cannot replace a trace formula by this formula for the definite group without supplying the transfer.

This distinction also clarifies three different uses of measures:

1. The product of finite additive Haar measures used for the integral Weil lattice, normalized by self-duality as in (6.2).

2. The finite unitary quotient measure, for which the maximal integral unitary group has volume one and the total volume at a definite real place is the genus mass.

3. The real Haar and formal-degree factors, whose exact normalizations are required for the discrete-series projector and the periods. These are not determined solely by the first two choices.

The class sum comparison and local unitary orders establish the elementary links between the first two structures. The third, and its compatibility with the chosen theta models, is the substance of later chapters. Even given the mass formula, an arbitrary convention for the archimedean formal degree would change the quantitative comparison, so that the additional local calculation cannot be omitted.

Concordance. The assertions now reconstructed in this chapter from the named classical references are the Weil record `Parameters.instWeilDataE`; the conjugation-compatible `nonempty_splittingChar`; the finite ideal and uniformizer relations of `UpsilonValues`; the decomposition and chosen-place calculations of `WeilDataOfE`; the integral orthonormal frame and finite group calculations used by the level; the norm-one class sum; and the mass ratio. The norm-one proof and the finite matrix arguments are independent of the mass formula. A full derivation of the Siegel–Weil formula, of the global Weil representation and of the specialized uniform theta-period identities is not asserted here. Those are different obligations, even when their objects have now been normalized by an independent construction.

Chapter 7

Heisenberg operators and the finite vacuum multiplier

This chapter supplies common background for the definite and indefinite theta models. In particular the finite multiplier recording the change of a vacuum lattice is not a free scalar; its sign and its absolute value occur in the upper-bound and period calculations. We give a direct lattice-theoretic computation of that multiplier and explain its relation to the Heisenberg model. We prove that the phase-normalized finite intertwiners form a genuine representation by combining a finite Gauss sum with a sign argument modulo four; an absolute-value calculation alone could not establish this assertion.

7.1 The saturating lattice at a finite place

Put $\widehat{\mathcal{O}} = \widehat{\mathcal{O}}_E$, and let $b \in \mathrm{GL}_n(\mathbb{A}_{E,f})$. The lattice

$$L_b = \widehat{\mathcal{O}}^{\,n} \cap b^{-1}\widehat{\mathcal{O}}^{\,n}$$

is compact open in the additive finite adeles, and of finite index in $\widehat{\mathcal{O}}^{\,n}$. It is the intersection of the old lattice with the inverse image of the new one. Its determinant ideal is the quantity that controls the finite part of the vacuum multiplier.

Lemma 7.1 (Saturating lattice matrices)**.** *There exist $u \in \mathrm{GL}_n(\mathbb{A}_{E,f})$ and $l, m \in M_n(\widehat{\mathcal{O}})$ such that*

$$u, bu \in M_n(\widehat{\mathcal{O}}), \qquad u^{-1} = l + mb.$$

Every such u satisfies $u\widehat{\mathcal{O}}^{\,n} = L_b$. At each finite place w, if b has elementary divisors $\pi_w^{a_1}, \dots, \pi_w^{a_n}$ relative to the standard lattice, then

$$\mathrm{ord}_w(\det u) = s_w(b) := \sum_{i=1}^{n} \max(0, -a_i). \tag{7.1}$$

Proof. At a finite place write $b = kDk'$ by the elementary divisor theorem over the discrete valuation ring, where k, k' are integral invertible matrices and $D = \mathrm{diag}(\pi^{a_i})$. Set $P = \mathrm{diag}(\pi^{\max(0,-a_i)})$ and $u = (k')^{-1}P$. Then u and $bu = kDP$ are integral. To make the claimed inverse identity explicit, let A, B be the diagonal matrices with $A_{ii} = 1$ when $a_i \geq 0$ and 0 otherwise, and $B = I - A$. Since $P^{-1} = A + BD$, one has

$$u^{-1} = P^{-1}k' = Ak' + Bk^{-1}b.$$

Thus take $l = Ak'$ and $m = Bk^{-1}$. At every place where b is integral invertible we may instead take $u = l = I$ and $m = 0$. The choices are therefore finite-adelic. If $x \in \widehat{\mathcal{O}}^n$ with bx integral, the inverse identity gives $u^{-1}x = lx + mbx$ integral. Conversely integrality of u and bu implies that uy lies in L_b for any integral y. This proves the lattice equality, and the explicit diagonal computation proves (7.1). It is independent of the choice of u since two bases of the same integral local lattice differ by a matrix of determinant a unit. □

The source uses the predicate `Sat` and the selected matrix `saturatingElem` in `Lib/Weil/Construction/FiniteMultiplier.lean`. The proof just given is the direct elementary divisor version of that construction. In particular there is a simple expression for every local determinant valuation, even when neither b nor its inverse is integral.

7.2 The multiplier and its transformation laws

Write $|a|_f = \prod_w q_w^{-\operatorname{ord}_w(a_w)}$ for the idele modulus, and let υ be Proposition 6.4's splitting character. In this section its restriction to finite ideles is denoted by the same symbol. Define

$$\chi_f(a) = \upsilon(a)|a|_f^{-1/2}, \qquad \Delta_f(b) = \chi_f(\det u).$$

By Lemma 7.1 and triviality of υ on finite units, this is well-defined. In particular

$$\Delta_f(b) = \prod_w (\upsilon(\pi_w)\sqrt{q_w})^{s_w(b)}, \qquad |\Delta_f(b)|^2 = \prod_w q_w^{s_w(b)}. \tag{7.2}$$

The product is finite, and the choice of local uniformizer is immaterial. This is the `glMultiplier` of the compact development.

Proposition 7.2 (Lattice multiplier identities)**.** *The multiplier has the following properties.*

1. *It is invariant under multiplication on either side by matrices in* $\mathrm{GL}_n(\widehat{\mathcal{O}})$.
2. $\Delta_f(b^{-1}) = \chi_f(\det b)\Delta_f(b)$.
3. $\Delta_f(b^*) = \overline{\Delta_f(b)}$, *where* $*$ *denotes conjugate transpose of the adelic matrix, including exchange of conjugate places.*
4. *It is multiplicative on block diagonal matrices. It is also multiplicative on a product* bb' *if at every place one of the two matrices is the identity.*
5. *For Haar measure satisfying* $\mathrm{vol}(\widehat{\mathcal{O}}^n) = 1$ *one has*

$$\mathrm{vol}(L_b) = |\Delta_f(b)|^{-2}, \qquad [\widehat{\mathcal{O}}^n : L_b] = |\Delta_f(b)|^2.$$

Proof. The first assertion is equivalent to invariance of the elementary divisors. For the second, the elementary divisors of b^{-1} are the negatives of those of b, in reverse order. Therefore

$$s_w(b^{-1}) - s_w(b) = \sum_i a_i = \operatorname{ord}_w(\det b).$$

Use (7.2) and the definition of χ_f. Transposition by itself does not change the elementary divisors, and conjugation sends the list at w to the list at $\bar{w}$. Since the two residue cardinalities agree and $\upsilon(\pi_{\bar{w}}) = \upsilon(\pi_w)^{-1}$ by (6.1), the third assertion follows. The fourth follows by addition of the local sums (7.1), respectively by separating the local factors in (7.2). Finally the index of $\pi^a\mathcal{O}_w$ in $\mathcal{O}_w$ is q_w^a for $a \geq 0$. The determinant sizes in Lemma 7.1 give the index formula, and Haar measure divides by that finite index on passage to the smaller lattice. □

Several identities used by the seesaw and the exchanged periods are immediate consequences. First, for g in the finite unitary group, $g^{-1} = g^*$ and $|\det g|_f = 1$, the latter because the determinant times its conjugate is one. Thus

$$\Delta_f(g^{-1}) = \overline{\Delta_f(g)} = \upsilon(\det g)\Delta_f(g). \tag{7.3}$$

If S is a set of places disjoint from its conjugate and g is trivial outside $S \cup \bar{S}$, let b equal its matrix at S and the identity elsewhere. Then $g = b(b^*)^{-1}$ and the supports are disjoint. Hence

$$\Delta_f(g) = |\Delta_f(b)|^2 \overline{\chi_f(\det b)}. \tag{7.4}$$

Indeed apply (2) and (3) and multiply. This identity explains why a single choice of one member of a split pair is sufficient in a local calculation: the other factor is determined by the inverse transpose and contributes the displayed conjugate factor. Replacing it by an independent local modulus would change the character.

As a one-dimensional example, suppose at a split pair g has values π^a and $\bar{\pi}^{-a}$, with $a \geq 0$. Then $s_w(g) = 0$ and $s_{\bar{w}}(g) = a$, so

$$\Delta_f(g) = \upsilon(\pi_w)^{-a} q_w^{a/2}, \qquad |\Delta_f(g)|^2 = q_w^a.$$

The same result follows from (7.4). At a conjugation-fixed place the sign in (7.2) is $(-1)^{s_w(b)}$; it is not a sign introduced by a choice of square root. It follows from the inert uniformizer value of the splitting character.

Equations (7.2)–(7.4) prove the identities `glMultiplier_inv`, `glMultiplier_star`, `glMultiplier_fromBlocks`, `latticeMeasure_inter_eq_glMultiplier`, `norm_finMultiplier_sq`, `upsilon_mul_finMultiplier`, and `finMultiplier_eq_of_supportedOn` in `Lib/Weil/Construction/FiniteMultiplier.lean`.

7.3 The finite vacuum and the Heisenberg cocycle

Let $V_f = \mathbb{A}_{E,f}^n$ with its standard Hermitian form $h(x, y) = x^* y$. It is convenient to work with the scaled lattice $L = \alpha\widehat{\mathcal{O}}^n$. Let ψ_f be the finite additive character of E, extended from the rational character by the trace. Set

$$\beta(x, y) = \psi_f(\tfrac{1}{2}\jmath h(y, x)).$$

We do not assume that the scalar 2 is a unit in the finite integers. It is invertible in the local fields, which is enough for the displayed definition. Bilinearity of the additive character makes β a bicharacter on the underlying additive group. Conjugation invariance of ψ_f and $\bar{\jmath} = -\jmath$ give

$$\beta(y, x) = \beta(x, y)^{-1}.$$

Moreover $\beta(x, x) = 1$, because $h(x, x)$ is fixed by conjugation and the trace of a purely imaginary element is zero. The latter trace assertion holds componentwise over each local base place, including split pairs.

The corresponding central extension has product

$$(x, z)(y, t) = (x + y, zt\beta(x, y)).$$

Associativity follows from the bicharacter identities, and the commutator is $\beta(x, y)^2$. The annihilator of L for this commutator pairing is L itself, by Lemma 6.6. Thus L is a maximal compact isotropic subgroup for the commutator, although β restricted to $L \times L$ may have values -1. This is the dyadic issue that must be respected when normalizing the vacuum.

Define on L the function

$$\lambda(x) = \psi_f(\tfrac{1}{2}c_0 \jmath h(x,x)), \qquad c_0 = (1-\jmath^{-1})/2.$$

It is a quadratic refinement on the lattice.

Lemma 7.3. *For $x, y \in L$ one has*

$$\lambda(x+y) = \lambda(x)\lambda(y)\beta(x,y).$$

If g is unitary and $x, gx \in L$, then $\lambda(gx) = \lambda(x)$.

Proof. Put $a = h(y,x)$ so that $h(x,y) = \bar{a}$. By conjugation invariance of the character,

$$\frac{\lambda(x+y)}{\lambda(x)\lambda(y)} = \psi_f(\tfrac{1}{2}c_0\jmath(a+\bar{a})) = \psi_f(\tfrac{1}{2}(c_0-\bar{c}_0)\jmath a).$$

Divide by $\beta(x,y) = \psi_f(\frac{1}{2}\jmath a)$ and use $c_0 + \bar{c}_0 = 1$. The ratio is $\psi_f(-\bar{c}_0\jmath a)$. Since $x, y \in L$, the element a belongs to $\alpha\bar{\alpha}\widehat{\mathcal{O}}_E$ and $\bar{c}_0$ is integral. The argument lies in the inverse different by (6.2), so its character is trivial. This proves the refinement. The second assertion is the equality of the Hermitian norms under g. □

The finite part of the source's `vacuumFn` in `Lib/Weil/Construction/AdelicHeisenberg.lean` is $\lambda\mathbf{1}_L$. This proves its essential lattice identity directly from the choice of the different idele and the trace-one element. In particular it would not suffice to use the indicator alone in a presentation of the Heisenberg bicharacter with the factor $1/2$.

7.4 The lattice model and the absolute vacuum coefficient

We describe the finite part in a standard induced model, rather than assume an arbitrary Hilbert-space normalization. Let $\mathcal{H}_L$ be the Hilbert space of functions f on V_f satisfying

$$f(y+\ell) = \lambda(\ell)\beta(y,\ell)f(y) \qquad (\ell \in L), \tag{7.5}$$

with square-summable values on the countable discrete quotient V_f/L. Use counting measure on that quotient. The absolute value of an equivariant function is independent of the representative by the unit modulus of all factors; hence the quotient norm is well-defined. Consistency of two translations in (7.5) is precisely the identity in Lemma 7.3, together with $\beta(\ell_1,\ell_2)^2 = 1$ on L. Define

$$(\rho(x)f)(y) = \beta(x,y)f(y-x).$$

For $\ell \in L$ the bicharacter terms show that this formula preserves (7.5), and it is an isometry because translation permutes the quotient. A direct calculation gives $\rho(x)\rho(z) = \beta(x,z)\rho(x+z)$, the cocycle convention of the source's `Heisenberg.Model`.

On L, we also have $\lambda(x)^2 = 1$, by integrality of c_0, self-duality and the definition of the finite character. The function $e = \lambda\mathbf{1}_L$ satisfies (7.5) by the refinement and is of norm one. For $\ell \in L$, $\rho(\ell)e = \lambda(\ell)e$ (use the refinement on the common support); if $x \notin L$ then $\rho(x)e$ has support in a different coset and is orthogonal to e. The vectors $\rho(x)e$ indexed by representatives of the quotient form an orthonormal basis, up to the irrelevant phases of those representatives. An element in $\mathcal{H}_L$ transforming by λ under all $\rho(\ell)$ has support in L: at any other coset, the equivariance formula and the action formula force the commutator character $\beta(\ell,y)^2$ to be identically one, hence $y \in L$. Its values on L are determined by its value at zero, so the space of such vectors is the single line $\mathbb{C}e$.

Let g be in the finite unitary group. It acts symplectically on V_f. Suppose U_g is a unitary intertwiner implementing g on the Heisenberg operators,

$$U_g \rho(x) U_g^{-1} = \rho(gx).$$

Existence and uniqueness up to phase follow from the standard Stone–von Neumann theorem, or directly by comparing the induced models of the two maximal compact isotropic subgroups L and gL. Here we compute the exact modulus of its vacuum coefficient without making a phase choice.

There is a direct identification with the integrable reproducing-kernel model in `Lib/Weil/Construction/Model.lean`. With $e = \lambda \mathbf{1}_L$ and $\mathrm{vol}(L) = 1$, its defining equation is

$$f(x) = \int_{V_f} f(y)\beta(y,x)e(x-y)\,dy.$$

If f satisfies (7.5), set $y = x - \ell$ on the support of the integrand. Then the lattice equivariance of f and the identities $\lambda(-\ell) = \lambda(\ell)$, $\lambda(\ell)^2 = 1$, and $\beta(\ell, x) = \beta(x,\ell)^{-1}$ show that the integrand is exactly $f(x)$. The integral has volume one. Conversely, replace x by $x + \ell$ in the reproducing equation; use the lattice equivariance of $e(x-y)$ and the bicharacter identity $\beta(y,\ell)\beta(x-y,\ell) = \beta(x,\ell)$. This yields (7.5). Thus the source's integrable finite model and the integrable lattice-equivariant functions agree, and the finitely supported coset functions are dense in their Hilbert completion. Since both implement the same group action on the Heisenberg operators, equality of the normalized vacuum coefficient identifies the finite intertwiners, not only their absolute coefficients.

Proposition 7.4. *With e as above,*

$$|\langle e, U_g e\rangle|^2 = [L : L \cap gL]^{-1} = |\Delta_f(g)|^{-2}.$$

Proof. On L the operators $\lambda(\ell)^{-1}\rho(\ell)$ form a genuine unitary representation, since the refinement cancels the cocycle on L. Integrating them against the probability Haar measure of L gives the orthogonal projection P onto the vacuum line just identified. Taking the matrix coefficient of P at $U_g e$, one obtains

$$\begin{aligned} |\langle e, U_g e\rangle|^2 &= \langle U_g e, P U_g e\rangle \\ &= \int_L \lambda(\ell)^{-1}\langle e, \rho(g^{-1}\ell)e\rangle\,d\ell. \end{aligned}$$

Here the inner product is linear in the second variable. The integrand vanishes off $L \cap gL$ by the coset support of the translated vacuum. On that intersection it equals $\lambda(\ell)^{-1}\lambda(g^{-1}\ell) = 1$ by the norm invariance in Lemma 7.3. The integral is therefore the reciprocal lattice index. Scaling by α commutes with g, and Proposition 7.2 applied to g^{-1} gives the last equality, using (7.3). □

The phase-normalized operator U_g is specified by $\langle e, U_g e\rangle = 1/\overline{\Delta_f(g)}$. Its inverse has vacuum coefficient $1/\Delta_f(g)$, so (7.3) and uniqueness up to a scalar give $U_g^{-1} = U_{g^{-1}}$. Uniqueness of an intertwiner still implies only $U_g U_h = c(g,h)U_{gh}$ for some scalar of absolute value one. We next prove that scalar equals one.

7.5 The Gauss-sum phase and the finite splitting

Extend the expression for λ to all of V_f (the refinement identity was proved only on L). Write $f_g = U_g e$ and $c_g = 1/\overline{\Delta_f(g)}$. For $a \in gL$, intertwining gives $\rho(a)f_g = \lambda(g^{-1}a)f_g = \lambda(a)f_g$,

using preservation of the Hermitian norm. At zero the eigenrelation yields $f_g(-a) = c_g\lambda(a)$; since the expression for λ is even, $f_g(a) = c_g\lambda(a)$. At any x in the support of f_g, comparison with the lattice-model eigenrelation for $a \in L \cap gL$ forces $\beta(a,x)^2 = 1$. Both lattices are self-dual for this commutator pairing. Their intersection has annihilator equal to their sum: the annihilator of a sum is the intersection of annihilators, and the double-annihilator identity for compact open lattices follows from perfect trace duality on finite lattice quotients. Hence f_g is supported in $L + gL$.

Write $x = a + b$ with $a \in gL$ and $b \in L$. By (7.5),

$$f_g(x) = c_g\lambda(a)\lambda(b)\beta(a,b) = c_g\lambda(x)\psi_f(\bar{c}_0\jmath h(b,a)). \tag{7.6}$$

For the second equality, by conjugation invariance of ψ_f, the ratio $\lambda(x)/(\lambda(a)\lambda(b))$ is $\psi_f((c_0 - \bar{c}_0)\jmath h(b,a)/2)$ and $1 - c_0 + \bar{c}_0 = 2\bar{c}_0$. This does not require $a \in L$.

The common support in the inner product of $f_{g^{-1}}$ and f_h gives the finite $\widehat{\mathcal{O}}_E$-module

$$Q = ((L + g^{-1}L) \cap (L + hL))/L.$$

Formula (7.6) and cancellation of $\lambda(x)$ show that

$$\overline{f_{g^{-1}}(x)}f_h(x) = \overline{c_{g^{-1}}}c_h F(x \bmod L)$$

where F is well-defined by common lattice equivariance, $F(0) = 1$, and all its values are roots of unity. More precisely F is the additive character of the difference of the two cross terms in (7.6). Scaling the two decompositions of x by $t \in \widehat{\mathcal{O}}_E$ scales this difference by $\bar{t}t$, so

$$\begin{aligned} F(tq) &= F(q) && \text{if } \bar{t}t = 1, \\ F(tq) &= 1 && \text{if } \bar{t}t = 0, \\ F(tq) &= \overline{F(q)} && \text{if } \bar{t}t = -1. \end{aligned} \tag{7.7}$$

Compute the normalized vacuum coefficient:

$$\begin{aligned} \frac{c(g,h)}{\Delta_f(gh)} &= \langle e, U_g U_h e\rangle = \langle U_{g^{-1}}e, U_h e\rangle \\ &= \frac{\sum_{q\in Q} F(q)}{\overline{\Delta_f(g)}\,\overline{\Delta_f(h)}}. \end{aligned} \tag{7.8}$$

This is the finite Gauss-sum relation. All lattices, modules and additive characters factor over finite places of F, treating the two components of a split pair together. Only finitely many factors are nontrivial. It suffices to show that the corresponding local cocycle is identically one.

Lemma 7.5 (The orbit sign test)**.** *Suppose a finite abelian group Q has no nonzero element of order two, and an automorphism i has $i^2 = -1$. Let $F : Q \to \mathbb{C}^\times$ have root-of-unity values, with $F(0) = 1$, $F(iq) = F(q)$, and $S = \sum_q F(q)$ real. If an integer $r \equiv 1 \pmod 4$ is nonzero and*

$$cr^a = Sr^b, \qquad a, b \geq 0, \quad |c| = 1,$$

then $c = 1$.

Proof. Every nonzero orbit of i has size four: if its size divided two, then $q = -q$, hence $q = 0$. Thus $S = 1 + 4A$ for an algebraic integer A, and the assumed equality gives $c = 1$ or -1. If $c = -1$, it implies $(r^a + r^b)/4 = -r^b A$, a rational algebraic integer and hence an integer. This is impossible since the numerator is 2 modulo 4. □

Theorem 7.6 (The finite normalized action)**.** *For the data of Chapter 6, the phase-normalized Heisenberg intertwiners satisfy $U_g U_h = U_{gh}$.*

Proof. At a split place write h as $(a, (a^t)^{-1})$ over the two copies of F_v. First suppose a^{-1} is integral. Then the second component of hL is contained in L, so Q is supported on the first side. With t the projection idempotent of that side one has $tq = q$ and $\bar{t}t = 0$. Equation (7.7) implies $F(q) = 1$ for every q, giving a positive integer sum.

For a split unitary element with first component b, the elementary divisor lists on the two sides are opposite. The phase in (7.2) is $\upsilon(\pi_w)^{-\operatorname{ord}_w(\det b)}$, a multiplicative character of b. Hence the multiplier ratio in (7.8) is positive real. The unit-modulus scalar $c(g, h)$ is therefore one for every g when a^{-1} is integral. The elements h with $c(g, h) = 1$ for all g form a subgroup by the cocycle identity $c(g, h)c(gh, k) = c(g, hk)c(h, k)$ (also apply it with $k = h^{-1}$). Integral unitary elements and diagonal elements with inverse integral lie in that subgroup. By the elementary divisor theorem they and their inverses generate the split group.

At a conjugation-fixed place w the quadratic local extension is unramified and $q_w = r^2$ with $r = q_v$. The residue characteristic is odd: a fixed place of E would restrict to a fixed place of K, and 2 splits in K. The reduction of $i = \sqrt{-1}$ has order four and its conjugate is its negative, so the nontrivial residue automorphism $x \mapsto x^r$ gives $r \equiv -1 \pmod 4$. The inert uniformizer value of υ is -1, whence the local multiplier (7.2) is an integer power of $-r$.

The local group Q has odd prime-power order. Multiplication by i squares to negation and preserves F by (7.7). By Lemma 6.1 choose a unit of norm -1; multiplication by it permutes Q and conjugates every value of F. Thus the sum of F is real. Equation (7.8) is of the form in Lemma 7.5, with base $-r \equiv 1 \pmod 4$ and nonnegative exponents $s_w(g) + s_w(h)$ and $s_w(gh)$. This proves $c(g, h) = 1$, completing both local cases and hence the global finite proof. □

The sign lemma and finite-lattice sum supply the mathematical content of `eq_one_of_orbits` in `Lib/Weil/Construction/GaussSumSign.lean` and `stdLattice.mem_cocycleRadical` in `Lib/Weil/Construction/ThetaFunctional.lean` for the specified fields, in the lattice model. The proof uses the standard irreducible Heisenberg intertwiner but computes its finite normalization independently. An absolute value of a Gauss sum would not suffice: the order-four orbits determine the final inert sign.

7.6 The real harmonic model and its basic normalization

The real part of the same oscillator has a particularly convenient holomorphic model. For a positive Hermitian space of complex dimension n, put on holomorphic polynomials the Fischer–Bargmann pairing

$$\langle P, Q \rangle = \pi^{-n} \int_{\mathbb{C}^n} \overline{P(z)} Q(z) e^{-|z|^2} \, d^{2n} z.$$

The constant polynomial 1 has norm one. For multi-indices a, b the monomials satisfy

$$\langle z^a, z^b \rangle = \delta_{a,b} \prod_j a_j!,$$

because angular integration kills distinct exponents and the radial integral at each coordinate is $\int_0^\infty t^{a_j} e^{-t} dt = a_j!$. Thus, in the Hilbert completion, $z^a / \sqrt{a!}$ is an orthonormal basis. The reproducing kernel is

$$K(z, w) = \sum_a \frac{z^a \bar{w}^a}{a!} = e^{z \cdot \bar{w}},$$

with locally uniform convergence by the exponential series. For a unitary matrix u, the action on polynomials by $P(z) \mapsto P(u^{-1}z)$ preserves the pairing, because both the Gaussian and Lebesgue measure are invariant. Tensoring with a determinant character of modulus one likewise preserves it. These elementary facts underlie the integral Fischer pairing and the compact harmonic coefficient formulas in Chapter 5.

The scale $s_\tau = (2\pi \operatorname{Im} \tau(\jmath))^{-1/2}$ fixed in Chapter 6.3 gives the norm-one vacuum wave function a Gaussian of the form $e^{-|z|^2/2}$ in the original coefficient variable, since $\pi \operatorname{Im} \tau(\jmath)|s_\tau z|^2 = |z|^2/2$. Its square therefore has the Bargmann density $e^{-|z|^2}$. For an indefinite unitary group the holomorphic and antiholomorphic variables are chosen according to the signs, and mixing them is the noncompact oscillator action. The compact calculation alone does not determine the noncompact multiplier or the full Flensted–Jensen coefficient. Those are studied in Chapter 22 in the normalization required by the filling system.

7.7 The rational theta functional in a polarized model

Finally, to explain the relationship between the Heisenberg model and adelic theta sums, consider a symplectic F-space with a rational Lagrangian splitting $X \oplus X^\vee$. For a Schwartz–Bruhat function f on $X(\mathbb{A}_F)$ define

$$\theta(f) = \sum_{x \in X(F)} f(x).$$

The sum is absolutely convergent: the finite part of a Schwartz–Bruhat function is compactly supported and locally constant, reducing the possible rational points to a finite union of lattices in the real vector space; the real part and all its derivatives decay faster than any polynomial. The Poisson identity

$$\sum_{x \in X(F)} f(x) = \sum_{x \in X^\vee(F)} \widehat{f}(x)$$

uses the self-dual Haar measure for the standard rational additive character. One proof is to reduce to the ordinary real lattice Poisson formula by decomposition into finite congruence cosets, followed by the finite character orthogonality of the dual lattice. The inverse different is exactly the trace-dual lattice used in that proof.

The three basic Schrödinger operations are changes of coordinates, multiplication by quadratic characters, and Fourier transform. For a rational invertible coordinate change $A \in \mathrm{GL}(X)(F)$, the change of variables factor in the oscillator is $|\det A|_{\mathbb{A}_F}^{1/2} = 1$ by the product formula, and the rational summation indices are permuted. For a rational symmetric form B on X, the operation multiplying by the full additive character of $\frac{1}{2}B(x,x)$ fixes each rational summand, because the full additive character is trivial on F. Fourier transform preserves the sum by Poisson. These operations generate the rational symplectic action in its standard global splitting, which follows from the elementary symplectic row reduction using Lagrangian bases. This is the usual proof of rational theta invariance in a Schrödinger model. When a unitary dual pair is embedded in this symplectic group and equipped with the same rational splitting, the calculation applies to its rational points too.

There are two comparisons one must still supply in the nonpolar presentation. First, the compact source constructs the theta functional from invariance under rational Heisenberg translations and normalizes it by a vacuum coefficient; existence and one-dimensionality of its continuous invariant functional are established in `Lib/Weil/Construction/ThetaFunctiona l.lean`. Second, it proves that the exact phase-normalized Weil operators fix this functional, using a doubling argument, `Space.theta_omega_ratEmbed` in `Lib/Weil/DualPair/PairSp`

`litting.lean`. The classical polarized calculation above proves the rational invariance in the standard model and explains the role of Poisson and product formula. To deduce the assertion for the precise nonpolar normalization one must verify the unitary intertwining identification including the real phase and theta-functional normalization. The finite phase has now been calculated in Theorem 7.6, but the global identification is not a consequence of that calculation alone. Its normalized mixed-model formulas are the input used in Chapter 16, where the actual orbit reduction and degenerate and nonsingular finite lattice calculations are developed.

Classical background and precise scope. For the existence and splitting of the adelic Weil representation, the classical sources are Weil's construction and Kudla's unitary dual-pair splittings ([15, 16]). The proofs in this chapter establish the finite saturation, all multiplier and index identities, the lattice quadratic refinement, absolute vacuum overlap, and finite normalized group action, and recall the usual polarized theta invariance. The remaining real-phase and theta-model-identification assertions are kept separate. They cannot be removed by proving only that the operators are unitary.

Chapter 8

Foundational theorem audit and universal coefficient comparison

In a conventional exposition a suitably precise reference is a proof of a known theorem, not a licence to cite a different theorem with similar terminology. This chapter checks several foundational interfaces appearing earlier and supplies additional proofs rather than requesting a quantitative strengthening of the literature. The conclusions concern the standard integral coefficient modules, the classical number-field and elliptic-curve facts, the finite condition in local Galois cohomology and integral characteristic-polynomial identities. They do not establish the integral period, automorphic, or quantitative geometric assertions by themselves.

8.1 The universal integral Schur basis

The integral coefficient statement in Section 5.6 and Proposition 13.15 is stronger than the existence of a basis of a complex irreducible representation. An isomorphism over $\mathbb{Q}$ between two integral coefficient models need not identify their reductions at the primes where the jet argument takes place. We establish the requisite universal basis and its coefficient-extraction property.

In the notation of `Lib/GLn/SchurModule/Nabla.lean`, the semi-invariance is the identity with *generic* upper triangular matrices, equivalently the identity after extension to every commutative algebra over the ground ring. In particular it is not an identity tested only on rational points over a finite field. This distinction is essential. For a partition $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$ with integral entries, let $P_{\lambda,R}$ be the submodule of $R[X_{ij}]$ consisting of multihomogeneous polynomials of column degrees $\lambda_1, \ldots, \lambda_n$. It is finite free over the arbitrary commutative ring R. Write $M_{\lambda,R}$ for its submodule satisfying the polynomial identity

$$f(XT) = \prod_j T_{jj}^{\lambda_j} f(X)$$

for generic upper triangular T. All the matrices and coefficient homomorphisms in this paragraph are defined over $\mathbb{Z}$.

Theorem 8.1 (Integral flag-minor basis, with base change)**.** *For each Young diagram of shape λ, with column lengths ordered nonincreasingly, choose the semistandard fillings with entries from $1, \ldots, n$: strictly increasing in their columns and weakly increasing in their rows. For such a filling A, let b_A be the product over its columns of the minors with the indicated rows and the first h columns, where h is the column length. The polynomials b_A form an R-basis of $M_{\lambda,R}$ for every commutative ring R. The basis is universal under base change, and its image*

is a direct summand of $P_{\lambda,R}$ by an integral coefficient-extraction map. Its character is the Schur polynomial.

Proof. The triangular substitution preserves the linear span of the first h columns, and acts on their exterior product through the determinant of the leading triangular block. Thus each b_A has the stated weight and belongs to the required module.

Take lexicographic order in the variables, ordering by column first, and within each column by increasing row index (the lower row index precedes and is larger than the higher one in the variable order). A minor with increasing row indices $i_1 < \cdots < i_h$ then has diagonal leading term

$$X_{i_1,1}X_{i_2,2}\cdots X_{i_h,h}$$

with coefficient 1: any nonidentity permutation first differing at column j puts a larger row index into that column. Therefore the product has unit leading coefficient. Its leading monomial records, for every row of the Young diagram, the multiset of entries in that row: the row of diagram index j contributes the variables with second index j. Since the entries in each such row are weakly increasing, these multisets determine the entire filling. Different semistandard fillings therefore have different leading monomials. Restricting the coefficient matrix to the leading monomials in their decreasing order gives a unit triangular square matrix. The span $L_{\lambda,\mathbb{Z}}$ of the b_A is a direct summand of the free module $P_{\lambda,\mathbb{Z}}$. Taking those coefficients and multiplying by the inverse of the unit triangular matrix gives an integral retraction onto the span. This argument proves independence and the split embedding over *every* commutative ground ring, not just a field.

We next show that after tensoring with any field k there are no more semi-invariants. Over its algebraic closure, restriction from the affine matrix space to GL_n is injective (localization at the determinant). Its image is in the space of regular functions on GL_n with the displayed right Borel transformation law. This is the space of sections of the associated homogeneous line bundle on the complete flag variety. In the convention $f(gb) = \chi(b)^{-1}f(g)$ the right upper Borel character defining the bundle is $-\lambda$; conjugating by the longest Weyl element gives the dominant lower Borel character $-w_0\lambda$. Its Weyl dimension is the same as that of λ (the index reversal leaves all the factors in the dimension formula unchanged). In arbitrary characteristic this dimension follows from Kempf vanishing and the Weyl character theorem for induced modules: Jantzen, *Representations of Algebraic Groups*, 2nd ed., Part II, §4.5 and §5.10, or Milne, *Algebraic Groups*, Theorems 22.50–22.52 (the 2022 author version is at https://www.jmilne.org/math/Books/iAG2022.pdf; the indicated theorem numbers and the all-characteristic formulation have been checked). Using GL_n rather than SL_n just retains the determinant character; this is harmless for dimension. Right multiplication by the whole triangular group, tested by the universal polynomial condition, implies exactly the required equivariance on this open set.

Here is a direct verification that the number and weights of the indicated tableaux equal this character. Put

$$a_j = \lambda_j + n - j, \qquad A_\lambda(x_1,\dots,x_n) = \det(x_i^{a_j}), \qquad A_0 = \prod_{i<j}(x_i - x_j).$$

For $z = x_n$, replace each column $j < n$ by that column minus $z^{a_j - a_{j+1}}$ times column $j+1$, simultaneously. This is a triangular change of determinant one. The last row is zero except at the last column. Divide each of the first $n-1$ rows of the resulting minor by $x_i - z$, using

$$\frac{x^{a_j} - z^{a_j - a_{j+1}}x^{a_{j+1}}}{x - z} = \sum_{b=a_{j+1}}^{a_j - 1} x^b z^{a_j - 1 - b}.$$

Multilinearity of the determinant gives

$$\frac{A_\lambda(x_1,\dots,x_{n-1},z)}{\prod_{i<n}(x_i-z)} = \sum_{\lambda_j\ge\mu_j\ge\lambda_{j+1}} z^{|\lambda|-|\mu|}A_\mu(x_1,\dots,x_{n-1}).$$

Indeed $b_j=\mu_j+n-1-j$ varies between a_{j+1} and a_j-1, and the remaining exponent of z is $\sum_j a_j-(n-1)-\sum_{j<n} b_j$. Dividing by the smaller Vandermonde determinant yields the branching recursion for A_λ/A_0, initially on the locus of distinct variables and then as a polynomial identity. In a semistandard filling the boxes filled by n form a horizontal strip; removing them leaves exactly an interlacing shape μ, and conversely filling the strip by n reconstructs the tableau. The tableau-generating polynomial obeys the very same branching recursion and the case $n=1$. This proves the character statement and gives the dimension equality after setting all the variables equal to one. Consequently the independent minor products form an entire basis of the semi-invariant space over every algebraically closed field, and hence over any field by scalar extension and coefficient comparison.

To pass from fields to general commutative rings, use the polynomial identity defining semi-invariance as a linear map $Q:P_{\lambda,\mathbb{Z}}\to P'$ between finite free abelian groups. Here P' is the group spanned by the finitely many monomials which occur in the substituted polynomial and in its proposed transformation, with both families of variables. The kernel after scalar extension is precisely $M_{\lambda,R}$. The prescribed column degrees follow also directly from diagonal substitutions and comparison of monomials in the generic diagonal variables, without dividing by their values. Choose a free complement C to $L_{\lambda,\mathbb{Z}}$. The restriction $Q|_C$ is injective after extension to $\mathbb{Q}$ and after reduction modulo every prime, by the preceding field argument. Its Smith normal form can therefore have no nonunit nonzero diagonal entry and must have full column rank. It has an integral left inverse. Hence it remains injective over every R, even a nonflat $\mathbb{Z}$-algebra with nilpotents. This shows $M_{\lambda,R}=L_{\lambda,R}$. □

For completeness, the same argument accommodates the colors and the variable number of columns in `PolyRep.exists_minorProduct_basis`. For a fixed color with m rows, extend the prescribed column degrees by zeros. If a nonzero degree lies beyond the m-th column, localize by the determinant of the generic first m columns. Subtract linear combinations of those columns from all the later ones using a unipotent upper triangular matrix over that localization. All later columns become zero, so the polynomial vanishes by multihomogeneity. The localization map is injective over any ring, since the determinant with its unit leading coefficient is a nonzero divisor in the polynomial ring. The semi-invariant module is consequently zero, agreeing with the absence of any strictly increasing tableau of column length greater than m. If all later degrees are zero, the polynomial is independent of those variables and the square case applies. For several colors the variables separate and each weight is imposed separately. The integral splittings can be applied successively over the polynomial coefficient ring in the other colors, or equivalently their tensor products can be applied; the result is the universal basis consisting of the products of the individual bases. This proves the existence and character assertions even for the more general colored row arrays of the source.

The two standard-basis citations in the previous manuscript were Akin–Buchsbaum–Weyman, *Schur functors and Schur complexes*, Adv. Math. **44** (1982), 207–278, Theorem II.2.16 (standard tableaux basis), and Theorem II.4.11 (decomposition filtration); and De Concini–Eisenbud–Procesi, *Young diagrams and determinantal varieties*, Invent. Math. **56** (1980), 129–165. The statements in II.2.16 and II.4.11 have been checked in the author-hosted article (https://jerzyweyman.com/wp-content/uploads/2022/11/Publ1-Schur-Functors-and-Schur-Complexes.pdf). They concern the Schur functor basis and direct-sum filtration; they are not on their own a proof that an arbitrary-ring kernel of a polynomial

substitution equals that span. The direct-summand and all-characteristic dimension argument above supplies that last step. The paper by De Concini–Eisenbud–Procesi is an alternative straightening-law reference, but nothing here depends on identifying a particular lemma there.

The source chooses an unspecified family of standard fillings by the choice operator, requiring universal independence, spanning of all minor products and the character identity. This does not require that its chosen family is literally the set of semistandard tableaux in Theorem 8.1. The stated conditions make any such choice an integral basis of the same module; in particular the transition between two choices and its inverse lie over $\mathbb{Z}$, by uniqueness of expansion over $\mathbb{Z}$. Because the semistandard model has an integral left inverse, the coordinate maps for the source's chosen basis do also. Thus if the coefficients of a polynomial expressed in this basis belong to an additive subgroup A of R, so do the extracted coordinates: the coefficient-extraction map is integer-linear and uses no division in R. This is the assertion `PolyRep.coord_mem_of_coeff_mem` in `Lib/GLn/SchurModule/SchurMatrix.lean`. The distinction between a coordinate model and an abstract isomorphism over $\mathbb{C}$ is thereby removed.

Corollary 8.2 (Rank change and the integral Schur expansion)**.** *Let $n \le n'$ and let all column multiplicities of heights strictly between n and n' be zero. As in the source, the basis in rank m uses columns of heights strictly less than m. Thus, if $n < n'$, the bigger diagram includes $\kappa(n)$ columns of height n which have been omitted from the smaller diagram. Put $e = \kappa(n)$ if $n < n'$, and $e = 0$ otherwise. For a matrix Z of shape $n' \times n$, the row-substituted polynomial has an expansion*

$$b_{A'}(ZY) = \sum_{A} c_{A'A}(Z) b_A(Y) \det(Y)^e$$

over the polynomial ring in the entries of Z, with $c_{A'A}$ integral polynomials in those entries. Specialization to every commutative ring is allowed, as is extraction of coordinates by an integer-linear map.

Proof. For a column of height less than n expand its transformed minor by Cauchy–Binet. For a column of height n, the product formula for two square minors factors out $\det(Y)$; there are exactly e such factors. The remaining products are in the weight module described in the preceding theorem, which provides the coefficients. For the coefficient-extraction assertion with the additional determinant factor, observe that the leading term of $\det(Y)^e$ is a monomial with coefficient one. The products $b_A \det(Y)^e$ therefore still have distinct leading monomials with coefficient one and an integral extraction map. For a different permissible universal basis, use the unimodular coordinate transition. The argument in each color separates, yielding the determinant product in the multi-color version. □

This is the exact rectangular identity and integral coordinate assertion in Proposition 13.15. In particular the Schur matrix identity `PolyRep.sum_schur_smul_nablaBasis`, its rank restriction and base change in `SchurMatrix.lean`, and the integral basis used in Section 5.6 can be treated as established, not additional specialized hypotheses. The numerical valuation bound on the entries of a particular theta section is a separate matter; it is not implied by this integral basis theorem.

8.2 The norm subgroup and the normalization of the splitting character

In Chapter 6 the arithmetic step is to use the norm subgroup for the quadratic CM extension E/F together with the local sign characters. One may use Neukirch, *Algebraic Number Theory*,

Grundlehren 322, Springer, 1999, Chapter VI, §5, Theorem (5.5), which identifies the quotient of the idele class group by the norms from a finite Galois extension with the abelianized Galois group. In the quadratic case, passing from idele classes to ideles gives exactly

$$J_F/(F^\times N_{E/F} J_E) \simeq \mathrm{Gal}(E/F).$$

Local-global compatibility of the reciprocity map (the same section) sends real nonnorms to the nontrivial element and at an unramified inert place a uniformizer to Frobenius; an integral unit maps to the identity. For a split place the local norm is surjective. Since the target has order two, the distinction between the reciprocity map and its inverse convention has no effect on these signs. Chapter VI, §4, Corollary (4.5) of the same book supplies the cyclic Hasse norm theorem for principal elements, should one wish to recognize an everywhere locally normed principal element as a global field norm. Both statements have been checked in the published text (`https://web.math.ucsb.edu/~agboola/teaching/2021/fall/225A/neukirch.pdf`).

These are precisely the norm facts used in Proposition 6.4 and Corollary 6.5. In particular they imply the index-two assertion and the actual finite-place signs of the splitting character. The extra hypothesis on roots of unity, which appears in the source's `nonempty_splitti ngChar`, is not a class-field-theory hypothesis: it is verified in Lemma 6.3. The subsequent extension of a circle-valued character from the product subgroup is proved in the text. The combination is an ordinary proof of the source's existence statement at `Lib/ClassField/H eckeCharacter/SplittingCharExists.lean`; no appeal to an unspecified adelic character with prescribed behavior is needed. Similarly the elementary class-group proof of Lemma 6.2 establishes the odd order and bijectivity of squaring that were called genus theory in the earlier CM chapter. Neither result proves any of the later quantitative trace identities by itself.

8.3 Classical CM and the discriminant normalization

The classical facts used in Proposition 12.1 and the local isogeny-degree argument are indeed available in the following precise form.

- J. H. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, GTM 151, Springer, 1994, Chapter II, Theorem 6.1 states that the j-invariant of a complex elliptic curve with complex multiplication is an algebraic integer. It applies to the $\mathcal{O}_K$-lattice of this construction and every conjugate. This is the precise location, in place of the previous broad reference to §6.
- J. H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd ed., GTM 106, Springer, 2009, Chapter VII, Proposition 5.5 gives potential good reduction if and only if j is integral over the local valuation ring. It is valid including residue characteristics 2 and 3.
- Silverman, *Advanced Topics*, Chapter II, Proposition 4.4, says that the reduction map on morphisms of elliptic curves with good reduction at the chosen finite place is injective and preserves degrees. The theorem allows inseparable reduction and makes no prime-to-residue-characteristic assumption on the degree.

The theorem and proposition statements have been checked in the books. To apply them choose a number field containing the normalized equation, the two-torsion point, and any finitely many endomorphisms or isogenies in question. This is possible: elliptic curves and their torsion points can be defined over number fields in the CM case; an isogeny is represented by the quotient by its finite torsion kernel followed by an isomorphism, and the invariant

differential fixes its complex multiplier. The chosen valuation extends to the number field, and a finite extension affords good reduction. In this extension the smooth proper group model is an abelian scheme, to which the endomorphisms extend by the good reduction Néron property. Thus the two factors of the 2-torsion at the split prime 2 are indeed finite flat; the zero-tangent factor is étale of rank two. Proposition 12.1 then gives the stronger statement about the unit discriminant of the *particular* marked cubic, rather than simply using potential good reduction of an unrelated model. Conversely the bound on the first unit coefficient in the good-reduction local degree computation follows by taking the local length at the lifted point in a reduced isogeny of degree q, and hence does not require the CM formal group to be identified with a chosen Lubin–Tate series. This distinction between local degree and total degree is relevant at the inert primes, where q is a square.

There is a related analytic reference whose normalization is worth proving directly. The unit-adjusted Chowla–Selberg calculation in Proposition 12.11 uses the first Kronecker limit formula for an area-normalized lattice, and a product formula for the discriminant. Different conventions concerning full and half periods in the literature can alter the constant in that assertion.

Proposition 8.3 (Kronecker limit with full periods and area)**.** *Let $L = \alpha(\mathbb{Z} + \tau\mathbb{Z})$ with $y = \Im\tau > 0$, area $A_L = |\alpha|^2 y$, and let $k_0 > 0$. In full-period notation the lattice invariants satisfy*

$$\Delta_L = g_2(L)^3 - 27g_3(L)^2 = (2\pi)^{12}\alpha^{-12}\eta(\tau)^{24}.$$

Then

$$(s-1)\sum_{\ell\in L\setminus\{0\}}\left(\frac{|\ell|^2}{k_0 A_L}\right)^{-s} = \pi k_0 + \pi k_0(s-1)\left(2\gamma_{\mathrm{E}} - \log\frac{A_L}{k_0\pi^2} - \frac{1}{6}\log|\Delta_L|\right) + O((s-1)^2).$$

Proof. The discriminant identity with generators $1, \tau$ is NIST DLMF (23.19.4), at https://dlmf.nist.gov/23.19.E4. Its convention is $g_2^3 - 27g_3^2$, as specified there and in (23.3.4); homogeneity of the lattice sums gives the factor α^{-12}. The eta product in this convention is $\eta(\tau) = e^{2\pi i\tau/24}\prod_{n\geq 1}(1 - e^{2\pi in\tau})$.

Write $\tau = x + iy$ and

$$Z(\tau, s) = y^s \sum_{(m,n)\neq(0,0)} ((m + nx)^2 + n^2y^2)^{-s}.$$

For $\Re s > 1$, separate the $n = 0$ terms and apply the Mellin integral and Poisson summation in m to the rest. The zero Fourier term is obtained by the one-dimensional Gaussian integral. In full,

$$Z(\tau, s) = 2y^s\zeta(2s) + \frac{2\sqrt{\pi}\,\Gamma(s - \frac{1}{2})}{\Gamma(s)}y^{1-s}\zeta(2s-1) + R(\tau, s).$$

To justify the remainder near one, use the Mellin integrals for the nonzero Fourier frequencies k:

$$\frac{\sqrt{\pi}}{\Gamma(s)}\int_0^\infty t^{s-3/2}\exp(-n^2y^2t - \pi^2k^2/t)\,dt.$$

The phase factor is $e^{2\pi iknx}$. These integrals (multiplied by y^s) converge and their series over $n, k \neq 0$ converges locally uniformly in a complex neighborhood of one, since $n^2y^2t + \pi^2k^2/t \geq 2\pi|nk|y$. Retaining half this exponential for the integral itself leaves an exponentially convergent double sum times a fixed polynomial in $|n|, |k|$; splitting the integral at $t = \pi|k|/(|n|y)$ gives that polynomial bound using the decay at 0 and ∞. The same estimate with $|\log t|$ proves holomorphy by differentiation.

At $s = 1$, the Fourier transform of $(u^2 + (ny)^2)^{-1}$ is $\pi/(|n|y)e^{-2\pi|kn|y}$, by an elementary residue calculation. Thus

$$R(\tau, 1) = 4\pi\Re \sum_{n,k\geq 1} \frac{e^{2\pi i n k\tau}}{n} = -4\pi \log \left| \prod_{k\geq 1} (1 - e^{2\pi i k\tau}) \right|.$$

Using $\psi(1/2) - \psi(1) = -2\log 2$ (from Gamma duplication), the second displayed term of $Z(\tau, s)$ is

$$\frac{\pi}{s-1} + 2\pi\gamma_{\mathrm{E}} - 2\pi\log 2 - \pi\log y + O(s-1).$$

The first term at one is $\pi^2 y/3$, which combines with $R(\tau, 1)$ to give $-4\pi\log|\eta(\tau)|$. Finally the lattice sum in the claim is $k_0^s Z(\tau, s)$. Multiplication by k_0^s adds $\pi k_0 \log k_0$ to its constant coefficient. Substitution of the discriminant identity now gives the formula as written. □

This proves the precise limit identity used in (12.12), corresponding to `kronecker_limit` in `Lib/CM/KroneckerLimit/LogTheta.lean`. The algebraic-unit part of Chowla–Selberg in Proposition 12.11 is not supplied merely by this limit formula; its proof there combines the class-product identity, integrality of the marked cubic, and integral isogeny multipliers. All these ingredients have conventional proofs in the earlier text. A reference to a usual product-of-periods Chowla–Selberg formula, without accounting for the unit whose conjugates normalize every period separately, would not justify the finite valuation conclusions. The soft normalized reciprocity step `CMLattice.complex_multiplication_theta` has also been reconstructed in Theorem 16.5. What remains is the identification of the normalized finite Weil vectors and real harmonic coefficient maps in the actual chain, as identified in Section 16.15; the classical CM references do not claim those identities.

8.4 The exact classical finite local condition

We clarify the status of Proposition 49.1. The relevant source definition is `crysCocycles` in `Lib/PadicHodge/Crystalline/Defs.lean`; a rational cocycle belongs to it if its image is a coboundary in $B_{\mathrm{cris}} \otimes V$. For the absolute Galois group of a finite extension $F/\mathbb{Q}_p$, and the usual Fontaine period ring, this is the inverse image in cocycles of the usual Bloch–Kato finite subspace. To be quite precise about continuity, the tensor relation gives every value of the cocycle in V as a difference $(g-1)y$ with $y \in B_{\mathrm{cris}} \otimes V$. Its components depend continuously in B_{dR} (the continuous action and the embedding of the crystalline ring are the usual period-ring facts). The components lie in the rational subspace, which has the induced p-adic topology, so the relation describes a continuous map to V. For a lattice, saturation under multiplication by a power of p follows immediately by rescaling the cobounding vector in $B_{\mathrm{cris}} \otimes V$. Thus the choice of this subset of cocycles instead of a quotient of cohomology causes no change of finite condition.

For a de Rham representation, the correct reference is S. Bloch and K. Kato, *L-functions and Tamagawa numbers of motives*, The Grothendieck Festschrift, vol. I, Birkhäuser, 1990, 333–400, Proposition 3.8 and Corollary 3.8.4. The paper is available at `https://math.stanford.edu/~conrad/BSDseminar/refs/BKTamagawa.pdf`; the exact sequences and hypotheses in these two statements have been checked. Corollary 3.8.4 asserts the classical exact sequence with terms

$$H^0(F, V), \quad D_{\mathrm{cris}}(V), \quad D_{\mathrm{cris}}(V) \oplus D_{\mathrm{dR}}(V)/\mathrm{Fil}^0, \quad H^1_f(F, V),$$

where the middle map is $(1 - \varphi, \mathrm{can})$. In the original corollary the term Fil^0 is kept in the source as a direct summand and then quotiented out; eliminating it gives precisely the

displayed sequence in Chapter 50. No assumption of dimension equality for $D_{\mathrm{cris}}(V)$ is used in this corollary; a *de Rham* representation is sufficient. Potentially crystalline suffices for de Rham here: after a finite extension the crystalline invariant basis maps to a de Rham invariant basis under $B_{\mathrm{cris}} \hookrightarrow B_{\mathrm{dR}}$, and finite Galois descent for the coefficient space of periods gives the de Rham property over the original local field. Equality between Hodge–Tate jumps and the graded de Rham filtration is the usual de Rham-to-Hodge–Tate comparison (e.g. Brinon–Conrad, *CMI summer school notes on p-adic Hodge theory*, Proposition 6.3.2, checked in the authors' version at `https://math.bu.edu/people/jsweinst/AWS/Files/BrinonConradPAdicHodgeTheory.pdf`). At the individual embeddings the sign convention can be checked on $\mathbb{Q}_p(j)$: its filtration jump is $-j$, and so is the weight defined by the local covariance of the semilinear maps in `HasHodgeTateWeight`.

Here is the passage to the lattice statement, recording the hypotheses that sometimes cause confusion. Let O be the integers of the coefficient field L_0, let T be free of rank r over O, and let $V = T[1/p]$. The exact sequence is linear over L_0, though the crystalline Frobenius is semilinear over the maximal unramified subfield of F. Thus Euler characteristic of the sequence gives

$$\dim_{L_0} H^1_f(F, V) = \dim_{L_0} H^0(F, V) + \dim_{L_0}(D_{\mathrm{dR}}(V)/\mathrm{Fil}^0).$$

The map $V \to B^1(F, V)$ has kernel $H^0(F, V)$, so the rational cocycle dimension is exactly r plus the last filtered quotient dimension. Every continuous rational cocycle on the compact Galois group becomes integral after multiplying by a sufficiently large power of p; this operation preserves the finite condition. Local cohomology finiteness for lattices and the coboundary quotient show finite generation over O. The cocycle module is torsion free as a submodule of the product of copies of the lattice, hence finite free. After passage to a splitting coefficient field, $F \otimes_{\mathbb{Q}_p} L_0 = \prod_\sigma L_0$, and its filtered quotient dimension is exactly the sum of the negative jump multiplicities. This proves the formula in Proposition 49.1, and in the applications the index $[\mathrm{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p) : G_F]$ is the local degree $[F : \mathbb{Q}_p]$. The group-level formulation in Lean allows a homomorphism onto an open Galois subgroup; for the local applications its action indeed factors through the subgroup. If an auxiliary abstract group is used with a potentially crystalline invariant spanning property, elements in the kernel act trivially on the representation because the period invariants span over B_{cris}; crystalline cocycles, being coboundaries after scalar extension, likewise vanish on that kernel. The same local argument then applies.

Proposition 3.8 of Bloch–Kato also states that $H^1_f(F, V)$ and $H^1_f(F, V^*(1))$ are exact mutual annihilators. In finite coefficients the cup pairing and the Euler–Poincaré order formula used in Lemma 50.2 are the usual local Tate duality and local Euler characteristic, e.g. J. S. Milne, *Arithmetic Duality Theorems*, 2nd ed., BookSurge, 2006, I, Corollary 2.3 and Theorem 2.8, checked in `https://math.stanford.edu/~conrad/BSDseminar/refs/MilneADT.pdf`. Characteristic zero of F makes their prime-to-char F hypothesis vacuous for finite p-primary modules. The trace pairing on O is used with a generator of the inverse different; this does not add a unit assumption about the ordinary trace. Passing from the rational orthogonality to integral cup products uses the torsion-free $H^2(F, \mathbb{Z}_p(1))$, with reduction giving the same pairing. For the finite cohomological Euler characteristic, $|O_F : mO_F| = m^{[F:\mathbb{Q}_p]}$ for m a power of p. The remaining rank and coboundary count is the purely finite calculation of Lemma 50.2. These results allow the classical local rank and exact-dual orthogonality interfaces to be treated as referenced and proved in the stated formulation.

They do *not* apply directly to lattice pairings only equivariant modulo p^a, to a congruence between unrelated crystalline lattices, or to the quantitative annihilation of such imperfectly dual cup products with a large range of Hodge weights. The indispensable structural and

quantitative period-ring comparison is a separate interface in Chapter 51. Citing the ordinary Bloch–Kato local condition cannot close it.

8.5 The integral characteristic-polynomial interface in Ribet's construction

The argument of Section 41.8 uses two integral forms of Amitsur's identity. We supply a proof valid over every commutative coefficient ring; in particular, passing to characteristic zero and applying Newton's identities to the particular coefficient algebra would not suffice.

Lemma 8.4 (An integral cyclic-word identity)**.** *Let $A_1, \dots, A_k$ be square matrices of equal size over a commutative ring L and let $t_1, \dots, t_k$ be central commuting formal variables. A word $w = i_1 \cdots i_d$ is primitive if it is not a proper positive power of a shorter word. Write $A_w = A_{i_1} \cdots A_{i_d}$ and $t^w = t_{i_1} \cdots t_{i_d}$. Then in $L[[t_1, \dots, t_k]]$,*

$$\det\left(1 - \sum_i t_i A_i\right) = \prod_{[w]\ \text{primitive}} \det(1 - t^w A_w), \tag{8.1}$$

where the product runs over cyclic rotation classes of primitive words, with any choice of representative for each class. Only finitely many factors contribute in any given total degree.

Proof. First take the matrices to have independent universal entries over $\mathbb{Z}$. Tensor this polynomial ring with $\mathbb{Q}$. Formal logarithms and the determinant derivative give $\log\det(1-U) = -\sum_{m\ge1} \operatorname{tr}(U^m)/m$ for a matrix U with entries of strictly positive total degree: introduce a further variable z and differentiate $\log\det(1 - zU)$, using $(1 - zU)^{-1} = \sum_{j\ge0} z^j U^j$, then integrate and set $z = 1$ degree by degree. In the expansion for $U = \sum_i t_i A_i$, each word of length m is a repetition of a primitive word of length $d \mid m$. A cyclic class of a primitive word of length d has exactly d distinct cyclic rotations, including when repeated m/d times; traces are invariant under those rotations. Hence the contribution of that class to the logarithm is

$$-\sum_{r\ge1} \frac{(t^w)^r \operatorname{tr}(A_w^r)}{r} = \log\det(1 - t^w A_w).$$

Summing proves equality of the two logarithms, and formal exponentiation proves the identity over the rational polynomial ring. Every coefficient of both sides before taking logarithms was an integral polynomial in the universal entries; thus their equality in the rational polynomial ring is their equality in the integral polynomial ring. Specializing the entries proves (8.1) over L, without any inverted integer. Rotation independence follows likewise from $\det(1 - XY) = \det(1 - YX)$ (use the two Schur-complement evaluations of $\det\begin{pmatrix} 1 & X \\ Y & 1 \end{pmatrix}$). This completes the proof. □

Proposition 8.5 (Characteristic coefficients on an integral spanning family)**.** *Let R, L be commutative rings, $R \to L$ a ring homomorphism, and $\rho : \mathscr{B} \to M_n(L)$ an R-algebra homomorphism, with $\mathscr{B}$ an associative unital R-algebra. Suppose $\mathscr{B}$ is spanned as an R-module by elements b for which all characteristic-polynomial coefficients of $\rho(b)$ belong to the image R' of R in L. Then the same is true of every $b \in \mathscr{B}$.*

Proof. Write $c_j(A)$ for the coefficient of z^j in $\det(1 - zA)$, with $c_0(A) = 1$. We prove $c_j(\rho(b)) \in R'$ simultaneously for all b, by induction on j. Express the current b as a finite R-linear combination of the stipulated generating elements b_i. Apply (8.1) to the $\rho(b_i)$

and substitute $t_i = zr_i$, where $r_i \in R'$ are the images of the coefficients in that linear combination. For a primitive word of length one every factor has coefficients in R' by the spanning hypothesis. For a primitive word of length $d \geq 2$, only coefficients $c_e(\rho(b_{i_1} \cdots b_{i_d}))$ with $de \leq j$ can contribute to the coefficient of z^j of the product. Apart from $e = 0$, these have $e < j$ and belong to R' by the induction hypothesis, whether or not the product word itself is one of the selected span generators. All scalar word monomials also belong to R'. The coefficient $c_j(\rho(b))$ therefore belongs to R', completing the induction. The coefficients of the characteristic polynomial are precisely those of this reversed determinant. □

Corollary 8.6 (Flattened block matrices)**.** *Under the hypotheses of Proposition 8.5, every characteristic coefficient of the image of a square block matrix over $\mathscr{B}$, flattened after applying ρ to its entries, belongs to R'. In particular this holds when $\mathscr{B} = R[G]$ is a group algebra and the given coefficient condition holds on the group elements.*

Proof. The block-matrix algebra is spanned by matrices $E_{ij}b$ with $b \in \mathscr{B}$. After flattening, the image of $E_{ij}b$ is $E_{ij} \otimes \rho(b)$. For $i = j$ its reversed characteristic polynomial is $\det(1 - z\rho(b))$; the other diagonal blocks of $1 - z(E_{ii} \otimes \rho(b))$ are identity matrices. For $i \neq j$ this reversed polynomial is 1, by block triangularity after permutation of the block labels. Thus these span generators satisfy the characteristic-coefficient hypothesis, and Proposition 8.5 applied to the flattened representation proves the assertion. For the group algebra itself, the group elements give the required module span. □

This is exactly the ring-theoretic generality of `amitsur_charpoly_mem_of_span` in `Lib/General/LinearAlgebra/PartialDerivComm.lean` and `amitsur_charpoly_mem_matrix` in `New/Selmer/RibetCramer.lean`. Apply it with $R = S$ the common subalgebra of coefficient functions on the eigensystem family and $L = \prod_{x \in Y} L_x$ the ambient coefficient ring. The group-algebra representation is obtained by taking the product of the individual representations and extending scalars to S. The hypothesis is precisely the family-valued characteristic-coefficient assumption used in Section 41.8, not an assertion that the individual entries of the representative matrices belong to S. Corollary 8.6 provides the characteristic coefficients needed for both polynomial adjugates in that section and therefore closes this integral algebra input. The arithmetic existence of the required family and of its relation matrices is a separate matter.

8.6 How to read the input labels in the preceding chapters

The volume retains several historical labels of the form *formal-source input* from the specialist-paper stage. Such a label need not indicate an independent gap: if the same assertion is proved at a later stage, the later proof is the appropriate justification. Conversely, a proof conditional on a geometric or integral compatibility hypothesis must retain that hypothesis as an open obligation. To avoid counting the same interface repeatedly, the accompanying source-anchor ledger records its position in the frozen 550-page version, which part of the mathematical argument it supports, and whether the earlier status is stale, conditional, a standard referenced theorem, or still open. A status of referenced means that both the theorem and its match to the application have been supplied; a citation found by title alone is not counted.

The foundational interfaces addressed specifically in this chapter have the following status.

Universal coefficient basis, colored semi-invariants and Schur expansion.
PROVED using the referenced all-characteristic Weyl character theorem, as in Theo-

rem 8.1 and Corollary 8.2; no arbitrary-ring flatness assertion is assumed. In particular the previous status of Proposition 13.15 is superseded.

Quadratic norm subgroup and splitting character.
REFERENCE for reciprocity and Hasse's theorem (the precise locations above); PROVED for the extra infinity type and local norm normalization in Chapter 6.

Classical CM reduction and degree.
REFERENCE for the three exact theorem statements in Silverman's books; PROVED for their application to the marked cubic and the first unit coefficient by the normalization and local-length arguments earlier. The class-group oddness argument is proved in Lemma 6.2.

Kronecker limit and unit-adjusted period.
PROVED for the normalized analytic limit above (with the checked modular-discriminant identity) and for the subsequent algebraic-unit adjustment in Proposition 12.11; the soft reciprocity theorem is proved in Chapter 16.

Classical finite local cohomology.
REFERENCE with exact proof bridges for the rank and exact-dual pairing in Proposition 49.1 and Lemma 50.2. Integral period-ring changes and imperfect duality remain separate obligations.

Integral Amitsur characteristic-polynomial input.
PROVED by the universal cyclic-word identity and its span and block consequences (Proposition 8.5 and Corollary 8.6). The assumption on the group-characteristic coefficient functions remains explicit.

These six items concern mathematics independent of the final quantitative analysis. The crucial lower-bound and spanning interfaces may also be split into smaller conventional theorems, but none of the conclusions of this chapter should be used to justify their remaining specialized hypotheses by analogy.

Chapter 9

The normalized real oscillator, global theta functional and mixed model

The finite part of the Weil construction was completed in Chapter 7, including the sign of the normalized finite cocycle. We now establish its real counterpart, the existence and uniqueness of the normalized adelic theta functional in the source's integrable model, the parabolic zero-term law, and rational invariance by doubling. These are mathematical proofs rather than appeals to an unspecified normalization of the Weil representation. We use ordinary Fourier duality, the Gaussian integral, and the uniqueness of the irreducible Heisenberg representation; the few places where a comparison with another geometric model is needed are indicated explicitly.

9.1 The real vacuum identity

Put $n = p + q$, $P = \mathrm{diag}(I_p, 0)$, $Q = I - P$, $S = P - Q$. The real unitary group is $G = \{g \in \mathrm{GL}_n(\mathbb{C}) : g^* S g = S\}$. Here and throughout this chapter $*$ denotes conjugate transpose. Use normalized Lebesgue measure $dx = \pi^{-n} d^{2n}x$, and define

$$e(x) = \exp(-x^* x/2), \qquad \beta(x, y) = \exp\big((y^* S x - x^* S y)/2\big).$$

These are exactly `realVac` and `realHeis` of `Lib/General/LinearAlgebra/HuaIntegral.lean`. The Heisenberg operators act on the reproducing space as

$$(\rho(a)v)(x) = \beta(a, x)\, v(x - a), \qquad \rho(a)\rho(b) = \beta(a, b)\rho(a + b).$$

In particular the sign in the translation is the one fixed in Chapter 7.

To check the vacuum normalization independently, define $z(x) = (\bar{x}_+, x_-)$, where the two blocks correspond to P and Q. Then z is a real-linear isometry of $\mathbb{C}^n$, and direct expansion gives

$$\rho(a)e(x) = e(x)e(a)\exp\big(z(a)^* z(x)\big). \tag{9.1}$$

Consequently the reproducing space at the real place consists, after completion, of the functions $e(x)F(z(x))$ with F in the usual holomorphic Fock space. The normalized pairing of monomials is $\langle z^I, z^J\rangle = \delta_{I,J} I!$. This follows by radial and angular integration in each complex coordinate, as in Chapter 7.6. The exponential series yields its reproducing kernel. The following stronger identity will be useful for the adelic Lagrangian-free construction.

Lemma 9.1 (An exact continuous rank-one projector)**.** *The real operators satisfy*

$$\int e(a)\rho(a)\rho(b)e\,da = e(b)e. \tag{9.2}$$

The same identity holds for the normalized finite vacuum of Chapter 7 and for products of these vacua.

Proof. In the real case write $u = z(a), v = z(b), w = z(x)$. Since z preserves measure, the integral at x is, by (9.1),

$$e(x)e(b)\exp(v^*w)\int_{\mathbb{C}^n}\exp(-u^*u - v_{!\mathrm{tr}}^*u + u^*w)\,du,$$

Indeed the cross term from $e(a+b)$ is $-\Re(u^*v)$, while $\beta(a,b)$ supplies $i\Im(u^*v)$; their sum is $-v^*u$. The one-dimensional Gaussian identity

$$\pi^{-1}\int_{\mathbb{C}} e^{-|u|^2+Au+B\bar{u}}\,d^2u = e^{AB}$$

is obtained by expanding the exponentials and integrating the monomials, or by real Gaussian integration followed by holomorphic continuation in A, B. In n coordinates this gives $\exp(-v^*w)$ and proves the assertion. All integrals are absolutely convergent by Gaussian decay.

For the finite lattice let L and λ be as in Section 7.3. On L, $\lambda^2 = 1$, and the operators $\lambda(a)\rho(a)$ form an ordinary unitary representation of L. Its normalized average over L is the projection onto the unique vacuum line, by the eigenrelations and self-dual annihilator computation of Section 7.4. The projection of $\rho(b)e$ onto this line is $e(b)e$, including $b \notin L$, when the two cosets have disjoint support. This gives the same integral identity with $e = \lambda\mathbf{1}_L$. The identity for products follows from Fubini. □

This is exactly the twisted-convolution condition in `Heisenberg.IsVacuum` of `Lib/General/Groups/ValuationPair.lean`. In particular, neither the real nor the finite part requires an unexplained normalization of the reproducing kernel.

9.2 The real phase from a Gaussian determinant

For $g \in G$ set $T(g) = P + Qg$ and $\Delta_\infty(g) = \overline{\det T(g)}$. If $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in the signature decomposition, then $\det T(g) = \det d$. The block d is invertible because $d^*d - b^*b = I_q$. Also $g^{-1} = Sg^*S$, so the negative block of g^{-1} is d^*. Hence $\Delta_\infty(g^{-1}) = \overline{\Delta_\infty(g)}$. This is the source's `realMultiplier`, not an arbitrary choice of the square root of a determinant.

We need the ordinary complex Gaussian formula in a slightly general form. A complex matrix A is accretive if $(A + A^*)/2$ is positive definite. Then

$$\int_{\mathbb{C}^m} e^{-x^*Ax}\,dx = (\det A)^{-1}. \tag{9.3}$$

For a positive definite Hermitian A this is the spectral theorem and the one-coordinate Gaussian integral. For any accretive $A = H + K$ with H Hermitian positive definite and K skew-Hermitian, both sides for $H + tK$ are holomorphic in t on the convex open set where the matrix is accretive, by locally uniform Gaussian domination. For imaginary t near zero the matrix is positive Hermitian. The one-variable identity theorem gives the formula at $t = 1$. Accretivity implies nonsingularity since x^*Ax has positive real part for $x \neq 0$. Thus no branch of a square root is involved.

Theorem 9.2 (Exact real multiplier cocycle)**.** *For $g, h \in G$, writing $k = gh$, one has*

$$\Delta_\infty(g)\Delta_\infty(h)\Delta_\infty(k^{-1}) \iint e(hx)e(ky)\beta(x,y)e(x+y)\,dx\,dy = 1. \tag{9.4}$$

Consequently the real Weil operator on the source reproducing space is

$$\omega(g)v = \Delta_\infty(g) \int v(y)\rho(gy)e\,dy$$

and these operators form a genuine representation of G.

Proof. In the double integral the exponent is $-(x,y)^* A(x,y)$ for

$$A = \begin{pmatrix} (h^*h + I)/2 & P \\ Q & (k^*k + I)/2 \end{pmatrix}.$$

Indeed the terms from $e(x+y)$ and $\beta(x,y)$ give the off-diagonal entries $(I+S)/2 = P$ and $(I-S)/2 = Q$. The real part of the exponent is $-(\|hx\|^2 + \|ky\|^2 + \|x+y\|^2)/2$, so A is accretive. We compute its determinant explicitly. Set

$$H = P + Qh, \quad K' = Pk + Q, \quad G' = Pk + Qh.$$

The factorization is

$$A = \begin{pmatrix} H & Q \\ 0 & K' \end{pmatrix}^* \begin{pmatrix} P & -Q \\ I & I \end{pmatrix} \begin{pmatrix} I & I \\ -H & -H + G' \end{pmatrix}. \tag{9.5}$$

For clarity, multiplication of the last two factors gives $\begin{pmatrix} H & P \\ Q(I-h) & K' \end{pmatrix}$. Multiplying by the first yields the diagonal blocks $H^*H = P + h^*Qh = (I + h^*h)/2$ and $(K')^*K' = k^*Pk + Q = (I + k^*k)/2$, using $h^*Sh = S$ and $k^*Sk = S$. The off-diagonal entries are $H^*P = P$ and

$$QH + (K')^*Q(I-h) = Qh + Q(I-h) = Q.$$

This verifies (9.5) entry by entry.

The determinant of the central factor is $\det(P+Q) = 1$, and that of the last factor is $\det G'$. Since $K' = T(k^{-1})k$ and $G' = T(g^{-1})k$, and $|\det k| = 1$ by the unitary equation, we find

$$\begin{aligned} \det A &= \overline{\det T(h) \det T(k^{-1}) \det k}\, \det T(g^{-1}) \det k \\ &= \Delta_\infty(h)\Delta_\infty(g)\,\Delta_\infty(k^{-1}). \end{aligned}$$

Substituting (9.3) proves (9.4).

For the representation assertion write $I_g v = \int v(y)\rho(gy)e\,dy$. This is bounded in the integrable model by the fixed L^1 norm of e, and a change of variables proves covariance $I_g\rho(a) = \rho(ga)I_g$. In particular

$$I_{k^{-1}} I_g I_h v = \int v(y)\rho(y)w\,dy, qquadw = I_{k^{-1}} I_g e.$$

In fact expand $I_h v$ first and move I_g and $I_{k^{-1}}$ past the coherent translations. The rank-one integration identity

$$\int v(y)\rho(y)w\,dy = w(0)v$$

holds on the coherent span and extends by boundedness. For $v = e$ this is Lemma 9.1; for $v = \rho(a)e$ substitute $y = t + a$ in the integral, obtaining $\rho(a) \int e(t)\rho(t)w\,dt$. Density of the coherent span gives it for all v. Thus the triple operator equals $w(0)$ times the identity. Expanding the remaining two integrals explicitly, and using evaluation at zero, yields

$$w(0) = \iint e(k^{-1}z)e(u)\beta(gu, z)e(z - gu)\,du\,dz.$$

Make the transformations $u = hx$, $z = k(x + y)$; both g, h and k preserve measure. The integrand becomes $e(x + y)e(hx)\beta(x, y)e(ky)$, since $gu = kx$ and $\beta(kx, k(x + y)) = \beta(x, y)$. This is exactly the double integral above. All changes and exchanges are justified by absolute Gaussian convergence. Multiplying by the three multipliers and using (9.4) gives $\omega(k^{-1})\omega(g)\omega(h) = 1$. Taking $g = h = 1$ is consistent with the reproducing operator $I_1 = 1$. Taking $h = 1$ gives inverses and the general formula the group relation.

The action preserves the Hilbert inner product. Indeed $\langle I_g v, w\rangle = \int \overline{v(y)}w(gy)dy$ by reproduction, and the analogous formula shows $I_g^* = I_{g^{-1}}$ on the integrable model. Since $\Delta(g^{-1}) = \overline{\Delta(g)}$, the normalized operator has adjoint equal to its inverse. Finally $[\omega(g)e](0) = \Delta(g) \int e(y)e(gy)dy$. The Gaussian formula applies to $(g^*g + I)/2 = \overline{T(g)}^*T(g)$, where the equality was verified above with $h = g$. Thus the overlap is exactly $1/\overline{\Delta(g)}$. □

Theorem 9.2 reconstructs `realAction` in `Lib/Weil/Fock/FockAction.lean`. This determinant factorization is the actual one used in the compact source, expressed in ordinary matrix notation. Together with Theorem 7.6 it closes the algebraic cocycle assertion of `adelicAction` in `Lib/Weil/Construction/ThetaFunctional.lean`: the adelic multiplier is the product of the local ones, and the vacuum measure is the product of the real Gaussian normalization and the finite lattice normalization after rescaling by `scaleIdele`. At all but finitely many finite places the multiplier is one. Products and the changes of variables are thus legitimate.

9.3 Adelic duality and summable vacuum correlations

We now address the global theta functional in the actual nonpolar model. Let V be a rational nonsingular Hermitian space equipped with the adelic frame of Chapter 6. In its framed coordinates write $W = \mathbb{A}_E^n$, $\Gamma = x^{-1}E^n$, and use the bicharacter

$$\beta(u, v) = \psi\big(\jmath h(v, u)/2\big),$$

where ψ is the standard full additive character of E (trace to the rational character), and h is the framed form. The character is trivial on rational scalars. It follows that $\beta(\gamma, \delta) = 1$ for $\gamma, \delta \in \Gamma$. The annihilator of Γ with respect to β^2 is exactly Γ.

We recall an elementary argument for the latter assertion. The additive character of $\mathbb{A}_E$ gives a perfect pairing with itself, and the annihilator of E is E. For the second statement, subtract from an adele in the annihilator a rational element so that its finite part belongs to the local trace-dual lattice $\widehat{\mathfrak{d}_E^{-1}}$; this is possible by additive weak approximation at finitely many denominator places, or the Chinese remainder theorem after clearing denominators. Testing the character against $\mathcal{O}_E$ forces its real part into the Euclidean trace-dual lattice, which equals the real embedding of $\mathfrak{d}_E^{-1}$ by a basis and its dual. Subtract that rational element as well. The remaining adele has zero real part and trivial finite character pairing against E. Since E is dense in $\mathbb{A}_{E,f}$ additively (again the Chinese remainder theorem), nondegeneracy of the local trace pairings makes the finite part zero. The full pairing is perfect by local self-duality and restricted products: the continuous character is trivial on the given integral

factors outside a finite set and is represented by a finite adele there, together with its real linear functional. Applying this coordinatewise to the invertible rational matrix of the pairing $\mathfrak{j}h$ and transporting by x proves the framed assertion. Similarly W/Γ is compact: for the standard rational lattice, first reduce the finite part into $\widehat{\mathcal{O}}_E^n$, then use a real fundamental parallelepiped for $\mathcal{O}_E^n$. Its character group is indexed by Γ via $a \mapsto \beta(\cdot, a)^2$.

The normalized product vacuum e on W has absolute value a positive-definite real Gaussian times the indicator of a compact open finite lattice. We use the following estimate, valid without any uniformity in the family of ranks and fields (none is needed in this step):

$$C_V := \sup_{a \in W} \sum_{\gamma \in \Gamma} |e(a+\gamma)| < \infty. \tag{9.6}$$

To prove this, fix the finite part of a. If no rational point contributes then the sum is zero. Otherwise choose one contributing point γ_0. All other contributing points differ from it by the rational points whose finite coordinates lie in the fixed finite lattice. These form, under the real embedding, a full Euclidean lattice: in the rational coordinates they lie between two fixed fractional integral lattices, since the adelic frame has integral invertible finite part away from finitely many places. Now the sum is bounded by the supremum of the periodized Gaussian over translations of that Euclidean lattice. This supremum is finite on a compact fundamental parallelepiped by uniform convergence; the number of lattice vectors in a ball of radius R is bounded by a constant times $(1+R)^{[E:\mathbb{Q}]n}$ by disjoint small balls. That proves (9.6).

9.4 Existence, uniqueness and normalization of theta

Let $\mathscr{S}_e(W)$ denote the integrable vacuum-coefficient model of the source, namely the continuous integrable functions v with

$$v(x) = \int_W v(y)\beta(y,x)e(x-y)\,dy = \left(\int_W v(y)\rho(y)e\,dy\right)(x). \tag{9.7}$$

It has the L^1 norm, and the Hilbert inner product inside its L^2 completion is conjugate-linear in the first variable. Since $|e| \le 1$, the point evaluations are bounded by the L^1 norm. The identity also shows that the span of the coherent vectors is dense in this norm: truncate the integral on a compact set, then approximate the continuous Banach-valued integrand by finite sums, using continuity of translations in L^1. The image of the kernel projector is closed in L^1 by its bounded integral kernel. These elementary facts justify the Bochner integrals on the model. We have

$$\langle \rho(y)e, v\rangle = v(y), \qquad P_e v := \int e(a)\rho(a)v\,da = v(0)e$$

by (9.7) and Lemma 9.1. All statements with P_e extend from the coherent span by boundedness.

Call a bounded linear functional ℓ on $\mathscr{S}_e(W)$ rationally invariant if $\ell(\rho(\gamma)v) = \ell(v)$ for $\gamma \in \Gamma$. Give W/Γ probability Haar measure. We prove the exact source claim.

Theorem 9.3 (The global theta functional)**.** *There is a bounded rationally invariant linear functional θ with*

$$\int_{W/\Gamma} |\theta(\rho(a)e)|^2\,d\bar{a} = 1.$$

The space of bounded rationally invariant linear functionals is one-dimensional. If θ is so

normalized, then for all v, w in the model,

$$\int_{W/\Gamma} \theta(\rho(a)v)\overline{\theta(\rho(a)w)}\, d\bar{a} = \langle w, v\rangle, \tag{9.8}$$

$$\theta(v)\overline{\theta(w)} = \sum_{\gamma\in\Gamma} \langle w, \rho(\gamma)v\rangle. \tag{9.9}$$

In particular $v \mapsto (a \mapsto \theta(\rho(a)v))$ is injective and isometric for the Hilbert norm and the quotient mean.

Proof. For every u, v in the model,

$$\sum_{\gamma\in\Gamma} |\langle u, \rho(\gamma)v\rangle| \le C_V \|u\|_1 \|v\|_1. \tag{9.10}$$

Indeed use (9.7) for u, v, expand the inner product into a double integral of coherent vectors and invoke Tonelli on absolute values. The coherent correlation has absolute value $|e(\gamma + a - b)|$ because of the Heisenberg product law. Estimate (9.6) gives the bound. Thus $\ell_u(v) = \sum_\gamma \langle u, \rho(\gamma)v\rangle$ is a bounded linear functional; it is rationally invariant because the bicharacter is trivial on $\Gamma \times \Gamma$ and the sum is absolutely convergent. Some ℓ_u is nonzero. Otherwise for every $a \in W$ take $u = v = \rho(a)e$; the sum would vanish, but it is the Fourier sum $\sum_\gamma e(\gamma)\beta(a,\gamma)^{\pm 2}$ (the sign has no effect on the constant term). Its quotient mean equals $e(0) = 1$ by orthogonality and absolute convergence, a contradiction.

Let ℓ, ℓ' be two bounded invariant functionals. For v, w define

$$F_{v,w}(a) = \ell(\rho(a)v)\overline{\ell'(\rho(a)w)}.$$

It is a continuous periodic function on W/Γ, since the two scalar factors from translating by a rational point cancel. Its mean is a bounded sesquilinear form invariant under every $\rho(a)$ by translation invariance of Haar measure. Every such form is a scalar multiple of the Hilbert inner product: if B is conjugate-linear in its first argument, invariance implies $B(\rho(a)v, w) = B(v, \rho(-a)w)$, whence

$$B(P_e v, w) = B(v, P_e w).$$

Here $e(-a) = e(a)$ is real. Thus $B(e, w) = B(e, P_e w) = w(0)B(e, e)$, and $B(\rho(a)e, w) = w(a)B(e, e)$. Integrate the coherent expansion for the first argument as in (9.7) to obtain $B(v, w) = B(e, e)\langle v, w\rangle$. In the convention of F this proves that its mean is $c\langle w, v\rangle$, with c the mean for $v = w = e$.

For nonzero ℓ take $\ell' = \ell$. The number c is positive: the coherent span is dense and the continuous function $\ell(\rho(a)e)$ is not identically zero. Rescaling by $c^{-1/2}$ gives a θ with $c = 1$ and proves (9.8).

For arbitrary ℓ, ℓ' use now Fourier inversion at zero for $F_{v,w}$. Its Fourier coefficients, up to replacing γ by $-\gamma$, are $c\langle w, \rho(\gamma)v\rangle$. Indeed the identity

$$\rho(a)\rho(\gamma) = \beta(a,\gamma)^2 \rho(\gamma)\rho(a)$$

and rational invariance identify multiplying $F_{v,w}$ by the character $\beta(a,\gamma)^2$ with $F_{\rho(\gamma)v,w}$. The coefficient is its previously calculated mean. By (9.10) the coefficients are absolutely summable, so the series converges uniformly and equals $F_{v,w}$ by uniqueness of Fourier coefficients of continuous functions on the compact abelian group. At zero this gives

$$\ell(v)\overline{\ell'(w)} = c \sum_\gamma \langle w, \rho(\gamma)v\rangle.$$

Specializing to θ proves (9.9). For any invariant ℓ choose w with $\theta(w) \neq 0$ and take $\ell' = \theta$. Comparison with (9.9) gives $\ell(v) = c\theta(v)$ for all v. This proves uniqueness. Injectivity and the norm assertion follow immediately from (9.8). □

This is a direct proof of `exists_isThetaFunctional`, `IsRatInvariant.quotMean_eq`, `IsRatInvariant.hasSum_modelInner`, and `IsThetaFunctional.eq_smul` in `Lib/Weil/Construction/ThetaFunctional.lean`. The sum used to produce an invariant functional need not equal a naively chosen rational theta sum on a nonpolar wave function; the proof constructs it within the exact integrable model and normalizes it by probability quotient mean.

Corollary 9.4 (Compatibility with orthogonal sum)**.** *If $W = W_1 \oplus W_2$ with rational lattices and vacuum data orthogonal products, normalized theta functionals satisfy*

$$\theta_W(v_1 \otimes v_2) = \zeta\theta_{W_1}(v_1)\theta_{W_2}(v_2), \qquad |\zeta| = 1.$$

The scalar is independent of the vectors. Moreover the normalized Weil operator of a block diagonal element is the tensor product of the two normalized operators.

Proof. The product functional on the coherent span extends continuously to the integrable model by

$$v \longmapsto \int v(a_1, a_2)\theta_{W_1}(\rho_1(a_1)e_1)\theta_{W_2}(\rho_2(a_2)e_2)\, da_1\, da_2.$$

Indeed the last two factors are uniformly bounded by the respective functional norms times the fixed vacuum L^1 norms. The reproducing identity shows that the integral equals the product on elementary tensors. Rational invariance follows on the dense coherent span, and the product quotient mean is one by Fubini. The theorem yields the scalar. For the operators the defining integral and multiplier factor over the two blocks (the finite block product is Proposition 7.2, and the real block determinants multiply). This proves the second assertion. □

9.5 The normalized hyperbolic Schrödinger model

Here is the remaining geometric comparison needed to calculate the exact parabolic scalar. Write a rational splitting of V as

$$V = lE^a \oplus V_0 \oplus kE^a, \qquad h(kb, lt) = b^*t,$$

with isotropic frames l, k, mutually paired by the identity, and residual form J_0. More explicitly $h(ku, lv) = u^*v$, so the Gram matrix on the two isotropic blocks is $\left(\begin{smallmatrix} 0 & I \\ I & 0 \end{smallmatrix}\right)$. We use the mixed Schrödinger form of the Heisenberg operator

$$R(lt + w + kb)F(x) = \psi\big(\jmath b^*(2x + t)/2\big)\rho_0(w)F(x + t), \qquad x \in \mathbb{A}_E^a. \tag{9.11}$$

Multiplying translations and multiplications verifies the Heisenberg relation directly; the w part is orthogonal to the first two. This is the formula `isMixedLaw_mixedTerm` in `Lib/Weil/MixedModel/ZeroTerm.lean`.

We first check the identification for an adapted splitting. At real places use the orthonormal basis of the hyperbolic Gram matrix given by the columns of

$$X_\infty = 2^{-1/2}\begin{pmatrix} I & -I \\ I & I \end{pmatrix}.$$

At finite places, the extension E/F is unramified and the hyperbolic integral form is isometric to the standard unimodular form: the local Gram–Schmidt proof in Chapter 6 supplies such an integral invertible matrix, including at two. For the finite additive normalization the form on $\alpha\widehat{\mathcal{O}}$ is self-dual, and scalar multiplication by α commutes with the Hermitian matrices. Thus in the hyperbolic coordinates the two halves of the normalized vacuum lattice are $\alpha\widehat{\mathcal{O}}^a$ in each block. The quadratic refinement on their product is trivial on each individual isotropic half and equals $\beta(lt, kb)$ on their sum, by Lemma 7.3.

Choose on the translation coordinate the measure for which the Schrödinger Fourier pairing is self-dual. With this choice the normalized wave vacuum is the product of a positive Gaussian on the real component and the indicator of $\alpha\widehat{\mathcal{O}}^a$ on the finite component, times the residual vacuum. Its constant factor is chosen so its Hilbert norm is one and its value at zero is positive on the hyperbolic factor. One can verify that its Heisenberg coefficient is exactly e. In real scaled orthonormal coordinates the character in the Heisenberg wave action is $\exp(2i\operatorname{Im}(b^*(x + t/2)))$ (a change of the isotropic sign replaces b by $-b$). That coefficient is the integral of the translate of $C\exp(-\|x\|^2)$ against itself with a unit-modulus phase. Completing the square shifts x by $t/2$, and the real Fourier transform of the Gaussian supplies

$$\exp(-\|t\|^2/2)\exp(-\|b\|^2/2);$$

the mixed term of the character cancels under that shift. In finite coordinates, translation has disjoint supports unless t is in the isotropic lattice. If t is integral, character orthogonality gives zero unless b is in its trace-dual lattice, which here is the same scaled lattice in the other half. In that case the value is precisely $\psi(\jmath b^* t/2)$, as required by the refinement. The residual coefficient is its vacuum. This verifies the equality of all vacuum coefficients.

Accordingly the unitary comparison can be constructed without choosing a phase of a Stone–von Neumann intertwiner: send every coherent vector $\rho(w)e$ of the nonpolar model to $R(w)$ of this normalized wave vacuum. All pairwise inner products agree by the coefficient calculation and the Heisenberg relation. The spans are dense in the respective Hilbert models. For the wave model the density follows from the irreducibility of translations and multiplications, or directly from Fourier uniqueness (a function orthogonal to every modulation of every translate of the vacuum vanishes, since the translates cover the finite variable and the Gaussian never vanishes at infinity). The comparison is thus an isometry. Its extension to evaluations of mixed functions at a fixed x in the integrable model is bounded: apply the coherent integral (9.7); on coherent wave vectors the hyperbolic factor is a translate of the bounded wave vacuum and the residual factor has fixed integrable norm. This observation justifies the use of the mixed zero map on the actual source normed model. For a general framed splitting, compare its hyperbolic and residual bases to the chosen adapted local basis by an adelic unitary isometry. Such a comparison exists because both Gram matrices agree, and its coordinates are integral at almost all places. Conjugation of the constructions by the exact normalized Weil action gives the same mixed Heisenberg formula. This amounts to composing the adapted wave map with a fixed bounded Weil operator on the integrable model. Multiplying the comparison map by a nonzero scalar does not change any of the covariance identities below.

9.6 The zero-term character on the actual parabolic

Let g preserve the isotropic space $k\mathbb{A}_E^a$ and act there by $a \in \mathrm{GL}_a(\mathbb{A}_E)$ (we use a also for the matrix, with its size understood in this paragraph). Its induced action on $k^\perp/k$ is g_0. We claim the exact coefficient from the source:

Theorem 9.5 (Zero-term law)**.** *For the mixed evaluation at zero, denoted* ev_0*, one has*

$$\mathrm{ev}_0\,\omega(g) = \upsilon(\det a)^{-1}|\det a|_{\mathbb{A}_E}^{1/2}\omega_0(g_0)\,\mathrm{ev}_0\,.$$

Here υ is the splitting character of Chapter 6, and ω_0 uses the identical residual vacuum convention. The same law holds for any nonzero scalar multiple of the mixed evaluation map.

Proof. First consider the Levi element $L(a, g_0) = \operatorname{diag}(a^{-*}, g_0, a)$ in the ordered splitting. In the wave model

$$F(x) \longmapsto |\det a|^{1/2}\omega_0(g_0)F(a^*x)$$

is a unitary intertwiner: the measure of the translation variable changes by $|\det a|$, and (9.11) verifies covariance. It differs from the exact normalized operator on the hyperbolic factor by a phase, determined by the vacuum overlap. Since the real wave vacuum is positive and its finite part is an indicator, the overlap of its dilate with itself is strictly positive: the finite intersection contains a compact open neighborhood of zero and the real Gaussian integral is positive.

Compute the phase of the required hyperbolic multiplier. At infinity the negative block in the coordinates X_∞ is $(a^{-*} + a)/2$; factor out a^{-*}, leaving the positive definite matrix $(I + a^*a)/2$. Hence $\overline{\det((a^{-*} + a)/2)}$ has phase equal to the inverse phase of $\det a$. At the finite part the hyperbolic frame is integral invertible, so bi-integral invariance and Proposition 7.2 give

$$\Delta_f(L(a, 1)) = \Delta_f(a^{-*})\Delta_f(a) = |\Delta_f(a)|^2\overline{\chi_f(\det a)}.$$

Its phase is again $\upsilon_f(\det a)^{-1}$. Together the two parts show $\Delta(L(a, 1))\upsilon(\det a) > 0$. The normalized overlap is $1/\overline{\Delta}$, with that same phase. Consequently the exact Levi action is the displayed unitary intertwiner multiplied by $\upsilon(\det a)^{-1}$. Tensoring with the residual operator and evaluating at zero proves the formula for the Levi.

For the unipotent radical describe it concretely: its elements $u(C, D)$ fix the k block, send lt to $lt + Ct + kDt$, and send w to $w - kC^*J_0w$. Hermitian invariance is equivalent to $D + D^* + C^*J_0C = 0$. Thus $B = C^*J_0C$ and $Z = D + B/2$ satisfy $Z^* = -Z$. The parameters C, Z describe the entire radical, by block multiplication. Define the wave-model operator

$$U(C, Z)F(x) = \psi(\jmath x^*Zx/2)\rho_0(-Cx)F(x).$$

It is unitary and fixes evaluation at zero. It intertwines the Heisenberg operators with their transforms by $u(C, D)$. For the k translations (multiplications) this is immediate. For $\rho_0(w)$ it is the identity obtained by commuting $\rho_0(-Cx)$ past $\rho_0(w)$, giving the factor $\psi(-\jmath w^*J_0Cx)$ prescribed by the new k coordinate. For the l translations the ratio of quadratic phases at $x + t$ and x is

$$\psi\big(\jmath(t^*Zx + t^*Zt/2)\big),$$

since ψ is invariant under conjugation and $\bar{\jmath} = -\jmath$. The residual products have bicharacter $\beta(Ct, -C(x + t))$, and substituting $D = Z - B/2$ makes the two sides of (9.11) identical. This proves covariance on generators and hence on all Heisenberg elements.

These operators form a representation of the radical: the product and the operator for the product group element are scalar multiples by irreducible Heisenberg uniqueness, and both preserve evaluation at zero, so the scalar is one. Similarly conjugating by any Levi intertwiner preserves their zero-evaluation normalization; it gives exactly the operator for the conjugated radical element. The exact normalized Weil representation thus differs on the radical from $U(C, Z)$ by a character η invariant under Levi conjugations. Conjugation by the Levi of tI, t a nonzero rational scalar, scales C by t and Z by t^2. On the central subgroup $C = 0$ the group law is addition in Z; taking $t = 2$ shows $\eta(0, 3Z) = 1$ for every Z. Multiplication by 3 is surjective on skew-Hermitian adelic matrices. Therefore the character is trivial on that subgroup. The quotient of the radical by it is the additive group of C matrices. The same argument with $2 - 1 = 1$ shows that the character is trivial on the quotient. This proves the normalized operator is $U(C, Z)$ and its evaluation scalar is one.

Finally the stabilizer is the semidirect product of the described radical and the Levi, by the elementary block equations for a unitary matrix preserving k. The result follows by multiplication. The covariance extends under the bounded evaluation map to the integrable model. For a nonadapted splitting, conjugate the entire calculation by the adapted frame isometries; both the Heisenberg formula and the Levi determinant are preserved in the specified k coordinates. Hence the same scalar holds for the general case. □

This is `zeroTerm_omega` in `Lib/Weil/MixedModel/ZeroTermLaw.lean`, once the mixed evaluation is identified with the source zero term. In the preceding proof the radical normalization was established by scalar conjugation rather than by assuming the phase of a unipotent element. The determinant factor $\chi_{\mathbb{A}}(\det a)^{-1}$ is exactly the source's $\upsilon(\det a)^{-1}|\det a|^{1/2}$.

9.7 The mixed-model expansion and equality with the source zero term

We give the remaining functional comparison, since the representation alone does not imply that the source's particular zero term is wave evaluation. Use a rational splitting; equation (9.11) holds in its rational coordinates, so the rational subgroup acts by translations and modulations on the hyperbolic factor, and rational Heisenberg translations on V_0. In this model consider the functional

$$F \longmapsto \sum_{x \in E^a} \theta_{V_0}(F(x)).$$

In an adapted model it is well-defined and bounded on the integrable model: for each coherent vector its wave support is a finite-adelic lattice translate times a translated real Gaussian, with residual factor a translate of the residual vacuum. The supremum of the corresponding sum of absolute values is finite by exactly the lattice-count/Gaussian argument of (9.6); use the boundedness of θ_{V_0} and then integrate the coherent expansion (9.7). The mixed expansion is consequently absolutely summable for every integrable vector. For a nonadapted splitting apply the same bound after the fixed bounded Weil operator just described; in particular no Gaussian support assertion for a nonadapted vacuum is required. This functional is invariant under rational translations in the first isotropic block (permute summands), rational modulations in the other (the full additive character on rational arguments is trivial), and the residual rational subgroup. Hence it is rationally invariant. It is nonzero: one may choose in the wave model a hyperbolic Gaussian packet with sufficiently small finite support and a residual vector of nonzero theta, or take a finite-adelic characteristic function and a real Gaussian with all rational contributions nonnegative. The product is in the integrable model, as the Gaussian and compactly supported locally constant functions have integrable coherent coefficients. By Theorem 9.3 this functional is a nonzero scalar multiple of θ_V. Rescaling the mixed comparison map by a common constant yields

$$\theta_V(v) = \sum_{x \in E^a} \theta_{V_0}(F_v(x)). \tag{9.12}$$

This choice of scalar also identifies the source zero-term map. Indeed integrate $\theta_V(\rho(\iota y + kb)v)$ against probability Haar measure on $\mathbb{A}_E^a/E^a$ in b. Formula (9.12), absolute summability, and orthogonality of characters give

$$\int_{\mathbb{A}_E^a/E^a} \theta_V(\rho(\iota y + kb)v)\, d\bar{b} = \theta_{V_0}(\rho_0(y)F_v(0)).$$

As a function of y, the right side is the source theta transform of the residual vector $F_v(0)$. The transform is injective by (9.8). Accordingly the uniquely defined residual zero-term vector

of the source is exactly $F_v(0)$. In particular this argument proves its existence, not just compatibility of two hypothetical evaluation rules. It also gives the source mixed-term map at x by translating with $\rho(lx)$, hence $v[x] = F_v(x)$. The representation identity of Theorem 9.5 therefore holds for the actual source zero term.

This proves both the formula `zeroTerm_apply` (read on the theta transform) and the assertion `hasSum_theta_mixedTerm` of `Lib/Weil/MixedModel/ZeroTerm.lean`. If the isotropic frame is changed by a rational invertible matrix, the probability quotient mean is unchanged under that linear map, since it is an automorphism of the compact group. If an isotropic direction is removed in two steps, the two means compose by Fubini. These observations give `zeroTerm_congr` and `zeroTerm_comp` in `Lib/Weil/MixedModel/AdaptedSplitting.lean`, for the same isotropic subspace and fixed residual identification. They are useful when different rational hyperbolic bases occur in the proof.

9.8 Rational invariance in the exact nonpolar representation

It remains to prove that rational unitary elements, not merely rational Heisenberg translations, fix the normalized theta functional. If a rational Hermitian space is hyperbolic and the element g preserves a rational maximal isotropic subspace, the claim follows at once from the mixed expansion and the zero-term law. Indeed the action on the full mixed wave model is

$$F(x) \longmapsto \upsilon(\det a)^{-1} |\det a|^{1/2} \psi(\text{rational quadratic expression in } x) F(a^* x),$$

by the Levi and radical calculations. For a rational g, the matrix a and radical coordinates are rational over E (use the rational splitting); $\upsilon(\det a) = 1$ by principality, and $|\det a|_{\mathbb{A}_E} = 1$ by the product formula. The phase on rational x is one. The sum in (9.12) is unchanged, as a^* permutes the rational points. This proves the rational stabilizer result `theta_omega_ratEmbed_of_mul_k` in `Lib/Weil/DualPair/PairSplitting.lean`.

Theorem 9.6 (Rational theta invariance by doubling)**.** *For any rational nonsingular Hermitian space V with the prescribed frames and every $\gamma \in U(V)(F)$,*

$$\theta_V(\omega(\gamma)v) = \theta_V(v).$$

Proof. Let $W = V \oplus (-V)$, a split hyperbolic Hermitian space. Its diagonal graph $L_0 = \{(v, v)\}$ and the graph $L_1 = \{(\gamma v, v)\}$ are maximal isotropic, with $(\gamma, 1)L_0 = L_1$. Choose a norm-one scalar $z \in E^\times$ with $z\bar{z} = 1$ that is not an eigenvalue of either 1 or γ. There are infinitely many norm-one scalars, for instance $(t + \sqrt{-d})/(t - \sqrt{-d})$ for rational t; only finitely many values are forbidden. The graph $L_2 = \{(zv, v)\}$ is then transverse to both L_0, L_1.

There is a rational unitary map u fixing L_2 pointwise and sending L_1 to L_0. To verify this, identify W with $L_0 \oplus L_2$, with the two isotropic blocks perfectly paired. Since L_1 is transverse to L_2, it is the graph of a map $B : L_0 \to L_2$. Isotropy of the graph means $B + B^* = 0$ after identification of the pairing with the standard hyperbolic one. The shear $(x, y) \mapsto (x, y - Bx)$ is unitary and has the desired properties. Thus u preserves L_2, and $u(\gamma, 1)$ preserves L_0. Both fix θ_W by the rational stabilizer argument. Since the normalized Weil operators are a genuine representation, $(\gamma, 1)$ fixes it as well.

Apply this to the tensor of v and any v' with $\theta_{-V}(v') \neq 0$. By Corollary 9.4, $\omega_W(\gamma, 1)$ acts on the tensor by $\omega_V(\gamma) \otimes 1$ and the theta functional is a nonzero common phase times the product. Cancel that phase and the nonzero factor $\theta_{-V}(v')$. This proves the theorem. □

The proof is the geometric doubling argument of `Space.theta_omega_ratEmbed`. It explains why a theorem about a polarization of the underlying symplectic vector space is not

being silently used to specify the phase of an anisotropic rational unitary element. The group property, zero-term formula and product formula are sufficient.

The exact product convention of the source. One can now choose the normalized theta phases consistently, not only up to arbitrary scalars. Choose a normalized theta on each of the standard signed lines; on the direct product of such lines choose their tensor product, which has norm one by Corollary 9.4. Every framed space with the same sign pattern is obtained from this signed standard space by a rational isometry and an adelic unitary change of coordinates, by Proposition 6.10. Transport the functional using the corresponding normalized Weil operator. If two such unitary changes carry its rational lattice to the standard rational lattice, their quotient preserves that lattice; in rational coordinates this is an actual rational unitary element. Theorem 9.6 shows that it fixes theta. Thus the transported normalization is independent of the isometry chosen. It is multiplicative for orthogonal sums, by choosing the product isometry, the product standard theta, and using the block-product action. This is exactly the choice of `exists_isThetaFunctional_signedSpace` and `Space.rawTheta` in `Lib/Weil/Construction/PairDefs.lean`. It supplies the exact scalar-one product law `theta_boxProd_source` and its target analogue used in the doubling calculations, rather than just an equality up to a constant phase.

9.9 Application to a unitary dual pair

For rational Hermitian spaces S, T, the Hermitian Hom form is $h(A, B) = \mathrm{Tr}(J_S^{-1} A^* J_T B)$, with the frames transported in the evident way. The pair of unitary elements (g, h) acts by $X \mapsto hXg^{-1}$, preserving the form by cyclicity of trace. In tensor coordinates its matrix is $h \otimes (g^{-1})^t$, which has determinant $(\det h)^{\dim S}(\det g)^{-\dim T}$. Thus restricting the just-constructed Weil action to the dual pair gives commuting actions, and twisting them by determinant characters trivial on rational points preserves rational theta invariance. These observations give the source's `G.onHom_mulVec` and the automorphy of the theta kernel in `PairSplitting.lean`.

For the exact character bookkeeping, let A, B denote the two characters of the norm-one ideles modulo rational norm-one elements, with

$$\omega_{A,B}(g, h) = A(\det h)B(\det g)^{-1}\omega(h \otimes (g^{-1})^t), \qquad C^{\mathrm{bc}}(x) = C(x/\bar{x}).$$

These are `pairChar`, `pairWeil` and the base-change convention in `Lib/Weil/Constructio n/PairDefs.lean`. Define $C_r(x) = C^{\mathrm{bc}}(x)\upsilon(x)^{-r}$, the source's `AutChar.spaceChar`. Let a be the dimension of an isotropic frame removed from the source, $r = \dim T$, m the map on that frame, and g_0 the residual source unitary element. Put $t = \det m$. The Levi acts on the dual frame by m^{-*}, so the induced action on the removed Hom coordinates is $h \otimes \bar{m}$, with determinant $d = (\det h)^a \bar{t}^r$. Also the determinant of the original g is $(t/\bar{t}) \det g_0$, by its triangular blocks. The source twists A by $\upsilon_{U(1)}^a$ before removing this frame. The ratio of the original pair character to the residual one is therefore

$$\upsilon(\det h)^a B^{\mathrm{bc}}(t)^{-1}.$$

By Theorem 9.5 the geometric zero term contributes $\upsilon(d)^{-1}|d|^{1/2}$, where $|\det h| = 1$, $|\bar{t}| = |t|$, and $\upsilon(\bar{t}) = \upsilon(t)^{-1}$. Multiplication gives exactly

$$B_r(t)^{-1}|t|^{r/2}. \tag{9.13}$$

If the isotropic frame is in the target, instead its induced Hom determinant is $d = t^r(\det h)^{-a}$, where h acts on the unchanged source of rank r. The original second character is $B\upsilon^a_{U(1)}$. The ratio of pair characters is $A^{\mathrm{bc}}(t)\upsilon(\det h)^{-a}$; multiplication by the geometric scalar gives

$$A_r(t)|t|^{r/2}. \tag{9.14}$$

This proves precisely the constants of `homSource_zeroTerm_pairWeil` and `homTarget_zeroTerm_pairWeil` in `Lib/Weil/DualPair/PairSplitting.lean`. The two formulas differ by inversion of the character but not of the positive modulus. There is no additional rank factor. The geometric computation also gives the nonzero-frequency source-term law by inserting a Heisenberg translation before taking the zero term, as in (9.11). Remaining concrete evaluations of particular joint-harmonic coefficient vectors are separate from this general theta normalization and are treated in Chapter 17.

9.10 Closure ledger for this chapter

Source assertion	Status	Exact justification or remaining obligation
`realAction`, `isVacuum_realVac`	Written proof	Lemma 9.1, Theorem 9.2; normalized Gaussian determinant and concrete projection.
`adelicAction`	Written proof	Product of Theorem 9.2 and Theorem 7.6; measure and block factorization.
`exists_isThetaFunctional`, `eq_smul`, quotient mean and Fourier inversion	Written proof	Theorem 9.3; summable coherent correlations and compact additive duality.
`boxProd`, orthogonal-sum theta normalization	Written proof	Corollary 9.4; fixed phase between separately normalized theta functionals is immaterial where a nonzero scalar is allowed.
`zeroTerm_omega`, `hasSum_theta_mixedTerm`	Written proof	Theorems 9.5 and Section 9.7; adapted hyperbolic model, scalar character and rational Fourier average.
`theta_omega_ratEmbed`	Written proof	Theorem 9.6, by a rational shear after doubling.
`homSource_zeroTerm_pairWeil`, `homTarget_zeroTerm_pairWeil`	Written proof	Section 9.9, formulas (9.13), (9.14); the source and target twist factors cancel against the geometric determinant character.

This closes the general nonpolar oscillator phase and theta-functional inputs at the level of the framed adelic model. We use standard uniqueness of the Schrödinger Heisenberg representation, which can also be proved by the coherent-vector construction where an explicit vacuum is given. Standard analytic facts used include Fourier inversion on compact abelian groups and the normed-space integrals indicated above. The specialized automorphic theta growth, joint-harmonic type and quantitative real-place comparisons are not consequences solely of this foundational calculation.

Chapter 10

Qualitative theta growth and convergence in the marked lattices

The exact oscillator action has now been established. For the unfolding and exchange of periods one also needs actual absolute convergence. The estimates in this chapter are qualitative in the field and the ranks: the constants may depend on the fixed datum and field. This is sufficient for the usual theta-lift and Fubini assertions, but does not replace the separate uniform estimates in the spanning theorem. We give an elementary height reduction and a small-translation proof of the rapid decay of the particular split-supported theta data, rather than infer it from an unqualified assertion about arbitrary theta kernels.

10.1 Lattice height and the compactness criterion

Use the notation of Chapter 6. Rank zero is trivial; set its height equal to one and consider positive rank below. For a framed rational nonsingular Hermitian space (V, J, x), an adelic unitary point g defines the rational integral lattice

$$L_g = \{v \in V(E) : (g^{-1}x^{-1}v)_f \in \widehat{\mathcal{O}}_E^n\}.$$

Its real embedding is given by $(g^{-1}x^{-1}v)_\infty$. Give the real product the maximum over the complex places of the ordinary positive Euclidean norm in the normalized signature coordinates. Set

$$\lambda_g = \inf_{v \in L_g \setminus \{0\}} \|(g^{-1}x^{-1}v)_\infty\|, \qquad H(g) = \max(1, \lambda_g^{-1}).$$

Equivalent Euclidean norms change the height by bounded factors. The choice of the maximum norm is convenient for the small isotropic-vector argument. The covolume of this real lattice is independent of g, with V, x fixed. In fact the change of finite lattice in the rational vector space is measured by the product of local determinant moduli, and the change of the real embedding by the real determinant modulus; for a unitary adelic element their product is one, as the idele determinant has norm one. This statement follows also by comparing the fractional lattices to a common lattice of finite index and applying the product formula. In particular the usual Euclidean covolume is a positive constant.

Lemma 10.1 (A lattice basis with polynomial height bound)**.** *There is a constant $C_V > 0$ such that L_g has n vectors v_i independent over E with*

$$\|(g^{-1}x^{-1}v_i)_\infty\| \le C_V H(g)^{2fn-1}, \qquad f = [E^+ : \mathbb{Q}].$$

Proof. Apply Minkowski's second theorem to the ordinary Euclidean lattice of real dimension $D = 2fn$ and a fixed symmetric convex unit ball (the maximum-of-Euclidean-norm balls may be used instead). Its product of successive minima is bounded above by a constant depending only on the ball and its covolume. Each of the first $D-1$ minima is at least $\lambda_g \geq 1/H(g)$. Thus the last minimum is at most $CH(g)^{D-1}$, yielding D independent vectors over $\mathbb{Q}$ with the claimed bound after a harmless enlargement of C. Their E-span is all V because a proper E-subspace has real dimension less than D, so one can select an E-basis from them. □

Here the standard external theorem is Minkowski's second theorem, in the usual form for a full lattice and a fixed symmetric convex body, e.g. Cassels [8]. No uniform arithmetic complexity estimate is claimed in this lemma.

Proposition 10.2 (Mahler compactness for these Hermitian quotients)**.** *For every fixed $M \geq 1$ there exists a compact subset C_M of the adelic unitary group such that every g with $H(g) \leq M$ has a left rational translate in C_M.*

Proof. By Lemma 10.1, choose the matrix r of an E-basis of V with integral finite normalized columns and real norm bounded in terms of M, V. Write $u = g^{-1}x^{-1}r$. Then the rational Gram matrix $B = r^*Jr$ equals u^*J_su in adelic coordinates, where J_s is the standard normalized form. Every entry of B is integral at finite places because the finite columns of u are integral; all its real embedding values are bounded by the real column bound. There are finitely many such matrices, by discreteness of the algebraic integers in their real embedding. Each is nonsingular since r is invertible.

For each matrix B that occurs choose one rational basis matrix r_B with that Gram matrix. Then $\gamma = rr_B^{-1}$ is a rational unitary element of V. Replace g by the left translate $g' = (x^{-1}\gamma^{-1}x)g$; one has

$$(g')^{-1}x^{-1}r_B = g^{-1}x^{-1}r = u.$$

The set of possible u with the stated real and finite bounds and $u^*J_su = B$ is compact in the adelic general linear group. To see invertibility and compactness explicitly, the determinant identity

$$\overline{\det u}\det u = \det B/\det J_s$$

holds at every place. At infinity the determinant of u has fixed nonzero absolute value at each embedding, so its inverse matrix is bounded by its adjugate formula. At finite places both determinant valuations in a conjugate pair are nonnegative and have sum bounded by the fixed valuation of $\det B$ (twice the valuation at a fixed place). Thus the inverse entries are uniformly bounded in a fixed fractional finite integral lattice and integral away from finitely many places. Both the matrices and inverses belong to compact products of bounded archimedean sets and compact finite integral lattices; the matrix equations are closed there. It follows that g' ranges over a compact set since the possible B, r_B form a finite set. □

This is the qualitative content of `Space.mahler_criterion` in `Lib/Adelic/Transference/MassBasic.lean`. It applies to indefinite signatures; definiteness is not assumed. In particular it includes the compactness used for the algebraic modular-form class sums. It does not give a uniform bound for the number of integral classes as the underlying field grows.

10.2 A bounded archimedean step that improves the height

Let g have $H(g) > 4$. The E-span of all vectors in L_g of real length less than one is totally isotropic. Indeed for any two such vectors v, w, their rational Hermitian pairing is an algebraic integer by the finite integrality condition. At each complex place its absolute value is less

than one by the usual Cauchy–Schwarz inequality applied to the normalized signature form. A nonzero algebraic integer has nonzero integral absolute norm and cannot satisfy all these strict inequalities. Hence the pairing is zero. The same holds for the entire E-span.

At each real place the images of this subspace form a complex totally isotropic subspace of the normalized signature form. Such a subspace can be carried by a compact block-diagonal unitary rotation to the span of vectors $(e_i^+ + e_i^-)$ in orthogonal positive and negative pairs: the projections to the positive and negative blocks are injective, their induced positive Gram matrices agree by isotropy, and orthonormalizing the common Gram matrix gives equal orthonormal lists in the two blocks. Extend these lists to orthonormal bases. On each paired two-plane the matrix

$$\begin{pmatrix} 5/4 & 3/4 \\ 3/4 & 5/4 \end{pmatrix}$$

is unitary for $\mathrm{diag}(1,-1)$ and has eigenvalues 2 and $1/2$ on the sum and difference vectors. It is the identity on the complement. Conjugating back gives a real unitary y_τ that multiplies every vector of the isotropic subspace by 2, has operator norm at most 2, and has inverse norm at most 2. Put $y_f = 1$. The family of all such y lies in a single compact subset, since there are finitely many possible numbers of hyperbolic pairs and the compact rotations vary in compact groups.

Set $g_1 = gy^{-1}$. The finite lattice is unchanged, and the real coordinates are multiplied by y. Vectors of length less than one are doubled at every place. Vectors of length at least one have new length at least $1/2$, since each inverse operator has norm at most two. Consequently

$$\lambda_{g_1} \geq \min(2\lambda_g, 1/2), \qquad H(g_1) \leq \max(H(g)/2, 2).$$

As $H(g) > 4$, this improves H by a factor of at least two. Iterate until the height is at most four. The number of steps is at most $1 + \log H(g)/\log 2$. Each step is right multiplication by the inverse of an element of the fixed compact family. Proposition 10.2 applies to the endpoint.

10.3 Moderate growth of the theta kernel

The Weil operator in the normalized integrable model has the bound

$$\|\omega(t)v\|_1 \leq |\Delta(t)|\,\|e\|_1\,\|v\|_1$$

by its explicit coherent integral. Both the real determinant and the finite multiplier are locally bounded; the latter is locally constant by its integral double-coset formula. The operator norms are therefore uniformly bounded for t in a compact subset of a framed adelic unitary group. The same is true for the restriction to a unitary dual pair by its continuous Hom action. All its determinant character twists have modulus one.

Theorem 10.3 (Qualitative moderate growth)**.** *For fixed framed spaces S, T, there are constants C, A such that every integrable datum v satisfies*

$$|\theta(\omega_{A_0,B_0}(g,h)v)| \leq C\|v\|_1\,(H_S(g)H_T(h))^A.$$

The constants may depend on the spaces, but not on g, h, v.

Proof. Apply the height-reduction steps of Section 10.2 separately to the two group variables, then translate their endpoints left by rational elements to reach compact subsets. The theta kernel is invariant under those rational left translations by Theorem 9.6; the determinant

characters are trivial on the rational determinants. The total operator norm in each direction is at most a fixed compact-endpoint bound times $B^{1+\log H/\log 2}$, for a fixed $B \geq 1$ bounding the operator norms of all individual compact steps. Multiplying the two estimates gives a power of the two heights. Finally apply the boundedness of θ from Theorem 9.3. □

This supplies `ThetaKernel.moderate_growth` of `Lib/Weil/DualPair/Lift.lean` by the qualitative version of `exists_rat_norm_le_height` in `Lib/Adelic/CuspForms/Peterson.lean`. Our power A need not be numerically uniform; it is used only to ensure the integrability of periods for each fixed field and rank.

10.4 Small real shifts in the Fock norm

Here is the analytic difference estimate used in the rapid-decay argument. Use the real Gaussian vacuum, the real Heisenberg translation $\rho(u)$ on the reproducing space, and the real positive Euclidean norm. For an integrable coefficient v write

$$M_N(v) = \int (1 + \|x\|)^N |v(x)|\, dx.$$

Lemma 10.4 (A finite difference of arbitrary order)**.** *For every integer $N \geq 1$ there exists C_N such that, if $\|u_i\| \leq t \leq 1$ for $1 \leq i \leq N$ and $M_N(v) < \infty$, then*

$$\|(1 - \rho(u_1)) \cdots (1 - \rho(u_N))v\|_1 \leq C_N t^N M_N(v). \tag{10.1}$$

Proof. The coherent expansion (9.7) implies, for partial derivatives ∂^α and nonnegative j with $|\alpha| + j \leq N$,

$$\int (1 + \|x\|)^j |\partial^\alpha v(x)|\, dx \leq C_{j,\alpha} M_N(v).$$

Indeed differentiation of the coherent vector $\rho(y)e(x) = \beta(y, x)e(x - y)$ gives a polynomial in $x - y$ and y of degree at most $|\alpha|$ times the same Gaussian and phase. Integrating with weight $(1 + \|x\|)^j$ bounds its norm by a constant times $(1 + \|y\|)^{|\alpha|+j}$. Fubini and differentiation under the coherent integral (or first truncation and convergence in the weighted derivative norm) give the assertion.

Write $\rho(u)$ in coordinates as translation by $-u$ times the phase $e^{iB(u,x)}$, where B is a fixed real bilinear alternating form. The difference product in (10.1) is, up to sign, the integral over $[0, 1]^N$ of the mixed derivative

$$\partial_{s_1} \cdots \partial_{s_N} \rho(s_1 u_1) \cdots \rho(s_N u_N)v.$$

The product inside is a translate by $-\sum_i s_i u_i$ times

$$\exp\left(iB\left(\sum_i s_i u_i, x\right) + i \sum_{i<j} s_i s_j B(u_i, u_j)\right).$$

Each derivative contributes a factor bounded by a constant times $\|u_i\|$ times a first derivative of v, or a polynomial of degree at most one in x and the bounded u_j. After all N derivatives, the terms are bounded in L^1 by a constant times $\prod_i \|u_i\|$ times a sum of the weighted derivative norms above, with total derivative order plus polynomial degree at most N. The translations are bounded because $t \leq 1$ and N is fixed. This proves (10.1), first for Schwartz vectors and then for all stated v by the weighted coherent estimates. □

The argument avoids differentiating the adelic variable. It is the real-coordinate finite difference mechanism needed below, in the integrable Heisenberg norm. In particular it applies to the polynomial Fock vectors, for which all the moments are finite.

10.5 The finite annihilation test

At a finite set of places disjoint from their conjugates one can polarize the split Hermitian matrix variables in the usual Schrödinger manner. A finite Schwartz–Bruhat datum Ψ on the chosen half then admits multiplication by any additive character of that half, implemented by a Heisenberg shift in the conjugate half. This is the finite component of (9.11), with the self-dual lattice and trace character already fixed. The other factors of the vector are held fixed. The following is the exact elementary principle needed in the source's `name_eq_zero_of_invariant`.

Lemma 10.5. *Let X, Y be adelic matrices such that the values $X_f^* \Psi_{mvar} Y_f$ are nonzero on the support of a compactly supported locally constant function Ψ of the selected matrix variables Ψ_{mvar}. Suppose L_0 is a bounded functional invariant under the Heisenberg translations by every adelic matrix of the form $XzY^* J_s$. Then L_0 vanishes on the vectors with finite name Ψ, regardless of their real component.*

Proof. In the split Schrödinger coordinates, translations supported on the complementary half multiply the selected-name vector at a point $A = \Psi_{\mathrm{var}}$ by the additive character of the pairing of z with $X_f^* A Y_f$. Because this last matrix is nonzero, local trace nondegeneracy supplies a finite z for which the character value differs from one. The pairing character is locally constant in A. On a small compact open neighborhood of each point of the support it is therefore a fixed scalar different from one. Compactness and total disconnectedness allow a finite disjoint compact open refinement of such neighborhoods and a corresponding decomposition of Ψ. Apply L_0 to each piece. Invariance under the selected translation says the value equals its product by that nontrivial scalar, hence it is zero. Summing proves the assertion. All translations act trivially on the real component since they have zero real coordinates. □

No assertion about the rank of the other coordinate is hidden here: the nonzero matrix condition is exactly what is needed to find a character distinguishing the finite name. The normalization factor in the additive character is a nonzero scalar and does not affect nondegeneracy. In the full-rank applications the nonzero condition follows from elementary matrix rank, as detailed next.

10.6 Decay on a full-rank split support

Fix a finite split-name datum and let the real datum v have all moments finite. We state the qualitative assertion with an explicit support condition. If S, T are the source and target spaces, suppose the selected finite name is supported on matrices which are surjective as maps from the normalized source coordinate space to the target at at least one fixed split-place factor. (The dual injectivity condition gives the reverse version.) Then for every N there are constants B, M, C depending on N, the fixed spaces and the fixed finite datum with

$$|\theta(\omega_{A_0,B_0}(g,h)(v \otimes \Psi))| H_T(h)^N \le C M_M(v) H_S(g)^B. \tag{10.2}$$

Here the notation $v \otimes \Psi$ includes the remaining fixed finite factors in the finite name. This is the qualitative version of `pairTheta_decay_of_onto` of `Lib/Weil/DualPair/KernelDecay.lean`; the transpose version uses injective support. It is not a uniform bound as the field, rank or finite name varies.

Proof. If $H_T(h) > 4$ take a nonzero short rational vector $d \in L_h(T)$ of real norm at most $2/H_T(h)$. Its Hermitian square is zero because it is an algebraic integer all of whose conjugates

have modulus less than one, as in Section 10.2. Choose an E-basis c of S made of vectors in $L_g(S)$ by Lemma 10.1. Write d', c' for their framed adelic coordinates at h, g. They are finite integral, c' is invertible, and

$$\|d'_\infty\| \, \|c'_\infty\| \le C_1 H_T(h)^{-1} H_S(g)^{D-1},$$

where D is the real lattice dimension of S. For the rational family of Hom vectors $dzc^* J_S$ as z ranges over rational row matrices, the adelic coordinates are $d'z(c')^* J_s$. Their Hermitian pairings vanish because d is isotropic. Thus their Heisenberg operators restrict to an ordinary additive representation of the adelic parameter group. Let

$$L = \theta \circ \omega_{A_0,B_0}(g, h).$$

By rational theta invariance the function of z given by $L(\rho(d'z(c')^* J_s)f)$ is periodic modulo the rational parameter lattice. Average it over that compact quotient to define a bounded functional L_0. It is invariant under translations of the entire parameter group by Haar invariance. On the support of Ψ the matrix $(d'_f)^* A c'_f$ is nonzero at the selected factor: d'_f is a nonzero column there because it is the image of a nonzero rational vector by invertible matrices; A is surjective, so its nonzero left functional does not vanish on every image; c'_f is invertible. Lemma 10.5 therefore gives $L_0(v \otimes \Psi) = 0$ for all real v.

Choose a positive integer M_0 such that the fixed finite name is fixed by every finite Heisenberg translation in $M_0 \widehat{\mathcal{O}}_E$ in the normalized Hom coordinates. Such an integer exists for a finite Schwartz–Bruhat vector: its translations and modulations are fixed in a small open additive subgroup, and the finite part is an ordinary vacuum outside finitely many places. Use a compact fundamental domain for the parameter adeles modulo rational parameters with finite coordinates in $M_0 \widehat{\mathcal{O}}_E$ and real coordinates in a fixed bounded set. This domain exists by the Chinese remainder reduction and a real lattice parallelepiped, as in Section 9.3. Since d', c' have integral finite coordinates, the associated finite translations lie in the subgroup fixing the datum. Thus for representatives in the compact domain the translation reduces to a purely real one on the real component, of size at most

$$t = C_2 H_T(h)^{-1} H_S(g)^{D-1}.$$

The identity $L_0(v \otimes \Psi) = 0$ holds for every real v, hence also after an arbitrary finite sequence of real Heisenberg differences, because its finite component is unchanged. Subtract the averaged identity from $L(v \otimes \Psi)$ and iterate. For any integer $K \ge 1$ this expresses $L(v \otimes \Psi)$ as the average of

$$L\big((1 - \rho(u_1)) \cdots (1 - \rho(u_K))v \otimes \Psi\big)$$

over independent parameter representatives, where all $\|u_i\| \le t$. All averages are legitimate by continuity and boundedness on the compact domain. If $t \le 1$, Lemma 10.4 and Theorem 10.3 give a bound

$$C_3 (H_T(h) H_S(g))^A t^K M_K(v).$$

Choose $K \ge N + A$ to obtain (10.2). If $t > 1$, then $H_T(h) < C_2 H_S(g)^{D-1}$; moderate growth alone implies (10.2) after increasing B, C. The case $H_T(h) \le 4$ is treated in the same fashion. The reverse assertion follows by using a short isotropic vector in the source and a full basis of the target, with the injective rank condition instead of the surjective one. □

The proof makes explicit why a finite full-rank support can remove a constant term, and why that removal gives arbitrary qualitative decay rather than a single small factor. It uses the split mixed finite model in the precise character convention of Chapter 9; any concrete vector must still be shown to possess the required support. It does not assert rapid decay of a general indefinite theta kernel without such a support or an appropriate vanishing hypothesis.

10.7 Convergence and transposition of theta periods

For fixed h, an automorphic function $y(g)$ decreasing faster than every power of $H_S(g)$ satisfies that $\theta(\omega(g,h)v)y(g)$ is bounded and integrable on the quotient, by Theorem 10.3 and finite covolume of the unitary quotient. For the latter use Getz–Hahn, [17, Theorem 2.6.3], in the number-field case: the adelic norm-one quotient of a smooth connected group has finite invariant volume. Here the unitary group over $F = E^+$ is smooth connected reductive, and it has no nontrivial F-rational character: over an algebraic closure its characters are powers of the determinant on GL_n, and the quadratic descent automorphism inverts that determinant. Thus its adelic norm-one subgroup is the entire group and the stated theorem applies to the quotient used here. The measure can be any fixed Haar normalization. This is a reference for finite volume only, not for the numerical estimates later in the book. The integral defining the theta lift thus converges absolutely. The same bound, uniform for h in a compact set, gives continuity by dominated convergence. Right translations and rational left invariance give lift equivariance by the ordinary quotient change of variables. If the finite datum satisfies the full-rank support in the preceding theorem, the resulting function inherits rapid decay in h: multiply (10.2) by $|y(g)|$ and integrate, absorbing the fixed power of $H_S(g)$ by rapid decay.

For two rapidly decreasing functions on the two quotients, their product against the common theta kernel is likewise absolutely integrable on the product quotient, since moderate growth is at most a fixed power in each variable. Fubini yields exact transposition of the two lift integrals in the conventions in which the theta pairing is bilinear. The identical reasoning permits an additional fixed moderate weight and a fixed group map whose height is at most a constant times the original height: absorb those fixed powers of the two heights into the two rapidly decreasing functions. These are the assertions `integrable_theta_mul`, `integrable_prod_weightFn`, and `integral_lift_eq_integral_descent` in `Lib/Weil/DualPair/Lift.lean`. They are qualitative; a new bound must be proved when the subsequent argument needs a uniform numerical estimate over the asymptotic data.

10.8 Status of the growth interfaces

Source assertion	Status	Justification
`Space.mahler_criterion`, qualitative rational norm bound	Written proof	Lemma 10.1, Proposition 10.2, Section 10.2, using classical Minkowski second theorem.
`ThetaKernel.moderate_growth`	Written proof	Theorem 10.3, exact normalized action and compact-step argument.
`name_eq_zero_of_invariant`	Written proof in the split-name model	Lemma 10.5; verify nonzero matrix condition for the specified finite name.
`pairTheta_decay_of_onto`, injective analogue	Qualitative proof	Section 10.6, with the explicit support hypothesis and finite tensor model; no estimate uniform in all arithmetic parameters is claimed.
Absolute convergence and lift transposition	Proof using a verified reference	Section 10.7, Getz–Hahn Theorem 2.6.3 with character hypothesis checked; rapid decay of the tested functions, moderate growth and Fubini.

This closes the basic convergence and covariance issues of the theta kernels in the model used in the earlier seesaw discussion. It does not prove independently that every automorphic function produced by a later construction has the full-rank support or the prescribed discrete-series type. Those are separate assertions about the actual chosen data, and the quantitative error terms require the stronger estimates addressed in the subsequent chapters.

Chapter 11

The convergent-range Siegel–Weil formula in the integrable model

The preceding chapters establish the vacuum-normalized Weil representation, its mixed model and the rational theta functional. We now give a literature bridge for one of the major remaining global formulas. Merely citing Siegel–Weil in a stable range would not suffice, because the source uses an integrable vacuum-coefficient Banach space, the probability quotient mean, and zero terms along rational Lagrangians. We check all three differences explicitly.

11.1 The classical theorem and the range used here

Let $F = E^+$, S be a split Hermitian space over E/F of rank $2a$, and T any nonsingular Hermitian space of rank $m > 2a$. Its rational Witt index r satisfies $2r \le m$, so $m-r \ge m/2 > a$. The Hermitian S can equally be viewed as a split skew-Hermitian space by multiplying its form by the fixed nonzero imaginary scalar. Its isometry group is unchanged. The associated symplectic pair is precisely the restriction of scalars of $\mathrm{Hom}_E(S,T)$ with the trace of the imaginary scalar times its Hermitian form; the fixed additive character of F can be chosen so that its trace pairing is the one used in Chapter 9.

We use the convergent-range classical unitary Siegel–Weil identity, with the following references and normalizations:

- A. Weil, *Sur la formule de Siegel dans la théorie des groupes classiques*, Acta Math. 113 (1965), no. 52, Theorem 5, in its Hermitian case. The theorem uses the Haar measure of mass one on the rational group quotient. The hypothesis on the rank in the quadratic-extension Hermitian case is the condition $m > 2a$ used here.
- The modern unitary formulation in S. Yamana, *The Siegel–Weil formula for unitary groups*, Pacific J. Math. 264 (2013), Theorem 2 (p. 237), with the absolutely convergent range $m > 2a$ also stated at the start of Section 3. In the notation of that theorem take the quadratic field case $D = E$, $\delta = 1$, and $n = a$. The standard section at the Siegel–Weil point $s_0 = (m-a)/2$ is the zero evaluation $\Phi \mapsto (\omega(g)\Phi)(0)$. Since $m - r > a$, the factor in the theorem is one, not the factor occurring in the small-rank regularized case.

In particular for the ordinary Schwartz–Bruhat Schrödinger vectors, the statement is

$$\int_{[U(T)]} \theta(\omega(1,h)\Phi)\, d\bar{h} = \sum_{\gamma \in P(F)\backslash U(S)(F)} (\omega(\gamma,1)\Phi)(0), \tag{11.1}$$

where P stabilizes a maximal isotropic frame, $d\bar{h}$ is Haar probability measure, and both expressions are absolutely convergent in this range. Formula (11.1) is the usual convergent formula, not a statement about analytic continuation past a pole. The original Haar normalization in Weil is relevant: replacing probability measure by an unnormalized Tamagawa measure would introduce its total mass. The checks in the remaining sections identify the exact terms in (11.1) with the nonpolar normalization; they also extend its assertion to the normed space of the source.

11.2 Identifying the splitting of the Weil representation

In the source, `splitWeil` of `Lib/SiegelWeil/Formula/ZeroTermHeight.lean` is the pair representation with characters $A = \upsilon_{U(1)}^{a}$, $B = 1$. (The declaration occurs in `ZeroTermHeig ht.lean` in the compact source.) A rational maximal isotropic frame of S yields the standard Schrödinger model on $T(\mathbb{A}_E)^a$. The action of $U(T)$ on this translation variable is exactly

$$(\omega(1,h)\Phi)(X) = \Phi(h^{-1}X). \tag{11.2}$$

Indeed its induced action on the relevant removed Hom frame has determinant $(\det h)^a$ and modulus one; the geometric scalar in Theorem 9.5 is therefore $\upsilon(\det h)^{-a}$, which cancels the explicit pair character. The coordinate change on the translation variable follows from the action $X \mapsto hXg^{-1}$ on Hom. Thus there is no residual determinant twist on the group of integration.

On the split group $U(S)$, the parabolic formulas of Chapter 9 give, for the matrix b acting on the chosen isotropic frame, a dilation scalar

$$\upsilon(\det b)^m |\det b|_{\mathbb{A}_E}^{m/2}.$$

The unipotent action is multiplication by the quadratic additive character of the Gram form in the translation variable. These are the standard Schrödinger formulas for the unitary split group with splitting character $\chi_T = \upsilon^m$ (if the dual isotropic frame rather than this frame is used to name the Levi, invert the corresponding determinant character together with the matrix variable). In Yamana's induced-representation convention of Section 1 the modulus is $|\det b|^{s+a/2}$, which equals $|\det b|^{m/2}$ at s_0. Moreover $\upsilon|_{J_F} = \eta_{E/F}$ by Proposition 6.4, so χ_T has the required restriction $\eta_{E/F}^m$ for that theorem. The additive characters agree on the unipotent formula by our choice of the trace pairing; the quadratic-form scalar can equivalently be absorbed in the nontrivial character of F, which is arbitrary in the theorem.

These parabolic formulas determine the entire split-group action, not just its zero evaluation. Indeed two genuine unitary splittings implementing the same Heisenberg action on an irreducible Schrödinger model differ, by Schur uniqueness, by a unitary character of the unitary group. Agreement on the parabolic implies triviality on its conjugates, hence on the whole split group. To see the last assertion locally, the group is generated by the Levi and the two opposite radicals by Gaussian elimination in a hyperbolic basis, or the relative Bruhat decomposition. For each exchanged hyperbolic plane the reflection is a product of three elementary upper/lower shears with skew-Hermitian parameters: the usual matrix product

$$\begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -u^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & u \\ -u^{-1} & 0 \end{pmatrix}$$

uses a nonzero invertible imaginary scalar u, both u and $-u^{-1}$ skew under conjugation. The matrices preserve the hyperbolic form, and plane permutations lie in the Levi. Relative Bruhat representatives are generated by these exchanges. At a split place the same assertion

is ordinary block Gaussian elimination in GL_{2a}. Thus the character is trivial on each local group. Finite-support elements at finite places together with the real factors are dense in the restricted product, so continuity gives triviality adelically. This proves compatibility of the normalized action with the standard one used in (11.1). In particular we do not infer the compatibility just from the fact that both actions are unitary.

11.3 The evaluation at a rational Lagrangian

Let K denote the distinguished isotropic a-plane of S. In the nonpolar model let $\mathrm{ev}_K(v)$ be the zero term, a scalar since the residual Hom space is zero. The mixed normalization of Section 9.7 is precisely that

$$\theta(v) = \sum_{X \in \mathrm{Hom}_E(E^a, T)(E)} F_v(X), \qquad \mathrm{ev}_K(v) = F_v(0),$$

where F_v is the wave function of v with its common fixed scalar. This scalar choice is immaterial to a linear identity on all wave functions, but is crucial for identifying the two source expressions on the same vector. If K' is another rational Lagrangian and γ is a rational isometry taking K' onto K, then the zero term along K' equals $\mathrm{ev}_K\, \omega(\gamma, 1)$. This follows from the rational invariance proved in Theorem 9.6, the expression as probability quotient mean, and the change of the rational frame coordinates. In particular it is independent of the choice of γ: a second choice differs by a rational parabolic element, whose zero-term factor is one by the product formula and principality of the determinant character. The rational unitary group acts transitively on maximal isotropic subspaces of a split Hermitian space by extending a basis to a hyperbolic basis (choose a dual frame and remove its Gram matrix by a skew-pairing correction). The cosets in (11.1) therefore enumerate exactly these Lagrangians. In particular there is no multiplicity or orbit-stabilizer factor in this enumeration.

It follows that on Schwartz–Bruhat vectors the two sides of (11.1) are already the source's probability mean and its sum of `Lagrangian.zeroTerm` of `ZeroTermHeight.lean`. There is still a substantive issue: the source asserts boundedness on the integrable model and summability of the *norms of the zero-term operators*. Pointwise equality on Schwartz functions alone would not prove that. We establish the required bounds next.

11.4 The maximum of a coherent wave vacuum

Let e be the normalized nonpolar vacuum on the full adelic Hom space. For any Lagrangian K' the corresponding wave vacuum has absolute value at an arbitrary translated point at most its absolute value at zero. Here is a direct verification, valid for a Lagrangian not necessarily adapted to the positive real coordinates. In its Schrödinger realization let F_e be the norm-one wave function; multiplying by a scalar to use the theta normalization does not affect the assertion. Along the modulation subgroup kb, its vacuum coefficient is

$$\langle F_e, R(kb) F_e \rangle = \int |F_e(X)|^2 \psi(\jmath b^* X)\, dX = e(kb).$$

The right side is a product of positive-definite real Gaussians in b and finite lattice indicators. The finite part on this isotropic subspace has phase one, since its Hermitian self-pairing is zero. It is the indicator of the intersection of the finite normalization lattice with the modulation subspace, a full compact open lattice there. Fourier inversion gives $|F_e(X)|^2$ as a product of positive-definite real Gaussians (with inverse covariance matrices) times indicators of the finite annihilator lattices, multiplied by a positive constant. The real

Gaussian transform follows from ordinary Gaussian integration and the finite transform from character orthogonality. The equality holds almost everywhere by Fourier uniqueness for the integrable density, and everywhere in its continuous realization in the wave model. The displayed function is maximized at zero. The coefficient evaluation at $X = 0$ of a coherent vector $\rho(a)e$ has the same absolute value as F_e at the corresponding translated point, by the mixed Heisenberg law. Hence

$$\sup_a |\operatorname{ev}_{K'}(\rho(a)e)| = |\operatorname{ev}_{K'}(e)|.$$

Using the coherent integral expansion of Chapter 9, we conclude

$$\|\operatorname{ev}_{K'}\|_{(L^1)^*} \le |\operatorname{ev}_{K'}(e)|. \tag{11.3}$$

The actual vacuum itself corresponds to a Schwartz–Bruhat vector in any fixed split rational Schrödinger model: compare its frame to an adapted frame as in Chapter 9, and apply a fixed linear symplectic transform to a Gaussian times a locally constant compactly supported wave, which preserves the Schwartz–Bruhat space by the elementary translation, quadratic-character, linear-change and Fourier operations. Therefore the absolute convergence of the classical Eisenstein series for this vector implies

$$\sum_{K'} |\operatorname{ev}_{K'}(e)| < \infty, \qquad \sum_{K'} \|\operatorname{ev}_{K'}\| < \infty$$

by (11.3). This is exactly the operator-norm summability required in `siegel_weil`; no assertion that mere pointwise convergence implies an absolutely summable operator series is being made.

11.5 A common integrable majorant for coherent translates

We can choose the reference hyperbolic coordinates on S with an adapted frame: its rational complete splitting is isometric to the standard hyperbolic model, and an adelic frame change gives a fixed bounded automorphism of the integrable Hom model. For that reference model the wave vacuum F_0 is a constant times a positive real Gaussian times a finite lattice indicator. For all h the pullback (11.2) preserves this shape of positive Gaussian times indicator, though its covariance and finite lattice may change. Any coherent translate in the Heisenberg variable changes the wave function by a translation and unit-modulus multiplication. We claim the uniform inequality

$$\left|\sum_{X \in T(E)^a} (\omega(1,h)\rho(b)F_0)(X)\right| \le \sum_{X \in T(E)^a} |F_0(h^{-1}X)| \quad \text{for every } b. \tag{11.4}$$

Indeed after discarding phases, the finite translated support either has no rational point, in which case its sum vanishes, or is a coset of the rational lattice selected by the unshifted finite support. The remaining real sum is a shifted positive Euclidean lattice Gaussian. For any positive-definite quadratic form Q_0 and full Euclidean lattice M, Poisson summation gives

$$\sum_{x \in M} e^{-Q_0(x+c)} = \frac{1}{\operatorname{covol}(M)} \sum_{\xi \in M^*} \widehat{e^{-Q_0}}(\xi) e^{2\pi i \langle \xi, c \rangle} \le \sum_{x \in M} e^{-Q_0(x)}$$

because the Gaussian Fourier coefficients are nonnegative, and all sums converge absolutely. This proves (11.4). Thus the absolute theta value of a normalized coherent vector is bounded by a common positive theta series of a fixed Schwartz vacuum, at every h. That positive

series is integrable by the classical Weil convergence theorem in the rank range checked above. A change from the reference adapted frame to another frame replaces the coherent expansion by one of a fixed bounded Weil transform of the data, and therefore preserves the existence of this bound with a constant.

Now every v in the integrable model is a Bochner integral of its coherent translates against $v(b)$; applying the bounded theta functional and the group action for each fixed h gives the corresponding ordinary integral. Inequality (11.4) implies

$$|\theta(\omega(1,h)v)| \le C\|v\|_1\Theta_0(h), \qquad \int_{[U(T)]}\Theta_0(h)\,dh < \infty$$

for a fixed positive majorant Θ_0. In this way the theta integral defines a bounded linear functional on the full integrable model, not only on Schwartz vectors. The coherent span consists of Schwartz–Bruhat wave functions and is dense in the integrable norm by the reproducing identity. By (11.3) the sum of zero-term operators also defines a bounded linear functional. The classical equality (11.1) holds on the coherent span, so it extends to the whole model.

Theorem 11.1 (Convergent-range formula in the source model)**.** *Under the source's hypotheses of a split rank-$2a$ source S, a rank-m target T with $2a < m$, and compatible Weil data, one has*

$$\sum_{K'}\|\,\mathrm{ev}_{K'}\,\| < \infty, \qquad \int_{[U(T)]}\theta(\mathrm{splitWeil}(1,h)v)\,dh = \mathrm{vol}([U(T)])\sum_{K'}\mathrm{ev}_{K'}(v)$$

for every integrable vacuum-coefficient vector v. The integral is absolutely convergent.

Proof. The preceding two sections prove the boundedness and summability in exactly the same source normed model; they extend the normalized formula from Schwartz vectors. Multiply the probability-mean identity by the volume of the fixed Haar quotient measure, which is finite by Chapter 10. This proves the displayed formula. When the split source has rank zero the model has a single scalar and the assertion reduces to the integral of a constant, so that case is immediate. □

This closes the statement `siegel_weil` of `Lib/SiegelWeil/Formula/SiegelWeilFormula.lean` using the classical formula with its verified hypotheses, rather than by reconstructing all its reduction-theoretic proof. It does not automatically supply every application of Rallis or a specialized global norm identity: the two data in a doubling comparison must still have the exact graph action, characters and measures, and the relevant period must converge. Those distinct obligations are discussed in Chapter 34.

11.6 The graph term in the doubled datum

For use in one such comparison we record the exact graph normalization. Put $V = \mathrm{Hom}(S,T)$, where now S need not be split, and let $V^- = \mathrm{Hom}(-S,T)$. Negating the source reverses the Hermitian form. In normalized coordinates this can be implemented by complex conjugation of coefficients and a coordinate change: at infinity keep the same coordinate with reversed sign, and at finite places multiply by the inverse of a finite unit idele of norm -1 (such an idele exists by Lemma 6.1). This is the source map `negSource` in `Lib/Weil/DualPair/PairSumNegation.lean`. The change reverses the Heisenberg bicharacter and conjugates the finite refinement; since the idele is an integral unit it preserves the lattice. It preserves the vacuum norms and the quotient measure and identifies the rational lattices in the two rational models with opposite signs. Hence the resulting anti-linear isometry D intertwines

the opposite Heisenberg translations. The conjugate of the transported functional on V^- is a rationally invariant theta on V of mean one. By Theorem 9.3 there is a scalar c of modulus one such that

$$\theta_{V^-}(cDf') = \overline{\theta_V(f')}.$$

This is the source's `exists_theta_negSource`; all choices of the phase are made once.

Form the double datum $f \boxtimes cDf'$ in $\mathrm{Hom}(S \oplus (-S), T)$, which is the definition `doubleDatum` in `Lib/SiegelWeil/RallisInner/SharpData.lean` (the declaration `SharpData.lean` in the compact source). For a rational unitary element δ of S, take the graph Lagrangian $K_\delta = \{(\delta u, -u) : u \in S\}$. With Gram matrix of S denoted by J_S, a dual isotropic frame is $(\delta; 1)(2J_S)^{-1}$ with the natural choice of signs. The mixed zero term is the probability mean over the corresponding additive quotient. The Hom variables along this graph, after unfolding the dual frame, are the simultaneous translations of the two factors. In rational matrix coordinates, write a parameter as X in $\mathrm{Hom}(S, T)$ by the invertible change

$$X \longmapsto J_T^{-1} X J_S$$

and then transport to the normalized coordinates of V. This is the Kronecker change $J_T^{-1} \otimes J_S^t$ in the source zero-term formula. It identifies the rational parameter lattices and hence preserves their probability quotient means. On the first factor the graph introduces the action of δ^{-1}; on the second it introduces the opposite-sign translation, which D converts into the conjugate theta value. These identities are also obtained by substituting the graph frame $(\delta; -1)$ and the above dual frame into the Heisenberg law (9.11); the two trace phases are opposite. Rational covariance moves the first rational unitary action from the theta functional into the datum, and its determinant character is one. Using the exact orthogonal-product convention of Chapter 9, the graph zero term is therefore

$$\int_{V(\mathbb{A})/V(E)} \theta_V(\rho(X)\omega(\delta^{-1}, 1)f)\overline{\theta_V(\rho(X)f')}, d\bar{X} = \langle f', \omega(\delta^{-1}, 1)f \rangle$$

by (9.8). This proves `zeroTerm_graphSplitting` in `SharpData.lean` with its normalized Hilbert pairing, not just up to a complex scalar. The scalar equality is why the graph part of the doubled sum can be written as an ordinary matrix coefficient. The further assertion that the other Lagrangian terms vanish on a specified datum remains a support problem; the graph calculation alone does not prove that vanishing or the final Rallis identity.

Part II

Upper bounds: CM jets, theta chains and Hecke repair

Chapter 12

The CM differential, theta normalization and integral frame

The purpose of this chapter is to establish the arithmetic preliminaries of the local jet theorem. A theta kernel is not an arbitrary analytic function on a complex vector space. Its translation law, the rational structures of the Hermitian spaces, and the lattice in the invariant differential have to be compatible. If one normalizes an individual complex theta function independently at every prime, the product formula used later in the proof is lost. Here we exhibit the single normalization and explain why its finite and infinite sizes agree with the requirements of the jet estimate. We give proofs for the local theta identities, the period scaling conditional only on classical CM and limit theorems and the normalization of the Hecke character, and the arithmetic tight-frame construction. At the end we explain the remaining Galois comparison that is needed to pass between finite Weil representations. The discussion uses d, K, E, N_1, Φ, f as in the admissible data. Write $N_1 = N$ temporarily within the frame construction; this N is *not* the auxiliary level prime of the proof.

12.1 An elliptic curve normalized at a point of order two

Let d be the prime discriminant parameter, $d \equiv 7 \pmod 8$, and set $c_0 = (1 + \sqrt{-d})/2$, so that $\mathcal{O}_K = \mathbb{Z}[c_0]$ and $c_0(1 - c_0) = (d + 1)/4$. In particular 2 splits in K. If Λ is homothetic to a fractional ideal of K, the quotient $\mathbb{C}/\Lambda$ has its natural $\mathcal{O}_K$ action. The two idempotents in $\mathcal{O}_K/(2) \simeq \mathbb{F}_2 \times \mathbb{F}_2$ act on its 2-torsion. There is a distinguished nonzero 2-torsion point t whose two projections are nonzero. Thus $2t \in \Lambda$, but neither $c_0 t$ nor $(1 - c_0)t$ belongs to Λ. In the source, such a t is part of the datum `CMLattice`.

Denote the Weierstrass functions of Λ by $\wp$, $\wp'$. Since t is a nonzero point of order two, $\wp'(t) = 0$ and $\wp''(t) \neq 0$. The second fact follows either from smoothness of the cubic or from the differential equation $\wp'^2 = 4\wp^3 - g_2\wp - g_3$: the cubic has distinct roots. On scaling Λ and t by u, the value of $\wp''(t)$ is multiplied by u^{-4}. We may therefore arrange $\wp''(t) = 2$. This fixes the differential up to a fourth root of unity. Choose a_1 with $a_1^2 = 12\wp(t)$. The shifted functions

$$X(z) = \wp(z + t) - \wp(t), \qquad Y(z) = \tfrac{1}{2}\big(\wp'(z + t) - a_1 X(z)\big)$$

give the affine equation

$$Y^2 + a_1 XY = X^3 + X. \tag{12.1}$$

Indeed $g_2 = 12\wp(t)^2 - 4$, so shifting the cubic by its root $\wp(t)$ gives $(2Y + a_1X)^2 = 4X^3 + a_1^2X^2 + 4X$. Its Weierstrass discriminant is $\Delta = a_1^4 - 64$ and $c_4 = a_1^4 - 48$. This is checked from $b_2 = a_1^2$, $b_4 = 2$, $b_6 = 0$, and $b_8 = -1$. The point $z = 0$ corresponds to $(0, 0)$ and the

point at infinity to $z \equiv t \pmod{\Lambda}$; z agrees with the invariant differential coordinate after translation by the point of order two. In particular $X = z^2 + O(z^4)$ and $Y = z + O(z^2)$. Differentiating gives

$$X' = 2Y + a_1X, \qquad Y' = 1 + 3X^2 - a_1Y, \qquad X'' = 6X^2 + a_1^2X + 2. \tag{12.2}$$

Proposition 12.1 (Unit discriminant in the chosen normalization)**.** *For every prime p and embedding of coefficients into an algebraically closed p-adic field, the normalized cubic* (12.1) *satisfies $|a_1|_p \le 1$ and $|\Delta|_p = 1$.*

Proof. We use two classical elliptic-curve facts: a CM j-invariant is an algebraic integer, and integrality of j implies potential good reduction (Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, II, §6, and *The Arithmetic of Elliptic Curves*, VII, Proposition 5.5). The normalized a_1 is algebraic, since $j = (a_1^4 - 48)^3/(a_1^4 - 64)$. Thus after a finite extension of the local field of definition we have a good Weierstrass model. All the endomorphisms and the point t can also be defined there, enlarging the extension if necessary. If $p \neq 2$, the nonzero point of order two has nonzero specialization, by finite etaleness of the 2-torsion. For $p = 2$ we use the distinguished choice of t. The complementary idempotents of $\mathcal{O}_K/(2)$ decompose the finite flat 2-torsion group scheme in the good model into two factors of rank two. The endomorphisms extend to the good model, and the induced idempotents act on its 2-torsion including the special fibre. On the one-dimensional tangent space of the special-fibre elliptic curve one of these two projectors is zero, the other the identity. The corresponding factor of rank two with zero tangent is etale. The projection of t onto this factor is a nonzero generic-fibre point, and specializes nontrivially. In particular t itself specializes nontrivially. This argument is the reason that choosing the point outside both CM kernels is essential: an arbitrary 2-torsion point on an ordinary curve might specialize to the identity.

We may work over the algebraically closed valuation field and translate the coordinates of our good Weierstrass model so that t is $(0, 0)$. Its affine coordinates are integral, since the reduction of the point is different from the point at infinity. Thus a good equation has the shape

$$y^2 + A_1xy = x^3 + A_2x^2 + A_4x$$

with integral coefficients; the vanishing of the constant term and of the term linear in y follow from the point and its order two. At $(0, 0)$ the partial derivative with respect to y vanishes, and the partial derivative with respect to x is $-A_4$. Smoothness of the integral model therefore implies that A_4 is a unit. A change $y = y' + sx$ eliminates the term in x^2 if $s^2 + A_1s = A_2$. All roots of this monic integral equation are integral. Scaling by a fourth root of A_4 then gives a good equation $y'^2 + A_1'x'y' = x'^3 + x'$. These operations are integral changes with unit leading scale. Finally the two normal forms of equation (12.1) over the algebraically closed local field are isomorphic preserving infinity and the distinguished point $(0, 0)$. A general change of Weierstrass coordinates with these two conditions has $x = u^2x'$ and $y = u^3y' + su^2x'$. Because the two coefficients of the term linear in x are both 1, one has $u^4 = 1$. Consequently the two discriminants differ by a unit, and $|a_1|_p \le 1$ follows from $a_1^4 = 64 + \Delta$. □

This proves the normalized integrality statement `CMLattice.norm_disc_eq_one` and `CMLattice.norm_a₁_le_one` in `Lib/CM/LocalHeight/IntegralParameter.lean` by a classical-model route instead of the formal source's explicit calculation with a chain of two-isogenies. We do not need the stronger assertion of reduction over any specified field of definition: the formal argument works after transferring coefficients to an algebraically closed p-adic field. The elementary good-model normalization also applies to all conjugate curves, since they carry the conjugate CM action and distinguished point.

12.2 Theta sections and their integral size

Put $A = \mathrm{area}(\Lambda)$. We must use the sigma function in the Hermitian normalization, not the unmodified classical Weierstrass product. If σ_W is the latter, with quasi-period η_l at $l \in \Lambda$, the usual quasi-period relation on an oriented basis of the lattice gives

$$\frac{\pi\bar{l}/A - \eta_l}{l} = c \qquad (l \in \Lambda \setminus \{0\})$$

for a single complex scalar c (this is the Legendre relation). Thus $\sigma(z) = e^{cz^2/2}\sigma_W(z)$ is an odd theta section with derivative one at zero, transforming at a period l by the sigma sign ($+1$ if $l \in 2\Lambda$, -1 otherwise) times $\exp(\pi(z\bar{l} + |l|^2/2)/A)$. This is precisely the choice of `PeriodPair.s` in `Lib/CM/Sigma/SigmaFunction.lean`. The classical addition and doubling identities are unchanged by multiplication by the indicated quadratic exponential; the exponential factors on their two sides agree. For a section of degree h use the Hermitian form $h\pi z\overline{w}/A$, and define the twisted translation

$$(U_y^h F)(z) = \exp\bigl(-h\pi(z\bar{y} + |y|^2/2)/A\bigr)F(z+y). \tag{12.3}$$

Let $\vartheta = U_t^1\sigma/(U_t^1\sigma)(0)$. It is even, satisfies $\vartheta(0) = 1$, and has its simple zeros at $t + \Lambda$. These facts follow from the sigma transformation law and $2t \in \Lambda$. The addition formula for the sigma function gives, off the indicated zeros and then by continuation,

$$\vartheta(z+w)\vartheta(z-w) = \vartheta(z)^2\bigl(\vartheta(w)^2 - X(z)\sigma(w)^2\bigr). \tag{12.4}$$

Another consequence is the identity $(\sigma(z)/\vartheta(z))^2 = X(z)$. To see this directly, use the addition identity at $w = t$ and the relation $2t \in \Lambda$ to move the translated arguments back by a period. The proportionality constant is one, as both sides begin with z^2. Write $G(z) = Y(z)\vartheta(z)^3$; the apparent pole at $t + \Lambda$ cancels because Y has a pole of order three there. Thus G is an entire section of degree three. The identities used below are

$$\vartheta(2z) = \vartheta(z)^4 - \sigma(z)^4, \tag{12.5}$$

$$\sigma(2z)^2 = \sigma(z)^2\vartheta(z)^2\bigl(4\vartheta(z)^4 + a_1^2\sigma(z)^2\vartheta(z)^2 + 4\sigma(z)^4\bigr), \tag{12.6}$$

$$G^2 + a_1\sigma^2\vartheta G = \sigma^6 + \sigma^2\vartheta^4. \tag{12.7}$$

For the first, specialize (12.4) to $w = z$; for the second, use $\sigma(2z)^2 = \wp'(z)^2\sigma(z)^8$ together with $\vartheta(z)^2 = (\wp(z) - \wp(t))\sigma(z)^2$ and the shifted cubic. The last is (12.1) multiplied by ϑ^6. As both sides are entire, the identities are valid even when a denominator used in their derivation vanishes. The sigma addition/translation identities are classical (Weil, *Elliptic Functions according to Eisenstein and Kronecker*, Springer, 1976); the exact formulas are also the declarations `theta_add_mul_theta_sub`, `sigma_two_mul_sq`, `phiThetaCube_sq_add`, `θ_two_mul` in `Lib/CM/Sigma/CurveTheta.lean`.

Write $H_p(z) = \max(|\sigma(z)|_p, |\vartheta(z)|_p)$ after applying one fixed embedding ι to the complex values. It might seem strange to use a p-adic norm of the value of a complex analytic function at every complex argument; no continuity of this operation is asserted. The point is that the following conclusions are algebraic identities between their values.

Lemma 12.2 (Exact doubling of the local size)**.** *One has $H_p(z) > 0$, $|G(z)|_p \le H_p(z)^3$ and $H_p(2z) = H_p(z)^4$ for all $z \in \mathbb{C}$.*

Proof. The zeros of σ and ϑ are disjoint, so the maximum is positive. Divide (12.7) by the sixth power of a value of modulus $H_p(z)$. The resulting equation is monic in the normalized

G, with integral coefficients by Proposition 12.1. It follows that $|G(z)|_p \le H_p(z)^3$. The two duplication formulas immediately give $H_p(2z) \le H_p(z)^4$. If the inequality were strict, choose a scalar C of modulus $H_p(z)$ and put $u = \sigma(z)/C$, $v = \vartheta(z)/C$, viewed through ι. Their maximum norm is one. Both right sides of (12.5) and (12.6), divided by C^4 and C^8 respectively, would reduce to zero in the residue field. The first gives $u^4 = v^4$; hence both are units. The second implies $8 + a_1^2(u/v)^2 = 0$ in that field. Squaring and using $(u/v)^4 = 1$ yields $a_1^4 = 64$ there. This contradicts the unit discriminant, even when $p = 2$ (then a_1 reduces to a unit). The opposite inequality follows. □

It is useful to compare this with the usual local canonical height. A generic Weierstrass model leads to a bounded-error doubling formula and to a limiting procedure. Here the carefully chosen model removes the error at every prime, exactly in the same normalization. For a section represented by a monomial $\vartheta^a\sigma^{2b}G^c$ of degree $h = a + 2b + 3c$, the bound is $|\vartheta^a\sigma^{2b}G^c|_p \le H_p^h$; this is why the distinguished coordinates use two units of degree for X and three for Y.

Lemma 12.3 (Spanning by monomial sections)**.** *The degree-h theta sections for the multiplier of ϑ are spanned by*

$$\vartheta^a\sigma^{2b}G^c, \qquad a, b, c \ge 0, \quad c \le 1, \quad a + 2b + 3c = h.$$

Proof. Divide a section by ϑ^h. The quotient is a meromorphic function on $\mathbb{C}/\Lambda$ with no pole except at t, of order at most h. In the coordinates (12.1), the punctured elliptic curve is affine, and its ring of regular functions is $\mathbb{C}[X, Y]/(Y^2 + a_1XY - X^3 - X)$. Every element is uniquely $P(X) + YQ(X)$. Alternatively this uniqueness and spanning follow by subtracting the leading Laurent coefficient at the pole successively, since X, Y have poles of orders two and three. The monomials X^b, YX^b have distinct pole orders $2b, 3 + 2b$, so the pole bound means all appearing monomials have order at most h. Multiplying by ϑ^h and using $X\vartheta^2 = \sigma^2$, $Y\vartheta^3 = G$ proves the assertion. This includes $h = 0, 1$, when there are only the constant and ϑ sections respectively. □

The analytic and algebraic arguments are interchangeable here. In Lean the spanning assertion is `CMLattice.sections_le_span_monomialSection`, in `Lib/CM/Sigma/PadicSeriesDefs.lean`, and is proved by dividing entire sections rather than invoking the affine description of the elliptic curve. We will use both the pole description and its degree bookkeeping.

12.3 Isogenies, their multipliers, and Tate's inequality

If $a\Lambda \subseteq \Lambda'$ is an isogeny of degree n, then $|a|^2A = nA'$. The principal polarizations on these curves are represented by the Hermitian forms above. Pullback of the second form is n times the first form. The multipliers in the source are chosen to be the canonical ones defined using c_0; on periods their formula is

$$\chi(l) = \exp(\pi i E(\bar{c}_0 l, l)),$$

where E is the integral Riemann form. The formula can be checked explicitly on $\Lambda/2\Lambda$. In a basis e_1, e_2 of its two split summands the symplectic pairing of the two basis vectors is one modulo two. The sigma sign of a class $ue_1 + ve_2$ is $(-1)^{u+v+uv}$, being one just for the zero class. Translation by the distinguished t introduces the sign $(-1)^{u+v}$, because its class after multiplication by two is $e_1 + e_2$. Their product is $(-1)^{uv}$. Projection by $\bar{c}_0$ onto one summand

followed by pairing with the original vector has value uv modulo two, giving the claimed formula. Because multiplication by a is complex-linear and commutes with $\bar{c}_0$, the pullback of this multiplier is χ^n. This is the elementary consequence `comp_mul_mem_sections` of `multiplier_eq_canonicalMultiplier`, in `Lib/CM/Sigma/CurveTheta.lean`. In particular it is important not to choose independently a different two-torsion point on the target curve. If F is a degree-h section on the target, $F(az)$ is a degree-hn section on the source.

Proposition 12.4 (Tate's local inequality)**.** *Let a be a CM isogeny as above and y a torsion point modulo Λ'. If $|F(w)|_p \le H'_p(w)^h$ for all w, then*

$$|(U_y^h F)(az)|_p \le H_p(z)^{hn} \quad (z \in \mathbb{C}). \tag{12.8}$$

Proof. All values are read through the same embedding. Put $A_y(w) = \exp(-\pi(w\bar{y}+|y|^2/2)/A')$ and define

$$R_a(z) = |A_y(az)|_p \frac{H'_p(az+y)}{H_p(z)^n}.$$

The function R_a is bounded. Indeed, choose $q > 0$ with $qy \in \Lambda'$. Raising each of the two functions $A_y(w)\sigma'(w+y)$ and $A_y(w)\vartheta'(w+y)$ to the power $2q$ (where primes here label the target curve's sigma and theta) makes them degree-$2q$ sections of the original multiplier: twisted translation by a point of order dividing q introduces a character of finite order dividing $2q$ on the period lattice. Pulling back by a yields degree-$2qn$ sections on the first curve. By Lemma 12.3 and the height bound for each monomial, each such section is bounded by a constant times H_p^{2qn}. Taking a maximum and a $2q$-th root proves the asserted boundedness. This uses finite-dimensional linear algebra, not continuity in the p-adic topology of the complex variable.

The identity $A_y(2w)A_y(2w+y) = A_y(w)^4$ follows by adding the exponents. Lemma 12.2 gives

$$R_a(z)^4 = R_a(2z)R_1(2az+y),$$

where R_1 is the same expression for the identity on the target curve. Let $M_a = \sup_z R_a(z)$; it is a finite positive real number, as is M_1. Then $M_1^4 \le M_1^2$ and hence $M_1 \le 1$. Likewise $M_a^4 \le M_a M_1$, so $M_a \le 1$. Raising $R_a(z) \le 1$ to the power h and using the hypothesis on F proves (12.8). □

This is the mechanism of `CMLattice.norm_translate_le_naiveHeight_pow` in `Lib/CM/LocalHeight/IntegralParameter.lean`. The proof only needs the boundedness of each ratio and the exact duplication identity. It is stronger than a bound allowing an unspecified factor depending on the order of torsion, which could not be paid uniformly when selecting tight frames at different primes.

12.4 The integral formal parameter and the Gauss principle

We use the translated coordinate $Y = \phi$ as local parameter at $z = 0$. It is important that $\mathfrak{o}[[\phi]]$ need not equal the integral series ring in the invariant differential z. In what follows $\mathfrak{o}$ is the valuation ring of the algebraically closed p-adic field, and all series are formal. Substitution $T \mapsto \phi(z)$ gives a subring $\mathfrak{R}_\phi \subset \mathbb{C}_p[[z]]$; because the linear term of ϕ is one, it is isomorphic to $\mathfrak{o}[[T]]$. From the equation

$$X = Y^2 + a_1 XY - X^3$$

there is a unique formal solution $X = Y^2 + \cdots$ over $\mathfrak{o}$ by coefficient recursion. Hence $X \in \mathfrak{R}_\phi$. Equation (12.2) shows that differentiation in the invariant differential preserves the ring: if

$F(T) \in \mathfrak{o}[[T]]$, then $(F(\phi(z)))' = F'(\phi(z))(1 + 3X^2 - a_1\phi(z))$. All factors are in the ring. This gives the first part of Proposition 13.3.

Lemma 12.5 (Gauss principle for finitely many polynomials)**.** *For nonzero polynomials $P_1, \dots, P_j$ over an algebraically closed valued field with infinite residue field, one can find x integral such that $|P_i(x)|_p$ equals the maximum of the absolute values of the coefficients of P_i for every i simultaneously.*

Proof. Divide each P_i by a coefficient of maximal absolute value and reduce the polynomials. None of the resulting polynomials in the residue field is zero. Choose a residue point outside the finitely many roots of their product, and lift it to the valuation ring. For several variables the same proof uses that a nonzero polynomial over an infinite field cannot vanish everywhere (induct on the number of variables). □

Proposition 12.6 (Integral coefficients of a bounded section)**.** *Let F be a degree-h section of the above multiplier and suppose $|F(z)|_p \le H_p(z)^h$ everywhere. Then $F = \vartheta^h(P(X) + YQ(X))$ for polynomials P, Q all of whose coefficients are p-adically integral. In particular the transported formal Taylor series of F belongs to $\vartheta^h\mathfrak{R}_\phi$. If F is even then $Q = 0$. For even h, one has $F(t) = P_{h/2}\sigma(t)^h$ when F is even, where $P_{h/2}$ denotes the coefficient in that degree.*

Proof. Write the quotient by ϑ^h as in Lemma 12.3. Let $D(X) = 4X^3 + a_1^2X^2 + 4X$. This polynomial has Gauss norm one: for odd p the coefficient 4 is a unit, and for $p = 2$ Proposition 12.1 gives that a_1 is a unit. Choose an integral x at which $|D(x)|_p = 1$ and both P, Q attain their coefficient maxima, using Lemma 12.5 and neglecting identically zero polynomials. Such a point can be chosen in the image of the complex coefficient field: it suffices to choose its residue in $\overline{\mathbb{F}}_p$ and lift algebraically. The associated points of (12.1) above x are conjugate under elliptic negation with Y coordinates y and $-a_1x - y$. They are integral by the monic quadratic equation, and their difference has norm one since $(2y + a_1x)^2 = D(x)$. By elliptic uniformization (in the shifted coordinate), choose $z, -z$ giving these two points. Here ϑ is nonzero, and $|X(z)|_p \le 1$ gives $H_p(z) = |\vartheta(z)|_p$ by $X = (\sigma/\vartheta)^2$. The hypotheses show that both

$$P(x) + yQ(x), \qquad P(x) + (-a_1x - y)Q(x)$$

are integral. Subtracting and dividing by the unit difference proves integrality of $Q(x)$, hence also of $P(x)$. The way x was chosen proves integrality of all coefficients. This proof works even at $p = 2$: we divide by the norm-one difference of the roots, not by 2. The formal integrality statement follows from $X, Y \in \mathfrak{R}_\phi$.

When F is even, $\vartheta(-z) = \vartheta(z)$ and $Y(-z) = -a_1X(z) - Y(z)$ give $(2Y + a_1X)Q(X) = 0$ identically. Thus $Q = 0$. Finally $\vartheta^hX^b = \vartheta^{h-2b}\sigma^{2b}$. The pole-order bound permits only $2b \le h$; at $z = t$ all terms with $2b < h$ vanish, proving the claimed formula. □

This is a reconstruction of `CMLattice.exists_eq_theta_pow_mul_of_norm_le`, including the exact even-coefficient assertion, from `Lib/CM/LocalHeight/IsogenyIntegrality.lean`. Applying Proposition 12.4 to the three sections ϑ, σ^2, G gives their pullbacks under an isogeny as ϑ^n (or its corresponding power) times elements of $\mathfrak{R}_\phi$. Dividing the pullbacks of G' and ϑ'^3 is permitted on the formal disc since $\vartheta'(0) = 1$ (here ϑ' is the *target theta function*, not differentiation). The factor in $\mathfrak{R}_\phi$ has constant coefficient one and therefore an integral inverse in the Y coordinate. Thus the target formal parameter pulled back by the isogeny is integral in the source parameter. Comparing their first coefficients shows that the scalar a of any isogeny between these normalized curves is integral at every p-adic embedding. We recover `rescale_translatedWpDeriv_mem` and `norm_le_one_of_isogeny`, with no loss from the degree of the isogeny.

Corollary 12.7 (Integral isogeny translates)**.** *Let $\delta \in \mathcal{O}_K$ be nonzero, put $n = \delta\bar{\delta}$, and suppose $\delta y \in \Lambda$. Then the Taylor series of $(U_y^1\vartheta)(\bar{\delta}z)$ lies in $\vartheta(z)^n\mathfrak{R}_\phi$.*

Proof. Pullback by multiplication by $\bar{\delta}$ yields the original multiplier to power n. Twisted translation of this pullback by $y/\bar{\delta}$ preserves it, since $n(y/\bar{\delta}) = \delta y \in \Lambda$ and the dual lattice contains Λ. Its degree is therefore n. Formula (12.3) identifies it with the function in the statement. Proposition 12.4 bounds its size by H_p^n. Apply Proposition 12.6. □

This proves the assertion (13.5) in the local jet chapter, using the classical theta multiplier identities as above. Notice that this corollary is a statement about a formal expansion near zero; it does not assert that the theta function is a globally analytic p-adic function.

12.5 Logarithms, norm-pair limits, and the formal height

As $\vartheta(0) = 1$, define the formal logarithm $\mathscr{L} = -\log\vartheta$. Evenness gives $\mathscr{L}(0) = \mathscr{L}'(0) = 0$. From the addition law (12.4), take the coefficient of w^2 and use $\sigma'(0) = 1$. On the left the coefficient is $\vartheta\vartheta'' - (\vartheta')^2$; on the right it is $\vartheta^2(\vartheta''(0) - X)$. Consequently

$$\mathscr{L}'' = L_2 + X, \qquad L_2 = -\vartheta''(0). \tag{12.9}$$

For $\delta \in \mathcal{O}_K \setminus \{0\}$, put $n = \delta\bar{\delta}$. The isogeny pullback at $y = 0$ proves that the series

$$\phi(\delta z), \qquad F_\delta = \exp(n\mathscr{L}(z) - \mathscr{L}(\delta z))$$

are in $\mathfrak{R}_\phi$. Since $F_\delta(0) = 1$, its inverse belongs to the same ring; differentiation and the logarithmic derivative give $D'_\delta \in \mathfrak{R}_\phi$ for $D_\delta = n\mathscr{L}(z) - \mathscr{L}(\delta z)$. The same statements hold trivially for $\delta = 0$.

Differentiating once more and evaluating at zero shows that $(n - \delta^2)L_2$ is integral. For $\delta = c_0$ and $\bar{c}_0$ the norms n coincide. Their difference is $(\bar{c}_0^2 - c_0^2)L_2 = -\sqrt{-d}L_2$. This gives the desired bound $|\sqrt{-d}L_2|_p \le 1$ in (13.3), including at the ramified prime.

View $\mathfrak{R}_\phi$ with the coefficientwise topology, not with a topology coming from a complex disc. Let Z denote the formal inverse of ϕ in the variable z. A series belongs to $\mathfrak{R}_\phi$ exactly when every coefficient after composition with Z belongs to the valuation ring. In particular $\mathfrak{R}_\phi$ is closed in the topology of pointwise convergence on coefficients. Each coefficient of $\phi(cz)$ and of $\exp(D\mathscr{L}(z) - \mathscr{L}(cz))$ depends continuously (indeed polynomially, over the coefficient field) on (c, D); the same is true of the derivative of the exponent. Thus the three integrality assertions extend to the closure of the pairs $(\iota\delta, \iota(\delta\bar{\delta}))$ in the product of the two p-adic fields. This proves the entirety of Proposition 13.3 from the preceding propositions. It is the normalization in `CMLattice.exists_log_theta`, `Lib/CM/KroneckerLimit/LogTheta.lean`, together with `rescale_translate_eq_theta_pow_mul`.

Lemma 12.8 (Prime-to-p root of unity dilation)**.** *If p is inert in K, let $q = p^2$; otherwise, let $q = p$. Let ζ be a $(q-1)$-st root of unity in the algebraically closed local field. Then (ζ, ζ^{p+1}) belongs to the norm-pair closure just described.*

Proof. In the inert case, choose an algebraic integer $a \in \mathcal{O}_K$ whose reduction at the given embedding equals that of ζ in $\mathbb{F}_{p^2}$. This is possible because $\mathcal{O}_K/(p) = \mathbb{F}_{p^2}$, and conjugation on this field is the p-power map. Thus the reduction of $\bar{a}$ is that of ζ^p. In the split or ramified case $q = p$, and one may choose a to be an ordinary integer with the specified reduction in $\mathbb{F}_p$; the two reductions then agree. Since q is a power of p, one has $\iota(a)^{q^h} \to \zeta$ and $\iota(\bar{a})^{q^h} \to \zeta^p$ as $h \to \infty$. Indeed if a unit is $\zeta(1 + w)$ with $v_p(w) > 0$, then the binomial theorem shows

$v_p((1+w)^p-1) \geq \min(v_p(w)+1, pv_p(w))$, and iteration tends to infinity. The same argument applies to the conjugate unit. The integers a^{q^h} yield the required limiting norm pairs. □

This proves `mem_normPairClosure_of_pow_eq` in `Lib/CM/KroneckerLimit/LogTheta.lean`. Note that in the inert case conjugation is genuinely a nontrivial Frobenius on the residue field. Treating it as the identity would produce the wrong sign in the theta couplings. This is also why using an arbitrary prime-to-p root of unity, rather than one of order dividing $q-1$, would not suffice.

The remaining height input in Chapter 13 may be expressed in conventional elliptic-curve terms. Assume that $a \in \mathcal{O}_K$ is nonzero, $a\bar{a} = q$ is odd, and denote multiplication by a on the curve by $[a]$. With the distinguished point t of order two, $[a](t) = t$: multiplication by a on $\mathcal{O}_K/(2) \simeq \mathbb{F}_2^2$ is invertible because its norm is odd, so it is the identity there. We consider the germ of the isogeny at the point t, as in the invariant coordinate z of the local jet theorem.

Proposition 12.9 (The degree upper bound for the first nonzero reduced coefficient)**.** *Let T be any integral formal parameter tangent to the identity in $\mathfrak{R}_\phi$ and let $Z = T^{-1}$ in the invariant differential coordinate. The first nonzero coefficient of the reduction of $g(u) = T(\iota(a)Z(u))$ has degree at most q, including when $p \mid q$.*

Proof. The equation (12.1) is a good Weierstrass model by Proposition 12.1; the endomorphism extends to the good integral elliptic curve. Its reduction is an isogeny of the same degree q. Here we use the classical good-reduction property that reduction preserves degrees of isogenies (e.g. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, II, Proposition 4.4). The local ring of the reduced curve at the point t is a smooth one-dimensional local ring; its completed local coordinate Y is a uniformizer, as $\partial(Y^2 + a_1XY - X^3 - X)/\partial X = -1$ there. The fibre of the isogeny above the reduced t has scheme-theoretic length q. At the point t its local length is the order of the image of the target uniformizer in the source formal coordinate; hence the order is finite and at most q. Since $T = \phi + O(\phi^2)$ is an integral change of uniformizer, the same is true of the order computed with T. Equivalently, g has a unit coefficient of degree at most q. □

This supplies the needed case of `CMLattice.exists_unit_coeff_degree_le` (`Lib/CM/FormalGroup/FormalHeight.lean`) by the ordinary formal-group interpretation of the local degree. It is essential that the bound is for a reduction modulo the maximal ideal, not for the first nonzero coefficient over the local field. The characteristic-zero linear coefficient is a, but it is a nonunit at the primes of interest. For instance an inert prime gives $a = p$, $q = p^2$, whereas the ramified prime admits $a = \sqrt{-d}$ and $q = p = d$. In Chapter 13 the symmetry under the group of order $q-1$ leaves only the first and the q-th coefficient through degree q; the first is a nonunit. The present bound therefore proves that the q-th coefficient is a unit, which is exactly the input used to normalize the leading Frobenius term of the inverse parameter.

Lemma 12.10 (A coefficient form of the Hasse invariant)**.** *Let p be odd. Suppose every $F \in \mathfrak{R}_\phi$ satisfies $|[z^{p-1}]F'|_p < 1$. Then $|[z^{p-1}]X|_p \geq 1$. Moreover, if $\chi_K(p) = 1$, the displayed hypothesis cannot hold.*

Proof. Write $p - 1 = 2k$. Assume for a contradiction that the coefficient in X has norm less than one. Since $1, X, X'$ belong to $\mathfrak{R}_\phi$, the same assertion for 1 is trivial (its positive-degree coefficient is zero). For each $n \geq 0$ the differential equation gives

$$(X^nX')' = (4n+6)X^{n+2} + (n+1)a_1^2X^{n+1} + (4n+2)X^n. \tag{12.10}$$

Indeed use $(X')^2 = 4X^3 + a_1^2X^2 + 4X$ and (12.2). For $0 \le n \le k-2$, the integer $4n+6$ is positive, even, and strictly less than $2p$, so it is prime to p. The hypothesis on derivatives and induction, using integrality of a_1, show that $[z^{p-1}]X^j$ has norm less than one for all $0 \le j \le k$. But $X = z^2 + O(z^4)$ has $[z^{p-1}]X^k = 1$. When $p = 3$ the supposed assertion for X contradicts this directly. Thus $|[z^{p-1}]X|_p \ge 1$.

To see the final statement, take the dilation $c = \sqrt{-d}$ with $n = d$. The series $(d\mathscr{L}(z) - \mathscr{L}(cz))'$ lies in the integral parameter ring, so its derivative has z^{p-1} coefficient of norm less than one. By (12.9) this coefficient is $(d - c^{p+1})[z^{p-1}]X$. If p splits in K, the reduction of c is in $\mathbb{F}_p^\times$, so $c^{p+1} \equiv c^2 = -d \pmod{\mathfrak{m}}$. The factor $d - c^{p+1}$ is then a unit as p is odd and $p \neq d$, a contradiction. □

The even prime is not nonsplit in these data, so Lemma 12.10 is precisely the relevant part of `CMLattice.one_le_norm_coeff_translatedWp_nonsplit` in `Lib/CM/FormalGroup/FormalHeight.lean`. This proof avoids an identification of the actual coefficient with a modular form. In the nonsplit calculation the gap in the invariant coordinate of the root-of-unity equivariant parameter makes the hypothesis automatic at an inert prime, giving the norm-one first exceptional coupling coefficient of the local normal form. Its sharpness is needed for the comparison in the determinant lower bound; for the upper bound it would not be sufficient just to know that all the indicated coefficients were integral.

12.6 A unit-adjusted Chowla–Selberg formula

An archimedean period for an elliptic curve generally carries no absolute p-adic coefficient normalization. The following logarithmic version of Chowla–Selberg is particularly useful because the normalizing factor is an algebraic unit, so it contributes nothing to finite valuations. We explain why it has this stronger property.

For a fractional ideal $\mathfrak{b}$ of K denote its norm by $\mathcal{N}(\mathfrak{b}) > 0$, with the usual multiplicativity under fractional ideal products. The area of $\lambda\mathfrak{b}$ is $|\lambda|^2\mathcal{N}(\mathfrak{b})\sqrt{d}/2$. For $h_K = |\mathrm{Cl}(K)|$, the class group has odd order by the classical genus theory of imaginary quadratic fields of prime discriminant: its quotient by squares has cardinality one. Thus squaring permutes its elements. Choose representatives $\mathfrak{a}_c$ for each class c. The ideals $\mathfrak{b}_c = \mathfrak{a}_{c^2}/\mathfrak{a}_c$ again represent every class once, and $\prod_c \mathfrak{b}_c = \mathcal{O}_K$ exactly. This last equality of fractional ideals, not merely of their ideal classes, is why the quotients of the chosen representatives are used.

Fix a normalized CM curve C of period lattice Λ. For every c choose a normalized curve C_c with period lattice $u_c\mathfrak{b}_c\Lambda$, where $u_c \in \mathbb{C}^\times$. Such a choice is possible by the normalization at t above. Every element of $u_c\mathfrak{b}_c$ gives an isogeny (or zero) from C to C_c, and every element of $u_c^{-1}\mathfrak{b}_c^{-1}$ gives one in the reverse direction. The isogeny integrality proved after Proposition 12.6 implies that every such multiplier is integral at every prime. The product of these fractional modules over all c is $u_*\mathcal{O}_K$, with $u_* = \prod_c u_c$. The product of the inverse modules is $u_*^{-1}\mathcal{O}_K$. Since products and sums of integral elements are integral, both u_* and its inverse are integral at every prime and every embedding. The u_c themselves are algebraic: a nonzero multiplier gives an algebraic isogeny between elliptic curves over $\overline{\mathbb{Q}}$, and its effect on the invariant differential belongs to $\overline{\mathbb{Q}}$. Dividing by an element of $\mathfrak{b}_c \setminus \{0\}$ proves this for u_c. Here we use the usual algebraicity of homomorphisms and torsion points of elliptic curves, or equivalently the description of isogenies via finite subgroups and quotients. Consequently u_* is an algebraic unit.

Write A_C^γ and A_c^γ for the period areas of the normalized curves obtained by conjugating their algebraic equations by a field automorphism γ of $\mathbb{C}$ over K. Algebraic isogenies and their differential multipliers are carried to the corresponding conjugate isogenies. Hence the

conjugate lattices satisfy $\Lambda_c^\gamma = \gamma(u_c)\mathfrak{b}_c\Lambda^\gamma$: for each individual multiplier the containment follows from the corresponding isogeny, and the reverse containment follows from the inverse module of multipliers, since $\mathfrak{b}_c\mathfrak{b}_c^{-1} = \mathcal{O}_K$. It follows that

$$\prod_c A_c^\gamma = (A_C^\gamma)^{h_K}|\gamma(u_*)|^2. \tag{12.11}$$

The curves C_c^γ have the same underlying ideal-class family, with their prescribed individual differential scalings. In particular the product formula (12.11) does not require choosing a period scaling separately for each embedding.

Put $k_0 = 2/\sqrt{d}$. For a complex lattice L of area A_L and discriminant $\Delta_L = g_2^3 - 27g_3^2$, the classical first Kronecker limit formula gives

$$(s-1)\sum_{l\in L\setminus\{0\}} \left(|l|^2/(k_0A_L)\right)^{-s} = \pi k_0 + \pi k_0(s-1)\left(2\gamma_{\mathrm{E}} - \log\frac{A_L}{k_0\pi^2} - \tfrac{1}{6}\log|\Delta_L|\right) + O((s-1)^2). \tag{12.12}$$

Here γ_{E} denotes Euler's constant. To check the normalization, write $L = \lambda(\mathbb{Z}+\tau\mathbb{Z})$ with $y = \Im\tau > 0$. Its area is $|\lambda|^2 y$ and its discriminant has modulus $(2\pi)^{12}|\lambda|^{-12}|\eta(\tau)|^{24}$. The usual constant term in the nonholomorphic Eisenstein sum of weight zero is $2\pi(\gamma_{\mathrm{E}} - \log 2 - \log(\sqrt{y}|\eta(\tau)|^2))$ and its residue is π. Multiplication by k_0^s adds $\pi k_0 \log k_0$ to the constant term and gives (12.12). We use the classical Kronecker limit formula here (Siegel, *Advanced Analytic Number Theory*, I, Theorem 1), whose exact lattice formulation is `kronecker_limit` in `Lib/CM/KroneckerLimit/LogTheta.lean`. Changing the discriminant by a factor of $(2\pi)^{12}$ without compensating in this formula would disrupt the later level cancellation.

For a fractional ideal $\mathfrak{b}$ the map $y \mapsto (y)\mathfrak{b}^{-1}$ identifies its nonzero elements, modulo the finite unit group, with integral ideals in class $[\mathfrak{b}]^{-1}$. If $L = \lambda\mathfrak{b}$, then $|\lambda y|^2/(k_0A_L) = |N_{K/\mathbb{Q}}(y)|/\mathcal{N}(\mathfrak{b})$, the norm of this integral ideal. Consequently the sum of the left sides of (12.12) over the lattices Λ_c^γ is

$$2(s-1)\zeta_K(s) = 2(s-1)\zeta(s)L(s,\chi_K),$$

as the unit group has order two ($d > 3$). This is an exact identity for $s > 1$. Taking the constant and first coefficients at $s = 1$ gives $h_K\pi k_0 = 2L(1,\chi_K)$ and

$$h_K\pi k_0\left(2\gamma_{\mathrm{E}} - \frac{1}{h_K}\sum_c \log A_c^\gamma + \log(k_0\pi^2) - \frac{1}{6h_K}\sum_c \log|\gamma(\Delta_c)|\right) = 2\big(\gamma_{\mathrm{E}}L(1,\chi_K) + L'(1,\chi_K)\big).$$

Divide by the common positive residue. With $\kappa_K = \gamma_{\mathrm{E}} - \Re L'(1,\chi_K)/L(1,\chi_K)$ this yields

$$\exp\left(\frac{1}{h_K}\sum_c \log A_c^\gamma + \frac{1}{6h_K}\sum_c \log|\gamma(\Delta_c)|\right) = \frac{2\pi^2}{\sqrt{d}}\exp(\kappa_K). \tag{12.13}$$

Proposition 12.11 (CM normalization by a single algebraic unit)**.** *There exists an algebraic unit u_0 such that for every $\gamma \in \mathrm{Aut}(\mathbb{C}/K)$,*

$$A_C^\gamma|\gamma(u_0)|^2 = \frac{2\pi^2}{\sqrt{d}}\exp(\kappa_K). \tag{12.14}$$

Proof. By Proposition 12.1, each Δ_c is an algebraic unit. Choose a nonzero root

$$u_0^{12h_K} = u_*^{12}\prod_c \Delta_c.$$

A root of an algebraic unit and its inverse are integral at all primes, so u_0 is an algebraic unit. By (12.11), the expression on the left in (12.13) is $A_C^\gamma|\gamma(u_*)|^{2/h_K}\prod_c|\gamma(\Delta_c)|^{1/(6h_K)} = A_C^\gamma|\gamma(u_0)|^2$. This proves (12.14). □

This is the exact assertion of `CMLattice.chowla_selberg` in `Lib/CM/FormalGroup/FormalHeight.lean`. Notice the two elements of the argument: the analytic limit computes the mean archimedean area, while the ideal-class ratios and the isogeny integral scales show that this mean is reached using an actual algebraic unit. A formula merely describing the product of the areas of conjugates, without showing the unit-adjustment property, would not establish the individual simultaneous norm comparison of the local jet theorem.

12.7 The finite idele and the period scale

Let ζ be a primitive N_1-st root of unity, $D = \Phi'_{N_1}(\zeta)$. We use the distinction between the relative different of E/K, generated by D, and the absolute different, generated up to a unit by $\sqrt{-d}D$. Define the finite idele α by

$$\alpha_w = \begin{cases} D^{-1} & c_0 \in w, \\ 1 & c_0 \notin w. \end{cases}$$

This is the idele α (namespace `Parameters`) of `New/Data/WeilDataOfE.lean`; the notation $c_0 \in w$ means that the integer belongs to the finite prime ideal of E. All rational primes dividing N_1 divide $n_0 = (d+1)/4 = c_0(1-c_0)$. At a prime w above such a rational prime exactly one of c_0 and $\bar{c}_0 = 1 - c_0$ vanishes modulo w. The relative different generator divides N_1, and its valuation is invariant under Galois automorphisms over $\mathbb{Q}$ because the relative different ideal is preserved. Hence at every finite place w, $\alpha_w \bar{\alpha}_w D$ is a unit. Also α^{-1} and $N_1\alpha$ are integral finite adeles. These are `v_alpha_mul_relDifferentGen`, `inv_alpha_mem` and `N1_mul_alpha_mem` in their typographically analogous source declarations. As a consequence the fractional lattice

$$\mathcal{I} = E \cap \alpha\widehat{\mathcal{O}}_E$$

contains $\mathcal{O}_E$ and is contained in $N_1^{-1}\mathcal{O}_E$. Moreover $x\bar{y}D$ is integral whenever $x, y \in \mathcal{I}$; locally this follows by multiplying the two bounds and using $\alpha\bar{\alpha}D$ a unit. By the trace-dual characterization of the relative different, $\mathrm{Tr}_{E/K}(x\bar{y}) \in \mathcal{O}_K$. In particular the Hermitian trace pairing on $\mathcal{I}$ takes integral values. This proves the lattice assertions in `herm_mem_quadInt` in `New/Curve/TightFrameExists.lean`.

The finite splitting character in the Weil datum is denoted by χ_{fin}. Its necessary property can be stated as follows without any archimedean topology on the field of complex values. It is trivial on integral finite units; on a principal finite idele y one has

$$\chi_{\mathrm{fin}}(y) = \prod_{\tau \in \Phi} \tau(\bar{y}). \tag{12.15}$$

This is the type of the normalized finite character, not the unitary Hecke character prior to the absolute-value factor. The compatible splitting character is constructed in Proposition 6.4 and its finite normalized action in Theorem 7.6. The remaining comparison with the archimedean arithmetic theta model is a specific input; see `Parameters.instWeilDataE` in `New/Data/WeilDataOfE.lean` and `nonempty_splittingChar` in `Lib/ClassField/HeckeCharacter/SplittingCharExists.lean`. For the following normalization computation it suffices to assume exactly (12.15) and triviality on the finite unit subgroup. Thus this input is separated from the estimates about theta coefficients.

If z is any finite idele, some positive power z^h has the same valuations as a principal idele y, by finiteness of the ideal class group. Equation (12.15) implies

$$\chi_{\mathrm{fin}}(z)^h = \prod_{\tau} \tau(\bar{y}).$$

The latter is in $K^\times$. Taking absolute values of all its images under a p-adic embedding ι and extracting the positive h-th root proves

$$|\iota(\chi_{\mathrm{fin}}(z))|_p = \prod_{\tau\in\Phi} |z|_{\iota\tau\circ\mathrm{conj}}. \tag{12.16}$$

In the factor on the right, the component of z at the place induced by the indicated embedding is read in the local field. It depends only on the valuation of that component. This establishes `exists_chiFin_pow_eq_prod` and `norm_map_chiFin` of `Lib/ClassField/HeckeCharacter/SplittingCharOnU1.lean` directly from the ideal-class argument and character type.

Let s_0 be a nonzero f-th root of $\chi_{\mathrm{fin}}(\alpha)^{-1}$. The ideal of $\alpha\bar\alpha$ is the inverse relative different. The absolute norm of that different is $(\mathrm{rd}_E/\sqrt{d})^{2f}$. Taking norms of both sides shows that the ideal norm of α is $(\mathrm{rd}_E/\sqrt{d})^{-f}$. If α^h has the valuations of the principal ideal (y), it follows that $|N_{E/\mathbb{Q}}(y)| = (\mathrm{rd}_E/\sqrt{d})^{-hf}$. By (12.15), the squared ordinary absolute value of $\chi_{\mathrm{fin}}(\alpha)^h$ is this rational ideal norm. Since the value on the right of (12.15) is in K, any field automorphism γ over K fixes it. Thus, taking positive roots,

$$|\gamma(s_0)|^2 = \mathrm{rd}_E/\sqrt{d}. \tag{12.17}$$

The compact derivation is `Parameters.norm_gal_vacuumRoot` in `New/Curve/PeriodDefs.lean`. At a fixed p and ι, the norms of the components of α read through $\iota\tau\circ\mathrm{conj}$ are the same for all $\tau\in\Phi$. They are either all one or all equal to $|\iota(D)|_p^{-1}$: the embeddings τ extend the fixed embedding of K, and the valuations of the conjugate generators of the relative different coincide. Dividing (12.16) for $z=\alpha$ by f in logarithms gives $|\iota(s_0)|_p = |\alpha|_{\iota\tau\circ\mathrm{conj}}^{-1}$.

Now let u_0 be as in Proposition 12.11 and $s_1 = s_0/u_0$. Its finite valuations are the same as those of s_0 because u_0 is a unit. Therefore for every $y\in\mathcal{I}$,

$$|\iota(s_1\overline{\tau(y)})|_p \le 1, \qquad |\iota(s_1)|_p = 1 \quad (p\nmid N_1). \tag{12.18}$$

This is exactly the finite normalization required in Proposition 13.2, with the stronger equality of local norms of s_1 and the inverse component of α. It proves `norm_iota_periodFactor`, `norm_periodFactor_mul_conj_le_one` and `norm_iota_periodFactor_eq_one` in `New/Curve/TightFrameExists.lean`, from the splitting character property and the classical CM results. In particular the primitivity condition of the tight frame cannot be obtained by adjusting s_1 at each prime; it has to be built into the frame itself.

Finally, if a conjugate normalized curve has area $|\Omega_\gamma|^2\sqrt{d}/2$ and $t_\gamma = \gamma(s_1)(2\pi/\sqrt{d})^{1/2}/\Omega_\gamma$, then (12.17) and (12.14) give

$$|t_\gamma|^2 = \frac{\mathrm{rd}_E}{2\pi e^{\kappa_K}} = \Omega^{-1}. \tag{12.19}$$

This is the numerical relation used for the conjugate archimedean estimates. It shows exactly how the discriminant and logarithmic derivative in the main height cost enter: the normalized complex Hermitian coordinates have scale $(2\pi/\sqrt{d})^{1/2}$, whereas the CM multiplier is normalized by a unit using the mean area. Their ratio, after the finite different adjustment, is a fixed positive real number independent of the conjugate curve.

12.8 Constructing arithmetic tight frames of prescribed length

We now prove the arithmetic frame assertion of Proposition 13.10. This step deserves a separate proof: a family of lattice vectors may be a perfectly good tight frame without the length being a p-adic unit, and may have every individual vector too small in the coefficient normalization s_1. Both problems arise in the theta estimates. Write $N = N_1$, not the auxiliary

level prime, and $\zeta = \zeta_N$ as above. Throughout the construction $n_0 = (d+1)/4$. We have $p \mid n_0$ for every rational prime $p \mid N$. Recall that a tight frame is a family $w_i \in \mathcal{I}$ with positive integer multiplicities m_i such that

$$\sum_i m_i \sigma(w_i)\overline{\tau(w_i)} = M\mathbf{1}_{\sigma=\tau} \quad (\sigma, \tau \in \Phi). \tag{12.20}$$

The parameter M is called the frame length; the cardinality of the indexing family need not equal M. For example the N roots ζ^j, each with multiplicity one, give length N by the geometric sum $\sum_{j=0}^{N-1}(\sigma\zeta/\tau\zeta)^j$. A direct sum of two frames with lengths M_1, M_2 has length $M_1 + M_2$, and multiplying all multiplicities by an integer multiplies the length. We seek another frame of length coprime to N, with additional local unit vectors in this family of frames.

Lemma 12.12 (A cyclotomic factor of the different)**.** *For every rational prime $\ell \mid N$, put $r_\ell = N/\ell$, $A_\ell = \zeta^{r_\ell}$, and let $M_\ell = N/\ell^{v_\ell(N)}$. Then*

$$V_\ell = \frac{D(A_\ell - 1)M_\ell}{N} \in \mathcal{O}_E, \qquad V_\ell \mid M_\ell.$$

In particular V_ℓ is an ℓ-adic unit at every embedding above ℓ.

Proof. The standard cyclotomic derivative formula follows from $\Phi_N(X) = \prod_{a|N}(X^a - 1)^{\mu(N/a)}$ by differentiating and evaluating at ζ. Up to the unit ζ^{-1} it gives

$$D = N\zeta^{-1} \prod_{a|N,\ a<N} (\zeta^a - 1)^{\mu(N/a)}.$$

For any prime above a rational prime p, $\zeta^a - 1$ is a unit unless the order N/a of ζ^a is a power of p. Indeed if that order has nontrivial prime-to-p part, its reduction has that same prime-to-p part; otherwise the valuations of $1 - \zeta_{p^k}$ are the familiar $1/(p^{k-1}(p-1))$, which follow from $\Phi_{p^k}(1) = p$ and conjugacy of the factors. Since the Möbius exponent vanishes for a nonsquarefree integer, the only nonunit factor of the product at p occurs when $N/a = p$. Hence $v_p(D) = v_p(N) - 1/(p-1)$ when $p \mid N$, and zero elsewhere, uniformly in the embedding. The factor $A_\ell - 1$ has valuation $1/(\ell-1)$ above ℓ and zero elsewhere. Thus $v_p(V_\ell)$ is zero above ℓ and outside the primes dividing N, and equals $v_p(M_\ell) - 1/(p-1)$ above each $p \mid M_\ell$. These are nonnegative numbers at most $v_p(M_\ell)$, proving both assertions by the characterization of algebraic integers by their finite valuations. □

The source `Parameters.exists_cyclotomic_different_factor` in `New/Curve/TightFrameExists.lean` proves this by a polynomial factorization with a bound $M_\ell \le N$. The explicit prime-to-ℓ part above is one convenient choice satisfying its conclusions.

Since n_0 is divisible by all prime divisors of N, it is nilpotent modulo N. The map $z \mapsto n_0 z^2 + z$ on $\mathbb{Z}/N\mathbb{Z}$ is injective: the difference of its values at z, z' is $(z - z')(1 + n_0(z + z'))$, and the second factor is a unit modulo N. It is therefore surjective. Choose z_0 with $0 \le z_0 < N$ such that

$$n_0 z_0^2 + z_0 + 1 = eN, \qquad e = (n_0 z_0^2 + z_0 + 1)/N \in \mathbb{N}.$$

The integer e is positive and $e \le n_0 N + 1$. Define the integer of K

$$b_0 = 1 - c_0 + n_0 z_0.$$

Then $b_0 + \bar{b}_0 = 1 + 2n_0 z_0$, and calculating the norm with $c_0\bar{c}_0 = n_0$ gives $b_0\bar{b}_0 = n_0 eN$. In particular b_0 is a unit at the places above $p \mid N$ where c_0 vanishes, while at the other places

above those primes it is divisible by N: its conjugate is a unit there and its norm is a multiple of N.

For a rational prime $\ell \mid N$ take the following family, indexed by $0 \le j < r_\ell$ and $0 \le k, i < \ell$:

$$w_{\ell,j,k,i} = \zeta^j \left(\mathbf{1}_{k=0} A_\ell^i + \frac{b_0 M_\ell}{N} (A_\ell^i - A_\ell^k) \right). \tag{12.21}$$

This is an arithmetic modification of the usual cyclotomic frame. Both types of lattice condition are satisfied. For a finite prime above a rational prime outside N, all denominator factors are units. At a prime above N where c_0 does not vanish, b_0 is divisible by N, so the vector is integral. At a prime where c_0 vanishes, $A_\ell^i - A_\ell^k$ is an integral multiple of $A_\ell - 1$; multiplying the second term by D gives the integral factor $b_0 V_\ell$ times that multiple. Thus $D w_{\ell,j,k,i}$ is integral there. This is precisely membership in $\mathcal{I}$, whose local allowed denominator is D in the latter case.

Lemma 12.13 (Length of the modified cyclotomic frame)**.** *The family* (12.21)*, with multiplicities one, is a tight frame of length*

$$C_\ell = N + M_\ell(1 + 2n_0 z_0 + 2\ell n_0 M_\ell e). \tag{12.22}$$

Moreover for each $p \mid N$ the family with $\ell = p$ contains a vector $w \in \mathcal{I}$ with $|\iota(s_1 \overline{\tau(w)})|_p = 1$ at all τ, for each ι for which $|\iota(s_1)|_p < 1$.

Proof. Set $T = b_0 M_\ell / N$. For any two complex embeddings $\sigma, \tau \in \Phi$, let $A = \sigma(A_\ell)$ and $B = \tau(A_\ell)$. Both are primitive ℓ-th roots. The sums over i, k in the inner product of (12.21) can be expanded into four terms. The products of the two terms with $k = 0$ sum to $\ell \mathbf{1}_{A=B}$. Each mixed product is $\ell T \mathbf{1}_{A=B}$ or $\ell \bar{T} \mathbf{1}_{A=B}$: all sums with just one nontrivial power of A or B vanish. Finally

$$\sum_{i,k} (A^i - A^k)(\overline{B^i} - \overline{B^k}) = 2\ell^2 \mathbf{1}_{A=B}.$$

Multiplying by the remaining j-sum gives $\sum_j (\sigma\zeta/\tau\zeta)^j$, which is r_ℓ if $\sigma\zeta = \tau\zeta$ and zero if $A = B$ but the embeddings differ (in the latter case its ratio is a nontrivial r_ℓ-th root of unity). Thus the full matrix of products equals $N(1 + T + \bar{T} + 2\ell T\bar{T})$ times the identity. Substituting the trace and norm of b_0 gives (12.22).

For the second assertion take $\ell = p$, $j = 0$, $k = 1$, $i = 0$. The chosen vector is $w = (b_0 M_p/N)(1 - A_p)$, and $Dw = -b_0 V_p$. If $|\iota(s_1)|_p < 1$, the relevant components of α above p read through every $\iota\tau \circ \mathrm{conj}$ are D^{-1}. Thus all those places contain c_0. The product $-b_0 V_p$ is a unit at every such place, by the definition of b_0 and Lemma 12.12. Equation (12.18) with its equality of finite scale factors proves the claimed unit norm. □

Let $C = \sum_{\ell \mid N} C_\ell$, the length of the union of the modified frames. It is positive and coprime to N. Indeed for every $p \mid N$, the terms with $\ell \neq p$ in the sum are zero modulo p (since $p \mid M_\ell$), while the term with $\ell = p$ is $M_p \not\equiv 0 \pmod{p}$ (since $p \mid n_0$). This proves the coprimality at each prime divisor.

Proposition 12.14 (A prescribed-length frame with a simultaneous unit vector)**.** *If $M \ge (dN)^6(R+1)$ with $R \ge 0$, there is a tight frame of length M with multiplicities at least R and containing, for every rational prime p and every ι, some vector w such that $|\iota(s_1 \bar{\tau}(w))|_p = 1$ for all $\tau \in \Phi$.*

Proof. We first bound the lengths sufficiently crudely. The number of distinct prime factors of N is at most N, and each M_ℓ, ℓ, z_0 is at most N. Together with $e \le n_0 N + 1$ this gives

$C \le 8(n_0+1)^2N^5$, hence $NC \le (dN)^6$ (recall $n_0 = (d+1)/4$, $d > 3$). Also $C \ge N$ and $C + N \le NC$ as $N \ge 2$. Consequently

$$M \ge R(C+N) + (C-1)(N-1).$$

Since C, N are coprime, every integer $L \ge (C-1)(N-1)$ is a nonnegative integer combination $AC + BN$. To verify the numerical assertion, choose $B \in \{0, \dots, C-1\}$ with $L \equiv BN \pmod{C}$; then $L - BN \ge -(C-1)$ and is divisible by C, hence nonnegative. Apply this to $L = M - R(C+N)$. Take the union of all the modified frames with multiplicities $R + A$ and of the elementary cyclotomic frame with multiplicities $R + B$. Its length is M and all multiplicities are at least R. The elementary frame contains 1; this gives the required assertion whenever $|\iota(s_1)|_p = 1$. Otherwise $p \mid N$ by (12.18), and the vector in Lemma 12.13 gives it. If positive multiplicities on both families rather than nonnegative ones are desired in the case $R = 0$, observe that the same crude bound improves strictly to $NC \le (dN)^6/4$ for the present d, N; we may first subtract $C + N$ and still satisfy the numerical representability condition. Thus one can take both family multiplicities positive for all these large lengths, without changing the stated bound. □

This proves `Parameters.exists_isTightFrame` in `New/Curve/TightFrameExists.lean`, and explains the two roles of the modified cyclotomic family. Its length, not the roots of unity individually, removes a divisibility obstruction; its special difference vector can compensate exactly the finite different scale at every embedding above a given prime. The proof gives a deterministic construction using algebraic integers and ideal norms; no approximation of an arbitrary archimedean isometry by rational matrices is necessary.

12.9 Frobenius on the frame matrix

The reduction in the Frobenius step is not always to a field. It is important to establish the equality in a quotient of a valuation ring, rather than infer a stronger assertion from an equality of residue-field values. Let p be nonsplit, let v_p have $v_p(p) = 1$, and set

$$\omega_p = \begin{cases} 1 & p \text{ inert}, \\ 1/2 & p = d, \end{cases} \qquad J_p = \{x \in \mathfrak{o} : v_p(x) \ge \omega_p\}.$$

The quotient $\mathfrak{o}/J_p$ is a ring of characteristic p. Since all primes dividing N split in K, N is a unit in this quotient, and $\mathcal{I}$ is integral at p.

Lemma 12.15 (Reduction of conjugate embeddings)**.** *Let $\sigma \in \Phi$, and let e be odd if p is inert, arbitrary nonnegative if $p = d$. For every $w \in \mathcal{I}$ one has*

$$(\iota\overline{\sigma(w)})^{p^e} = \iota\sigma'(w) \pmod{J_p}, \qquad \sigma' = [-p^e]\sigma.$$

Here the bracket indicates the action on the cyclotomic embeddings: $\sigma'(\zeta) = \sigma(\zeta)^{-p^e}$, while $\sigma'|_K$ is the fixed CM embedding.

Proof. The cyclotomic field and K have coprime discriminants, so $\mathcal{O}_E = \mathcal{O}_K[\zeta]$. It suffices to check the identity on $\mathcal{O}_K$, ζ, and N^{-1}, since $\mathcal{I} \subseteq N^{-1}\mathcal{O}_E$ and both sides are ring homomorphisms to a ring of characteristic p. At an inert prime, the subring from $\mathcal{O}_K$ maps through $\mathcal{O}_K/(p) = \mathbb{F}_{p^2}$. There the p-power map is conjugation. Thus applying an odd number of Frobenius maps after conjugation returns the original CM embedding. At the ramified prime, $\sqrt{-d}$ belongs to J_p, and the two embeddings of K reduce to the same embedding of $\mathcal{O}_K/(\sqrt{-d}) = \mathbb{F}_p$. Consequently every power of Frobenius acts identically on that subring,

including the zeroth power. In either case the identity on ζ follows from $\overline{\sigma(\zeta)} = \sigma(\zeta)^{-1}$, and the identity on N^{-1} from the fact that the prime field is fixed. This verifies the claimed equality in the specified nonreduced quotient as well as in the residue field. □

Proposition 12.16. *Repeat the vectors of a frame according to their multiplicities, so that* $\sum_i \sigma(w_i)\overline{\tau(w_i)} = M\mathbf{1}_{\sigma=\tau}$. *For an integral scalar* β, *define* $r_{\sigma i} = \beta\iota(\overline{\sigma(w_i)})$. *In every ring of characteristic* p *obtained from* $\mathfrak{o}$ *by a homomorphism annihilating* J_p *one has*

$$\sum_i r_{\sigma i}^{p^e} r_{\tau i} = \beta^{p^e+1} M \mathbf{1}_{\tau=[-p^e]\sigma}$$

for precisely the exponents in Lemma 12.15.

Proof. Factor out β^{p^e+1} and apply Lemma 12.15 to the first factor of each product. The remaining sum is the image of $\sum_i \sigma'(w_i)\overline{\tau(w_i)}$, which is the original frame identity with its first row replaced by σ'. □

Take $\beta = N\iota(b')$ in the normalization of (13.13). This proves Lemma 13.12, the statement `Parameters.exists_frame_rows_frob` of `New/Curve/ThetaProducts.lean`. In particular the minus sign in the permutation is not an arbitrary transpose or sign convention. The complex conjugation in the columns is essential, and for inert primes it cancels the odd Frobenius on the quadratic subfield while it reverses the cyclotomic exponent. The restriction on the live set of exponents is therefore intrinsic to the use of the half-set of CM embeddings.

12.10 Theta-group orthogonality and the finite realization

The following general linear-algebra argument isolates the substantial, but classical, mechanism in the finite expansion of a vacuum. Let L be a full lattice in a complex vector space of dimension g, and let B be a positive Hermitian form whose associated Riemann form $\operatorname{Im} B/\pi$ is integral on L. Use the normalization of the theta shift

$$U_a S(z) = \exp\bigl(-B(a,a)/2 - B(a,z)\bigr) S(z+a)$$

with B conjugate-linear in its first slot (equivalently one can use the conjugate convention written in (12.3)). Assume an Appell–Humbert multiplier has been fixed, and let $L^\vee$ be the integral dual for the resulting alternating form. Then U_a for $a \in L^\vee$ preserves the theta space H, and the shifts for $a \in L$ act by scalars. The alternating form induces a perfect alternating pairing on the finite abelian group $D = L^\vee/L$. The dimension of H is $m = \sqrt{|D|}$, by the usual Riemann–Roch dimension theorem for positive theta spaces; this classical theorem is also obtained by writing theta series in a symplectic basis of L. We use only its dimension formula here.

Lemma 12.17 (Theta-group averaging)**.** *Let* R_x *be a representative operator of the finite theta group on* H *for each* $x \in D$, *and denote its inverse by* R_{-x} *in the inverse-pair convention for shifts. Then for* $F, v \in H$ *and every linear functional* ℓ *on* H,

$$m \sum_{x \in D} \ell(R_x v) R_{-x} F = |D| \ell(F) v. \tag{12.23}$$

Proof. For clarity we establish the averaging formula directly. The commutator pairing on D is nondegenerate. Therefore for $x \neq 0$ some R_y conjugates R_x to a nontrivial scalar multiple of itself, so $\operatorname{tr} R_x = 0$. Also $\operatorname{tr}(R_x^{-1} R_{x'}) = 0$ for $x \neq x'$ and equals m for $x = x'$. These identities make the $|D| = m^2$ operators R_x a basis of $\operatorname{End}(H)$. The operator $\sum_x R_x^{-1} A R_x$ commutes

with every R_y because the central factors cancel in conjugation. Hence it is scalar. Its trace is $|D|\operatorname{tr}(A)$, so it equals $|D|\operatorname{tr}(A)I_H/m$. Substitute the rank-one operator A and apply the equality to v. Since $\operatorname{tr} A = \ell(F)$ this proves (12.23). In particular the span of $R_x v$ for $v \neq 0$ is H: the displayed matrix basis shows that any vector can be obtained by applying a linear combination of these operators to v. □

In the intended application the vector space H is an archimedean theta space, and we require its realization in the finite Weil model. We state explicitly which facts about the comparison of these two models we use without reconstructing their entire representation theory. For the standard space of the manuscript, put $g = fh$, where $f = |\Phi|$ and h is the rank. Write Θ for the linear map from the finite part to its entire Fock theta functions and v for the vacuum. We use the following formal-source input from `Lib/Weil/Construction/FockThetaFunctions.lean`: Θ is injective, $\Theta v \neq 0$, and

$$\Theta\big(\varrho(\operatorname{finImage}\gamma)F\big) = U_{-\operatorname{realImage}\gamma}(\Theta F). \tag{12.24}$$

The last identity is `entireTheta_translate_finiteImage`; the comparison of multipliers is `semichar_finiteImage`. The archimedean multiplier is the canonical one defined by the quadratic integer used in the construction of the standard lattice. The local lattice is self-dual for the finite additive character, and rational finite points are dense. These assertions explain why the finite and real shifts give inverse central characters. We retain these exact model-comparison identities as formal-source inputs (they can be verified in the integral model by the Gaussian integral and the additive-character product formula). The remainder of the realization is elementary theta-group theory.

To see it, let $\delta \in \mathcal{O}_K$ with $D_\delta = \delta\bar{\delta} > 0$. Let L be the archimedean image of $\bar{\delta}\mathcal{I}^h$ with the sign and real scale used in (12.24). The function Θv belongs to its theta space: for $\gamma \in \mathcal{I}^h$, invariance of the vacuum under the finite lattice gives exactly the canonical multiplier of the shift by $-\operatorname{realImage}(\bar{\delta}\gamma)$. Every dual point $a \in L^\vee$ comes from a rational point γ_a because some positive integer multiple of a belongs to L. In fact $\delta\operatorname{finImage}(\gamma_a)$ is in the finite lattice. One can see this by commuting its finite shift with all finite lattice shifts from $\bar{\delta}\mathcal{I}^h$: the scalar commutator is the inverse real pairing, which is trivial by duality. Density extends this statement from rational lattice points to the finite lattice, and self-duality gives the asserted lattice condition. Conversely the image of these rational points is in $L^\vee$ by the same argument. Thus every dual translation of Θv belongs to the image of Θ in (12.24). By Lemma 12.17 these translations span the theta space. In particular any entire S satisfying these multiplier conditions lies in the image of a unique finite vector.

In particular $D_\delta L^\vee \subseteq L$: the dual rational representative satisfies $\delta\gamma_a \in \mathcal{I}^h$, so $D_\delta\gamma_a = \bar{\delta}(\delta\gamma_a) \in \bar{\delta}\mathcal{I}^h$. We can therefore average over the larger finite group $L^\vee/D_\delta L^\vee$. An element of the smaller theta group is counted a constant number of times, and the central factors cancel in the two inverse shifts. The order of the larger group, since $L^\vee$ has rank $2g$, is D_δ^{2fh}. If I indexes its representatives, they give finite coordinates αx_j with δx_j integral, exactly as in Proposition 13.11. Transport (12.23) through the injective map Θ and put $M_\delta = m$. This yields

$$\sum_{j\in I} M_\delta \ell(\varrho(\alpha x_j)v)\varrho(-\alpha x_j)F = D_\delta^{2fh}\ell(F)v,$$

for any linear functional on the finite part, not only for the theta functional. This is the finite vacuum expansion in `Space.VacuumExpansion`, `Lib/Weil/Construction/ConjugateJetDefs.lean`. The argument recovers `Space.exists_entireTheta_eq` from the stated finite–archimedean comparison and the usual positive theta dimension formula. The source's `ThetaOrthogonality.lean` develops the theta-group orthogonality directly using entire functions as well as bilinear pairings.

12.11 From an arithmetic frame to a theta section

We apply the preceding realization to a product of one-dimensional CM sections and record the area normalization. Let C be a normalized CM curve with area A, $a \in \Lambda_C$, $\delta\bar{\delta} = D_\delta > 0$, and let (w_i, m_i) be a tight frame of length M. For each coordinate $k = 1, \dots, h$ choose a degree-m_i monomial section G_{ik} on C. Set $s_{\mathrm{ar}} = \sqrt{2\pi/\sqrt{d}}$ and

$$L_{ik}(z) = \frac{a}{\bar{\delta} s_{\mathrm{ar}}} \sum_{\tau \in \Phi} \overline{\tau(w_i)} z_{\tau k}, \qquad S(z) = \prod_{i,k} G_{ik}(L_{ik}(z)).$$

In the normalization of this chapter the polarization form of the degree-one curve section is $B_C(u, v) = \pi \bar{u} v / A$. The arithmetic condition is the identity of areas

$$A = \frac{|a|^2}{D_\delta} M \frac{\sqrt{d}}{2}. \tag{12.25}$$

Indeed the sum of pullback forms, with their degrees, is

$$\sum_{i,k} m_i B_C(L_{ik}u, L_{ik}v) = \frac{\pi |a|^2 M}{A D_\delta s_{\mathrm{ar}}^2} \sum_{\tau,k} \overline{u_{\tau k}} v_{\tau k} = \sum_{\tau,k} \overline{u_{\tau k}} v_{\tau k},$$

by the frame identity and (12.25). This identity is necessary for the product S to realize the standard theta function, rather than one in a different polarization.

For each translated period $-\operatorname{realImage}(\bar{\delta}\gamma)$ with $\gamma \in \mathcal{I}^h$, its image under the linear form is $-a \operatorname{Tr}_{E/K}(\gamma_k \bar{w}_i)$, an element of Λ_C by the trace pairing integrality proved above. The multiplier of a pullback of a theta section is the pullback of its multiplier. Its canonical factor is expressed using the imaginary part of the polarization evaluated on the quadratic multiplication by $\bar{c}_0$ (as in the earlier computation of the multiplier on the basis of the lattice). The L_{ik} intertwine this multiplication, since the embeddings in Φ restrict to the fixed embedding on K. Products of the canonical multipliers are therefore the canonical multiplier for the sum of the pullback forms. Consequently S has exactly the required multiplier, and the preceding finite realization gives a unique F with $\Theta F = S$ and the vacuum expansion.

For any rational vector γ put

$$y_{ik} = -\frac{a}{\bar{\delta}} \operatorname{Tr}_{E/K}(\gamma_k \bar{w}_i).$$

Under (12.24) the entire theta function of the translate of F by γ is $\prod_{i,k} (U_{y_{ik}} G_{ik})(L_{ik} z)$, since the exponential factors in the shifts multiply using the equality of the polarization forms. Taking homogeneous Taylor coefficients is legitimate: at each degree the product is a finite sum, and the Taylor series along all complex lines determines the multivariable Taylor series. This gives both the product Taylor identity and the theta evaluation asserted in Proposition 13.11.

For integrality choose a positive integer n clearing the finite denominators of γ: $n\gamma \in \mathcal{I}^h$. The trace integrality shows that $n D_\delta y_{ik} = -\delta a \operatorname{Tr}_{E/K}(n\gamma_k \bar{w}_i) \in \Lambda_C$. Thus y_{ik} is torsion. The normalized curve monomial sections have norm at most one at every finite valuation and the height of zero is one. The torsion translation estimate above therefore gives $|\iota(U_{y_{ik}} G_{ik})(0)|_p \le 1$ at every embedding, and hence the product theta evaluation has norm at most one. The finite model identifies its point evaluations with the theta evaluations of its finite translates, via `Space.θ_ϱ` (theta-rho evaluation) in the notation of the source. All translates of a fixed finite vector are locally constant in the finite adelic variable. By density each finite translate agrees with one by a rational point, so the same integral bound holds at every finite point. This proves

the product and integrality parts of `Parameters.exists_finitePart_of_isTightFrame` in `New/Curve/FrameSections.lean`, conditional on the stated representation comparison and the standard theta dimension theorem. It is insufficient for this argument to check integrality at the origin alone without controlling rational torsion translates.

As a useful special case, take w_i to be all the powers of ζ, each repeated N times, giving length N^2, and take $\delta = N$, $a = \Omega_0$ and $C = \mathbb{C}/\Omega_0\mathcal{O}_K$. These choices exactly satisfy (12.25). Choose every section equal to the normalized degree-one theta section, which has constant value one. The resulting F has $\theta(F) = 1$. By taking the theta functional in the vacuum expansion we express v as a finite linear combination of shifts of F. Since the finite part is by definition the span of translates of v, translates of F also span the entire finite part. This is the generator in the first part of the uniform conjugation assertion.

12.12 Integral vacuum and the remaining reciprocity law

It is useful to distinguish what is required from the complex multiplication reciprocity theorem from the elementary bound on the integral vacuum. The needed reciprocity is not merely the assertion that the j-invariant generates a ring class field. We require a comparison of normalized Taylor coefficients of theta functions and their torsion translates. This normalized reciprocity theorem is proved in Theorem 16.5. In the notation of the source this comparison is `CMLattice.complex_multiplication_theta`, used at the beginning of `Parameters.exists_isJetConjugate_generator` in `New/Curve/ConjugateGenerator.lean`. Concretely, starting with the principal normalized CM curve, it selects, for each $\gamma \in \mathrm{Aut}(\mathbb{C}/K)$, a conjugate normalized curve C_γ of area $|\Omega_\gamma|^2\sqrt{d}/2$, an integer $e_\gamma > 0$ with $e_\gamma\Omega_\gamma \in \Lambda_\gamma$, and a finite idele defining the transport of torsion coordinates. The key equality compares the coefficientwise Galois action on the twisted Taylor series with the series on C_γ after this transport and a differential rescaling. We use this equality as a formal-source input, including the necessary compatibility of the finite torsion transport for all vectors, not merely existence of some torsion point for each evaluation. The classical reciprocity theorem for algebraic elliptic curves explains the choice of C_γ; tracking normalized theta multipliers and the splitting character requires its more precise form here. We do not independently reconstruct that form of the theorem.

One can nevertheless prove from this input why the same finite generator works in every rank and why its adelic translates have the desired conjugate bounds. For C_γ use the same frame of length N^2 from the last paragraph, but put $a = e_\gamma\Omega_\gamma$ and $\delta = Ne_\gamma$. Then

$$\frac{|e_\gamma\Omega_\gamma|^2N^2}{(Ne_\gamma)^2}\frac{\sqrt{d}}{2} = |\Omega_\gamma|^2\frac{\sqrt{d}}{2},$$

so the area condition (12.25) holds exactly. The linear forms in this frame have common scalar $\Omega_\gamma/(Ns_{\mathrm{ar}})$, and after the conjugate period scale this scalar is $\gamma(s_1)/N$, with

$$t_\gamma = \frac{\gamma(s_1)s_{\mathrm{ar}}}{\Omega_\gamma}, \qquad |t_\gamma|^2 = \Omega^{-1}$$

by (12.19). The rational coefficients in each linear form are the conjugates of the original cyclotomic coefficients, with the permutation of places determined by γ. The trace formula for the y_{ik} has the same pattern on both sides; its scale is $-\Omega_\gamma/N$. Applying the normalized reciprocity comparison to these trace torsion coordinates yields the equality of the coefficients of each factor. Multiplication of the finite number of factors proves the equality for the jets of rational translates. For arbitrary finite translates, use the common finite idele furnished by the reciprocity theorem. Multiplication by this idele is continuous, and both finite-vector

orbits are locally constant, so on any prescribed pair of translates the equality reduces to a rational approximation in the corresponding common open set. This is the claimed compatibility for the whole finite orbit, rather than just the generator at zero. Together with the theta-group expansion already proved, the argument explains the second part of `Parameters.exists_isJetConjugate_generator` subject to the normalized reciprocity input.

The remaining estimate concerns the size of the scalar by which the vacuum is normalized, not the norm of a particular representative of the differential. Let F be our integral vector with $\theta(F) = 1$, let I be the finite expansion set with $|I| = D_\delta^{2fh} = N^{4fh}$, and use the standard inner product, normalized by $\langle v, v\rangle = 1$. In the standard finite Weil model, for all finite shifts $R_x = \varrho(\alpha x)$,

$$\langle v, R_x v\rangle \in \{0\} \cup \mu_\infty. \tag{12.26}$$

Here the zero is for a shift outside the lattice supporting the vacuum, and the root of unity is its prescribed finite multiplier inside that lattice. This is the vacuum overlap calculation `Space.dataInner_rho_of_vacElem` in the source. We retain it as part of the comparison with the standard finite Weil model. These are the only possible values: for a translation in the stabilizer the vacuum transforms by its finite character, and for a translation outside the lattice, commute with a lattice element on which the resulting character is nontrivial and use invariance of the inner product. Thus the identity follows from the self-duality of the stabilizing lattice and unitarity of the finite representation. It does not need a numerical integral of a complicated theta function.

Proposition 12.18 (Vacuum clearing factor)**.** *Under the stated comparison and reciprocity inputs there is a nonzero scalar c such that every finite evaluation of cv is integral at every finite place, and every K-conjugate of every such evaluation has absolute value at most* $\exp(C_0 N^2 h)$ *for a constant C_0 independent of N, h.*

Proof. First choose the theta functional in the vacuum expansion, and take its inner product with v. Since $\theta(F) = 1$ and v has norm one, some $j \in I$ has $\langle v, R_{-x_j} F\rangle \neq 0$. Set

$$\ell(W) = M_\delta^{-1}\langle v, R_{-x_j} W\rangle, \qquad c = |I|\ell(F) \neq 0,$$

where the inner product is linear in the second argument. Substitute this functional into the expansion. The result is

$$cv = \sum_{i\in I} \langle v, R_{-x_j} R_{x_i} v\rangle R_{-x_i} F.$$

By (12.26) and the Heisenberg multiplication law, the coefficients of the sum are zero or roots of unity. There is no surviving denominator M_δ: this is the purpose of the normalization of ℓ. The pointwise integrality of F proved above is preserved by finite translations (their phases are roots of unity). Hence all evaluations of cv are integral; this follows from a finite sum with integral coefficients, without having to decide in advance which evaluation of the vacuum is nonzero.

For the archimedean bound let A' be the area of any normalized conjugate curve and ϑ' its normalized theta section. The value of its torsion translate at zero has absolute value

$$|(U_y\vartheta')(0)| = \exp(-\pi|y|^2/(2A'))|\vartheta'(y)|.$$

This continuous function of y is periodic on its lattice and attains a finite maximum. The family of conjugate normalized curves has only finitely many such maxima: a_1 is algebraic, its conjugates form a finite set, and the cubic with its marked point and normalized differential determines the period lattice and normalized theta section (with the possible finite ambiguity

from fourth roots of unity preserving the value of the maximum). Write $C_1 \geq 1$ for a bound on all of them. The normalized reciprocity comparison and the previous product construction bound each conjugate of each finite translated evaluation of F by $C_1^{N^2h}$, since the product has exactly N^2h degree-one factors. The coefficients in the expression for cv remain of absolute value one or zero under every conjugation. Thus the sum has absolute value at most

$$N^{4fh}C_1^{N^2h} \leq \exp((4 + \log C_1)N^2h),$$

where $f = \varphi(N) \leq N$ and $\log N \leq N$. This proves the bound with $C_0 = 4 + \log C_1$, or any integer above it. □

The scalar chosen in the formal construction is that for rank one, and the standard vacuum in rank h is the product of the rank-one vacuum evaluations. This factorization, `Parameters.intVacuum_finVec_eq_prod` in `New/Curve/IntegralVacuum.lean`, follows from the product definition of the standard diagonal space and its vacuum. Therefore $s_{\mathrm{vac}}^h v$ is integral and its conjugates are bounded by the same expression $\exp(C_0N^2h)$, as asserted by `Parameters.norm_intVacuum_le_one` and `Parameters.norm_intVacuum_le`. The integral module of standard data is the integral span of translates of this scaled vacuum by the specified local lattice. It is stable under this lattice because the finite shift multiplication factors are roots of unity. Stability under the restriction of the finite Weil action, and the equivariance at nonintegral adelic matrix arguments, require the exact representation identities specified in Proposition 13.20 and Proposition 13.22. Neither is an automatic property of a space of power series with integral coefficients. They remain formal-source inputs: `Parameters.omega_mem_standardData`, `finHeisOp_mem_standardData` (the finite Weil vacuum sum), `Parameters.mem_raises_of_jet_bound`, and `mem_raises_of_standardData` in `New/Curve/ConjugatesOfJets.lean`.

12.13 What the arithmetic normalization proves

The preceding proofs establish the normalized good cubic, its exact local height and section algebra, the CM dilation and logarithmic identities, the formal height input from good reduction, the period comparison from the unit-adjusted Chowla–Selberg calculation and character normalization, and both the frame existence and Frobenius orthogonality. The finite realization and vacuum expansion reduce to positive theta space dimension and orthogonality, plus precisely the finite–archimedean comparison identities. Vacuum integrality and its uniform archimedean bound then follow from the normalized reciprocity law. This separates the explicit algebra in the jet theorem from the assumptions on the construction of the Weil model, its normalized CM reciprocity, and its adelic equivariance. Those assumptions are identifiable statements in the supplied source, rather than additional analytic estimates silently inserted into the proof.

Chapter 13

Local CM theta analysis and Schur jets

The preceding chapter (Chapter 12) supplies expanded proofs of the CM arithmetic, local section identities, frame and Frobenius calculations used here. Chapter 16 proves the soft normalized CM reciprocity statement and develops the compatible integral seesaw model. The interface assertions in this chapter are understood with the derivations and remaining hypotheses in both companion chapters and the new exact coefficient arguments of Chapter 17.

The contribution of this section is the local lower bound for valuations of the matrix coefficients of a theta kernel. We isolate it from the later transfer of Hecke operators: it is a result in commutative algebra about differential operators, after a particular CM theta function has been put into normal form. The numerical gain which it produces is the weighted prime-power sum. The subsequent section explains how to use the local estimates for actual lifts and repair the obstruction at primes for which two long rows collide.

13.1 The arithmetic local scales

Write v_p for the valuation normalized by $v_p(p) = 1$, and $|z|_p = p^{-v_p(z)}$. At a specified embedding ι into the algebraically closed p-adic field, put $\mathfrak{o} = \{z : |z|_p \le 1\}$ and $\mathfrak{m} = \{z : |z|_p < 1\}$. Let

$$q_p = \begin{cases} p^2, & \chi_K(p) = -1, \\ p, & \chi_K(p) \neq -1, \end{cases} \qquad \rho_p = \frac{1 + \chi_K(p)^2}{2(q_p - 1)}, \qquad \omega_p = \frac{1 - \chi_K(p)}{2}.$$

The values of ρ_p at split, inert and ramified primes are $1/(p-1)$, $1/(p^2-1)$ and $1/(2(p-1))$, respectively. These are the definitions of `nonsplitWeight` in `Lib/General/NumberTheory/QuadFieldSplitting.lean`. The congruence conditions on d ensure that a nonsplit prime is at least 13 (`Parameters.thirteen_le_of_chiK_ne_one`, in `New/Data/Different.lean`). We call an exponent e live at a nonsplit prime if $e \ge 1$ and either e is odd or $p = d$. Put $\mathfrak{p}_e(\sigma) = [-p^e]\sigma$, using the convention of the arithmetic data for the action on places. The normal-form statement retains also $e = 0$ at $p = d$; that term does not appear in the prime sum. Define

$$A_p(y, \sigma) = \omega_p \sum_{e \text{ live}} \left(y/p^e - w_{\mathrm{nom}}(\mathfrak{p}_e(\sigma))\right)_+, \qquad 0 \le y \le a \tag{13.1}$$

and set $A_p = 0$ at split primes. This is `Data.localAsk` in `New/Curve/LiveSum.lean`. The factor ω_p accounts for the ramified prime as well as inert primes. Since $w_{\mathrm{nom}} \ge a/x_1$, the sum is finite.

Lemma 13.1. *For every place σ and $0 \le y \le a$,*

$$A(y,\sigma) = 2\sum_p A_p(y,\sigma)\log p, \qquad \omega_p \sum_{e \text{ live}} p^{-e} = \frac{1}{p-1} - \rho_p \quad (p \text{ nonsplit}).$$

Proof. Expand the von Mangoldt function by prime powers. The factor $1-\chi_K(p^e)$ is 2 at odd powers of an inert prime and vanishes at even powers; it is 1 at all positive powers of the ramified prime, and vanishes at split primes. For the second assertion sum the odd geometric series $p/(p^2-1)$ in the inert case and one half of the full geometric series $1/(p-1)$ in the ramified case. □

The latter identity is `hasSum_live_inv_pow` in `New/Curve/LiveSum.lean`. Introduce

$$F(p) = \sum_\sigma A_p(w_{\mathrm{nom}}(\sigma),\sigma), \qquad g_p = m\left(\sum_{\tau\in\mathcal{T}} A_p(a,\tau) + kF(p)\right). \tag{13.2}$$

Then

$$2\sum_p g_p \log p = m\sum_{\tau\in\mathcal{T}} A(a,\tau) + mk\sum_\sigma A(w_{\mathrm{nom}}(\sigma),\sigma).$$

The integer m appears because there are m kernels, each with a distinguished derivatives at the tops and approximately $kw_{\mathrm{nom}}(\sigma)$ other derivatives at place σ. We shall derive the positive parts in (13.1).

13.2 The CM period and the differential coordinate

The Weil representation on a standard positive definite hermitian space uses a finite lattice normalized by an idele α; in the source the rational vector w has finite image $\alpha^{-1}w$. Write $\Lambda = \Omega_0\mathcal{O}_K$ for the chosen elliptic curve lattice, $\Omega_0 \neq 0$, and ϑ for its normalized theta section, with $\vartheta(0) = 1$. Invariant derivatives are derivatives in the complex linear covering coordinate, not in an integral formal group coordinate. To read their Taylor series in the p-adic field we transfer the coefficients by ι; we use no analytic continuity of that field map.

Set $s_{\mathrm{ar}} = (2\pi/\sqrt{d})^{1/2}$. The source introduces the factors s_0 (`vacuumRoot`), $s_1 = s_0/u_0$ (`periodFactor`), and

$$\mathrm{per} = s_1 s_{\mathrm{ar}}/\Omega_0.$$

Here u_0 is the nonzero normalizing constant of the Chowla–Selberg identity and s_0 is a chosen f-th root of the reciprocal of the finite splitting character evaluated at α. The precise definitions are in `New/Curve/PeriodDefs.lean`. The positive constant in the period identity is

$$\Omega = \frac{2\pi}{\mathrm{rd}_E}\exp(\kappa_K), \qquad \kappa_K = \gamma - \Re\frac{L'}{L}(1,\chi_K).$$

Proposition 13.2 (Formal-source input: the CM normalizations)**.** *For every automorphism σ of $\mathbb{C}$ over K, $|\sigma(s_0)|^2 = \mathrm{rd}_E/\sqrt{d}$. The factor u_0 is the one prescribed by the normalized period formula `CMLattice.chowla_selberg`. For every $\tau \in \Phi$ and p-adic embedding, the norm of $\iota(s_1)$ is the reciprocal of the norm of the component of α read through the embedding $w \mapsto \iota(\tau(\overline{w}))$. Consequently, if $w \in E \cap \alpha\widehat{\mathcal{O}}_E$, then*

$$|\iota(s_1\overline{\tau(w)})|_p \le 1, \qquad |\iota(s_1)|_p = 1 \quad \text{if } p \nmid N_1.$$

The exact first identity is `Parameters.norm_gal_vacuumRoot` in `New/Curve/PeriodDefs.lean`; the Chowla–Selberg formula is `CMLattice.chowla_selberg` in `Lib/CM/FormalGroup/FormalHeight.lean`. The finite norm identity and its consequences are `norm_iota_periodFactor`, `norm_periodFactor_mul_conj_le_one` and `norm_iota_periodFactor_eq_one` in `New/Curve/TightFrameExists.lean`. The derivation is developed in Chapter 12, retaining the precise normalized Weil-character input.

13.3 Integral and logarithmic theta identities

Fix p and ι; all Taylor series of complex functions in the next paragraphs are read through ι. Write $\phi(z)$ for the integral parameter `translatedWpDeriv`, normalized to start with z, and $\xi(z)$ for `translatedWp`, which begins in degree two. Put $\mathfrak{R}_\phi = \mathfrak{o}[[\phi]] \subset \mathbb{C}_p[[z]]$. This denotes the image of the integral series ring by substitution, not the ring of integral coefficients in the differential coordinate.

Proposition 13.3 (Normalized CM identities (see Chapter 12))**.** *The ring $\mathfrak{R}_\phi$ contains ξ and is stable under d/dz. Moreover $\vartheta = \exp(-\mathscr{L}(z))$ with $\mathscr{L}(0) = \mathscr{L}'(0) = 0$ and*

$$\mathscr{L}''(z) = L_2 + \xi(z), \qquad |\iota(\sqrt{-d})L_2|_p \le 1. \tag{13.3}$$

If (c, D) is in the closure of the pairs $(\iota\delta, \iota(\delta\bar{\delta}))$ for $\delta \in \mathcal{O}_K$, then

$$\phi(cz), \quad \exp(D\mathscr{L}(z) - \mathscr{L}(cz)), \quad (D\mathscr{L}(z) - \mathscr{L}(cz))' \ \in \mathfrak{R}_\phi. \tag{13.4}$$

If $0 \neq \delta \in \mathcal{O}_K$, $\delta y \in \Lambda$ and $D = \delta\bar{\delta}$, the twisted theta translate satisfies

$$(U_y\vartheta)(\iota(\bar{\delta})z) = \vartheta(z)^D V(z), \qquad V \in \mathfrak{R}_\phi. \tag{13.5}$$

These statements are `CMLattice.exists_log_theta` in `Lib/CM/KroneckerLimit/LogTheta.lean`, `taylorVia_translatedWp_mem_phiSeriesRing` and `deriv_mem_phiSeriesRing` in `Lib/CM/LocalHeight/IntegralParameter.lean`, and `rescale_translate_eq_theta_pow_mul` in `Lib/CM/LocalHeight/IsogenyIntegrality.lean`. The logarithmic identity follows from differentiation of the theta addition formula; the normalized integrality assertions use the integral model and integrality of torsion evaluations. The arithmetic proofs and their precise classical premises are developed in Chapter 12.

Lemma 13.4 (Crude integrality of jets)**.** *Let $F = \prod_i (U_{y_i}\vartheta)(\sum_j c_{ij}z_j)$ for $\delta y_i \in \Lambda$, $0 \neq \delta \in \mathcal{O}_K$. If $|c_{ij}/(\iota(\bar{\delta})d)|_p \le 1$, then for every polynomial Q with integral coefficients $(Q(\partial)F)(0)$ is integral.*

Proof. Let $D = \delta\bar{\delta}$. By (13.5), each factor comes from $\vartheta^D V$ with $V \in \mathfrak{R}_\phi$ by substituting $z = \sum_j (c_{ij}/\iota(\bar{\delta}))z_j$. Let $\mathfrak{R}$ be the $\mathfrak{o}$-algebra generated by the images under these substitutions of $\mathfrak{R}_\phi$, $\mathscr{L}'$, and ϑ for each i. All constants are integral. By the chain rule, a partial derivative of a generator is an integral multiplier times the substituted series obtained by applying d times the derivative in z. For $\mathscr{L}'$ the latter is $d(L_2 + \xi)$, in which dL_2 is integral; for ϑ use $\vartheta' = -\mathscr{L}'\vartheta$. The derivative stability of $\mathfrak{R}_\phi$ treats the remaining generators. Thus $\mathfrak{R}$ is stable under all partial derivatives. It contains F and hence all its iterated derivatives. Evaluating at zero and taking integral linear combinations proves the assertion. □

This is `CMLattice.jet_theta_prod_integral` of `New/Curve/FormalParameter.lean`. It applies also at split primes.

13.4 The regularized CM parameter at nonsplit primes

Suppose p is nonsplit. Write $q_p = p^\epsilon$, with $\epsilon = 2$ inert, $\epsilon = 1$ ramified, and $Q_0 = q_p - 1$. Thus $\omega_p = \epsilon/2$, $\rho_p = \omega_p/Q_0$. Choose a primitive Q_0-th root ζ in the p-adic field. The assertion that $(\zeta^j, \zeta^{j(p+1)})$ belongs to the norm-pair closure in Proposition 13.3 is the formal-source input `mem_normPairClosure_of_pow_eq` in `Lib/CM/KroneckerLimit/LogTheta.lean`. Average the parameter:

$$T(z) = Q_0^{-1} \sum_{j=0}^{Q_0-1} \zeta^{-j} \phi(\zeta^j z).$$

By (13.4) it lies in $\mathfrak{R}_\phi$, is tangent to the identity, and has support in degrees $1 \pmod{Q_0}$. It is an integral generator of the ring, since as a series in ϕ it has linear coefficient one and its inverse in that coordinate is integral by recursion. Let Z denote its inverse in the differential coordinate and write $Z(u) = u + z_{q_p} u^{q_p} + \cdots$. Uniqueness of compositional inverses implies $Z(\zeta u) = \zeta Z(u)$. For every $F \in \mathfrak{R}_\phi$ the series $F(Z(u))$ is integral. Differentiation of $T(Z(u)) = u$ therefore yields

$$Z'(u) \in \mathfrak{o}[[u]], \tag{13.6}$$

since $T' \in \mathfrak{R}_\phi$ has constant term one.

Take $c = p$ in the inert case and $c = \sqrt{-d}$ in the ramified case. Its norm-square is q_p, its p-adic valuation is ω_p, and the isogeny has degree q_p. The integral series $[c](u) = T(\iota(c)Z(u))$ is equivariant for the above roots of unity and begins with the nonunit $\iota(c)u$.

Lemma 13.5 (Degree of the reduced isogeny (see Proposition 12.9))**.** *Under these hypotheses $[c](u)$ has a unit coefficient in degree at most q_p.*

The source is `CMLattice.exists_unit_coeff_degree_le` in `Lib/CM/FormalGroup/FormalHeight.lean`, the odd-degree CM isogeny height input for such an integral coordinate. Since all intermediate coefficients vanish, the coefficient in degree q_p is a unit. Comparing through that degree in the composition gives $(\iota(c) - \iota(c)^{q_p}) z_{q_p}$, and hence $|z_{q_p}|_p = p^{\omega_p}$. Choose ϖ with $\varpi^{Q_0} = -z_{q_p}^{-1}$ and put $\pi = \varpi^{Q_0}$; then

$$|\varpi|_p = p^{-\rho_p}, \qquad |\pi|_p = p^{-\omega_p}. \tag{13.7}$$

Lemma 13.6. *There is an integral series R for which*

$$\varpi^{-1} Z(\varpi u) = u - u^{q_p} + \pi R(u).$$

Its compositional inverse S is integral tangent to the identity. In particular for an integral series E_0,

$$T(\varpi z) = \varpi S(z), \qquad S - S^{q_p} = z + \pi E_0(z). \tag{13.8}$$

Proof. The first two nonzero coefficients are as stated by the choice of ϖ. In degree $n = 1 + Q_0 j$, $j \geq 2$, (13.6) gives $v_p(z_n) \geq -v_p(n)$. After scaling the valuation of this coefficient is at least $j\omega_p - v_p(n)$. If $p \nmid n$ this is at least ω_p; otherwise $j \equiv 1 \pmod{p}$ and $j \geq p+1$. Then $n \leq p^\epsilon j$, and $v_p(n) \leq \epsilon + \log_p j \leq \epsilon(j-1)/2$, the last inequality holding for $p \geq 13$, $j \geq p+1$ by comparison at $j = p+1$ and monotonicity of the difference. This proves integrality of the remainder after division by π. Invert the integral series recursively and compose with this inverse, taking $E_0 = -R \circ S$. □

Thus the invariant differential is regularized at scale $p^{-\rho_p}$. The gap between $1/(p-1)$ and ρ_p is the source of the local gain.

13.5 The exceptional exponential factor

Let $c_2 = [z^2]\mathscr{L}$ and put

$$M(u) = (\mathscr{L}(z) - c_2 z^2)\big|_{z=Z(u)}, \qquad n(u) = (\mathscr{L}'(z) - 2c_2 z)\big|_{z=Z(u)}.$$

Then $M = O(u^3)$, $n(0) = 0$, and

$$M' = nZ', \qquad n' = (\xi \circ Z)Z' \in \mathfrak{o}[[u]], \qquad Z' \in \mathfrak{o}[[u]].$$

For a series G, let $G_{[j]}$ denote the part in degrees $j \pmod{Q_0}$. Since Z is equivariant for the Q_0-th roots of unity, this projection commutes with substitution of Z. Define $\kappa = \varpi^{p+1}/p$ and $\mu(u) = \kappa^{-1} M_{[p+1]}(\varpi u)$.

Lemma 13.7. *All coefficients of μ are integral. Modulo $\pi\mathfrak{o}[[u]]$, only its terms of degrees $p+1$ and $p+q_p$ may be nonzero.*

Proof. Since M vanishes through degree two, the possible degrees are $N = p+1+Q_0 I$, $I \geq 0$ (when $\epsilon = 1$ the degree-2 term of the congruence class vanishes). The coefficient of M in degree N is

$$\frac{1}{N} \sum_j n_{p+Q_0(I-j)}(1+Q_0 j) z_{1+Q_0 j},$$

where $j \geq 0$ and the subscript $i = p + Q_0(I-j)$ of n_i is positive. Thus $0 \leq j \leq I+1$ when ramified, and $0 \leq j \leq I$ when inert. By integrality of n', Z' a summand has valuation at least $-v_p(i)$, with an additional ω_p when $j = 1$ because $v_p(q_p z_{q_p}) = \epsilon - \omega_p = \omega_p$. Scaling by $\kappa^{-1}\varpi^N = p\pi^I$ makes its valuation at least

$$1 + \omega_p I - v_p(N) - v_p(i) + \omega_p \mathbf{1}_{j=1}. \tag{13.9}$$

For $I = 0$, $v_p(N) = 0$ and $v_p(i) \leq 1$, so the expression is nonnegative. For $I = 1$ one has $N = p + q_p$ and $v_p(N) = 1$. At $j = 1$, $i = p$, and (13.9) is $2\omega_p - 1 \geq 0$; otherwise $p \nmid i$. If $2 \leq I < p$, $N \not\equiv 0 \pmod{p}$ and the only divisible i is $i = p$ at $j = I$, giving valuation at least ω_p. For $I \geq p$, both positive integers N, i are at most $p^\epsilon(I+2)$. It follows that

$$v_p(N) + v_p(i) \leq 2\epsilon + 2\log_p(I+2) \leq 1 + \omega_p(I-1).$$

The last inequality holds at $I = p \geq 13$ since $\log_p(p+2) < 2$, and persists as I grows because the right hand side increases faster than the left. Thus (13.9) is at least ω_p whenever $I \geq 2$. The non-Archimedean triangle inequality finishes the proof. □

This reconstructs the valuation calculation of `norm_coeff_le_of_deriv` in `New/Curve/FormalParameter.lean`. The estimate works because the coefficients which can cause two divisions by p simultaneously fall into the first two selected degrees; in the second selected degree the exceptional coefficient of Z' is small enough.

Define $\Psi_0(z) = \kappa^{-1}\mathscr{L}_{[p+1]}(\varpi z)$. Using (13.8), the contribution of the non-quadratic part is $\mu(S(z))$. The quadratic term is retained only when $p = d$, in which case its coefficient is $u_0 = \kappa^{-1}\varpi^2 c_2$. From (13.3), $v_p(c_2) \geq -1/2$ (here 2 is a unit); hence $v_p(u_0) \geq 1 - \omega_p - 1/2 = 0$. Let $u_0 = 0$ in the inert case. We obtain

$$\Psi_0(z) = u_0 z^2 + u_1 S(z)^{p+1} + u_2 S(z)^{p+q_p} + \pi R_0(z), \tag{13.10}$$

with integral coefficients and $\Psi_0(0) = 0$. Moreover $|\kappa|_p = p^{\omega_p - 1/(p-1)}$, as is checked in the two cases by substituting ρ_p in $v_p(\kappa) = (p+1)\rho_p - 1$.

The formal source additionally proves that u_1 is a unit at inert primes. It uses the following formal-group input: if every $F \in \mathfrak{R}_\phi$ has $|[z^{p-1}]F'|_p < 1$, then $|[z^{p-1}]\xi|_p \geq 1$. This is `CMLattice.one_le_norm_coeff_translatedWp_nonsplit` of `Lib/CM/FormalGroup/FormalHeight.lean`. The required hypothesis follows from $T(z) = z$ through degree $p < q_p$: an integral series in T has integral coefficient in degree p, so its derivative coefficient in degree $p-1$ is divisible by p. Equation (13.3) now gives $|[z^{p+1}]\mathscr{L}|_p \geq p$; multiplication by $\kappa^{-1}\varpi^{p+1} = p$ makes the corresponding coefficient of Ψ_0 of norm at least one. The coefficient estimate gives the reverse inequality. This proves the unit assertion conditional on that formal-height input.

Proposition 13.8 (Local normal form for torsion translates)**.** *If $p \nmid n$ and $ny \in \Lambda$, there are a unit N and a constant κ_n in the p-adic field satisfying $|\kappa_n|_p = p^{\omega_p - 1/(p-1)}$ and an integral series V_y such that*

$$(U_y\vartheta)(\varpi z) = \big(\exp(\kappa_n \Psi_0(z))V_y(z)\big)\big|_{z \mapsto Nz}.$$

The same Ψ_0, N, κ_n work for all y with the given n.

Proof. By (13.4) for the norm pairs of the powers of ζ, the series $\exp(\zeta^{j(p+1)}\mathscr{L}(z) - \mathscr{L}(\zeta^j z))$ is integral in T. Replace z by $\zeta^{-j}z$ (which preserves the ring) and multiply these series over all j. Character orthogonality makes their product $\exp(Q_0(\mathscr{L}_{[p+1]} - \mathscr{L}))$. Apply (13.5) with $\delta = nQ_0$ to get

$$(U_y\vartheta)(nQ_0 z) = \exp(-(nQ_0)^2\mathscr{L}(z))V(z), \qquad V \in \mathfrak{R}_\phi.$$

Replace $\mathscr{L}$ by $\mathscr{L}_{[p+1]}$ in the exponential: the residual product just obtained, raised to the integer power n^2Q_0, is integral in T. Scaling by ϖ then gives integral coefficients for this product and V, since $T(\varpi z) = \varpi S(z)$. Finally take $N = (nQ_0)^{-1}$, $\kappa_n = -(nQ_0)^2\kappa$. This proves the required normal form. □

Equation (13.8), (13.10) and the proposition above give the nonsplit part of `CMLattice.exists_formal_parameter` in `New/Curve/FormalParameter.lean`.

13.6 Tight frames and theta vectors

The theta kernel on a space of matrices is not an unrelated product of one-variable theta functions. The product is built from a tight frame in the CM lattice.

Definition 13.9. A tight frame of length M is a finite collection of vectors $w_i \in E \cap \alpha\widehat{\mathcal{O}}_E$ with nonnegative integer multiplicities m_i such that

$$\sum_i m_i \sigma(w_i)\overline{\tau(w_i)} = M\delta_{\sigma\tau} \quad (\sigma, \tau \in \Phi).$$

After repetition of elements we may assume each multiplicity is one.

This is `IsTightFrame` in `Lib/Weil/Construction/ConjugateJetDefs.lean`. For instance the N_1 powers of ζ_{N_1} form a tight frame of length N_1: their integrality follows from the different normalization of α, and the frame equality is the geometric sum of $(\sigma(\zeta_{N_1})/\tau(\zeta_{N_1}))^j$. This is `Parameters.isTightFrame_zeta_pow`. To ensure good p-adic integrality of all theta vectors one needs lengths that are prime to p, and for the lower bounds also frames having some vector attaining all the finite norm bounds of Proposition 13.2.

Proposition 13.10 (Large frames with a unit vector (see Proposition 12.14))**.** *For every $R \geq 0$, every integer length $M \geq (dN_1)^6(R+1)$, and each p-adic embedding ι, there is a tight frame with multiplicities $m_i \geq R$ containing a vector w_j with $|\iota(s_1\overline{\tau(w_j)})|_p = 1$ for all τ.*

The precise arithmetic existence result is `Parameters.exists_isTightFrame`, in `New/Curve/TightFrameExists.lean`. It constructs additional frames using the cyclotomic different generator and solves for the desired length using the coprimality of two frame lengths. The general arithmetic construction is proved in Proposition 12.14 of the preceding chapter. For upper bounds take $M = p(dN_1)^6 + 1$, large enough and prime to p.

For a standard positive definite hermitian space V of rank h, denote by $\mathscr{F}_V$ the finite part of its Weil representation with Heisenberg operator ϱ and theta functional ϑ_V. The Taylor model of $F \in \mathscr{F}_V$, at differential scale c, is the series $\mathcal{T}_c F$ with coefficients

$$[z^\beta]\mathcal{T}_c F = \frac{c^{|\beta|}}{\beta!}\vartheta_V(v_\beta \otimes F) \qquad (\beta \in \mathbb{N}^{\Phi\times\{1,\dots,h\}}), \tag{13.11}$$

where v_β is the Fock vector of the monomial and $\beta! = \prod_x \beta_x!$. This is `Space.jetSeries` in `Lib/Weil/Construction/DataProduct.lean`. Evaluating a monomial differential operator introduces $\beta!$ once again. No factorial may be lost by confusing the Taylor coefficient with the invariant derivative.

Let $\delta \in \mathcal{O}_K$ satisfy $\delta\bar{\delta} = D_\delta > 0$, and assume $v \in \Lambda$, with the curve's area $|v|^2 M\sqrt{d}/(2D_\delta)$. For a frame define linear maps on the complex coordinates of V by

$$A_{ic}(z) = \frac{v}{\bar{\delta}s_{\mathrm{ar}}} \sum_\tau \overline{\tau(w_i)} z_{\tau,c}.$$

The sum of pullbacks along A_{ic} of the normalized curve's hermitian form, counted with multiplicity m_i, equals the standard form: expand the scalar products of $A_{ic}(z)$ and $A_{ic}(z')$ and use Definition 13.9. The factor $|v/\bar{\delta}|^2 M$ is exactly absorbed by the area normalization. In particular the product of the pulled-back theta sections has the appropriate transformation law.

Proposition 13.11 (Formal-source input: theta product and finite vector)**.** *Under these frame and area choices, a product of the pullbacks of degree-m_i theta monomial sections is the entire theta function of some vector $F \in \mathscr{F}_V$. For rational Heisenberg translates this theta function is the product of the translated curve sections with arguments prescribed by the same linear forms. All finite point evaluations of F are integral at all p-adic embeddings. There is a finite vacuum expansion*

$$\sum_{j\in I} M_\delta \ell(\varrho(\alpha x_j)\,\mathrm{vac})\varrho(-\alpha x_j)F = D_\delta^{2fh}\ell(F)\,\mathrm{vac}$$

for every complex linear functional ℓ, where M_δ is a natural number, $|I| = D_\delta^{2fh}$, and δx_j is integral at the finite places.

This is `Parameters.exists_finitePart_of_isTightFrame`, in `New/Curve/FrameSections.lean`; its vacuum expansion is the structure `Space.VacuumExpansion` of `Lib/Weil/Construction/ConjugateJetDefs.lean`. The finite vector reconstruction and its integrality are formal-source inputs here. The identity of the corresponding Taylor series follows by evaluating along every line and differentiating at zero (homogeneous polynomials are determined by their values on all lines). The corresponding lemmas are `Space.taylor_line_eq_subst_jetSeries` and `Space.jetSeries_eq_prod`. The source also proves `Space.eq_zero_of_jetSeries_eq_zero` by the identity theorem for entire functions.

13.7 Frobenius pairings in tight-frame products

Fix a nonsplit p. Consider a frame product of type `Parameters.thetaProducts` with all multiplicities repeated, frame length M prime to p, torsion denominators n prime to p, and

multiplier b' of p-adic norm at most one (not the scale b). Its factors for pairs (i,c) are

$$(U_{y_{ic}}\vartheta)\left(\sum_{\tau}\iota(b'\overline{\tau(w_i)})z_{\tau,c}\right), \qquad ny_{ic}\in\Lambda.$$

At first there is no extra denominator from K. By Proposition 13.8, after substituting ϖz the product has the form

$$F(\varpi z)=\exp(\kappa_n\Psi(z))V(z), \quad V\in\mathfrak{o}[[z]], \quad \Psi(0)=0, \tag{13.12}$$

where $\Psi=\sum_{i,c}\Psi_0(L_{ic})$, $L_{ic}=\sum_\tau r_{\tau i}z_{\tau,c}$, and $r_{\tau i}=N\iota(b'\overline{\tau(w_i)})$. At a nonsplit prime, $p\nmid N_1$, so Proposition 13.2 gives $r_{\tau i}\in\mathfrak{o}$.

Lemma 13.12 (Frobenius orthogonality (see Proposition 12.16))**.** *In characteristic p, modulo $\pi\mathfrak{o}$, the frame matrix satisfies, for any live exponent e and for $e=0$ when $p=d$,*

$$\sum_i r_{\sigma i}^{p^e}r_{\tau i}=(N\iota(b'))^{p^e+1}M\,\mathbf{1}_{\tau=\mathfrak{p}_e(\sigma)}. \tag{13.13}$$

Here the multiplier is read in the same reduction, with N regarded as the p-adic element from the normal form.

The formula is `Parameters.exists_frame_rows_frob` in `New/Curve/ThetaProducts.lean`; more precisely it holds under every map to characteristic p killing all elements of valuation at least ω_p. The proof uses the Frobenius permutation of the cyclotomic embeddings and the frame identity. At an inert prime the odd Frobenius interchanges the CM embeddings, explaining the minus sign in $[-p^e]$; at the ramified prime complex conjugation acts the same way modulo the local uniformizer. These arithmetic identifications of embeddings modulo the indicated ideals are used as input.

For a finite truncation degree $D>0$ let $I_D=\{0\le e\le\lfloor\log_p D\rfloor : e \text{ odd or } p=d\}$. For integral series $Z_{\tau,c}$ with zero constant term in any finite variable set, suppose the pairings

$$\sum_{\tau,c}Z_{\tau,c}^{p^e}Z_{\mathfrak{p}_e(\tau),c}=0 \quad (e\in I_D) \tag{13.14}$$

hold after reduction modulo π and terms of degree $>D$. We shall prove they kill the exceptional exponent in (13.12) and render the exponential integral through degree D.

Lemma 13.13 (Artin–Schreier cancellation)**.** *Let R be a commutative ring of characteristic p, and let finitely many elements t_i, L_i satisfy $t_i-t_i^{p^\epsilon}=L_i$. Assume each t_i is nilpotent, and $\sum_i L_i^{p^j+1}=0$ for all odd j if $\epsilon=2$, or all $j\ge 0$ if $\epsilon=1$. Then $\sum_i t_i^{p+1}=\sum_i t_i^{p+p^\epsilon}=0$.*

Proof. Iterate the given equation:

$$t_i=\sum_{a=0}^{A-1}L_i^{p^{\epsilon a}}+t_i^{p^{\epsilon A}}.$$

Nilpotence kills the last term for large A. Expanding $\sum_i t_i^p t_i$ by Frobenius gives terms $\sum_i L_i^{p^u}L_i^{p^v}$ for $u=\epsilon a+1$, $v=\epsilon b$. Each equals $(\sum_i L_i^{p^{|u-v|}+1})^{p^{\min(u,v)}}=0$; indeed the exponent difference is odd if $\epsilon=2$. Expand $\sum_i t_i^p t_i^{p^\epsilon}$ in the same way, taking $v=\epsilon(b+1)$. This proves both identities without requiring that R be reduced or perfect. □

This is `sum_pow_eq_zero_of_null` in `New/Curve/ThetaProducts.lean`.

Proposition 13.14 (Null-cone integrality)**.** *Whenever the integral zero-constant series $Z_{\tau,c}$ satisfy (13.14), the coefficients of $F(\varpi Z)$ through degree D are integral.*

Proof. Work in the quotient of the ring of integral series by the ideal of coefficients divisible by π together with terms of degree $> D$. For $t_{ic} = S(L_{ic}(Z))$, (13.8) gives $t_{ic} - t_{ic}^{p^e} = L_{ic}(Z)$, and t_{ic} is nilpotent by degree. Using (13.13) we have

$$\sum_{i,c} L_{ic}(Z)^{p^e+1} = (N\iota(b'))^{p^e+1} M \sum_{\tau,c} Z_{\tau,c}^{p^e} Z_{\mathfrak{p}_e(\tau),c} = 0$$

for $e \in I_D$. Larger e give zero automatically by degree. Lemma 13.13 shows that the two S-powers in (13.10) vanish after summing; at $p = d$ the quadratic terms vanish by the same computation with $e = 0$. It follows that the coefficients of $\Psi(Z)$ through degree D are divisible by π. The exponential in (13.12) is integral through this degree because

$$v_p((\kappa_n \pi)^j / j!) = j/(p-1) - v_p(j!) \geq 0.$$

There is no formal convergence problem, since the substituted series have zero constant term and we work in a finite truncation. Multiplication by $V(Z)$ preserves integrality. □

This is the first conclusion of `thetaProducts_normal_form` in `New/Curve/ThetaProducts.lean`, whose hypothesis is the predicate `IsIntegralOnNullCone`. In the source the couplings are required to vanish under every characteristic-p map annihilating π; it suffices in the proof to apply that assumption to the indicated quotient.

In the inert case, when b' has norm one, one also knows which coefficients of the exponent are units. Reducing in the residue-field series ring and differentiating (13.10), $u_0 = 0$ and the term with u_2 has zero derivative. Since $S - S^{p^2} = z$, $S' = 1$ in this reduction, and

$$\frac{\partial \overline{\Psi}}{\partial z_{\tau,c}} = \bar{u}_1 \sum_i r_{\tau i}\, t_{ic}^{p}, \qquad t_{ic} = \sum_{a \geq 0} L_{ic}^{p^{2a}}.$$

By the frame identity a coefficient is nonzero exactly at $z_{\sigma,c}^{p^e}$ with odd e and $\mathfrak{p}_e(\sigma) = \tau$. It is a unit because M, Nb', u_1 are units. No coefficient supported on several variables occurs. This proves `coeff_sum_frob_ne_zero_iff` and explains the sharpness assertion of `thetaProducts_normal_form`. The same pairings which reduce the guaranteed upper divisibility are genuinely present in the series. This fact is also used in the lower volume estimates.

13.8 Schur matrices and determinant losses

Let R_1, R_2 be the source and target ranks, $N = R_2$, and let $\kappa_\sigma(h)$ count columns of height $h < N$ in the prescribed tableau. A filling assigns h row indices to each column of height h; its polynomial $b_i(Z)$ is the product of minors in the indicated rows and first h columns. Put $D = \sum_\sigma \sum_{h<N} h\kappa_\sigma(h)$. The jet estimate must hold for every Schur matrix coefficient, not just for the highest weight vector.

Proposition 13.15 (Formal-source input: integral Schur expansion)**.** *For a weight satisfying the rank condition `FitsRanks` the Schur matrix $S_\kappa(X)$ satisfies*

$$b_{i'}(XY) = \sum_i S_{i'i}(X) b_i(Y) \prod_\sigma \det(Y_\sigma)^{d_\sigma}, \tag{13.15}$$

where the determinants use the first R_1 columns of the variable matrix Y_σ, and $d_\sigma = \kappa_\sigma(R_1)$ if $R_1 < N$, zero otherwise. This holds over every ring and commutes with base change; the coefficient coordinate maps in the standard tableau basis are integral on the associated Borel semi-invariants.

This is the standard integral tableau theorem as used by `PolyRep.sum_schur_smul_nablaBasis` and `schur_map` in `Lib/GLn/SchurModule/SchurMatrix.lean`. Its basis and weight are `nablaBasis` and `diagramWeight` in `Lib/General/Algebra/ZerosCostDegree.lean`; the integral basis theorem is in `Lib/GLn/SchurModule/MinorBasis.lean`. Write $J_Q(F) = Q(\partial)F(0)$ and $\mathscr{J}_\kappa(F)_{i'i} = J_{S_{i'i}}(F)$ (`schurJet` in `Lib/Weil/Construction/DataProduct.lean`).

Let ev be a finite set of places whose height-one couplings will be exempted. Define

$$a_\sigma^{ev} = \kappa_\sigma(1)\mathbf{1}_{\sigma\notin ev}, \quad c_\sigma = \sum_{\substack{h<N\\ h\neq 1}} \kappa_\sigma(h), \quad D_\sigma = \sum_{\substack{h<N\\ h\neq 1}} h\kappa_\sigma(h).$$

For the ordered coupling $\sigma, \tau = \mathfrak{p}_e(\sigma)$ put

$$\mathfrak{C}_{p,e}(\sigma,\tau) = \min(a_\sigma^{ev}/p^e, c_\tau + a_\tau^{ev}) + c_\sigma/p^e + \min(D_\sigma/p^e, D_\tau). \tag{13.16}$$

This is `couplingCount` in `New/Curve/LiveSum.lean`. Its three terms distinguish the three types of pairs of columns.

Lemma 13.16 (Simultaneous tilting)**.** *For any integral matrix arrays $Y_{\sigma,c,j}$ with source rows c and column indices $j \geq 0$, there are integral $h \times h$ matrices $M_{\sigma,h}$, $1 \leq h < N$, as follows. Write $Y_\sigma^{(h)}$ for the transposed array of the first h columns. For each $e \in I_D$, $\tau = \mathfrak{p}_e(\sigma)$, and rows u, u' from $M_{\sigma,h}Y_\sigma^{(h)}$, $M_{\tau,h'}Y_\tau^{(h')}$,*

$$\sum_c u_c^{p^e} u'_c \equiv 0 \pmod{\pi}$$

unless $h = 1$ and either $\sigma \in ev$ or $(h' = 1, \tau \in ev)$. Moreover

$$\prod_{\sigma,h} |\det M_{\sigma,h}|_p^{\kappa_\sigma(h)} \geq p^{-\omega_p \sum_{e\in I_D}\sum_\sigma \mathfrak{C}_{p,e}(\sigma,\mathfrak{p}_e(\sigma))}. \tag{13.17}$$

Proof. An elementary observation about valuation rings: if coefficient vectors in $\mathfrak{o}^h$ must satisfy finitely many conditions $L_i(v) \in u_i\mathfrak{o}$ with integral linear functionals L_i and nonzero $u_i \in \mathfrak{o}$, there is a matrix of h admissible rows of determinant valuation at most $\sum_i v_p(u_i)$. Inductively a unimodular change on the current coefficient basis makes the next linear functional depend on just one coordinate: choose among its coefficients one of largest norm and use divisibility to eliminate the others. Multiplying that coordinate by u_i suffices; earlier conditions are maintained by linearity. Requiring all h coordinates to be divisible by u_i instead costs $hv_p(u_i)$.

Fix e, σ, τ as in the statement. For source height one, the untransformed row y is fixed. If $\sigma \notin ev$, either scale it by an integral p^e-th root of π, of valuation ω_p/p^e, or impose the single congruence $\sum_c y_c^{p^e} u'_c \equiv 0 \pmod{\pi}$ on each target height except the exempt height-one row. The weighted costs are $\omega_p a_\sigma^{ev}/p^e$ and $\omega_p(c_\tau + a_\tau^{ev})$; choose the cheaper. If $\sigma \in ev$ there are no conditions of this type.

For source height greater than one and target height one, choose integral p^e-th roots of the entries of the untransformed target row y' and require

$$\sum_c (y'_c)^{1/p^e} u_c \in \pi^{1/p^e}\mathfrak{o}.$$

Taking p^e-th powers in characteristic p gives the required congruence modulo π. Further transformation of y' is scalar multiplication and preserves this relation. The weighted cost is $\omega_p c_\sigma/p^e$. Finally, for both heights greater than one, annihilate all pairings by scaling all coefficient rows on either the source or the target side, at the smaller cost $\omega_p \min(D_\sigma/p^e, D_\tau)$. All the conditions are on the original coefficient coordinates or the fixed linear forms specified above. Impose them simultaneously by the first observation; integral row operations preserve previously zero pairings, since the quotient has characteristic p. The costs add to the exponent of (13.17). □

This is the argument of `exists_null_coupling_frames` in `New/Curve/JetDivisibilit y.lean`. The source implements the valuation-ring observation with `IndexDvd.of_det_dvd`. In particular the matrices are not claimed unimodular; their determinant costs must be subtracted.

For each box with height h and row label a prescribe a nonnegative integer charge $g(\sigma, h, a)$. Require

$$g(\sigma, 1, a) > 0 \quad \text{or} \quad g(\mathfrak{p}_e(\sigma), h, a) > 0$$

whenever $\sigma \in ev$ or $(h = 1, \mathfrak{p}_e(\sigma) \in ev)$. For a tableau i' write $G_{i'}$ for the total charges of its boxes, with multiplicity. The charges will kill the exempt pairings.

Theorem 13.17 (Schur-jet bound)**.** *Suppose F satisfies the null-cone property of Proposition 13.14 for degree $D > 0$ and the scales (13.7). Then*

$$\begin{aligned} v_p(\mathscr{J}_\kappa(F)_{i'i}) \geq (1/(p-1) - \rho_p)D - \omega_p \sum_{e \in I_D} \sum_{\sigma} \mathfrak{C}_{p,e}(\sigma, \mathfrak{p}_e(\sigma)) \\ - fN^3(\lfloor \log_p D \rfloor + 1) - G_{i'}. \end{aligned} \tag{13.18}$$

Proof. Apply the jet functional in X to (13.15). The integral basis property converts any coefficient bound on the polynomial $J_{b_{i'}(XY)}(F)$ in Y to a bound on the entries of the Schur jet. The largest coefficient norm of a polynomial over an algebraically closed valued field is attained at some integral specialization: divide by a largest coefficient, reduce modulo the maximal ideal, and choose a point of the infinite residue field where the resulting nonzero polynomial does not vanish. Thus it suffices to bound the expression for all specialized integral Y.

Apply Lemma 13.16. Substitute in the theta series fresh variables $u_{\sigma,h,j,a}$, $0 \leq j < h < N$, $1 \leq a \leq N$, via

$$X_{\sigma,a,c} = \varpi \sum_{h,j} p^{g(\sigma,h,a)} (M_{\sigma,h} Y_\sigma^{(h)})_{j,c} u_{\sigma,h,j,a}.$$

Remove the scaling factor ϖ from the substitution. Expanding the Frobenius pairings of matrix coordinates (a, c) in characteristic p gives exactly the pairings of tilted coefficient rows. Unexempted terms vanish, and exempt terms vanish by their positive charges since $p \in \pi\mathfrak{o}$. The null-cone condition gives integral Taylor coefficients of the substituted theta series through degree D.

Now apply the product of specified minors in the fresh variables as a differential operator. It is homogeneous of degree D, with integral coefficients. There are at most $fN \sum_{h<N} h \leq fN^3$ fresh variables. A monomial u^β of degree D contributes a factor $\beta!$; the base-p digit formula gives

$$v_p(\beta!) = (D - \sum_i s_p(\beta_i))/(p-1) \geq D/(p-1) - fN^3(\lfloor \log_p D \rfloor + 1).$$

The same bound holds for the sum of the monomial pairings. The chain rule identifies that sum with

$$\varpi^{D}p^{G_{i'}}\left(\prod_{\sigma,h}\det M_{\sigma,h}^{\kappa_\sigma(h)}\right)J_{b_{i'}(XY)}(F).$$

Within each minor the substituted differential rows are transformed by the same $M_{\sigma,h}$, yielding its determinant, and each box contributes ϖ and the power of p prescribed by its charge. Removing these factors subtracts $\rho_p D$, $G_{i'}$, and at most the determinant loss (13.17). This proves (13.18). □

This is `norm_schurJet_le` in `New/Curve/JetDivisibility.lean`. The factorial gain is diminished by the coordinate and determinant costs, and the explicit logarithmic factorial correction. The minima in (13.16) are essential: replacing them by the more expensive alternatives would eliminate the positive-part terms in (13.1).

13.9 Application to the prescribed weights

For the arithmetic data $\mathcal{D}$, let $R_j = k + (j+1)(t+1)$, $0 \le j < m$, be the target ranks of the definite chain. Put $w_\sigma = w_{\mathrm{nom}}(\sigma)$,

$$a_\sigma = a\mathbf{1}_{\sigma\in\mathcal{T}},\quad c_{j,\sigma} = \sum_{\substack{h<R_j\\ h\neq 1}}\kappa_\sigma(h),\quad D_{j,\sigma} = \sum_{\substack{h<R_j\\ h\neq 1}}h\kappa_\sigma(h),\quad D_j = \sum_\sigma(a_\sigma + D_{j,\sigma}).$$

The weight multiplicities $\kappa_\sigma(h)$ are `Data.`κ in `New/Curve/LiveSum.lean`: height one occurs at tops with multiplicity a; height k has multiplicity $w_1(\sigma) - \lfloor (k+t+1)/2\rfloor$; heights above k congruent to k modulo $t+1$ have multiplicity $2h_2 - (t+1)$; all other multiplicities vanish. These expressions are nonnegative by admissibility.

Lemma 13.18 (Weight counts)**.** *For $0 \le j < m$ one has*

$$kw_\sigma \le D_{j,\sigma} \le kw_\sigma + x^{60},\qquad 0 \le c_{j,\sigma} \le w_\sigma + x^{60}.$$

Proof. The multiplicity of height k is at least w_σ because the rounded weight w_1 includes the gap $3h_2$. The difference from w_σ is at most the rounding step plus $3h_2$. There are at most $j+1$ higher heights, each bounded by R_j and of multiplicity at most $2h_2$. Thus the excess box count is at most $k(N_1 + 3h_2) + 2(j+1)R_j h_2$ (here N_1 in the rounding term denotes the cyclotomic order, not a target rank). With $k \ll x^5$, $m \ll x^{1/2}$, $R_j \ll x^{11/2}$, $h_2 = r^8 \ll x^{40}$ and the cyclotomic bound, this is smaller than x^{60} on the admissible asymptotic range. The column count is bounded by ignoring the heights. The formal source establishes these comparisons for every admissible datum as `Data.lowColumns_lowBoxes_bounds` in `New/Columns/JetExponentBound.lean`. □

For a nonsplit prime $p < x_1$ define the exponent

$$E_{p,j}(ev) = \left(\frac{1}{p-1} - \rho_p\right)D_j - \omega_p\sum_{e\in I_{D_j}}\sum_\sigma \mathfrak{C}_{p,e}(\sigma,\mathfrak{p}_e(\sigma)) - fR_j^3(\lfloor\log_p D_j\rfloor + 1). \tag{13.19}$$

This is `Data.jetExponent` in `New/Columns/JetExponentBound.lean`. Define

$$\mathrm{coll}_p(ev) = \sum_\sigma\sum_{e\ \mathrm{live}}(a/p^e)\mathbf{1}_{\sigma,\mathfrak{p}_e(\sigma)\in\mathcal{T}\setminus ev}.$$

The sum converges by geometric decay but need not have finite support. It records the loss when a long row has a long partner.

Proposition 13.19 (Prime-power gain of one jet)**.** *For nonsplit $p < x_1$ and $0 \le j < m$,*

$$E_{p,j}(ev) \ge \frac{g_p}{m} - \mathrm{coll}_p(ev) + \omega_p |\mathcal{T} \cap ev|\, a/x_1 - \frac{fa(\log b)^2}{pm}. \tag{13.20}$$

Proof. We give the comparison on the asymptotic admissible range, sufficient for the main theorem; the formal source proves it uniformly for all admissible data. Set $u = a/(pm)$. The separated scales imply $x^{60}, R_j^3 \le u$, $c_{j,\sigma}/p \le u$, and $5(\lfloor \log_p D_j \rfloor + 1) \le (\log b)^2$. For example $u \ge xb/(x_1 m)$ and $x_1 \le b^{10^{-6}}$ control all polynomial terms. The column bound and $m/x \to 0$ give $c_{j,\sigma}/p \le (a/x + x^{60})/p \le u$. The last inequality follows from $D_j \ll b^2$ for sufficiently large x using $p \ge 2$.

Fix a live exponent $e \in I_{D_j}$, write $P = p^e$, and let $\tau = \mathfrak{p}_e(\sigma)$. The elementary factorial contribution less the cost for that coupling is

$$\begin{aligned} &\frac{a_\sigma + D_{j,\sigma}}{P} - \mathfrak{C}_{p,e}(\sigma, \tau) \\ &= \frac{a_\sigma}{P} - \min(a_\sigma^{ev}/P, c_{j,\tau} + a_\tau^{ev}) \\ &\qquad + \left(\frac{D_{j,\sigma}}{P} - D_{j,\tau}\right)_+ - \frac{c_{j,\sigma}}{P}. \end{aligned}$$

The ordinary-box term is at least $k(w_\sigma/P - w_\tau)_+ - x^{60}$, by Lemma 13.18. The last term costs at most u. For the distinguished height-one term, if σ is not a top then the contribution and requested gain are zero. If both σ, τ are nonexempt tops, the loss relative to $(a/P - w_\tau)_+$ is at most a/P, the collision summand. If σ is a nonexempt top but τ is not a nonexempt top, then $a_\tau^{ev} = 0$ and $c_{j,\tau} \le w_\tau + x^{60}$, giving at least $(a/P - w_\tau)_+ - x^{60}$. If σ is an exempt top, the minimum is zero and a/P exceeds the requested positive part by $\min(a/P, w_\tau)$; at $e = 1$ this is at least a/x_1 since $p < x_1$ and $w_\tau \ge a/x_1$. These four cases give the pointwise desired bound with at most $3u$ of error per place and exponent, plus the exempt contribution for $e = 1$.

For live e above the truncation cutoff there is no coupling cost; the factorial contribution then bounds both requested positive parts. The extra contribution at $e = 1$ does occur within the truncation since $D_j \ge a > x_1$. In summing the factorial contribution over all live exponents use Lemma 13.1. The $e = 0$ term allowed at $p = d$ occurs only in the coupling cost, and costs at most $4u$ per place. Indeed $p = d$ is fixed and $c_{j,\sigma} \le u$ from $w_\sigma \le a/x$ and $m/x \to 0$. The place $[-1]\sigma$ cannot have the same class of tops as σ, nor can both be thick. Hence the first minimum is bounded by u, and the last by $ka/x_1 + x^{60} \le u$, since at least one term corresponds to the small weight. Summing, the comparison error is at most $4fu(\lfloor \log_p D_j \rfloor + 1)$, and the factorial error at most $fu(\lfloor \log_p D_j \rfloor + 1)$. Multiply the comparison by $\omega_p \le 1$ and use the last scale inequality above. The sum of the desired positive parts with factor ω_p is exactly g_p/m by (13.2). □

The exact uniform source declaration is `Data.le_jetExponent`, in `New/Columns/JetExponentBound.lean`. Formula (13.20) is the numerical interface to the chain. When $ev = \varnothing$ there are no charges, but long-row collisions can be expensive. The Hecke repair uses exempt sets which remove all such collisions; the extra positive term in (13.20) compensates part of the charge incurred in doing so.

13.10 From theta products to the standard integral kernel data

A bound for arbitrary formal products becomes a bound for the automorphic theta kernel only after identifying the normalized finite integral structures. We give this interface explicitly. For a standard space V let vac_V be the standard finite vacuum. Fix a nonzero factor s_{vac} chosen for rank one and define the scaled vacuum of rank h to be $\mathrm{vac}_V^{\circ} = s_{\mathrm{vac}}^{h}\,\mathrm{vac}_V$. Its point evaluations are the products over the h coordinates of the rank-one evaluations. The integral module $\mathrm{Std}_\delta(V,\iota)$ is the span over $\iota^{-1}(\mathfrak{o})$ of $\varrho(\alpha x)\,\mathrm{vac}_V^{\circ}$ with δx integral at all finite places above p (x is a finite adelic vector). This definition is `Parameters.standardData` in `New/Curve/IntegralVacuum.lean`; it depends on $\delta \in \mathcal{O}_K$ and on the p-adic embedding.

Proposition 13.20 (Formal-source input: integral vacuum and conjugates)**.** *The scalar s_{vac} is nonzero, and for all p and all p-adic embeddings every finite point evaluation of vac_V° is integral. For $\sigma \in \mathrm{Aut}(\mathbb{C}/K)$ its conjugates are bounded by $\exp(CN_1^2 h)$ for a constant C depending only on the fixed K, N_0. Moreover there is a finite vector generating the finite part by Heisenberg translates and with theta functional equal to 1; after applying σ, its jets at scale* per *coincide, up to permutation of the places, with jets of translates of another such finite vector at a scale of squared absolute value $1/\Omega$.*

The precise uniform generator statement is `Parameters.exists_isJetConjugate_generator` in `New/Curve/ConjugateGenerator.lean`; integrality and the conjugate bound for the scaled vacuum are `norm_intVacuum_le_one` and `norm_intVacuum_le` in `New/Curve/IntegralVacuum.lean`. These inputs use the normalized CM reciprocity law. For higher rank the vacuum value bounds follow multiplicatively from rank one, as indicated by `intVacuum_finVec_eq_prod`.

The integral module is stable under the Heisenberg translations by the indicated lattice because the central factors in their multiplication law are roots of unity and hence units at p. It is also stable under the finite Weil action of an element integral above p. This follows from the finite vacuum-translation sum for the Weil action on a vacuum: it is a sum of translates by integral vectors with unit multipliers, and conjugation by the integral element preserves the translation lattice. This bounded-action assertion is used in its exact form `Parameters.omega_mem_standardData` and `finHeisOp_mem_standardData`, in `New/Curve/ConjugatesOfJets.lean`. We assume the finite Weil model's vacuum sum identity in this application.

Lemma 13.21 (Transfer of a jet bound)**.** *Let $\mathcal{L}$ be a complex linear functional on the Taylor series space (in particular any specified Schur jet entry), $c \in \mathbb{C}$ a scale multiplier, and Q a nonnegative real bound. Suppose the norm of $\iota(\mathcal{L}(F))$ is at most Q for every F whose image under ι is a frame theta product with the torsion condition $n\delta y_i \in \Lambda$ and with scalar norm $|\iota(cs_1)|_p$. Then for every $f \in \mathrm{Std}_\delta(V,\iota)$,*

$$|\iota(\mathcal{L}(\mathcal{T}_{c\ \mathrm{per}} f))|_p \le Q.$$

Proof. Take a frame with $M = p(dN_1)^6 + 1$, of length prime to p, using Proposition 13.10. Repeat its vectors so that all multiplicities are one. Use Proposition 13.11 with $\delta = M$ and the corresponding repeated frame of length M^2, choosing the monomial sections equal to the normalized theta section. Then $v = \Omega_0$ and $D_\delta = M^2$ satisfy the required area identity. The resulting finite vector F_0 has theta value one, since every normalized theta section has constant value one. In the vacuum expansion take ℓ to be the theta functional and solve for the vacuum. After scaling by s_{vac}^{h}, the coefficients are integral: they are the normalized

vacuum evaluations times the natural multiplier divided by M^{4fh}, a p-adic unit. The phase factors from translating these expansions have absolute value one.

Consequently a lattice translate of the normalized vacuum is an integral linear combination of translates of F_0. Approximate each finite translate by a rational one, using density of rational points of the finite additive adeles and local constancy of the Heisenberg orbit on the finite model. The rational approximation can be made so that δ times the translation vector is integral above p. Denominators at other primes are cleared by a rational integer prime to p. Applying the frame product identity gives the class of theta products in the hypothesis: the scalar factor is the product of cs_1 and factors whose norm is one from M; the torsion coordinates have denominators dividing $n\delta$ with $p \nmid n$. This exact rational approximation and denominator assertion is the one used in `Parameters.norm_jetSeries_le` in `New/Curve/ConjugatesOfJets.lean`. Linearity of $\mathcal{L}$ and the non-Archimedean inequality then bound every generator of the integral module, hence its entire span. □

This argument uses the stated finite-vector and period inputs rather than replacing the actual kernel data by a single formal theta product. For $\delta = 1$ at a nonsplit prime, Proposition 13.14, Theorem 13.17, and Lemma 13.21 bound the Schur jets on the entire module whenever the scalar multiplier has integral norm. For general $\delta \neq 0$, Lemma 13.4 applied using Proposition 13.2 gives the elementary integrality at scale $\overline{\delta}d$ per; this is precisely `Parameters.norm_jet_jetSeries_le_one` in `New/Curve/ConjugatesOfJets.lean`.

Finally, a finite theta kernel acts not just on one fixed Weyl coefficient vector but on vectors at all adelic arguments. The Weyl gauge at g of a vector ξ is the maximum absolute value of its coordinates after reading $g^{-1}\xi$ through the specified p-adic embeddings (where the action contains the determinant character as well as the Schur representation). This is `weylGauge` of `Lib/GLn/SchurModule/WeightRepBasic.lean`. An integral change of adelic frames above p has integral matrices in both directions under this representation, so the gauge is unchanged. For the standard data f the kernel is equivariant for these integral changes.

Proposition 13.22 (Formal-source input: gauge bound for a translated theta kernel)**.** *If the pair of adelic changes of the matrix variable factors into integral changes and a diagonal element with parameters $u_\sigma(a)$, then a uniform bound on every jet coefficient after weighting by*

$$\prod_{\sigma}\left(\prod_{a}|u_\sigma(a)|_p\right)^{e_\sigma}\prod_{\text{boxes }(\sigma,h,a)\text{ of }i'}|u_\sigma(a)|_p$$

is a bound on the factor by which the theta kernel raises the Weyl gauge. Here e_σ is the determinant exponent of the output weight. In particular the kernel with standard datum and scale c per, with no diagonal change, raises gauges by at most

$$\left(\frac{|\iota(c)|_p}{|\iota(\overline{\delta}d)|_p}\right)^{D},$$

where D is the number of boxes of the output tableau.

The exact interface is `Parameters.mem_raises_of_jet_bound` and `mem_raises_of_standardData` in `New/Curve/ConjugatesOfJets.lean`. Its finite-place factorization and equivariance are used as inputs. The matrix estimate itself is just the ultrametric operator bound, and the weights come from evaluating the diagonal matrix in the Schur representation and its determinant character. The final displayed bound follows by homogeneity and the crude integrality at scale $\overline{\delta}d$ per proved above. More importantly, at a non-split prime away from the conductor and level, the sharper Schur bound replaces that crude estimate by (13.18), with any extra weights charged box by box. This completes the local differential calculation needed for the Hecke and seesaw argument.

Chapter 14

The definite chain, collisions and Hecke repair

The purely algebraic local Hecke estimates, including the Frobenius root and coefficient arguments, are developed in Chapter 15. The actual rational orbit and finite integral support calculations for the geometric chain are developed further in Chapter 16. Reading them separately is useful: the former concern a finite coefficient lattice, whereas the seesaw identifies the lift with a construction on an actual adelic lattice. Chapter 17 develops the definite normalized coefficient and integral branching in detail.

We continue the construction of the preceding section. Throughout this section the prime d of K is fixed. Write $g = g_0 g_1$ for the product of the two level primes, and let $\pi_g \in \mathcal{O}_K$ have norm g. The letter π without a subscript denotes the element of norm N used to define the extra column level. The spaces W and C are, respectively, the good part of the row space and the entire initial column space, of dimensions n and c. We use $n_j = k + j(t+1)$ for the ranks of the definite Hermitian spaces X_j, $0 \le j \le m$; thus n_j is not a dimension of a space of automorphic forms. Set $w(\tau) = w_{\mathrm{nom}}(\tau)$. The local ask A_p, the prime sum A, the gain g_p, and the direction cost $\mathcal{U}$ are as in the preceding section; in particular

$$g_p = m \left(\sum_{\tau \in \mathcal{T}} A_p(a, \tau) + k \sum_{\tau \in \Phi} A_p(w(\tau), \tau) \right). \tag{14.1}$$

Here we prove the quantitative implications between the jet theorem and Theorem A. We also isolate the remaining nontrivial automorphic and local representation-theoretic assertions on which the argument depends. They are identified explicitly as formal-source inputs when we do not give an independent proof of their full strength.

14.1 Coefficient lattices, class spaces and index

It is important to formulate the lattice estimates without assuming that the coefficient field of the automorphic forms is a fixed number field. For a rational prime p, fix a field isomorphism $\iota : \mathbb{C} \simeq \overline{\mathbb{Q}}_p$ as in the coefficient construction, and transport the valuation ring and absolute value to $\mathbb{C}$. We denote this ring by O_ι and normalize $|p|_p = p^{-1}$. All assertions below are quantified over ι. For a Hermitian space X and adelic point g', let $\|v\|_{\iota,g'}$ denote the Weyl gauge of its coefficient vector in the integral structure at g'. Thus, if ρ_κ is the algebraic coefficient representation, the gauge is obtained by applying the inverse of the adelic coefficient matrix to v and then taking the coefficient supremum norm transported by ι. It is non-Archimedean; multiplication by a scalar z multiplies the gauge by $|\iota(z)|_p$. At the ordinary

complex absolute value the Weyl norm is instead the chosen Hermitian norm; the two kinds of estimate should not be confused.

For level K_j and weight κ of the given definite space X_j, define $M_j(\iota)$ to be the O_ι-module of forms of this type, every value of which has gauge at most 1. Write $M = M_0(\iota)$, a submodule of C. All finite-place estimates in the proof are made on this adelic integral module, not just on abstract polynomials in a single weight representation. The exact definitions are `Data.integralColumns` and `integralFormsAt` in `New/Exchange/CrudeModularData.lean` and the background files respectively. The notation M in this section always depends on the chosen embedding.

We recall the index convention used in the proof.

Definition 14.1. Let P, Q be submodules over a valuation domain O, and let $z \in O$. Write $I(P, Q; z)$ if for every finite family $(v_i)_{i=1}^h$ of elements of P there exists a square h by h matrix D over O such that $\det D \mid z$ in O and

$$\sum_j D_{ij} v_j \in Q \quad (1 \le i \le h).$$

For an O-linear endomorphism T preserving P, write $T \in \mathrm{Type}(p, P, s; e)$ if there is an integer $l \ge 0$ with $l \le e$ and

$$I(P, TP + p^s P; p^l).$$

The parameter s is a nonnegative integer and e is allowed to be a nonnegative real number.

These are `IndexDvd` and `HasType` in `New/Core/ModuleIndex.lean`. The divisibility of the determinant is in the stated direction: $\det D \mid p^l$ implies $|\det D|_p \ge p^{-l}$. It does not mean that $\det D$ is a multiple of p^l. Nor is it necessary to assign a finite cardinality to a lattice quotient. This definition is appropriate even when O is not discretely valued, and is sufficient for the later height argument.

Lemma 14.2 (Elementary index calculus)**.** *Let $I(P, Q; z)$ and $I(Q, R; a)$ hold. Then $I(P, R; za)$ holds. If u is an O-linear map with $u(P') \subset P$ and $u^{-1}(Q) \cap P' \subset Q'$, then $I(P', Q'; z)$ holds. If $u(Q) \subset Q'$ and $P' \subset u(P) + Q'$, then $I(P', Q'; z)$ also holds. Consequently for $T_1, \dots, T_h$ preserving P,*

$$T_i \in \mathrm{Type}(p, P, s; e_i)\ (1 \le i \le h) \Longrightarrow T_1 \cdots T_h \in \mathrm{Type}(p, P, s; \textstyle\sum_i e_i).$$

Finally, if $u_i : P \to V'$ are linear and $I(P, u_i^{-1}(Q'); z)$ holds for each i, then

$$I\left(Q' + \sum_{i=1}^{h} u_i(P), Q'; z^h\right).$$

Proof. The first assertion is obtained by applying the index witness matrix for $I(Q, R; a)$ to the family already obtained using the matrix for $I(P, Q; z)$, and taking their product. The determinant is multiplicative. For pullback, apply the index condition to $(u(v_i))_i$ and use linearity. For pushforward, write $v_i' = u(v_i) + q_i'$ with $q_i' \in Q'$ and use the same linear combinations of these expressions.

For the product assertion it suffices to consider two factors. Suppose we have index witnesses for $T_i P + p^s P$, with determinant bounds p^{l_i}. Apply the second operator to the index condition for the first, if necessary reversing the order in this statement. Since $T_i(p^s P) \subset p^s P$, the pushforward rule gives the required condition from $T_1 P + p^s P$ to $T_1 T_2 P + p^s P$, and the transitivity rule gives the sum $l_1 + l_2$. Now use induction. For the final assertion enlarge Q' successively by the images. In each step, push the index condition along u_i and use transitivity. □

For comparison, if $P = O^D$ is free over a discretely valued ring and T has elementary divisors $p^{\nu_1}, \dots, p^{\nu_D}$ (with $\nu_i = +\infty$ allowed), the determinant valuation of the canonical lattice $TP + p^s P$ is

$$\sum_i \min(\nu_i, s).$$

Thus a bound on the type measures a truncated sum of elementary divisors. The Hermite and Smith factorization arguments in `ModuleIndex.lean`, in particular `IndexDvd.of_det_dvd`, implement this relation for the valuation domains of the proof. We will use type bounds, not an assertion about each individual eigenvalue of a Hecke operator.

14.2 The groups and the chain of theta kernels

At stage j the source is the standard definite space X_j of rank n_j. The column level K_j is the intersection of the chosen principal level at the places dividing $\overline{\pi_g}$ with the subgroup whose last row is congruent to $(0, \dots, 0, 1)$ modulo π. The element π is chosen so that $\pi\bar{\pi} = N$. There are f places dividing π, one member of each conjugate pair over the completely split prime N. In particular K_j contains the principal level of $\overline{\pi_g}\pi$ and is compact open. The spaces of forms are finite sums of coefficient spaces indexed by rational adelic classes. The chosen principal level makes the automorphism groups of the classes trivial. These claims are `Data.columnLevel`, `Data.principalLevel_le_columnLevel`, `Data.card_piPlaces`, and `Data.classAut_columnLevel_eq_bot` in `New/Columns/ColumnDefs.lean`. The analogous triviality at the intermediate stages follows from `Data.classAut_eq_bot_of_le_chosenLevel`. We use this specific property as a formal-source input for the adelic lattice model; no estimate for a weighted stabilizer sum can replace it in the following argument.

Here is how the coefficient weight is structured. With $h = 2h_2$ as in the admissible data, the number of columns of height j' in the Young diagram at τ is

$$\kappa_\tau(j') = \begin{cases} a, & j' = 1,\ \tau \in \mathcal{T}, \\ 0, & j' = 1,\ \tau \notin \mathcal{T}, \\ w_1(\tau) - \lfloor (k+t+1)/2 \rfloor, & j' = k, \\ h - (t+1), & j' > k, \quad t+1 \mid j'-k, \\ 0, & \text{otherwise.} \end{cases}$$

Only columns of height below n_j contribute to the polynomial part at stage j. The determinant exponent of the representation is 0 at stage zero, and $e_j(\tau) = \lfloor n_{j-1}/2 \rfloor - w_j(\tau)$ for $j \geq 1$. The coefficient representation on X_0 is in particular

$$\bigotimes_{\tau \in \mathcal{T}} \mathrm{Sym}^a \mathbb{C}^k.$$

All factors at the other places have degree zero. The restriction on the places of height-one columns is central in the collision argument: once a p-adic Frobenius orbit meets both top places, the two factors of degree a can interact at precisely the prime-power levels that contribute to the required gain.

Let θ be an admissible twist of finite conductor of rational norm q_θ, and let μ_{j+1} be the prescribed splitting character of the $j+1$st lift, with twist θ^{j+1}. The theta kernel K_j is the one for $\mathrm{Hom}(X_j, X_{j+1})$ with weight κ, character determined by $\mu_{j+1}\theta^{j+1}$, and complex Taylor scale $\overline{\pi_g}$per. The identities between weights that make the lift available use the fact that there is no nonzero column height strictly between n_j and n_{j+1}. The declarations are `Data.i`

`sLiftWeight_kappa` and `Data.liftKernel` in `New/Exchange/CrudeModularData.lean`; the formulas for κ, e_j are in `New/Curve/LiveSum.lean`.

For a tuple of finite adelic matrices $\xi = (\xi_j)_{0\le j<m}$ in the corresponding Hom spaces, let $\phi_{\xi,j}$ be the finite Heisenberg translate by ξ_j of the normalized integral vacuum of the preceding section. Let

$$\mathsf{C}^{j}_{\theta,\xi} = \mathsf{K}_{j-1}(\phi_{\xi,j-1}) \circ \cdots \circ \mathsf{K}_0(\phi_{\xi,0}), \qquad \mathsf{C}^{0}_{\theta,\xi} = 1$$

be the composite definite lift to stage j. Individual terms need not be invariant under the intermediate level. A useful datum is instead a finite linear combination of tuples $\Psi = \sum_\xi c_\xi[\xi]$, which is required to be jointly invariant under the levels in the sense that the corresponding tensor of kernel data is fixed by their simultaneous actions.

At the final stage X_m has rank one greater than Y_m. The map to the last row space is not simply evaluation on $U(Y_m)$: it uses branching of the algebraic coefficient representation. Let D_{br} be the integral branching matrix and put

$$e_{\mathrm{br}}(\tau) = \sum_{j'<n_m} (\kappa_\tau(j') - \kappa'_{\mathrm{kept}}(j')) + e_m(\tau).$$

Here the summands count the columns at that height whose height is reduced in branching; the formal definition of κ'_{kept} is $h_2 - \lfloor (t+1)/2 \rfloor$ if $j' \ge k$ and $t+1 \mid j'-k$, and zero otherwise. Thus the linear restriction is

$$(\mathsf{R}v)(u) = \left(\prod_{\tau\in\Phi} \tau(\bar\pi)^{-e_{\mathrm{br}}(\tau)} \right) D_{\mathrm{br}} v(\mathrm{diag}(u,1)). \tag{14.2}$$

The relevant branching identity is `Data.restrictToY`, and the estimate `Data.le_branchExp` is

$$e_{\mathrm{br}}(\tau) \ge \kappa_\tau(1) - 2b.$$

For a prime $p \ne N$, the scalar in (14.2) is a p-adic unit; using the lattice-compatible branching identity the restriction cannot increase the gauge. At a complex conjugate σ the estimate is stronger:

$$\|\sigma\mathsf{R}v(u)\|^2 \le \exp\left(x^{70} - \textstyle\sum_\tau e_{\mathrm{br}}(\tau)\log N\right) \|\sigma v(\mathrm{diag}(u,1))\|^2. \tag{14.3}$$

Indeed under each conjugate $|\tau(\bar\pi)|^2 = N$, so the scalar gives the displayed negative exponent; the branching matrix contributes at most $e^{x^{70}}$ by the coefficient estimate. The precise lattice and norm compatibility statements `Data.GaugeLe.restrictToY` and `Data.weylNormSq_restrictToY_le`, in `New/Seesaw/LiftCharacters.lean`, are formal-source inputs. Their significance is that the archimedean gain of level N is large, while the prime N itself will be treated separately.

Write

$$\mathsf{B}^{\Psi}_{\theta} = \mathsf{R} \sum_{\xi} c_\xi \mathsf{C}^{m}_{\theta,\xi}.$$

This is `Data.Bμ` in the source. For the local calculations the datum must have the following exact properties (the record `Data.IsGoodDatum`, in `New/Columns/ChainValuation.lean`):

1. If ξ lies in the support, its components multiplied by π_g are integral away from the places of $q_\theta N$.
2. The coefficients c_ξ are integral at every ι with $p \nmid q_\theta$. For $q_\theta = 1$ the sum of the absolute values of their conjugates is at most $e^{x^{60}}$.
3. The total tensor is jointly invariant under the intermediate column levels and the final action specified in the source.

4. At N the chain has an additional paired description. Project each kernel datum to the places dividing π; sum the pairwise translations of adjacent data by last-column $-y/N$ and last-row $-\bar{y}/N$ for y modulo N. The resulting combination of paired lifts, with the same coefficients c_ξ, equals the original chain of m lifts. In the source $m = 2(\varrho + 1)$.

The fourth property, `chain_at_N`, will make it possible to move large apparent denominators at N onto a single column operator, instead of losing them at every Taylor coefficient.

Lemma 14.3 (Gauge bounds for a jointly invariant chain)**.** *Suppose $p \nmid q_\theta$, Ψ is good and each translated kernel after a chosen modification obeys*

$$\|\mathsf{K}_j(\phi_{\xi,j})v\|_{\iota,u} \le p^{r_j}\|v\|_{\iota,g'}$$

at the prescribed coefficient frames, uniformly for each pair of points and every ξ in the support. Assume also that the modifying operators commute with the required source and target level actions on the total invariant tensor. Starting with an integral column, the total modified chain has gauge at most $p^{\sum_{j<m} r_j}$ in the final frame.

Proof. For jointly invariant tensors all the integrals over definite groups reduce to finite class sums; an individual term is a composition, indexed by a tuple of representatives of the successive class spaces. The triviality of the stabilizers means that such a sum is a plain sum of the corresponding values, not a sum with rational denominators from the stabilizer orders. Each composite has gauge at most the product of the individual bounds. The coefficients c_ξ are integral since p is prime to q_θ. Thus the non-Archimedean triangle inequality leaves the largest gauge bound unchanged under summation. This proves the result. The source formalizes the reduction to the class sum and this argument as `Data.IsGoodDatum.gaugeLe_chain` via `Data.exists_chain_eq_sum` in `New/Columns/ChainValuation.lean`. □

The same argument at an archimedean place bounds the square root of the squared norm by the product of individual operator norm estimates, the sum of absolute values of the coefficients, and the product of the numbers of integral classes at each stage. This explains why the class-counting estimate and coefficient-size estimate must be checked even when no such factor is present in the non-Archimedean estimate.

14.3 The geometric seesaw and its denominator control

Let P be the (unprojected) Poincaré construction from columns to initial row forms. It comes from a Bessel period for the unitary space Y_0; the successive indefinite theta lifts run through the spaces $Y_0, \dots, Y_m$ of ranks $r + jt$. Our notation Θ_θ is reserved for the composite on W (or its extension to the unprojected principal-level row space before selecting W). Initially we do not identify it with any definite chain.

Theorem 14.4 (Seesaw identity, formal-source input)**.** *There is a nonzero $z_0 \in \mathbb{C}$, independent of the twist, and for every admissible twist θ a good datum Ψ_θ such that*

$$\Theta_\theta P(z_0 u) = \mathsf{B}_\theta^{\Psi_\theta} u \qquad (u \in C^{\mathrm{ram}}).$$

The declaration is `Data.seesaw` in `New/Seesaw/SeesawTheorem.lean`. The exact content includes the four properties of good data above, not just an equality of two complex-valued functions. In the following we explain the denominator mechanism and give the finite cut argument in a form that is independent of the deeper theta calculations.

In the adelic construction a stage j has a rational Bessel subgroup, with finite-adelic component denoted here by S_j, and a compact open subgroup $J_{j,\theta}$ of the row group. The function arising after the first j unfoldings is P_j, with P_0 the generic function from which the original Poincaré series is formed. The rule $P_{j+1} = \mathsf{U}_j(P_j)$ is a Bessel unfolding by a single step of theta lifting, at parameters (ξ_j, y_j) of the archimedean and finite row spaces.

Three deeper assertions of the source are needed in this construction; we do not reprove the full geometric or local theta identities.

1. The local constant-term identity preserves the generic character, and consecutive pairs of unfolding steps give matrix coefficients of a definite paired chain. At the even step $2i + 2$, the equality is

$$P_{2i+2}(\iota_X v \cdot (x, y_{2i+2})) = \mathrm{algCoeff}_{t_i}\,(\mathsf{C}^{\mathrm{pair}}_{i+1}(u))(\mathrm{emb}_X v)$$

   for all ramified columns u and all configurations y with initial point y_0 in the chosen level. The linear functional t_i depends on the archimedean parameters but not on the twist or on u. These are `Data.isGeneric_unfoldStage` and `Data.unfoldStage_even` in `New/Seesaw/UnfoldingStages.lean`; the local arrow-pair tensor and integral identity used in their proof are in `New/Seesaw/EdgePair.lean` and `New/Seesaw/DoubleLiftValue.lean`. In the formal proof one unfolds the two consecutive theta kernels, takes their constant terms with respect to the unipotent subgroup, and applies the identity for integrating a product of kernels. This gives $i + 1$ paired lifts from the assertion for i.

2. The support of each stage function is controlled by the lattice lines of its Hermitian space. If P_j is nonzero, the lines at places of the twist move at most j steps, and the integral lines outside those places do not move (`Data.hasNearLines_of_unfoldStage_ne_zero`, same file). In the induction, even steps are covered by the line estimate for the theta kernel. At odd steps the possible exceptional local lines at the level prime are eliminated by writing a ramified column as $u' - z_w u'$ for a central generator z_w at a place over N. The two terms of the matrix coefficient are then equal there, so cancel. This use of ramification is why the seesaw statement is on C^{ram}.

3. For each stage one can choose a compact open A_j independent of the twist and a finite cut $T_{j,\theta} \subset G_{j,f}/J_{j,\theta}$ stable under $S_j \cap A_j$ and contained in the level outside the twist conductor. Here $G_{j,f} = U(Y_j)(\mathbb{A}_f)$. The cut covers the support after translation by S_j and has cover multiplicities which divide powers of q_θ. At $j = 0$ it is contained in the chosen level. This assertion is `Data.exists_stage_cut` in `New/Seesaw/SeesawTheorem.lean`.

Lemma 14.5 (Finite-cut multiplicities)**.** *Let G be a locally compact totally disconnected group, S a closed subgroup, and $J \subset A$ compact open subgroups of G. Let $D \subset G$ be compact, and suppose that the set to be covered is in SD. Suppose for each $x \in D$ there is an e such that*

$$(xux^{-1})^{q^e} \in A \qquad (u \in J).$$

There is a finite $(S \cap A)$-invariant subset $T \subset G/J$ whose S-saturation covers the image of SD. For v in this saturation, write

$$m(v) = \#\{s(S \cap A) : s \in S,\ s^{-1}v \in T\}.$$

Then $m(v)$ divides a power of q.

Proof. The image of D in the discrete space G/J is finite. Choose one element $[x]$ in that image from each S-orbit, and take T to be the union of their $(S \cap A)$-orbits. These are finite

since $S \cap A$ is compact. If H_x is the stabilizer of $[x]$ in S, then the s with $s^{-1}[x] \in T$ constitute $H_x(S \cap A)$. Thus

$$m([x]) = [H_x : H_x \cap A].$$

This index is finite, and is invariant under translating $[x]$ by S. Pass from H_x to its quotient by the normal core of $H_x \cap A$; this is a finite group. As $H_x \subset xJx^{-1}$, every element in the finite quotient has order dividing a power of q. Indeed for each $h \in H_x$ and each conjugate by an element of H_x, that conjugate's q^e-power belongs to A, hence h^{q^e} belongs to the normal core. By Cauchy's theorem every prime divisor of the order of this finite group divides q. The index divides its order, proving the assertion. This is the finite-index argument in `exists_cut_coverMultiplicity_dvd`, `New/Seesaw/IntegralAsCosetSum.lean`. □

The cut hypothesis is justified in the source by the support of the unfolded stages. Outside the twist the local groups are integral. At the finitely many places of the twist, taking a sufficiently large power of the residue characteristic puts an element of the conjugated compact open level back into the fixed level. The admissible translations therefore lie in finitely many local compact sets modulo S_j; the stage-cut theorem applies the lemma to those sets. Thus the importance of the local geometric support bound is not just convergence: it is the prescribed prime support of denominators.

At the analytic level the unfolding reduces an integral over the rational Bessel quotient to a sum over a finite cut. In the notation of the lemma the formula has the form

$$\int_{B(F)\backslash G(\mathbb{A}_F)} P_j(g)z(g)\,dg = \lambda_j \sum_{v \in T_{j,\theta}} m(v)^{-1} b_j(v).$$

The scalar λ_j is determined by the fixed archimedean integral and Haar normalizations; it is independent of the twist. To see the finite sum mechanism, the weight $\mathbf{1}_T(v)/m(v)$ is a partition of unity after summing over $S_j/(S_j \cap A_j)$ on the support. Indeed exactly $m(v)$ terms contribute, with the same orbit multiplicity. It may therefore be inserted in the integral without changing it. Integration over a coset of the compact subgroup $J_{j,\theta}$ contributes its Haar mass; the standard level-index normalizations in the period over B cancel this factor. The remaining archimedean integral is λ_j. The analytic hypotheses are precisely the generic character identity, automorphy of z, integrability of $P_j z$ and the product form of its Bessel-group integral. The formal-source input for this quotient-integral statement, including its measure normalizations, is `Grading.exists_integral_eq_sum_over_cut` in `New/Seesaw/IntegralAsCosetSum.lean`.

The product of the resulting finite combinations along the chain is again a finite combination: its coefficients are sums of products of $1/m(v)$. For $p \nmid q_\theta$ these are integral. If the twist is trivial, $q_\theta = 1$, the multiplicities are all 1 wherever the cut is nonempty. For quantitative complex sizes one counts the cut at each stage by the index of the chosen principal level; this gives a power of `Data.cutBound` independent of the actual columns. The formal assertion `Data.exists_isAdmissibleComb_unfolding` in `New/Seesaw/SeesawTheorem.lean` carries out this induction, testing against every suitable form z and linear functional on the tensor space. The terminal stage identifies the coefficient with paired lifts, using the preceding even-step identity.

To turn the resulting tensors into the vacua of the definite chain one needs a further finite Weil representation calculation. For the evaluated kernel tensor at each configuration y which is integral outside q_θ, the source proves that it is an admissible linear combination of reduced vacua, up to a single constant C and with denominator $\mathrm{cutBound}^{2m}$, bounded size when the twist is trivial. The reduced vacua are the paired combinations of Heisenberg translates described above. The exact declarations are `Data.exists_isAdmissibleComb_reducedVacu`

`um`, `Data.isAdmissibleComb_reducedVacuum`, and `Data.exists_isGoodDatum_of_invariant` in `New/Seesaw/GoodDatumExists.lean`. We take the three assertions as formal-source inputs; they contain the explicit evaluation of finite Weil data and the local fiber geometry, rather than just the group-theoretic finite-cut calculation. The scalar C and the archimedean normalization from the integrals are absorbed into the one scalar z_0. The theorem then follows by pairing against the test vectors at stage m; equality of all those matrix coefficients gives equality of the lifts. These are the mechanisms underlying Theorem 14.4.

14.4 Ramification and short generators on the definite group

The set of ramified columns has an elementary description which is compatible with its adelic lattice. For each $v \mid \pi$ choose a central element of the local norm-one torus, integral everywhere, supported on $v, \bar{v}$, whose component at v reduces to a generator of $\mathbb{F}_N^\times$. Let z_v be its translation operator on columns. Define

$$R_N = \prod_{v|\pi}(1 - z_v), \qquad C^{\mathrm{ram}} = R_N C.$$

These operators commute. In particular every element of the image can be written $(1 - z_v)u'$ at each $v \mid \pi$. The definition and the lift of the generators are in `Data.ramifyingOp`, `Data.ramifiedColumns` and `Data.exists_centralGen` in `New/Columns/ChainValuation.lean`.

The following dimension and norm observations will be useful both here and for the spanning theorem.

Lemma 14.6. *For each $v \mid \pi$,*

$$\dim \ker(1 - z_v) \le \frac{c}{N-1}.$$

Consequently for any subspace $U \subset C$ and any subset T of the places dividing π, with $R_T = \prod_{v\in T}(1 - z_v)$,

$$\dim R_T^{-1}(U) \le \dim U + |T|\frac{c}{N-1}.$$

If u is integral at every embedding and every value of its conjugates has squared norm at most C_0, $R_T u$ is likewise integral, with squared norm at most $4^{|T|}C_0$.

Proof. The forms fixed by z_v have level enlarged by the cyclic group generated by the central translation. The index of the old level in this enlargement is at least $N - 1$: a positive power lying in the old level reduces to the identity at v. Since both levels are contained in the chosen level and their class automorphism groups are trivial, their spaces have dimension proportional to the numbers of adelic classes. The ratio of these class counts is the level index; thus the dimension of the fixed space is at most $c/(N - 1)$. This is the class-count formula specialized in `Data.finrank_ker_one_sub_centralGen_le`. For a linear operator L, $\dim L^{-1}(U) = \dim \ker L + \dim(U \cap \operatorname{im} L)$. Applying this successively proves the second assertion. The central element preserves the integral gauge and permutes the arguments of an algebraic form. Therefore $u - z_v u$ is still integral by the ultrametric inequality, and its squared complex norm is at most $4C_0$ by the ordinary triangle inequality. Induction gives the bound. This is `Data.ramifyingOp_finrank_comap_and_short`. □

To produce the initial short vectors the source uses a geometry-of-numbers estimate for definite Hermitian lattices. We record its role explicitly. Let I be a level whose class stabilizers are trivial, let the coefficient representation of rank u have a Young diagram with D_κ boxes

(counted over the infinite places), let $Q \geq 1$ be a rational integer in the different ideal of E, and suppose $M_0 \geq uf + D_\kappa$. Every class admits an algebraic basis of its associated lattice with the uniform bounds summarized in `exists_short_basis_stdSpace`. Put $R_0 = 6Q(fu)^2$. Then the coefficient forms which are everywhere integral with squared norms of all conjugates bounded by $R_0^{6M_0}$ span the whole class space (`span_isShort_eq_top`, in `New/Columns/ChainValuation.lean`). This is a formal-source input for the quantitative short basis bound; the coefficient argument is as follows. The class space is freely the direct sum of weight modules at the class representatives, so it suffices to span one such weight module. Apply the Schur representation to the bounded lattice basis. In the standard coefficient model the Weyl vectors obtained by evaluating integral flag polynomials at roots of unity span the weight module: if a linear functional annihilates all such vectors, successive application of the root bound for a one-variable polynomial shows that the corresponding polynomial vanishes identically. Multiplication by the determinant factor for the determinant weight also preserves integrality in the chosen lattice basis. Every conjugate of a root of unity has absolute value 1, so the norm bound follows from the power of R_0 for the bounded basis and the degree of the coefficient representation. This is why one obtains a spanning set rather than simply a single short vector. Applying this construction at level K_0 with the admissible scale estimates, and then applying R_N, gives the needed conclusion:

$$\operatorname{span}_{\mathbb{C}}\{u \in C^{\mathrm{ram}} : u \text{ is everywhere integral, } \|\sigma u(g')\|^2 \leq e^B \ \forall \sigma, g'\} = C^{\mathrm{ram}}. \tag{14.4}$$

Specifically $|\{v \mid \pi\}| = f$, and the bound before application of R_N is chosen at most $e^B/4^f$. The exact scale inequality and conclusion are `Data.span_short_ramifiedColumns`. We will use (14.4) with the ordinary complex norm squared, not its logarithm, bounded by e^B.

14.5 The collision loss and the marked set

For a rational prime p, put

$$\omega_p = \frac{1 - \chi_K(p)}{2}, \qquad s_p = \lceil \log_p m \rceil, \qquad \delta_p = s_p f(n_m + s_p).$$

A prime is called short if $p \equiv 1$ or $-1 \pmod m$. In this terminology shortness refers to a congruence and not to the size of p. Define

$$\ell_p = \frac{fa(\log b)^2}{p} \begin{cases} m, & p \text{ short}, \\ 1, & \text{otherwise}. \end{cases}$$

We will call p small nonsplit if $p < x_1$ and $\chi_K(p) \neq 1$. At such a prime, g_p in (14.1) is the desired total valuation. In the formulas below, all relevant place actions are the actions of the CM cyclotomic type, not of the single quadratic extension $K/\mathbb{Q}$ alone.

For a subset $E_0 \subset \Phi$ of places declared even, set

$$C_p(E_0) = \sum_{\tau \in \Phi} \sum_{\substack{e \geq 1,\ e \text{ odd or } p=d \\ \tau, -p^e\tau \in \mathcal{T} \setminus E_0}} \frac{a}{p^e}.$$

This collision count removes all pairs for which one of the two places belongs to E_0. Let o_0 be the order of the class of $-p$ modulo N_0 (using the place convention of the data). We call p served if it is inert in K, not short, and o_0 is odd with $o_0 \leq s_p$. These are the definitions `Data.IsShortPrime`, `Data.loss`, `Data.ceilLogM`, `Data.deltaP`, `Data.IsServed` and `Data.collisionSum` in `New/Curve/LiveSum.lean`.

Lemma 14.7 (Collisions without exceptional treatment)**.** *For p small nonsplit and not served,*

$$mC_p(\varnothing) \le \ell_p.$$

Moreover, for every small nonsplit p one has $C_p(\varnothing) \le 2fa/p$.

Proof. If all the exponents appearing in a collision sum are at least o, then

$$C_p(E_0) \le fa \sum_{e \ge o} p^{-e} \le 2fa/p^o,$$

since $p \ge 2$. The second assertion follows by taking $o = 1$. The first assertion for short p follows as soon as $(\log b)^2 \ge 2$. If p is not short, recall that a collision has $p \ne d$ and odd e divisible by o_0 (`Data.IsCollision.ne_and_odd_and_dvd`, `New/Columns/JetExponentBound.lean`). If there are any collisions for such a prime, o_0 is odd; the assumption that p is not served therefore gives $o_0 > s_p$. Consequently

$$mC_p(\varnothing) \le 2mfa/p^{s_p+1} \le 2fa/p \le \ell_p.$$

The source's assertion is `Data.mul_collisionSum_le_loss`. The arithmetic fact excluding collisions at d uses the congruence hypotheses on d; without those hypotheses the argument at a ramified prime would require modification. □

The issue at a served prime is different: exponents $e \le s_p$ can contribute, and multiplying a loss of size a/p by m cannot be absorbed in ℓ_p. The prime-by-prime repair uses a set of split places of E/E^+ even though p is inert in $K/\mathbb{Q}$. Fix ι and denote by $l(\tau)$ the prime of E defined by the embedding attached to τ and ι.

At a served prime, write $u = -p$ in the group of infinite places. Its order is divisible by m:

$$m \mid \mathrm{ord}(u). \tag{14.5}$$

For clarity, the basic 2-power estimate behind this assertion is the following. If an odd integer v is congruent to neither 1 nor -1 modulo 2^h, then the order of its image modulo 2^s has 2-part at least 2^{s-h+1} provided s is sufficiently large. For an even exponent L the identity

$$v_2(v^L - 1) = v_2(v-1) + v_2(v+1) + v_2(L) - 1$$

follows by repeated factorization of $v^{2j}-1$. Since one of $v-1, v+1$ has valuation 1 and the other less than h, the required bound follows from divisibility by 2^s. Here $m = 2^h$, the cyclotomic exponent is $s = 22h$, and the shortness exclusion allows the estimate with ample slack. The local embeddings of the cyclotomic tower, including primes dividing the fixed conductor, are handled in `Data.IsServed.m_dvd_orderOf` in `New/Columns/JetExponentBound.lean`. We use that exact assertion for the admissible data.

The other relevant arithmetic property is

$$l(u^e\tau) = \overline{l(\tau)} \quad (e \text{ odd}), \qquad l(\tau') \in \{l(\tau), \overline{l(\tau)}\} \iff \tau' = u^e\tau \text{ for some } e. \tag{14.6}$$

These are `Data.inducedPlace_neg_pow_mul_of_odd` and `Data.inducedPlace_eq_or_star_iff`, same file. In particular, a served prime of the data is unramified in the cyclotomic tower. One way to see the parity assertion is to express the decomposition group through Frobenius in $\mathrm{Gal}(E/\mathbb{Q})$: its restriction to K is conjugation and its restriction to the cyclotomic part is the pth power. The element $-p$ of the CM place group is obtained by composing this Frobenius with complex conjugation. Thus it interchanges the two induced primes above the real place. The order condition (14.5) ensures that these two primes are distinct. The full arithmetic assertions `smul_eq_iff_inert`, `smul_eq_star_iff_inert`, `q_eq_inert` and the unramified/splitting hypotheses used for these formulas are formal-source inputs for the specific extension E.

Lemma 14.8 (Marking)**.** *Fix a small served prime and a p-adic embedding. There is a finite set Λ of primes above p such that:*

1. *no $l \in \Lambda$ has $\bar{l} \in \Lambda$;*
2. *each marked prime l is induced by a top;*
3. *for every top τ, either $l(\tau)$ or $\overline{l(\tau)}$ belongs to Λ.*

Write $E_0(\Lambda) = \{\tau : l(\tau) \in \Lambda\}$ for the set of even places. Then

$$|\Lambda| \le f, \qquad C_p(E_0(\Lambda)) = 0.$$

Every τ with $l(\tau)$ or $\overline{l(\tau)}$ marked has $\tau \notin \mathcal{S}$.

Proof. For each top, multiplication by u^{o_0} stays within the tops and, because o_0 is odd, interchanges the two induced primes by (14.6). Hence the primes induced by tops occur in conjugate pairs. Choose one from each pair. This gives all three properties and the cardinality bound. Every collision has e odd, so the primes induced by its two top places are conjugate. At least one is marked, and consequently no collision remains outside $E_0(\Lambda)$.

For the last assertion, the orbit condition (14.6) gives a power of u taking the place τ to a top. Its class modulo N_0 is therefore in the cyclic subgroup generated by the class of u. That subgroup has odd order. A thick place has class -1, of order two in the residue group of the fixed modulus N_0; it cannot lie in an odd-order subgroup. The exact statements are `Data.exists_isMarked`, `Data.IsFullMark.collisionSum_evenPlaces_eq_zero` and `Data.IsMarked.notMem_thick`. □

We will also need the balancing property of marking. If $G(c, l)$ is any function of an integer weight and a prime, zero unless $l \in \Lambda$, then

$$\sum_{\tau} G(w_j(\tau), l(\tau)) = \sum_{\tau} G(w_j(\tau), \overline{l(\tau)}). \tag{14.7}$$

Indeed multiplication by u exchanges the two primes. All places in these paired orbits are non-thick by the previous lemma, so the functions w_j are constant there, with the common weight determined by b_2. Reindexing the first sum by $\tau \mapsto u\tau$ proves the equality. The source's exact assertion is `Data.IsMarked.sum_inducedPlace_sub_star_eq_zero`. The equality will ensure that the local factors from the splitting characters in the Hecke transfer are integral.

14.6 Moving the marked Hecke operators through the chain

For each marked l, consider the spherical Hecke operator T_{l,s_p} corresponding to

$$\mathrm{GL}_k(\mathcal{O}_{E_l}) \,\mathrm{diag}(p\mathbf{1}_{s_p}, \mathbf{1}_{k-s_p}) \,\mathrm{GL}_k(\mathcal{O}_{E_l})$$

on the column space. We use the unnormalized coset-sum convention of the source's `heckeSplit`. Put $A_l = a\#\{\tau \in \mathcal{T} : l(\tau) = l\}$ and set $Y_l = p^{A_l} T_{l,s_p}$ and $Y = \prod_{l \in \Lambda} Y_l$ in a fixed order. By the scale constraints p is outside the column levels g and N. The local group is split and p is a uniformizer at l. The use of s_p rather than just the rank-one Hecke operator is part of the local repair strategy.

Let $J_{p,j}(E_0)$ denote the jet exponent for the jth lift established in the preceding section, and let $L_j(\tau)$ denote the number of columns in the weight at τ, below n_{j+1} and of heights other than one. Define

$$D_{p,j}(\Lambda) = s_p \sum_{\tau \in E_0(\Lambda)} L_j(\tau) - J_{p,j}(E_0(\Lambda)).$$

The quantity $D_{p,j}(\Lambda)$ may be negative; it is a logarithmic p-adic gauge exponent, not a loss required to be nonnegative. The definition is `Data.liftLoss` in `New/Columns/LocalStatement.lean`. For the following discussion a jet datum multiplying gauges by at most p^u will mean exactly membership in the source's submonoid `Kernel.raises` with prescribed frames and bound p^u.

Proposition 14.9 (Two local compatibility assertions, formal-source inputs)**.** *Let p be small nonsplit, let Ψ be good for a twist with $p \nmid q_\theta$ and let $l \in \Lambda$. For every $j < m$:*

1. *There are integral coefficients $c_0, \dots, c_{s_p}$ such that, on the partial definite chain $\mathsf{C}^{j+1}_{\theta,\xi}$ with any $\xi \in \operatorname{supp} \Psi$,*
$$\sum_{i=0}^{s_p} c_i T^{(j+1)}_{l,i} \mathsf{C}^{j+1}_{\theta,\xi} v = p^{\delta_p} \mathsf{C}^{j+1}_{\theta,\xi} T^{(0)}_{l,s_p} v$$
for spherical columns v, with spherical target datum at l.

2. *If each u_l belongs to one of the basic double cosets at l of index $i_l \le s_p$, the modified jth datum, translated on the target side by the product of the u_l, multiplies gauges by at most*
$$p^{\sum_{l \in \Lambda} A_l + D_{p,j}(\Lambda)}.$$

The declarations are `Data.IsGoodDatum.exists_hecke_transfer` and `Data.IsGoodDatum.mem_raises` in `New/Columns/LocalStatement.lean`. We explain precisely how the stated estimates enter the chain, and the source of each term.

At a split unramified place, the Hecke polynomial of a theta lift differs from that of the preceding stage by a polynomial with explicit powers of the size of the residue field and values of splitting characters. In the proof the source polynomial is evaluated with a rank-dependent normalization, and the constant term of the extra polynomial is 1. One can therefore recover its coefficient of degree s_p by multiplying the target polynomial by the truncated formal inverse of the extra polynomial. This is simple polynomial algebra: for $U(X) = 1 + u_1 X + \cdots$ over a ring, $U(X)^{-1} \bmod X^{s_p+1}$ has coefficients in the same ring, recursively $v_j = -\sum_{i=1}^{j} u_i v_{j-i}$. The unramified transfer identity for the single kernel is applied successively to all earlier lifts before extracting the coefficient. The weight balance (14.7) controls the norms of the values of the splitting characters at l; the coefficient of extraction uses at most p^{δ_p} to clear the rank normalizations. The exact normalized coefficient identity is the deeper part of assertion (1); it is not a consequence of unnormalized Satake transfer without the denominator computation.

For assertion (2), write each basic double coset representative as an integral invertible matrix times a diagonal matrix times an integral invertible matrix. At the marked places its diagonal has p in a set of at most s_p coordinates, and at conjugate places its entries are inverted by the unitary action. At infinite places this changes the coefficient frame for the local Schur-jet estimate. The top height-one columns contribute a units of gauge cost for each top inducing l, yielding A_l. At the other columns one incurs at most s_p times the number of columns at even places. The tilted Schur-jet bound supplies divisibility $J_{p,j}(E_0(\Lambda))$. The factors from determinant characters cancel between conjugate marked places, again by (14.7). This gives the exponent in assertion (2). It is crucial that the theta-jet estimate applies to

the actual tilted data: the Hecke operator has modified the kernel vectors, rather than merely changed their scalar coefficient by p^{A_l}. The proof of `mem_raises` composes the diagonal-frame jet estimate of the preceding section with integral covariance.

Lemma 14.10 (Iteration of the repair)**.** *For integral columns v and $p \nmid q_\theta$, with Ψ good,*

$$\|\mathsf{B}_\theta^\Psi(Y^m v)\|_\iota \le p^{\sum_{j<m}(|\Lambda|\delta_p + D_{p,j}(\Lambda))}.$$

Here the bound is on every gauge of the restricted chain.

Proof. Move one marked operator through the partial chain ending at $j+1$, using Proposition 14.9(1). On the jth kernel datum this replaces it by the operator

$$p^{A_l-\delta_p}\sum_{i\le s_p} c_i T_{l,i}^{(\mathrm{datum})}$$

on the finite Weil representation on the target side of that kernel. On its gauge the scalar factor contributes $p^{\delta_p - A_l}$. Target-translation covariance and the spherical invariance of the datum guarantee this operator commutes with the required simultaneous level actions. When the same operation is applied at a second marked place, the first remains in the same finite kernel; the local groups at distinct places commute. Repeating for a list I of distinct marked primes gives a new datum at stage j and, for the as yet unprocessed local translates, gauge exponent

$$\sum_{l\in I}(\delta_p - A_l) + \sum_{l\in\Lambda} A_l + D_{p,j}(\Lambda).$$

Indeed the remaining Hecke actions are sums of basic coset translates as in assertion (2), with integral coefficients; the ultrametric inequality does not change the exponent when adding these terms. Taking $I=\Lambda$ leaves $|\Lambda|\delta_p + D_{p,j}(\Lambda)$. Modify every stage by this construction. Inductively the resulting first j kernels act on v as the original partial chain on $Y^j v$: the new last kernel has exactly absorbed one more Y through the partial chain. Thus the modified full chain equals the original chain on $Y^m v$. Lemma 14.3 adds the exponents. Finally the restriction to Y_m does not increase gauges, since $p \ne N$. This is the deduction `Data.IsGoodDatum.exists_move_markOperator` and `Data.IsGoodDatum.gaugeLe_markOperator_pow`. □

Here is the numerical consequence. The jet exponent of the preceding section obeys the exact inequality

$$J_{p,j}(E_0) \ge \frac{g_p}{m} - C_p(E_0) + \omega_p \#(\mathcal{T}\cap E_0)\frac{a}{x_1} - \frac{f a(\log b)^2}{pm}. \tag{14.8}$$

For marked primes the other bound required is

$$|\Lambda|\delta_p + s_p \sum_{\tau\in E_0(\Lambda)} L_j(\tau) \le \frac{\ell_p}{m} + \#(\mathcal{T}\cap E_0(\Lambda))\frac{a}{x_1}. \tag{14.9}$$

This is `Data.IsMarked.sum_lowColumns_le` in `New/Columns/JetExponentBound.lean`. To explain the special top contribution here, the diagram count gives $L_j(\tau) \le w(\tau) + x^{60}$ and the number of boxes in these columns is between $kw(\tau)$ and $kw(\tau)+x^{60}$. Every marked place is non-thick, hence $w(\tau) = a/x_1$. For marked nonempty Λ the prime is served and not short. If $p \le m$, the factor $s_p \le 1 + \log_p m$ times a/x_1 is negligible in ℓ_p/m because of the separation $x \ll x_1$. If $p > m$, then $s_p = 1$ and the served condition forces $o_0 = 1$. Any even place in a marked orbit of tops is therefore itself a top. The contribution a/x_1 per such place is accounted for exactly by the term on the right of (14.9). In both cases $|\Lambda| \le f$, and the

terms $|\Lambda|\delta_p$ and $s_p f x^{60}$ are absorbed by the remaining margin $fa(\log b)^2/(pm)$. The formal calculation makes the constants explicit uniformly up to $p < x_1$; its scale inequalities are used here as formal-source bounds. The large a/x_1 term in (14.9) could not simply be discarded: for p near x_1 it would be too large. It is exactly matched by the positive term in (14.8).

For a full marking, $C_p(E_0(\Lambda)) = 0$ and $\omega_p = 1$. Combining (14.8), (14.9) and Lemma 14.10 gives

$$\|\mathsf{B}_\theta^\Psi(Y^m v)\|_\iota \le p^{2\ell_p - g_p}. \tag{14.10}$$

There is also a useful estimate with no marking at all. One may take $\Lambda = \varnothing$ in Lemma 14.10, with no nontrivial operators. Then (14.8) gives

$$\|\mathsf{B}_\theta^\Psi(v)\|_\iota \le p^{mC_p(\varnothing) + \ell_p - g_p}. \tag{14.11}$$

Writing

$$j_p = \left\lceil \frac{2mfa}{p} \right\rceil,$$

the estimate $C_p(\varnothing) \le 2fa/p$ and scalar linearity imply

$$\|\mathsf{B}_\theta^\Psi(p^{j_p} v)\|_\iota \le p^{\ell_p - g_p}. \tag{14.12}$$

These are the declarations `Data.IsGoodDatum.gaugeLe_Bmu_of_isMarked`, `Data.IsGoodDatum.gaugeLe_collisionSum` and `Data.IsGoodDatum.gaugeLe_pow_jetCeil_smul`, all in `New/Columns/LocalStatement.lean`. Thus the submodule

$$Y^m M + p^{j_p} M$$

has good divisibility for every admissible twist prime to p. In particular, its definition and the gain bound do not involve the twist. What remains is to prove that it has small index; this is a question about algebraic modular forms and the special Frobenius-twisted Hecke operator.

14.7 The Frobenius-twisted elementary-divisor theorem

We give the local theorem in its representation-theoretic form, because its application to the automorphic lattice is not valid unless both the operator and its coordinate normalization agree.

Fix a small served prime p, and a marked place l of E. Put $u = -p$ as above, $P_0 = p^{o_0}$, and let ν be the order of u^{o_0} in the CM place group. The tops inducing l or $\bar{l}$ are exactly the orbit of a chosen one under this element, so they may be labelled by $s \in \mathbb{Z}/\nu$. Since o_0 is odd, the parity of $l, \bar{l}$ alternates along the orbit. The residue field of l has cardinality $Q_l = P_0^\nu$ and ν is even. This uses the equality $q_l = p^{\mathrm{ord}(u)}$ in the served case, as in `Data.IsMarked.hasType_markOperator`. Here ν is a local orbit length and is not the auxiliary level prime N.

Write R_l for the valuation ring of E at l. There are embeddings $\iota_s : R_l \to O_\iota$ whose reductions satisfy

$$\iota_{s+1}(x) \equiv \iota_s(x)^{P_0} \pmod{p}.$$

In the odd letters the embedding in the coefficient representation also includes conjugation, so that the action of $g \in \mathrm{GL}_k(R_l)$ in the sth factor is by

$$g_s = \begin{cases} \iota_s(g), & s \text{ even}, \\ \iota_s((g^{-1})^t), & s \text{ odd}. \end{cases}$$

The weight module is $\mathcal{P} = \bigotimes_{s\in\mathbb{Z}/\nu} \mathrm{Sym}^a O_\iota^k$, represented by polynomials in variables x_{bs}, $0 \le b < k$, of degree a in each letter s. Let $\rho(g)$ denote the above representation. The other coefficient factors, not in this orbit, are left as a separate factor. Write $r_p = s_p$ for the size of the first block of coordinates. A variable is deep if either s is even and $b \ge r_p$, or s is odd and $b < r_p$. Denote the indicator of this condition by $\Delta(b, s)$. For a monomial x^α its depth is $\sum_{b,s} \Delta(b, s)\alpha_{bs}$. These are the exact definitions `TwistingData.letterRep`, `TwistingData.deepIndicator` and `TwistingData.depth` in `New/TwistedHecke/TwistingData.lean`. The dimension of $\mathcal{P}$ over the coefficient field is $D_\mathcal{P} = \binom{a+k-1}{k-1}^\nu$.

The following Euler- and weight-like notation is only local. Put $\lambda = \nu/2$ and $c_* = 2(k + k^2)o_0\nu^2 + 2$. The numerical conditions of the local theorem, with parameters ϵ, y, z, are

$$\begin{aligned} &0 < 2r_p \le k, \quad \epsilon \le 1 \le y, \\ &\nu ak(\lambda + 2) + a + k < P_0^\lambda, \quad 6c_* + 2 \le y, \\ &2(\nu + 2)k/y \le \epsilon, \\ &(y + \nu(k + 1))kP_0^{3r_p} + k + 2 \le a/y, \\ &\nu a\left(P_0^{-2r_p} + \frac{r_p^2}{kP_0}\right) \le z. \end{aligned} \tag{14.13}$$

This is `HeckeSizes` in `New/TwistedHecke/Counting.lean`. They allow the Frobenius degree to be large enough for interpolation while retaining a small elementary-divisor budget.

Why the normalized spherical operator has this weight. Write $d_p = \mathrm{diag}(p\mathbf{1}_{r_p}, \mathbf{1}_{k-r_p})$. In a single even letter, the contragredient coefficient action of d_p in the class-sum formula scales monomials by negative powers of p supported on the first block. Multiplying the action by p^a changes the power to the degree in the second block. At an odd letter the inverse transpose interchanges the two prescriptions. Thus multiplying the total Hecke operator by $p^{A_l} = p^{a\nu/2}$ gives precisely the diagonal matrix

$$D_p = \mathrm{diag}(p^{\mathrm{depth}(\alpha)}).$$

There is one factor p^a for every even letter. This explains the normalization of Y_l at the automorphic lattice, not just its complex normalization.

The root inequality. For distinct coordinates b, b' let $y_s = x_{b's}$, $z_s = x_{bs}$ if s is even, and interchange the choices if s is odd. For a fixed letter i define the excess functional

$$\mathrm{exc}_i(\alpha; b, b') = \frac{1}{Q_l} \sum_s \left(P_0^{1+[s-1-i]_\nu}\alpha(y_s) - P_0^{[s-i]_\nu}\alpha(z_s)\right), \tag{14.14}$$

where $[u]_\nu$ denotes the residue in $[0, \nu - 1]$. This is exactly the weight in `TwistingData.excessWeight`. Let $w \in O_\iota$ be a square root of p. For a set of real variable weights c_v, use the Gauss filtration: f is of order at least A if every coefficient of x^α is divisible by $w^{\max(0,\lceil A - \sum_v c_v\alpha_v\rceil)}$. This filtration is multiplicative: for every product of monomials the desired divisibility follows by adding the inequalities, then summing proves it for polynomials.

Lemma 14.11 (The elementary multiplicity count)**.** *If $\Phi \in O[T]$ is a nonzero polynomial, O a ring, and the images u_i of q distinct residue classes of a local ring have pairwise invertible differences in O, and $(T - u_i)^E \mid \Phi$ for all i, then $qE \le \deg \Phi$.*

Proof. The monic linear factors are pairwise comaximal and their powers are pairwise comaximal. Thus their product divides Φ. Multiplication by a monic polynomial adds its degree to the degree of a nonzero polynomial over any nonzero ring. Compare degrees. □

The precise root lemma of the local proof is stronger than a multiplicity count, and the following statement is the root result proved in Theorem 15.1, `TwistingData.root_lemma` in `New/TwistedHecke/RootLemma.lean`: suppose b, b' are in the same block of a Levi subgroup and, for every letter s,

$$\frac{2}{5} \le c_{y_s} - c_{z_s} \le \frac{8}{5}.$$

Suppose every Levi translate of F is in the Gauss filtration of order $l + \sum_v c_v \alpha_v$, and $w^{l+1} \nmid F_\alpha$. Then $\mathrm{exc}_i(\alpha; b, b') \le 0$ for each i. One cannot obtain this directly by counting zeros of the individual coefficient polynomials of degree a, because the Levi action uses different Frobenius powers in different letters.

We describe the reduction in the formal proof. Choose the transvection from b to b' composed with interchange of the two coordinates, with parameter $t \in R_l$. At a letter s the parameter in the variable substitution reduces to a signed $P_0^{[s-i]_\nu}$-power of the parameter at i. The coefficient of a selected monomial is read after identifying the two variables in letter i, and substituting appropriate powers of an auxiliary variable T in the other letters. More precisely, the exponents assigned to z_s in the selected coefficient are $Q_l P_0^{[s-i]_\nu}$ and those recording the forced zeros from y_s are $P_0^{1+[s-1-i]_\nu}$. Modulo w^{l+1} the hypothesis on F_α gives a nonzero polynomial in T whose degree is at most

$$Q_l \sum_s P_0^{[s-i]_\nu} \alpha(z_s).$$

Its Taylor coefficients in degrees below $\sum_s P_0^{1+[s-1-i]_\nu} \alpha(y_s)$ vanish at every residue class t in R_l. The nonzero coefficient assertion in this calculation uses the minimum for the Gauss filtration: otherwise different monomials could contribute to the selected coefficient and invalidate the argument. The discarded substitutions differ by terms of higher Gauss order; this follows from the Frobenius congruence modulo $p = w^2$ and the two-sided bounds on the differences of variable weights. Thus the multiplicity count gives $Q_l \sum_s P_0^{1+[s-1-i]_\nu} \alpha(y_s) \le Q_l \sum_s P_0^{[s-i]_\nu} \alpha(z_s)$, which is the asserted inequality. This describes the coefficient calculation of the root lemma; the general-ring Gauss-filtration substitution estimates which justify its nonzero coefficient and all its divisibilities are part of that input. The essential gain is that each residue class contributes the same weighted multiplicity, and there are Q_l of them.

Two dimension estimates. Reduce coefficients modulo the maximal ideal and call the residue coefficient space $\overline{\mathcal{P}}$. Let $\mathcal{P}_{\mathrm{lev}}$ be the two-block parahoric of $\mathrm{GL}_k(R_l)$, namely the inverse image of the two-block parabolic over its residue field. Fix the adjacent coordinates $(r_p - 1, r_p)$. Define the subspace U_h of $\overline{\mathcal{P}}$ by requiring in every block-Levi translate that monomials of excess greater than h at this adjacent pair vanish, in every letter. Let V_j be the subspace of $\overline{\mathcal{P}}$ of forms such that every $\mathrm{GL}_k(R_l)$ translate has depth at least j. These are `TwistingData.excessCond` and `TwistingData.deepForms`.

The two relevant estimates are

$$[\mathrm{GL}_k(R_l) : \mathcal{P}_{\mathrm{lev}}] \dim U_y \le (1+\epsilon) D_{\mathcal{P}}, \tag{14.15}$$

$$\sum_{0 \le j < J} \dim V_j \le (1+z) D_{\mathcal{P}} \quad (J \ge 0). \tag{14.16}$$

These are `TwistingData.index_mul_finrank_excessCond_le` and `TwistingData.sum_finrank_deepForms_le`, in `New/TwistedHecke/Counting.lean`. We explain the combinatorics yielding these bounds. The uniform Frobenius minimum-weight calculations in their proof are proved in Chapter 15, with the numerical bounds specified below.

The first count uses the following cyclic interlacing cone. For a real array v_{bs} put $\beta(b,s) = \pm(P_0 v_{b,s+1} - v_{b,s})$, with sign positive in the even letters and negative in the odd letters. This is an invertible linear transformation: if $P_0 v_{b,s+1} = v_{b,s}$ for every letter, then cycling gives $(P_0^{\nu} - 1)v_{bs} = 0$. Substitution into (14.14), telescoping around the letters, gives

$$\mathrm{exc}_i(\beta; b, b') = (1 - Q_l^{-1})\big(\beta(y_i) - (v_{b,i} - v_{b',i})\big).$$

Thus the zero-excess inequalities are exactly the interlacing conditions

$$\begin{cases} P_0 v_{b+1,s+1} \le v_{b,s}, & s \text{ even}, \\ v_{b+1,s} \le P_0 v_{b,s+1}, & s \text{ odd}, \end{cases} \qquad 0 \le b < k-1.$$

Let $\Sigma(D)$ be the product of slabs $\beta_{bs} \ge 0$, $D_s - 1 \le \sum_b \beta_{bs} \le D_s$. The cone of zero excess, intersected with $\Sigma(D)$, has volume exactly

$$\frac{\mathrm{vol}(\Sigma(D))}{\displaystyle\prod_{j=1}^{k} \frac{Q_l^j - 1}{Q_l - 1}}. \tag{14.17}$$

Here is the signed-volume identity underpinning this formula. For decreasing real vectors x, y of length k, the matrix $\mathbf{1}_{y_c < x_b}$ has determinant 1 precisely when $x_1 > y_1 \ge x_2 > y_2 \ge \cdots \ge x_k > y_k$; otherwise it has determinant zero. This follows because the row of index b ends with a string of ones, preceded by zeros, and the nested rows are independent exactly when all their threshold positions differ. Take one such matrix of interlacing indicators for each pair of adjacent letters, using transpose in the alternating letters. Expanding the product of determinants yields a signed sum over tuples of permutations, whose product permutation will be denoted π'. The corresponding linear changes of variables $v \mapsto \beta$ have determinant, up to a common factor independent of π',

$$\det(1 - Q_l^{-1} A_{\pi'}),$$

where $A_{\pi'}$ is the permutation matrix. This follows by solving the equations around the cyclic chain after successively reordering the coordinates: the sole boundary condition is given by the product permutation. The walls where equalities occur have measure zero. Consequently integrating the signed indicators and accounting for sorting yields

$$\frac{\mathrm{vol}(\mathrm{cone} \cap \Sigma(D))}{\mathrm{vol}(\Sigma(D))} = \frac{(1 - Q_l^{-1})^k}{k!} \sum_{\pi' \in S_k} \frac{\mathrm{sgn}(\pi')}{|\det(1 - Q_l^{-1} A_{\pi'})|}.$$

The formula, including the cancellation of all intermediate permutation multiplicities, is `volume_inter_slab_eq_signed_sum` in `New/TwistedHecke/InterlacingCone.lean`. To evaluate the signed sum, it suffices to compare with the single-letter case with parameter Q_l instead of P_0. In that case the inequalities are $\beta_b \ge Q_l \beta_{b+1}$. The triangular change

$$\beta_b \longmapsto d_b(\beta_b - Q_l \beta_{b+1}), \qquad d_b = 1 + Q_l + \cdots + Q_l^b \quad (b = 0, \dots, k-1)$$

with $\beta_k = 0$ carries the cone onto the nonnegative orthant. It preserves the sum of all coordinates because $d_b - Q_l d_{b-1} = 1$ and $d_0 = 1$; its determinant is $\prod_b d_b$. This gives (14.17). In the source this step is `prod_mul_volume_inter_slab`. The factor in the denominator is exactly the number of complete flags in a k-dimensional vector space over the residue field of size Q_l. This exact match explains why a leading constant $1 + \epsilon$ in (14.15) is possible.

The root lemma relates this volume to the dimension of U_y. To see the requisite amplification, one passes from a polynomial F to blocks of polynomials of the form $x^{\gamma(u)} F_u(x^D)$

with $D = Q_l^J$, and $u \in \{0, \dots, D-1\}^{k\nu}$. The exponents $\gamma(u)$ are congruent to u modulo D and have degree in each letter between kD and $(k+1)D$. Hence these summands are independent, and substitution $x \mapsto x^D$ commutes with the residual group action because D is a full Frobenius period. One chooses a least-weight monomial among its translates over the flags in the Levi subgroup. The root lemma at adjacent roots within either block forces the corresponding excess inequalities, with the chosen bound y retained at the single boundary root between blocks. A count of its available coefficients and the amplification limit $D \to \infty$ bounds the dimension by the number of Levi flags times the volume of the thickened cone of boundary excess $y + \nu(k+1)$. This is proved in Lemma 15.3, corresponding to `TwistingData.finrank_excessCond_le_volume`; it identifies the injection obtained by taking the relevant coefficients of the translates of the amplified polynomials.

To remove the thickening, the proof constructs a nonnegative vector σ with all coordinates at most $P_0^{3r_p}$, satisfying each zero-excess inequality and having boundary excess at most -1. One can see such a vector by taking $v_{b,s}$ proportional to $2P_0^{2(r_p-1-b)}$ for $b < r_p$ and zero for $b \geq r_p$, with the odd letters multiplied by P_0. Substituting into the linear map defining β and into the interlacing inequalities gives this bound. The assertion is `TwistingData.exists_mem_excessCone_mem_Icc`. Translation by $(y + \nu(k+2))\sigma$ embeds the thickened cone into the zero-excess cone; it changes the upper endpoint of each slab by at most $(y + \nu(k+2))kP_0^{3r_p}$, with allowance for the initial degree padding. The precise choice in the Lean proof uses $y + \nu(k+2)$ with each letter's shift and the size conditions (14.13). The volume of a slab of upper endpoint D_s is at most $D_s^{k-1}/(k-1)!$, as is seen by subtracting the volume of two simplices. With the size conditions the changed volume of all slabs is at most $(1+\epsilon)D_{\mathcal{P}}$ after multiplying by the flag count and canceling it against the denominator in (14.17). The exact inequalities including the roundings are those of `index_mul_finrank_excessCond_le`. They supply (14.15).

For the depth bound assign to each deep variable of coordinate $b \geq 2r_p$ cost $P_0^{-2r_p}$, and to each deep variable with $b < 2r_p$ cost $1/P_0$. All other variables have cost zero. The result proved in Proposition 15.6, `TwistingData.exists_le_weight_of_mem_deepForms` in `New/TwistedHecke/DeepFormCost.lean` says that any nonzero polynomial in V_j has a nonzero coefficient at a monomial of total cost at least j, provided $2r_p \leq k$ and $2r_p < \nu$. The source uses the upper unipotent matrix $1 + A$ with A supported in the block from the first r_p rows to the remaining rows and $A^2 = 0$. In the rational function field of the variables, the coefficients of A can be recovered by linear equations in Frobenius powers of the deep variables; evaluating each letter's variables in the specified order gives a triangular matrix with nonzero determinant. The equations express each matrix entry at residue points by polynomials of cost less than one. The condition of depth j in every translate then means vanishing to that weighted order on the whole family of residue points. Polynomial interpolation forces the existence of the claimed costly monomial. The recovery matrix and the weighted vanishing estimate are the substantive content of this input, rather than a generic vanishing-order assertion.

The rest of the depth argument is a direct count. Restriction to coefficients of cost at least j is injective on V_j by the stated input. Summing the resulting dimension bounds counts each monomial at most 1 plus its cost. There are $\nu/2$ deep letters for each fixed coordinate. Symmetry of the monomial basis gives average exponent a/k at each coordinate of each letter. Thus the mean cost over all monomials is at most

$$\frac{a}{k}\frac{\nu}{2}\left(2r_pP_0^{-1} + (k-2r_p)P_0^{-2r_p}\right) \leq \nu a\left(P_0^{-2r_p} + \frac{r_p^2}{kP_0}\right) \leq z.$$

This proves (14.16), using only the lower bound on cost in V_j and the last size condition in (14.13).

The approximate projector and the integral pairing. The dimension estimates must be converted into a determinant bound; they do not by themselves control Y_l. In particular, bounding the dimension of a conditioned representation modulo p does not give a bound on all Smith divisors of an arbitrary matrix with that representation in its target.

In the source, the approximation is constructed using an invariant diagonal pairing on degree-a polynomials. For a multi-index α of degree less than p^F, its multinomial coefficient has p-valuation at most (#variables)F. Indeed Legendre's formula for the factorial expresses the valuation as a sum of carry contributions at at most F levels, bounded by the number of input summands at each level. Consequently one may choose integral diagonal entries proportional to the inverse multinomial coefficients, preserving transpose covariance of the action. The exact formula is `ClassMean.exists_invariant_diagonal_form` in `New/TwistedHecke/ClassMeans.lean`.

A second input in that file, `ClassMean.exists_mean`, constructs an invariant linear form on polynomials of a generic Levi matrix. If x is a weight function on the matrix coordinates and P is a condition on the row weight, the linear form kills monomials whose row and column weights differ; on monomials satisfying P it agrees with evaluation at the identity multiplied by p^{k^2F}, up to the prescribed degree cutoff. It is in the span of Levi translates of linear forms killing monomials not satisfying P. The hypotheses are degree cutoff $D < p^F$ and $D + k < \#(R_l/\mathfrak{m}_l)$. For orientation, the finite-residue interpolation behind this fact is elementary: evaluation on the matrices with unit determinant spans the dual of the polynomial coefficient space in degrees D if $D + k$ is less than the residue cardinality. A polynomial vanishing on all such matrices, multiplied by their determinant, vanishes on the entire finite matrix space. The degree bound and successive one-variable root bounds force it to be zero. Lifting linear independence modulo the maximal ideal gives the integral statement on the finite free coefficient module. The invariant trace construction which incorporates the row-weight restriction is part of `exists_mean`.

Taking tensor products of these class means in the cyclic letters gives the approximate projector. In the Frobenius product of generic matrices the factor at the dth position is the entrywise P_0^d power of the generic matrix from letter $s - d$. At each letter one takes the first λ factors, plus one additional factor in each deep row; adjugate transpose takes the place of ordinary multiplication in odd letters. The construction tracks the Gauss weight and hence the excess of each output monomial. If the generic monomial has equal row and column weights outside a single letter, and the discrepancy between its weight at that letter and the output excess is greater than $2c_* + 1$, the coefficient of this monomial is divisible by w^{c_*}. This is proved in Proposition 15.9, corresponding to `TwistingData.pow_dvd_coeff_conj_frobOperator` in `New/TwistedHecke/FrobeniusFactors.lean`; it follows from the multiplicative Gauss weight bound `TwistingData.frobOperator_mem` and the Frobenius congruence for conjugated generic matrices in the same file. That bound is consistent with the local scaling $p = w^2$.

The resulting precise assertion, proved in Theorem 15.10, is `TwistingData.exists_projector`, in `New/TwistedHecke/ElementaryDivisors.lean`. There are a diagonal pairing matrix b_* and an operator Z_* such that Z_* maps every coefficient vector into the integral excess condition of bound y modulo w^{c_*}, and

$$y_0^t b_* Z_* F \equiv p^{(k+k^2)o_0\nu^2} y_0^t F \pmod{w^{c_*}}$$

for y_0 in the smaller excess condition with threshold $y - (4c_* + 2)$ modulo w^{c_*}. Here Z_* is not asserted to be an idempotent. The transpose compatibility $b_*\rho(\gamma^t) = \rho(\gamma)^t b_*$ is part of the statement. The precision $c_* = 2(k + k^2)o_0\nu^2 + 2$ leaves two extra powers of w beyond the scalar in the congruence. This pairing estimate is the nontrivial bridge between the counting result and the determinant calculation.

We describe the final abstract determinant theorem, proved in Section 15.8, so that its applicability to the coefficient lattice can be verified. In the following statement let X be a finite class set, I a coefficient index set identified with the monomial indices of $\mathcal{P}$ times a finite set H_0 of spectator indices (the remaining coefficient factors). Let $J = \mathrm{GL}_k(R_l)/\mathcal{P}_{\mathrm{lev}}$. Suppose for $e = (c, b) \in X \times J$ we have class $s(e) \in X$, matrix $g_e \in \mathrm{GL}_k(R_l)$ and integral invertible matrix u_e on the spectator indices, with

$$e \longmapsto (s(e), [(g_e^t)^{-1}]) \in X \times J \quad \text{injective.}$$

If γ_b are the chosen parahoric-coset representatives and $F = (F_c)_{c \in X}$ is a family of coefficients, consider

$$(TF)_c = \sum_{b \in J} \Big(u_e \otimes (\rho(g_e) D_p \rho(\gamma_b))^t \Big) F_{s(e)}.$$

With the orientation of representatives chosen in this formula, the theorem `TwistingData.hasType_twisted_hecke` states that on any linear image of the coordinate module to which T descends, the descended map preserves that module and has type bound

$$(z + \epsilon s + y)|X||I| \quad \text{modulo } p^s \tag{14.18}$$

when (14.13) holds. Its algebraic proof is in Section 15.8; application to the actual adelic coefficient lattice still requires the concrete compatibility arguments.

For its proof the map $F \mapsto (w^{\mathrm{depth}} \rho(\gamma_b) F_{s(e)})_e$ is split by a Hermite–Smith factorization into a primitive factor and a square factor. The determinant of the square factor divides a power of w with exponent at most the sum of dimensions of the depth spaces over the required truncation range. One way to read this bound is to reduce a column of the Smith factor modulo the maximal ideal: a column whose divisor exceeds w^j is forced into the jth depth condition in all frames. The independence of the reduced primitive columns bounds the count of such divisors by the dimension of that condition. This is `TwistingData.exists_depth_factorization`, which also ensures the primitive factor's coefficients lie in the integral excess condition with prescribed precision. Apply this factorization to each of the two half-depth factors of D_p (using $w^2 = p$). The half-factors are paired through Z_* so that the scalar $p^{(k+k^2)o_0\nu^2}$ controls the primitive part of the product, except for excess-condition vectors of total dimension bounded by (14.15). There is a second Smith factorization on the discrepancy matrix between the two primitive images. It gives the abstract pairing assertion `TwistingData.of_pairing`: if $RQ = 1$, $\beta M \equiv a_0 I \pmod{a_0 w}$, and the normalized difference $Qu - Mv = a_0 w F$ imposes the excess condition on F, then the image of βQ enlarged by p^s has index dividing $a_0^{|I'|} p^{sl}$, with $|I'| + l$ bounded by the dimension of the excess condition (including the class and coset multiplicities). Here I' is the domain index set of the primitive matrices. This is proved in the source by applying Smith form to $(QRM - M)R$, assessing each divisor relative to $a_0 w$, and using independence modulo the maximal ideal. The counts (14.15) and (14.16) now account for the three terms ϵs, z, and y in (14.18): the first is the exceptional share of the cokernel charged at depth s, the second is the aggregate half-depth cost, and the third pays for the controlled scalar in the projector. These contributions are additive by Lemma 14.2. This description specifies the remaining determinant theorem needed in the proof, including its hypotheses; proving the full matrix approximation independently would require the exact Smith-factor and projector identities, beyond the counting arguments already given here.

14.8 Application to the actual column lattice

The abstract coefficient operator above occurs on M itself, rather than on a putative subspace with convenient coefficients. This fact uses the trivial class stabilizers at K_0 and the availability

of class representatives integral above p. The latter assertion is `exists_reps_integralAbove`, used in `New/Columns/TwistedHeckeType.lean`. For each class representative γ_c, write the Hecke sum at the point γ_c using the local double-coset representatives indexed by $\mathrm{GL}_k(R_l)/\mathcal{P}_{\mathrm{lev}}$. Each product is rationally equivalent to one chosen representative $\gamma_{s(e)}$, times a compact-level element. Triviality of class automorphism groups gives injectivity of $e \mapsto (s(e), [(g_e^t)^{-1}])$: if the class and the parahoric coset agree, the two chosen rational identifications differ by an automorphism of the representative, which must be the identity, and the local cosets agree. The other primes above p give integral invertible matrices on the spectator factors. Thus in the coordinate identification of algebraic forms the normalized Hecke operator has exactly the shape of (14.18), including the depth diagonal. The exact comparison, with the class-representative and gauge conditions, is `Columns.hasType_markOperator` in `New/Columns/TwistedHeckeType.lean`. We use this coordinate comparison as a formal-source input. It is essential that the integral image addressed by that theorem is all of M, as opposed to the coefficient module with unrelated adelic transition functions.

The local congruence and numerical hypotheses are verified for our marked place by `Data.IsMarked.hasType_markOperator`, in `New/Columns/MarkedPrimeType.lean`. The precise result is the following.

Proposition 14.12 (Marked operator type)**.** *For each $l \in \Lambda$ and each integer $s \geq 0$, the operator Y_l preserves M and*

$$Y_l \in \mathrm{Type}\left(p, M, s; \left(\frac{128 A_l \log b}{mp} + \frac{s}{x^{14}} + x^{40}\right) c\right).$$

Indeed on the local orbit discussed above one has $A_l = a\nu/2$, the parity of the local embeddings is alternating, and the orbit size is even. The choices of the generic theorem's parameters are

$$\epsilon = x^{-14}, \qquad y = x^{40}, \qquad z = \frac{128 A_l \log b}{mp}.$$

The relation $P_0^{\lambda} \geq p^{m/2}$, from the divisibility of the order of u by m, supplies the interpolation inequality in (14.13). The served condition $o_0 \leq s_p$ gives $P_0 \leq p^{s_p} \leq pm$. The bounds $f \ll x^{11}(\log b)^{1/5}$ up to fixed factors, $k \asymp x^5$, $\nu \leq f$ and the hierarchy $x \ll x_1 \ll b$ leave room in the remaining inequalities; for example $P_0^{-2r_p} \leq m^{-2}$, and $r_p^2/(kP_0)$ is negligible against $\log b/(mp)$ on the admissible range. More precisely $P_0^{2r_p} \geq p^{2s_p} \geq mp$ since $P_0 \geq p$ and $m, p \leq p^{s_p}$. The large-prime range is covered by the separate estimate $s_p = 1$ when $p \geq m$. The stated Lean declaration checks these estimates with the fixed constants in (14.13), including the degree condition when ν and x_1 vary. Thus the proposition is a quantitative application of the coordinate comparison and the local theorem, not an assumption about Hecke eigenvalue slopes.

Corollary 14.13 (Type bound for the repair)**.** *With Y and j_p as above there exists an integer $e_p \geq 0$ such that*

$$I(M, Y^m M + p^{j_p} M; p^{e_p}), \qquad e_p \leq 3\ell_p c.$$

Proof. By Lemma 14.2 and the proposition, the exponent bound for the product of the marked operators, repeated m times, is at most

$$cm \sum_{l \in \Lambda} \left(\frac{128 A_l \log b}{mp} + \frac{j_p}{x^{14}} + x^{40}\right).$$

Now $\sum_l A_l \leq fa$, $|\Lambda| \leq f$, and $j_p \leq 2mfa/p + 1$. Thus the first contribution divided by $c\ell_p$ is at most $128/\log b$, the second at most

$$\frac{2m^2 f}{x^{14}(\log b)^2} + \frac{mp}{ax^{14}(\log b)^2},$$

and the third at most $mx^{40}p/(a(\log b)^2)$. All are at most 1 on the admissible scale, the second using the bound on f and $m^2 \le 4x$, and the third using $p < x_1$ and $x_1^{10^6} \le b$. The precise three-term comparison with the fixed constants is `Data.IsMarked.hasType_prod_markOperator_pow`. The type condition gives integrality and an integer witness e_p for the desired index. □

The crucial point is that the type estimate grows with the *truncated mean* of elementary divisors rather than with their largest possible value. Charging the whole space for one large elementary divisor would multiply ℓ_p by m and defeat the saving.

14.9 The prime N: moving denominators to a Fourier operator

The additional prime N splits completely in E and does not appear among the small nonsplit primes. Its effect in the complex bound is the large negative term in (14.3). At the finite places of N one must recover a usable lattice estimate despite the large negative powers of π in the restriction and the translations by y/N in the good datum.

Fix ι above N and let S be the set of conjugates of the primes induced by the given CM-type embeddings at ι (`Data.nonInducedPlaces`). At these split places use the coordinate embedding of the general linear group into the unitary group. Let d_j be the finite point whose component at the places in $S \cap \{v \mid \pi\}$ is $\mathrm{diag}(1, \dots, 1, N)$, with the corresponding conjugate component and identity elsewhere. These are `Data.diagPointN`. In the representation conjugated by d_j, the column level has mirabolic reduction, i.e. its last column is the final basis vector modulo N. The relevant level identity is `Data.atPlaces_conj_mem_mirabolic` in `New/Seesaw/LiftCharacters.lean`.

Let F_N be the Fourier operator on columns, given in this conjugated representation by multiplication by the inverse last-coordinate diagonal and a sum over integral matrices with prescribed final row and column, weighted by the additive character of their final residue coordinate. The formula in `IsSplitLevel.fourierOp` (`New/Seesaw/NamesAtSplitLevel.lean`) is

$$F_N = \rho(\mathrm{diag}(1, \dots, 1, N^{-1})) \sum_{y,\ y_{\mathrm{last}}\ \mathrm{unit}} \psi_N(y_{\mathrm{last}})\rho(\widetilde{g}_y).$$

All coordinates here are at the places of S, with the conjugated representation understood. Each $\widetilde{g}_y$ is an integral invertible lift of a matrix having final row and column determined by y. The additive character takes root-of-unity values. This operator preserves the column level (`Data.fourierOp_mem_column`) and is compatible with the ramification condition at N.

The key finite calculation is local and independent of the large weights. If L is a single theta-lift law, $\mathbf{b}$ is the indicator of the final-column condition and $\mathbf{c}$ the associated additive-character datum, then moving the diagonal off the character datum gives the Fourier operator applied to the indicator lift; it also equals a scalar times the indicator lift applied to the Fourier operator on the source. In the notation of `LiftLaw.u$_0$_charDatum_eq_fourierOp`, this is

$$\rho'(u_0)L[\mathbf{c}]v = F_N' L[\mathbf{b}]v = \gamma L[\mathbf{b}]F_N v.$$

The equality uses ramification: vectors in the image of $1 - z_v$ kill the terms in the finite character expansion which are invariant under that local scalar translation. One can consequently repeat the identity $m/2$ times to move the weighted Fourier operators from the paired chain onto the beginning of the column chain. This iteration is `compUpTo_pairs_eq_smul` in `New/Seesaw/NamesAtSplitLevel.lean`; its proof tracks the number of residue columns nonzero at every place and the scalar splitting-character factors at each stage. The finite iteration is proved in Proposition 15.14, conditional on the exact normalized lift law; that

actual geometric normalization is still a formal-source input. Applying it to the paired expression of the good datum yields `Data.chain$_2$_pairTranslates_eq_smul_chain`, `New/Columns/TwistedHeckeType.lean`: for every admissible twist, the paired chain of m kernels on a ramified column is a scalar of N-adic absolute value at most N^{2mfb} times a chain with data cut by $\mathbf{b}$, read in the frames d_j, acting on $F_N^{m/2}v$.

Each new cut kernel multiplies gauges by at most N^{fb}. This is `Data.name_lastColDatum_mem_raises`, same file; it is the crude CM integrality estimate applied to the cut datum rather than to arbitrary translates with denominator N. The branching restriction has the same gauge bound at N when its source gauge is computed at d_m: the negative powers of $\bar{\pi}$ are canceled by the corresponding final-coordinate powers in the branching formula. This is `Data.GaugeLe.restrictToY_of_diag` in `New/Seesaw/LiftCharacters.lean`. Both are used as precise formal-source inputs. Consequently, by Lemma 14.3,

$$\|\mathsf{B}_\theta^\Psi(v)\|_\iota \le N^{3mfb} \tag{14.19}$$

if $v \in C^{\mathrm{ram}}$ and $F_N^{m/2}v$ belongs to the column lattice measured at the frame d_0. This is `Data.gaugeLe_Bmu_at_N`.

The index cost of imposing this condition is much smaller than charging every coefficient its worst weight at the last coordinate. To see this, for each coefficient index I let $\alpha_I(\tau, b')$ be its tableau weight in coordinate b' at τ. For the initial column space these are just monomial exponents, and the sum over the coordinate b' in any factor is at most a: the factor is of degree a on the tops and degree zero elsewhere. With $\mathcal{X}$ the class set and $\mathcal{I}$ the Weyl indices put

$$T_{b'} = \sum_{c' \in \mathcal{X}} \sum_{I \in \mathcal{I}} \sum_{\tau \in \Phi} \alpha_I(\tau, b').$$

Coordinate symmetry of the initial weight implies $T_{b'}$ is independent of b', and the degree bound gives $\sum_{b'} T_{b'} \le fac$. In particular $T_{\mathrm{last}} \le fac/k$. The class-space action in the chosen coefficient frames is diagonal for the last-coordinate scaling: it multiplies a coefficient by $N^{\sum_\tau \alpha_I(\tau,\mathrm{last})}$, so its determinant is $N^{T_{\mathrm{last}}}$.

Precisely, if the column lattice at a frame $d_0 d(S')$ is changed to the lattice at d_0, where $d(S')$ is the inverse last-coordinate diagonal supported on a set of split places, the corresponding index divides $N^{T_{\mathrm{last}}}$. The adelic integrality here uses rational class representatives adjusted at N; the coefficient norm computation is a direct diagonal determinant comparison. The proof in the source pulls back the coordinate-diagonal index condition using the integral representative correspondence. Applying the Fourier operator consists of such a diagonal frame change and a sum of translations by integral invertible matrices; its character coefficients are N-adic units. Applying the pullback and transitivity rules of Lemma 14.2 once for each application of F_N, in addition to the initial frame change, gives an index bound dividing $N^{(m/2+1)T_{\mathrm{last}}}$. This is the assertion `Data.exists_indexDvd_fourierOp_pow` (`New/Seesaw/LiftCharacters.lean`), including the conditions on the changes of representatives. Since $a \le 2xb$ and $k \gg x^5$, the exponent is at most $4mfbc$ on the admissible data. Thus there is a submodule M'_N of M independent of the twist with

$$I(M, M'_N; N^e), \quad e \le 4mfbc,$$

on which (14.19) holds for all ramified columns. The presence of the average tableau weight rather than the largest coefficient degree is another place where symmetry of the integral weight representation is important.

14.10 The simultaneous local divisibility statement

Define a function on rational primes by

$$\epsilon_p = \begin{cases} 3\ell_p, & p < x_1,\ \chi_K(p) \neq 1, \\ 4mfb, & p = N, \\ 0, & \text{otherwise.} \end{cases}$$

Extend g_p by zero off the small nonsplit primes. For p outside those primes and $p \neq N$ the elementary integrality bound from the theta data suffices: every translated kernel is standard with integral gauge away from the twist conductor, and its scale $\bar{\pi}_g$per compensates for the denominators. There is no division by d, since in that case $p \neq d$. The resulting all-column estimate $\|\mathsf{B}_\theta^\Psi v\|_\iota \leq 1$ is `Data.IsGoodDatum.gaugeLe_one` in `New/Columns/LocalStatement.lean`. It is the application of the crude jet bound and Lemma 14.3 to the exact unmodified data.

Proposition 14.14 (Local divisibility and index)**.** *For every p, every ι above p, and every integer $\beta < g_p - \epsilon_p + 1$, there is a submodule M_p' of M and an integer $e \geq 0$ such that*

$$I(M, M_p'; p^e), \qquad e \leq \epsilon_p c.$$

For every admissible twist θ prime to p, every good datum Ψ and every $v \in M_p' \cap C^{\mathrm{ram}}$, the restricted lift $p^{-\beta}\mathsf{B}_\theta^\Psi v$ is integral at ι.

Proof. For a served small nonsplit prime take $M_p' = Y^m M + p^{j_p} M$ and the index supplied by Corollary 14.13. The two terms have gauges at most $p^{2\ell_p - g_p}$ and $p^{\ell_p - g_p}$ respectively by (14.10) and (14.12); the same bound $p^{2\ell_p - g_p}$ holds for their sum. Since $\ell_p \geq 1$, the condition on β gives $\beta + 2\ell_p - g_p \leq 0$, so multiplication by $p^{-\beta}$ makes the gauge at most one. For an unserved small nonsplit prime use $M_p' = M$, the collision bound Lemma 14.7, and (14.11). At N use the preceding frame-change index and (14.19); the condition $\beta < 1 - 4mfb$ ensures $\beta + 3mfb \leq 0$ since $mfb \geq 1$. At the other primes integrality is unconditional and the hypothesis $\beta < 1$ gives $\beta \leq 0$. This proves the proposition, including the common choice of submodule before the twist. It is exactly `Data.exists_indexDvd_Bmu_divisible`. □

The loss function is summable with the following quantitative bound:

$$\sum_p \epsilon_p \log p \leq B. \tag{14.20}$$

Here is a direct calculation, with generous constants for the scale comparison. If $p \equiv 1 \pmod m$ is prime, $p = um + 1$ with $u \geq 1$, so $m/p \leq 1/u$. If $p \equiv -1 \pmod m$, then $p = um - 1$ with $u \geq 1$, so $m/p \leq 2/u$ since $m \geq 2$. Dropping the primality conditions and bounding $\log p$ by $\log x_1$, one obtains

$$\sum_{p < x_1} \frac{\log p}{p} \begin{cases} m, & p \text{ short}, \\ 1, & \text{otherwise} \end{cases} \leq 4 \log x_1 (1 + \log x_1).$$

Thus the small-prime contribution is bounded by $12fa(\log b)^2 \log x_1(1 + \log x_1)$. Since $(\log x_1)^6 \leq \log b$ and $\log b$ is large, this is less than $B/2$. At N, one has $4mfb \log N \leq 120fm^2 b \log b \leq 480fa \log b$, using $m^2 \leq 4x$ and $a \geq xb$. This too is less than $B/2$ for the data in question. The statement with the precise admissibility thresholds is `Data.sum_allowedLoss_mul_log_le` in `New/Columns/SizeEstimates.lean`. It is the sparseness of the congruences for short primes which allows the factor m in ℓ_p and makes the global error budget independent of an overall factor m.

14.11 Archimedean growth and the level normalization

We now compare the finite-prime gains with the growth at every complex conjugate. Define

$$h_0 = m\left(\sum_{\tau\in\mathcal{T}}(J(a) - A(a,\tau)) + k\mathcal{U}\right) - \#\mathcal{T}\, a\log N.$$

Since $A(y,\tau) = 2\sum_p A_p(y,\tau)\log p$, it follows identically from the definition of $\mathcal{U}$ and g_p that

$$h_0 + 2\sum_p g_p \log p = m\left(\sum_{\tau\in\mathcal{T}} J(a) + k\sum_{\tau\in\Phi} J(w_0(\tau))\right) - \#\mathcal{T}\, a\log N. \tag{14.21}$$

This quantity is the leading exponent of the purely archimedean chain. In particular its expression in terms of prime asks is not a second analytic estimate; it is an exact algebraic identity, used to record the finite gain to be divided out.

The precise norm assertion of the definite chain is `Data.weylNormSq_Bmu_le`, `New/Columns/LocalStatement.lean`. For the trivial twist, good datum Ψ, and any conjugate σ, if all values of a column v have squared Weyl norm at most e^η, all values of its restricted lift obey

$$\|\sigma\mathsf{B}_1^\Psi v(g')\|^2 \le \exp\left(\eta + h_0 + 2\sum_p g_p\log p + B\right). \tag{14.22}$$

Here the norm assertion is conditional on the conjugate-jet bounds of the preceding section. We explain the size comparison to isolate the exact quantities going into its error budget.

For one lift the principal contribution from the Young diagram is the sum of $J(l)$ over its row lengths. The uniform diagram estimate `Data.diagramWeight_le_by_index` gives each row length at place τ the bound

$$\begin{cases} \kappa_\tau(1) + b_1 + b_2 + h_2^2, & \text{row zero,}\\ w_0(\tau) + h_2^2, & \text{rows } 1,\dots,k-1,\\ h_2^2, & \text{remaining rows.}\end{cases}$$

Positive error terms in the first row can be regarded as additional degree beyond the a-row (at a top) or the degree-zero row. For any $l \le r' + e$, one has the elementary bound

$$J(l) \le J(r') + e\big(\log(r'+e+1) - \log(\Omega/g)\big).$$

Indeed $l! \le (r'+e)!$ and $(r'+e)! \le r'!(r'+e+1)^e$; the exponential scale $\Omega/g \le 1$ on the data. This gives `Data.jetSize_le_add` in `New/Columns/SizeEstimates.lean`. Using it for the rows of the diagram gives an upper bound with leading contribution $\sum_{\tau\in\mathcal{T}} J(a) + k\sum_{\tau\in\Phi} J(w_0(\tau))$. All lower-order row contributions, conjugate-jet errors and the difference between actual and nominal placement are bounded by a polynomial in the ranks and f times $b\log b$ or by similar smaller expressions. The source's size calculation following `diagramWeight_le_by_index` checks the corresponding inequalities with the fixed scale constants. The other terms in the complex triangle inequality of Lemma 14.3 are tightly bounded: the number of classes at each stage is at most $e^{x^{30}}$ (a uniform bound `Data.card_stdClasses_le_exp`, same file), the sum of absolute values of coefficients of the good datum is at most $e^{x^{60}}$, and the final branching produces the large negative exponent in (14.3), with error at most $x^{70} + 2fb\log N$. Since $\log N \ll m\log b$, the latter error is bounded within the budget. Combining the per-lift factorial estimates with this branching calculation gives the right side of (14.21), plus at most B of error. The exact inequality (14.22), including the conjugation uniformity and class-count normalization, is a formal-source input at the quantitative scale being used.

To see how the level N is chosen, retain the notation ℓ_{CS} for the Chowla–Selberg constant and put

$$\overline{w} = \frac{1}{f}\sum_{\tau} \log w(\tau), \quad G_0 = \log(g\,\mathrm{rd}_E/(2\pi)), \quad M_0 = a(\log a - 1 - \ell_{\mathrm{CS}} - \overline{w}).$$

Thus $\Omega = e^{\ell_{\mathrm{CS}}} 2\pi/\mathrm{rd}_E$, $\mathcal{R} = \sum_{\tau\in\mathcal{T}}(A(a,\tau) - M_0)$ and $J(a) = \log(a!) - a\log(\Omega/g)$. The elementary Stirling upper bound for $a \geq 1$ is

$$\log(a!) \leq a\log a - a + 1 + \tfrac{1}{2}\log a.$$

It follows from the monotonicity of the Stirling remainder with the constant fixed at $a = 1$, or by the standard Stirling bound with positive explicit error. Substituting in $J(a) - M_0$ gives

$$J(a) - M_0 \leq a(\overline{w} + G_0) + 1 + \tfrac{1}{2}\log a.$$

By construction of the level window

$$\log N \geq m(\overline{w} + G_0) - G_0 - \tfrac{15}{2}\log b.$$

Consequently

$$\begin{aligned} h_0 &= H - m\mathcal{R} + \#\mathcal{T}\{m(J(a) - M_0) - a\log N\} \\ &\leq H - m\mathcal{R} + \#\mathcal{T}\left\{m(1 + \tfrac{1}{2}\log a) + a(G_0 + \tfrac{15}{2}\log b)\right\} \\ &\leq H - m\mathcal{R} + B. \end{aligned} \tag{14.23}$$

The last estimate uses $\#\mathcal{T} \leq f$, $G_0 = O(\log b)$ and $m \leq 2\sqrt{x} \ll a$, with the large $\log b$ admissibility margin. This is precisely the calculation in `Data.heightPerRow_le` (`New/Endgame/ShortAndEnough.lean`). The level is chosen to remove the large contribution $m\#\mathcal{T}a(\overline{w} + G_0)$, leaving the prime remainder itself.

14.12 Projection, product formula scaling and Theorem A

The good projection e_0 on the finite level quotient G is normalized by $|G|$. Write $P_s = |G|e_0 sP(z_0\cdot)$ with z_0 from Theorem 14.4. The equality at the level of lifts is translation by finite compact elements followed by the same group-ring projection: the value of the lift of $|G|e_0 sz$ at g' is

$$\sum_{u\in G} (|G|e_0(u))\,(\Theta_\theta z)(g'\widetilde{us}).$$

This is `Data.iota_goodProjTranslate_apply` (`New/Rows/MonomialCoefficientVector.lean`). The coefficient $|G|e_0(u)$ is an algebraic integer: the group-ring central idempotents are projections onto specified irreducible types and their coefficients are characters multiplied by dimensions, divided by $|G|$. Finite group characters take algebraic-integer values. Thus the normalized projection preserves integrality at every finite place, including twists prime to that place. This yields `Data.goodProjTranslate_mem_integralForms`. For the trivial twist its complex norm contributes at most $e^{x^{99}}$ times the previous squared-value norm bound when converted to the squared class-sum norm. Indeed the absolute values of the coefficients of $|G|e_0$ are bounded in terms of the finite group order, and the number of classes is bounded as above. The statement with exact constant is `Data.normSqAt_goodProjTranslate_le`, same file.

Theorem 14.15 (A: short forms)**.** *There is a positive rational q such that, with $Q_s = qP_s$, the everywhere-integral columns $u \in C^{\mathrm{ram}}$ for which*

$$\|\Theta_1 Q_s u\|_\sigma^2 \le \exp(H - m\mathcal{R} + L) \qquad (s \in G,\ \sigma \in \mathrm{Aut}(\mathbb{C}))$$

span C^{ram}. For every finite $J \subset G$ and every finite family y of elements of the images $Q_s(C^{\mathrm{ram}}_{\mathrm{int}})$, $s \in J$, the logarithmic index of y in the twisted integral structures is at most

$$|J|cB.$$

Proof. Take the spanning set of (14.4). Put $h_p = \lfloor \epsilon_p - g_p \rfloor$ and $q = \prod_p p^{h_p}$, with the product finite by support of the loss and gain functions. In particular

$$\log q = \sum_p h_p \log p, qquad |\iota(q)|_p = p^{-h_p}.$$

For each p the integer $-h_p = \lceil g_p - \epsilon_p \rceil$ satisfies the strict hypothesis of Proposition 14.14. Therefore on M'_p the factor q times the restricted lift is integral for every admissible twist prime to p. The upper bound (14.22) with $\eta = B$ now gives

$$\begin{aligned} \log \|\Theta_1 Q_s u\|_\sigma^2 &\le 2\log q + h_0 + 2\sum_p g_p \log p + 2B + x^{99} \\ &\le h_0 + 2\sum_p \epsilon_p \log p + 2B + x^{99} \\ &\le H - m\mathcal{R} + 5B + x^{99} \le H - m\mathcal{R} + L. \end{aligned}$$

The last inequality uses the numerical hypothesis of the data, $L = B \log b$ and x^{99} negligible compared to B. A zero norm already satisfies the desired bound; the logarithmic calculation is applied to nonzero norms.

For the index, fix p and ι. On $M \cap C^{\mathrm{ram}}$ pull back the condition $I(M, M'_p; p^e)$ of Proposition 14.14 by inclusion; thus its determinant bound is unchanged. Under every particular Q_s the restricted submodule has integral lifted rows by the seesaw and the normalized projection. If M'' is the inverse image of the integral lifted rows under Θ_θ, this gives $I(M \cap C^{\mathrm{ram}}, Q_s^{-1}(M''); p^e)$. The last part of Lemma 14.2 therefore shows that any family from the specified union of $|J|$ images has a square coefficient matrix over O_ι turning its lifts into integral forms, with determinant dividing $p^{|J|e}$. In particular, its p-adic absolute value is at least $\exp(-|J|c\epsilon_p \log p)$. The numerical costs are independent of ι and of the twist, though the matrices need not be. This exactly meets the definition of logarithmic index in the row space: there is one function of p bounding the costs for every embedding and every twist prime to p. Finally (14.20) gives an upper bound $|J|cB$ for the sum of the costs over every finite set of primes. This proves the second part and completes the proof from the stated inputs. The source declaration is `Data.short_forms`, in `New/Endgame/ShortAndEnough.lean`. □

We emphasize which part of the mathematical reconstruction is still an input. The theorem's last deduction uses only the explicitly stated local divisibility, conjugate-size and index bounds, and their proofs as far as given above. The deepest local subtheorems are the CM jet normal form and Schur bound of the preceding section, the genuine seesaw with uniformly integral finite data, and the Frobenius-twisted Hecke divisor theorem `TwistingData.hasType_twisted_hecke`, with its root inequality and approximate projector. The presentation above specifies the actual versions needed, including the level and gauge compatibilities. They are formal-source inputs, not consequences merely of the usual theta correspondence or Satake parameter formulas.

Chapter 15

The local Hecke estimate in full detail, and the integral seesaw

The exceptional-prime correction of Chapter 14 is not an estimate for generic eigenvalues of a Hecke algebra. It is an estimate for the elementary divisors of a particular integral operator on the full column lattice, followed by an application to its image in ramified columns. The distinction matters: a small characteristic polynomial, or a bound for a normalized spherical function, need not give the required estimate on an integral image. In this chapter we reconstruct the main finite-dimensional theorem of `New/TwistedHecke/ElementaryDivisors.lean`, explaining in particular the integral root, flag-counting and approximate-projector arguments. We finish with some detailed geometric and integral computations needed at the extra level prime. The notation in the Hecke part is local: k is the column rank, n_0 the polynomial degree in each letter (equal to a in the application), r_0 the size of the first block, and $\nu = 2\lambda$ the length of the local Frobenius orbit. These should not be confused with the dimension $n = \dim W$, the global parameter r, or the auxiliary prime N.

15.1 The cyclic coefficient representation and the two filtrations

Let O be a valuation domain of characteristic zero, p a rational prime which is not a unit in O, and fix $w \in O$ with $w^2 = p$. Let R be a local ring with residue cardinality $q = P^\nu$ for $P = p^{d_0} \geq 5$. For each letter $s \in \mathbb{Z}/\nu$ there are ring maps $\iota_s : R \to O$ with

$$\iota_{s+1}(x) \equiv \iota_s(x)^P \pmod{p}.$$

Let a choice of alternating parity on $\mathbb{Z}/\nu$ be fixed. For $g \in \mathrm{GL}_k(R)$ define $g_s = \iota_s(g)$ in an even letter and $g_s = \iota_s(g^{-t})$ in an odd letter. We use the action ρ on the polynomial module generated by the monomials of degree n_0 in each block $X_{b,s}$, given by $gX_{b,s} = \sum_c (g_s)_{cb} X_{c,s}$. Its dimension is

$$M_0 = \binom{n_0 + k - 1}{k - 1}^\nu.$$

For a boundary between coordinates $r_0 - 1$ and r_0, let $\delta(b, s)$ be one on the deep variables (even s and $b \geq r_0$, or odd s and $b < r_0$), and zero otherwise; put

$$d(\alpha) = \sum_{b,s} \delta(b, s)\alpha_{b,s}, \qquad D_w = \mathrm{diag}(w^{d(\alpha)}).$$

The two-block Levi L consists of invertible block-diagonal matrices; let $\mathcal{P}$ be the inverse image of the lower block-triangular parabolic over the residue field. The formula for the group is precisely the convention used in `TwistingData.p_dvd_letterMatrix_of_mem_parahoric`, where an entry with block-colour $i < j$ is zero mod the maximal ideal (even letter).

For distinct indices b, b' write $y_s = X_{b',s}$, $z_s = X_{b,s}$ in the even letters, and $y_s = X_{b,s}$, $z_s = X_{b',s}$ in the odd letters. For a letter i define

$$E_i(\alpha; b, b') = \frac{1}{q}\left(\sum_s P^{1+[s-1-i]_\nu}\alpha(y_s) - \sum_s P^{[s-i]_\nu}\alpha(z_s)\right), \tag{15.1}$$

where $[s-i]_\nu$ is the residue in $[0, \nu-1]$. Write $E_i(v)$ for the weight of a variable v, so that $E_i(\alpha) = \sum_v E_i(v)\alpha(v)$. The exceptional summand at $s = i$ in the first sum has coefficient q, not 1. All weights $E_i(v)$ lie in $[-1/P, 1]$. This follows because the second exponent is at most $\nu - 1$ and the first is at most ν. If O is replaced by its residue field K_0, put U_h equal to the subspace whose every Levi translate has zero coefficients at monomials of excess above h for *some* i, using the boundary $b = r_0 - 1, b' = r_0$. Likewise V_j consists of vectors whose every full-group translate has zero coefficients at depth less than j. These definitions correspond to `excessCond` and `deepForms` of `TwistingData.lean`. We also use U_h mod w^e for the integral congruence condition in which forbidden coefficients must belong to (w^e) instead of being zero.

We will repeatedly use a Gauss filtration. If a variable v has real weight u_v, say a coefficient polynomial F is in filtration at least A if its coefficient of exponent α belongs to $(w^{\max(0,\lceil A - u\cdot\alpha\rceil)})$. This definition makes sense over an arbitrary commutative ring. Its groups are nested, and the product of elements of levels A, A' has level $A + A'$ because the product of their coefficient divisors is sufficient for the sum inequality. Two successive Gauss filtrations (first for a new variable T and then for the old variables) amount to using the additive array of weights of all variables. If two substitutions on polynomial variables each give weight at least u_v, and their difference on v has weight at least $u_v + \eta$, with $\eta > 0$, they induce filtration-preserving maps and their difference raises filtration by η. Indeed the difference on any monomial is a sum obtained by changing one factor at a time, and each summand contains a changed factor of the extra weight. Linearity gives the general statement. This proves the substitution lemma `aeval_sub_mem_of_gaussFiltration` used repeatedly in `RootLemma.lean`.

15.2 The root estimate, including its coefficient extraction

The short argument using multiplicity of zeros in Chapter 14 can be made completely explicit. It is crucial to use a sufficiently large weight separation, rather than directly asking a low-degree polynomial over the residue field to detect all the twists. Here is the complete substitution calculation in `TwistingData.root_lemma`.

Theorem 15.1 (Frobenius root lemma)**.** *Fix $b \neq b'$ in the same block of any Levi, a real weight c_v on all the variables and a coefficient $F_\alpha \in O$. Suppose that for every s*

$$\frac{2}{5} \leq c(y_s) - c(z_s) \leq \frac{8}{5}.$$

Suppose $l \geq 0$, $F_\alpha \notin (w^{l+1})$, and every translate of F by this Levi is in Gauss filtration of level $A = l + c\cdot\alpha$. Then for every letter i, $E_i(\alpha; b, b') \leq 0$.

Proof. Fix i and renumber cyclically so that its relative index is zero; set $t_s = [s-i]_\nu$. Thus $t_i = 0$. Write

$$\omega = \frac{c(y_i) - c(z_i)}{q}, \quad m(z_s) = qP^{t_s}, \quad e(y_s) = P^{1+[s-1-i]_\nu},$$

with $m = e = 0$ off their indicated variables. In particular $e(y_i) = q$. Let π collapse the pair y_i, z_i to z_i and leave every other variable fixed. Define a substitution χ with new variable T by

$$\chi(v) = \pi(v) T^{m(v)}.$$

For $s \neq i$ put $\epsilon_s = -1$ if the parity of s equals that of i, and $\epsilon_s = 1$ otherwise. Put $\epsilon_i = -1$. Define a second substitution ψ by

$$\begin{aligned}
&\psi(z_s) = z_s T^{qP^{t_s}} - \epsilon_s y_s T^{P^{t_s}}, && s \neq i,\\
&\psi(y_s) = y_s - \epsilon_s z_s, && s \neq i,\\
&\psi(z_i) = z_i T^q, && \psi(y_i) = z_i,
\end{aligned}$$

and $\psi = \chi$ on all other variables.

First give T weight $-\omega$ and give the old variables weight $c(v) + \omega m(v)$. The weight of $\chi(v)$ is exactly $c(v)$, since the two weights at the collapsed pair coincide after the adjustment. Each correction term in $\psi(v) - \chi(v)$ has weight at least $c(v) + 1/20$: for $v = z_s$, $s \neq i$, the gain is

$$c(y_s) - c(z_s) - \omega P^{t_s} \geq \frac{2}{5} - \frac{8}{5P} \geq \frac{2}{25} > \frac{1}{20},$$

and for $v = y_s$, $s \neq i$, its gain is

$$c(z_s) + \omega q P^{t_s} - c(y_s) \geq \frac{2P}{5} - \frac{8}{5} \geq \frac{2}{5}.$$

Consequently $\chi(F)$ and $\psi(F)$ have filtration at least A in these weights, and their difference at least $A + 1/20$. Define a scalar-coefficient polynomial $\Phi(T)$ by extracting the coefficient of the old monomial $X^{\pi_* \alpha}$ from $\psi(F)$ and reducing it modulo (w^{l+1}). Put $M = m \cdot \alpha$. Both terms of filtration A at this old exponent have T^j coefficient divisible by w^{l+1} whenever $j > M$, since its required w-order is $l + \omega(j - M) > l$. At $j = M$ the extra filtration of the difference means that its coefficient is also zero mod w^{l+1}. In $\chi(F)$ the old exponents determine all exponents of F except the split between z_i, y_i; then the value of M determines that split uniquely, because changing the exponent of z_i by one changes M by q. The coefficient of T^M is therefore $F_\alpha \bmod w^{l+1} \neq 0$. We have proved

$$\Phi \neq 0, \qquad \deg \Phi \leq M. \tag{15.2}$$

Next give T weight $+\omega$ and the old variables weight $c(v) - \omega e(v)$. Again the adjusted weights of the two collapsed variables agree. For $x \in R$ use the Levi element which first makes the transvection on the root corresponding to (b, b'), with parameter chosen as $-x$ when i is even and x when i is odd, and then interchanges those two basis vectors. For this element g_x the substitution in the sth letter sends $z_s \mapsto y_s$ and $y_s \mapsto z_s + \epsilon_s \iota_s(x) y_s$. This follows by writing the two-by-two transvection matrix: on odd letters its inverse transpose reverses the sign. Compose its action with ψ and the translation $T \mapsto T + \iota_i(x)$.

We verify filtration preservation with respect to these second weights. At i the image of z_i is z_i, and that of y_i is

$$z_i((T + \iota_i(x))^q - \iota_i(x)).$$

Modulo p the expression in parentheses is T^q, by the Frobenius congruence iterated around the full cycle. The main term has weight $c(y_i)$, and the error (coefficients divisible by $p = w^2$)

has weight at least $c(z_i)+2 \geq c(y_i)$. For $s \neq i$, the image of z_s is $y_s - \epsilon_s z_s$ and each term has weight at least $c(z_s)$, by the first inequality in the preceding calculation. The image of y_s is congruent modulo p to

$$z_s T^{qP^{t_s}} - \epsilon_s y_s T^{P^{t_s}}.$$

Indeed $(T+\iota_i(x))^{P^{t_s}} \equiv T^{P^{t_s}} + \iota_s(x)$ and $(T+\iota_i(x))^{qP^{t_s}} \equiv T^{qP^{t_s}} + \iota_s(x)$ modulo p. The constant terms cancel against the terms from the transvection. The first displayed main term has weight $c(z_s)+\omega q P^{t_s} \geq c(y_s)$, the second has weight exactly $c(y_s)$. The error has weight at least the smaller of $c(z_s)+2$ and $c(y_s)-\omega P^{t_s}+2$, so it too has weight at least $c(y_s)$. All other variables are unchanged. This proves the required filtration preservation without any assumption that the valuations are discrete.

For the desired Taylor expansion of $\psi(F)$, apply the proved filtration-preserving composite $(T \mapsto T+\iota_i(x)) \circ \psi \circ g_x$ to $g_x^{-1}F$, which satisfies level A by hypothesis. Its resulting polynomial is exactly the translate $\psi(F)(T+\iota_i(x))$. At the selected old monomial, the coefficient of T^j now has w-order at least

$$l + \omega(e \cdot \alpha - j).$$

For $j < e \cdot \alpha$ this is strictly greater than l. Thus the polynomial Φ of (15.2) has Taylor coefficients zero in degrees below $E = e \cdot \alpha$ at $\iota_i(x)$ for every $x \in R$.

Choose lifts of all q residue classes of R. Their pairwise differences are units in R, so their images remain units in $O/(w^{l+1})$. The polynomials $(T-\iota_i(x))^E$ are pairwise comaximal, each divides the nonzero polynomial Φ, and their monic product has degree qE. Hence $qE \leq M$ (multiplication by a monic polynomial adds its degree even over a ring with zero divisors). Substitution of the definitions of m, e gives exactly (15.1) with nonpositive sign. This proves the theorem. □

The proof only uses ring identities, the specified congruences, and quotient rings by powers of w. In particular it also applies after reduction to a field of characteristic p, with $w = 0$, $l = 0$ and the convention $w^0 = 1$. The proof illustrates why the small adic precision $p = w^2$ is sufficient: its weight 2 is strictly larger than every possible loss in the variable weights. The root lemma is a simultaneous condition on all possible least monomials in all frames. It does not say that all coefficients of an arbitrary polynomial already obey the excess inequalities.

Example 15.2. Take two letters, one even and one odd, and the boundary between coordinates 0 and 1. The two inequalities at a least monomial include

$$q\alpha(y_0) + P\alpha(y_1) \leq \alpha(z_0) + P\alpha(z_1), \qquad q\alpha(y_1) + P\alpha(y_0) \leq \alpha(z_1) + P\alpha(z_0).$$

Even if $\alpha(y_0)$ is small relative to the total degree in the even letter, it may violate the first inequality. The cyclic coefficients are a reflection of the two different embeddings of the same residue field: treating the letters as independent finite fields would lose precisely these constraints.

15.3 The dimension bound from flags and convex volume

We now calculate the dimension of U_h over a field K_0 of characteristic p with the indicated reduction of the twisting data. Because the critical coefficient is asymptotic to the reciprocal of the degree of the local Hecke correspondence, it is worth keeping the leading volume constant.

For $h \geq 0$, let $\mathscr{C}_h$ be the cone in the exponent space $\mathbb{R}^{k\nu}$ defined by the excess inequalities for adjacent coordinates:

$$E_i(\beta; b, b+1) \leq 0 \ (b+1 \neq r_0), \qquad E_i(\beta; r_0-1, r_0) \leq h$$

for every letter i. Its intersection with the nonnegative orthant and with the sets where the coordinate sums are in bounded intervals is a compact convex polytope. Write $\Sigma(D)$ for the product of slabs $\beta_{bs} \geq 0$, $D_s - 1 \leq \sum_b \beta_{bs} \leq D_s$. Let I denote the Iwahori subgroup (lower triangular reduction) of $\mathrm{GL}_k(R)$. Write $F_L = [L : L \cap I]$ for the number of full flags in the two Levi blocks.

Lemma 15.3 (Amplification by a full Frobenius period)**.**

$$\dim U_h \leq F_L \operatorname{vol}(\mathscr{C}_{h+\nu(k+1)} \cap \Sigma(n_0 + k + 1)).$$

Here the endpoint in the slab is constant in every letter and the measure is ordinary coordinate volume.

Proof. Choose representatives g_j for $(L \cap I)\backslash L$ including the identity. Take $M = q^u$ with u any positive integer. For each array $v = (v_{bs})$ with $0 \leq v_{bs} < M$, define $\gamma(v)$ by adding

$$M\left(k - \left\lfloor \frac{\sum_b v_{bs}}{M} \right\rfloor\right)$$

to the coordinate $b = r_0 - 1$ in each letter s and leaving the other coordinates unchanged. The exponent $\gamma(v)$ is congruent to v modulo M coordinatewise, and its degree in each letter lies in $[kM, (k+1)M)$. Make the following map on $M^{k\nu}$ copies of U_h:

$$(F_v)_v \longmapsto F = \sum_v X^{\gamma(v)} F_v(X^M).$$

It is injective, since exponents for different v are distinct modulo M, while for a fixed v distinct monomials remain distinct. Raising coefficients in any letter action matrix to the Mth power leaves them unchanged: M is a power of the cardinality of the original residue field and the reduced actions have entries in its image. By the Frobenius identity, linear substitution in $F_v(X^M)$ therefore commutes with the letter action. This is the necessity of choosing a *full* Frobenius period, not just $M = p^u$ of arbitrary exponent.

Suppose $F \neq 0$. Among all nonzero terms of all $g_j F$, choose one exponent α with least weight relative to the array $c(X_{b,s}) = b$ in an even letter and $-b$ in an odd letter. Every translate by an element of L has all terms of weight at least $c \cdot \alpha$: such a translate is hg_j with $h \in L \cap I$. The action of h cannot decrease this weight. Indeed in an even letter all nonzero matrix entries in the residue field have row index at least column index, and in an odd letter the reverse inequality holds after inverse transpose. Thus the chosen coefficient meets the hypotheses of the root lemma over K_0, with $w = 0, l = 0$ and with the Levi L for any adjacent pair lying in the same block. For each such pair the weight separation is exactly one. All its excess inequalities are consequently nonpositive.

For the boundary pair, any nonzero coefficient of any $g_j F$ comes from an exponent of the form $M\beta + \eta$, where β is in the support of a Levi translate of some F_v, and η has degree less than $(k+1)M$ per letter. This follows by expanding each summand of the substituted polynomial; nonzero coefficient means at least one such term contributes. Thus $E_i(\beta) \leq h$ and $E_i(\eta) \leq \nu(k+1)M$ because all variable excess weights are at most one. Hence $E_i(\alpha/M) \leq h + \nu(k+1)$ at the boundary. The degree of α/M is in $[n_0 + k, n_0 + k + 1]$ in each letter. It follows that it belongs to the polytope in the lemma.

Therefore the coefficient restrictions at the lattice points α/M of this polytope of all the finitely many $g_j F$ give an injective linear map. If their values all vanished, the above least nonzero coefficient would yield a contradiction. Its target has dimension at most F_L times the number of lattice points in the polytope of mesh $1/M$. Comparing dimensions gives

$$M^{k\nu} \dim U_h \leq F_L \#((M^{-1}\mathbb{Z}^{k\nu}) \cap \mathscr{C}_{h+\nu(k+1)} \cap \Sigma(n_0 + k + 1)).$$

Divide by $M^{k\nu}$ and let $u \to \infty$. The lattice-point Riemann sums converge to volume: the set is bounded and its boundary is contained in finitely many affine hyperplanes, unless it lies in such a hyperplane, in which case both the limiting sums and the volume are zero. This proves the bound. □

This is the proof of `TwistingData.finrank_excessCond_le_volume`, `New/TwistedHecke/Counting.lean`. It is a pure dimension argument and does not require the irreducibility of the symmetric-power representations.

The exact volume in the basic cone can also be calculated without an asymptotic estimate. In the following calculation we use the sign $\epsilon(s) = 1$ on even letters, -1 otherwise. Introduce an invertible linear change

$$\beta_{b,s} = \epsilon(s)(Pv_{b,s+1} - v_{b,s}).$$

The determinant is nonzero because in the kernel the equations cycle to $(P^\nu - 1)v_{b,s} = 0$. A direct telescoping gives

$$E_i(\beta; b, b') = (1 - q^{-1})(\beta(y_i) - (v_{b,i} - v_{b',i})). \tag{15.3}$$

For clarity, set $i = 0$. The excess can be written as $(1-q^{-1})\beta(y_i)+q^{-1}\sum_s P^{[s-i]_\nu}(\beta(y_s)-\beta(z_s))$. Now $\beta(y_s) - \beta(z_s) = P(v_{b',s+1} - v_{b,s+1}) - (v_{b',s} - v_{b,s})$, in either parity. The cyclic sum is $(q-1)(v_{b',i} - v_{b,i})$. This proves the identity. Nonnegative β satisfying $\mathscr{C}_0$ correspond to

$$\begin{cases} v_{b,s} \le Pv_{b,s+1}, & Pv_{b+1,s+1} \le v_{b,s}, \quad s \text{ even}, \\ Pv_{b,s+1} \le v_{b,s}, & v_{b+1,s} \le Pv_{b,s+1}, \quad s \text{ odd}. \end{cases}$$

Away from the equality hyperplanes this forces all vectors v_s to be decreasing in b. In particular the signs of v_s do not enter the counting, since only the adjacent comparisons do.

Lemma 15.4 (Cyclic interlacing volume)**.** *For all endpoints D_s,*

$$\operatorname{vol}(\mathscr{C}_0 \cap \Sigma(D)) = \frac{\operatorname{vol}\Sigma(D)}{\prod_{j=1}^{k}(q^j - 1)/(q-1)}.$$

Proof. Ignore hyperplanes of equalities (measure zero). For decreasing vectors x, y let $C(x, y)$ be the matrix whose (b, c) entry is $\mathbf{1}(y_c < x_b)$. Its rows have nested supports in the final segments of the coordinate set. Its determinant is one for strict interlacing $x_1 > y_1 > x_2 > \cdots > x_k > y_k$, and zero otherwise. Indeed all rows must be distinct and nonzero, so their lengths must be exactly $k, k-1, \ldots, 1$. This gives the claimed determinant and condition.

Apply this identity to (Pv_{s+1}, v_s) in an even letter and (v_s, Pv_{s+1}) in an odd letter, transposing the former matrix if needed to give indices at s and $s+1$ consistently. The product of these determinants on sorted vectors is the indicator of the interlacing cone with nonnegative β. It is invariant under permutation of the coordinates at any single letter, because that permutation affects two consecutive determinants and their signs cancel. We may integrate over all vectors, subject only to the condition on the coordinate sums, and divide by $(k!)^\nu$. On expanding the determinants, each permutation tuple $(\sigma_s)_s$ gives the signed indicator of positivity in the linear transformation

$$(L_\sigma v)_{b,s} = \epsilon(s)(Pv_{b,s+1} - v_{\sigma_s(b),s}).$$

The sums of coordinates of $L_\sigma v$ in a letter are independent of σ, so the integral of the indicator in the sum equals $\operatorname{vol}\Sigma(D)/|\det L_\sigma|$. To calculate the determinant, put A_s for the

permutation matrix associated to σ_s, in the convention $A_s v_s = (v_{\sigma_s(b),s})_b$. Ignore the sign factors, which are units. The equations give $v_{s+1} = P^{-1}(A_s v_s + \beta_s)$; successive elimination of $v_1, \dots, v_{\nu-1}$ contributes powers of P whose product is $P^{k(\nu-1)}$. The remaining block for v_0 is $PI - P^{-(\nu-1)} A_{\nu-1} \cdots A_0$; multiplying its determinant by those factors gives precisely $|\det(qI - A_\pi)|$, where π is the product permutation. Replacing the v-coordinates by the original β-coordinates multiplies the result by $|\det L_1| = (q-1)^k$. For any fixed π there are $(k!)^{\nu-1}$ permutation tuples of product π and the product of their signs equals $\operatorname{sgn}\pi$. The volume ratio is thus

$$\frac{(q-1)^k}{k!} \sum_{\pi \in S_k} \frac{\operatorname{sgn}\pi}{|\det(qI - A_\pi)|}.$$

This expression depends only on q and k, not on the number of letters or their parities. Evaluate it by the same one-letter determinant formula with parameter q (the all-positive parity in that one letter is legitimate for this measure calculation). Then the cone is $\beta_b \geq q\beta_{b+1}$, $\beta_k = 0$ for numbering $b = 0, \dots, k-1$. Let $d_b = (q^{b+1} - 1)/(q-1)$. The triangular linear map $\beta \mapsto (d_b(\beta_b - q\beta_{b+1}))_b$ carries that cone onto the nonnegative orthant, preserves coordinate sum, and has determinant $\prod_b d_b$. Hence the volume ratio is its reciprocal, giving the formula. □

This proves the determinant-volume identities in `New/TwistedHecke/InterlacingCone.lean`. For example, if $k = 2$ the volume fraction $1/(q+1)$ is exactly the reciprocal of the number of lines over $\mathbb{F}_q$. For higher k the denominator is the number of complete flags, obtained recursively by choosing a line in the first quotient and then a complete flag of the quotient space.

We now need the thickened cone with nonzero allowance at the boundary. Define $u_b = 2P^{2(r_0-1-b)}$ if $b < r_0$, and $u_b = 0$ otherwise. Use $v_{b,s} = u_b$ on even letters and $v_{b,s} = Pu_b$ on odd letters. Then $(L_1 v)_{b,s}$ equals $(P^2 - 1)u_b$ in even letters and zero in odd letters. Call this vector σ. It has nonnegative coordinates bounded by P^{3r_0} (indeed $2P^{2r_0} \leq P^{3r_0}$). Since

$$u_b = P^2 u_{b+1} + 2\mathbf{1}(b+1 = r_0),$$

substitution in (15.3) shows $\sigma \in \mathscr{C}_{-1}$. At an even letter the inner expression is $P^2 u_{b+1} - u_b$, giving -2 on the boundary, zero elsewhere. At an odd letter it is $-P(u_b - u_{b+1})$, which is nonpositive and at most $-2P$ on the boundary. The factor $1 - q^{-1}$ leaves the boundary bound at most -1. This is the explicit vector of `exists_mem_excessCone_mem_Icc`.

Put $h' = h + \nu(k+1)$. Translating every exponent in the polytope of Lemma 15.3 by $h'\sigma$ embeds it in the zero-excess cone, while the endpoint of the slab in letter s increases by at most $h'kP^{3r_0}$. Suppose $\varepsilon \leq 1$ and the size assumptions (14.13) hold (with $a = n_0, y = h$). Then the new endpoints are at most

$$n_0 + k + 1 + h'kP^{3r_0} \leq n_0(1 + 1/h).$$

The volume of a nonnegative slab with endpoint D in dimension k is $(D^k - (\max(D - 1, 0))^k)/k! \leq D^{k-1}/(k-1)!$. Together the shifted slabs have volume at most

$$\left(\frac{n_0^{k-1}}{(k-1)!}\right)^{\nu} e^{\nu(k-1)/h} \leq (1+\varepsilon)M_0,$$

since $n_0^{k-1}/(k-1)! \leq \binom{n_0+k-1}{k-1}$, $\nu(k-1)/h \leq \varepsilon/2$, and $e^{v/2} \leq 1 + v$ for $0 \leq v \leq 1$. Finally $[\mathrm{GL}_k(R) : \mathcal{P}]F_L$ is the number of complete flags: reduction identifies the cosets of $\mathcal{P}$ with subspaces of dimension the appropriate block size, and a full flag refining the fixed block is a pair of flags inside the two blocks (the unipotent part lies in I). Lemmas 15.3 and 15.4 therefore give the needed bound

$$[\mathrm{GL}_k(R) : \mathcal{P}] \dim U_h \leq (1+\varepsilon)M_0. \tag{15.4}$$

This is the quantitative count `index_mul_finrank_excessCond_le` with exactly the needed leading constant.

15.4 Deep coefficients: a weighted Nullstellensatz and its application

The volume argument controls the dimension of U_h. The other loss in the Hecke theorem is the cumulative dimension of the depth spaces V_j. To obtain it one needs a genuinely different argument: the obstruction is high-order vanishing at a large family of twisted unipotent points. We prove the multiplicity-weighted version of the Nullstellensatz first.

Lemma 15.5 (Weighted vanishing)**.** *Let K be a field, $u_1, \dots, u_l$ independent variables, and $\kappa_1, \dots, \kappa_l$ positive integers. Give u_i cost $1/\kappa_i$. Let $Z \subset K^l$ be a finite set of at least $\prod_i \kappa_i$ distinct points. For every i suppose there is a polynomial g_i of cost strictly below one agreeing with $u_i^{\kappa_i}$ at all points of Z. If a nonzero polynomial F vanishes to order at least d at all points of Z, then some monomial of F has cost at least d.*

Proof. The assertion is clear for $d = 0$. Set $h_i = u_i^{\kappa_i} - g_i$, which vanish on Z. The polynomials

$$u^\alpha \prod_i h_i^{\beta_i}, \qquad 0 \le \alpha_i < \kappa_i,$$

have distinct initial monomials $u^{\alpha+\kappa\beta}$ of cost $\sum \alpha_i/\kappa_i + |\beta|$; all remaining terms have lower cost. They form a basis of the polynomial ring over K: independence follows by taking largest cost (and distinct initial terms at that cost), and spanning follows by subtracting the appropriate basis elements from the terms of maximal cost. This terminates because the cost is discrete, being a rational number with bounded denominator. In particular the polynomials of cost below d lie in the span of basis elements with $|\beta| < d$.

Consider the quotient of the polynomial ring by the intersection of the dth powers of the maximal ideals of the points in Z. These ideals are pairwise comaximal, so by the Chinese remainder theorem the quotient has dimension $|Z|\binom{d+l-1}{l}$. Since all basis elements with $|\beta| \ge d$ vanish to the indicated order, the quotient is spanned by the images of the $(\prod_i \kappa_i)\binom{d+l-1}{l}$ remaining basis elements. The cardinality hypothesis implies equality and makes their images independent. A nonzero polynomial of cost below d cannot belong to the intersection. This is the desired statement. □

The argument is the ordinary division proof of `exists_le_cost_of_vanishing`, in `Lib/General/Algebra/ZerosCostDegree.lean`. It also shows why merely using the degree of a polynomial on a finite grid would not suffice: certain variables have extremely different costs, and the grid is not a product in those coordinates.

Proposition 15.6 (Everywhere deep forms have a costly coefficient)**.** *Suppose $2r_0 \le k$ and $2r_0 < \nu$. Give each deep variable of index $b < 2r_0$ cost P^{-1}, each deep variable of index $b \ge 2r_0$ cost P^{-2r_0}, and all other variables cost zero. Every nonzero member of V_j has a nonzero monomial of cost at least j.*

Proof. Put $k = 2r_0 + h_0$. We use upper unipotent matrices $1 + A$ with entries A_{ab} only in the following positions: the r_0 paired positions $(a, r_0 + a)$, $0 \le a < r_0$, and all $r_0 h_0$ positions (a, b) with $b \ge 2r_0$. In particular $A^2 = 0$, and $1 + A$ is invertible. Index the positions by (a, c), $0 \le a < r_0$, $0 \le c \le h_0$, where the second coordinate of the matrix entry is $r_0 + a$ for $c = 0$ and $k - c$ otherwise. Denote the resulting pair of matrix indices by $\rho(t), \sigma(t)$ for a position t.

Take each entry $A_{\rho(t),\sigma(t)}$ to be a lift of a residue parameter $u_t \in \mathbb{F}_q$ and write $a_s(t) = \bar{\iota}_s(u_t)$ for its residue in K_0. The Frobenius congruences give $a_{s+e}(t) = a_s(t)^{P^e}$.

Let K be the field of rational functions in the shallow variables over K_0 (one may harmlessly adjoin independent indeterminates also for the original deep variables). In the new polynomial algebra over K, keep only the deep variables as variables and regard the others as scalar coefficients. Define a point $z(u)$ by

$$z(u)_{b,s} = \sum_{\sigma(t)=b} a_s(t) X_{\rho(t),s}, \qquad s \text{ even},\ b \geq r_0,$$

$$z(u)_{b,s} = -\sum_{\rho(t)=b} a_s(t) X_{\sigma(t),s}, \qquad s \text{ odd},\ b < r_0.$$

Thus $\rho(1-A)$ sends a deep variable to the corresponding variable minus its coordinate of $z(u)$, and fixes shallow variables. This follows directly from g_s being the ordinary matrix in even letters and the inverse transpose in odd ones. Since $\rho(1+A)F$ has depth at least j, writing $F = \rho(1-A)\rho(1+A)F$ shows that $F \in \mathfrak{m}^j_{z(u)}$ for every u.

We claim that the coordinates satisfy the hypotheses of Lemma 15.5 with $\kappa_{b,s} = P$ for the deep variables $b < 2r_0$ and $\kappa_{b,s} = P^{2r_0}$ for the other deep variables. To prove this, it suffices to recover every $a_i(t)$ as the value at $z(u)$ of a polynomial of cost less than one, uniformly in u. For fixed i and each paired position $(a, 0)$, take the coordinate of z in letter i corresponding to the deep root variable of $(a, r_0 + a)$, with exponent one. For the other position (a, c) choose $j_t \in \{2a, 2a+1\}$ such that the letter $i - j_t$ is even, and use the coordinate $z^{P^{j_t}}_{k-c,i-j_t}$. Its cost is $P^{j_t - 2r_0} < 1$, since $j_t \leq 2r_0 - 1 < \nu$. The paired coordinates also cost less than one.

At the points, these expressions are linear combinations of the $a_i(t')$ by the Frobenius identities. Their coefficient matrix over the rational function field is invertible. Here is a direct check: specialize each shallow variable $X_{b,s}$ to one if $b \geq \lfloor [i-s]_\nu/2 \rfloor$, and to zero otherwise. For a row indexed by (a, c) with $c > 0$ its only nonzero coefficients correspond to (a', c) with $a' \geq a$, since $\lfloor j_t/2 \rfloor = a$; its diagonal coefficient is one. For a row $(a, 0)$ in an even i only the corresponding paired matrix entry contributes, with coefficient one. For odd i only entries of row a contribute (the paired position and the far columns), with sign minus one, and its diagonal coefficient is minus one. In the lexicographic order on (a, c) the specialized matrix is upper triangular with unit diagonal (up to signs), and hence its determinant is nonzero. This proves the invertibility claim before specialization.

The claim now follows: after raising any coordinate of $z(u)$ to the indicated $\kappa_{b,s}$ and applying Frobenius, the result is a linear combination of residue parameters $a_{s+1}(t)$ or $a_{s+2r_0}(t)$ over the rational function field. Replace each by its polynomial expression of cost below one. The points are distinct, because recovering $a_i(t)$ from the coordinates recovers u_t (the map on residue fields is injective). Their cardinality is $q^{r_0(h_0+1)}$. The product of the costs' denominators is exactly this: at each letter there are r_0 deep variables among the first $2r_0$ coordinates, giving $P^{\nu r_0}$; the h_0 far coordinates are deep in half the letters, giving $P^{2r_0 h_0 \nu/2}$. This proves the required cardinality. Lemma 15.5 applies to the polynomial in the deep variables over the rational function field. Its high-cost coefficient must come from at least one monomial of the original F with the same deep exponents. The shallow variables have zero cost. This proves the proposition. □

This reconstructs `exists_le_weight_of_mem_deepForms` in `New/TwistedHecke/DeepFormCost.lean`, including the recovery matrix in the proof. The triangular specialization is essential: there cannot be a separate parameter for every deep variable because that would eliminate the gain P^{-2r_0} in the estimate.

The restriction to costly coefficients is injective on V_j: a nonzero vector in its kernel would contradict Proposition 15.6. Each monomial is counted at most 1 plus its cost when one sums

these dimensions over $j \ge 0$. The mean exponent of any specified variable among degree-n_0 monomials in a letter is n_0/k, by coordinate symmetry. Hence the mean cost is

$$\frac{n_0\nu}{2k}\left(\frac{2r_0}{P}+\frac{k-2r_0}{P^{2r_0}}\right) \le \nu n_0\left(P^{-2r_0}+\frac{r_0^2}{kP}\right) \le z$$

under (14.13). The same size assumptions imply $2r_0 < \nu$: indeed $P^{3r_0} \le n_0 < P^{\lambda}$, hence $3r_0 < \lambda$. We obtain, for every J,

$$\sum_{0\le j<J} \dim V_j \le (1+z)M_0. \tag{15.5}$$

Thus both dimension bounds needed in the Hecke matrix theorem are consequences of explicitly proved algebraic estimates. We have not appealed to a semisimplicity statement in characteristic p.

15.5 Integral diagonal pairings and class means

Although the dimension bounds were proved in characteristic p, our application concerns an integral matrix. It is not enough to assert that a desired projector exists after reduction mod p. We need an integral approximation to it, with an explicit scalar normalization. The source constructs this approximation from invariant linear functionals on polynomials in a generic matrix. We spell out their construction.

For a finite set of coordinates S and monomial exponent α of total degree $a_0 < p^F$, the multinomial $m_\alpha = a_0!/\prod_i \alpha_i!$ divides $p^{|S|F}$ in O. Indeed the part of the integer prime to p is a unit in the local valuation ring. Legendre's formula expresses the exponent of p in m_α as

$$\sum_{j\ge 1}\left(\left\lfloor\frac{a_0}{p^j}\right\rfloor-\sum_i\left\lfloor\frac{\alpha_i}{p^j}\right\rfloor\right).$$

Terms for $j \ge F$ vanish, and each other term is at most $|S|-1$, by the floor inequality. Therefore $q_\alpha = p^{|S|F}/m_\alpha$ belongs to O. These numbers give the reciprocal-multinomial diagonal inner product for the symmetric power action. More precisely for any $S \times S$ matrix G over O, let L_G act on polynomials by substituting $X_b \mapsto \sum_c G_{cb}X_c$. The Fischer pairing is diagonal with $\prod_i \alpha_i!$ on X^α. Expanding powers of linear forms (or checking the identity by the binomial formula) gives

$$\langle L_G X^\beta, X^\alpha\rangle = \langle X^\beta, L_{G^t}X^\alpha\rangle.$$

Multiplying by $p^{|S|F}/a_0!$ in the fraction field and using the integral values yields

$$q_\alpha[X^\alpha]L_G(X^\beta) = q_\beta[X^\beta]L_{G^t}(X^\alpha). \tag{15.6}$$

This is `ClassMean.exists_invariant_diagonal_form`. In particular it holds integrally for noninvertible G too.

We also need the elementary interpolation span over the residue field. Let d_1 be a homogeneous degree and Δ a nonzero homogeneous polynomial of degree d_2 over R with a unit value. Suppose $d_1 + d_2 < q$. Then the evaluations on points where Δ is a unit span the full dual of the homogeneous polynomial module of degree d_1 over O. Indeed a vector in the kernel of all those evaluations over the residue field gives a polynomial F for which $F\bar{\Delta}$ vanishes at every $\mathbb{F}_q$-point. Applying the one-variable root bound in each variable successively makes this polynomial zero, because its degree in each variable is below q. As $\bar{\Delta}$ is not

identically zero, $F = 0$. This argument also works over a residue coefficient extension, by the same root bound with points in the embedded residue subfield. Therefore a finite subset of the evaluation vectors has invertible determinant mod the maximal ideal and lifts to a basis over O. We will only apply this strict inequality case of `ClassMean.eq_top_of_eval_monomial`; the source includes equality in total degree for its homogeneous version.

For the next construction fix the generic $k \times k$ matrix X, a real weight array x_b on its indices, and a subset $T \subseteq \mathbb{R}$ (not necessarily an interval). For a monomial in matrix entries let its row weight be the sum of weights x_b of its row indices counted with multiplicities, and column weight defined similarly. Put $C_g F(X) = F(\iota(g)^{-1} X \iota(g))$ for g in the Levi. We use C_g for this algebra automorphism, not for the column space.

Proposition 15.7 (Weighted conjugation mean)**.** *Suppose $D < p^F$, $D + k < q$. There is an integral linear form Λ_T on the polynomial ring in X with properties:*

1. *It is invariant under C_g for all $g \in L$.*
2. *Its value on monomials of different row and column weights is zero, and its value on a monomial X^α with row weight in T is*
$$\Lambda_T(X^\alpha) = \begin{cases} p^{k^2 F} X^\alpha(1), & |\alpha| \le D, \\ 0, & |\alpha| > D. \end{cases}$$
3. *It belongs to the O-span of forms $\ell \circ C_g$, where ℓ vanishes on monomials whose row weight does not belong to T.*

Proof. The problem splits by homogeneous degree; construct the form in one degree $a_0 \le D$ and extend by zero on higher degrees. Let $U = (O^k)^{\otimes a_0}$ with its tensor basis $e_{t_1,\dots,t_{a_0}}$. Denote by $Y(X)$ the block-diagonal part of X^t (transpose included in this definition). For any homogeneous dual functional ℓ define $S(\ell) : U \to U$ by applying ℓ to each entry of $Y(X)^{\otimes a_0}$. This is an O-linear map in ℓ. The evaluation vectors at block-invertible matrices span the dual by the interpolation argument with polynomial $\Delta = \det(Y(X))$ of degree k, which has unit value at the identity. Therefore the image of S is the O-linear span of the tensor powers of the transposes of matrices in the Levi. Write this image as $\mathscr{A}$; it is an algebra, since the product of two such matrix evaluations is another such evaluation.

Let U_T be the $\mathscr{A}$-submodule generated by the tensor basis vectors e_t for which $\sum_j x_{t_j} \in T$. It is a finitely generated torsion-free O-module, hence finite free. (One proof of the standard valuation-ring fact is pivot elimination on finitely many generators of a submodule of a free module: an entry dividing the other finitely many entries in its row can be split off and induction applies.) The action of $\mathscr{A}$ on U_T consequently has a trace. For each exponent α of degree a_0 let $K(X^\alpha) = q_\alpha[X^\alpha]$ be a functional on the homogeneous polynomials, with q_α as in (15.6) for $|S| = k^2$. Define

$$\Lambda_T(F) = \mathrm{Tr}_{U_T}(S(K(F))|_{U_T}).$$

Equation (15.6), applied to the matrix substitution under conjugation, gives covariance of $S(K(F))$ under the tensor action of a Levi element. With $C_g(F)(X) = F(g^{-1}Xg)$ the transpose of the substitution on entry variables is $X \mapsto g^{-t} X g^t$. On the transposed block matrix $Y(X)$ this induces $Y \mapsto gYg^{-1}$. Thus $S(K(C_g F)) = S(\mathrm{ev}_{g^t}) S(K(F)) S(\mathrm{ev}_{g^{-t}})$, with ι implicit in the matrix entries. Since the Levi is transpose-stable, this makes Λ_T invariant by cyclicity of trace.

Suppose the variable monomial has unequal row and column weights. The matrix $S(K(X^\alpha))$ on U sends a basis vector of weight ω to one of weight $\omega + \delta$, where δ is the column minus row

weight of α. Here $\delta \neq 0$. Since the set of tensor weights is finite, sufficiently large positive powers of this matrix are zero. Its restriction to the stable U_T is nilpotent, hence has zero trace (check over the fraction field). This proves the first part of property 2. If instead the row weight lies in T, the matrix annihilates all e_t not in the indicated generating set and has image in U_T. Its trace on U_T equals its trace on all U; again this may be checked after passing to the fraction field and choosing a complement to U_T. The trace of the universal tensor matrix is $(\sum_b X_{bb})^{a_0}$, so extracting the coefficient of X^α and multiplying by q_α gives $p^{k^2F} X^\alpha(1)$. This gives the remaining equality in property 2.

To prove the span assertion, choose a basis (z_j) of U_T and let λ_j be its dual. Express each z_j as a finite sum of terms $cS(\mathrm{ev}_g)e_t$ with e_t of allowed weight, possible by the definition of U_T and the evaluation-span assertion. The trace is a sum of expressions

$$c\lambda_j(S(K(F))S(\mathrm{ev}_g)e_t).$$

Move $S(\mathrm{ev}_g)$ to the left by the covariance just proved. Each term is now of the form $\ell(C_{g'}F)$ for an integral functional ℓ, where $g' = g$ or g^{-1} as appropriate; ℓ vanishes on forbidden row weights because $S(K(X^\alpha))e_t = 0$ there. This proves property 3. Summing the finitely many homogeneous constructions completes the proof. □

This is the trace construction of `ClassMean.exists_mean`, `New/TwistedHecke/ClassMeans.lean`. The name "mean" need not indicate averaging with an invertible group order. The bounded power of p replaces division by the order of a large finite group, and the trace construction works in the integral setting. Replacing it by the ordinary group average would destroy the denominator estimate.

15.6 Frobenius products of matrices and the approximate projector

Continue with the same local representation; we now take the particular adjacent pair $b = r_0 - 1, b' = r_0$. Write

$$x_b = \frac{\mathbf{1}(b = r_0) - \mathbf{1}(b = r_0 - 1)}{P^\lambda}, \qquad \nu = 2\lambda.$$

Use a generic matrix X_j for each letter $j \in \mathbb{Z}/\nu$. Put $\Psi_{s,d} = X_{s-d}^{(P^d)}$, taking the entrywise power. The product

$$H_{s,m} = \Psi_{s,m-1} \cdots \Psi_{s,0}$$

for $m = \lambda + \delta(b, s)$ supplies row b of a generic coefficient action matrix A_s: row b of $H_{s,m}$ in an even letter, row b of its transposed adjugate in an odd letter. Let

$$\mathscr{F}(\Psi) = \bigotimes_s \mathrm{Sym}^{n_0} A_s.$$

The row-dependent product length is essential. In the formal source it is `TwistingData.frobOperator` in `New/TwistedHecke/FrobeniusFactors.lean`. If all generic matrices are conjugated by the corresponding letter embedding of $g \in L$, the induced operator changes by conjugation by $\rho(g)$ (with the order determined by the substitution on functions). Indeed at fixed s both $H_{s,\lambda}$ and $H_{s,\lambda+1}$ undergo ordinary conjugation. The selection of rows commutes with the block diagonal g_s, because δ is constant on each block; and the identity $\mathrm{adj}(GHG^{-1})^t = G^{-t}\,\mathrm{adj}(H)^t G^t$ gives the inverse-transpose rule in the odd letters. Finally the symmetric-power action respects products. This proves the exact covariance `frobOperator_map`. We distinguish this exact invariance from the approximate compatibility of conjugation with entrywise Frobenius powers.

Lemma 15.8 (Weights of products)**.** *Let $\mathcal{F}^u$ be any decreasing multiplicative filtration on a commutative coefficient ring. Let $\alpha_{s,d}(b)$ be real weights with $\sum_b \alpha_{s,d}(b) = 0$, and $|\alpha_{s,0}(b)| \le u$, $u \ge 0$. Suppose*

$$(\Psi_{s,d})_{bc} \in \mathcal{F}^{\alpha_{s,d+1}(b)-\alpha_{s,d}(c)-u} \qquad (0 \le d \le \lambda).$$

Put $\widetilde{x}(b,s) = \alpha_{s,\lambda+\delta(b,s)}(b)$ in an even letter and its negative in an odd letter. Then every entry in row β of $\mathscr{F}(\Psi)$ has filtration at least

$$\widetilde{x} \cdot \beta - \nu n_0 k(\lambda+2)u. \tag{15.7}$$

Proof. Expanding the product $H_{s,m}$ entry by entry and applying the multiplicative filtration gives $(H_{s,m})_{bc} \in \mathcal{F}^{\alpha_{s,m}(b)-\alpha_{s,0}(c)-mu}$, because the intermediate indices' weights telescope. This is at least $\alpha_{s,m}(b)-(m+1)u$. In an even letter this bounds the row of A_s. In an odd letter each cofactor of the transposed adjugate in row b is a signed sum of products using every row of $H_{s,m}$ except b. The sum-zero assumption on $\alpha_{s,m}$ gives filtration at least $-\alpha_{s,m}(b)-(k-1)(m+1)u$. Both estimates are at least $\widetilde{x}(b,s) - k(\lambda+2)u$. In the symmetric power, a contribution to the coefficient of the output monomial β_s uses exactly $\beta_s(b)$ entries from row b of A_s; the total loss per letter is at most $n_0k(\lambda+2)u$. Taking the product over all letters gives (15.7). □

This is the abstract filtration lemma `TwistingData.frobOperator_mem`. In particular the degrees in the variables of any one generic matrix which occur in $\mathscr{F}(\Psi)$ are at most $D = \nu n_0 k(\lambda+2)P^\lambda$: apply the same argument to the degree filtration of negative weights, or simply note that each symmetric-power term uses at most $n_0k(\lambda+1)$ factors of degrees at most P^λ in a letter, and then sum over the ν letters. This slightly coarse bound is sufficient for the interpolation cutoff in Proposition 15.7.

Now let $g \in L$ and replace the generic matrices used above with

$$\Psi_{s,d} = \iota_s(g)\big(\iota_{s-d}(g^{-1})X_{s-d}\iota_{s-d}(g)\big)^{(P^d)}\iota_s(g^{-1}). \tag{15.8}$$

Modulo p this is exactly $X_{s-d}^{(P^d)}$, by the coefficient Frobenius congruence and the fact that entrywise raising to a power of p is a ring homomorphism over characteristic p. In particular the error terms have coefficients divisible by $p = w^2$ and are homogeneous of degree P^d.

Proposition 15.9 (The weighted coefficient estimate)**.** *Let η be an exponent in the generic matrices, with restriction η_j to X_j. Assume its row weight equals its column weight at every $j \ne i$. Put t_i equal to its row weight at i, with weights x_b above. Assume $\nu n_0 k(\lambda+2) \le P^\lambda$. If the output exponent β satisfies*

$$|E_{i+\lambda}(\beta; r_0-1, r_0) - t_i| > 2c+1,$$

then every coefficient of X^η in row β of $\mathscr{F}(\Psi)$ for (15.8) is divisible by w^c.

Proof. Let $e \in \{-1,1\}$ be the sign of the indicated difference and put

$$A_a(b) = \frac{e}{2}\frac{P^a x_b}{q}, \qquad u = \frac{1}{2P^\lambda}.$$

Give the variable $(X_j)_{bc}$ weight $A_\nu(b)$ if $j = i$, and $A_{[j-i]_\nu}(b) - A_{[j-i]_\nu}(c)$ otherwise. These weights, denoted ω_{jbc}, satisfy $|\omega_{jbc}| \le P^{-\lambda}$. Set $\alpha_{s,d}(b) = A_{d+[s-d-i]_\nu}(b)$. Then $|\alpha_{s,0}(b)| \le u$ and their sum over b is zero. Consider the Gauss filtration on generic polynomials for the variable weights ω and coefficient parameter w.

The main monomial $(X_{s-d})_{bc}^{P^d}$ of the entry of (15.8) has weight at least $\alpha_{s,d+1}(b)-\alpha_{s,d}(c)-u$. For $j = s-d \ne i$, $d+1+[j-1-i]_\nu = d+[j-i]_\nu$, so its weight is exactly $\alpha_{s,d}(b) - \alpha_{s,d}(c)$.

For $j = i$ the weight is $A_{d+\nu}(b)$ and the desired expression may further subtract $A_d(c)$; since $d \le \lambda \le \nu$, its absolute value is at most u. This proves the main bound in both cases. An error monomial has coefficient divisible by w^2, and degree P^d, hence has Gauss order at least $2 - P^{d-\lambda} \ge 1$. Every $\alpha_{s,d}(b)$ and $\alpha_{s,d+1}(b)$ in this range has absolute value at most $1/2$ (the largest exponent is at most $d + \nu$), so its Gauss order is again sufficient. Lemma 15.8 applies.

For the special adjacent pair, the deep root variable y_s has $\delta = 1$ and z_s has $\delta = 0$, and the $\widetilde{x}$ of that lemma is precisely $(e/2)$ times the variable weight of $E_{i+\lambda}$. This is immediate by substituting $d = \lambda + 1$ for y_s and $d = \lambda$ for z_s, with the odd sign introduced by inverse transpose. At every other variable both sides vanish. On the generic exponent η the weight sum is $(e/2)t_i$: the other letters vanish by the row-column equality hypothesis. The filtration loss in (15.7) is at most $1/2$. Thus the required order of the coefficient is at least

$$\frac{e}{2}(E_{i+\lambda}(\beta) - t_i) - \frac{1}{2} > c.$$

In particular it is divisible by w^c, proving the proposition. □

This gives the coefficient calculation of `pow_dvd_coeff_conj_frobOperator` without an appeal to generic representation theory; the only input is the Frobenius congruence modulo p.

Let now $a_* = (k + k^2)d_0\nu^2$, $c_* = 2a_* + 2$, and assume $D + n_0 + k < P^\nu$ with the stronger size condition of (14.13), namely $\nu n_0 k(\lambda + 2) + n_0 + k < P^\lambda$. In particular $D + k < P^\nu$ and $n_0 < P^\nu = p^{d_0\nu}$. Take $F = d_0\nu$ in the diagonal pairing and the conjugation means, and take the homogeneous cutoff D. Let b_β be the product over letters of the integral reciprocal-multinomial diagonal entries q_{β_s}. Each divides $p^{kF\nu}$ in O (it is a quotient of that power by an integer multinomial product). Put $b'_\beta = p^{kF\nu}/b_\beta$, another integral diagonal entry. In tensor notation let B and B' denote these two diagonal matrices in the local coefficient space. They are distinct from the global budget B of Chapter 4; in this subsection the distinction is made by attaching a matrix argument, as in BF. They satisfy $BB' = p^{kF\nu}I$ and $B\rho(g^t) = \rho(g)^t B$ by (15.6).

Theorem 15.10 (Approximate projector)**.** *For every real h, there is an integral operator Z on the coefficient module such that*

$$ZF \in U_h \bmod w^{c_*} \quad \textit{for every } F,$$

and, for every $y_0 \in U_{h-(4c_*+2)} \bmod w^{c_*}$ and every F,

$$y_0^t BZF \equiv p^{a_*} y_0^t F \pmod{w^{c_*}}. \tag{15.9}$$

Proof. In every letter take the means of Proposition 15.7 for the weight x_b. Write Λ^{all} for the one with every row weight allowed, Λ^{hi} for that allowing row weight above $h - (2c_* + 1)$, and $\Lambda^{\mathrm{lo}} = \Lambda^{\mathrm{all}} - \Lambda^{\mathrm{hi}}$. These are invariant for conjugation in their letter, and vanish on unequal row-column weights. The last mean vanishes on row weights above $h - (2c_* + 1)$, since the first two agree there. Apply the tensor of the Λ^{lo} to the coefficients of the generic matrix $\mathscr{F}(\Psi)$ with $\Psi_{s,d} = X_{s-d}^{(P^d)}$, and multiply on the domain by B'. This defines Z.

To verify the first assertion, translate by any $g \in L$ and look at an output row β with excess above h in letter $i + \lambda$. Move the translate into the generic matrix using exact covariance and the invariant means; this changes the generic matrix to (15.8) and introduces only an integral change of the input. A monomial surviving the means has equal row and column weights at all letters, and row weight at i at most $h - (2c_* + 1)$. The difference in Proposition 15.9 is strictly greater than $2c_* + 1$. Therefore the output coefficient is divisible by w^{c_*}. We have obtained the first assertion for all Levi translates, not just for the chosen coordinates.

For the congruence, expand $\bigotimes_s(\Lambda^{\mathrm{all}} - \Lambda^{\mathrm{hi}})$. The all-term is $p^{k^2F\nu}$ times evaluation at the identity matrices, because the generic degrees are bounded by D. There the operator $\mathscr{F}$ is the identity. The contribution to the left of (15.9) is consequently $p^{k^2F\nu}y_0^tBB'F = p^{a_*}y_0^tF$.

Every other term contains at least one distinguished factor Λ^{hi}, say in letter i. By the span property of Property 3 in Proposition 15.7, express that factor as a linear combination of forms supported on row weights above $h-(2c_*+1)$ after a Levi conjugation. Invariance of the other factors permits the same conjugation in every letter. Move it off the generic matrices by covariance. The transpose of the conjugating coefficient action can be moved off B using $B\rho(g^t) = \rho(g)^tB$. Thus the new testing row vector is a Levi translate of y_0 with an additional diagonal factor B. Its coefficients at output rows of excess greater than $h-(4c_*+2)$ are zero modulo w^{c_*}; multiplication by B cannot spoil this vanishing. For all other output rows, the difference between the row weight supported in the distinguished generic factor and the output excess is strictly larger than $2c_*+1$. The nondistinguished factors still vanish on unequal row-column weights because they are invariant means. Proposition 15.9 applies again, so the entire contribution is zero modulo w^{c_*}. This proves (15.9). □

In `TwistingData.exists_projector` the cutoff constant is called c, here c_*. The estimate requires that the test vector lie in the smaller condition, not only that the output lie in the larger condition. This shift by $4c_*+2$ is precisely what makes the pairwise argument work after changing frames.

15.7 Primitive half-depth matrices

We pass from the preceding polynomial estimates to a determinant estimate for a sum over classes. First observe that the normalized half-depth operator is well-defined on all frames associated to the two-block parahoric. If $h \in \mathcal{P}$, any ordinary (even-letter) entry in the prohibited off-diagonal block is divisible by p under ι_s. Indeed an element x in the maximal ideal of R satisfies $1-x^{q-1}$ a unit; the full-cycle congruence $\iota_s(x)^q - \iota_s(x) \in (p)$ implies $\iota_s(x) \in (p)$. The same is true for the opposite prohibited entries in an odd letter by applying this to h^{-1} and transposing. On the coefficient variables, conjugating h_s by the single-letter diagonal with entries $w^{\delta(b,s)}$ therefore gives integral coefficients: a factor w^{-1} can occur only in the prohibited block, where it is canceled by a factor of p. Taking symmetric powers and then their tensor product gives

$$D_w\rho(h)D_w^{-1} \in \mathrm{GL}_{M_0}(O) \qquad (h \in \mathcal{P}). \tag{15.10}$$

Invertibility follows by the same assertion for h^{-1}. This is the half-depth version of the parahoric lemma of `TwistingData.lean`.

Let X be a finite class set and E_0 a finite set of frames with entries $x_e \in X$, $g_e \in \mathrm{GL}_k(R)$ such that $(x_e, g_e^{-1}\mathcal{P})$ covers X times the parahoric cosets. Allow a finite spectator index H_0, which need not undergo the nontrivial Frobenius action; equivalently replace X by $X \times H_0$ in the next construction. Consider the rectangular matrix

$$J : F = (F_x)_{x\in X} \longmapsto (D_w\rho(g_e)F_{x_e})_{e\in E_0}.$$

If all coordinates of JF are divisible by some scalar $a \in O$, the same is true of $D_w\rho(g)F_x$ for *any* x, g. This follows from surjectivity on the coset set and (15.10): if $g^{-1}\mathcal{P} = g_e^{-1}\mathcal{P}$, then $g = hg_e$ with $h \in \mathcal{P}$. In particular, J is injective over the fraction field of O, since all powers of w are nonzero and each x appears.

Proposition 15.11 (Primitive factorization)**.** *Write $J = Qd$ where d is square and Q is left invertible over O. Such a factorization exists and, with the maximal depth denoted by x_0, it can be chosen with*

$$\det d \mid w^{|X| \sum_{0 \le j < x_0} \dim V_j}.$$

Moreover every frame component of QF satisfies $U_h \bmod w^c$ whenever $h \ge 2c$ (at the boundary pair $r_0 - 1, r_0$).

Proof. Choose a Smith factorization of the full-column-rank matrix J over O. Such a factorization exists also for a nondiscrete valuation domain: select an entry among the finite matrix entries dividing all the others, clear its row and column by integral operations, and continue. Thus $J = Qd$ with Q primitive and d invertible over the fraction field of O. Denote its elementary divisors by t_i. There is an integral change of basis in the domain with basis vectors F_i for which the coordinates of JF_i are all divisible by t_i. By the all-frame divisibility just proved, t_i divides $w^{d(\alpha)}[X^\alpha]\rho(g)F_i(x)$ for every x, g, α. Since $d(\alpha) \le x_0$, taking $g = 1$ shows t_i divides every coordinate of $w^{x_0}F_i$. In an invertible basis each column has a unit coordinate, so $t_i \mid w^{x_0}$.

Let j_i be the smallest integer j with $t_i \mid w^j$. If $j < j_i$ then in every frame t_i divides the depth-weighted coefficient, yet t_i does not divide w^j. For any monomial of depth below j, therefore, the coefficient in that frame is a nonunit: indeed $w^{d(\alpha)} \mid w^j \mid t_i$ and the quotient $t_i/w^{d(\alpha)}$ cannot be a unit. Reducing these basis vectors modulo the maximal ideal gives independent vectors satisfying the depth condition in every frame, so

$$\#\{i : j_i > j\} \le |X| \dim V_j.$$

Multiplying divisors and summing this inequality over $j < x_0$ gives the asserted bound for $\det d$.

It remains to prove the excess condition on the primitive factor. The previous divisibility $t_i \mid w^{x_0}$ implies $w^{x_0}d^{-1}$ is integral; hence every column vector QF can be expressed as $w^{-x_0}JF'$ with F' in the integral domain of J. For each x, g and exponent α the inequality

$$w^{x_0} \mid w^{d(\alpha)}[X^\alpha]\rho(g)F'_x \tag{15.11}$$

holds by the all-frame property. Fix a letter i, and use the real variable weights $c(v) = \delta(v) - E_i(v)/2$ where E_i is for the boundary pair. For the two root variables the difference of weights is

$$c(y_s) - c(z_s) = 1 - (E_i(y_s) - E_i(z_s))/2 \in [2/5, 8/5],$$

using the explicit range $[-1/P, 1]$. For any fixed class x, all its full-group translates $\rho(g)F'_x$ have Gauss level at least x_0 with these weights. If not, choose a nonzero term with w-order l, a nonnegative *integer* defined by the largest power of w dividing that coefficient, and $l + c \cdot \alpha < x_0$. One can choose such a term with globally minimal integer order plus weight: for each exponent the set of attained finite integer orders below its violating threshold is finite, with a minimum, and there are finitely many monomials. Infinite valuation in powers of w cannot violate a finite bound. By the root lemma for the full group as Levi, that monomial must satisfy $E_i(\alpha) \le 0$. But (15.11) then forces

$$l + c \cdot \alpha = l + d(\alpha) - E_i(\alpha)/2 \ge x_0,$$

a contradiction. The Gauss inequality for all coefficients follows.

In a primitive frame the coefficient of QF is that of $w^{-x_0}D_w\rho(g_e)F'$. The Gauss inequality implies that a term of excess above $2c$ in letter i is divisible by w^c. Finally, D_w commutes with the block-Levi action: monomials are transformed within blocks, so their deep degree in each

letter is preserved. Thus the same conclusion applies in every further Levi translate of each frame. This gives membership in U_h mod w^c for $h \geq 2c$. The proof works with an arbitrary finite spectator index since every frame can be treated separately in those coordinates. □

This is the depth factorization `TwistingData.exists_depth_factorization`. Notice the use of the full-group root lemma in the second part, despite the final condition being only in block-Levi translates.

15.8 The Smith pairing lemma and the local Hecke theorem

We recall that the index condition $I(M, M'; z)$ of Chapter 14 means that every finite family in M can be made a family in M' by a square integral matrix whose determinant divides z. In a finite free module it suffices to exhibit such a matrix on a basis: more generally the condition on an arbitrary family then follows by pulling back under the map from its free coefficient module, as in Lemma 14.2. In particular a square matrix with determinant dividing z and image contained in M' gives the condition. We need the following consequence of Smith elimination.

Lemma 15.12 (Pairing rectangular factors)**.** *Let $Q, M : O^D \to O^{D'}$ be integral matrices, Q left invertible with inverse on its image R_Q, and $\beta : O^{D'} \to O^D$ integral. Suppose u_0, w are nonzero, w is a nonunit, and*

$$\beta M \equiv u_0 I \pmod{u_0 w}.$$

Suppose that for all v, v', F satisfying $Qv - Mv' = u_0 wF$, the image of F over the residue field belongs to a fixed vector subspace S of the reduced target. Then there is an integer $l \geq 0$ with $D + l \leq \dim S$ such that the submodule $\mathrm{im}(\beta Q) + zO^D$ has index dividing $u_0^D z^l$ for every $z \in O$.

Proof. Identify the target with the direct sum of the primitive image of Q and a complement; the latter is finite free over the valuation ring. Then Q is the first inclusion, and write $M = \binom{M_1}{M_2}$ relative to that decomposition. Make a Smith factorization of M_2 by changing the basis of the complement and the source of M, so it is diagonal rectangular with elementary divisors t_j, including zero divisors for missing pivots. Change the output basis of β compatibly with the new basis of the source of M; this preserves the congruence $\beta M \equiv u_0 I \bmod u_0 w$.

Let J be the set of positions with nonzero t_j dividing $u_0 w$ and $l = |J|$. The space S contains the reductions of all columns of Q (take $v = u_0 w e_j, v' = 0$). For $j \in J$ choose $v' = (u_0 w / t_j) e_j$, $v = M_1 v'$: the resulting vector F is (up to sign) the corresponding complement basis vector. These are independent modulo the image of Q, so $D + l \leq \dim S$.

For a position $j \notin J$, the divisor t_j is zero or is divisible by $u_0 w$ (valuation-domain comparability). The vector $\beta Q M_1 e_j = \beta M e_j - \beta\binom{0}{M_2 e_j}$ is congruent to $u_0 e_j$ mod $u_0 w$. For $j \in J$ instead use the vector $u_0 z e_j$, which lies in zO^D. These vectors are all in the desired submodule. Factor u_0 out of each of their columns and factor z from the columns $j \in J$. The determinant of the remaining matrix is congruent to one modulo w, and is therefore a unit. The index condition follows. This argument is independent of the complement coordinates, which were only changed by invertible integral matrices. □

This is `TwistingData.of_pairing`; the complement-block proof is an equivalent way to read the Smith factorization of the discrepancy matrix $(QR_Q M - M)R_Q$ used by the formal development.

We can now prove the precise local elementary-divisor estimate from Chapter 14. Let X be the finite class set, I_0 the spectator index set, and $V = O^{X \times I_0 \times \mathcal{M}}$, where $\mathcal{M}$ indexes the

monomials in the cyclic coefficient representation. Thus $D = \dim V = |X||I_0|M_0$. Write J_0 for the parahoric coset set. For each $e = (x, b) \in X \times J_0$ there are $s(e) \in X$, $g_e \in \mathrm{GL}_k(R)$ and an invertible spectator matrix u_e on O^{I_0}, with $e \mapsto (s(e), (g_e^t)^{-1}\mathcal{P})$ bijective (injectivity suffices since the finite sets have the same cardinality). Let T be the local Hecke matrix whose block from $s(e)$ to x is

$$u_e \otimes (\rho(g_e)D_w^2\rho(\gamma_b))^t,$$

where γ_b are the specified representatives with inverses representing the cosets J_0. Contributions with repeated class pairs are summed. This is the matrix of (14.18) in its chosen orientation. Define two half-depth maps into the frame module

$$(J_1F)_e = D_w\rho(g_e^t)F_{s(e)},$$
$$(J_2F)_e = D_w\rho(\gamma_b)F_x,$$

with trivial action on the spectator coordinates. Both sets of frames cover the requisite cosets. Write $J_1 = Qd$, $J_2 = Q'd'$ as in Proposition 15.11; both Q, Q' have integral left inverses. In the frame module let U be the block-diagonal operator acting as u_e on the spectator factor in frame e, and put $\mathbb{B}$ for the repeated diagonal matrix B of Theorem 15.10. Then by the transpose identity and $D_wB = BD_w$ one has an exact matrix identity, with B also repeated over the domain:

$$TB = J_2^tU\mathbb{B}J_1 = d'^t(Q'^tU\mathbb{B})Qd. \tag{15.12}$$

To check it, the contribution from frame e on the right, before the primitive factorizations, is $u_e \otimes \rho(\gamma_b)^tD_wBD_w\rho(g_e^t)$. Moving B to the right using its transpose covariance gives exactly the required block times B. This calculation is why the automorphic Hecke operator must be identified with the specific transpose action rather than merely an abstract endomorphism with the same eigenvalues.

Set $\beta = Q'^tU\mathbb{B}$ and choose a left inverse R' of Q'. Let Z of Theorem 15.10 act in each frame on each spectator factor, and define the rectangular matrix $M = U^{-1}ZR'^t$. The second primitive factor has every frame component in $U_{h-(4c_*+2)}$ mod w^{c_*}: indeed $h \geq 6c_* + 2$ by the size hypotheses, so this threshold is at least $2c_*$. Therefore the approximate projector gives, by summing over the finitely many frames and spectator coordinates,

$$\beta M = Q'^t\mathbb{B}ZR'^t \equiv p^{a_*}Q'^tR'^t = p^{a_*}I \pmod{w^{c_*}},$$

where $a_* = (k + k^2)d_0\nu^2$ and $c_* = 2a_* + 2$. Both Q and M have components in U_h mod w^{c_*}; for M this follows from the projector and the fact that U^{-1} acts only on spectator coordinates. If $Qv - Mv' = p^{a_*}wF = w^{c_*-1}F$, cancellation of w^{c_*-1} in the integral congruences means that F satisfies the excess condition modulo w, hence modulo the maximal ideal. We can take for S in Lemma 15.12 the direct sum of copies of U_h in every frame and spectator coordinate. The flag estimate (15.4) gives

$$\dim S = |X||J_0||I_0| \dim U_h \leq (1 + \varepsilon)D.$$

Lemma 15.12 yields an index condition for $\mathrm{im}(\beta Q) + p^sV$ with exponent at most $(a_* + s\varepsilon)D$ in p. Left and right multiplication by the square factors in (15.12) adds their index costs. Each determinant divides $w^{|X||I_0|\sum_j \dim V_j}$, whose sum of two exponents is at most $2(1 + z)D$ by (15.5); using $w^2 = p$ the corresponding extra exponent is $(1 + z)D$. The determinant pullback, pushforward and composition rules apply even after adding p^sV: an integral matrix sends p^sV into itself, and passing to the image can only reduce the required determinant. Consequently

$$I(V, TBV + p^sV; p^{(a_*+1+z+s\varepsilon)D})$$

in the sense that an integer exponent bounded by the displayed real exponent suffices. The $a_* + 1$ term is $c_*/2 \le h$, so the bound is at most $(z + \varepsilon s + h)D$. If $f : V \to M'$ is any integral linear map under which the Hecke action descends to Y on $f(V)$, then $f(TBV) \subseteq Yf(V)$ and the pullback rule transfers the same estimate to $Yf(V) + p^s f(V)$. The action preserves $f(V)$ by its very intertwining equation. This completes a mathematical proof of `TwistingData.hasType_twisted_hecke`, including the leading constants of the dimension and index estimates.

In the global application $n_0 = a$, $r_0 = s_p$ is the rank of the marked Hecke block, $d_0 = o_0$ and the local residue field has cardinality $q = P^\nu$. The spectator factor consists of the weight variables belonging to other places and letters, and all computations above act trivially on their monomial indices except for the integral matrices u_e. The injective correspondence of local frame labels, verified on the actual column class model, is the local-to-global formal-source identification `Data.IsMarked.hasType_markOperator` in `New/Columns/MarkedPrimeType.lean`. The present proof discharges the abstract coefficient-matrix theorem; it should not be read as supplying the local-to-global identification of g_e and u_e if the chosen level or weight representation is changed.

15.9 The finite Fourier operator at the level prime

The arithmetic prime p in the preceding argument is a prime of the base of the column coefficient lattice. We now work at the different prime N, the auxiliary split level prime of the chain. Unlike the twisted Hecke estimate, the Fourier operation at this prime is based on a small finite sum with no divisions by the size of a residue field. We give the argument with all the scalar factors; it supplies a useful test of the integrality of the denominator trade described in Chapter 14.

Let S be the finite chosen set of split places of E above N, $\Lambda = \prod_{w\in S} \mathcal{O}_{E_w}$, $F_S = \Lambda/N\Lambda = \prod_{w\in S} \mathbb{F}_N$. All definitions below refer to the places in S, not to their conjugates. Write V_j for a representation of $G_j = \mathrm{GL}_{k_j}(\prod_{w\in S} E_w)$, with action π_j, and let e_j denote the last standard basis vector of rank k_j. An F_S-valued coefficient vector is called everywhere nonzero if its image in $\mathbb{F}_N^{k_j}$ is nonzero at every $w \in S$; this is stronger than being a nonzero element of $F_S^{k_j}$. A scalar in F_S is a unit if each component is nonzero. The finite additive character used below is exactly

$$\psi_N(t) = \psi_0(-\,\mathrm{residueOverN}(t)),$$

where the w-component of $\mathrm{residueOverN}(t)$ is the chosen integer representative of t_w divided by N, and all components outside S vanish.

A residue function is extended to local matrices by zero outside the integral matrices. In rank k_{j+1} by k_j define three such functions:

$$b_y(X) = [\overline{X}e_j = y], \qquad e_z(X) = [e_{j+1}^t \overline{X} = z^t],$$
$$c_z(X) = \psi_N(e_{j+1}^t \overline{X} z).$$

Here y has length k_{j+1}, z length k_j, brackets denote an indicator, and $b = b_{e_{j+1}}$, $e = e_{e_j}$. Put $u_j = \mathrm{diag}(1, \dots, 1, N^{-1})$ at every place in S. These are the finite data `lastColDatum`, `lastRowDatum`, `charDatum`, and `diagNInv` in `New/Seesaw/NamesAtSplitLevel.lean`. For any rectangular local matrix X, one has the exact identity

$$e(u_{j+1}^{-1} X u_j) = b(X). \tag{15.13}$$

Indeed, in blocks the transformation sends $\left(\begin{smallmatrix} A & v \\ w & d \end{smallmatrix}\right)$ to $\left(\begin{smallmatrix} A & v/N \\ Nw & d \end{smallmatrix}\right)$. For its result to be integral with last row congruent to $(0, \dots, 0, 1)$ requires A, w, d integral, $v \in N\Lambda^{k_{j+1}-1}$ and $d \equiv 1$

(mod N); these say just that $b(X) = 1$. This proves the identity also when the matrix is not integral. Omitting that case would not suffice for the kernel covariance calculation.

A lift law consists of a linear map $f \mapsto L_j[f]$ from the space of local functions to linear maps $V_j \to V_{j+1}$, together with a scalar γ_j satisfying

$$\pi_{j+1}(B)L_j[f(\overline{B}\cdot\overline{A})]\pi_j(A) = L_j[f], \qquad A \in \mathrm{GL}_{k_j}(\Lambda), \tag{15.14}$$

$$\pi_{j+1}(u_{j+1})L_j[e] = \gamma_j L_j[b]\pi_j(u_j). \tag{15.15}$$

The notation in (15.14) also requires $B \in \mathrm{GL}_{k_{j+1}}(\Lambda)$ and f a residue function. This is precisely the structure `IsSplitLevel.LiftLaw`. The existence of the law for a concrete Weil kernel, with its correct normalization γ_j, is a separate assertion about kernel equivariance and Haar factors (see `New/Seesaw/LiftCharacters.lean`); we use that formal-source input when applying the algebra below. In particular (15.13) is the support calculation in the second law, not by itself a determination of its scalar.

Call a vector $v \in V_j$ ramified at S if, for each $w \in S$, there is an integral scalar unit a_w whose reduction is 1 at each other place, such that $v \in \mathrm{im}(1 - \pi_j(a_w I))$. This weaker formulation than a fixed character eigenspace is the exact one used in the formal proof. It immediately implies an annihilation rule. If a residue function is unchanged by multiplication of its matrix argument by every unit scalar supported at w, then $L_j[f]v = 0$: choose a_w witnessing ramification and apply (15.14) with $A = a_w I$, $B = I$. The operator $L_j[f](1 - \pi_j(a_w I))$ is zero. The same argument works for any function invariant under the specific selected a_w.

For $y \in F_S^{k_j}$ with y_{last} a unit, select an integral invertible K_y whose reduction has both last row and last column given by y (with the row transposed). Such a choice is possible: for $y = (v, t)$ the matrix

$$\begin{pmatrix} I + vv^t/t & v \\ v^t & t \end{pmatrix}$$

has determinant t, by the Schur complement formula, and satisfies both conditions. Lifting arbitrary entries to Λ preserves invertibility since its determinant is a residue unit. Define

$$\mathcal{F}_j = \pi_j(u_j) \sum_{\substack{y \in F_S^{k_j} \\ y_{\mathrm{last}} \in F_S^\times}} \psi_N(y_{\mathrm{last}})\pi_j(K_y). \tag{15.16}$$

This operator is defined using a chosen set-theoretic section; its action on the indicated invariants is what matters, not an assertion of independence of its choices on the whole representation. In particular, $\mathcal{F}_j$ commutes with the central scalar operators used to define ramification.

Proposition 15.13 (Sliding the finite Fourier operator)**.** *For a ramified vector v and a lift law as above,*

$$\pi_{j+1}(u_{j+1})L_j[c_{e_j}]v = \mathcal{F}_{j+1}L_j[b]v = \gamma_j L_j[b]\mathcal{F}_j v. \tag{15.17}$$

Proof. The function $c_{e_j}(X) = \psi_N(\overline{X}_{\mathrm{last,last}})$ is the sum over all bottom right entries $t \in F_S$ of $\psi_N(t)[\overline{X}_{\mathrm{last,last}} = t]$ on integral matrices. If t is a nonunit, there is a place w with $t_w = 0$. The displayed indicator is unchanged under central scalars supported at w. The annihilation rule shows its lift vanishes on v, so we may restrict the sum to t a unit. We may refine the remaining indicators either according to their entire last column or to their entire last row, obtaining

$$L_j[c_{e_j}]v = \sum_{y_{\mathrm{last}} \in F_S^\times} \psi_N(y_{\mathrm{last}})L_j[b_y]v = \sum_{z_{\mathrm{last}} \in F_S^\times} \psi_N(z_{\mathrm{last}})L_j[e_z]v.$$

The two sums have respectively vectors of target and source rank as their indices. By (15.14),

$$L_j[b_y] = \pi_{j+1}(K_y)L_j[b], \qquad L_j[e_z] = L_j[e]\pi_j(K_z),$$

since $b_y(\overline{K}_y X) = b(X)$ by the last-column condition and $e_z(X\overline{K}_z) = e(X)$ by the last-row condition. For example, the latter condition is equivalent to $\mathrm{lastrow}(X)\overline{K}_z = z^t = e_j^t\overline{K}_z$, hence to $\mathrm{lastrow}(X) = e_j^t$. Inserting these function identities in (15.14) gives exactly the two displayed lift identities. Applying $\pi_{j+1}(u_{j+1})$ to the column expansion gives the middle member of (15.17). Applying it to the row expansion and using (15.15) gives the final member. □

We have used precisely the two lift laws and ramification, with no averaging over the residue field. This is the proof of `IsSplitLevel.LiftLaw.u0_charDatum_eq_fourierOp`.

Proposition 15.14 (The paired chain)**.** *Suppose the consecutive lifts $L_0, L_1, \dots$ satisfy lift laws with constants $\gamma_0, \gamma_1, \dots$. Let d_i be arbitrary scalars and define the paired lift from rank k_{2i} to rank k_{2i+2} by*

$$\mathcal{P}_i = d_i\pi_{2i+2}(u_{2i+2}) \sum_{y \in F_S^{k_{2i+1}}} L_{2i+1}[c_y]L_{2i}[b_y].$$

Write $B_j = L_{j-1}[b]\cdots L_0[b]$ and $B_0 = 1$, with the standard last-column datum of the correct rank in every occurrence. For h pairs and a ramified initial vector v,

$$\mathcal{P}_{h-1}\cdots\mathcal{P}_0 v = \left(\prod_{0\le i<h} d_i q_i \prod_{0\le l\le 2i+1} \gamma_l\right) B_{2h}\mathcal{F}_0^h v, \qquad q_i = \prod_{w\in S}(N^{k_{2i+1}} - 1). \tag{15.18}$$

Proof. If y is not everywhere nonzero, b_y is invariant under central scalar units at a place with $y_w = 0$ and its lift on a ramified vector vanishes. For an everywhere nonzero y choose any K of middle rank with last column reducing to y; at every w a nonzero vector can be completed to a basis. Equivariance gives

$$L_{2i}[b_y] = \pi_{2i+1}(K)L_{2i}[b], \qquad L_{2i+1}[c_y]\pi_{2i+1}(K) = L_{2i+1}[c_{e_{2i+1}}],$$

the latter because $c_y(X) = c_{e_{2i+1}}(X\overline{K})$. Each surviving summand therefore has the same value; there are q_i of them. It follows from (15.17) that on a ramified vector $\mathcal{P}_i$ acts as

$$d_i q_i \mathcal{F}_{2i+2} L_{2i+1}[b]L_{2i}[b].$$

The ramification condition is preserved both by $\mathcal{F}_j$ and by $L_j[b]$. For the latter, use (15.14) with $A = a^{-1}I$, $B = aI$: the transformed matrix is unchanged. Hence $L_j[b]$ carries an image of $1 - \pi_j(aI)$ into the corresponding image of $1 - \pi_{j+1}(aI)$. On these vectors (15.17) gives $\mathcal{F}_j B_j = (\prod_{l<j}\gamma_l)B_j\mathcal{F}_0$, by induction on j. Starting with $h = 0$, induction on h now proves (15.18). This is the formula `IsSplitLevel.compUpTo_pairs_eq_smul`. □

Notice that each q_i is an N-adic unit. There is an important asymmetry between the constants q_i and γ_j: the former are known units directly by counting, whereas the latter require their precise Weil/kernel normalization. One must not replace the q_i by the count $N^{k_{2i+1}|S|} - 1$, which would overlook the several residue factors. In the arithmetic application the normalized scalars from the kernels are combined with the d_i in the formal-source theorem `Data.chain2_pairTranslates_eq_smul_chain` (subscript 2 in the source) of `New/Columns/TwistedHeckeType.lean`. The formula over local residue functions alone does not compute those scalars.

Lemma 15.15 (The mirabolic preservation calculation)**.** *For a fixed rank let $\mathcal{M} \subset \mathrm{GL}_k(\Lambda)$ be the matrices with last column congruent to the last basis vector. Let $g \in \mathcal{M}$ and let T be an endomorphism of a representation π. Suppose a vector v has the following property: $\pi(A)T\pi(B)v = v$ for all A, B with $AgB \in \mathcal{M}$. Then $T\mathcal{F}v = \mathcal{F}v$.*

Proof. Put $K = u^{-1}gu$. Both K and K^{-1} are integral, with reduction having last row e^t: apply (15.13) to g and g^{-1}. Thus $y \mapsto y' = \overline{K}y$ permutes the vectors with unit last entry and preserves y_{last}. For these vectors set $A = (uK_{y'})^{-1}$, $B = uK_y$. The reduction of $K_{y'}^{-1}KK_y$ fixes the last basis vector, so $AgB \in \mathcal{M}$. The hypothesis yields $T\pi(uK_y)v = \pi(uK_{y'})v$. Summing with coefficient $\psi_N(y_{\mathrm{last}}) = \psi_N(y'_{\mathrm{last}})$ proves the lemma. □

This is `IsSplitLevel.mirabolic_fix_fourierOp`. In the actual initial column space the assertion that the conjugated column level lies in this mirabolic is `Data.atPlaces_conj_mem_mirabolic` of `New/Seesaw/LiftCharacters.lean`; it includes the conjugate-place inverse-transpose condition. Using that assertion and the precise compatibility of the representation π with the smooth right action, the lemma is the preservation step of `Data.fourierOp_mem_column`. One still has to preserve integrality in arithmetic frames, which is stronger than preservation of the space of columns.

Example 15.16. If $N = 5$, two places occur in S, and a middle column has rank two, there are 25^2 residue vectors in total. Only $(25-1)^2 = 576$ enter the surviving paired sum. An individual Fourier operator of rank two, on the other hand, sums over $(5 \cdot 4)^2 = 400$ vectors with last entry a unit. Although neither count is the total field size, the paired count is prime to 5, exactly as required. This illustrates why the row/column distinction in Proposition 15.13 cannot be replaced by a sum over all vectors regardless of ramification.

15.10 Geometry and integrality in passing to a seesaw

The local calculation just proved answers a conditional question: if the data of two consecutive lifts are the indicated residue functions, what operator must be transferred to the end of the chain? Showing that the actual period can be reduced to that situation is a separate geometric and arithmetic question. We record the logical order of these reductions so that they are not concealed inside the local Fourier formula.

For a sequence of adjacent ranks, the lift $L_j[f]$ is an integral transform with a matrix parameter. The sum over global rational vectors that occurs in the theta kernel is invariant under simultaneous changes of two frames; one does not have the right to integrate each stage against a separate independent copy of a rational orbit. Instead one first uses the jointly invariant product of kernels on the full chain. Fold the odd-rank groups into the even-rank groups by integrating once over their common stabilizers. The relevant source statements are `Data.unfoldStage_even`, `Data.hasNearLines_of_unfoldStage_ne_zero`, and `Data.exists_stage_cut`, as described in Chapter 14. These are formal-source inputs here: they include the explicit global orbit classification and supports of the named Weil kernels.

At a finite cut one may replace integration over a compact class space by a finite weighted sum, provided the integrand is constant on the chosen classes. In particular if C is a compact domain with a finite disjoint partition $C = \coprod C_a$ and f is constant $f(a)$ on each measurable C_a, then

$$\int_C f = \sum_a \mathrm{vol}(C_a) f(a).$$

If a stabilizer acts with a finite orbit of size s_a on the chosen finite classes, the resulting multiplicity is s_a only when the volumes and values are equal on that orbit. More generally its contribution is the sum of those volumes times the common value. Applying this identity iteratively to the joint product of kernels requires a simultaneous cut, not one chosen independently for every appearance of a repeated class variable. The needed invariant, admissible finite cut is the formal-source input `Data.exists_isAdmissibleComb_unfolding`; its arithmetic expression by integral class frames is the input `Data.exists_isGoodDatum_of_invariant`. These are exactly the steps where one must establish finite support before redistributing scalar constants. The Gauss-sum calculation above can then be applied to the resulting chain of named finite data. It cannot replace the cut construction.

For clarity about the lattice index which remains after this reduction, suppose a coefficient module is a direct sum over a finite class set of integral polynomial coefficient representations. Let T_{last} denote the sum, over classes, monomials and embeddings contributing to the level prime, of the last-coordinate exponent. For homogeneous degree a in any given letter of rank k, the monomial basis is permuted by permuting the k coordinates; the sum of the exponents in all coordinates is a times the number of monomials. Consequently the sum in just the last coordinate is a/k times that number. The same statement holds with independent spectator letters, by multiplying their numbers of monomials. If D denotes the full dimension, f the number of relevant letters, the contribution of inverse level diagonal operators on this ambient space is bounded by faD/k. Transpose or conjugate letters are handled by taking the negative part of the diagonal exponent, so the resulting bound remains an upper estimate for denominator exponents. This is the determinant-counting mechanism behind the $1/k$ saving in the index bound for the Fourier operator.

To see how a denominator estimate composes, let L be an integral lattice of rank D over a valuation ring, and $g_1, \dots, g_s$ invertible endomorphisms of its fraction space. Suppose L_i is a lattice containing both the intermediate image and the appropriate integral frame lattice, and that the elementary divisors of the required change of frame at step i have total negative part at most b_i. A denominator-removing sublattice exists with determinant cost at most $\sum_i b_i$. Indeed, for a diagonal matrix of powers of a uniformizer, replace each basis vector whose exponent is negative by that vector multiplied by the inverse negative power; the resulting sublattice has the claimed determinant. Integral changes of bases preserve the determinant ideal, and successive intersections or inverse-images of sublattices add at most their determinant costs by the index rules of Chapter 14. This proves the same assertion for arbitrary matrices with these elementary divisors. Finite sums of frame operators must first be treated on a common cut or common lattice where all the summands act; in particular the argument is not a license to multiply the bound by no factor without first proving the simultaneous class-frame assertion.

In the source the resulting cost for the power of the level Fourier operator includes the extra change back to the reference class frame. The arithmetic statement `Data.exists_indexDvd_fourierOp_pow` in `New/Seesaw/LiftCharacters.lean` does this uniformly, using integral representatives of the CM classes and a product of inclusion/index conditions. It is a formal-source input here beyond the elementary weight-sum and composition calculation just given. The power transferred by (15.18) is $m/2$; the extra frame change is why the denominator allowance includes $m/2+1$. Likewise, the passage to the branching sublattice and its comparison with the non-isotypic column representation require the specified archimedean coefficient and complex CM character estimates, not finite-vector counting alone.

The principal improvement in this chapter is thus the proof of the abstract twisted Hecke estimate, including its nonsemisimple finite-field component and integral approximation, and the proof of the two Fourier-sliding identities. The identification of those operators with the specific marked-prime columns, and the arithmetic and geometric machinery realizing the

seesaw, are still explicitly identified formal-source inputs. Keeping the distinction makes it possible to reuse the local estimate independently while leaving the original arithmetic proof's normalization intact.

Chapter 16

Normalized CM reciprocity and the geometry of the iterated seesaw

The passage from an adelic chain with varying signature to a chain on definite groups is the essential geometric step in the upper bound. There are two different issues. First, the finite vectors in the definite chain must be related to the arithmetic theta series of Chapter 12, and the relation must hold under conjugation of the coefficients. Secondly, the data of the definite chain must be obtained with coefficients integral away from the twist conductor, not simply as an arbitrary element of a complex tensor product. We give proofs of the normalized reciprocity statement, the rational orbit calculation underlying one unfolding, the rigidity of integral lattice lines, and the finite-cut calculation including its exact multiplicities. We then explain the assembly of these results into the two-edge evaluation and iterated seesaw. The conclusions of the latter construction distinguish the elementary reductions established here from the remaining specialized comparison of finite and infinite Weil models.

16.1 A marked algebraic curve and its sections

We first treat the normalization which was left open in Chapter 12. Use $K = \mathbb{Q}(\sqrt{-d})$ with d prime and $d \equiv 7 \pmod 8$, and write $\omega = (1 + \sqrt{-d})/2$. To avoid confusion with the Weil action, denote a field automorphism fixing K by γ. For a CM lattice C let L be its period lattice, t its distinguished half-period and $a_1 = a_1(C)$ its scalar in the equation

$$Y^2 + a_1 XY = X^3 + X.$$

As in the preceding CM chapter, write θ_C for the even theta function of degree one normalized to have value one at zero. Write s_C for the canonical Hermitian sigma section, $G_C = \phi_C \theta_C^3$ and $s_C^2 = \xi_C \theta_C^2$, where ξ_C, ϕ_C are the translated elliptic coordinates. In particular the two functions s_C^2, G_C are entire. It is essential that the sigma function here is the one with the Hermitian quasiperiod, not the uncorrected Weierstrass product. We use the analytic uniformization theorem for elliptic curves and the algebraicity of morphisms of elliptic curves, in their classical characteristic-zero forms.

The marked origin. The uniformizing coordinates are

$$\xi_C(z) = \wp_L(z + t) - \wp_L(t), \qquad \phi_C(z) = \tfrac{1}{2}(\wp_L'(z + t) - a_1 \xi_C(z)).$$

The point with uniformizing coordinate zero is the affine point $(0, 0)$, while the point with coordinate t is the point at infinity. We take $(0, 0)$ as the origin of the group law, translating

the usual group law of the projective cubic by that rational point. This is an algebraic group law. The invariant differential is $dX/(2Y + a_1X)$, whose pullback is dz by the identity for ξ'_C. In particular the scalar multiplier of an isogeny is its multiplier on this differential. Conjugation of the equation and of the rational functions defining an isogeny transports this differential and changes the multiplier a to $\gamma(a)$. No application of γ to an analytic exponential is involved.

Lemma 16.1 (Marked conjugate curve)**.** *There is a CM lattice D of the same discriminant $-d$, with $a_1(D) = \gamma(a_1(C))$. If C_1, C_2 and their conjugates D_1, D_2 are so marked, an isogeny given analytically by multiplication by a from C_1 to C_2 is transported to multiplication by $\gamma(a)$ from D_1 to D_2, with the same degree. Its action on the distinguished half-periods is transported as well.*

Proof. Conjugate the cubic and its nonzero discriminant. Uniformize it, using the differential just specified to fix the analytic scale, and choose the distinguished half-period to be the point at infinity in the translated group law. The explicit conversion to Weierstrass coordinates gives the required equalities for $\wp(t), g_2, g_3$. To check the extra CM conditions, use that on a complex elliptic curve a homomorphism of tori is algebraic: its pullback on meromorphic functions is an inclusion of elliptic function fields, hence a morphism of the projective nonsingular curves. The endomorphisms $a \in \mathcal{O}_K$ conjugate to endomorphisms on D with analytic multiplier $\gamma(a) = a$. Consequently $\mathcal{O}_K L_D \subset L_D$. The two endomorphisms $\omega, \bar{\omega}$ do not kill the half-period of C by the definition of the marked lattice; they cannot kill its transported point, since the conjugation on points is bijective and preserves all group operations and identities. This proves the half-period condition on D. Conjugation of an isogeny preserves the degree, for example by the induced extension of function fields, and its differential gives its analytic multiplier. It also preserves the condition that the image of the marked point equals the origin. □

This is a convenient conventional proof of the algebraic part of `CMLattice.exists_galConjugate` and `conj_isogeny_mem_lattice`, in `Lib/CM/MainTheorem/ConjugateCurve.lean`. The formal source implements the conjugation of isogenies by comparing sections. Both approaches have to retain the specified differential and half-period; a statement just about the j-invariant does not suffice.

The canonical multiplier. Let A_L be the area of the lattice, let $H_L(z, w) = \pi z\bar{w}/A_L$, and let E_L be the integral alternating Riemann form, normalized so that its values on a positive lattice basis are ± 1. The multiplier of θ_C on the lattice is $\chi_C(l) = \mathrm{sgn}_\sigma(l) \exp(2\pi i E_L(l, t))$. Here the sigma sign is $+1$ on $2L$ and -1 on the other classes. There is a useful expression for the multiplier not involving t:

$$\chi_C(l) = (-1)^{E_L(\bar{\omega}l, l)}. \tag{16.1}$$

Indeed $\mathcal{O}_K/(2) \simeq \mathbb{F}_2 \times \mathbb{F}_2$, and $\omega, \bar{\omega}$ are complementary idempotents modulo 2 because their sum is 1 and $\omega\bar{\omega} = (d+1)/4$ is even. The lattice is a rank-one projective $\mathcal{O}_K$-module, so $L/2L$ is the sum of the two one-dimensional eigenspaces. The alternating form on this quotient is perfect. Thus the right side of (16.1) is negative precisely on the element with both nonzero components. As ωt and $\bar{\omega} t$ are not in L, that element is the class of $2t$. The sigma sign and the alternating pairing with $2t$ give the same rule on the four elements of $L/2L$. This proves (16.1).

Let multiplication by a be an isogeny $C \to C_2$ of degree N. Then

$$H_{L_2}(az, aw) = NH_L(z, w), \qquad E_{L_2}(az, aw) = NE_L(z, w).$$

Inserting $\bar{\omega}l$ and l into these equalities shows from (16.1) that $\chi_{C_2}(al) = \chi_C(l)^N$. A degree-$j$ section of C_2, pulled back by multiplication by a, is therefore a degree-jN section of C. This is the full character compatibility of `CMLattice.comp_mul_mem_sections` in `Lib/CM/Sigma/CurveTheta.lean`. Its proof does not require the isogeny to have odd degree.

Lemma 16.2 (Section transport and point evaluation)**.** *Let $S_m(C)$ be the sections of degree m. Put*

$$R(z) = \frac{\mathrm{Tay}_0\,\theta_D(z)}{\gamma(\mathrm{Tay}_0\,\theta_C(z))} \in 1 + z\mathbb{C}[[z]].$$

There is a γ-semilinear isomorphism $\mathcal{T}_m : S_m(C) \to S_m(D)$ satisfying

$$\mathrm{Tay}_0\,\mathcal{T}_m f = R^m \gamma(\mathrm{Tay}_0\, f). \tag{16.2}$$

At any conjugate pair of points $P \in C(\mathbb{C})$ and $P^\gamma \in D(\mathbb{C})$, take lifts z, z' to their period planes. A nonzero scalar λ depending on the lifts but not on m or f satisfies

$$(\mathcal{T}_m f)(z') = \lambda^m \gamma(f(z)). \tag{16.3}$$

In particular transport preserves the divisor and multiplicities of the zero set on the elliptic curve.

Proof. First divide f by θ_C^m. The quotient is elliptic and meromorphic, with its only possible pole at the marked half-period t. In elliptic coordinates every such function is a unique expression

$$P(\xi_C) + \phi_C Q(\xi_C), \qquad 2\deg P \le m, \quad 3 + 2\deg Q \le m;$$

the inequality for P is omitted if $P = 0$, and likewise for Q. To see this, functions regular away from t are the coordinate ring of the affine cubic. Its relation is monic quadratic in Y, so the displayed expressions are unique. Their pole orders at t are respectively even and odd and cannot cancel, since ξ_C has a pole of order two and ϕ_C one of order three. Conversely each displayed monomial times θ_C^m is entire and has the required multiplier. This also proves the monomial spanning statement in the source.

Apply γ to the coefficients of P, Q and use the corresponding functions of D, multiplied by θ_D^m. This defines $\mathcal{T}_m$. The formal Taylor coefficients of ξ_C, ϕ_C are recursive rational polynomial expressions in $a_1(C)$: the differential equations

$$\xi' = 2\phi + a_1\xi, \qquad \phi' = 1 + 3\xi^2 - a_1\phi, qquad\xi(0) = \phi(0) = 0$$

determine every coefficient in characteristic zero. Hence their Taylor series are carried to those on D. This proves (16.2).

Off the marked point use $\lambda = \theta_D(z')/\gamma(\theta_C(z))$. The algebraic functions ξ, ϕ have the conjugate values at the two points, so (16.3) follows. At the marked point use the alternative generators s_C^2, G_C of degrees two and three. Both are nonzero there and $G_C^2 = s_C^6$ there, by the cubic identity. Set $a = s_D(z')^2/\gamma(s_C(z)^2)$ and $b = G_D(z')/\gamma(G_C(z))$. Then $b^2 = a^3$ and $\lambda = b/a$ gives $\lambda^2 = a$, $\lambda^3 = b$. All monomial sections are products of θ_C, s_C^2, G_C, so this gives the same evaluation property at the marked point. The algebraic expressions identify the sections of $\mathcal{O}(mt)$; their divisors, including vanishing orders at t, are preserved under algebraic conjugation. □

The source analogues are `thetaQuotient`, `taylorAtZero_monomialSection`, and `exists_conjugate_section` in `Lib/CM/LocalHeight/IsogenyIntegrality.lean`. Observe why the unknown factor R^m is allowed at this stage. The isomorphism of section rings is algebraic, but the analytic trivializations by normalized theta functions may differ. The following rigidity argument is what removes this factor.

Proposition 16.3 (Rigidity of the normalized theta Taylor series)**.** *For the conjugate CM lattice of Lemma 16.1, one has*

$$\mathrm{Tay}_0\,\theta_D = \gamma(\mathrm{Tay}_0\,\theta_C).$$

Proof. For an isogeny $a : C \to C_2$ of degree N, the section $F(z) = \theta_{C_2}(az)$ belongs to $S_N(C)$. Its divisor is the inverse image of the marked point under the isogeny, with each point simple. By Lemmas 16.1 and 16.2, $\mathcal{T}_N F$ has that same divisor on the conjugate curve as $\theta_{D_2}(\gamma(a)z)$. The ratio of these two sections has no pole or zero, so is constant on a compact elliptic curve. At zero both equal one, by (16.3) with the normalized lift zero (or directly by (16.2)); the constant is one. Taking Taylor series gives $R_2(\gamma(a)z) = R_1(z)^N$.

Take $C = C_2$ and the two endomorphisms $a = 2$ and $a = \omega$. If $R \neq 1$, let $j \geq 1$ be the first positive index with coefficient $r_j \neq 0$. The first relation, $R(2z) = R(z)^4$, gives $2^j = 4$, hence $j = 2$. The second relation gives $\omega^2 = \omega\bar{\omega}$. Since $\omega \neq 0$ and $\omega \neq \bar{\omega}$, this is impossible. Notice that the degree used here is the actual norm, not a degree inferred from an action on a formal parameter. This proves $R = 1$. □

The result is `CMLattice.thetaQuotient_eq_one` in `Lib/CM/MainTheorem/ConjugateCurve.lean`. Conjugation of the normalized theta coefficients is much stronger than conjugation of the cubic. This proof has no appeal to a theorem on transcendental periods.

16.2 Simultaneous torsion translations and finite-adelic reciprocity

The next argument is needed because the upper bound does not involve only the theta function at the origin. It involves finite-adelic translates of the generating vector. For y in the complex period plane set

$$U_y\theta_C(z) = \exp\bigl(-\pi(z\bar{y} + |y|^2/2)/A_L\bigr)\theta_C(z + y).$$

The addends in the exponent are not individually algebraic, so coefficient conjugation is not to be applied to them separately.

Proposition 16.4 (One multiplier for a finite set of translates)**.** *Assume $L_C = \Omega_0\mathcal{O}_K$. Given a finite subset $S \subset K$, there is $v \in L_D$ such that for every $x \in S$*

$$\gamma(\mathrm{Tay}_0\,U_{x\Omega_0}\theta_C) = \mathrm{Tay}_0\,U_{xv}\theta_D. \tag{16.4}$$

Proof. Choose a positive integer n with $nS \subset \mathcal{O}_K$. Let $u = \Omega_0/(2n^2)$ and choose a complex lift w on D of the point conjugate to u on C. As $2n^2u \in L_C$, the conjugate point is killed by $2n^2$; hence $v = 2n^2w \in L_D$. The scalar w is chosen once for all $x \in S$. The point nxu is transported to nxw because $nx \in \mathcal{O}_K$ and the marked CM action is preserved.

Fix $x \in S$, set $y = x\Omega_0$, $y' = xv$, and compare the two functions

$$F_C(z) = U_y\theta_C(nz), \qquad F_D(z) = U_{y'}\theta_D(nz).$$

They are sections of degree n^2 for their respective curves. Indeed the ordinary theta pullback has degree n^2, and the extra multiplier from translation is $\exp(2\pi i E_L(nl, y)) = 1$ since $ny \in L_C$ (and similarly $ny' \in L_D$). Their zero divisors are $nz + y \equiv t_C \pmod{L_C}$ and $nz + y' \equiv t_D \pmod{L_D}$. These divisors correspond under γ: the points y, y' correspond because $y = 2n(nx)u$, $y' = 2n(nx)w$. Therefore $\mathcal{T}_{n^2}F_C = cF_D$ for a nonzero constant c.

To determine c, evaluate at $z_0 = -nxu$, $z_0' = -nxw$. These are conjugate points and $nz_0 = -y/2$, $nz_0' = -y'/2$. The exponential factor of the shift is exactly one there. Since θ is even,

$$F_C(z_0) = \theta_C(nz_0), \qquad F_D(z_0') = \theta_D(nz_0').$$

The ordinary pullback of theta is carried exactly to the ordinary pullback of the conjugate theta by the isogeny calculation in Proposition 16.3. Thus the common scale λ^{n^2} in (16.3) cancels, proving $c = 1$ if the value is nonzero. If it vanishes, then $y/2$ and $y'/2$ are the distinguished half-periods modulo their respective lattices. In this case $y \in L_C$, $y' \in L_D$, with classes $2t_C, 2t_D$ modulo twice the lattice. By (16.1) their multipliers are -1. Hence at zero $F_C(0) = F_D(0) = -1$. The normalized scale at zero is one, and again $c = 1$.

Proposition 16.3 makes $R = 1$ in (16.2). Taking Taylor series of the equality of sections and undoing the rescaling by n proves (16.4). Every choice made before fixing x works on the whole finite set. □

This proves `CMLattice.exists_taylorVia_twistedTranslate_theta` in `Lib/CM/MainTheorem/ConjugateCurve.lean`. The special evaluation at the negative half-translate is important: using a value at zero alone can fail if the shift moves the zero divisor there. Also note that the choice of v may depend on the finite set S; it has not been assumed to be a homomorphism of adeles.

Theorem 16.5 (Soft normalized CM reciprocity)**.** *There exist $\Omega \neq 0$ and an integer $e > 0$ with*

$$A_{L_D} = |\Omega|^2\sqrt{d}/2, qquade\Omega \in L_D.$$

For each finite $S \subset K$ one can choose $a_S \in K$ such that

$$\gamma(\mathrm{Tay}_0\, U_{x\Omega_0}\theta_C) = \mathrm{Tay}_0\, U_{x\Omega a_S}\theta_D \quad (x \in S).$$

Moreover the net a_S indexed by finite subsets ordered by inclusion has a cluster point in the finite adeles of K.

Proof. Here use the elementary odd-class-number argument of Chapter 6.2, which makes the square map on the class group bijective. Write $L_D = u\mathfrak{c}^2$ for a nonzero complex number u and a nonzero integral ideal $\mathfrak{c}$ (choose a representative of the square root of the lattice ideal class and absorb the principal factor in u). Set $e = \mathcal{N}\mathfrak{c}$ and $\Omega = eu$. The area of the lattice of an integral ideal of K is its norm times $\sqrt{d}/2$. Thus the claimed area identity follows from multiplicativity of the ideal norm. Since $e \in \mathfrak{c}$, one has $e\Omega = e^2u \in L_D$.

Use Proposition 16.4 for each S and write the resulting lattice element v as $u\beta_S$ with $\beta_S \in \mathfrak{c}^2$. Then $a_S = \beta_S/e$ works. Every a_S belongs to the fixed compact subset $e^{-1}\widehat{\mathcal{O}}_K$ of the finite adeles. The closures of the tails of the net in this compact set have the finite intersection property; their intersection is nonempty. This gives the cluster point. □

The theorem is exactly the soft statement `CMLattice.complex_multiplication_theta` in `Lib/CM/FormalGroup/FormalHeight.lean`. It does not assert an identification of the cluster point with the Artin symbol. Elliptic uniformization and the classical marked-curve comparison are standard external inputs, but the differential, section and scalar normalization are now derived. If a finite-adelic translate is locally constant in its adelic translation variable, its value on the net agrees with its value at the cluster point along a cofinal subnet: for each fixed finite set of torsion tests, the equality already holds on the corresponding tail. This is the passage needed by the normalized generating-vector construction of Chapter 12. It avoids the invalid assertion that arbitrary complex Taylor series are continuous in the finite-adelic topology before the finite Weil realization has been made. The precise comparison of the finite and infinite theta models is still used at that point.

16.3 Removing one string: the Hermitian linear algebra

We now turn to the geometry behind the passage to the definite chain. All the gradings in this section are those of `Grading` in `Lib/Weil/Periods/DegreeAndUnipotent.lean`. Temporarily let E be the CM coefficient field, with fixed field E^+ under the involution. A graded Hermitian space of string length s is

$$V = X \oplus \langle e_0, \dots, e_{s-1} \rangle_E,$$

where X is the positive-definite space with identity matrix. String vectors pair to 1 with their reflected partner e_{s-1-i}, except that the middle vector, if s is odd, has norm $-N$. All other pairings are zero, and the string is orthogonal to X. Here N is the positive split level prime from Chapter 4; the Hermitian form itself is defined over E^+. Let J_s be this matrix, and use the grades

$$\mathrm{gr}(X) = 0, \qquad \mathrm{gr}(e_i) = s - 1 - 2i.$$

The reflection of a string index is written $i^* = s - 1 - i$; reflection on X fixes each chosen basis index. For a matrix M from a graded space to another, write $\deg M \geq a$ if $M_{vu} = 0$ whenever $\mathrm{gr}(v) < \mathrm{gr}(u) + a$.

The adjoint M^* is $J_s^{-1} \bar{M}^t J_{s'}$ and sends a matrix of degree at least a to a matrix of degree at least a in the opposite direction. In fact each nonzero entry is paired with one whose row and column grades are the negatives of the original column and row grades. Products add the lower bounds on degree. These elementary properties apply both to rational matrices and to adelic ones.

Lemma 16.6 (The graded Bessel subgroup)**.** *Let U_s be the unitary group, let U_s^+ consist of the $g \in U_s$ such that $\deg(g-1) \geq 1$, and let B_s consist of those unitary matrices for which*

$$ge_i - e_i \in \bigoplus_{\mathrm{gr}(v) > \mathrm{gr}(e_i)} Ev \quad \text{if } \mathrm{gr}(e_i) \geq 0.$$

Then $B_s = U_s^+ \rtimes U(X)$, and equivalently $g \in B_s$ if $\deg g \geq 0$ and its degree-zero part acts as identity on the string and arbitrarily unitarily on X. All these identities hold also over the adeles and the finite adeles.

Proof. In the defining condition of B_s, preservation of the positive flag, and the identity on its successive lines, are immediate. The annihilators of that flag under the Hermitian form form the complementary negative filtration; unitarity implies their preservation and gives the inverse adjoint action on the negative graded pieces. If s is odd, the grade-zero quotient is $X \oplus Ee_{(s-1)/2}$. The middle string vector is fixed; its norm is nonzero, so the orthogonal complement X is preserved in that quotient. If s is even, the grade-zero quotient is simply X. Altogether this proves the description of the degree-zero part, and multiplication identifies it with a unitary matrix x acting on X. Then $gx^{-1} \in U_s^+$. Conversely any product of the stated kind satisfies the defining condition. The arguments on filtrations apply over each completion, and to the restricted product because all the relevant matrix identities preserve integrality at almost all places. □

Both groups are closed. The rational points of B_s are cocompact in its adelic group: for U_s^+ this follows by the positive-degree filtration, whose successive groups are additive vector groups over E^+, and the compactness of $E^+ \backslash \mathbb{A}_{E^+}$. One can choose a compact set of representatives successively using the filtration and lifts of compact subsets of the vector groups. The quotient for $U(X)$ is compact by anisotropy and the finiteness of its class set, as in Chapter 5. The combination gives a compact set of representatives for B_s. In the formal source

this is `Grading.isCocompact_ratBessel` in `Lib/Weil/Periods/BesselIntegral.lean`; no assertion of compactness of the full quotient of the indefinite group is made.

Write $V' = X' \oplus \langle e'_0, \dots, e'_{s-2} \rangle$ for the next space, with the same N and with parameter $c \in (E^+)^\times$ (for our chain $c = -(jg\varpi)^{-1}$). Define $T : V \to V'$ to kill X and the final string vector, and to send e_i to $t_i e'_i$, where, if $0 \le i < s-1$,

$$t_i = \begin{cases} 1, & s \text{ odd and } 2i+1 < s, \\ -Nc, & s \text{ odd and } 2i+1 = s, \\ c, & s \text{ odd and } 2i+1 > s, \\ c, & s \text{ even and } 2i+1 < s, \\ 1, & s \text{ even and } 2i+1 > s. \end{cases} \tag{16.5}$$

This is `Grading.stringMap` in `Lib/Weil/Periods/FiniteBesselGroup.lean`. In particular T is homogeneous of degree -1. Let S_s send each e_i to ce_{i+1}, with extra factor $-N$ at degree zero and $-Nc$ at degree one, and kill X and e_{s-1}. It has degree -2.

Lemma 16.7 (The two moment-map identities)**.** *For the string maps in* (16.5)*,*

$$T^*T = S_s, \qquad TT^* = S_{s-1}.$$

If $g \in B_s$ and $h \in B_{s-1}$, then $\deg(hTg - T) \ge 0$.

Proof. Put $h_i = -N$ at the middle string index and $h_i = 1$ otherwise; similarly put h'_i for the target. Directly from the Gram matrices, the adjoint sends e'_i to

$$\bar{t}_{s-2-i} \frac{h'_{s-2-i}}{h_{s-2-i}} e_{i+1}.$$

Consequently the coefficient of e_{i+1} in T^*Te_i is $t_i t_{s-2-i} h'_{s-2-i}/h_{s-2-i}$, because each t_i is fixed under conjugation. If neither of the paired indices is a middle index, it equals c. In the odd case an exceptional factor $-N$ at degree zero occurs, and in the even case the target middle index gives $-Nc^2$ at degree one. These are exactly the indicated coefficients; the same calculation on V' gives TT^*. Finally, the degree-zero parts of g, h are identity on string coordinates, and T kills all other coordinates. In expanding $hTg - T$, each surviving term therefore contains a positive-degree factor from h or g, and hence has degree at least zero. □

For example, in two consecutive ranks with string length 3 and 2, T sends $e_0 \mapsto e'_0$, $e_1 \mapsto -Nce'_1$, $e_2 \mapsto 0$. Its adjoint sends $e'_0 \mapsto ce_1$, $e'_1 \mapsto e_2$. Composing gives $ce_1, -Nce_2, 0$ on the old string, and $-Nc^2 e'_1, 0$ on the new one. These are not the moment-map identities for a string with the middle Gram entry 1. It is the middle factor N that will reappear in the local lattice restriction.

16.4 The surviving rational orbit

For maps $w : V \to V'$ consider the Heisenberg bicharacter in rational coordinates

$$\beta(w, z) = \psi\left(\tfrac{j}{2} \operatorname{tr}(z^* w)\right)$$

with the additive character and purely imaginary j used in the global Weil model. We suppress the trace to the rational adeles inside ψ. For the purposes of this calculation β has the usual skew perfect pairing after taking the ratio with its reverse. Its precise half-factor is important

in relating the phase of the two moment maps. Let d be target weights with $d(v^*) = -d(v)$. The survival condition at weights d is:

$$u \in U_s^+, \quad \deg(u-1) \geq 2, \quad \deg_d(w(u-1)) \geq 1 \quad \Longrightarrow \\ \beta(Tu, T) = \beta(wu, w). \tag{16.6}$$

Here $\deg_d$ uses the old source grades and target weights d. This is exactly `Grading.Survives` in `Lib/Weil/Periods/UnfoldingStep.lean`. We prove the linear algebra responsible for its consequences. It is useful to do so separately from the analytic Fourier expansion.

Lemma 16.8 (An Eichler test)**.** *Suppose w satisfies (16.6). Fix an old string index a and let $k = e_{a^*}$. Suppose that a vector v has support in positive grades at least $\max(1, \mathrm{gr}(a) + 2)$, that wv has support in target weights $> \mathrm{gr}(a)$, and that for every nonzero component of wk and every nonzero component of v the target weight plus source grade is positive. Then*

$$(T^*Tv)_a = (w^*wv)_a.$$

The same statement holds for adelic w, v under the indicated support assumptions.

Proof. Both v and k are isotropic: v lies in the positive-grade string and k is isotropic unless a is the middle string index. If a is the middle string index, k is not isotropic; this case is handled by the correction below. Moreover $\langle k, v\rangle = 0$ because $v_a = 0$. For any adelic scalar z, let $k_z = zk$ and put

$$M = vk_z^* - k_z v^*, \qquad u = 1 + M - \tfrac{1}{2}\langle k_z, k_z\rangle vv^*.$$

Here vectors are read as column maps and the star is the Hermitian adjoint. Direct multiplication gives $M^* = -M$, $M^2 = -\langle k_z, k_z\rangle vv^*$, $M^3 = 0$, and hence $u^*u = 1$. (This convention for the sign follows by writing $u = 1 + M + M^2/2$.) The degree of M is at least two, and that of M^2 at least four. The two assumed support properties show that wM has degree at least one relative to d; the term wM^2 has that degree as well by multiplication by M. Thus (16.6) applies.

If k is isotropic, the quadratic term is absent. If not, apply the character identity separately to z and $-z$, and divide; the term quadratic in z cancels. In either case the identity for all z implies that the difference $\langle wk, wv\rangle - \langle Tk, Tv\rangle$ vanishes: the linear character in z is trivial for every adelic z only if its coefficient is zero, by the perfect adelic additive pairing. More explicitly the trace of $(w^*w - T^*T)(vk_z^* - k_z v^*)$ is the difference of $\bar{z}\langle k, (w^*w - T^*T)v\rangle$ and its conjugate, and multiplication by the nonzero purely imaginary j detects both components when z varies. The entry in the reflected position is paired by the nonzero Gram value with the coordinate at a. This proves the claim. □

The rank-one test is the mechanism of `Grading.eq_of_survives`. The correction in the formula for u is necessary if the reflected basis vector has nonzero norm. Our applications include precisely this possibility.

Define $\mathrm{Agree}(w, T)$ to mean that $w_{ba} = T_{ba}$ whenever the source grade $\mathrm{gr}(a)$ is positive and the target grade $\mathrm{gr}(b)$ is smaller. Notice that w need not agree with T at positive or zero relative degree, nor at all entries in a negative source column.

Proposition 16.9 (Rational normalization of the surviving orbit)**.** *Let w be a rational map $V \to V'$ satisfying (16.6) for the zero target weight. Then a rational unitary matrix h on V' makes hw agree with T.*

Proof. Write V_+ for the strictly positive-grade string subspace. First $w|_{V_+}$ is injective. If not, take a nonzero $v \in V_+$ killed by w, and let i be its largest string index with nonzero coefficient. Set $a = i + 1$; this string index still exists since e_i has positive grade. The support of v lies at grade at least $\mathrm{gr}(a) + 2$, and the zero target-weight assumptions of Lemma 16.8 hold (the condition on wv is vacuous, while all components of v have positive grade). By Lemmas 16.8 and 16.7, the a-component of T^*Tv is zero. But it is the nonzero coefficient of the shift of v_i. This is a contradiction.

For two positive string indices i, l, apply the lemma to $v = e_i$, $a = l^*$. The target-weight condition on wv is automatic since $\mathrm{gr}(a) < 0$. The support condition on wk follows as all target weights are zero and v has positive grade. We obtain

$$\langle we_l, we_i \rangle = \langle Te_l, Te_i \rangle.$$

The restrictions of w and T are injective (the latter is clear from its nonzero string coefficients), so they give an isometry between the two subspaces of V'. The classical Witt extension theorem over the field E, valid also for degenerate subspaces of a nondegenerate ambient Hermitian space in characteristic zero, extends it to a rational unitary matrix. In particular all the positive string columns can be made equal to those of T, which is stronger than agreement. □

Lemma 16.10 (Stabilizer of agreement)**.** *Suppose w agrees with T, $g \in B_s$. Then hwg agrees with T if and only if $h \in B_{s-1}$.*

Proof. On a positive string column e_i, g sends it to itself plus higher positive string columns. Thus wge_i is $t_i e'_i$ plus a vector of grade at least $\mathrm{gr}(e_i)$: this uses (16.5), whose grade drop is one. If $h \in B_{s-1}$, the degree property of Lemma 16.6 immediately gives agreement. Conversely start from the highest nonnegative target string grade, and proceed downwards over the nonnegative target string indices k. Each e'_k corresponds to a positive source string index k, with grade one greater, and $t_k \neq 0$. By induction, h already fixes every higher target string column modulo still higher grades. Applying the agreement of hwg to the source column corresponding to k, the higher terms therefore cannot contribute at grades at most $\mathrm{gr}(e'_k)$. The coefficient t_k is nonzero, so $he'_k - e'_k$ is supported in grades strictly greater than $\mathrm{gr}(e'_k)$. This is exactly the defining condition of B_{s-1}. □

Proposition 16.11 (The final surviving class)**.** *If w agrees with T and satisfies* (16.6) *for the actual target grades $d = \mathrm{gr}'$, then $w - T$ has degree at least zero.*

Proof. All negative-degree entries in positive source columns already agree. For the remaining entries prove by induction on increasing target grade $r < 0$ that $w_{ba} = T_{ba}$ for source grade $\mathrm{gr}(a) > r$ and target string row b of grade r. (There are no other rows of negative grade.) Use the target reflected row b^* of grade $-r$ and let the source string vector v be the one with grade $-r + 1$ sent by T to te'_{b^*}, $t \neq 0$. By agreement, $wv = te'_{b^*}$ plus terms of grade greater than $-r$.

Fix a source column a with $r < \mathrm{gr}(a) \leq 0$. If it is a string column, put $k = e_a$, and in the Eichler test use the reflected index a^*. If it is in X, use the same calculation with the corresponding norm-one basis vector $k = e_a$. The latter vector also has grade zero, so the proof of the lemma with its quadratic correction applies unchanged. The vector v has grade $-r + 1 \geq -\mathrm{gr}(a) + 2$; every component of wv has target grade at least $-r > -\mathrm{gr}(a)$. By the induction hypothesis, all entries of wk in grades less than r equal those of Tk. Such entries of Tk vanish, since T either kills the column or drops its grade by one. Thus every nonzero component of wk has grade at least r, and its grade plus that of v is at least one. The Eichler test gives $\langle wk, wv \rangle = \langle Tk, Tv \rangle$. Terms of wv above grade $-r$ pair only with grades below r,

where wk vanishes. Thus the equality of pairings forces equality in row b, because t and the paired Hermitian Gram entry are nonzero. This completes the induction. □

These propositions derive the rational-orbit part of the one-step unfolding. Their source anchors are `Grading.exists_agreesWithT`, `Grading.agreesWithT_mul_iff`, and `Grading.hasDegreeGe_of_survives`, all in `Lib/Weil/Periods/UnfoldingStep.lean`. The two uses of survival, at zero weights and at true weights, are different: applying only the first gives too many matrix entries and too large a stabilizer.

16.5 The one-step period transfer

We explain the analytic argument in sufficient detail to state precisely which Weil realization it uses. A rational symplectic vector space in the construction is $\mathrm{Hom}(V, V')$, with alternating form given by the trace and j; it is paired by the reflection of both indices. The maps of strictly positive relative degree form an isotropic subspace. Its annihilator is the subspace of nonnegative degree, because each coordinate of relative degree a pairs with a coordinate of degree $-a$. This continues to hold for target weights d which are odd under reflection. Choose a rational symplectic splitting into these positive coordinates, their dual negative coordinates, and the degree-zero part.

In the resulting mixed Schrödinger model, write $F(x, z)$ for a Schwartz function with x in the negative-coordinate space and z in a remaining Lagrangian of the degree-zero part. The mixed-model theta functional uses the rational sum over both sets of coordinates. The constant term at the rational matrix w, denoted $C_d(f; w)$, is the integral of the theta functional of the translated datum over the adelic positive-coordinate group modulo its rational points. Fourier expansion gives

$$\Theta(f) = \sum_{[w]} C_d(f; w), \tag{16.7}$$

where $[w]$ runs over rational maps modulo nonnegative-degree maps. To verify this, first use $w = 0$: integrating the mixed-model theta sum in the positive variables kills every nonzero dual rational frequency. At general w, its negative-coordinate part shifts that frequency; the coset label is precisely the list of negative coordinates. The degree-zero translation acts by the rational Heisenberg operator and does not change the theta functional, and the nonnegative positive translation changes the phase trivially on a rational w. This proves that $C_d(f; w)$ is constant on the indicated rational coset and establishes (16.7).

If $w - w'$ has strictly positive relative degree (now not necessarily rational), the Heisenberg multiplication law and a change of variable in the compact period integral give

$$C_d(f; w') = \beta(w, w') C_d(f; w). \tag{16.8}$$

For rational elements $g \in B_s$, h in the target parabolic preserving the weight filtration, conjugation of the positive-coordinate integration variables by their action on Hom gives

$$C_d(\omega(g, h) f; w) = C_d(f; h^{-1} w g). \tag{16.9}$$

The quotient measures on additive groups here have total mass one; hence no determinant Jacobian is left by rational substitution. The splitting characters of the rank twists are trivial on rational determinants. The identical formula for adelic grade-raising unipotents on the two sides follows from the mixed-model action: their determinant on the isotropic subspace is one and their remaining diagonal action is trivial. These are the standard mixed-model identities, corresponding to `Grading.constTerm_congr_rat`, `Grading.constTerm_eq_homBi`

`character_mul`, and `Grading.constTerm_mul` in `Lib/Weil/Periods/UnfoldingStep.lean`. They express the needed normalization of the Schrödinger model, not a statement of analytic continuation of a theta series. The finite–archimedean identification of the particular datum of our chain with this realization is a separate matter.

One can also take partial Fourier expansions using different target weights. In particular take zero target weights and actual target grades. In either expansion retain only the terms agreeing with T in the earlier sense. The fixed negative-coordinate positions are precisely those satisfying simultaneously

$$0 < \mathrm{gr}(a), \qquad \mathrm{gr}'(b) < \mathrm{gr}(a).$$

Those positions occur in both sets of negative coordinates. Their dual positive-coordinate positions form a common isotropic subgroup of the two choices. Summing over all other negative coordinates in either model is the same as averaging the theta functional with the character prescribed by T only on this common isotropic subgroup, by ordinary Fourier inversion on the compact rational quotient. This proves that the two retained partial sums agree; the observation is the content of `Grading.tsum_constTerm_agreesWithT`. It is often a useful shortcut to avoid choosing a model in which all Fourier coordinates are written out at once.

Let P be a continuous function invariant under $B_s(E^+)$ and transforming under degree-at-least-two unipotents by

$$P(ug) = \beta(Tu, T)P(g) = \psi\left(\tfrac{j}{2}\,\mathrm{tr}(S_s u)\right) P(g).$$

It is called generic here only relative to this specific string and character. The theta kernel is evaluated on a fixed datum Φ by the twisted Weil action $\omega(g, h)$. Define

$$P'_x(h) = \int_{[B_s]} P(rx)\, C_{\mathrm{gr}'}(\omega(rx, h)\Phi; T)\, dr, \qquad [B_s] = B_s(E^+)\backslash B_s(\mathbb{A}_{E^+}). \tag{16.10}$$

This is the function `Grading.unfoldStep`.

Theorem 16.12 (One step of the global period unfolding)**.** *In the normalized mixed model just described, let y be a continuous rapidly decreasing automorphic function on $U(V')$. With the compatible quotient measures of the theta integral and subgroup periods, the identity is*

$$\begin{aligned}\int_{[B_s]} P(rx) \left(\int_{[U(V')]} \Theta(\omega(rx, h)\Phi) y(h)\, dh \right) dr \\ = \int_{B_{s-1}(E^+)\backslash U(V')(\mathbb{A}_{E^+})} P'_x(h) y(h)\, dh.\end{aligned} \tag{16.11}$$

The unfolded function is generic for the next string and its own shift.

Proof. Use the expansion (16.7) first at zero target weights. Partition its rational labels into orbits of $B_s(E^+)$. For a particular label w with failure of the survival condition choose a degree-at-least-two unipotent u as in (16.6) for which the ratio of the two characters is not one. The integral of that label, after unfolding its rational orbit, is an integral on the quotient by its rational stabilizer. Left translation by u multiplies the integrand by

$$\beta(Tu, T)/\beta(wu, w)$$

by genericity and (16.8)–(16.9). One must justify this step, since left multiplication by an arbitrary adelic element need not be a map on a rational stabilizer quotient. The following

general argument does so. For a continuous, nonzero, left-Γ-invariant integrand F on a locally compact group G, let H consist of the elements h for which $F(hz) = \chi(h)F(z)$ for some $|\chi(h)| = 1$ and all z. By evaluating at a point where F is nonzero, H is seen to be a closed subgroup with continuous unitary character χ; it contains Γ. The orbit quotient integral is absolutely integrable. Disintegrate it in stages, first on $\Gamma\backslash H$ with left Haar measure, then on $H\backslash G$ with the compatible quotient density. On each fibre the map $h \mapsto hu$ is well defined on $\Gamma\backslash H$ and scales its measure by a positive real modular factor. But it scales the integrand by $\chi(u)$ because $F(hux) = \chi(h)\chi(u)F(x)$. As $\chi(u) \neq 1$ and has absolute value one, it cannot equal that positive modular factor. The fibre integral is zero, hence so is the total integral. This proves the required vanishing without a normalizer assumption on u. It is the quotient-measure argument of `integral_quotMeasure_eq_zero_of_twist` in `Lib/Adelic/QuotientMeasure/CocompactIntegrable.lean`.

By Proposition 16.9, the remaining rational labels are carried to the set of labels agreeing with T by rational matrices of the target unitary group. Lemma 16.10 identifies the stabilizer within that group as the rational Bessel subgroup. Unfold the rational cosets of the target automorphic integral: a sum over translates indexed by the rational Bessel subgroup quotient of the rational target group converts the integral on $[U(V')]$ to the integral on $B_{s-1}(E^+)\backslash U(V')(\mathbb{A}_{E^+})$. After this unfolding it remains to sum the agreed labels at zero target weights. Replace the partial Fourier sum by the equal partial sum at actual target grades, as explained above. The same unipotent character argument applies to that partial sum. Proposition 16.11 shows that its sole surviving class is T modulo the nonnegative-degree subgroup. This is exactly the integrand of (16.10).

The invariance of P'_x under rational B_{s-1} follows from (16.9) and Lemma 16.7. For a degree-at-least-two unipotent v of the target, $vT - T$ has positive relative degree. The phase of (16.8) for that change is the character of the next shift: the identity $TT^* = S_{s-1}$ and cyclicity of trace give $\psi(j\,\mathrm{tr}(S_{s-1}v)/2)$. This is next-stage genericity.

The integrals and sums involved are absolutely convergent under the usual Schwartz-theta growth estimates and rapid decay stated here. Indeed a partial Fourier transform takes Schwartz functions to Schwartz functions, and their absolute rational sums converge uniformly for data in bounded sets of Schwartz seminorms, since the finite support restricts to a lattice of bounded denominators and archimedean Schwartz decay beats its polynomial counting bound. One may use a compact set of representatives for $[B_s]$ when bounding the factor depending on r. On the remaining quotient, the rational theta sums, and the sum of absolute values of their partial Fourier coefficients, are bounded by a power of the adelic reduction height of h, by the standard moderate-growth estimate for the Weil representation on Schwartz seminorms. Multiplication by a rapidly decreasing y makes this integrable using reduction theory for unitary groups. These are the same analytic hypotheses as the single-step result of B. Wang Peng Jun, *Global theta lifting and automorphic periods associated to nilpotent orbits*, J. Number Theory 271 (2025), 122–149, Proposition 3.2 and Theorem 4.5, in its unitary hook setting. The proof uses this classical moderate-growth input rather than any quantitative bound in the varying parameters of GRH. The estimates justify the changes of order made above. For a purely algebraic finite-place test function the corresponding summations are finite already in the coefficient variables. □

This is the structure of `Grading.integral_besselGroup_unfold` and `isGeneric_unfoldStep`. The geometric orbit computation, including the Witt step and the single surviving class, has been reconstructed above. The theorem uses the ordinary normalized mixed-model identities and its qualitative moderate-growth theorem. The quantitative uniformity in the upper bound will instead come from the integral coset cuts below. In particular, the one-step period identity alone does not imply the iterated seesaw with its good finite datum.

16.6 The glued lattice and integral Witt extension

Here is the algebra of the lattices used in that quantitative step. Write $\widehat{O} = \prod_w O_{E,w}$ for the integral finite adeles. A finite adelic lattice of full rank is a matrix translate of $\widehat{O}^n$ in rational coordinates; its dual is taken with respect to the given Hermitian form. The field in the construction contains a primitive 5040th root of unity, the field hypothesis named `StablyUnram`. In particular the quadratic CM involution is unramified or split locally: if it were a nontrivial inertia element at a place above a rational prime p, it would be trivial on roots of unity of order prime to p. One can choose an odd prime dividing 5040 and different from p, whose primitive root is not fixed by complex conjugation. This contradiction proves the assertion. Norm-one units and matrices should nevertheless be treated separately at a split pair of places; the involution there exchanges the two factors rather than acting within either one.

Choose a finite idele π with $\pi\bar{\pi} = N$, supported with valuation one at precisely one place over each split pair above N. For a finite idele a and a unitary idele u define the basis $B_s(a, \pi, u)$ as follows. If s is even it is the identity on X and multiplies the first half of the string by u/a and the second half by $u\bar{a}$. If s is odd it multiplies every string vector by u, and, writing e for the last vector in X and z for the middle string vector, replaces e by

$$\bar{\pi}^{-1}e + uz/N.$$

It does not change the other X-vectors. This is the matrix `gradedLatticeBasis` in `New/Seesaw/LatticeAndTorsionDefs.lean`.

Lemma 16.13 (The glued basis)**.** *The Gram matrix of this basis is integral and unimodular. Its string diagonal entries in the upper triangular flag order are u, for odd s, and $u/a, u\bar{a}$ on the respective halves for even s.*

Proof. In the even case every pair of reflected string vectors has pairing $\overline{u/a}u\bar{a} = 1$, just as before. In the odd case the only changed Gram block is the pair consisting of the new X-vector and the new middle string vector. Their Gram matrix is

$$\begin{pmatrix} 0 & -1 \\ -1 & -N \end{pmatrix}.$$

Indeed the new X-vector has norm $N^{-1} - N/N^2 = 0$, and its pairing with uz is -1. This block has determinant -1 and integral inverse. All other Gram blocks are unchanged. The described diagonal entries follow directly from the basis. For the flag order put all the positive strings first, then the middle string if present, then the X-block, then the negative strings; the added middle entry in the last X-column lies earlier in the flag order. □

At a nonsplit unramified place a unimodular Hermitian lattice admits an orthonormal basis. To see this, some diagonal value of its reduction is nonzero, or a pair of off-diagonal entries is nonzero and an appropriate linear combination of the two basis vectors has unit norm, by surjectivity of the residue trace. Split off a unit line and induct. Rescale the resulting unit diagonal entries by norms of local units, using the surjectivity of the norm of an unramified quadratic extension on units. At a split pair the two factors are in perfect duality, so choose a basis on one and the dual basis on the other. Taking these choices at all places gives a finite adelic frame x_f with $x_f^* J_s x_f = 1$ spanning the glued lattice. At an archimedean place the signature of J_s has exactly $\lfloor (s+1)/2 \rfloor$ negative directions. The reflected pairs are diagonalized by sum and difference divided by $\sqrt{2}$, and the middle vector, if present, is rescaled by $N^{-1/2}$. This gives a real frame of the prescribed signs; for $N = 1$ its matrix in the original

coordinates can be chosen unitary for the Euclidean inner product. These are the existence statements of `isUnimodular_gradedLatticeBasis_gram` and `exists_frame_gradedGram` in `New/Seesaw/FramesForY.lean`. They apply to the intermediate spaces as well as the endpoints.

Lemma 16.14 (An integral Hermitian extension lemma)**.** *Suppose F and F' are n by r matrices over $\widehat{O}$ whose columns admit integral left inverses, and $F^*F = (F')^*F'$ for the standard finite Hermitian form. There exists an integral unitary h with $hF = F'$.*

Proof. We give a local proof, since it clarifies the need for the unramified condition at residue characteristic two. In the nonsplit case the residual extension is quadratic over finite fields. The reductions of both frames are injective and define an isometry between their subspaces, including possibly degenerate subspaces. The Hermitian Witt extension theorem over finite fields gives a residual unitary carrying one to the other. It is enough to lift it to an exact unitary with the same prescribed frame action.

Use a uniformizer ϖ_+ of the fixed field, also a uniformizer of the extension. Suppose a matrix h_n has $h_nF \equiv F'$ and $h_n^*h_n \equiv 1 \pmod{\varpi_+^n}$. Let $\Delta = (F' - h_nF)/\varpi_+^n$ and $D = (1 - h_n^*h_n)/\varpi_+^n$, read modulo the uniformizer. We seek $h_{n+1} = h_n(1 + \varpi_+^n A)$ with

$$A\bar{F} = \bar{h}_n^{-1}\bar{\Delta}, qquad A + A^* = \bar{D}.$$

The compatibility condition is

$$\bar{F}^*\bar{D}\bar{F} = \bar{F}^*(\bar{h}_n^{-1}\bar{\Delta}) + (\bar{h}_n^{-1}\bar{\Delta})^*\bar{F},$$

which follows by expanding equality of the two Gram matrices mod ϖ_+^{n+1}. Pick a residual scalar τ with $\tau + \bar{\tau} = 1$ by surjectivity of the trace (this is valid even in characteristic two). Then $A_0 = \tau\bar{D}$ has the required symmetrization. The remaining prescribed action of $A - A_0$ on the column space of $\bar{F}$ is skew-adjoint there by the displayed compatibility. Such a partial skew-adjoint map extends to the entire ambient space: choose any linear complement to the column space and define the corresponding sesquilinear form on pairs of column-space and complementary vectors by the prescribed map and skew symmetry; on the two complementary variables take it to be zero. The nondegenerate ambient form identifies this extended form with a skew-adjoint endomorphism. Thus an A exists. Lift its entries arbitrarily and iterate; completeness gives the exact unitary matching the two frames.

At a split pair the module is a pair of free modules R^n, R^n on which the involution exchanges factors. The residual action is an arbitrary invertible g on one and g^{-t} on the other. With reduced frames $F = (P, Q)$ and $F' = (P', Q')$, equality of Gram means $Q^tP = (Q')^tP'$. We need $gP = P'$ and $(Q')^tg = Q^t$. Here P, P' are injections and $Q^t, (Q')^t$ are surjections. The prescribed map between the images of P, P' identifies their intersections with the kernels of the respective surjections. Extend that identification to an isomorphism between the two full kernels; choose representatives of the remaining quotient vectors compatible with the common image of Q^tP in the quotient. This gives an invertible g with both properties. The same successive lifting argument works over the split ring, now using trace-one element $(1, 0)$ and complementary bases in the two factors (the linear compatibility is the equality of the prescribed paired values). Alternatively one can make the kernel and quotient basis choices integrally by taking primitive frames and matching the common elementary divisors of Q^tP. This proves the local assertion at every place. All the resulting h are integral, so their product is a finite-adelic unitary element. □

In this proof the residual use of the Hermitian field Witt theorem is elementary classical linear algebra. The lifting of the frames is essential; reducing a rational-unitary Witt matrix

modulo a prime would not prove the required existence of an integral one. This is a conventional proof of `witt_Ohat_of_orthonormal` in its more general primitive form the integral Witt lemma for $\widehat{O}$ in `Lib/Adelic/Witt/SpaceOperations.lean`; we will refer to the latter also as the integral Witt extension lemma.

There is a further useful congruence inclusion. Embed $U(X)$ by fixing all string vectors. If s is even, every integral unitary on X preserves the glued lattice. In the odd case suppose the last row of the matrix h on X is congruent to the last row of the identity modulo π. The inverse satisfies the same row congruence (multiply $(h-1)$ on the right by h^{-1}), and by unitarity this is the congruence of the last column of h modulo $\bar{\pi}$. In the odd glued basis the X-part has the last coordinate scaled by $\bar{\pi}^{-1}$; the Hermitian conjugate row is thus scaled by π^{-1}. For the endomorphism $h-1$ on V, all the entries of

$$B_s^* J_s (h-1) B_s$$

are then integral: the only one in which both denominators occur is the last diagonal X-entry, divisible by $\pi\bar{\pi} = N$ since these ideals are coprime at each local place. The contributions involving the extra string coordinate vanish as $h-1$ kills the string. Because $B_s^* J_s B_s$ is unimodular, $B_s^{-1} h B_s$ and its inverse are integral. This proves the inclusion of the subgroup fixing the last row modulo π into the lattice stabilizer. Restricting further at the chosen primes gives the subgroup `Data.columnLevel`. It proves the assertion of `Grading.inclXFin_mem_Kf` in `New/Seesaw/StandardLatticeLevel.lean`.

16.7 Adapted bases and equality of lattice lines

Use the flag order used above, grouping all indices of X into a single block. If $L_x = x_f x \widehat{O}^n$ is the lattice associated to $x \in U(V)(\mathbb{A}_f)$ in framed coordinates, an adapted basis B is a matrix of generators upper block triangular in this flag order. Its string diagonal entries are finite ideles β_i. Such a basis exists: locally intersect the lattice with every subspace of the ordered flag. These intersections are saturated in the lattice, hence the successive quotients are free over the discrete valuation ring. Choose and lift bases of these quotients; this gives an integral basis adapted to the flag. At almost all places the lattice is already standard, so one obtains a compatible finite-adelic basis. In particular $B^{-1} x_f x$ is integral invertible. The valuation of β_i is independent of the adapted basis, since the change of bases is triangular and integral invertible, and string diagonal entries multiply under triangular products. More intrinsically it is the valuation of the rank-one lattice induced on the ith string graded piece. If $L_{x'} \subset L_x$ then $\operatorname{ord}_w \beta_i' \geq \operatorname{ord}_w \beta_i$ (inclusions induce inclusions on each flag quotient); for equality of local lattices the valuations agree. The source uses ideleLogVal $= -\operatorname{ord}_w$.

Let $h_i = 1$ except for the middle string entry of odd length, where $h_i = -N$. The following formula controls the Gram of the lifted flag:

$$(B^* J_s B)_{ij} = h_i \bar{\beta}_i \beta_j \, [j = s-1-i], \qquad j \leq s-1-i \tag{16.12}$$

for string indices i, j. Indeed in the product the ith column is supported only in string indices $\leq i$ and possibly in X if i lies later in the flag. If $i + j < s-1$ no paired string indices can contribute; if $i + j = s-1$ only the diagonal pair contributes. At least one of the two indices in the displayed range lies in the nonnegative-grade string and therefore has no X-component. This proves the formula even when the other string vector lies after the X-block.

Proposition 16.15 (Bessel double cosets determined by the nonnegative lines)**.** *If two such lattices $L_x, L_{x'}$ have equal string diagonal valuations at all places on the nonnegative-grade string, then $x' k x^{-1}$ belongs to the finite Bessel subgroup for some framed integral unitary k.*

Proof. Rescale the corresponding adapted basis columns of B' by integral units to make their diagonal ideles exactly those of B. This is possible because equality of valuations means the ratios of diagonals are integral idelic units. The matrices $F = (x_f x)^{-1}B$ and $F' = (x_f x')^{-1}B'$ are integral invertible. Retain their columns from the nonnegative string. The resulting two frames have integral left inverses and their Gram matrices coincide by (16.12), since the finite frame identifies J_s with 1 and x, x' are unitary.

Use Lemma 16.14 to send one frame to the other by an integral unitary k. The matrix of $x'kx^{-1}$ in rational coordinates consequently sends every retained column of B to the corresponding column of B'. By the flag-triangular form with the same diagonal, it fixes each of these standard basis vectors modulo higher string grades. One sees this inductively along the columns: earlier string columns have already been treated and the coefficient of the current line is the same. Lemma 16.6 identifies the resulting element as a member of the Bessel subgroup. □

The source anchors are `Grading.exists_isAdaptedAt`, `Grading.IsAdaptedAt.gram_string_entry`, `Grading.IsAdaptedBasis.ideleLogVal_le`, and `Grading.IsAdaptedAt.exists_mem_besselFin` in `New/Seesaw/EqualLinesOrbit.lean`. This proposition is stronger than a mere compactness criterion: it gives a representative for the double coset from just finitely many string-line valuations, not from all the off-diagonal entries of an arbitrary basis.

16.8 Local supports of zero terms

We recall the finite-coordinate normalization used in the support estimate. The product finite lattice $O^n = \widehat{O}^n$ is self-dual for the additive pairing $\psi_0(\bar{x}^t y)$ of the Heisenberg representation in its standard finite coordinates. This means that if $\psi_0(\bar{x}^t y) = 1$ for every integral x, then $y \in O^n$; products over places reduce the assertion to self-duality for the local additive character of conductor O_w. The frame of each graded spaces is related to rational coordinates by

$$w \longmapsto \alpha^{-1} x_f'^{-1} w x_f$$

on the matrix Hom space (in vector notation). The finite idele α incorporates the different and character normalizations; see Chapter 6. For the following support lemma all finite vectors are already in these normalized coordinates.

Let L_0 denote the finite part of a zero-term map along an isotropic frame k in the source space of the finite Heisenberg representation. We use its standard mixed-model property $L_0\rho(kz) = L_0$ for all adelic z. Denote by M^* the conjugate transpose in finite coordinates.

Lemma 16.16 (A support criterion from finite translation invariance)**.** *Let M, M' be matrices with $M(M')^* = 1$, and let the finite datum f be fixed by $\rho(Mx)$ for all $x \in O^n$. If $L_0\rho(y)f \neq 0$, then there is $\ell \in O^n$ with*

$$k^*(y - M'\ell) = 0.$$

Proof. The matrix $(M')^*k$ is a full-rank matrix over the finite adeles. Choose an invertible column matrix D such that $X = (M')^*kD$ is an integral primitive frame. Locally this is just a basis of the saturated lattice in the column space, and almost everywhere the given frame is already integral. For any integral vector z, one has $MXz = kDz$. The operator $\rho(kDz)$ fixes f by the assumed invariance, and disappears after applying L_0. Commuting it past $\rho(y)$, and using the finite Heisenberg relation, gives

$$L_0\rho(y)f = \psi_0((kDz)^*y)L_0\rho(y)f.$$

Since the displayed vector is nonzero, the phase is one for every $z \in O^r$. But $kD = MX$, so this says $X^*M^*y \in O^r$ by self-duality. The primitive frame X has an integral left inverse; hence its adjoint maps O^n onto O^r. Choose $\ell \in O^n$ with $X^*\ell = X^*M^*y$. Since $X^* = D^*k^*M'$ and $M'M^* = 1$, the equality is $D^*k^*(y - M'\ell) = 0$. The matrix D is invertible, giving the claim. □

This is exactly `zeroTermFin_support` in `New/Seesaw/EdgePair.lean`. A zero term of a finite datum need not vanish whenever its translated argument is nonintegral: only its coordinates dual to the isotropic frame are tested. Replacing this criterion by the stronger and false full integrality condition would destroy the sharp estimates.

There is also a zero-term formula for the vacuum. Write i for the finite embedding of the degree-zero part, and let $B = ig + kc$ be an integral matrix, where g is unitary on the degree-zero part and c is arbitrary. The vacuum v_0 on this part is determined up to scalar by invariance under the standard integral finite shifts (one-dimensionality of its fixed line). The vector $\omega(g)^{-1}L_0v_{\mathrm{vac}}$ is on that line, by the mixed-model equivariance and the integrality of B: translating by $iz + kcg^{-1}z$ has the same zero-term action as translating by iz, and the first vector fixes the incoming vacuum when z is integral in the transformed coordinates. Hence $L_0v_{\mathrm{vac}} = C\omega(g)v_0$ for some scalar C. For a general translate $\rho(y)v_{\mathrm{vac}}$ the support lemma with $M = M' = 1$ supplies $\ell \in O^n$ with $y - \ell$ orthogonal to k if the result is nonzero. Decompose this difference as $iw + kb$ in the mixed-model splitting. Its zero term is the translate of L_0v_{vac} by w. Because $B^*(y - \ell) = g^*w$, and $B^*\ell$ is integral, the finite shifts and their commutation phases now give

$$L_0\rho(y)v_{\mathrm{vac}} = C\zeta\,\omega(g)\rho(B^*y)v_0, \qquad \zeta = 0 \text{ or a root of unity.} \tag{16.13}$$

The root of unity is the product of a finite number of values of the finite additive character and semicharacter. In particular its magnitude after any algebraic conjugation is one when nonzero, and it is integral at all finite places. The precise source is `zeroTermFin_finHeisOp_vacElem` in `New/Seesaw/EdgePair.lean`. This proof uses the fixed vacuum line and the elementary mixed-model covariance, but not a computation of the real vector in the zero term.

16.9 A one-step bound for the lattice lines

Let the old adapted basis have string diagonals β_i and the next one β'_i. Let Q be the chosen split places of the twist conductor (one from each conjugate pair occurring), and write

$$A_w = \mathrm{ord}_w\, a, qquad D_w = A_w - A_{\bar{w}}.$$

For an integer n define the baseline for a string by

$$B_{s,n,i}(w) = nD_w + \begin{cases} 0, & s \text{ odd}, \\ -A_w, & s \text{ even},\ i < s/2, \\ A_{\bar{w}}, & s \text{ even},\ i \geq s/2. \end{cases} \tag{16.14}$$

These are the additive valuations of the diagonals of $B_s(a, \pi, (a/\bar{a})^n)$. In particular

$$B_{s,n,i}(w) + B_{s,n,s-1-i}(\bar{w}) = 0. \tag{16.15}$$

The condition called `Grading.HasNearLines` is that the difference between the actual valuation and (16.14) is bounded by k above $Q \cup \bar{Q}$ and is zero everywhere else. At stage zero it follows immediately when nonzero support lies in the Bessel orbit of the standard lattice: multiplying its adapted basis by a Bessel element leaves all diagonal entries unchanged. This is the base case `Grading.hasNearLines_zero` in `New/Seesaw/StandardLatticeLevel.lean`; the first Poincaré datum supplies the needed support condition.

Lemma 16.17 (Inequality extracted from a nonzero constant term)**.** *Let f be a finite datum invariant by all finite shifts with parameter $b\widehat{O}^n$, with b a finite idele. Suppose the constant term at T of its Weil translate at two adjacent lattice points is nonzero. Then, with $s' = s - 1$ and $0 \le i < s'$, one has for every place w*

$$\operatorname{ord}_w \beta'_i \le \epsilon_i \operatorname{ord}_w c + \operatorname{ord}_{\bar w} b - \operatorname{ord}_w \alpha - \operatorname{ord}_{\bar w} \beta_{s-1-i}, \tag{16.16}$$

$$\epsilon_i = \begin{cases} 1, & s' \text{ odd and } i \le (s'-1)/2, \\ 1, & s' \text{ even and } i > (s'-1)/2, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Use the finite support lemma, with the finite unitary action of the pair of lattice points and with $M = bg$, $M' = \bar b^{-1} g$. The product $M(M')^*$ is the identity. The isotropic frame consists of the strictly positive relative degree coordinates, so its annihilator imposes that the negative-degree part of the rational frequency agrees with the negative-degree part of an integral matrix in the translated frames. On returning to the rational Hom coordinates this gives a matrix W satisfying

$$\deg(W - T) \ge 0, qquad \alpha^{-1}\bar b\,(x'_f x')^{-1} W x_f x \in \mathrm{Mat}(\widehat{O}).$$

The factor α^{-1} is exactly the inverse scale in the finite Hom coordinates used in Lemma 16.16. The involution on b comes from $M' = \bar b^{-1} g$. Replacing either frame by an adapted basis does not change integrality. Thus

$$\alpha^{-1}\bar b\,(B')^{-1} W B \in \mathrm{Mat}(\widehat{O}).$$

Look at row i, column i in the string-to-string block. Since the source column has grade one more than this target row, the equality $W - T$ of degree at least zero gives $W_{ii} = t_i$. The matrices B and $(B')^{-1}$ are upper triangular for the flag. In the product $(B')^{-1} W B$, the contributing intermediate source grades are at least the grade of e_i and the intervening target grades are at most that of e'_i; this forces all such terms to lie in the negative-degree part of W, where it equals T. The only possible diagonal string contribution is then at row i, column i. More exactly, the integral dual basis of the lattice B is $J_s^{-1}(B^{-1})^*$; its diagonal at the ith string coordinate is $h_i^{-1}\bar\beta_{s-1-i}^{-1}$ with h_i the paired Gram coefficient. As the lattice is self-dual, we may use this basis for the domain. This gives the required integral scalar

$$\frac{\bar b}{\alpha}\,\frac{t_i}{h_i \beta'_i \bar\beta_{s-1-i}}. \tag{16.17}$$

In the odd source case the coefficient t_i/h_i equals 1 on the first half and c on the second half including the old middle index (the two factors $-N$ cancel). For even source it is c on the first half and 1 on the second. In both cases this is c^{ϵ_i}. Taking additive valuations of (16.17) proves (16.16). □

For clarity about the middle index in this calculation, the dual basis may be taken with its reflected column order, namely the columns of $J_s^{-1}(B^{-1})^*$ indexed by reflected strings. This is again upper triangular in the original string order by Hermitian reflection. Both these bases generate the same lattice, since $B^* J_s B$ is integral unimodular. Formula (16.17) is the scalar calculated in `Grading.HasNearLines.step` of `New/Seesaw/LinesStep.lean`.

Proposition 16.18 (The velocity of adapted lines)**.** *Assume that the source lattice satisfies the near-line condition with error k and exponent n, that $n' = n$ when s is odd and $n' = n - 1$ when s is even. Assume also*

$$\operatorname{ord}_w c = A_w + A_{\bar w}, \qquad \operatorname{ord}_w b \le \operatorname{ord}_{\bar w}(\alpha/a) + [w \in Q],$$

and suppose the next lattice has

$$\operatorname{ord}_w \beta'_i + \operatorname{ord}_{\bar{w}} \beta'_{s'-1-i} \geq 0. \tag{16.18}$$

Then a nonzero constant term at T forces the next lattice to satisfy the near-line condition with exponent n' and error $k+1$.

Proof. In (16.16) substitute the bound on the valuation of b and the near-line condition at the reflected index of the source. Using (16.15) gives

$$\operatorname{ord}_w \beta'_i \leq B_{s,n,i}(w) + \epsilon_i(A_w + A_{\bar{w}}) - A_w + (k+1)[w \in Q \cup \bar{Q}].$$

The first three terms are exactly $B_{s',n',i}(w)$. Indeed, for odd s the subtraction changes the first half of the next baseline by $-A_w$ and the second by $A_{\bar{w}}$, while $n' = n$. For even s the old half contributions cancel with the relevant two terms to give $(n-1)D_w$ in both halves. This proves the upper bound. Apply the same upper bound at the reflected index and conjugate place, then use (16.18) and (16.15) to obtain the opposite bound. Off the two supports all errors vanish. □

Most of (16.18) follows from (16.12): the Gram on an adapted basis is integral and a reflected pair has coefficient 1. The only apparent exception is the middle string index for odd length at a place dividing N, where integrality yields just the lower bound -1. The source eliminates this exceptional case by using ramified columns, rather than accepting a step at the level prime.

For the actual modulus in the construction, $a = \alpha/\bar{\lambda}$ where λ is the chosen finite idele generating the chosen-prime ideal, and b generates that ideal times the product of the places in Q. Therefore the valuation of b is exactly the valuation of $\overline{\alpha/a} = \lambda$ plus $[w \in Q]$. This is the inequality used in Proposition 16.18. Moreover the finite residue datum is a sum of translated vacua with translation parameters t satisfying $\bar{b}t \in \widehat{O}^h$. Each such vector is fixed by shifts $\rho_f(bx)$, x integral: the vacuum is fixed since b is integral, and the commutation phase with the translate is trivial by integrality of $\bar{b}t$ and self-duality of the finite pairing. The same holds for the sum. Thus the invariant-shift hypothesis of the finite support lemma is compatible with the level modulus; it does not require the translate itself to be an integral vector. These are the idele and modulus calculations of `New/Trace/GoodIdempotent.lean` and the residue expansion used in `hasNearLines_of_unfoldStage_ne_zero`.

Lemma 16.19 (The exceptional middle line)**.** *For s odd, if the middle pair sum of valuations equals -1 at a split place w of N, there is a representative in the same Bessel–integral-unitary double coset whose lattice admits the embedded full local maximal compact subgroup of $U(X)$ at w. Consequently, if a nonzero generic stage function has the property that it vanishes on all such representatives, then* (16.18) *holds.*

Proof. At the pair $(w, \bar{w})$ write the lattice in one local factor; its other local factor is the dual under the Hermitian form. Form a diagonal lattice at w as follows. On the positive and middle string lines, choose the valuations of the given adapted basis at w. For a negative string index j, choose the negative of the original valuation on its positive reflected partner at $\bar{w}$. Take the identity lattice on X. The dual at $\bar{w}$ has the correct valuations on the nonnegative string by these choices; for the middle string its valuation is minus the valuation at w minus one, since that Gram entry has valuation one. By the assumed defect this is exactly the original middle value at $\bar{w}$. Elsewhere keep the old lattice. This specifies a valid self-dual lattice, hence a point in the finite unitary group (use local dual bases at the split pair). Now use Proposition 16.15, since the nonnegative lines agree. The diagonal lattice admits the claimed local compact subgroup by its identity lattice on X. The asserted vanishing condition excludes this possibility. □

The ramification condition on the initial column is precisely designed to obtain the vanishing used in the lemma. At an even stage, an unfolded value is a coefficient of a paired definite lift. The action of a local scalar z at a place of N on the input column is intertwined with its action on this coefficient by the pair Weil equivariance; the relevant splitting twists are unramified there. Every ramified column lies in the image of $1-\rho(z)$ for each such local generator. If the output test value is invariant under z because the full local compact is contained in the stabilizer, it vanishes on this image. Thus, once the even-stage coefficient identity and its equivariance have been established, Lemma 16.19 applies at the odd-length stages. This is the exact use of `exists_eq_sub_centralGen`, `scalarFin_centralGen_mem` in `New/Columns/ChainValuation.lean` and `hasNearLines_of_unfoldStage_ne_zero` in `New/Seesaw/UnfoldingStages.lean`. The geometric part of the implication is proved here; the particular even-stage coefficient identity is discussed below.

16.10 Finite cuts with prime-supported multiplicity

There is an entirely group-theoretic argument which explains why this bound on lines is enough. Let H be a locally compact group, $J \subset A \subset B$ subgroups, where J, A are compact open, and let S be a closed subgroup. Let D be a compact subset of B such that for every $d \in D$ there is an integer $n_d \geq 0$ with

$$(dud^{-1})^{q^{n_d}} \in A \quad (u \in J). \tag{16.19}$$

Let a desired support $W \subset H$ satisfy $W \subset SD$. For a finite subset T of H/J stable under $S \cap A$ define

$$\mathrm{mult}_T(xJ) = \#\{r(S \cap A) \in S/(S \cap A) : r^{-1}xJ \in T\}.$$

It is constant on S-orbits.

Theorem 16.20 (Finite cut lemma)**.** *Under the preceding hypotheses there is a finite $S \cap A$-stable $T \subset B/J$ such that W/J is covered by ST, and the multiplicity on T and on W/J is nonzero and divides a power of q.*

Proof. The image of D in the discrete set H/J is finite. Choose one representative in this image of each distinct S-orbit meeting it. Call the resulting finite set C, and take $T = (S \cap A)C$. This is finite because the image of a compact group on a discrete set is finite. It lies in B/J and is stable as required. Every point of W/J is in the S orbit of one of the chosen representatives.

For a representative $dJ \in C$, put $U = S \cap dJd^{-1}$. Its multiplicity is exactly

$$[U : U \cap A].$$

Indeed the only points of T in that S-orbit are $(S \cap A)dJ$. The elements r with $r^{-1}dJ \in T$ form $U(S \cap A)$, and passage to right cosets of $S \cap A$ gives the asserted index. It is finite because U is compact and A open. Let V be the normal core of $U \cap A$ in U. The target index divides the order of the finite group U/V. By (16.19), for every $v \in U$ its q^{n_d}th power belongs to $A \cap U$, and the same is true for every conjugate of v in U. Thus $v^{q^{n_d}} \in V$. All element orders in U/V divide a power of q. By Cauchy's theorem every prime factor of its group order is a prime factor of q; hence its order divides some power of q. This proves the assertion, including the case $q = 1$ when the multiplicities must all equal one. □

This is `exists_cut_coverMultiplicity_dvd` in `New/Seesaw/IntegralAsCosetSum.lean`. Notice that $U \cap A$ was not assumed normal. Passing directly to a quotient group

by it rather than its core would invalidate the argument. The same cut gives a partition of unity along the S-orbits with their Haar measure. Normalize Haar measure dr of S by $\mathrm{vol}(S \cap A) = 1$, and define

$$\phi_T(x) = \frac{\mathbf{1}_T(xJ)}{\mathrm{mult}_T(xJ)}$$

with zero value off T. Then

$$\int_S \phi_T(r^{-1}x)\, dr = 1 \quad (x \in W). \tag{16.20}$$

Indeed the integrand is constant of value $1/\mathrm{mult}_T(xJ)$ on a union of exactly that many disjoint left cosets of $S \cap A$; invariance of the multiplicity along S was used. The formula requires no assertion of normality of the small level in H. It is this partition of unity, not counting of representatives without a weight, that enters the quotient periods.

Construction of the required compact set. For stage $k > 0$ let W be the support constrained by the near-line bound. At each place outside $Q \cup \bar{Q}$, the adapted valuations of any such point equal the fixed baseline, which is the valuation pattern of $x = 1$. At the remaining finitely many places they lie in fixed intervals of length $2k + 1$. The number of possible tuples of nonnegative string valuations at those places is therefore finite. For each tuple that actually occurs, choose one finite group point having that tuple, and replace its components outside $Q \cup \bar{Q}$ by the identity. This operation preserves the unitary condition, since the indicated set of places is stable under the involution. The adapted valuations at a given place depend only on the local point, not on choices elsewhere: they are the valuations of the graded flag quotients of the local lattice. Thus the modified point has the desired tuple at those places and the baseline elsewhere. By Proposition 16.15, every element of W belongs to the Bessel orbit of one of these finitely many points times K_0. Taking D to be their union of right K_0-cosets gives a compact set integral away from q. In particular this argument does not require a diagonal representative on the middle string line at a twist prime; such a representative need not have the two desired valuations. When the twist is trivial one simply takes $D = K_0$. For stage zero the initial Poincaré support lies in the S-orbit of the chosen level above g, so take that level as D and A.

Above a twist prime the subgroup J at that stage belongs to the first principal congruence group in the local lattice coordinates, which is pro-p at a residue characteristic p dividing q. For every fixed representative d, conjugation of sufficiently deep congruence subgroups by d is contained in the local maximal compact A; the finite quotient of the first principal group by any such deeper congruence subgroup is a finite p-group. Therefore some power of q as in (16.19) sends all elements at these places into the desired subgroup. Away from the twist primes no power is needed by integrality and the choice of A (the first stage has A equal to the chosen level and all other stages can use the maximal compact). At the first stage $d \in A$, $J \subset A$, and the condition holds even with exponent one. For stage $k > 0$, the embedding of the indicated column subgroup in A was proved above using the glued lattice. This supplies the compact set and prime support conditions of `Data.exists_stage_cut` from the near-line estimate, including the first-stage support and equality-of-line arguments. There is no need to bound the number of its cosets uniformly for nontrivial twists: only finiteness and prime support matter.

The measure factor. Suppose an integrable stage product $F = Py$ on the rational Bessel quotient of the ambient group is supported in W in its finite variable. In the compatible quotient disintegration, insert the partition of unity (16.20). This replaces integration over

the ambient finite quotient of the Bessel group by integration over the finite cut, with weight ϕ_T. Suppose the periods over the Bessel group on the cut factor as

$$\int_{[B_s]} F(r(x_\infty, x_f))\, dr = [K_0 : J], a(x_\infty) b(x_f J).$$

The compact-open K_0 is normalized to have mass one. Every J-coset has measure $[K_0 : J]^{-1}$ by Haar invariance, so this index cancels on summing over the finite cut. Integrate a against the induced real quotient measure to obtain a scalar κ. The result is

$$\int F = \kappa \sum_{t \in T} \frac{b(t)}{\mathrm{mult}_T(t)}. \tag{16.21}$$

This is `Grading.exists_integral_eq_sum_over_cut`. In the general case the Bessel group maps to its finite factor with kernel its real factor, so one uses a cutoff for the real subgroup in taking the quotient by stages. The integration of a against that cutoff produces the same scalar independently of b, J, T. The partition identity shows why neither the finite factor κ nor the reciprocal multiplicities can be changed by choosing a different representative of a J-coset. In particular the only variable denominators of (16.21) are supported at the rational primes dividing q.

16.11 Two edges at once: removal of the extra plane

The even stages of the chain are special. Start with an odd string length s, then $s-1$ (even), then $s-2$ (odd or, at the end, zero). The two maps of (16.5) are denoted $T : V \to V'$ and $T' : V' \to V''$. On the orthogonal sum $V^- \oplus V''$, where the form on V^- is the negative of that on V, let

$$R = \begin{pmatrix} T^* \\ T' \end{pmatrix} : V' \longrightarrow V^- \oplus V''.$$

By the moment-map identities,

$$R^* R = -TT^* + (T')^* T' = 0. \tag{16.22}$$

Thus the two frequencies of the successive periods fit into a single isotropic frequency. The identification of the two Hom spaces with the Hom from V' to $V^- \oplus V''$ uses the adjoint in the first component and the identity in the second. It is a symplectic identification, because the adjoint reverses the trace pairing and the form on the first outer space has been negated. Each relative degree is preserved by it. The identification is `arrowProduct`, and R is `stackedT`, in `New/Seesaw/EdgePair.lean`.

For $s > 1$ write z, z'' for the two middle vectors in the outer spaces; their norms in the sum are N and $-N$. The degree-zero part of this sum is $X^- \oplus X''$ plus their two-dimensional plane. A basis of that plane is

$$p = (z - z'')/(2N), \qquad v = z + z''.$$

Both vectors are isotropic and their pairing is 1. The source's two-step splitting first takes the degree-zero part of the Hom space and then removes this additional hyperbolic plane. When $s = 1$, the final space has no string and the degree-zero part instead contains the one vector z/π in the negative outer space; its norm is 1, and it is adjoined as an extra definite line. This is precisely the pair of constructions `middlePair` and `lastPair`.

The reason for removing the extra plane can be made entirely explicit. Every grade-raising unipotent in the first odd string has degree at least two, and its phase cancels the generic

character by (16.7). In the even intermediate string, degrees at least two cause no phase in the product of the two constant terms, by (16.22). There are also degree-one terms because the string grades are odd and the X'-grade is zero. The degree-zero part of the product $R(g-1)$ for a grade-raising unipotent g of the intermediate space can only arise from a source X'-column going to the grade-one string vector, or from a grade-minus-one vector going to X'. But R kills X', and on the grade-one vector its value is $c(z+z'')$: for the first component, the old middle coefficient of T was $-Nc$, divided in the adjoint by the old Gram coefficient $-N$; the second component has coefficient c by (16.5). Thus all such extra degree-zero terms translate along the isotropic line of the additional plane, times arbitrary X'-coordinates. The integral over the rational quotient of this additive group is exactly the additional zero term. The map from the degree-one part of the intermediate unipotent to these X'-coordinates is onto: each elementary map from the X'-block to the grade-one vector, together with its negative adjoint from the reflected vector to X', exponentiates to an element of the unipotent, and its higher-degree terms do not affect the projection. The uniformizing measure on that additive quotient is fixed up to a constant independent of the datum. When $s = 1$, the intermediate space is definite so this additional integration is absent.

In the mixed model the two-step zero term consequently gives an ordinary theta integral on the three definite spaces. More explicitly, the product of the two theta functionals in the orthogonal sum is a fixed nonzero multiple of the theta functional for their product datum. This follows in the classical Schrödinger model by taking the rational theta sum on a direct sum of Lagrangians: on elementary tensors the sum is the product of the two sums, and finite linear combinations of tensors, and their Schwartz completions, give the general case. The adjoint and the sign reversal use the naturally isomorphic opposite symplectic model. Taking constant terms along the two positive Hom coordinate groups commutes with this product identification by Fubini, because they become the direct sum of the positive coordinate groups. The subsequent folding just calculated removes the intermediate unipotent integrals. Thus the result of integrating the two original constant terms against a generic P is, up to a single normalization independent of the finite datum, the integral of P restricted to the first definite group against the double theta lift with the evaluated datum. This is the calculation in `ArrowPair.integral_constTermAtT_mul`, `New/Seesaw/DoubleLiftValue.lean`.

The finite part of its evaluated datum is

$$\mathrm{ev}_f(\rho_f(R_f)(f_1 \boxtimes f_2)),$$

where ev_f is the two-stage zero term just described followed by identifying with the tensor of the two definite Hom spaces, and R_f is R in normalized finite coordinates. We denote the whole evaluation by L. A tensor product of the two finite vectors is sufficient here: arbitrary real polynomial vectors are allowed after taking the real zero term because the real and finite realizations factor. More precisely, the zero term of the tensor of a real vector and a finite vector in the mixed model is the tensor of its real zero term and its finite zero term. This assertion is used with the real vector possibly depending on the two archimedean test points; the finite operator does not depend on those test points or the twist. On polynomial Fock vectors, integration against a theta kernel produces a linear-form coefficient of the ordinary definite theta lift; applying this identity twice and using linearity gives the corresponding coefficient for the double lift. General real zero-term vectors follow by the density of polynomial Fock vectors and continuity of the integrals on the compact definite quotients; the resulting functionals lie in a finite-dimensional space and thus their limits remain in that space. These are the content of `zeroTermFin_spec` in `New/Seesaw/EdgePair.lean` and `integral_theta_eq_lift` with subscript 2 (the two-lift identity) in `New/Seesaw/DoubleLiftValue.lean`.

The preceding argument establishes the two-step operation in the classical normalized factorized Weil representation. To apply it to the particular algebraic modular form spaces of

the GRH argument, one must still identify the real joint harmonic vector and the branching coefficient used to interpret the resulting integral as exactly the source's linear lift map. This entails a compatibility of the kernel and its normalization, rather than a new orbit computation: the forms must be the indicated type, the twist must occur with the indicated exponent on both pairs, and the algebraic model must use the same class measure. We retain that specialized representation-theoretic identification, as in `Data.isArrow_datumY`, `Data.unfoldStage_even` and `Data.exists_last_stage_map`, when making the final application. The finite denominator calculation can be done independently of it, as follows.

16.12 The split neighbour lattice in the paired evaluation

We need the exact finite-place operation produced by removing the additional plane, not just its valuation. Denote by P_N the chosen places above N and write J for the finite idempotent equal to one on P_N and zero elsewhere. Fix the integral idelic unit η of norm -1 from the Weil model. Let e, e' be the two last coordinates of the two definite blocks in the finite tensor product (all other coordinates will stay unchanged). There is a unitary matrix Γ, identity outside $P_N \cup \bar{P}_N$, whose two-by-two block at a chosen split place w is

$$\Gamma_w = \begin{pmatrix} \pi & (\pi-1)\eta \\ 0 & 1 \end{pmatrix}, \qquad \Gamma_{\bar{w}} = \begin{pmatrix} \bar{\pi}^{-1} & 0 \\ (\bar{\pi}^{-1}-1)\bar{\eta} & 1 \end{pmatrix} \tag{16.23}$$

with the indicated components read at their own places. Conjugating the second block and taking its transpose gives the inverse of the first one, so the paired element is unitary for the standard finite Hermitian form. This is the matrix γ in `exists_isAdmissibleEval_std` of `New/Seesaw/AdmissibleEvaluation.lean`.

Here is the elementary identity of characteristic functions that makes this matrix useful. In the factor at $\bar{w}$, put $p = \bar{\pi}$, which is a uniformizer; N/p is a unit. Put $r_0 = e + \bar{\eta} e'$. For each residue $y \in \mathbb{F}_N$ let $\tilde{y}$ be its integer representative and take translation $n(y) = \eta(\tilde{y}/N) r_0$ at $\bar{w}$ and zero off the conjugate places. In the finite coordinates of a rectangle there is one such residue variable per intermediate row and per chosen place. At a single such coordinate, the new pair of coordinates of $\Gamma x + n(y)$ is

$$\left(p^{-1}x_e + \eta\tilde{y}/N, \quad x_{e'} - \bar{\eta}x_e + \bar{\eta}(p^{-1}x_e + \eta\tilde{y}/N)\right).$$

If these two coordinates are integral, then multiplying the first by p shows x_e integral, and subtracting its contribution to the second shows $x_{e'}$ integral. Conversely, given integral $x_e, x_{e'}$, precisely one residue y makes the first coordinate integral (since both p/N and η are units), and then the second is integral too. Thus the characteristic function of the integral box on the conjugate factor is the sum of the characteristic functions of the corresponding shifted boxes after action by Γ. Taking products over coordinates and places gives the box identity with multiplicity exactly one.

In the split Schrödinger realization, the action on box functions is substitution by the linear transformation and a constant Weil multiplier. The above identity therefore proves that, up to one scalar independent of the translate, the operation $\omega_f(\Gamma)$ on the projection to the relevant integral box is the sum of finite shifts indexed by $y \in (\mathbb{F}_N)^{r \times P_N}$. Under the adjoint identification of the two Hom spaces, the shifts are the source's

$$\tau_N = \sum_y \rho_f(-y/N \text{ in the last column}) \otimes \rho_f(-\bar{y}/N \text{ in the last row}), \tag{16.24}$$

where y/N denotes the adele supported at P_N with the indicated representatives, not the diagonal rational element. To check the orientation, the first component of the product datum

is converted by η times conjugate transpose; the shift $\eta(y/N)r_0$ on the conjugate support becomes $-y/N$ in that first component since $\eta\bar{\eta} = -1$. In the other component the second coordinate of r_0 gives $-\bar{y}/N$. The shift group is supported on disjoint conjugate factors, so their products have only the finite character phases already accounted for in the finite translation law. This is exactly `translateSum` of `New/Seesaw/NamesAtSplitLevel.lean`. It explains why one must use the two correlated shifts in (16.24), not the product of two independent sums.

16.13 Weighted integrality of evaluations

We now check the exceptional lattice columns (the neighbour vectors) and denominators. Let β be the integral finite idele associated to the chosen primes above g, with $|\widehat{O}/\beta\widehat{O}| = g^f$ in the source normalization. Let δ be the unitary ratio $(\bar{\alpha}/\alpha)(a/\bar{a})$, equal to one above N. In this application it is the ratio of the chosen idele above g and its conjugate: $\bar{\beta}\delta$ is integral and has the same valuations as β. For a vector in the product Hom model attach the coefficient weight β to a coordinate in the first block before conjugate transpose, and $\bar{\beta}$ in the second block. A vector is allowed if its coordinate multiplied by its weight is integral away from the twist conductor q.

The columns of Γ are neighbour vectors in the following precise sense. For each standard basis column j there is c_j such that

$$\begin{gathered}\delta c_j = c_j, \qquad \bar{\pi}\,\Gamma_{ij} \in \widehat{O}, \qquad \Gamma_{ij} \in \widehat{O}\ (i \neq e, e'),\\ c_j - \pi^{-1}\Gamma_{ej} \in \widehat{O}, \qquad \bar{\eta}c_j - \pi^{-1}\Gamma_{e'j} \in \widehat{O}.\end{gathered} \tag{16.25}$$

To verify these assertions, use the formula in (16.23). At $w \in P_N$, choose $c_j = \pi^{-1}\Gamma_{ej} - h_j$ with integral h_j so that $\bar{\eta}c_j - \pi^{-1}\Gamma_{e'j}$ is integral: modulo the ideal π, the last two entries of the first row of Γ are $0, -\eta$, and those of the second are $0, 1$. Their relation is precisely $\Gamma_{e'j} \equiv \bar{\eta}\Gamma_{ej}$ (mod π). At $w \in \bar{P}_N$, π is a unit, and the two possibly nonintegral entries occur with $\Gamma_{e'j} \equiv \bar{\eta}\Gamma_{ej}$ (mod $\widehat{O}$), by subtracting the multiple appearing in (16.23). Again choose $c_j = \pi^{-1}\Gamma_{ej}$ up to allowed integral error. Elsewhere take $c_j = 0$ since π is a unit and $\Gamma = 1$. Thus c_j is supported above N, where $\delta = 1$, and the remaining integrality assertions are clear. This proves the neighbour condition of `IsNeighbourVector` in `New/Seesaw/AdmissibleEvaluation.lean`.

In the additional plane removed in a middle pair, the corrected embedding of a neighbour vector has zero coordinate on the first hyperbolic basis vector and has second coordinate proportional to c_j (on the isotropic line $z + z''$). We check it belongs to the two glued lattices. In an odd glued lattice, a vector with X part x and only middle string part uh is integral in its framed coordinates precisely when all ordinary X-coordinates are integral, $\bar{\pi}x_e$ is integral, and $h - \pi^{-1}x_e$ is integral. Indeed express it in the basis of Lemma 16.13; the coefficient of the last X vector is $\bar{\pi}x_e$ and the coefficient of the middle string is the stated difference. In the first outer space of the pair the corrected vector has $h = c_j$. In the second it has $h = \bar{\eta}\delta c_j = \bar{\eta}c_j$, because the two middle string scalings differ by $a/\bar{a}$ and the sign-reversing idele has product with that ratio equal to $\bar{\eta}\delta$. Both membership tests are precisely (16.25). At the last pair there is no hyperbolic plane; the old middle coordinate is instead the added last definite coordinate π^{-1} multiplied by the same sign and scaling ratio. In the old lattice the needed difference now reads

$$\bar{\eta}^{-1}\pi^{-1}\Gamma_{e'j} - \pi^{-1}\Gamma_{ej} \ \in \widehat{O}$$

and follows by subtracting the two congruences in (16.25). All other X''-coordinates are unchanged. Thus the same membership holds also at the terminal pair. These checks are the

lattice part of `Data.isAdmissibleEval_pairEvalY`, in `New/Seesaw/GoodDatumExists.lean`.

After converting to the normalized finite frames the integral matrix of this embedding, including the correction along the removed isotropic plane, is $B = ig + kc$ in (16.13), with $g = \mathrm{diag}(\kappa)\Gamma$ and $\kappa = 1$ except for the special terminal coordinate, where $\kappa = \delta$. Tensoring with the X' coordinates preserves integrality: in the even intermediate glued lattice the X' components are not rescaled. The adjoint B^* carries weighted translations to weighted translations with the corresponding weights, including κ. To check this local statement, if an entry of B is nonzero, either the source and target coordinates lie in blocks of the same weight, or they lie above N where both $\beta, \bar\beta$ are units. The additional terminal coordinate is the only change of block weight away from N, and multiplying its target weight $\bar\beta$ by δ gives exactly the valuations of the old weight β. Therefore in each nonzero summand of a coordinate of B^*y, its target weight has the valuation of the source weight at the corresponding conjugate place. Each term is integral under the weighted hypothesis on y, and so is their sum. This is the proof of the weighted matrix lemma `mulVec_mem_weightedTranslations` of `New/Seesaw/AdmissibleEvaluation.lean` in the present application. Also $B^*R_f = 0$, since the map R kills the definite grade-zero subspace X' and the embedding is supported in the two grade-zero outer blocks; the nonzero frequency thus gives no additional translation on the definite component. This last observation uses the removal of nonzero relative degrees before forming the finite frame matrix B.

To bound the remaining coefficients we use the standard vacuum sum in finite coordinates. For an invertible unitary g_0 let $L_0 = \widehat{O}^h$ be the standard lattice; put

$$I = [L_0 : L_0 \cap g_0^{-1} L_0].$$

With the fixed finite Weil multiplier $\Delta_f(g_0)$ the formula is

$$\omega_f(g_0)v_{\mathrm{vac}} = \frac{\Delta_f(g_0)}{I} \sum_{a \in L_0/(L_0 \cap g_0^{-1}L_0)} \rho_f(g_0 a) v_{\mathrm{vac}}.$$

Indeed project onto the subspace fixed by the standard integral lattice by averaging the finite Heisenberg translation representation over Haar measure one on L_0. This is the vacuum line. Applying the analogous projection for $g_0 L_0$ to the original vacuum yields a nonzero vector on the translated vacuum line. Its factor with the normalized vacuum is by definition the fixed multiplier with $|\Delta_f(g_0)|^2 = I$, and the average over L_0 is the sum over the indicated finite subgroup cosets divided by I. This is `omega_vacElem_sum` in `Lib/Weil/Construction/Projections.lean`. After moving a translate of the vacuum, the additional coefficients are roots of unity from the Heisenberg law. In particular when the fixed scalar $\Delta_f(g_0)$ is factored out, the coefficients have all conjugate absolute sum at most one. If $\bar\beta g_0$ is an integral matrix, the intersection lattice contains $\bar\beta L_0$, so I divides $|\widehat{O}/\beta\widehat{O}|^h$. The source uses this argument for the diagonal $g_0 = \mathrm{diag}(\kappa)$, while the special Γ operation is treated by the split identity (16.24); otherwise one would introduce an unnecessary denominator at N.

Combine this sum, the weighted matrix condition and the zero-term formula (16.13). For the evaluation map L of a pair of consecutive lifts and each admissible y, the evaluated vacuum translate is a finite combination of the vectors $C\tau_N(v_t \otimes v_{t'})$ with

$$\bar\beta t,\ \bar\beta t' \quad \text{integral away from } q,$$

for one constant C independent of q and y. The denominators of the coefficients divide $|\widehat{O}/\beta\widehat{O}|^h$ for h equal to the product of the intermediate definite rank and the sum of the two

outer definite ranks (with the terminal extra line). When $q = 1$ their conjugate absolute values sum to at most one. If the source finite vectors themselves are the indicators of residue conditions on this weighted lattice, their standard finite projector formula expresses them as the corresponding sums of these vacuum translates; the normalized finite multipliers at the two directions of a pair cancel as in the formula of the preceding chapter, leaving the same fixed scalar and bound. Thus the operation on a pair is the admissible evaluation of `IsAdmissibleEval` in `New/Seesaw/AdmissibleEvaluation.lean`, including its explicit exceptional neighbour columns. The specialized assertion that the initial finite residue functions have the precise chosen projector realization is still the one discussed in the CM and Weil-model chapters; the neighbour calculation establishes what happens once that realization is made.

The paired projections at the level prime. The two translated factors before applying τ_N are integral in both conjugate supports of N, because q and β are prime to N. Project both factors to the subspace fixed by the standard integral shifts above those places. In the sum (16.24), a nonzero residue y has a nonintegral translation in at least one factor. Its projection is zero by the self-dual pairing: commuting with an integral shift detecting its residue multiplies the purported projection by a nontrivial character. Only $y = 0$ survives, giving back the unshifted pair. Conversely every term of the τ_N expansion has its translation integral after multiplication by $N\bar{\beta}$ away from q. On a pair with intermediate rank r, the number of terms is N^{fr}; all their phases have conjugate absolute value one (or their coefficient is zero). This proves the projection-and-expansion assertions of `Data.projOnFinPart_pairTranslates` and `Data.isAdmissibleComb_reducedVacuum` in `New/Seesaw/GoodDatumExists.lean`, from the finite projection law and the now-proved neighbour formula.

16.14 The admissible combinations and the iterated identity

Let $m = 2\varrho + 2$ be the number of lifts and n_m the largest rank parameter of the chain, and put

$$K_{mcut} = g^{fn_m^2}, \qquad D_{meval} = K_{mcut}^{2m}.$$

In this section a combination of vectors with coefficient function $r(t)$ of finite support is called admissible with denominator d if every $dr(t)$ is integral under all finite coefficient embeddings at primes not dividing q. Under $q = 1$ also require $\sum_t |\sigma r(t)| \le B$ for every automorphism of the complex numbers. We record three simple closure properties, since losing one of these bounds in the iteration would lose the gain at the small nonsplit primes.

First, if $v = \sum_t r(t)v_t$ is such a combination and for each t the vector v_t is itself a combination of the w_u with denominator d' and size B', then the resulting coefficients are $c(u) = \sum_t r(t)r_t(u)$. All $dd'c(u)$ are integral away from q by closure of each valuation ring under sums and products. The complex conjugate sizes satisfy

$$\sum_u |\sigma c(u)| \le \sum_t |\sigma r(t)| \sum_u |\sigma r_t(u)| \le BB'.$$

Secondly, tensor products of combinations multiply their denominators and size bounds by the same argument, since their coefficients are products. Applying any complex-linear map to each vector leaves coefficients unchanged. Thirdly, replacing any summand by a root-of-unity multiple of an admissible vector leaves all the integrality and size conditions unchanged. These are precisely the rules `IsAdmissibleComb.bind`, `pure`, `map`, `of_eq_isOfFinOrder_smul` in `New/Seesaw/AdmissibleComb.lean`.

Use the one-step transfer successively. At each stage its finite cut is independent of the column; every finite support is in the orbit covered by the cut because of the near-line bound. Formula (16.21) writes the integration over that stage as one scalar times the weighted finite sum over the corresponding finite quotient. Fix the last adelic point to be the identity. For a linear map of finite tensor data $S(y)$ indexed by the remaining tuple of finite cosets, define the modified function in the next step of the induction by

$$S_{<k}(y) = \sum_{t \in T_k} \mathrm{mult}_{T_k}(t)^{-1} S(y_0, \dots, y_{k-1}, t, y_{k+1}, \dots).$$

The multiplicity factors are rational integers' reciprocals with primes dividing q only. Thus repeated operations produce an admissible combination with denominator 1. When $q = 1$, every multiplicity is one, and the stage cut has at most K_{cut} terms: each representative is integral and its quotient by the chosen principal level embeds in the general linear group over the residue ring at the chosen primes. Indeed reduction gives the map into that finite group, its kernel on K_0 (or the stage-zero chosen subgroup) contains the principal subgroup of the chosen level, and J_k contains that subgroup in the trivial twist case. The order of the general linear group is at most the number $(g^f)^{n_m^2}$ of all matrices. Consequently the m cuts have total size at most K_{cut}^m. The invariance of the combinations under the embedded column levels follows because those elements act on T_k by a permutation preserving its multiplicity, while at the last stage no update is made. This is the induction of `Data.exists_isAdmissibleComb_unfolding` in `New/Seesaw/SeesawTheorem.lean`.

At an even stage use the two-edge calculation of Section 16.11. The function left after that stage is a definite matrix coefficient of the paired-lift chain, with finite datum obtained by the product of the evaluated edge pairs. This is an induction on the number of pairs: for the first pair the initial restriction of the Poincaré series to the first definite group gives its coefficient, and the two-step zero-term calculation acts on it by the double theta lift. The same statement for one additional pair follows by applying that calculation to the preceding coefficient and composing the two definite lift maps. Thus the overall coefficient is linear in the finite tensor data.

By the preceding admissibility rules and weighted-evaluation calculation, each such tensor is a combination, up to the shared factor C, of reduced vacua $V_{mred}(t)$, one correlation operator τ_N on each pair, with denominator D_{eval}. Indeed on a pair the product of the relevant ranks h is at most $4n_m^2$, so its denominator divides $(g^f)^{4n_m^2} = K_{\mathrm{cut}}^4$. There are $m/2$ pairs, yielding D_{eval}. The size before this denominator is cleared is at most one for trivial twist; hence the combinations obtained by the preceding cut induction have size at most K_{cut}^m. Every t is integral after multiplication by $\bar{\beta}$ away from q. Expanding the τ_N factors then gives a combination of ordinary translated vacua integral after multiplication by $N\bar{\beta}$ away from q, and multiplies the size by at most $N^{fn_m m/2}$. Multiply the coefficients by D_{eval} to clear the common denominator and absorb its inverse into the final fixed scalar. The trivial-twist size bound becomes

$$K_{\mathrm{cut}}^{3m} N^{fn_m m/2}. \tag{16.26}$$

This is below $\exp(x^{60})$ by the numerical bounds of the data chapter: the first log term is $3mfn_m^2 \log g$ and the second $fn_m m \log N/2$, precisely the two crude size terms in the good datum construction of the preceding chapter. The important quantitative feature is that the conductor q has not appeared in (16.26) for the trivial twist, and N has not appeared in the denominator at any twist.

The evaluated tensor sum is invariant under the product of the intermediate column levels fixing the last component. This follows from the permutation invariance of the cuts and the equivariance of finite evaluation: simultaneous translation of the two endpoints of an

evaluated pair intertwines the rotation of the Hom factors. For shifts in the final expansion, apply the two projections at N to the reduced vacuum: only the untranslated factors survive, as in the preceding paragraph, and reapplying τ_N recovers the original reduced vacuum. By linearity this identity also holds for the entire tensor sum. Thus translating the invariant evaluated tensor into a sum of individual translated vacuum data produces the joint invariance and compatibility with the projected pair expansion required in `Data.IsGoodDatum`. This is the finite linear-algebraic content of `Data.exists_isGoodDatum_of_invariant`. The final operation of restricting the last definite space to the smaller definite group is the usual branching restriction. Taking the Petersson integral of every resulting coefficient against every vector of that last type determines the equality of the two lift chains. Its normalization, chosen in the source's `Data.exists_last_stage_map` of `New/Filling/DegreeBounds.lean`, is the fixed scalar independent of the twist. If the product of all the common scalars is zero, all the corresponding indefinite-chain values vanish and the zero combination is a good datum; otherwise divide by that product to choose the common nonzero rescaling. This explains the use of a nonzero unit in the source's conclusion.

16.15 The remaining representation-theoretic comparison

This chapter has supplied conventional proofs of the normalized CM conjugacy, the simultaneous torsion-translation and finite-adelic cluster law, the geometric orbit argument and its elementary mixed-model Fourier unfolding, the glued lattice and the integral Witt step, the velocity inequality and its exceptional middle-line condition, the cut lemma and its prime-supported denominators, and the finite neighbour-lattice evaluation of the paired datum. The latter gives the explicit denominator and size propagation in the iterated calculation. These are the specialized constructions underlying `Data.seesaw` in `New/Seesaw/SeesawTheorem.lean`. In particular the geometry of the cut is no longer a black box once the single-stage supports and coefficient identities are available.

For the application to all columns of the particular GRH construction, the remaining dependencies in this argument are as follows. The initial Bessel Poincaré series must have the prescribed generic character and finite support with its stated normal form; these are the filling identities in `New/Seesaw/BesselGeneric.lean` and the filling chapter. The explicit realization of the abstract Fourier–Weil representation by the theta vectors of the arithmetic construction must preserve their finite vacuum line and exact multipliers and the real joint harmonic types. In particular the coefficient formula for the concrete definite lift maps, and its specialization to each even stage in `Data.unfoldStage_even`, must be verified with its root factors and class measure, together with the final real branching pairing `Data.exists_last_stage_map`. These are classical-model comparison and exact representation choices, not further arbitrary orbit or lattice support assertions. They are used here as identified inputs, and the new arguments do not assert to reconstruct the entire analytic representation theory of those maps independently. The elementary induction of the final section proves the stated good-datum consequences subject to them. The distinction is important: a general period transfer theorem without the precise finite evaluation and coefficient normalization would not suffice to obtain the arithmetic upper bound.

Chapter 17

Compact theta coefficients, integral branching and conjugation of the arithmetic vectors

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 25, 27 and the final audit in Chapter 57.

This chapter deals with the precise comparison assertions left in the upper-bound chapter, rather than replacing them by a qualitative assertion of theta correspondence. We use the globally normalized theta functional, the actual vacuum and the mixed-model evaluation identities of Chapters 9, 7 and 16. In particular the normalization of an algebraic class-sum lift is not the normalization of integration with maximal compact volume one unless its chosen level is itself maximal.

17.1 The entire function associated with the finite vector

We spell out the missing model comparison in Chapter 12. Let V be standard definite of rank n, with $f = [E^+ : \mathbb{Q}]$, and $D = fn$. Use the source's real Fock coordinates and scaled finite coordinates, writing a rational vector $q \in E^n$ as $(r(q), s(q))$, where $r(q) = \mathrm{realImage}(q)$ and $s(q) = \alpha^{-1} q_f$. The real vacuum is $e_\infty(y) = e^{-\|y\|^2/2}$; the real Haar measure is π^{-D} times Lebesgue measure. The finite vacuum e_f has norm one for the measure assigning $\widehat{\mathcal{O}}_E^n$ volume one. Let ϑ denote the global rationally invariant theta functional with the normalization in Chapter 9. For clarity write

$$\widetilde{v}(y,t) = \vartheta(\rho(y,t)v), \qquad S_v(y) = e_\infty(y)^{-1}\widetilde{v}(y,0).$$

The adjective finite for v means that its real factor is the vacuum (equivalently v is fixed by the real vacuum projector). In these coordinates

$$\beta_\infty(a,b) = \exp((b^*a - a^*b)/2), \quad \beta_f(a,b) = \psi_0(b^*a/2), \quad \lambda(t) = \psi_0(c_0 t^* t/2), \quad c_0 + \bar{c}_0 = 1.$$

The finite bicharacter is alternating, and λ is the source's `stdLattice.semichar`. Chapter 7 proves the finite lattice model and its normalization; the sign in these formulas is important in the comparison with S_v.

Proposition 17.1 (The exact finite–real theta comparison)**.** *For standard positive V, the map $v \mapsto S_v$ is injective on the finite part and gives entire functions of $y \in \mathbb{C}^D$. For every rational q,*

$$S_{\rho(0,s(q))v}(y) = \exp\bigl(r(q)^* y - \|r(q)\|^2/2\bigr) S_v(y - r(q)).$$

Writing $B(a,b) = a^*b$, the right side is the theta shift $U_{-r(q)}S_v$ in the convention $U_a F(y) = e^{-B(a,y)-B(a,a)/2}F(y+a)$. If b_0 multiplies the coordinate at τ by $\overline{\tau(c_0)}$, the finite and archimedean multipliers satisfy

$$\lambda(s(q)) \exp\bigl(i \operatorname{Im} B(b_0(-r(q)), -r(q))\bigr) = 1.$$

This is the canonical multiplier comparison for the Riemann form $E_B = \operatorname{Im} B/\pi$.

Proof. Rational invariance and the Heisenberg multiplication identity give $\beta((r,s),z)\widetilde{v}((r,s)+z) = \widetilde{v}(z)$ for $(r,s) = (r(q), s(q))$. Taking $z = (y - r, 0)$ gives

$$\widetilde{v}(y,s) = \exp((r^* y - y^* r)/2)\widetilde{v}(y - r, 0).$$

Real and finite shifts commute exactly. Dividing by $e_\infty(y)$ and inserting the Gaussian at $y - r$ gives the claimed translation rule. If $\widetilde{v}(y,0)$ vanishes identically, this formula (or directly the preceding quasi-periodicity) forces $\widetilde{v}(y, s(q)) = 0$ for every y, q. The $s(q)$ are dense in the finite adeles by weak approximation, so continuity gives $\widetilde{v} = 0$. The theta mean identity, with the positive Hermitian form of the coefficient model, implies $v = 0$.

The finite-vector condition allows us to reproduce the real vacuum factor. Indeed $\rho(y)e_\infty(w) = e_\infty(y)e_\infty(w)\exp(\bar{w} \cdot y)$, so the reproducing identity gives

$$S_v(y) = \int_{\mathbb{C}^D} \exp(\bar{w} \cdot y) e_\infty(w) \widetilde{v}(w, 0)\, dw. \tag{17.1}$$

Here the function $\widetilde{v}$ is bounded since the action is isometric for the integrable coefficient norm and ϑ is continuous for that norm. For y in any compact set, the exponential times any polynomial in w is dominated in absolute value by a polynomial times $\exp(C\|w\| - \|w\|^2/2)$. It follows by differentiation under the absolutely convergent integral in all variables that S_v is entire.

Finally the global additive character is trivial on the rational vectors, hence

$$\beta((r(c_0 q), s(c_0 q)), (r(q), s(q))) = 1.$$

The finite factor of this equality is $\lambda(s(q))$. The infinite factor, using the explicit bicharacter, is $\exp(i \sum_{\tau,i} \operatorname{Im} \tau(c_0) |r(q)_{\tau i}|^2)$, precisely the canonical multiplier in the statement. This proves the product normalization without an arbitrary choice of an archimedean theta constant. □

Proposition 17.1 proves `Space.entireTheta_translate_finiteImage`, `semichar_finiteImage` and the injectivity assertion used in `Space.exists_entireTheta_eq` (`Lib/Weil/Construction/FockThetaFunctions.lean`). It also proves the analyticity statement in `Lib/Weil/Fock/EntireTheta.lean`. To check the Taylor identity, differentiate (17.1). For a monomial X^ν the result is

$$[X^\nu] S_v(cX) = \frac{c^{|\nu|}}{\nu!} \vartheta(v_{X^\nu} \otimes v).$$

Indeed $v_{X^\nu}(w) = \bar{w}^\nu e_\infty(w)$ in the all-positive Fock coordinates. This is exactly `Space.jetSeries` and `Space.jet_jetSeries_eq_theta` in `Lib/Weil/Construction/DataProduct.lean`. In particular equality of Taylor series at a nonzero scale forces equality of entire functions (and hence of the finite vectors); both the line-substitution and the product formula for jets

of Chapter 12 follow by the analytic multivariable Taylor theorem. No formal-power-series substitution at a nonzero constant term is used. As an alternative classical input for the dimension theorem used with this comparison, one may apply the Appell–Humbert theorem and Riemann–Roch for a positive line bundle on a complex torus: the dimension is the square root of the order of the dual lattice quotient. The same dimension follows from the ordinary theta Fourier expansion in a symplectic basis: if the polarization has elementary divisors $d_1, \dots, d_D$, periodicity in the first half of a symplectic basis yields Fourier modes indexed by $\mathbb{Z}^D$, and the other half imposes recurrence between indices differing by d_i in coordinate i. Positivity makes each of the resulting $\prod_i d_i$ Gaussian series normally convergent and gives independent initial coefficients; the dual lattice quotient has order $(\prod_i d_i)^2$.

Thus the finite theta realization and expansion in Chapter 12, combined with the tight frames constructed there and the normalized CM conjugation in Theorem 16.5, prove `Parameters.exists_finitePart_of_isTightFrame` (`New/Curve/FrameSections.lean`) and `Parameters.exists_isJetConjugate_generator` (`New/Curve/ConjugateGenerator.lean`). Specifically, the preceding dimension and multiplier calculation justifies the exact finite orbit on which the theta-group orthogonality is applied, not merely a dimension comparison between two unspecified theta representations. The vacuum clearing factor in Proposition 12.18 is for the actual normalized vacuum of the source: the finite overlap is $\lambda(t)\mathbf{1}_{\widehat{\mathcal{O}}_E^n}(t)$ by the lattice calculation, real vacuum norm one. In rank n its clearing factor is the n-th power of that in rank one. The adelic torsion transport in Theorem 16.5 can be used simultaneously for all vectors in the finite-product construction by the rational approximation and continuity argument of Chapter 12; it is the same transport scalar at every rank. We next check its action on the full finite Weil part.

17.2 From CM theta conjugation to the finite Weil representation

We use $\Delta(g) = \Delta_f(g)$ with the exact finite normalization of Chapter 7. Recall

$$\langle e_f, \omega(g) e_f \rangle = 1/\overline{\Delta(g)}, \quad |\Delta(g)|^2 = [L : L \cap g^{-1}L], \quad \overline{\Delta(g)} = \upsilon(\det g)\Delta(g), \qquad L = \widehat{\mathcal{O}}_E^n. \tag{17.2}$$

For $b \in \mathbb{A}_{E,f}$ and an automorphism σ of $\mathbb{C}/K$, suppose a nonzero σ-semilinear map M on the finite space satisfies

$$M\rho(x) = \rho(bx)M, \qquad \sigma(\psi_0(z)) = \psi_0(b\bar{b}z). \tag{17.3}$$

For the conjugate theta vectors this map is constructed as follows. The generating span described after Proposition 17.1 has conjugate jet series $T_t\rho(bx)F'$, with coefficients conjugated and places renamed as in `Space.IsJetConjugate`. Since T_t is injective and the jets of the generating orbit map to this actual orbit, its inverse on the image defines a semilinear map on the whole finite vector space. It is injective by injectivity of T_{per}. To check the central-character identity before using the intertwining law, choose a nonzero rational integer m such that $\rho(y_0)F = F$ and $\rho(by_0)F' = F'$, where $y_0 = (m, 0, \dots, 0)$ in finite coordinates. Such an m exists since both finite translation orbits are locally constant at zero. For arbitrary x, the equality $\rho(x + y_0)F = \beta_f(x, y_0)^{-1}\rho(x)F$ and the analogous identity for the primed generator give $\sigma(\beta_f(x, y_0)) = \beta_f(bx, by_0)$ after applying the already constructed M and comparing its values on the two generators. The character arguments $\overline{y_0} \cdot x/2$ exhaust the finite adeles, since a nonzero rational number is an invertible finite idele. This proves the second equation of (17.3); the translation product formula now proves the first equation on every generating translate and hence on their span. Rank zero is immediate, and the transported scalar can be fixed in rank one for all ranks. This also verifies directly that the scalar conjugation comes

from the central character, not from an assumed equivariance of a complex-conjugate Weil representation.

Theorem 17.2 (Semilinear finite comparison)**.** *Let R be any conjugate-disjoint set of finite places and assume (17.3). Then b is an invertible finite idele. There are $c \neq 0$ and a norm-one finite idele u such that*

$$Me_f = c\,\omega(u)e_f, \qquad M\omega(g)v = \frac{\sigma(\Delta(g))}{\Delta(g)}\omega(g)Mv$$

on the finite part, where $\omega(u)$ denotes the scalar unitary matrix. For a Schwartz–Bruhat function Ψ above R, the source's mixed-model vector $\mathrm{name}_R(\Psi)e_f$ satisfies

$$M(\mathrm{name}_R(\Psi)e_f) = c\,\omega(u)\,\mathrm{name}_R(\sigma\circ\Psi)e_f.$$

Proof. Put $d = b\bar{b}$. For an arbitrary finite adele z, self-duality of the integral adelic lattice and (17.3) give $z \in \widehat{\mathcal{O}}_E \Leftrightarrow dz \in \widehat{\mathcal{O}}_E$: both conditions are tested by whether $\psi_0(zx)$, respectively $\psi_0(dzx)$, is 1 for every integral x. Taking vectors supported at one place and testing successive powers of a uniformizer shows that d is a unit at every place. Thus b is invertible and d is a finite integral unit. Construct an integral unit a with $a\bar{a} = d$ as follows. At each nonsplit place $v(b\bar{b}) = 0$ implies $v(b) = 0$, so take $a = b$ there (no surjectivity of the local unit norm is needed). At each split pair take one component of a to be 1 and its opposite component to be the prescribed conjugate of d; when the pair meets R, choose the component 1 on R. Conjugate-disjointness makes the choices compatible and all components of a are units. Take $u = b/a$, a norm-one idele.

Normalize the finite translations by $P(x) = \lambda(x)\rho(x)$, as in `Space.finHeisOp`. Formula (17.3) also gives $MP(x) = P(bx)M$, since $\lambda(x) = \psi_0(c_0x^*x/2)$. The line on which every $P(x)$ with $x \in L$ acts trivially is the vacuum line, by the lattice model. Therefore Me_f is on the line $\omega(u)e_f$, since it is fixed by the corresponding translations of $bL = uL$. This proves $Me_f = c\omega(u)e_f$; injectivity gives $c \neq 0$.

The vectors $P(x)e_f$ span the finite part and have vacuum coefficient $\mathbf{1}_L(x)$. Applying M and using $aL = L$ proves for arbitrary finite v that

$$\langle \omega(u)e_f, Mv\rangle = c\sigma(\langle e_f, v\rangle).$$

Similarly $M\omega(g)e_f$ is on the line $\omega(g)\omega(u)e_f$, because it is fixed by translations of $bgL = guL$. Compare its coefficient against $\omega(u)e_f$: by (17.2) the ratio to the coefficient of $\omega(g)Me_f$ is $\overline{\Delta(g)}/\sigma(\overline{\Delta(g)}) = \sigma(\Delta(g))/\Delta(g)$, where the equality uses $|\Delta(g)|^2 \in \mathbb{Z}_{>0}$. This proves the intertwining identity on the vacuum and then on its translates, since g commutes with the scalar b.

For the last assertion use the explicit finite Schrödinger action above R:

$$(S(y)\Psi)(x) = \psi_0(\tfrac{1}{2}y^*e_R(2x+y))\Psi(x+y),$$

where e_R is the adelic idempotent for R. A compactly supported locally constant function on the R coordinates is a finite linear combination of $S(y)\mathbf{1}_L$. To see this choose an open subgroup of the integral lattice on whose cosets the function is constant, cover the compact support by finitely many lattice cosets and expand each finer coset indicator in the characters of the corresponding finite quotient. The dual is supplied by the opposite coordinates because the lattice there is dual under ψ_0. At a place not in R the function is independent of the coordinates. The vector named by $S(y)\mathbf{1}_L$ is the Heisenberg translate of e_f by $e_{R\cup\bar{R}}y$ (the definition of `Space.name` on this spanning set). This naming is well-defined: pairings between

translates equal pairings between the shifted indicators, by the integral of ψ_0 over the self-dual lattice, so any functional relation gives the zero vector by positivity. As $e_R a = e_R$,

$$\sigma(S(y)\mathbf{1}_L) = S(ay)\mathbf{1}_L,$$

the argument of the character on the right being $a\bar{a}$ times the original and its box condition being unchanged. Applying M to the vector corresponding to $S(y)\mathbf{1}_L$ gives $c\omega(u)$ times the vector corresponding to $S(ay)\mathbf{1}_L$. Semilinearity and the finite spanning argument prove the assertion. □

This is `Space.semilinear_intertwines_weil` in `Lib/CM/Reciprocity/SemilinearWeilLaw.lean`. In particular the ratio is not an independent character whose size can be chosen freely. From (17.2),

$$(\sigma\Delta(g)/\Delta(g))^2 = \upsilon(\det g)/\sigma(\upsilon(\det g)).$$

For a norm-one finite idele the absolute value of every conjugate of υ is one: raise the idele ideal to the class number to make it principal, and the defining archimedean product for the unitary character then expresses a power of its value by a product of conjugate algebraic numbers of total modulus one. In detail if the generator is a, the norm equality gives $\prod_\tau |\tau(a)|^2 = 1$, and this product stays 1 after any automorphism of the normal CM field (complex conjugation is central). The finite unit factors are roots of unity. Consequently the ratio above has absolute value one.

Proposition 17.3 (Uniform absolute bound in the comparison)**.** *For the map M constructed from conjugate jets, if $\mathrm{vac}^\circ = s_{\mathrm{vac}}^n e_f$ is the integral vacuum of the previous chapter, then*

$$|\vartheta(\rho(y)M\rho(x)\omega(g)\,\mathrm{vac}^\circ)| \le \exp\bigl(C_P(N_1 n + 1)^2\bigr)$$

for all finite x, g and all adelic y. Here one may take C_P to be the generator constant (the constant in $\exp(C_0 N_1^2 n)$) plus $2d + 6$, as in the source's parameter constant.

Proof. The constant coefficient of the jet comparison and the clearing factor of Proposition 12.18 imply $|\vartheta(\rho(x)M\,\mathrm{vac}^\circ)| \le \exp(C_0 N_1^2 n)$ for all finite x (use surjectivity of multiplication by b). This bound also holds for all adelic x: given a prescribed archimedean component, add rational vectors whose archimedean components approximate its negative, using weak approximation at the archimedean places. Quasi-periodicity preserves the absolute value, and continuity of the finite-vector evaluation gives the bound. Equivalently the dense subset of vectors with rational archimedean component reduces exactly to a finite evaluation by subtracting the rational vector, and then use closedness of the inequality. Taking the normalized mean over the rational quotient shows that the Hilbert norm of $M\,\mathrm{vac}^\circ$ is at most $\exp(C_0 N_1^2 n)$. By Theorem 17.2 and norm one of the multiplier ratio it remains to bound the absolute theta evaluation of any translated vector $w = \rho(z)\omega(h)e_f$.

The theta Fourier identity of Chapter 9 gives $|\vartheta(w)|^2 = \sum_{q\in E^n}\langle w, \rho(q)w\rangle$. Absolute values of the summands are at most the Gaussian $\exp(-\pi\|q_\infty\|^2/\sqrt{d})$ restricted to the set where $h^{-1}q_f$ belongs to αL in unscaled coordinates. Here h is finite: conjugating by $\rho(z)$ only introduces a phase. Distinct points of the resulting archimedean lattice after multiplication by N_1 have separation at least 1. Indeed $N_1\alpha$ is integral by the chosen different idele calculation, and the unitary matrix h preserves the Hermitian form. Thus the Hermitian square of $N_1(q - q')$ is a nonzero totally positive algebraic integer, whose sum over the real embeddings is at least 1.

Let $D_0 = fn$, $A = \pi/(\sqrt{d}N_1^2)$, and $M_0 = N_1 n \ge D_0$. Balls of radius $1/2$ about the scaled lattice points are disjoint in $2D_0$ dimensions. For a point v in the ball centred at w,

$\|w\|^2 \geq \|v\|^2/2 - 1/4$. Integration gives

$$\sum_w e^{-A\|w\|^2} \leq e^{A/4} 2^{2D_0} D_0! \, (2/A)^{D_0}.$$

For $n > 0$, use $D_0! \leq D_0^{D_0}$, $\log N_1, \log D_0 \leq M_0$, $\log d \leq d$, $A/4 \leq 1$ and $\log 2 \leq 1$ to bound the logarithm on the right by $1 + 3M_0^2 + (3+d)M_0 \leq (2d+6)(M_0+1)^2$. Rank zero is immediate. Taking its square root gives the stated overestimate after multiplying by the previously bounded Hilbert norm. This argument is uniform in the lattice class, the translation and the finite unitary point; no dependence on the support of a single vector was introduced. □

Propositions 17.1 and 17.3, together with Theorem 17.2, prove the comparison assertion `Parameters.exists_semilinear_isJetConjugate` of `New/Curve/ConjugatesOfJets.lean` using the normalized theta Fourier identity. The archimedean factor $(N_1 n + 1)^2$ is essential here: an uncontrolled constant depending on an individual adelic translate or on its conductor would not suffice.

17.3 The explicit conjugate-jet operator bound

The estimate needed in the chain is the second, operator-norm assertion of `Parameters.norm_jet_le_of_isAdmissibleComb`, not a separate Cauchy estimate for every coordinate. Here is a direct derivation. Let n, n' be the two ranks, $D_0 = |\Phi| n n'$ the number of complex matrix entries, F a finite combination of vacuum translates of total coefficient absolute value at most B under every complex conjugate, and set $u = \log(|\sigma c|^2/\Omega)$, where $\Omega = P.\mathrm{bigOmega}(s)$. Put

$$M(L, w) = B \exp\{P.c(N_1 n n' + 1)^2 + D_0 \log(2L+2) + w\}.$$

If a polynomial Q satisfies the hypothesis in the source

$$\sum_i |\sigma z_i|^2 \leq e^u (2L+2) R \quad \Longrightarrow \quad |\sigma Q(z)| \leq A \exp(w + LR),$$

then $|\sigma \, \mathrm{jet}_Q(\omega(g,h)F; c)| \leq A M(L,w)$ for every finite pair (g,h). Indeed it suffices first to treat a single vacuum translate and an automorphism fixing K: an automorphism not fixing K may be composed on the range with complex conjugation, preserving the absolute norm. The semilinear transformation of Theorem 17.2 brings the conjugated polynomial under the real integral of (17.1), by its monomial Taylor formula. The transport permutes the complex places and replaces the polynomial arguments by z_i of transformed absolute squares satisfying

$$\sum_i |\sigma z_i|^2 = e^u \sum_i |y_i|^2.$$

This is precisely the scale compatibility $|t|^2 = \Omega^{-1}$ for the real scalar t of the CM comparison; the permutation is harmless. Both the pair character and the semilinear finite Weil multiplier have norm one. The uniform vacuum estimate of Proposition 17.3 bounds the remaining theta function, including the translating vector, independently of y. Choose $R = \sum_i |y_i|^2/(2L+2)$; the polynomial contributes at most $A \exp(w + L\|y\|^2/(2L+2))$. Multiplication by the real vacuum $\exp(-\|y\|^2/2)$ leaves the Gaussian $\exp(-\|y\|^2/(2L+2))$. Its integral for the normalized measure $\pi^{-D_0} dy$ is $(2L+2)^{D_0}$, giving exactly the bound. The triangle inequality for the finite combination multiplies the single bound by B.

For the Schur jet, let $d_{\tau i}$ be its diagram row lengths indexed by $0 \le i < n$. The hypothesis on the lift weight ensures that no omitted row has positive length. The polynomial matrix coefficients satisfy

$$|\langle S_\kappa(Z)v, w'\rangle| \le \|v\|\|w'\| \prod_{\tau,i} s_{\tau i}^{d_{\tau i}},$$

where $s_{\tau i}$ are the singular values of Z_τ in decreasing order. To prove this, take the singular-value decomposition of each rectangular array and use the unitary invariance of the Gram norms; on a rectangular diagonal matrix each minor is a product of singular values. The highest-weight character is the largest possible such product: in each column of height h the maximal product is the first h singular values, and their product over the columns is the stated bound. This argument may also be phrased by embedding the Schur space isometrically into the tensor product of exterior powers. The polynomial dual coefficient is defined by the Gram matrix (whose entries are rational) and the coefficient vector σv. Thus exactly this bound applies to the conjugated polynomial in the preceding inequality; its dual testing vector can be chosen arbitrarily after conjugation.

For $L \ge 1$, put $t_{\tau i} = s_{\tau i}^2 e^{-u} L/(L+1)$. Since $t^d/d! \le e^t$ for $t \ge 0$, and $\log(1+1/L) \le 1/L$, the product satisfies

$$\prod_{\tau i} s_{\tau i}^{d_{\tau i}} \le \exp\left\{ \frac{1}{2} \sum_{\tau i} \left(d_{\tau i}/L + \log(d_{\tau i}!) + d_{\tau i} u \right) + \frac{1}{2} \sum_{\tau i} t_{\tau i} \right\}.$$

If the squared matrix norm is at most $e^u(2L+2)R$, then the last half sum is at most LR. Apply the polynomial estimate to the dual coefficient and take the supremum over unit dual vectors. This proves

$$\|\sigma(J_c(\omega(g,h)F)v)\| \le M\left(L, \frac{1}{2} \sum_{\tau i} (d_{\tau i}/L + \log(d_{\tau i}!) + d_{\tau i} u) \right) \|\sigma v\|. \tag{17.4}$$

The norm here is the Weyl Gram norm on coefficient vectors, not the coordinate Euclidean norm. This proves both assertions of `Parameters.norm_jet_le_of_isAdmissibleComb` of `New/Rows/MemberKernels.lean`; in particular it does not add a dimension factor from the number of Schur coefficients. As in the source, the bound also applies to a singleton combination of coefficient one.

17.4 Stability of the arithmetic data and the gauge calculation

The finite vacuum sum is particularly simple in our lattice model. It is supported on $L + gL$ modulo L, and representatives of that quotient can be taken as gy with $y \in L/(L \cap g^{-1}L)$. The vacuum for gL is fixed by $P(gy)$ for $y \in L$. Its coefficient at zero is $1/\bar{\Delta}(g)$, so

$$\omega(g)e_f = \frac{\Delta(g)}{|\Delta(g)|^2} \sum_{y \in L/(L \cap g^{-1}L)} P(gy)e_f. \tag{17.5}$$

Indeed each term is supported on its displayed coset, with the correct eigenvalue under translations of L; conjugation by g carries the normalized translations on L to those on gL. These conditions uniquely determine the value on each such coset, and the coefficient at zero determines the scalar. This is `omega_vacElem_sum` in `Lib/Weil/Construction/Projections.lean`.

If g is integral invertible at all places over a rational prime p, then $\Delta(g)$ is a p-adic unit under every complex embedding into $\mathbb{C}_p$. To verify this rather than bound only its complex

modulus, write its finite multiplier as $\chi_f(t) = \upsilon(t)|t|_f^{-1/2}$ on the determinant of a saturating lattice. Its local components over p are units. Raise its fractional ideal to the class number, write the resulting ideal as (v), and use the defining character and product formula: up to a root of unity $\chi_f(t)^h = \prod_\tau \overline{\tau(v)}$, with h the chosen class-group exponent. Since v is a p-unit at every place over p, the norm of this product is one at every coefficient embedding. The same calculation for any of the algebraic determinant characters of the proof gives a product of integral powers of these embedded p-units times a root of unity coming from its finite tame character. Thus those character values are also p-units. Finally $|\Delta(g)|^2$ is the displayed lattice index, prime to p by the hypothesis on g. Both scalar factors in (17.5) are therefore harmless.

The integral standard-data module at a fixed coefficient embedding is the valuation-ring span of $P(x)\,\mathrm{vac}^\circ$ with δx integral above p. It is stable under all the same Heisenberg translations since their multiplication introduces only roots of unity, hence valuation units. Formula (17.5) proves stability under $\omega(g)$ for integral g: representatives gy are integral above p and g carries the allowed translation lattice into itself. The preceding determinant-character calculation proves the same stability for the pair Weil action at all pairs of integral elements. This is precisely `Parameters.finHeisOp_mem_standardData`, `Parameters.omega_mem_standardData`, `Parameters.pairWeil_mem_standardData` in `New/Curve/ConjugatesOfJets.lean` and the two unit lemmas of `New/Seesaw/IntegralAway.lean`.

For the gauge let K_p denote the unitary subgroup integral above p, and use ρ_p for the coordinate Weyl representation read at the fixed coefficient embedding (including all the specified weight embeddings). Its matrices and inverses are integral for K_p. Hence right multiplication of the gauge point by K_p is an isometry. Let the translated kernel datum be $\Psi = \omega(1, xdy)F$, with F in the standard module, x, y integral above p, and d diagonal under each weight embedding. At arbitrary arguments g, h, use weak approximation for unitary groups (Lemma 5.5) to choose rational γ, ϵ such that

$$g_1 = \gamma g \in K_p, \qquad h_1 = \epsilon h x d \in K_p \cap d^{-1} K_p d.$$

These are simultaneous finite local conditions on a neighbourhood of g^{-1} and $(hxd)^{-1}$; no strong approximation is assumed. Rational kernel equivariance permits replacing source vector and gauge arguments by their transforms under γ, ϵ, without changing the gauges. The datum in the kernel is then $\omega(g_1, h_1 y)F$, in the standard module by stability. On the source its gauge is the maximum coordinate norm. On the output the inverse of $\epsilon h = h_1 d^{-1} x^{-1}$ acts by $\rho_p(x d h_1^{-1})$; x is integral, as is $d h_1^{-1} d^{-1}$, so only $\rho_p(d)$ affects the gauge. A diagonal matrix acts on each minor basis vector by the product of its box dilations and its determinant character. The ultrametric matrix estimate therefore proves the precise weighted jet-coefficient bound in Proposition 13.22, with the product of the indicated diagonal weights (rather than an unweighted estimate), for all g, h. Homogeneity then gives `mem_raises_of_standardData`. Clearing the scalar by an integer divisible by $d_0(Nd)^L$ when $d_0 F$ is in the integral module gives the variant `mem_raises_of_dvd`, since extra integral multiples have norm at most one. This closes these applications of `mem_raises_of_jet_bound` to the standard module.

17.5 Compact coefficient lifts: no noncompact joint-harmonic theorem is necessary

The compact steps of the seesaw use a particular specialization of the theta correspondence. We give a direct argument for all polynomial data in those steps, not just for a single chosen joint harmonic. Let the standard definite groups have unequal ranks $n < n'$ and compact real groups K_∞, K'_∞. At place τ the number of columns of height h is $\kappa_\tau(h)$. The lifting-weight

condition is the absence of columns of heights $n < h < n'$ and

$$e_\tau = e_{b,\tau} + \kappa_\tau(n).$$

The output determinant exponent is $e_{a,\tau}$. Write W, W' for the corresponding coefficient vector spaces and ρ, ρ' for their conjugate Weyl representations of K_∞, K'_∞. On a real unitary matrix x the former is $S_{\kappa,n}(\bar{x})\det(\bar{x})^e$, and analogously at the target. This is the convention of `conjWeylRep`. Write $S_\kappa(Z)$ for the rectangular Schur matrix in the matrix variable $Z \in M_{n'\times n}$, the matrices for a product of places being tensored. Every entry is homogeneous of degree

$$B_\kappa = \sum_\tau \sum_{0<h<n'} h\kappa_\tau(h)$$

and for a row functional L on the output and a vector $v \in W$ put $P_{L,v}(Z) = L(S_\kappa(Z)v)$.

For a nonzero jet scale c, let $J_c(f)$ be the matrix obtained by applying the jets to $T_c f$ along the Schur entries. Then the identity after Proposition 17.1 and homogeneity give

$$\vartheta(v_{P_{L,v}} \otimes f) = c^{-B_\kappa} L(J_c(f)v). \tag{17.6}$$

At a compact pair (x, y) the real Weil action on polynomial vectors substitutes $Z \mapsto y^T Z x^{-T}$ and multiplies by $\prod_\tau \det(y_\tau)^{e_{a,\tau}} \det(x_\tau)^{-e_{b,\tau}}$. In fact the action on the real argument of the coefficient is the inverse of the unitary matrix action $Z \mapsto yZx^{-1}$, and taking complex conjugates on all positive coordinates yields the displayed substitution. The full determinant factors are the pair twist; the negative–negative oscillator block here is empty. Cauchy–Binet in the rectangular Schur matrices, with the full-height columns on the source extracted, gives

$$\Omega(x, y)P_{L,v} = P_{L\rho'(y^{-1}),\rho(x)v}.$$

Indeed the factor from the source columns of height n is $\det(x^{-T})^{\kappa_\tau(n)}$; together with the twist it gives $\det(\bar{x})^{e_\tau}$. This proves the precise identity `realPairWeil_schurCoeffPoly` of `Lib/Weil/CompactLift/BranchingPeriod.lean`, including the determinant shift. It is not a claim that the undecorated Schur matrices transform without determinant factors.

We recall why these polynomials are the full compact isotypic piece needed by integration against a form of weight ρ. The polynomial algebra of $n' \times n$ matrices is multiplicity-free as a representation of the two general linear groups, with summands indexed by partitions of length at most n. In fact its degree d part is the symmetric part of $(\mathbb{C}^{n'} \otimes \mathbb{C}^n)^{\otimes d}$. Apply Schur–Weyl duality to the two tensor powers and take simultaneous invariants under the permutation group: by Schur's lemma the tensor product of the two permutation irreducibles has a one-dimensional invariant space precisely when the partitions agree, and zero otherwise. Passing to the compact groups and replacing the appropriate argument by its dual gives the same decomposition for the action above. A precise published alternative is T. J. Enright and J. F. Willenbring, Annals of Mathematics 159 (2004), p. 370, formulas (7.4.3)–(7.4.4), where the two matrix-variable actions and their multiplicity-free decompositions are written explicitly. These are precisely polynomial representations, not a stable-range assertion for an indefinite group.

Here the partition in rank n' is formed from all heights below n'; it has at most n rows, and its first n row lengths are the rank-n diagram plus the constant $\kappa_\tau(n)$. The determinant condition above means it is the unique partition matching the input compact type. The coefficient map $(L, v) \mapsto P_{L,v}$ is nonzero on this component (evaluate the product of leading minors at the standard rectangular inclusion matrix). It is equivariant by the preceding calculation. The component is irreducible for the product of compact groups, so its image is exactly that summand. This also identifies the source's chosen all-positive joint harmonic:

its highest-weight line is the product of leading minors, of the weight just computed. The one-dimensional highest-weight line follows from multiplicity one and ordinary highest-weight theory, and the product evaluates to one at the standard rectangular inclusion after compatible row order. Any nonzero normalized joint harmonic of that type is therefore a nonzero scalar times the corresponding $P_{L,v}$. The scalar in the compact lift theorem is allowed to depend on that choice, not on the finite datum, the automorphic form, the twist parameter with the fixed weights, or the points.

Theorem 17.4 (Exact compact class-sum lift for arbitrary polynomial datum)**.** *Let $K \subset K_0$ be the specified finite source level, T a linear functional on W, and let $F_{\phi,T}(x,g) = T(\rho(x^{-1})\phi(g))$ be the scalar coefficient of an algebraic form. Denote the algebraic class-sum lift at level K by $k_f\phi$, namely*

$$(k_f\phi)(h) = \int_{\Gamma\backslash G_f}^{\operatorname{vol} K=1} J_c(\omega(g,h)f)\phi(g)\,dg.$$

For every polynomial P there exists a linear functional T' on W', depending on P, T, c, K and the fixed weights, such that the actual theta-integral lift by $v_P \otimes f$ of $F_{\phi,T}$ is $F_{k_f\phi,T'}$. For $P = P_{L,v}$ one can take

$$T' = \frac{c^{-B_\kappa} T(v)}{\dim W\,[K_0 : K]} L. \tag{17.7}$$

The result remains valid for every vector in the integrable real Fock model in place of v_P.

Proof. All rational quotients for the definite groups are compact, and the integrands for fixed finite data are continuous with open finite stabilizers. For a polynomial $P_{L,v}$ the compact-real substitution identity and (17.6) express the kernel as

$$c^{-B_\kappa} L\rho'(y^{-1}) J_c(\omega(g,h)f)\rho(x)v.$$

Now average the product of this kernel and $T(\rho(x^{-1})\phi(g))$ over the real compact group with probability Haar measure. Schur averaging applied to the rank-one operator vT gives $T(v)I_W/\dim W$, so the result is $c^{-B_\kappa}T(v)(\dim W)^{-1}L\rho'(y^{-1})J_c(\omega(g,h)f)\phi(g)$. The remaining finite quotient integral has volume $K_0 = 1$ and is therefore $[K_0 : K]^{-1}$ times the class integral used to define $k_f\phi$. This gives (17.7).

For other polynomials use the multiplicity-free decomposition above. The orthogonality relation for two unequal compact irreducible types gives zero on every summand except the one spanned by the $P_{L,v}$. On that summand sum the functionals (17.7). This gives a single T' simultaneously for all twists with the prescribed weights, all finite data and all source algebraic forms; the projection depends only on the real compact actions.

To extend to the integrable real model, its bounded integrated pairing with the global theta functional is continuous. Polynomial Gaussian vectors are dense: the translated Gaussian is $e_\infty(y)e_\infty(w)\exp(\bar{w}\cdot y)$, whose Taylor polynomials converge in the integrable norm by domination by $e^{C\|w\|}e_\infty(w)$. Finite linear combinations of translates of the vacuum are dense from the reproducing integral (approximate its Bochner integral by finite simple sums in the integrable norm). The compact isotypic projection has finite-dimensional range, spanned by the indicated polynomial component, hence is closed. Since it is bounded as an integral over the compact group and vanishes on the complementary polynomial types, its range remains that same finite-dimensional component. The same proof therefore applies to the projected real vector. This proves the full assertion without a limiting choice of functionals depending on f. □

For the particular normalized `jointHarmoniceaebll'11` in the source, the same argument gives its scalar β of `exists_lift_ofAlgForm_eq_smul`. Namely every polynomial on this

definite arrow is harmonic for the lowering operators of both groups (their real Lie algebras are compact, and in the all-positive block the noncompact contraction terms are absent). The prescribed compact highest weights from the two HC parameters, with the lifting-weight condition, select precisely the component $P_{L,v}$ above. Its highest-weight space for the two compact groups has dimension one, and the vector P_{L_+,v_+} using their indicated highest coefficient vectors is nonzero by rectangular Schur evaluation. Thus the specified nonzero joint highest vector differs from it by a scalar, fixed by the chosen archimedean normalization and independent of the finite data, the level, and the twist with those weights. Inserting this scalar in (17.7) gives the source's asserted $\beta/[K_0 : K]$. This establishes `exists_lift_ofAlgForm_eq_smul` and `isThetaLift_thetaKernel` of `Lib/Weil/CompactLift/RallisFormula.lean` directly for the definite steps. The usual identification between the type-projected automorphic forms and algebraic forms in `Lib/Adelic/CuspForms/AlgebraicForms.lean` follows by inverting the displayed matrix coefficient: average $F(x, g)\rho(x)v_+$ over the compact probability measure and multiply by $\dim W$. For the normalized highest vector v_+ the result is precisely $\phi(g)$ (the Schur projection acts as $I_W/\dim W$). In particular this identifies the real coefficient map by the same Gram form, the precise volume convention, and the conjugate Weyl action used in the rest of the monograph.

17.6 The even-stage coefficient identification

The precedingly calculated result also applies to the two-stage integral in `New/Seesaw/DoubleLiftValue.lean`. Recall that the arrow product on the Hom spaces is obtained by transposing the first arrow by Hermitian adjoint and stacking it with the second; its target is the space with the first sign reversed. This is an isometry of the Hermitian Hom spaces, with the first bicharacter reversed according to its sign. The Heisenberg representation transports accordingly. The theta functional of the product is a nonzero scalar times the product of its two theta functionals: both are rationally invariant on the product, their nonzero nature follows from the theta mean identity, and uniqueness of the invariant functional gives the scalar. The norm-one mean normalization gives a scalar of modulus one if all three quotient means are normalized compatibly; it suffices here that it is independent of the finite factors and weights. Consequently for separated real polynomial vectors on the two arrow factors, the theta-integral expression of $\texttt{integral_theta_eq_lift}_2$ equals, up to this single scalar, the composition of two compact integrals as in Theorem 17.4. The automorphism interchanging the first variables and their conjugates is complex-linear on polynomial coefficients when one changes the signs of that factor in the Fock model. Thus polynomial data in the stacked variable space are linear combinations of products of polynomials in the disjoint arrow-variable sets. Finite tensors are spanned by pure tensors as well. Applying the result twice and adding the resulting functionals proves $\texttt{integral_theta_eq_lift}_2$ with one functional T'' on the second target, simultaneously for all the indicated data and parameters. For a general integrable real vector one may take the compact isotypic projections just used in the proof of Theorem 17.4 on both polynomial factors. Their ranges in the product are finite dimensional (the tensor product of the prescribed finite summands), hence closed, and polynomial density gives the same result on all integrable vectors. Compactness of the two definite rational quotients and continuity of the theta functions justify the interchange of their integrations.

Now consider the iterated stages P_k of `Data.unfoldStage`. The rational orbit, finite support and arrow-pair evaluation were proved in Chapter 16. For the first even step the initial generic Bessel function restricted to the definite middle group has exactly its stated normal form, so its dependence on the column is a linear functional of that column's compact coefficient vector; the initial finite translate is the one in the normal form when the stage-zero

point belongs to the principal level. This is the Bessel normal-form and admissibility input of Chapter 22; the assertion here is conditional on that input for the actual datum, and does not itself prove existence of the coherent Bessel packet. Suppose inductively P_{2i} is a compact coefficient of the first i paired lifts. The one-stage integral identities of Chapter 16, applied twice, express the evaluation at the next even group using the geometric arrow-pair zero term and its finite evaluation pairEvalY. The archimedean vector so obtained is determined by the fixed real evaluation points and the two arrows (its possible common scalar is fixed there); the finite vector is exactly the evaluated datum of the chapter. By the compact result of the preceding paragraph the integral of this real vector is one linear functional of the double algebraic lift. It is independent of the finite stage points, the twist with prescribed weights and the column except for the prescribed fixed real evaluation points. Induction proves the equality of functions

$$P_{2i+2}(\mathrm{incl}_X(g)(x, y_{2i+2})) = F_{\mathrm{chain}_2^{i+1}(\text{evaluated datum})\phi,T''}(\mathrm{emb}_X\, g)$$

under precisely the arrow-pair, stage, and initial-level conditions of `Data.unfoldStage_even` in `New/Seesaw/UnfoldingStages.lean`. If a common normalization scalar is zero, the same equality holds with the zero functional. Crucially one does not require that each unprojected individual arrow datum be a highest joint harmonic. The arbitrary-real-vector compact calculation absorbs all the archimedean projections into the final functional and is sufficient for this use of the seesaw. The genuine noncompact discrete-series type and Bessel normal-form existence remain a separate dependency in the spanning part, not part of this compact argument.

17.7 Integral branching with a controlled small diagram

We prove the precise integral map used at the end of the chain, keeping track of why very long first rows of the diagram need not contribute to its norm. The result also applies to the interlacing maps used elsewhere in the monograph. All polynomials, tableaux and Gram forms in the following proof are those of Chapter 5; for several places take the products in disjoint polynomial variables and the corresponding tensor-product vector spaces.

Let the larger rank be $N = n + 1$, and write its diagram as $\kappa = \gamma + \kappa'$. Let the smaller diagram have column multiplicities $\kappa_0(h) = \gamma(h + 1) + \kappa'(h)$ for $1 \le h < n$, and determinant exponent $e_0 = e + \kappa'(n)$. Assume $\kappa'(h) = 0$ for $h < q$. Put

$$L_0 = \sum_{\tau} \sum_{q \le h < n} h\kappa_{0,\tau}(h), \qquad b_0(\tau) = e_\tau + \sum_{1 \le h \le n} \gamma_\tau(h).$$

The number L_0 measures only the small remaining high columns. Let ev_Y denote the evaluation column of the source's integral minor basis at a matrix Y, and write $P_v(X) = \mathrm{ev}_X^T\, Gv$ for its Gram-polynomial map. The Gram-polynomial correspondence, tensor embedding and hook number used below were proved in Chapter 5.

Theorem 17.5 (Controlled integral branching map)**.** *There is a nonzero integer matrix B such that over every characteristic-zero field*

$$B\rho_{\kappa,e}(\mathrm{diag}(g, s)) = \left(\prod_{\tau} s_\tau^{b_0(\tau)}\right) \rho_{\kappa_0,e_0}(g)B,$$

where here $\rho = S_\kappa \det^e$ denotes the algebraic weight. Moreover B spans the intertwiner space for the compact subgroup $\prod_\tau U(n)$, and in the two specified Hermitian Gram norms

$$\|B\|_{\mathrm{op}} \le (L^L)^3 \qquad (L \ge L_0)$$

with $0^0 = 1$. In particular this is the bound on the exact integral map, not merely on a rationally scaled map.

Proof. Use the product polynomial $F(Y)$ of leading minors of the large array, taking the added row first (with the other rows in order) for the γ columns and last for the κ' columns. Each factor is a Gram polynomial of a normalized extremal evaluation vector for its own weight. The Cartan isometric embedding thus gives $F = P_\xi$ with $\|\xi\| \le 1$: the product of Gram polynomials corresponds to the adjoint of the isometric embedding applied to the tensor product of their vectors. The polynomial is nonzero.

First we determine the necessary small integer multiplier. On the smaller rank, write f for the cyclic row order that starts with row 2 and ends with row 1. Define

$$P(X) = P_{v_{\gamma(\cdot+1),f}}(X)\, P_{v_{\kappa',\mathrm{id}}}(X)$$

using the diagrams below height n. Here each Gram polynomial of an extremal evaluation vector is the product of its indicated leading minors, with the row order specified. This is a polynomial of weight κ_0 and normalized extremal factors. Let β consist of the columns of $\gamma(\cdot+1)$ of height less than q, put $\alpha = \kappa_0 - \beta$, and let $C : W_{\kappa_0} \hookrightarrow W_\beta \otimes W_\alpha$ be the integral split Cartan embedding. It sends ev_X to $\mathrm{ev}_X \otimes \mathrm{ev}_X$; its integer matrix rows are the coordinates of products of basis minor polynomials. The Gram kernel factorizes into the product of the two Gram kernels by the defining product of minors, proving the isometry on evaluations, which span. Every semistandard tableau of the combined shape can have its columns partitioned into the prescribed heights of the two summands: keeping their original order gives a semistandard tableau of each shape, since each column is strictly increasing and each remaining row is a subsequence of a weakly increasing row. Hence every combined basis polynomial is a product of the corresponding two basis polynomials. In the matrix of C these rows are precisely the coordinate rows of an identity matrix, giving an integral left inverse. The product polynomial P corresponds to a vector w with $\|w\| \le 1$.

For each $s < q$ take the set of the first s rows in the cyclic order. Every kept column has height at least q and therefore includes these rows among its first standard rows (they are among rows $2, \dots, s+1$). The γ factors attain the maximum possible number of occurrences of those rows. The torus weight of P therefore attains the full bound $\sum_h \kappa_0(h) \min(h, s)$ on every such set. In the tensor product $W_\beta \otimes W_\alpha$, the bound is the sum of the individual bounds. Hence in every weight term of Cw both bounds must be attained. For the factor β, which consists only of columns of height less than q, equality for all $s < q$ forces precisely the first h rows of the cyclic order in each column of height h. This factor is thus the one-dimensional extremal weight line, and $Cw = v_{\beta,f} \otimes z$ for some $z \in W_\alpha$.

By the isometric Gram-polynomial identity, $P = P_{v_{\beta,f}} P_z$, so canceling the nonzero first factor shows that P_z is exactly the product polynomial with the β factors removed. It has integer polynomial coefficients and is in the integral Schur polynomial module (its expansion in the integral basis is given by the straightening identities). By Lemma 5.7, $h_\alpha P_z$ is the Gram polynomial of an integer coordinate vector: the Fischer matrix is integral and satisfies $G_\alpha H_\alpha = h_\alpha I$. The integral left inverse of C proves that cw is an integer coordinate vector for $c = h_\alpha$. The degree of α is L_0 and the hook bound of Chapter 5 gives $1 \le c \le (L^L)^3$. This is the denominator calculation in `exists_smul_weylKernel_mul_eq_toPoly`. It explains the absence of the potentially much larger degree of β from the bound.

We next construct B itself, not just its adjoint on one weight line. For a smaller square array X and larger square array Y set

$$K(X, Y) = F(\mathrm{diag}(X^T, 1)Y).$$

If the last row of Y is zero beyond its first entry, expansion of each moved-row minor along that row gives the product of its first entry, the determinant factor $\det(X)^{\kappa'(n)}$, and the smaller polynomial P evaluated after row substitution by the first n rows of Y. More explicitly, write A for the first n rows and first n columns of Y in the specified row order, and u for its added-row first entry. Then in a single place one has

$$K(X,Y) = u^{\sum_h \gamma(h)} \det(X)^{\kappa'(n)} \det(A)^{\kappa'(n)} P(A^T X).$$

Indeed expansion of a moved minor of height h in its first row leaves the subarray of $X^T A$ with first $h-1$ rows and columns $2,\dots,h$; transposition gives the indicated leading minor of $A^T X$ in cyclic order. The unmoved minors have first h rows and columns; the unmoved minor of height n is $\det(X)\det(A)$. The signs here use exactly the order with the added row first for moved columns. Since cP has an integral Gram vector, Schur equivariance under the integral matrix of this substitution yields $cK(X,Y) \in \det(X)^{\kappa'(n)} \operatorname{im}(P_\cdot)$ with coefficients in the polynomial ring of the entries of Y, in that special form of the last row.

For an arbitrary generic array put $u = Y_{N,1}$. Multiply each column of Y by u and for columns $j > 1$ subtract $Y_{N,j}$ times the first column of the original array. The transformed last row has zeros beyond its first entry; the right-hand column operation is upper triangular with all diagonal entries u. Every leading minor of height h is thus multiplied by u^h. Hence the transformed polynomial cK is the original cK multiplied by the monomial $u^{\sum h\kappa(h)}$. Since membership in the displayed integral Gram-polynomial submodule can be checked coefficientwise in the entries of Y, multiplying by this monomial and shifting the coefficient indices back proves the same membership for cK itself. There is no division by an integer or by a polynomial value. The same construction in each place proves the assertion for their product.

As a function of Y, this polynomial belongs to the integral Schur polynomial module of the larger diagram, since the leading minors have exactly its right-upper-triangular semi-invariance. Extracting coefficients in its integral minor basis leaves the previously stated smaller Gram-polynomial membership in every coefficient. For a justification of this extraction over an arbitrary coefficient module, the standard minor basis has leading monomials of coefficient ± 1 distinguished by the standard tableaux; successively subtracting the coefficient of a leading monomial is an integral coordinate extraction on its span. Equivalently this is the integral spanning-and-coordinate lemma for the basis used in Chapter 5. Consequently an integer matrix B satisfies the identity of polynomials

$$cK(X,Y) = \left(\prod_\tau \det X_\tau^{\kappa'_\tau(n)} \right) \operatorname{ev}_X^T G_0 B \operatorname{ev}_Y .$$

This is the source's `exists_int_branching_of_weylKernel` (`Lib/GLn/Branching/Branching.lean`).

Evaluate this identity after substituting $Y \mapsto \operatorname{diag}(g,s)Y$. In F the last-row scalar appears exactly $\sum_h \gamma(h)$ times; the remaining substitution on the right is $X \mapsto g^T X$. The determinant factor gains $\det(g)^{\kappa'(n)}$ and the Gram-polynomial transformation gives $S_{\kappa_0}(g)B$. Comparing evaluation vectors (which span over every characteristic-zero field) proves the equivariance identity in the theorem including the determinant twist. Taking $X = I$ gives $B^* v_{0,+} = c\xi$ under the two Gram forms. Indeed the Gram-polynomial identity becomes the polynomial cF and all matrices and chosen extremal vectors are real in these coordinates, so the bilinear transpose identity gives the stated Hermitian adjoint. Since BB^* commutes with the smaller irreducible unitary representation, it is a nonnegative scalar times the identity. Therefore $\|B\| = \|B^* v_{0,+}\| \le c$ (the smaller extremal evaluation vector has norm one). Also $cF \neq 0$, so $B \neq 0$.

For multiplicity one it is enough to apply the ordinary compact branching rule: the partitions interlace because for the γ columns the height falls by one and for the κ' columns it stays fixed, so the removed boxes form a horizontal strip. To see the branching rule directly, use the semistandard tableau character formula. Upon adjoining the variable for the largest row label, all boxes carrying that label form a horizontal strip; removing them leaves a standard semistandard tableau of the smaller diagram. Thus the character is the sum over interlacing smaller diagrams of their characters times the prescribed power of that variable, each once. By independence of irreducible characters and complete reducibility of compact representations, every such summand has multiplicity one. This remains true under determinant twists and for a product of embeddings. Restriction to the compact subgroup without the final scalar still gives multiplicity one for this particular smaller determinant weight, since the scalar exponent is forced by total degree. Thus B spans the compact intertwiner space, finishing the proof. □

This gives the precise integral matrix, scalar character, and L^{3L} norm in `exists_branching_matrix`. The row order in the construction is the one encoded by `newRowOrder` in `Lib/GLn/Branching/BranchingVector.lean`; changing it by the index equivalence permutes the integral bases and leaves the Gram formula invariant. For example the value q can be the initial rank k in `Data.branching`; the numerous smaller columns outside the kept diagram are all in the extremal factor β and therefore do not enter the bound. The same argument applies to the branching pairing between opposite weights after the integral duality isometry of the coefficient spaces, as in the definition `exists_branching_pairing`.

17.8 The last stage as an actual compact branching integral

Let the smaller group be the standard group $U(Y_m)$ and the larger group $U(X_m)$, identified by the index equivalence of `Data.lastIndexEquiv` and its block embedding j. For the finite form ψ on X_m and a linear functional t on its coefficient space, consider its scalar coefficient restricted to $j((x,h),1)$. A scalar form on the smaller group is of the form $F_{\phi,L_+}(x,h)$ from Theorem 17.4. The integrand of the last stage is its conjugate multiplied by $t\rho'(j(x)^{-1})\psi(j(h))$. On the compact real group the conjugate of the first coefficient is $\langle \phi(h), \rho(x)v_+\rangle_0$. Consequently its compact integral against the second coefficient is contraction with the matrix

$$A_t = \int_{U(Y_m)_\infty} \rho(x)v_+ \, t\rho'(j(x)^{-1})\, dx,$$

where the measure is probability Haar measure. The integral exists and changing variables immediately gives $A_t\rho'(j(x)) = \rho(x)A_t$. Multiplicity one in Theorem 17.5 therefore gives $A_t = C_tB$, with the same integral B chosen for the branch map. The finite quotient integral supplies exactly the class integral of $\langle \phi(h), B\psi(j(h))\rangle_0$ at the relevant level. Its measure differs from maximal compact volume one only by the known index, which is independent of ϕ, ψ. The prescribed scalar $\prod_\tau \tau(\bar\pi)^{-\,\mathrm{branchExp}(\tau)}$ in `Data.restrictToY` is nonzero and independent of the forms, so it can be inserted into $R\psi$ and its inverse into C_t. This proves the source identity `integral_algFormCoeff_blockEmb`, and the last-stage pairing

$$\int_{[U(Y_m)]} F(j(k))\,\overline{x(k)}\,dk = C_t'\langle x, R\psi\rangle$$

for every x of the prescribed type. Here R is the displayed restriction followed by the integral branching map and the coefficient-to-form identification of Theorem 17.4; its formula is independent of t. This is the second and more precise assertion of `Data.exists_last_stage_map` in `New/Filling/DegreeBounds.lean`. The first assertion follows from the definitions of

the analytic and algebraic end maps: $\iota(\vartheta, y)$ is the algebraic form of the analytic last-stage chain, and $B^{\Psi}\phi$ is the same restriction restrictToY applied to the algebraic chain. The coefficient isomorphism of Theorem 17.4 takes R to that restriction, so equality of the two analytic forms gives equality of their algebraic coefficient forms. The source uses a scalar C'_t whose possible vanishing is harmless for this pairing; it does not use an unproved nonzero value of an individual branching period.

At the p-adic level the identity in Theorem 17.5 also controls the matrix in padded gauges at the added row: for a scalar s in the added coordinate it introduces exactly the scalar $s^{b_0(\tau)}$, not a norm loss of the entire larger diagram. This is the source's mechanism for the local normalization by $\bar{\pi}^{-\,\mathrm{branchExp}}$ and its cancellation with the last-level powers. Together with the preceding compact coefficient calculation it discharges the last-stage comparison, still using the actual geometric and finite data supplied to it.

17.9 Uniform counting of the definite classes

We supply the counting estimate `Space.card_classes_le` in `New/Core/ClassCountBound.lean`, since mere compactness gives no useful uniform bound. Write $L_g = E^n \cap g\widehat{\mathcal{O}}_E^n$ for the self-dual Hermitian lattice of a finite adelic unitary element g. Let N_1 belong to the different of E. In the archimedean Euclidean space $\mathbb{C}^{fn}$ the dual of $2N_1^{-1}L_g$ for the real Euclidean pairing is $N_1\mathfrak{D}_E^{-1}L_g$ by the trace-dual identity (the factors of two cancel). It is contained in L_g; every nonzero vector of L_g has squared length at least one, as the trace of a nonzero totally positive algebraic integer. The covering-radius estimate proved in Chapter 22 therefore gives a covering radius at most $(2fn)^2$ for $2N_1^{-1}L_g$. If fewer than n E-linearly independent lattice vectors have already been chosen, choose an archimedean vector of length $2(2fn)^2$ perpendicular to the real span of their E-linear subspace. The covering-radius estimate finds a vector of $2N_1^{-1}L_g$ within $(2fn)^2$ of it, necessarily outside that span, of norm at most $3(2fn)^2$. Scaling back gives a further vector of length at most $6N_1(fn)^2$. Induction produces $Y_g \in \mathrm{GL}_n(E)$ with $A_g = g^{-1}Y_g$ integral at every finite place and

$$|\tau(Y_{g,ij})| \le R_0 = 6N_1(fn)^2$$

for every embedding and entry. This proves `Space.exists_short_basis_stdSpace`. The rank-zero case needs only the single class and is immediate.

We also use scalar approximation with the same bound. For a finite adele a and a prescribed archimedean vector z, weak approximation gives a rational point r with $a - r$ integral at all finite places (all but finitely many integrality conditions are already automatic). Apply the covering-radius estimate for $2N_1^{-1}\mathcal{O}_E$ to $(z - r)/N_1$ and scale back; the adjustment is in $2\mathcal{O}_E$. This gives $v \in E$ with $a - v$ integral and $|z_\tau - \tau(v)| \le 4N_1 f^2$. In particular for the entries of A_g^{-1} and $z = 0$ obtain a rational matrix Z_g with

$$A_g^{-1} - Z_g \text{ integral}, \qquad |\tau(Z_{g,ij})| \le 4N_1 f^2.$$

Define the code of the class to consist of the two rational matrices $H_g = Y_g^* Y_g$ and $H_g Z_g$. The first is integral since $H_g = A_g^* A_g$ at all finite places; the second is integral since $A_g Z_g = I - A_g(A_g^{-1} - Z_g)$ is integral. Their conjugate entries have absolute value at most R_0^3: the respective bounds nR_0^2 and $n^2R_0^2(4N_1f^2)$ are at most R_0^3. If two codes coincide, then $H_{g'} = H_g$, $Z_{g'} = Z_g$, and $\gamma = Y_{g'}Y_g^{-1}$ is rational unitary. Moreover

$$g'^{-1}\gamma g = A_{g'}A_g^{-1} = A_{g'}Z_g + A_{g'}(A_g^{-1} - Z_g)$$

is integral, since $A_{g'}Z_g = A_{g'}Z_{g'}$ is integral. Its inverse, its Hermitian adjoint, is also integral. Hence the classes coincide.

For distinct algebraic integers the difference has some complex embedding of absolute value at least one, by the field norm. Consequently open product balls of complex radius $1/2$ in the f embeddings about them are disjoint. If all embedding absolute values are at most R_0^3, these balls lie in the product ball of complex radius $R_0^3 + 1/2$. Its relative volume gives at most $(1 + 2R_0^3)^{2f}$ integers. Applying this to the $2n^2$ entries in the injective code proves

$$\#\mathrm{StdClasses}(K_0) \le (1 + 2R_0^3)^{4fn^2}.$$

At a level containing the principal level of rational modulus G, the maximal-compact double cosets split into at most $[K_0 : K(G)] \le G^{2fn^2}$ cosets each, by reduction of their matrices over $\mathcal{O}_E/G$. This proves the requisite level bound with no assumption that the reduction is surjective.

For clarity we check the exponents used in `Data.card_stdClasses_le_exp_mul_log`. Put $X = \log x$ and $\ell = \log b$. The scale imposes

$$10^{12} P.c \le \log \ell \le X/100, \qquad 10^6 X \le \log x_1 \le 10^{-6}\ell.$$

The magnitudes of $N_1 = P.N_0 \,\mathrm{towerDeg}(s)$ and f give $N_1, f \le x^{11.1}$: the bounds before rounding are $e^{P.c} x^{11} \ell^{1/5}$ (use $P.N_0 \le P.c$ for the former). The rank satisfies $n_m \le e^{12} x^{5.5}$, hence even $2n_m \le x^{5.6}$. The extremely large lower bound on X absorbs every fixed real constant in these inequalities. Thus $R_0 \le x^{45}$ and $4fn^2 \le x^{23}$. Also $\log(1 + 2R_0^3) \le 136X \le x$, so the logarithm of the preceding class bound, including the level index, is at most $x^{24}(1 + \log G) \le x^{29}(1 + \log G)$. In particular for $G = g_{\mathrm{prod}} N$ the source magnitudes give $\log g_{\mathrm{prod}} \le 22\ell$ and $\log N \le e^{32}\sqrt{x}\ell$. As $\ell \le x^{1/100}$, these imply $1 + \log G \le x$ and prove the bound $e^{x^{30}}$ at the actual column levels. This proves both class-count assertions in `New/Columns/SizeEstimates.lean`.

17.10 The exact size budget along the lift chain

We finish the arithmetic check in `Data.weylNormSq_Bmu_le` of `New/Columns/LocalStatement.lean`, using the conjugate-jet bound, class count and integral branch just proved. Retain $X = \log x$, $\ell = \log b$ and write $\mathcal{B} = fa\ell^3$ for the error budget. The source bounds $xb \le a \le exb$, $m \le 2\sqrt{x}$, $b_1 \le e^2 b$, $b_2 \le e^2 bx/x_1$, $h_2 \le e^{25} x^{40}$ and $h \le e^{27} x^{40}$. In particular $h_2, h \le x^{40.1}$, $n_m h_2^2 \le x^{86} \le b$, and $b_1 + b_2 + n_m h_2^2 \le 16b$. Since $\Omega \in [x^{-13}, 1]$ and $\log g_{\mathrm{prod}} \le 22\ell$, the jet scale parameter satisfies $u = \log(g_{\mathrm{prod}}/\Omega) \le 22\ell + 13X$.

At stage j of one translated chain the datum is a single translate of the integral vacuum, to which (17.4) applies with $B = 1$, $L = a + 1$, $c = \bar{\pi}_G$. All $d_{\tau i}$ are at most $2a$: the row bounds are

$$d_{\tau 0} \le \kappa_\tau(1) + b_1 + b_2 + h_2^2, \quad d_{\tau i} \le w_0(\tau) + h_2^2 \ (0 < i < k), \quad d_{\tau i} \le h_2^2 \ (i \ge k),$$

with $w_0(\tau) \le a$ and the error in any row at most a. Here $\kappa_\tau(1) = a$ at a top and zero otherwise. The leading terms of the logarithm of the squared norm in (17.4) are $\sum J(d_{\tau i})$, $J(l) = \log(l!) + lu$. For integers $l \le r + e$ with $r + e \le 2a$,

$$J(l) \le J(r) + e(\log(2a + 1) + u) \le J(r) + 24e\ell.$$

The first inequality uses $u \ge 0$ and $(r + e)! \le r!(r + e)^e$; the second follows from the bounds on a, u and $10^{12} X \le \ell$. Use baseline $r = \kappa_\tau(1)$ in row zero, $w_0(\tau)$ in the next $k - 1$ rows and zero elsewhere. Enlarging the principal contribution by $J(w_0(\tau)) \ge 0$ gives exactly the convenient upper bound

$$\sum_{\tau, i} J(d_{\tau i}) \le \sum_{\tau \in \mathcal{T}} J(a) + k \sum_{\tau} J(w_0(\tau)) + 384 fb\ell.$$

The geometric sum of the small row errors is at most $f(b_1 + b_2 + n_m h_2^2) \le 16fb$ per stage. No main row has been charged to the error budget.

The remaining square-norm exponent in (17.4) per singleton term is at most

$$2P.c(N_1 nn' + 1)^2 + 2fnn' \log(2a + 4) + 2fn_m.$$

Indeed the contribution of $\sum d_{\tau i}/L$ is at most $2fn_m$. For $n, n' \le n_m \le x^{5.6}$, $N_1, f \le x^{11.1}$, $P.c \le \log \ell \le X$ and $\log(2a + 4) \le 2\ell \le 2x^{1/100}$, the first term is at most x^{46} and the other two terms are at most x^{24} each. Their sum is at most x^{47}. The product of class sums along m stages adds at most $2mx^{30}$ to the squared logarithmic norm, because the class-sum weights are reciprocal stabilizer orders and hence at most one. Joint invariance of the complete datum permits choosing the same finite class representatives in the expanded sum; after expansion, regardless of whether a single summand is itself invariant, each evaluation is a finite iterated class sum and is bounded by the product of the stage bounds. This is the elementary expanded-sum argument of `exists_chain_eq_sum`; no exchange with an infinite family of class choices is needed. The final sum over the support of the good datum adds at most $2x^{60}$, since the sum of absolute values of its conjugated coefficients is bounded by $e^{x^{60}}$ for the trivial twist (one of the actual hypotheses of `IsGoodDatum`).

For the final restriction, the kept branching degree satisfies $L_0 \le fn_m^3 h$: there are at most n_m heights per place, their heights are at most n_m, and after the first $k-1$ small heights have been separated from the diagram the prescribed column multiplicities are at most $n_m h$. This is the kept-column definition in `ColumnDefs.lean` (the inequality uses the reduced-half-gap bound). Consequently $L_0 \le x^{68}$, and Theorem 17.5 gives $\log \|B\|^2 \le 6L_0 \log L_0 \le x^{70}$. Its matrix entries are integers, and the branch scalar $\prod_\tau \tau(\bar\pi)^{-\operatorname{branchExp}(\tau)}$ has squared absolute value under every conjugate equal to $N^{-\sum_\tau \operatorname{branchExp}(\tau)}$, because $\pi\bar\pi = N$. This proves `Data.weylNormSq_restrictToY_le`. The elementary weight bound $\operatorname{branchExp}(\tau) \ge \kappa_\tau(1) - 2b$ therefore yields the main correction $-|\mathcal{T}|a \log N$ and error $2fb \log N$. In the following calculation even the weaker $\log N \le x^{0.6}$ suffices.

Altogether, apart from an input squared logarithmic norm η, the principal exponent is

$$m\left(\sum_{\tau\in\mathcal{T}} J(a) + k\sum_\tau J(w_0(\tau))\right) - |\mathcal{T}|a \log N.$$

Check explicitly that the remaining terms fit the claimed budget:

$$\begin{aligned} E &\le 384mfb\ell + mx^{47} + 2mx^{30} + 2x^{60} + x^{70} + 2fbx^{0.6} \\ &\le 768\sqrt{x}\, fb\ell + x^{75} + 2fbx^{0.6}. \end{aligned}$$

Here $x^{75} \le b$ by $\ell \ge 10^{12}X$. Dividing the last bound by $fa\ell^3 \ge fxb\ell^3$ gives at most $768/(\sqrt{x}\,\ell^2) + 1/(x\ell^3) + 2/(x^{0.4}\ell^3) < 1$, since $\ell > 1$ and the scale implies $x > 10^8$. Thus the remaining exponent is at most $\mathcal{B}$. The exact identity (14.21), obtained by expanding the prime asks and collecting their forced parts, expresses the displayed principal exponent as $h_0 + 2\sum_{p\in\mathrm{nonsplitPrimes}} \mathrm{gain}(p) \log p$. It follows that

$$\|\sigma \mathsf{B}_1^\Psi v(g')\|^2 \le \exp\left(\eta + h_0 + 2\sum_{p\in\mathrm{nonsplitPrimes}} \mathrm{gain}(p)\log p + \mathcal{B}\right).$$

This proves (14.22) with the actual constant-one budget and the uniformity in the complex conjugate. It uses the exact good-datum hypothesis, in particular its invariance and coefficient size, and is not a proof of the existence of those data. The point of retaining the small degree in branching and the row singular-value estimate in the jet norm is that replacing either by a bound on the whole diagram would no longer satisfy the last displayed budget.

17.11 Status of the upper-bound proof obligations

Here *proved* denotes the conventional argument for the specified source interface, assuming its explicit hypotheses and results previously proved or referenced in this monograph. It does not denote a new formal compilation of the Lean project.

PROVED — finite–real arithmetic theta realization.
`Space.entireTheta_translate_finiteImage`, `semichar_finiteImage`, `Space.exists_entireTheta_eq`, `Space.jet_jetSeries_eq_theta`: positive standard space, rational vectors and nonzero jet scale. Entire-function realization, its injectivity, multiplier law and exact Taylor normalization are proved in Proposition 17.1. This supplies the missing map in Chapter 12.

PROVED — conjugate generators and finite conjugacy.
`Parameters.exists_finitePart_of_isTightFrame`, `Parameters.exists_isJetConjugate_generator`, `Parameters.exists_semilinear_isJetConjugate`: use the positive CM theta lattice, marked-section reciprocity of Theorem 16.5, and the integral vacuum. Theorem 17.2 and Proposition 17.3 extend the generator identity to all finite translates and Weil operators, agreeing with their actual multipliers and uniform norm. The classical elliptic-curve comparison required for these arguments has the independent references in Section 8.3; it is not a consequence of the finite-model calculation.

PROVED — upper-bound definite coefficients.
`exists_lift_ofAlgForm_eq_smul`, `isThetaLift_thetaKernel`: unequal definite ranks, matching lifting weights, nonzero c, and prescribed compact types. Theorem 17.4 gives the exact coefficient and level factor using compact Schur orthogonality. If instead one uses the multiplicity-free Cauchy decomposition as a reference, Enright–Willenbring, Annals of Mathematics 159 (2004), p. 370, (7.4.3)–(7.4.4), gives the same two polynomial actions and decompositions. This is not a reference for indefinite real theta lifts.

PROVED — even stages and last-stage comparison subject to actual Bessel input.
$\texttt{integral_theta_eq_lift}_2$, `Data.unfoldStage_even`, `Data.exists_last_stage_map`: Sections 17.6 and 17.8 prove the analytic–algebraic comparisons including product normalization, real coefficient map, class measures and final branching pairing. The initial Bessel normal form and existence of the admissible coherent packet are separate dependencies (Chapter 22); these must be discharged for the actual initial datum, not assumed merely because some Poincaré series exists. Noncompact harmonic/type identification is not proved by the compact argument.

PROVED — integral and quantitative branching.
`exists_smul_weylKernel_mul_eq_toPoly`, `exists_int_branching_of_weylKernel`, `exists_branching_matrix`: horizontal-strip interlacing and the lower bound q on kept heights; Theorem 17.5 proves the integral equivariant map with its norm in the Gram convention and multiplicity-one assertion. This is stronger than citing the complex multiplicity-one branching rule alone.

PROVED — arithmetic stability and the upper-bound size.
`Parameters.finHeisOp_mem_standardData`, `Parameters.omega_mem_standardData`, `Parameters.pairWeil_mem_standardData`, `Parameters.mem_raises_of_standardData`, `Parameters.mem_raises_of_dvd`: valuation at a coefficient embedding, standard integral module and the prescribed diagonal gauges. The finite vacuum sum and weak-approximation argument establish the stable lattice and gauge bound. `Parameters.norm_jet_le_of_isAdmissibleComb`, `Space.card_classes_le`, `Data.card_stdClasses_le_exp_mul_log`,

`Data.card_stdClasses_le_exp`, `Data.weylNormSq_restrictToY_le`, `Data.weylNormSq_Bmu_le`: the proved class count, jet operator bound and chain estimate of Sections 17.3, 17.9, 17.10 use the admissible scales and good-datum hypothesis as stated, yielding the exact $fa(\log b)^3$ error and all conjugates.

OPEN in this chapter — distinct Bessel/existence obligations.
This chapter does not prove the noncompact Bessel Fourier coefficient and coherent-packet construction needed on input to the even-stage argument. It also does not prove existence of good data independently of the geometric transfer and initial generic support: those belong to the seesaw and filling chapters and their own closure statuses. Such existence must not be inferred from a bound conditional on a good datum.

Part III

Spanning: Bessel and Poincare series

Chapter 18

Formal degrees, elliptic trace comparison and the row dimension

There is a global representation-theoretic issue in comparing the row space (on a unitary group which is not definite) with a space of algebraic modular forms on a definite group. A comparison of dimensions of the full spaces would be false without hypotheses; in particular the discrete series at infinity cannot simply be replaced by its compact counterpart. The source uses the central projection onto the good finite types, and compares only the corresponding Hecke traces. We give the exact formal-degree normalization, explain the trace comparison by changing the signs of elliptic classes, and prove the finite algebraic step that isolates the good traces. The construction of the actual discrete-series kernel and its reproducing property, the classical discrete-series character formula on the elliptic Cartan, and the Galois-theoretic compatibility of the automorphic labels are distinguished from the deductions proved here.

18.1 Parameters and the compact-dual Haar measure

For a rank n unitary group put $J = \mathrm{diag}(I_p, -I_q)$, with $p + q = n$, and write $G_J = U(J)$; its maximal compact subgroup K_∞ is $U(p) \times U(q)$. The notation K_∞ in this chapter is not the auxiliary imaginary quadratic number field. Let $\lambda = (\lambda_i)_{i=1}^n$ be the Harish–Chandra parameter, and $l = 2\lambda$ the doubled integer vector used in the source. For an admissible regular parameter one has $l_i \equiv n - 1 \bmod 2$. Its decreasing rearrangement $\lambda_1^\downarrow > \cdots > \lambda_n^\downarrow$ is used for the compact form, with highest weight

$$\mu_i = \lambda_i^\downarrow - \rho_i, \qquad \rho_i = \frac{n + 1 - 2i}{2}.$$

In particular μ_i are integers and $\mu_i \geq \mu_{i+1}$, since the positive differences of the λ_i are integers. The compact representation has dimension

$$\dim V_\mu = \prod_{i<j} \frac{\lambda_i^\downarrow - \lambda_j^\downarrow}{j - i} = \frac{\prod_{l_i > l_j} (l_i - l_j)/2}{\prod_{a=0}^{n-1} a!}. \tag{18.1}$$

Indeed the Weyl character of μ on the diagonal torus is the quotient $\det(z_i^{\mu_j + n - j}) / \det(z_i^{n-j})$. Its limit at the identity is the displayed ratio: set $z_i = e^{t u_i}$ for distinct u_i and expand the two determinants in powers of t. In the numerator the first nonzero term has total degree $0 + 1 + \cdots + (n-1)$ and equals the product of the Vandermonde determinants of (u_i) and of $(\mu_j + n - j)$, divided by $\prod_{a=0}^{n-1} a!$. The corresponding factor for (u_i) and the factorials cancel in the ratio. Finally $\prod_{i<j}(j - i) = \prod_{a=0}^{n-1} a!$. This is the polynomial `weylDimensionPoly`

in `Lib/DiscreteSeries/FlenstedJensen/RepresentationDefs.lean`. A simultaneous additive shift of all parameters changes the determinant character but leaves (18.1) unchanged.

We must identify the scalar implicit in the Haar measure. On G_J the form $Jg^{-1}dg$ is skew-Hermitian. Its Lebesgue density uses the n diagonal imaginary coordinates and the real and imaginary coordinates of the upper triangular off-diagonal entries. Divide this left-invariant density by the volume of the compact group $U(n)$ in the corresponding coordinates at $J = I_n$. The result is the compact-dual measure of the source. Explicitly a Cayley chart is

$$g(Y) = (I + \tfrac{1}{2}J^{-1}Y)(I - \tfrac{1}{2}J^{-1}Y)^{-1}, \qquad Y^* = -Y.$$

Putting $A = I - \frac{1}{2}J^{-1}Y$, its logarithmic derivative in a skew-Hermitian direction dY is

$$Jg^{-1}dg = A^{-*}\, dY\, A^{-1}.$$

This follows by differentiating $g(Y) = 2A^{-1} - I$, and using $Y^* = -Y$ and $J^* = J = J^{-1}$. The real determinant of congruence $Y \mapsto C^*YC$ on the real space of skew-Hermitian matrices has absolute value $|\det C|^{2n}$. To see this, use the singular-value decomposition of C: the unitary factors preserve the Euclidean form on that space and have determinant of absolute value one. For $C = \operatorname{diag}(c_1, \dots, c_n)$ with $c_i > 0$, the diagonal coordinates are multiplied by c_i^2, while each off-diagonal pair of real coordinates is multiplied by c_ic_j. The product is $\prod c_i^{2n}$. The Jacobian in the Cayley chart therefore equals $|\det A|^{-2n}$, which is precisely `cayleyDensity` of `Lib/DiscreteSeries/Haar/CompactDualMeasure.lean` for $J^2 = 1$. The chart omits the locus where $g + I$ is singular, a proper real-analytic locus of measure zero. The volume normalization is thus an exact comparison of two Haar measures, not merely a proportionality up to an unknown real scalar.

For $J = I_n$ the volume of $U(n)$ in these Lie algebra coordinates equals

$$\operatorname{vol}_0 U(n) = \prod_{a=1}^{n} \frac{2\pi^a}{(a-1)!}.$$

One may compute it successively using the first-column map $U(a) \to S^{2a-1}$, whose fiber is $U(a-1)$. At the identity, the first-column tangent coordinates consist of one diagonal imaginary coordinate and the $a - 1$ pairs of off-diagonal real coordinates; these are the ordinary Euclidean tangent coordinates of the unit sphere, of density one with respect to the indicated Haar volume element. The fiber coordinates are exactly the smaller skew-Hermitian coordinates. Invariance under $U(a)$ and the invariant quotient measure then give the recurrence multiplying by the ordinary sphere area $2\pi^a/(a-1)!$ (from polar coordinates in the Gaussian integral on $\mathbb{C}^a$). This matches `volU` and `isProbabilityMeasure_haarU` in `Lib/DiscreteSeries/Haar/HaarMass.lean`. In particular the measure of the compact form in this normalization is one.

18.2 Evaluation of the constant in the formal degree

For an irreducible unitary discrete series π let v be any unit vector. Its formal degree with respect to the preceding Haar measure is

$$d_\pi = \left(\int_{G_J} |\langle v, \pi(g)v \rangle|^2 \, dg \right)^{-1}.$$

The independence of v follows from Schur orthogonality for a square-integrable irreducible representation: integrating a rank-one operator against conjugation gives an operator commuting with the representation, and polarization gives the stated coefficient orthogonality. In

the compact case with probability measure this reduces to the elementary formula $d_\pi = \dim \pi$. The formal degree changes by the inverse of the scalar by which Haar measure is changed.

We use the classical Harish–Chandra formal-degree theorem: for a fixed real unitary group and Haar measure, its regular discrete series of parameter λ satisfy

$$d_\pi = C_J \prod_{i<j} |\lambda_i - \lambda_j|,$$

with a constant independent of the parameter (Harish–Chandra, *Discrete series for semisimple Lie groups II*, Acta Math. 116 (1966), in particular the formal-degree formula). Instead of relying on a convention for C_J, we evaluate it using one scalar holomorphic discrete series. This is also a check of the factor two implicit in the doubled parameter. The classification of the scalar holomorphic discrete series and the lowest K_∞-type formula $\Lambda = \lambda + \rho_n - \rho_c$ are the standard classical statements used in this calculation.

For $p, q > 0$ use the graph coordinate $Z \in M_{q,p}(\mathbb{C})$ on G_J/K_∞, where $I_p - Z^*Z > 0$. If g has blocks $(A, B; C, D)$, its action is $Z' = (C + DZ)(A + BZ)^{-1}$, and

$$I_p - Z'^*Z' = (A + BZ)^{-*}(I_p - Z^*Z)(A + BZ)^{-1}.$$

For the compact dual Grassmannian $U(n)/K_\infty$, the same graph coordinate uses $I_p + Z^*Z$ instead; it covers the complement of the subspaces whose projection to $\mathbb{C}^p$ is noninvertible, a subset of measure zero. Differentiate the graph action to obtain

$$dZ' = (D - Z'B)\, dZ\, (A + BZ)^{-1}.$$

A block determinant calculation gives $\det(D - Z'B)\det(A + BZ) = \det g$; one can multiply g on the right by the block lower triangular matrix with $(2,1)$ block Z and diagonal identities and then use the Schur complement. The absolute complex determinant of g is one. Thus the real Jacobian of $dZ \mapsto dZ'$ has absolute value $|\det(A + BZ)|^{-2n}$. Consequently the invariant quotient densities on the two graph charts are respectively

$$c \det(I - Z^*Z)^{-n}\, dZ, \qquad c \det(I + Z^*Z)^{-n}\, dZ.$$

Here the constant c is the same for both: at $Z = 0$ the horizontal tangent matrices in the two groups correspond to the same complex off-diagonal block dZ, differing only by the sign in the opposing block, hence with the same volume in the skew-Hermitian coordinates of the preceding subsection. The fiber K_∞ has the same tangent density on both sides, and we use its probability measure for both disintegrations. Since the compact quotient has total mass one,

$$c^{-1} = \int_{M_{q,p}(\mathbb{C})} \det(I + Z^*Z)^{-n}\, dZ.$$

Take an integer $k > n$, and the scalar weighted Bergman representation on the ball $I - Z^*Z > 0$, with squared norm proportional to

$$\int |f(Z)|^2 \det(I - Z^*Z)^{k-n}\, dZ.$$

The usual holomorphic action on functions is $f \mapsto \det(A' + B'Z)^{-k} f(g^{-1} \cdot Z)$, where $(A', B'; C', D')$ are the blocks of g^{-1}. Preservation of this norm follows from the two transformation formulas just given. The constant functions form the K_∞-type $(\det)^k \boxtimes 1$. Rotation $Z \mapsto e^{it}Z$ in the second block of K_∞ projects a holomorphic function onto its constant term by averaging: expand its power series in the scalar variable e^{it}, or apply Cauchy's integral formula on each complex line. Every nonzero invariant closed subspace contains the constants,

since one may move a point where a nonzero function is nonzero to the origin and then average over these rotations. Applying the same argument to an invariant orthogonal complement proves irreducibility. With the constant function normalized to norm one, projection onto that function is evaluation at zero. Its coefficient at g is therefore $\overline{\det A}^{-k}$ (up to the convention on which argument of the inner product is linear). In particular its squared absolute value is $|\det A|^{-2k} = \det(I - Z^*Z)^k$ for $Z = g \cdot 0$.

We evaluate the two matrix integrals explicitly. For real $s > p + q - 1$ and $t > -1$ put

$$I_+(s) = \int_{M_{q,p}(\mathbb{C})} \det(I + Z^*Z)^{-s}\, dZ, \qquad I_-(t) = \int_{Z^*Z<I_p} \det(I - Z^*Z)^t\, dZ.$$

Writing the last column of Z as z and the rest as Z_0, let $B = I \pm Z_0 Z_0^*$. The determinant identity

$$\det(I \pm ZZ^*) = \det B\,(1 \pm z^*B^{-1}z)$$

follows by factoring out B and taking the determinant of a rank-one perturbation of the identity. The domains correspond to all z in the plus case and to $z^*B^{-1}z < 1$ in the minus case. Substitute $z = B^{1/2}u$, whose real Jacobian is $\det B$. In the plus case the remaining exponent is $-(s-1)$ and the u integral is $\pi^q\Gamma(s-q)/\Gamma(s)$. In the minus case the remaining exponent is $t+1$ and the u integral is $\pi^q\Gamma(t+1)/\Gamma(t+q+1)$. These scalar integrals follow by radial integration and the beta integral, with the sphere area computed above. Iterating column by column gives

$$I_+(s) = \pi^{pq} \prod_{j=0}^{p-1} \frac{\Gamma(s-q-j)}{\Gamma(s-j)}, \tag{18.2}$$

$$I_-(t) = \pi^{pq} \prod_{j=0}^{p-1} \frac{\Gamma(t+1+j)}{\Gamma(t+q+1+j)}. \tag{18.3}$$

The chosen ranges guarantee absolute convergence at every stage. We used $\det(I \pm Z^*Z) = \det(I \pm ZZ^*)$, which follows by applying the determinant formula to the two block decompositions of a matrix with identity diagonal blocks.

In the Bergman realization the squared coefficient integral is finite and equals $I_-(k-n)/I_+(n)$, by disintegration along G_J/K_∞. Thus the representation is square-integrable and its degree is

$$\frac{I_+(n)}{I_-(k-n)} = \prod_{i=1}^{p} \frac{(p-i)!}{(n-i)!}\frac{(k-i)!}{(k-q-i)!}.$$

For this holomorphic discrete series the parameter with positive noncompact roots $e_i - e_{p+j}$ is, by the displayed lowest-type relation,

$$\lambda_i = k + \frac{p-q+1}{2} - i \quad (1 \le i \le p), \qquad \lambda_{p+j} = \frac{n+1}{2} - j \quad (1 \le j \le q).$$

Here ρ_c is half the sum of positive roots within each of the two blocks, and ρ_n is half the sum of the cross-block roots. Since $k > n$, these λ_i are indeed strictly decreasing in the indicated order. The differences within the two blocks are exactly those of ρ. The cross-block differences are $k - q - i + j$, and their ratios to the corresponding differences of ρ multiply to

$$\prod_{i=1}^{p}\prod_{j=1}^{q} \frac{k-q-i+j}{p+j-i} = \prod_{i=1}^{p} \frac{(k-i)!}{(k-q-i)!}\frac{(p-i)!}{(n-i)!}.$$

This is exactly the degree just computed. Harish–Chandra's parameter-independent proportionality now determines C_J for all regular discrete series of G_J: it is $(\prod_{i<j}(j-i))^{-1}$. If $p = 0$ or $q = 0$ the conclusion follows instead directly from compact Schur orthogonality and (18.1).

Proposition 18.1 (Exact formal-degree normalization)**.** *For a regular discrete series of $U(p,q)$ with doubled parameter l, with respect to the compact-dual measure used in the source, one has*

$$d_l = \dim V_\mu = \mathrm{WeylDim}(l),$$

where μ and the right hand side are given by (18.1)*. Thus the formal degree is independent of the signature for the corresponding compact infinitesimal character.*

For this use of the classical formal-degree theorem, it remains essential that the actual source coefficient `HCParam.dsCoeff` at the regular parameter be the normalized coefficient of the indicated discrete series (the Flensted–Jensen identification in `Lib/DiscreteSeries/FlenstedJensen/`). The preceding calculation fixes the constant in the classical degree formula; it does not prove that an unrelated function belongs to this representation. The degree equality is the first part of `Orbital.harish_chandra_degree_and_orbital` in `Lib/DiscreteSeries/Character/FormalDegree.lean`.

18.3 The exact elliptic orbital character identity

For a normalized unit vector v in the discrete series, set

$$e_l(g) = d_l\,\overline{\langle v, \pi_l(g)v\rangle}.$$

The source makes this choice with its highest coefficient; in particular it is a *minimal* convolution idempotent, not the character distribution of the representation. Indeed the coefficient orthogonality with degree d_l gives $e_l * e_l = e_l$ and $e_l(g^{-1}) = \overline{e_l(g)}$. These two identities also explain why taking the trace of a projection onto this type counts one coordinate per occurrence of the representation, rather than its full K_∞-type dimension.

We invoke the following precise classical orbit-character statement on the regular elliptic compact Cartan. For $z = (z_1, \dots, z_n)$ with pairwise distinct complex numbers of modulus one, write t_z for the corresponding diagonal element. Use on the centralizer torus $U(1)^n$ its probability Haar measure. Let

$$O_l(z) = \int_{U(1)^n \backslash G_J} e_l(g^{-1} t_z g)\, dg.$$

For the very regular parameters used in the source this integral exists. The stable sum is

$$(-1)^{pq} \sum_{\sigma \in S_n} O_l(z \circ \sigma) = p!\, q!\, \overline{\chi_\mu(t_z)}. \tag{18.4}$$

Here χ_μ is the compact Weyl character and the Haar measure on G_J is the compact-dual measure. This is the second assertion of `Orbital.harish_chandra_degree_and_orbital`; the relevant classical result is Harish–Chandra's discrete-series character formula on the compact Cartan (Acta Math. 116 (1966), Theorem 16, with the orbital-integral character formula of part I). We retain (18.4) in this precise normalization as the elliptic orbit-character input. Replacing the integral by a character value at each individual permutation would be false. Its *sum* has the asserted compact character and the sign $(-1)^{pq}$.

The integrand has been stated in the actual source convention. Haar integration over the whole noncompact group against a centralizer-invariant function would in general not be suitable; at regular elliptic elements the centralizer is compact of volume one, so the integral may equivalently be written on the whole group, as in the source. The stable sum divides by $p!\,q!$, because within each of the two signature subspaces the roots may be permuted by the compact subgroup. That division will be exactly the number of orderings of each signed orbit in the trace calculation below.

18.4 Absolute convergence and a direct ellipticity argument

The trace comparison in the formal source has explicit decay hypotheses. This is important for using ordinary integral kernels instead of an abstract pseudo-coefficient and a truncated trace formula. Write $F = E^+$, $f = [F : \mathbb{Q}]$, and

$$R(g) = \prod_{\tau \in \Phi} \|g_\tau\|_{\mathrm{op}} \quad (g \in G_\infty).$$

Then $R(g) \geq 1$, $R(g^{-1}) = R(g)$ and $R(gg') \leq R(g)R(g')$ by the identity $g_\tau^{-1} = J_\tau g_\tau^* J_\tau$. Let $\mathcal{O} \subset G_f$ be compact, with the condition that every $h \in \mathcal{O}$ has irreducible characteristic polynomial at some finite place of E. We consider the real product kernel $e = \prod_\tau e_{l_\tau}$. The precise `IsCuspKernel` hypothesis of order

$$B = 16[E : \mathbb{Q}](n+1)^3$$

provides $e * e = e$, a bound $|e(g)|R(g)^B \leq C$, integrability of $|e(g)|R(g)^{B+1}$, and a compact bump reproducing e. The specialized proof of this final kernel construction for the actual large parameters is part of the real discrete-series analysis; here we give the deductions from these properties.

Lemma 18.2 (Cyclic vector and power-packing bound)**.** *Let V be a framed rational Hermitian space of rank n, with fixed finite standard lattice and real norm as above. For every adelic point x and $T \geq 1$, the number of rational unitary elements γ such that*

$$(x^{-1}\gamma x)_f \in \mathcal{O}, \qquad R((x^{-1}\gamma x)_\infty) \leq T$$

is at most $C_{\mathcal{O},E,n} T^{[E:\mathbb{Q}](n+1)^3}$, independently of x.

Proof. Compactness of $\mathcal{O}$ in the finite adeles supplies a single positive rational integer c with ch carrying the standard lattice into itself for all $h \in \mathcal{O}$. Such an integer is obtained by bounding the finitely many denominators of the coordinates, since a compact set in the restricted product lies in a finite union of restrictions with all remaining coordinates integral. Use the adelic frame of V together with x to pull this standard finite lattice back to a lattice L_x in E^n, and pull the real Euclidean product norm back at infinity. Under that real norm the discrete full lattice in $\mathbb{R}^{[E:\mathbb{Q}]n}$ has a positive first minimum δ_x; choose a nonzero vector u of norm at most $2\delta_x$.

For each indicated γ, the vectors $c^j\gamma^j u$ $(0 \leq j \leq n)$ belong to L_x and have norm at most $2\delta_x(cT)^n$. The characteristic polynomial of γ over E is irreducible: a factorization of a monic polynomial over the field would remain proper after extension to every local field, contradicting the hypothesis on $\mathcal{O}$. Consequently $u, \gamma u, \dots, \gamma^{n-1}u$ form an E-basis. Indeed the annihilator of the nonzero cyclic vector in $E[X]$ is generated by a divisor of the irreducible characteristic polynomial (use Cayley–Hamilton), so must equal it. If two such γ, γ' give the same tuple $(c^j\gamma^j u)_{j=0}^n$, they act identically on that basis and are equal.

The number of lattice points in a Euclidean ball of radius $2\delta_x(cT)^n$ is at most $(1 + 4(cT)^n)^{[E:\mathbb{Q}]n}$: place disjoint open balls of radius $\delta_x/2$ around the points and compare total Euclidean volume to the enlarged ball. There are at most the $(n+1)$-st power of this number for the tuples. The resulting exponent $[E : \mathbb{Q}]n^2(n+1)$ is at most the exponent in the assertion. All constants are independent of the first minimum and hence of x. □

Summation in dyadic ranges of T now shows that $\sum_\gamma |k((x^{-1}\gamma x)_\infty)|\, \mathbf{1}_{\mathcal{O}}((x^{-1}\gamma x)_f)$ is uniformly bounded in x for any function k with $|k(g)|R(g)^B$ bounded. The finite-volume

assertion for a rational unitary quotient is the standard arithmetic-group covolume theorem (Borel–Harish-Chandra; the center here is anisotropic). Thus the sum is absolutely integrable over the rational quotient. Grouping by rational conjugacy classes, and replacing a sum over the translates of a class by the integral over its rational centralizer quotient, is justified by absolute convergence. This proves the relevant convergence assertion `Space.orbitalIntegral_unfold` with its explicitly larger exponent B. In particular it applies not just to e but also to the real majorant R^{-B}.

We next prove the special Selberg principle needed here from convolution idempotence and that convergence. Let γ have irreducible characteristic polynomial and an eigenvalue λ at an archimedean place with $|\lambda| \neq 1$. Fix a nonzero left eigenvector (a row) Q there. Every element a of its rational centralizer acts on that row by a scalar $\chi(a)$: the relevant eigenline in the dual space is one dimensional since the characteristic polynomial is separable. Put $H(x) = \log \|Qx_\tau\|$ for an adelic point expressed in the chosen real frame. Then $H(ax) = H(x) + \log|\chi(a)|$, so differences $H(xy) - H(x)$ for $y \in G_\infty$ descend to the centralizer quotient. Moreover

$$|H(xy) - H(x)| \leq \log R(y) \leq R(y), \qquad H(xd) - H(x) = \log|\lambda|,$$

where $d = (x^{-1}\gamma x)_\infty$ acts only in the real factor (the last equality uses only the real coordinates). The first inequality follows by the operator norm bound, applied in both directions using $R(y^{-1}) = R(y)$.

Let I be the orbital integral of $e(d)\mathbf{1}_{\mathcal{O}}((x^{-1}\gamma x)_f)$ on the centralizer quotient. In the integral write $e(d) = (e * e)(d)$ and insert the identity

$$\log|\lambda| = (H(xy) - H(x)) + (H(xd) - H(xy)).$$

The integral associated to the first term is

$$J_2 = \int_x \int_y e(y)e(y^{-1}d)(H(xy) - H(x))\, \mathbf{1}_{\mathcal{O}}((x^{-1}\gamma x)_f)\, dy\, dx.$$

It is absolutely convergent: by submultiplicativity the integrand in absolute value is bounded by

$$CR(d)^{-B}|e(y)|R(y)^{B+1}\mathbf{1}_{\mathcal{O}}((x^{-1}\gamma x)_f),$$

which is integrable by the packing bound and the moment hypothesis. The second term has the same integral by the change of variables $(x, y) \mapsto (xy, y^{-1}d)$. This transformation preserves product measure: first translate x on the right by y, then invert and translate y by a fixed group element depending on the new x. All relevant groups are unimodular and the rational centralizer quotient has invariant measure. The finite indicator is unchanged. Thus

$$\log|\lambda|\, I = 2J_2.$$

Alternatively use the triple convolution $e * e * e = e$ and the three successive increments along x, xy, xyz, xd. The first increment has integral J_2 after integrating the convolution in z; its absolute convergence follows from the bound

$$CR(d)^{-B}|e(y)|R(y)^{B+1}|e(z)|R(z)^{B}\mathbf{1}_{\mathcal{O}}((x^{-1}\gamma x)_f).$$

The cyclic change of variables $(x, y, z) \mapsto (xy, z, z^{-1}y^{-1}d)$ preserves measure by the same argument, and permutes the increments. It therefore proves convergence of the other two and equality of their integrals with J_2. Hence $\log|\lambda|\, I = 3J_2$. Subtracting the two identities gives $J_2 = 0$ and then $I = 0$. This proves precisely the required assertion: a nonzero orbital integral on this irreducible finite support can only come from a rational element whose complex eigenvalues all have modulus one. It is `Space.elliptic_of_integral_ne_zero` in `Lib/LocalGL/TraceFormula/TraceComparison.lean`. No noncompact orbital integral is silently equated to a compact orbital character.

18.5 Changing the signatures of the rational elliptic classes

The source changes real signatures while keeping the finite local groups and the parameter fixed. The elementary global algebra behind this operation uses the norm theorem, not strong approximation. Here are the details.

Let $\gamma \in \mathrm{GL}_n(E)$ have irreducible characteristic polynomial and suppose it preserves a rational Hermitian form J and has only eigenvalues on the unit circle at infinity. Its rational centralizer is the multiplicative group of the field $B_\gamma = E[\gamma]$: the cyclic-vector argument identifies a commuting linear map with multiplication by its value on a cyclic vector. The adjoint involution with respect to J on this centralizer induces an involution $\dagger$ on B_γ extending conjugation of E and satisfying $\gamma^\dagger = \gamma^{-1}$; this does not depend on the particular preserved form because these prescriptions determine it. At each embedding of B_γ into $\mathbb{C}$ extending the chosen CM type, the image of γ^{-1} is the complex conjugate of the image of γ. It follows on generators that $\dagger$ is complex conjugation on every embedding. Thus B_γ is a CM field, with totally real fixed subfield B_γ^+. The rational unitary centralizer is precisely $\{a \in B_\gamma^\times : a^\dagger a = 1\}$, independent of the preserved form.

A frame class may be described by a pair (x_f, γ) consisting of a finite adelic invertible matrix and the rational matrix, modulo simultaneous rational conjugacy (left transformation of the frame and conjugation of γ) and right multiplication of x_f by the fixed standard finite unitary group. The pair is admissible if some rational J preserved by γ satisfies $x_f^* J x_f = I$. This condition determines J uniquely when it exists. For fixed γ, after choosing numbered roots at every complex place, a frame class has a sign on every eigenline: eigenlines with distinct eigenvalues are mutually orthogonal, and each is nonisotropic because the Hermitian form is nondegenerate. Denote the total sign assignment by $\epsilon_{\tau,i}$.

Lemma 18.3 (Norm change of a signed elliptic frame)**.** *Any prescribed changes $\delta_{\tau,i} = \pm 1$ of those signs whose global product is $+1$ can be realized by a bijection of these frame classes. It commutes with their centralizers, preserves their finite positions $x_f^{-1}\gamma x_f$, and preserves the finite centralizer embeddings into the standard finite group.*

Proof. The sign changes are indexed by the real embeddings of B_γ^+. The global quadratic norm theorem for B_γ/B_γ^+ gives $a \in (B_\gamma^+)^\times$ with exactly these signs and a finite idele z of B_γ with $az^\dagger z = 1$ in the finite adeles. Indeed the infinite idele of the prescribed signs with finite part 1 has quadratic Artin symbol equal to their product, so lies in the principal-times-norm subgroup. The real norms are positive, yielding the stated sign of the principal term. This argument is the same as in Corollary 6.5, and does not require B_γ/B_γ^+ to be unramified.

View a and z as multiplication operators in the centralizer algebra and put

$$J' = Ja, \qquad x'_f = z x_f.$$

The first is Hermitian and preserved by γ, since a is adjoint-fixed and commutes with γ. At the finite places,

$$(x'_f)^* J' x'_f = x_f^* J z^\dagger a z x_f = x_f^* J x_f = I.$$

At infinity a acts by the scalar of sign $\delta_{\tau,i}$ on each eigenline. Hence the resulting form has exactly the required modified signs. The rational matrix γ is unchanged, the finite position $(x'_f)^{-1}\gamma x'_f$ equals the original one, and every element of its centralizer commutes with z, so its finite position also remains unchanged.

For the assertion about classes, fix the characteristic polynomial and identify all its cyclic rational matrices with one companion matrix by rational conjugation. The multipliers a, z can then be chosen once for each fixed sign assignment. The conjugations identifying a rational

matrix with the companion differ by an element of its centralizer, which commutes with both multipliers. Thus the construction is well-defined on frame classes. Its inverse uses a^{-1} and z^{-1}, proving bijectivity. □

This is the mechanism of `exists_hasSignType_mul_iff` in `Lib/LocalGL/TraceFormula/ChangingSigns.lean`. In the comparison of signatures (p, q) constant over Φ, every choice of q negative eigenlines at each real place differs from the positive assignment by a sign vector of product $(-1)^{qf} = +1$, since f is even. Lemma 18.3 therefore gives all the signed classes from the positive ones with their finite positions and centralizers preserved. Conversely every contributing signed class is obtained uniquely from its positive counterpart for each specified assignment. Equivalence of rational Hermitian spaces with the same local signatures and frames follows from Landherr's theorem, as explained in Chapter 6. This identifies those classes in any of the fixed global spaces used for the trace computation. In particular it is not necessary to assume that all indefinite unitary spaces with arbitrary local invariants have the same automorphic spectra.

18.6 The trace of the elliptic finite test functions

Let $\mathcal{V}_s(K_f)$ be the principal-level row-type space on a framed space of signature s, with all real parameters fixed. Here K_f denotes the principal level at the chosen places (a compact open subgroup of the standard finite maximal group). For $h \in G_f$, consider the compression of right translation by h to this fixed-level space; it is represented by the finite test function $1_{hK_f}/\operatorname{vol}(K_f)$. If h normalizes K_f this is the ordinary translation on the fixed-level space; more generally one may compose with the orthogonal projection to the level space. We use the source's reproducing-kernel property of the actual cusp kernel: the real idempotent and the finite-level averaging operator have a finite-rank automorphic reproducing kernel whose range is precisely $\mathcal{V}_s(K_f)$ (see `Lib/Adelic/CuspForms/ReproducingKernel.lean`, `Space.exists_kernel_form`). This is an identification of the kernel with the actual rapid cusp forms, not merely an application of the formal-degree polynomial.

The ordinary kernel trace, on multiplying by $\operatorname{vol}(K_f)$, is

$$\int_{[G_s]} \sum_{\gamma \in G_s(F)} e((x^{-1}\gamma x)_\infty) \mathbf{1}_{hK_f}((x^{-1}\gamma x)_f)\, dx.$$

For example it follows by expressing a reproducing kernel in any orthonormal basis of its finite-dimensional range, composing with translation, and integrating its diagonal. The equality with the rational sum is the periodized kernel identity; the normalization by $\operatorname{vol}(K_f)$ comes from averaging over the level. Off the diagonal the periodizing sum converges on compact sets: compact finite support bounds the rational matrix entries in a fixed denominator lattice on any such set, and the archimedean norm bound reduces counting to a Euclidean lattice point estimate of dimension $[E : \mathbb{Q}]n^2$. The exponent B above is more than sufficient for this estimate. For the diagonal and the integral above, the stronger uniform estimate of Lemma 18.2 applies under the irreducibility hypothesis on the coset. Thus regrouping by rational conjugacy classes and applying the ellipticity argument of the preceding subsection is legitimate.

Here is the remaining comparison for the elliptic part. Fix a positive definite frame class with its rational γ and the finite frame. Its centralizer at each real place is $U(1)^n$, and the finite centralizer embedding and finite position are the same for all the sign assignments obtained from Lemma 18.3. The nonarchimedean factors of the orbital integral are therefore unchanged. More formally, take product Haar measures on the centralizer adelic group, choosing at the

real places its probability torus measures; its rational subgroup and the quotient by it are independent of the signed frame by the centralizer description above. Disintegrating over the adelic centralizer and its quotient factors every geometric integral as a common centralizer volume times its finite orbital integral times a product of real orbital integrals. This is just invariant quotient integration: on changing the real eigenbasis, the conjugation identifies the centralizers with the same probability compact torus. Thus none of these factors introduce signature-dependent real constants. The disintegration is justified by the absolute convergence proved above.

At each real place, the sum over the choices of q negative eigenlines is the sum over S_n divided by $p!\,q!$: permutations within the positive and negative blocks are represented by the compact group and give the same orbital integral. Formula (18.4) makes this sum equal to $(-1)^{pq}\overline{\chi_\mu(t_z)}$, the compact orbital integral with the indicated sign. Across the f real places the total factor is $(-1)^{pqf} = 1$. Hence summing the orbital contributions in the signed space yields the very same number as summing them in the positive space. This argument proves signature independence of the trace for every such finite test coset under the stated cusp kernel and elliptic character input:

$$\operatorname{tr}_{\mathcal{V}_s(K_f)} T_h = \operatorname{tr}_{\mathcal{V}_{s'}(K_f)} T_h. \tag{18.5}$$

The proof follows the mechanism of `Space.exists_trace_eq` in `Lib/LocalGL/TraceFormula/TraceComparison.lean` (compare Labesse, *Changement de base CM et séries discrètes*, Thm. 5.4). The two genuine representation-theoretic inputs here are the real orbital character identity (18.4) and the identification and decay of the actual reproducing cusp kernel. We have given independently the algebraic movement of the framed orbits, the convergence mechanism, and the idempotent proof of elliptic support. The trace identity applies to the exact finite irreducible-coset support, not to all finite test functions indiscriminately.

18.7 Primitive labels and the finite cubic correction

The passage from (18.5) to equality of traces on the good subspaces is a separate finite argument. We detail it, including why it does not amount to assuming equality on the whole principal-level space.

Recall the regular labels c for $\mathrm{GL}_r(k)$: monic squarefree degree r polynomials with nonzero constant term. The Green character χ_c and its central idempotent are written

$$e_c = \frac{\chi_c(1)}{|\,\mathrm{GL}_r(k)|} \sum_h \overline{\chi_c(h)}\, h.$$

If c is not a label the source puts $e_c = 0$. When c is irreducible of degree r, χ_c is the cuspidal Green character of the corresponding regular finite-extension character, and c is primitive when its roots generate $\mathbb{F}_{q^r}^\times$, $q = |k|$. We use the classical finite-general-linear-group character facts: these regular labels give pairwise orthogonal central projectors; for an irreducible label its cuspidal character vanishes unless the matrix characteristic polynomial is a power of an irreducible polynomial; twisting that character by a determinant character amounts to scaling the roots of its label by the corresponding base-field character name. These are Green's character table facts (J. A. Green, *The characters of the finite general linear groups*, 1955), used in `Lib/LocalGL/Green/FrobeniusOrbits.lean` and `New/Trace/BadShare.lean`. The important elementary consequence is the following.

Lemma 18.4 (Cubic avoidance with unchanged nonelliptic coefficients)**.** *Suppose $r = 3^a$ for $a \geq 1$, $9 \mid q-1$, and S, T are two sets of primitive labels, each containing at most one*

element. Given any label c there is a label c' such that the coefficients of e_c and $e_{c'}$ agree at every h whose characteristic polynomial is reducible, and if c is primitive then c' is primitive and outside $S \cup T$.

Proof. If c is not primitive, take $c' = c$. Otherwise let β be a primitive root of c in $\mathbb{F}_{q^r}$, and put $D = q^r - 1$. The three elements $a\beta$ for $a^3 = 1$ in $\mathbb{F}_q^\times$ are all generators of $\mathbb{F}_{q^r}^\times$: express a as $\beta^{jD/3}$, so $a\beta$ has exponent $1 + jD/3$ relative to the generator, which is coprime to D because $9 \mid D$. Indeed every prime divisor of D divides $D/3$. In particular each has degree r over k. Their Frobenius orbits are distinct. If two were equal, β^{q^i-1} would be a nontrivial cube root of unity for some $0 \le i < r$. But $q^i - 1$ is either zero or lies strictly between zero and $D/3$ (use $q > 3$); it cannot be congruent to $D/3$ or $2D/3$ modulo D. Thus these give three distinct primitive labels. They are the cubic determinant twists of the original label, by the classical compatibility just stated. One of them is not in the at most two excluded labels.

At any h of reducible characteristic polynomial where the original cuspidal character is nonzero, that polynomial is P^b for an irreducible P and $b > 1$ dividing r. Therefore $3 \mid b$. Its determinant $(-1)^r P(0)^b$ is a cube, the sign being a cube as well. The cubic determinant character is thus one at h. At all other reducible characteristic polynomials the original character and its twist both vanish. The dimension is unchanged by twisting, so the coefficients of their central idempotents agree at every such h. □

This proves the content of `exists_labelIdem_eq_off_irreducible` used in the source apart from the standard Green character formulas. Notice the distinction between the *label* polynomial of a representation and the characteristic polynomial of a chosen matrix h. The equality in Lemma 18.4 concerns the latter being reducible, not the former.

At a fixed Hecke eigensystem α write $S_w(\alpha, V)$ for the primitive labels occurring at a chosen place w on the principal-level space of the framed group V. This means the corresponding e_c is nonzero on the α eigenspace. The source proves that this set has at most one element for the actual row and single-lift source spaces, not for arbitrary unitary groups. We record why the unusual primitive condition allows weak Galois compatibility rather than full equality of local Langlands parameters.

Lemma 18.5 (A finite rigidity implication)**.** *Let c, c' be primitive labels of degree r over $\mathbb{F}_q$, and suppose a common attached Galois representation has an inertia element whose operator is annihilated by both polynomials obtained from c, c' by adjoining further linear factors. Assume all the further roots have order dividing M, with $q - 1 > M$, under the same faithful multiplicative embedding of the residue algebraic closure into the coefficient field. If the determinant of the operator is the product of the roots of the first enlarged polynomial, then $c = c'$.*

Proof. Two distinct irreducible polynomials of the same degree have disjoint sets of roots. If $c \neq c'$, every eigenvalue of the common operator, since it is killed by both polynomials, must therefore be among roots of order dividing M. Consequently its determinant has order dividing M. On the other hand the product of the roots of c is the norm of a generator of $\mathbb{F}_{q^r}^\times$, and has exact order $q - 1$ in $\mathbb{F}_q^\times$: it is $\beta^{(q^r-1)/(q-1)}$. Multiplying by roots of order dividing M cannot change it into an element of order dividing M, since that subgroup is closed under division. This contradicts the asserted determinant identity. □

In the actual application the primitive roots may first be scaled by a scalar of order dividing $M = M_\omega$. They stay primitive when $M^2 \mid q - 1$: relative to a generator of $\mathbb{F}_{q^r}^\times$ the exponent of such a scalar is divisible by $(q^r - 1)/M$, which in turn is divisible by every prime divisor of $q^r - 1$. Thus the exponent of the scaled generator is coprime to $q^r - 1$, as in the proof

for cubic scaling. Scaling by this same scalar is injective on labels. The further roots of the attached polynomial arise from determinant characters of additional blocks and have order dividing M. The local-global statement used by the source is `Attached.isLabelledAt` in `Lib/GaloisRep/LocalGlobal/LabelCompatibility.lean`; it gives a killing polynomial and the determinant identity at a suitable common tame exponent away from the coefficient prime, not necessarily equality of characteristic polynomials. The tame exponent one is used in `Data.occurringLabels_subsingleton` of `New/GaloisRep/LabelUniqueness.lean`. Weak attachment and the equality of the semisimple attached Galois representation for the two eigenforms follow there from the spherical Hecke system and its density compatibility. The actual injective auxiliary theta lifts and the parahoric intertwining are needed so that this local statement applies to the primitive label on the original space. These geometric and Galois-theoretic inputs are retained at their stated strength; Lemma 18.5 proves the finite implication once they have been established. In particular the condition `HasRigidLabels` for `spaceY0` and `spaceWj` is not a formal consequence of the existence of an abstract Hecke system.

18.8 Equality of all good Hecke traces and determinants

Let $\mathcal{G} = \prod_w \mathrm{GL}_r(k_w)$ be the chosen finite-level group. The good idempotent of the data is

$$e_0 = \sum_{\mathbf{c}\ \text{good}} \bigotimes_w e_{c_w}.$$

A good family consists of allowed regular labels at every chosen place and at least one primitive label above each of the two chosen rational primes. It is an orthogonal central projector. Let V, V' be two of the spaces of principal-level forms of the proof, with the same archimedean parameter and (possibly) different signatures. Denote their actions of the full away-from-chosen-places Hecke algebra $\mathcal{H}$ by $P(a), P'(a)$. In the source this algebra may be larger than the spherical commutative algebra: it consists of families of endomorphisms induced, across the groups, by finite group-algebra combinations of translations trivial at the chosen places (`Data.heckeAlgebra`, `New/Selmer/FitTwists.lean`). It commutes with the chosen finite group action. Its spherical operators at the split places away from the level commute and act normally, so their joint eigenspaces give a direct sum on the finite-dimensional spaces. The following argument does not require every element of the larger algebra to act diagonally.

Let $\mathcal{F}$ be the union of the finitely many occurring joint eigensystems on V, V'. For each $\alpha \in \mathcal{F}$ choose, for each distinct $\beta \in \mathcal{F}$, a spherical Hecke operator $T_{x(\beta)}$ on which they have distinct eigenvalues. The product

$$p_\alpha = \prod_{\beta \neq \alpha} \frac{T_{x(\beta)} - \beta(x(\beta))}{\alpha(x(\beta)) - \beta(x(\beta))} \quad \text{in } \mathcal{H}$$

acts as identity on the α-part and zero on the other parts in both spaces. This explicit interpolation proves the finite simultaneous projection `Data.exists_hecke_proj_eigenPart`. It is harmless that one eigensystem might occur on just one of the spaces.

Assume the rigidity property $|S_w(\alpha, V)|, |S_w(\alpha, V')| \leq 1$. At each w and for each local label c use Lemma 18.4 with these two sets to choose $c'(\alpha, w, c)$. Form the correction

$$E_\alpha = e_0 - \sum_{\mathbf{c}\ \text{good}} \bigotimes_w e_{c'(\alpha, w, c_w)}.$$

For each good tuple, a primitive factor is replaced by one not occurring on the α-parts of *either* space; the corresponding tensor operator kills both parts. Hence on their α eigenspaces E_α

and e_0 have the same action. On the other hand every coefficient of E_α at a tuple of matrices having reducible characteristic polynomials at all chosen places vanishes by Lemma 18.4, applied factor by factor to the corresponding summands. Thus its support consists of matrices having an irreducible characteristic polynomial at at least one chosen place. These are two completely different properties: the first concerns which representations the replacement labels act on, while the second concerns the geometric support of the group-algebra element.

For any $a \in \mathcal{H}$, cyclicity of trace and the fact that a commutes with the chosen-level action yield

$$\mathrm{tr}_{e_0 V}(P(a)) = \mathrm{tr}_V(P(a)e_0) = \sum_{\alpha \in \mathcal{F}} \mathrm{tr}_V(P(a p_\alpha)E_\alpha),$$

and likewise for V'. In the first equality the trace on the full space is that of the action on the image of the projection, with zero on its kernel. For the second one, sum the projections p_α to the identity and then use the equality of E_α and e_0 on that part; both commute with p_α. No commutation of a with all p_α is needed for this formula.

Each operator on the last sum is a finite linear combination of translations by a product of an away-from-chosen-places adelic element and a lift of a tuple in the support of E_α. The full right coset by the principal level has irreducible characteristic polynomial at one of the chosen places where the tuple does. Indeed its reductions there have the same monic irreducible characteristic polynomial, so the lifted characteristic polynomial is irreducible over the local field by reduction of monic integral factors (Gauss's lemma over the discrete valuation ring). The away operator is trivial there and cannot affect this fact. Thus every summand may be treated by the elliptic trace comparison (18.5). We obtain

$$\mathrm{tr}_{e_0 V}(P(a)) = \mathrm{tr}_{e_0 V'}(P'(a)) \qquad (a \in \mathcal{H}). \tag{18.6}$$

This is the argument of `Data.trace_heckeOnGood_eq` in `New/Trace/GoodFormTransfer.lean`. It proves the finite projection step exactly, assuming the explicit hypotheses of the geometric comparison and the actual rigidity assertion, rather than making a stronger unsupported transfer claim.

Taking $a = 1$ proves equality of dimensions. Taking all powers of a in (18.6) proves equality of the characteristic polynomials (by Newton's identities over $\mathbb{C}$) and hence

$$\det_{e_0 V}(P(a)) = \det_{e_0 V'}(P'(a)). \tag{18.7}$$

In particular if a joint spherical eigensystem is absent from one good space it is absent from the other: interpolate the projector onto its part on the union of the finite eigensystems and take its trace, which is the dimension of that part. These implications reproduce `det_heckeOnGood_eq` and `goodForms_inf_eigenPart_eq_bot`. The determinant equality, not a mere numerical dimension comparison, is also used to compare the Rallis operators of the row chain and the auxiliary single lifts in the lower bound.

18.9 The precise row-dimension inequality

We finally apply these comparisons to `Data.finrank_rows_le_mass`. For the row group the doubled parameter at $\tau \in \Phi$ is

$$l_\tau(i) = 2\,\varsigma_\tau(\mathrm{basisNumber}(i)),$$

where ς_τ is the row parameter in the data, a strictly decreasing sequence. Thus the decreasing position of i in l_τ is exactly basisNumber(i) (the number of entries larger than it), independent

of the ordering chosen for the basis indices. This is `decrPosition_paramW` in `New/Filling/RowDimension.lean`. If the corresponding dominant diagram is κ_τ with central shift e_τ, the Harish–Chandra parameter associated to that highest weight is

$$l_\tau^{\downarrow}(i) = 2(e_\tau + \kappa_\tau(i)) + r - 1 - 2i, \qquad 0 \le i < r.$$

Equivalently $\kappa_\tau(i) + e_\tau$ is $l_\tau^{\downarrow}(i)/2 - \rho_i$ with zero-based $\rho_i = (r-1-2i)/2$. This recovers `hcParameter` and `IsHCParam.lowestKType_eq` in `Lib/Adelic/CuspForms/WeightParameter.lean`. The compact tensor coefficient representation has dimension

$$d_\infty = \prod_{\tau \in \Phi} \dim V_{\mu_\tau} = \prod_{\tau} d_{l_\tau}.$$

The last equality is the exact compact-dual degree formula of Proposition 18.1. The local Bessel system in the source is required to use these same degrees at each real place (the parameter-matching component of `D.filling_fits`); thus d_∞ equals the number called `Filling.besselDegree`, and not a multiple of that number.

For the definite group with these compact real representations, forms of the minimal real type correspond to algebraic modular forms with this tensor coefficient representation. One can see the isomorphism directly: if v_μ is the unit highest vector in that irreducible representation and f is a vector-valued finite automorphic form, the scalar coefficient at (x_∞, x_f) is obtained by pairing $f(x_f)$ with the translate of v_μ by x_∞ in the appropriate contragredient convention. The rational automorphy follows from the transformation of f by the coefficient representation, and compact Schur orthogonality gives the reproducing identity. Conversely integrate the scalar form against these translates of the highest vector and multiply by d_∞ to recover the vector-valued form. The two maps are inverse by Schur orthogonality and cyclicity of the highest vector; the latter follows because its orbit linearly spans a nonzero invariant subspace of the irreducible representation. Finite-level invariance is preserved. This is the compact equivalence `Space.algFormEquiv` in `Lib/Adelic/CuspForms/AlgebraicForms.lean`.

At the chosen principal level in rank r all rational stabilizers are trivial by the neatness calculation; the index into the standard finite maximal group is $|\mathcal{G}|$. The mass of its finite class groupoid is $|\mathcal{G}|\mathfrak{m}_r$, by the level-index identity proved in Chapter 6, and since the stabilizers are trivial this is exactly the number of classes. An algebraic form may be assigned an arbitrary coefficient vector independently at each such class. Thus the dimension of the full definite principal-level space equals

$$|\mathcal{G}|\,\mathfrak{m}_r\,d_\infty.$$

Its good subspace can have smaller dimension. By (18.6) for $a = 1$, the dimension of the actual row good space W equals the dimension of that definite good subspace. Therefore

$$\dim_{\mathbb{C}} W \;\le\; |\mathcal{G}|\,\mathfrak{m}_r\ \mathrm{besselDegree}\,. \tag{18.8}$$

This is precisely the inequality stated in the source as `Data.finrank_rows_le_mass`. In the notation of the spanning chapter it is the bound by the mass times the local formal degrees times the number of level translates. No factor of dimension of the lowest K_∞-type appears: the chosen archimedean projection is the rank-one matrix coefficient, as explained above. Nor does one need equality of the full indefinite and definite principal-level spaces.

To summarize the remaining dependencies of this particular chapter: the elementary coordinate measure, matrix integral, compact degree calculation and finite projection arguments are independent conventional proofs. The formal-degree proportionality and discrete-series elliptic character identity are classical real representation-theoretic inputs; the precise match of the actual source kernel to them, with its uniform decay and reproducing property, must

be verified for the constructed row parameters. The weak Galois attachments, local label compatibility and actual auxiliary injective lifts supplying the rigidity hypothesis must also be established. They are used only at the stated interfaces here. Once those inputs are supplied, the signature comparison and exact dimension inequality follow by the arguments above without introducing an additional global dimension-transfer assumption.

Chapter 19

A trace-class derivation of the normalized elliptic comparison

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 25 and the final audit in Chapter 57.

In Chapter 18 the exact elliptic stable orbital identity was retained as a separate interface. Here we prove its normalization and its stable sum directly from ordinary discrete-series coefficient orthogonality, the character theorem and the determinant-decay bound proved for the explicit coefficient earlier. This isolates a narrower remaining obligation: identification of the source's explicit Flensted–Jensen coefficient with the normalized highest vector of the intended discrete-series representation, including the exact representation and differentiation properties. One may not invoke Harish-Chandra's character formula on an arbitrary kernel without that identification.

19.1 A uniform local orbital majorant

Let $G = U(p,q)$, $n = p+q$, with Haar measure as in Chapter 18. We use normalized probability measure on its compact Cartan $T = U(1)^n$. Suppose the coefficient $\psi(g)$ satisfies, for a fixed exponent γ,

$$|\psi(g)| \le C\|g\|_{\mathrm{op}}^{-\gamma}. \tag{19.1}$$

For the source coefficient the large exponent is supplied by Lemma 22.5, from the noncompact gaps. The exact numerical thresholds in the source are more than enough for the following estimate whenever $\gamma > n^3$.

Lemma 19.1. *Let z range over a compact subset of the regular torus (the eigenvalues z_i are distinct). If $n \ge 2$ and $\gamma > n^3$, then*

$$\int_G |\psi(g^{-1}t_z g)|\,dg$$

is uniformly bounded. The orbital integral of a continuous coefficient is continuous in z on this regular torus.

Proof. The cases $p = 0$ or $q = 0$ are compact and immediate. Otherwise let P be projection onto the positive signature coordinates and $S = 2P - I$. For each regular z there is a polynomial $A_z(X)$ of degree at most $n-1$ with $A_z(z_i) = 1$ for the positive coordinates and

zero elsewhere. Lagrange interpolation shows that its coefficients are uniformly bounded on the specified compact set: the denominators are products of nonzero eigenvalue differences, bounded away from zero. Then

$$g^{-1}Pg = A_z(g^{-1}t_z g), \qquad g^{-1}Sg = Sg^*g.$$

The second equality uses $g^{-1} = Sg^*S$. Therefore

$$\frac{\|g\|_{\mathrm{op}}^2 - 1}{2} \le \|g^{-1}Pg\|_{\mathrm{op}} \le C'(1 + \|g^{-1}t_z g\|_{\mathrm{op}})^{n-1}.$$

Combining with (19.1) gives a majorant $C''\|g\|_{\mathrm{op}}^{-2\gamma/(n-1)}$ for $\|g\| \ge 2$, uniformly in z. A unitary-for-S matrix and its inverse have the same operator norm, so the norm is at least one.

For completeness this majorant is integrable. The compact-dual matrix-ball chart of Chapter 18 has invariant quotient density proportional to $\det(I - Z^*Z)^{-n}$ on the bounded matrix ball; the compact group has measure one. If $\|g\|_{\mathrm{op}} \le R$, the singular values of the corresponding Z are at most $\sqrt{1 - R^{-2}}$: in a Cartan block $Z = \tanh(a)$ and $\|g\| \ge \cosh(a)$. Thus $I - Z^*Z \ge R^{-2}I$, and the density is at most R^{2n^2} times a fixed constant. The Haar volume of the operator norm ball is consequently at most CR^{2n^2}. Summing over dyadic annuli proves integrability of $\|g\|^{-u}$ for $u > 2n^2$. The assumed bound on γ implies $2\gamma/(n-1) > 2n^2$. Dominated convergence gives continuity. □

The same proof gives a locally uniform orbital majorant when the element t ranges over a compact subset of the regular elliptic locus of G, not merely the diagonal torus. Indeed every such t has a neighborhood in this locus in which it is conjugate to a varying regular compact Cartan element by a locally bounded conjugator. This follows from the derivative of the conjugation map: since the eigenvalue differences are nonzero, its derivative on the tangent space modulo the Cartan centralizer is invertible onto the noncentral directions, and the torus variables give the remaining directions. The inverse function theorem gives local sections. A finite subcover of a compact support yields bounded conjugators; change variables in the Haar integral. This justifies the following use of Fubini for test functions supported in the regular elliptic locus.

19.2 The degree factor in the orbital trace

Theorem 19.2 (Orbital integral of a normalized coefficient)**.** *Let π be an irreducible unitary discrete series of G, v a unit vector, and*

$$\psi(g) = \langle v, \pi(g)v\rangle, qquade(g) = d_\pi\overline{\psi(g)}$$

where d_π is its formal degree for the given Haar measure. Suppose (19.1) holds for some $\gamma > n^3$. Then at every regular elliptic compact Cartan element

$$\int_G e(g^{-1}t_z g)\,dg = \overline{\Theta_\pi(t_z)}. \tag{19.2}$$

Here Θ_π is the ordinary analytic character on the regular locus. The quotient integral over the centralizer has exactly this value when the centralizer is given probability Haar measure.

Proof. Schur orthogonality for the discrete series gives the resolution

$$d_\pi \int_G |\langle w, \pi(g)v\rangle|^2, dg = \|w\|^2$$

for all Hilbert vectors w. For any trace-class operator A this implies

$$\operatorname{Tr} A = d_\pi \int_G \langle \pi(g)v, A\pi(g)v\rangle \, dg. \tag{19.3}$$

To verify absolute convergence and the equality, write a trace-norm convergent sum $A = \sum_i s_i |u_i\rangle\langle w_i|$ with orthonormal u_i, w_i and $\sum_i s_i < \infty$. Cauchy–Schwarz and the resolution bound each integrated product in absolute value by s_i/d_π. Then interchange sum and integral, and use the polarized resolution for each term; its value equals the trace of that rank-one operator.

Take $A = \pi(f) = \int f(t)\pi(t)\,dt$, for $f \in C_c^\infty(G)$ supported on the regular elliptic locus. The standard character-distribution theorem applies: $\pi(f)$ is trace class and its trace is integration against Θ_π. For the trace-class premise see Wallach, [18, Theorem 95], which states the result for irreducible unitary representations of real reductive groups; the ordinary analytic character here follows from the classical discrete-series character theorem discussed in Chapter 23. Substituting A in (19.3) and applying Fubini (justified by Lemma 19.1 and its uniform version on compact elliptic support) gives

$$\int f(t)\Theta_\pi(t)\,dt = \int f(t) d_\pi \int_G \psi(g^{-1}tg)\,dg\,dt.$$

Both functions of t are continuous on the regular elliptic locus by the character theorem and the majorant. Equality for all such test functions gives equality pointwise there. Taking complex conjugates proves (19.2).

The integrand is left invariant under the compact centralizer in the variable g, so integrating over G equals integrating over the quotient by the centralizer with its probability Haar measure. No extra formal-degree or K-type dimension factor is needed. □

19.3 The stable Weyl sum

For a regular Harish-Chandra parameter λ in the source convention, choose the positive roots ordered by λ, write ρ for their half sum, and let $W = S_n$, $W_K = S_p \times S_q$. The compact highest weight is $\mu = \lambda - \rho$. The parity and regularity conditions of the source ensure this is integral and dominant in the chosen ordering. With $D(z) = \sum_{w\in W} \epsilon(w) e^{w\rho}(z)$ (or the equivalent product of roots), the discrete-series character formula on the regular compact Cartan is

$$\Theta_\lambda(z) = (-1)^{pq} \frac{\sum_{w\in W_K} \epsilon(w) e^{w\lambda}(z)}{D(z)}.$$

We use the classical Harish-Chandra character theorem, Part II, Acta Math. 116 (1966), Theorem 16, with its compact Cartan formula and the ordering fixed as above; the matching reference and notation are discussed in Chapter 23. Although half-integral coordinates can be used in numerator and denominator, their quotient is single valued by the stated parity. One may equivalently multiply both by a common half-determinant to work in integral Laurent monomials.

For clarity the summation is finite and purely algebraic. Replacing z by $z \circ \sigma$ multiplies the denominator by $\epsilon(\sigma)$ and sends the numerator monomials to their permuted monomials. Hence

$$\sum_{\sigma\in W} \Theta_\lambda(z\circ\sigma) = (-1)^{pq}|W_K| \, \frac{\sum_{w\in W} \epsilon(w) e^{w\lambda}(z)}{D(z)} = (-1)^{pq} p!q!\, \chi_\mu(z).$$

The last equality is the compact Weyl character formula. Applying Theorem 19.2 and complex conjugating gives exactly

$$(-1)^{pq} \sum_{\sigma \in S_n} \int_G e(g^{-1} t_{z\circ\sigma} g)\, dg = p!q!\, \overline{\chi_\mu(z)}.$$

This is the stable orbital assertion in `Orbital.harish_chandra_degree_and_orbital` of `Lib/DiscreteSeries/Character/FormalDegree.lean`, once the source explicit coefficient is identified with the indicated unit vector of the discrete series. The first formal-degree assertion uses the Haar calibration of Chapter 18. In particular the stable formula is not a conjectural transfer principle or an extra trace normalization. For each individual permutation the character is that of the corresponding discrete series, not the compact character; only the signed sum is the compact character.

Closure status. The exact factor in the normalized stable orbital comparison is proved from (i) the actual coefficient realization as a discrete-series matrix coefficient, (ii) the determinant decay already supplied, and (iii) the ordinary character and trace-class theorems. The missing input is (i), together with any additional analytic differentiation and rigidity requirements for the actual constructed representation in the source. No assertion of the full row transfer follows before those hypotheses and the finite-label conditions are checked. This result nevertheless removes the stable Weyl-sum and Haar quotient normalization as independent unresolved interfaces.

Chapter 20

Quantitative spanning by projected Bessel–Poincaré series

Chapters 21 and 22 supply the kernel, Schur, Laplace, finite-volume, Poisson, real-peak and sieve arguments underlying the estimates used here. Statements that also require uniform geometric input retain that input unless it is proved in these companion chapters; the elementary rank deduction is independent of the details of this analysis. Chapters 19, 23 and 24 close further exact analytic and geometric steps, with their remaining uniform hypotheses explicitly tracked.

The preceding bound for the columns is useful only if the resulting rows span a sufficiently large space. A merely qualitative nonvanishing or spanning theorem does not suffice: the integral-index bound is proportional to the number of translates used. We give the quantitative statement, develop the relative trace calculation on which it rests, and prove the finite-dimensional deduction. This section separates out the analytic and arithmetic estimates that are specialized to the large ranks, levels and weights of this proof. The term *Formal-source input* below denotes a result extracted from the indicated formal declaration; it is not asserted to have been independently reproved here. We retain the actual constants when they are used in another estimate. The Hilbert space arguments and the finite-field sieve deductions are given directly.

20.1 Statement and finite-dimensional framework

Let C denote the column space of algebraic modular forms, C^{ram} its subspace ramified at the chosen auxiliary prime, W the row space, and $\mathfrak{G}$ the finite group of level translations. Put $c = \dim_{\mathbb{C}} C$ and $n = \dim_{\mathbb{C}} W$. The maps $P_s : C^{\mathrm{ram}} \to W$ are the projected translates defined above, before the common rational rescaling of Theorem A. This rescaling has no effect on the image of any map.

Theorem 20.1 (Quantitative spanning, Theorem B)**.** *For admissible data $\mathcal{D}$ we have $c, n > 0$. There exists a finite subset $T \subset \mathfrak{G}$ for which*

$$|T|c \le 5n\log(1+b^2), \qquad \dim\Big(\sum_{s\in T} P_s(C^{\mathrm{ram}})\Big) \ge (1-4b^{-2})n.$$

This is `Data.enough_forms` in `New/Endgame/ShortAndEnough.lean`. We will deduce it from a local-to-global moment calculation and a finite-dimensional variance inequality, rather than take `Data.enough_forms` as an input. Notice the first inequality uses c, the dimension

of the entire column space: the actual index estimate of A costs at most cB per translate. It is not enough to bound $|T| \dim C^{\mathrm{ram}}$.

For an arbitrary subspace $J \subset W$, with fraction $\alpha = \dim J/n$, the operative intermediate result is

$$\dim(P_s D_N)^{-1}(J) \le ((1+\alpha)/2 + b^{-2})c \quad \text{for some } s \in \mathfrak{G}. \tag{20.1}$$

Here $D_N : C \to C^{\mathrm{ram}}$ is the ramifying operator of the previous section, viewed as a map onto its image. When J has dimension appreciably less than n, the dimension of the new image modulo J is correspondingly large. There are two analytical tasks: construct the original Poincaré map on C, and control its translates after discarding bad finite types. The ramification operator imposes only a small additional cost on the source rank.

20.2 Finite level, central projection, and the transfer bound

Write E^+ for the maximal totally real subfield of E. The row forms before projection belong to a finite-dimensional Hilbert space V of cusp forms on $U(Y_0)$ of the prescribed archimedean type and of principal level at the chosen split places of g_0, g_1. Its inner product is the Petersson pairing. The group of finite level translations is

$$\mathfrak{G} = \mathfrak{G}_0 \times \mathfrak{G}_1, \qquad \mathfrak{G}_i = \prod_{v \mid g_i \text{ chosen}} \mathrm{GL}_r(\mathbb{F}_v),$$

acting via a unitary representation ρ. In each product there are precisely f factors with residue cardinality g_i. Conjugate places are omitted from the indexing because the local unitary condition recovers the data at the conjugates. For an irreducible complex character ξ of a finite group G the element

$$e_\xi = \frac{\xi(1)}{|G|} \sum_{g \in G} \overline{\xi(g)}[g]$$

is a self-adjoint central idempotent. If $e = \sum_{\xi \in F_0} e_\xi$ for a set of irreducible characters, its coefficient at the identity is the Plancherel mass $\mu(F_0) = \sum_{\xi \in F_0} \xi(1)^2/|G|$. Consequently $\mu(F_0)$ is the fraction of the regular representation occupied by these isotypic spaces; it need not be the analogous fraction of V. Distinguishing these assertions is essential in the filling proof, which bounds the trace of the discarded part on the particular image of the Poincaré series, rather than deducing it from the discarded proportion in an arbitrary representation.

At a chosen place with residue field $\mathbb{F}_q$ the regular labels used in the proof are monic squarefree polynomials of degree r with nonzero constant term. Let M_ω denote the excluded torsion order, as in the definition of the level characters. A label is allowed if it has no root $x \in \mathbb{F}_q$ of $x^{M_\omega} = 1$. A primitive label is an irreducible polynomial with one (and hence all) roots of multiplicative order $q^r - 1$. The good tuples are tuples of allowed labels for which at least one label is primitive. The definitions are `IsPrimitiveLabel` and `goodTuples` in `New/Trace/BadShare.lean`; the corresponding idempotents are `labelIdem`, `tupleIdem` and `Data.e0` in the finite character theory and `New/Seesaw/IndexTypes.lean`. Let e_i be the good central projection in $\mathbb{C}[\mathfrak{G}_i]$ and $R = \rho(e_0 \otimes e_1)$. (Here e_0 denotes the first factor's projection; the code uses `Data.e0` for their product.) Our row space is $W = \operatorname{im} R$. The two factors are both used: the goodness requirement contains a primitive label above *each* g_i.

Lemma 20.2 (Plancherel deficit)**.** *Let $1 < r < q$ and $M_\omega \ge 1$. Suppose $0 \le \varepsilon \le 1$ and $L_0 \ge 0$ satisfy*

$$(1 + M_\omega + 2r)/q \le \varepsilon, \qquad 2r(2r \log q + 1)L_0 \le f.$$

Then the coefficient at the identity of $1 - e_i$ is at most $f\varepsilon + e^{-L_0}$. In particular, for our data, the complementary coefficient is at most $b^{-8}/2$ for each i.

Proof. The representation-theoretic input is the degree estimate of `labelChar_degree_bounds` for the Green characters, used in `bad_share_le`: each regular label contributes Plancherel mass at least

$$\mu_0 = (1 - r/q)q^{-r}.$$

We take this degree formula as the *Formal-source input* from the finite general linear group character theory (see `New/Trace/BadShare.lean` for the use of this bound). The rest is counting. There are q^r monic polynomials of degree r. For each of at most $1 + M_\omega$ prohibited linear roots (including zero), there are q^{r-1} polynomials divisible by the corresponding linear factor. The nonsquarefree polynomials belong to the union of the families Q^2P, where Q, P are monic and $1 \le \deg Q \le r/2$, $\deg P = r - 2\deg Q$. For fixed degree j there are at most q^{r-j} choices, so a crude bound for the total is rq^{r-1}. Hence allowed labels have total uniform lower mass at least

$$(1 - r/q)(1 - (1 + M_\omega + r)/q) \ge 1 - \varepsilon.$$

Primitive roots are precisely the $\varphi(q^r - 1)$ generators of the multiplicative group of the degree-r field. Such a generator is not contained in a proper subfield, so each of their minimal polynomials is irreducible of degree r and corresponds to exactly r generators. The number of primitive labels is therefore $\varphi(q^r - 1)/r$. The elementary totient lower bound used in the source is $\varphi(u) \ge u/(2\log u + 1)$ for $u \ge 2$ (`le_totient_mul_log`). Thus the uniform mass q^{-r} assigned to these labels is at least $1/[2r(2r\log q + 1)]$. The mass of nonprimitive labels with respect to the weight μ_0 is at most their mass with respect to q^{-r}, whence it is at most $1 - L_0/f$. On the product of f factors, the mass lower estimate for all allowed labels is $(1-\varepsilon)^f$, and the uniform lower mass of the allowed tuples that are not primitive anywhere is bounded above by $(1 - L_0/f)^f$. Thus the mass of the good tuples is at least $(1-\varepsilon)^f - (1 - L_0/f)^f$. Subtracting from one, and using Bernoulli's inequality and $(1-u)^f \le e^{-fu}$, proves the claimed bound.

For the last specialization one takes $L_0 = 9\log b$. The size inequalities used in `Data.ramifiedColumns_fill_rows`, together with $g_i \ge b^{10}$ and f sufficiently smaller than the level, give $f\varepsilon + b^{-9} \le b^{-8}/2$. These exact numerical inequalities on the data are the *Formal-source input* `bad_share_le` as instantiated in `New/Filling/FillingTheorem.lean`. The counting proof explains the two separate contributions: excluded labels are individually rare, while failure to contain even one primitive label is exponentially rare across the many factors. □

The second input is a bound on the actual number of target rows. Let $d_\infty = \prod_{v\in\Phi} d_v$ be the product of the local formal degrees (defined through the Bessel systems below) and let $\mathrm{mass}_r(E)$ be the mass of the rank-r positive definite unimodular genus, with finite stabilizer weights. Define

$$n_0 = |\mathfrak{G}| d_\infty \,\mathrm{mass}_r(E). \tag{20.2}$$

Proposition 20.3 (Dimension transfer (conditional derivation in Chapter 18))**.** *One has $n \le n_0$. More precisely the trace of every element of the common split unramified Hecke algebra on good forms of the same rank, level and compatible archimedean Harish–Chandra data is independent of the unitary form, in the rigid-label situation of this proof.*

The statements are `trace_heckeOnGood_eq`, `det_heckeOnGood_eq` and `finrank_rows_eq_finrank_goodForms` in `New/Trace/GoodFormTransfer.lean`, and `Data.finrank_rows_le_mass` in `New/Filling/RowDimension.lean`. For the dimension computation, the primitive-label conditions and the local rigidity lemma in `New/Trace/LabelRigidity.lean` are needed in the trace formula, as well as the existence of projectors onto the finite Hecke eigenspaces. The proof transfers the row forms to the definite member of rank r, where good forms are a subspace of the full space of algebraic modular forms. Here the class stabilizers

are trivial at the chosen level. The class sum weighted by the reciprocal group orders is $|\mathfrak{G}|\,\mathrm{mass}_r(E)$ by the level-index identity; the dimension of the finite-dimensional algebraic representation at infinity is d_∞ in the transfer normalization. Multiplying gives (20.2). The nontrivial passage between signatures is not a formal consequence of dimension equality and cannot be omitted; Chapter 18 proves the trace comparison and dimension deduction from its explicitly stated elliptic orbital and finite rigidity hypotheses, including the precise formal-degree normalization. The same trace equality is used again to compare auxiliary lifts with the main chain in the lower-bound argument.

20.3 Hermitian geometry of the Bessel subgroup

We describe the objects for the unfolding calculation. In this section use $h = m/2 - 1$ and $s_0 = h + 1$ (not the dimension n of row forms). The admissible rank $r = k + 2h + 1$ is the rank of the split extension of the definite rank-k space X_0. It has ordered basis of h hyperbolic pairs e_i, e_i^* $(0 \le i < h)$, an orthonormal basis of X_0, and one middle vector z with norm $-N$. Thus the Gram matrix in the rational basis is the sum of the hyperbolic planes, I_k, and $(-N)$. Denote this form by Y. Let U be the unipotent radical of the stabilizer of the isotropic flag spanned by successive initial e_i, with its standard Bessel character, and let H_0 be the subgroup isomorphic to $U(X_0)$ acting as the identity on the frame and its dual. The Bessel subgroup is $S_B = UH_0$. Its frame is $F = (e_0, \dots, e_{h-1}, z)$; U^0 is the pointwise frame stabilizer in U. The coordinate form of these groups, including the flag, is the `Shape` construction of `New/Filling/EichlerFlag.lean` and the groups in `Lib/Adelic/Poincare/BesselSubgroup.lean`.

At finite places put $N = \pi\bar{\pi}$, interpreted as an idelic factorization if necessary, and modify the basis of $X_0 \oplus Ez$ by replacing its selected vector e_* with $v_* = e_*/\bar{\pi} + z/N$. Its self-pairing is $1/N - 1/N = 0$ and its pairing with z is -1; the 2×2 middle Gram block becomes

$$\begin{pmatrix} 0 & -1 \\ -1 & -N \end{pmatrix}$$

with determinant -1. All other middle basis vectors retain their original pairings; the hyperbolic planes are unchanged. Thus the lattice Gram matrix is unimodular. These are the formulas of `latMiddleGram` and `gram_besselLatticeBasis` in `New/Filling/EichlerFlag.lean`. Let K_f be the stabilizer of this lattice away from the infinite places, and $K(g)$ the kernel of reduction to the above finite group. The column level is the inverse image of $K(g)$ under inclusion of H_0; this is one of the hypotheses in the filling record. Let $\mathcal{C} = H_0(E^+)\backslash H_0(\mathbb{A}_f)/K_c$ and h_u be the chosen images of representatives of classes $u \in \mathcal{C}$.

Proposition 20.4 (Formal-source input: global filling datum)**.** *The admissible data give a filling system in the sense of* `Filling` *and* `Filling.IsAdmissible` *of* `New/Filling/Setup.lean`*, with the prescribed r, k, h, N, g_i, b, whose column stabilizers are trivial, reduction onto $\mathfrak{G}$ is surjective, and whose Bessel coordinate spaces at each infinite place identify with the column weight representation. The resulting column coordinates give an isomorphism*

$$C \simeq \bigoplus_{u \in \mathcal{C}} \mathbb{C}^I, \qquad I = \prod_{v \in \Phi} I_v.$$

In particular $c = |\mathcal{C}||I| > 0$.

The coordinate identification is `Data.columnCoords` in `New/Seesaw/BesselGeneric.lean` (the coordinate model is built in `New/Filling/ColumnCoords.lean`); the filling construction

is `New/Filling/FillingData.lean` and the admissibility check is `New/Filling/FillingAdmissible.lean`. Positivity follows since the class of the identity and every weight coordinate set are nonempty. The dimension formula also follows directly from the double-coset description when stabilizers are trivial. For the local-to-global estimates below we record several of the other numerical hypotheses, rather than replace them by unspecified implied constants. If ϖ is the purely imaginary generator of the absolute different and $\kappa_v = \Re(gv(\varpi)/(2\pi i))$, then

$$1 \le |\kappa_v| \le b^{23}, \quad |v(\varpi)| \le b, \quad N \le b^{60 s_0}, \quad b^{10} \le g_i \le b^{11}.$$

Also $h \ge 10^5$, $1000(2h+2)^9 \le k \le f$, $s_0^7 \le k$ and $(200 f k \operatorname{rd}_E)^{1000} \le b$. These are much stronger than the hypotheses of the elementary rank argument. They enter below in estimates which would be false uniformly for arbitrary families of lattices of varying dimension.

The normalized archimedean basis at v is scaled by $\kappa_v^{h-i}/\sqrt{N}$ on e_i, its inverse on e_i^*, and $1/\sqrt{N}$ on z. On X_0 the basis is unchanged. The middle norm is thereby normalized to -1. Put

$$l_{vi} = \frac{\kappa_v^{2(h-i)}}{N} \quad (0 \le i \le h), \qquad \mathcal{L} = \prod_{v \in \Phi} \prod_{i=0}^{h} l_{vi}^{k+2(h-i)}.$$

For a rational element γ of $U(Y)$ define the frame-product matrix $Z(\gamma) = (\langle F_i, \gamma F_j \rangle)_{0 \le i,j \le h}$, and write $D_i(\gamma)$ for its leading principal minor of order i. The discriminant separating the open and smaller cells is

$$\mathfrak{d}(\gamma) = D_{h+1}\overline{D_{h+1}} - N^2 D_h \overline{D_h}. \tag{20.3}$$

The normalized archimedean matrix Z_v has entries $\sqrt{l_{vi} l_{vj}}\, v(Z(\gamma)_{ij})$ up to signs of the real scales (the scales themselves are chosen real, not assumed positive); its normalized discriminant is $|\det Z_v|^2 - |D_h(Z_v)|^2$. Direct expansion shows that it equals $(\prod_i l_{vi})^2 v(\mathfrak{d}(\gamma))$, since the ratio between the square of the products of the full scalings and of the first h scalings is $l_{vh}^2 = N^{-2}$. The definitions are `Filling.frameProducts`, `frameScaleSq`, `termDisc` in `Setup.lean`, and `frameDisc` in `EichlerFlag.lean`. The nonnegativity of this discriminant for actual frame products and the strict inequality in the full open cell are geometric consequences of the positive definiteness of the complement of the hyperbolic planes, recorded in `NearTerms.lean`.

This particular frame is relevant both to the archimedean kernels and to counting at the finite level: the leading minors act as pivots. It is important that one does not have to count all pairs of rational points of the full rank-r group indiscriminately. Coherent pairs are sorted by common frame data and by a linear relation between the two rank-k middles; then the deepest strata admit a local orbital calculation. The smaller strata are controlled by means of the same discriminant and congruences.

20.4 The archimedean Bessel system

For each $v \in \Phi$ the group at the normalized real place is the unitary group $U(k+h, h+1)$ in the split basis just described. Denote it by G_1. We use Haar measures in the normalization of `BesselSystem.lean`; in particular the measure on $U\backslash G_1$ is the quotient by the specified unipotent measure and is not a probability measure. The compact type of $H_0(\mathbb{R}) \simeq U(k)$ is identified with the prescribed column representation (of dimension $|I_v|$). The archimedean parameter of the target representation has large gaps: the local construction uses a gap at least r^7 and each doubled parameter has absolute value at most $3a$. The finite-dimensional column type has weight $(\kappa_v^{\mathrm{col}}, 0, \dots, 0)$ in the convention of the proof: its only potentially nonzero entry is the one corresponding to the marked first column of order a at the tops. This weight should not be confused with the real scaling constant κ_v just introduced.

The representation is described in Lean via a reproducing kernel rather than via a choice of realization of a discrete-series Hilbert space. An invariant kernel of formal degree d_v, acting by convolution, reproduces the functions of its type. Its integral against the Bessel character constructs a relative kernel; the associated map on the compact type is normalized to have prescribed L^2 norm. The source constructs a full function and a cutoff with unusually strong uniform bounds despite the changing representation and rank. Here are the exact properties that will be used.

Definition 20.5 (Quantitative Bessel system)**.** For a real place v, let I_v be a nonempty finite set, let $D_v = |I_v|$, and let $d_v > 0$. Let W_v (full) and $W_v^\flat$ (cut) be continuous functions on G_1 with values in $\mathbb{C}^{I_v}$, equivariant on the left under U through the fixed character and under H_0 through the specified irreducible unitary representation. Define the relative size

$$\mathrm{bs}(y) = \inf_{s \in S_B} (1 + \|sy\|_{\mathrm{op}}).$$

The following conditions are part of the quantitative Bessel system:

1. The full function is integrable on $U\backslash G_1$, its squared L^2 norm is D_v, and the cut has squared norm at most D_v and vanishes where $\mathrm{bs}(y) > b^{6s_0+2}$. Moreover
$$\|W_v^\flat - W_v\|_2^2 \le b^{-10r^2} D_v.$$

2. There are a measurable left U-invariant majorant A and a finite measure μ on G_1 such that each of the two norms is bounded by A, and in the region where the functions differ their norms are bounded by $b^{-3r^3 f} A$. In addition
$$A(y) \le \mu\{g : \mathrm{bs}(yg) \le b^{6s_0+2}\}, \qquad \mu(G_1) \int_{U\backslash G_1} A(y)\, dy \le b^{2r^3} D_v.$$

3. For each word α in the matrix-coordinate directions, with $|\alpha| \le 2s_0^2$, there is a continuous derived Bessel function $W_{v,\alpha}$. Its coordinate right differentiation relation is the ordinary chain rule along a curve $\gamma'(t) = \gamma(t)X$, namely the derivative of $W_{v,\alpha}(\gamma(t))$ is the sum of the coordinates X_{ij} times $W_{v,(i,j)\alpha}(\gamma(t))$. For every $P \ge b^5$ and every integrable scalar ϕ on G_1, the derived Schur bound is
$$\int_{U\backslash G_1} \left\| \int_{G_1} \phi(y) W_{v,\alpha}(\gamma y)\, dy \right\|^2 d\gamma \le P^{2|\alpha|} \frac{D_v}{d_v} \|\phi\|_2^2. \tag{20.4}$$

4. The Bessel correlation of the cut, after the changes of coordinates below, belongs to a tower of kernels with explicit Fourier bounds, called a sieve system. Its bottom constant is $(1 + b^{-10})/d_v$.

These are precisely `IsCutPair`, `HasSchurBound`, `BesselSystem` and `HasBandLimits` of `New/Filling/BesselSystem.lean`. The word "derived" refers here to differentiation, not to derived categories. Condition (2) is stronger than L^2 approximation alone: after summing translates over a growing lattice, a small L^2 error does not automatically imply a small contribution to the second moment.

Proposition 20.6 (Formal-source input: existence of local systems)**.** *There exist quantitative Bessel systems with the prescribed formal degrees, type, and column-coordinate representations at all $v \in \Phi$. The cutoff system satisfies the Fourier bounds of* (20.6) *below, and the open-cell estimate of* (20.7) *below.*

The existence assertion is `Data.exists_besselSystem` in `New/Filling/FillingData.lean`, obtained from `HasPeakAt.exists_cut_isCutPair` and `IsBesselFunction.hasSchurBound_of_convolve_eq` in `New/RealPlace/BesselCut.lean` and `hasBandLimits_of_isCutBessel` in `New/RealPlace/BandLimits.lean`. The local construction obtains a peak of a Bessel convolution by lifting between unitary pairs with separated parameters; localization and lower bounds on the peak are what permit normalization by the peak value. The gap and the bound on the highest parameters are used both in the tail of the kernel and to control iterated right derivatives. For example the derivative estimate for a reproducing kernel of rank r, highest doubled parameter bounded by A_0, permits a Schur constant P at least $(r+1)^4(A_0+r+1+2s_0^2)$; the admissible data absorb this in b^5. In reproducing-kernel notation the reason for the Schur estimate is that convolution is an orthogonal projection up to the factor d_v. After normalization and differentiation, its operator norm is bounded by $d_v^{-1/2}$ times the norm of the differentiated vector. The nontrivial uniform part is bounding those vectors and obtaining the stated peak and cutoff, parts of the above formal-source input; neither follows merely from the existence of a discrete-series representation.

Lemma 20.7 (Orthonormality of the full coordinates)**.** *For each v and all $i, j \in I_v$,*

$$\int_{U\backslash G_1} W_{v,i}(y)\overline{W_{v,j}(y)}\,dy = \delta_{ij}.$$

Proof. The integral matrix $T = \int W_v(y)W_v(y)^* dy$ is a Hermitian positive semidefinite operator on the coordinate space. Left invariance of the quotient measure under the compact group H_0 and Bessel equivariance show that T commutes with its irreducible unitary representation. Schur's lemma gives $T = aI$ with $a \geq 0$. Taking traces and using condition (1) gives $aD_v = D_v$, so $a = 1$. This is the argument of `BesselSystem.orthonormal` in `New/RealPlace/BesselOrthonormal.lean`. □

We record the sharper local correlation bounds because they fix the constant in the global moment. For any vector-valued W on G_1 let

$$J_W(\gamma) = \int_{G_1} W(y) \otimes W(\gamma^{-1}y)^*\,dy$$

when this integral is defined. For $W = W_v^\flat$, the full open-cell Weyl representative with head $p = (p_0, \ldots, p_h)$ is denoted w_p. Here $p_i \neq 0$ are the successive pivot ratios in elimination of the frame-product matrix; for $i = h$ we have $p_h = \langle z, \sigma z\rangle$ for the residual middle element $\sigma \in U(X_0 \oplus \mathbb{C}z)$. Let $V(U(j))$ be the volume of the compact unitary group in the source normalization and put

$$c_G = \frac{V(U(k))}{V(U(r))}\frac{2\pi^k}{(k-1)!}, \qquad c_{\mathrm{band}} = \frac{V(U(k))}{V(U(r))}\pi^{(2h+1)k}.$$

For matching top entries $|p_h| > 1$ define the normalized correlation tensor

$$T_v(\sigma, p) = \frac{\sqrt{c_G}\,(|p_h|^2-1)^{(k-1)/2}}{D_v}\prod_{i<h}|p_i|^{k+1+2(h-i-1)}\,J_{W_v^\flat}(w_p\sigma). \tag{20.5}$$

The normalization of the tensor has a particular Jacobian meaning: the powers in the product are the half-densities of the successive rank-one steps in the open cell. In the last step the residual middle group contributes the volume of a sphere of dimension $2k-1$ and the radial factor $(|p_h|^2-1)^{k-1}$. This is the real factor that will compensate the global finite-place quadric count.

The top kernel of the local sieve system is the inner product of the tensors (20.5), when the two top entries both come from σ. Hence the top kernel is positive semidefinite for common top entry. The frequency bound for the system has the following explicit majorant. Write

$$\lambda_a(u) = u^a \min(1,u)^6, \qquad \Lambda_i(p) = \prod_{j<i} \lambda_{s_0^6}(|p_0 \cdots p_j|^{-1}).$$

In the integral of one common entry at level i the nonzero Fourier coefficient, at frequency ξ, is bounded in absolute value by

$$\beta_v(\xi)\Lambda_i(p)\Lambda_i(q), \qquad \beta_v(\xi) = c_{\mathrm{band}} b^{2000 s_0^9} \lambda_{k-s_0^7}(|\xi|^{-1}). \tag{20.6}$$

The same bound in the unequal-entry relation (with the next head weights) is used below. These are the actual constants in `HasBandLimits`. The large exponent $k - s_0^7$ is critical; a bound merely asserting rapid decay for fixed parameters would not provide the required powers after multiplying over f places. The exponent 6 in $\min(1,u)^6$ is a separate damping term used for summability of frequencies after applying the product formula. The construction of these kernels and the bounds (including the identity of the top kernel with the correlation) is the formal-source input `hasBandLimits_of_isCutBessel` and the preceding real-place calculations in `New/RealPlace/BandLimits.lean` and `LocalTheorem.lean`.

A second local estimate is used for the terms that cannot be treated by the sharp main-term calculation. Let $Z_v(\gamma)$ denote the real-place frame-product matrix and let $\mathfrak{d}_v = |\det Z_v|^2 - |D_h(Z_v)|^2 > 0$. If $\rho_0 \geq b$ and the box in which all frame coordinates have norm at most ρ_0 meets its translate by γ, define

$$M_v(\rho_0,\gamma) = \left(\frac{\rho_0^{9s_0}}{\mathfrak{d}_v}\right)^{10^5 s_0^4} \frac{D_v^2}{d_v c_G (\mathfrak{d}_v/2)^{k-1}}. \tag{20.7}$$

The open-cell estimate (*Formal-source input*: `BesselSystem.open_cell` in `New/RealPlace/OpenCellBound.lean`) asserts the existence of a compact integration set, an averaging parameter in $U^0 H_0$, and a measure of total mass at most M_v, such that the averaged squared norm of the integrals against $W(\gamma ty)W(ty)^*$ is at most the integral of the squared modulus of the test function against this measure. The compact integration set contains the required translates of the box, almost surely in the averaging parameter. This is a local operator bound with an explicit discriminant pole, not a pointwise estimate for W. The normalization in (20.5) is intentionally different: in the sharp term one preserves cancellation between Fourier phases, while in (20.7) one uses upper bounds allowing losses. The large exponent on $\mathfrak{d}_v$ is compensated globally by discriminant bounds and congruence counts.

20.5 Construction and exact unfolding of the Poincaré map

Using the coordinate isomorphism of Proposition 20.4, endow C with its orthonormal coordinate basis $e_{u,i}$ for $u \in \mathcal{C}$, $i \in I$. Write $W(y)_i = \prod_v W_{v,i_v}(y_v)$ and similarly $W^\flat$ for the cut. The superscript $\flat$ now denotes the approximate function, not the auxiliary cutoff on unipotent cosets. The product Bessel functions live on the product of the normalized infinite groups, denoted G_∞. For any rational element γ, use γ_∞ and γ_f for its images in these normalized coordinates and in the finite lattice coordinates, respectively.

For $u \in \mathcal{C}$, let $U_u(g)$ be the rational unipotents with $h_u^{-1}\gamma_f h_u \in K(g)$. The image at infinity is a discrete group. Take a measurable nonnegative class cutoff β_u on G_∞ with

$$\sum_{v \in U_u(g)} \beta_u(vy) = 1. \tag{20.8}$$

Such a function may be obtained by taking a measurable fundamental domain, or by normalizing a nonnegative locally finite function that is positive somewhere on each orbit. The choice in the source is expressed as a cutoff for a discrete subgroup. The arithmetic assertion of discreteness, and the existence of a cutoff with the specified compatibility, is `exists_cutoff_classUnip` in `New/Filling/Setup.lean`. The measures are chosen so that the covolume of $U_u(g)$ in the infinite unipotents is a constant $\nu > 0$ independent of u. Explicitly, put $d_U = h(h+k)$ and $D_E = |\operatorname{disc}(E)|$. With $v_f = \operatorname{vol}(\widehat{\mathcal{O}}_E)$ in the self-dual finite measure of the source, the covolume is

$$\nu = \frac{(g^{2d_U+h}/N^k)^f}{v_f^{d_U}} \frac{D_E^{h/4}}{\mathcal{L}}. \tag{20.9}$$

This is `Filling.unipCovolume`. To see the origin of the exponent, the unipotent filtration has a middle step of size $k+1$ and two strings of sizes i, i for each $0 \le i < h$; the sum of their sizes is $\sum_i (2i+k+1) = h(h+k)$. Each such adelic coordinate has congruence density and a self-dual covolume, while the normalized archimedean change of basis supplies the Jacobian $\mathcal{L}$. The norm term N^{-kf} occurs in the middle lattice modification. The exact quotient formula (20.9), including independence of u, is the *Formal-source input* `Filling.lintegral_classCutoff_eq_unipCovolume` of `New/Filling/UnipotentCovolume.lean`; the numerical factors are those of `Setup.lean`. For an integrable function F on the unipotent quotient, the cutoff identity implies

$$\int_{G_\infty} \beta_u(y) F(y)\, dy = \nu \int_{U_\infty \backslash G_\infty} F(y)\, dy$$

provided F is invariant on unipotent orbits: integrate first over the left unipotent and then use (20.8) to replace that integral by its quotient covolume. This is the quotient calculation used repeatedly in the unfolding.

For (u, i) let $\phi_{u,i}$ be the seed given by $\beta_u(y) W_i(y)$ at infinity times the indicator that $h_u^{-1} g_f \in K(g)$, in the common coordinates. A fixed common adelic coordinate change is understood (the map called `Filling.Fits.finiteEquiv` and the inclusion of its fixed η in `New/Filling/PoincareSeries.lean`); it has no effect on the formulas for relative positions since it cancels from them. The Poincaré map $Q : C \to V$ is the periodization

$$Qz(g) = \sum_{u,i} z_{u,i} \sum_{\gamma \in G(E^+)} \phi_{u,i}(\gamma g). \tag{20.10}$$

When summing first over unipotents, the periodization replaces the cutoff β_u by 1 on the permitted finite coset, by (20.8) and the Bessel character being trivial on $U_u(g)$. This identifies (20.10) with the Poincaré series on the Bessel quotient; note that the source function called `cutBessel` in this construction contains the full W and the cutoff β_u, not the cut $W^\flat$.

To state the convergence and unfolding precisely, put

$$\operatorname{pos}_{u,u'}(\gamma) = h_u^{-1} \gamma_f h_{u'}, \qquad \mathcal{E}(u, s, u') = \{\gamma : \operatorname{pos}_{u,u'}(\gamma) \in K_f, \operatorname{red}(\operatorname{pos}_{u,u'}(\gamma)) = s\}.$$

Here K_f is the lattice stabilizer used in the filling datum, with reduction kernel $K(g)$. For a set T of rational terms define the amplitude matrices (coordinates $I \times I$)

$$A(\gamma; u, u') = \int_{G_\infty} \beta_u(y) \beta_{u'}(\gamma_\infty^{-1} y) W(y) \otimes W(\gamma_\infty^{-1} y)^*\, dy, \tag{20.11}$$

$$A_T(u, s, u') = \sum_{\gamma \in T \cap \mathcal{E}(u,s,u')} A(\gamma; u, u'). \tag{20.12}$$

Define $A_T^\flat$ with $W^\flat$ in place of W.

Proposition 20.8 (Formal-source input: convergence and reproduction)**.** *For every u, s, u' the sums in (20.12) are absolutely convergent in the finite-dimensional tensor space. For fixed u, s the sum over u' and the corresponding rational terms of the integrated norm difference of the full and cut kernels, with the product of the two cutoffs as in (20.11), is at most $\nu b^{-k^3 f}/2$. The sum of the norms in the inner periodization in (20.10) is bounded uniformly in g. The Poincaré map has image in V.*

The first assertions are `amplitudeAbs_lt_top_and_cut_le`, `summable_amplitude` in `New/Filling/AmplitudeSums.lean`; the last two are `Fits.tsum_cutBessel_le`, `Fits.poincareMap` in `New/Filling/PoincareSeries.lean`. They are not consequences of pointwise boundedness of the seed alone. The majorant in Definition 20.5(2), together with the bounded box partition at infinity and lattice estimates, is used to prove convergence before interchanging the integrals. The reproducing-kernel condition and integration on the Bessel quotient then put the Poincaré series in the space of the required archimedean cusp type and principal finite level.

Lemma 20.9 (Exact coefficient identity)**.** *In the Petersson normalization there is a constant $v_0 = \mathrm{vol}(K(g)) > 0$ such that*

$$\langle Qe_{u,i}, \rho(s) Qe_{u',j} \rangle = v_0 A_{\mathrm{all}}(u, s, u')_{i,j}. \tag{20.13}$$

Here and below the pairing uses the convention in which the order of i, j agrees with (20.11); if one takes the opposite convention, both sides are conjugated/transposed accordingly.

Proof. Unfold the first periodization in the quotient Petersson integral and expand the second into its defining series. This gives a single sum of integrals over the whole adelic group, of the product of a seed of class u with a translated conjugate seed of class u'. Absolute convergence and Fubini are justified by Proposition 20.8 and the integrability statement `Fits.integral_poincare` in `New/Filling/PoincareSeries.lean`. After left translating the finite integration variable by the class representative h_u^{-1}, its range is $K(g)$. Normality of $K(g)$ in K_f implies that the second indicator is identically 1 on this range precisely when the relative element has reduction s, and otherwise identically zero. Hence the finite integral contributes v_0 and restricts the rational terms to the set $\mathcal{E}(u, s, u')$. The archimedean part of the same integral is (20.11); the two class cutoffs remain in it. Thus the integral is precisely (20.13). This is the unfolding carried out by `Fits.petersson_poincare` in `New/Filling/PoincareSeries.lean`. □

Set $P = (v_0\nu)^{-1/2} Q$. Then $P^* \rho(s) P$ in the coordinate basis has entries $\nu^{-1} A_{\mathrm{all}}(u, s, u')_{i,j}$. Projecting by R gives the translates of the row map after restriction to the ramified space, up to their fixed nonzero scalar. It is the normalized map P that will enter the variance estimate, whereas the integral-index theorem uses the actual unscaled map of Theorem A.

Lemma 20.10 (Identity term)**.** *The amplitude of the Bessel subgroup at $s = 1$ is exactly*

$$A_{S_B}(u, 1, u')_{i,j} = \nu \delta_{u,u'} \delta_{i,j}.$$

In particular the Bessel-subgroup contribution to P^*P is the identity operator.

Proof. The elementary reduction property used here is `Filling.eq_and_mem_classUnip_of_besselS` in `New/Filling/NearTerms.lean`: a term in S_B over $(u, 1, u')$ forces the two column classes to be equal and belongs to $U_u(g)$. This follows by projecting to H_0 and using the definition of the column classes and triviality of their automorphism groups at K_c (the precise compatibility with the modified lattice and the level is the cited formal-source input). Thus only

$u = u'$ can contribute. On summing over the unipotents, covariance of W and (20.8) eliminate the second cutoff. The remaining integral is ν times the integral of $W_i\overline{W_j}$ on the unipotent quotient. Use the product over v of the orthonormality relation in Lemma 20.7. This proves the claimed identity (`amplitudeSum_besselS_one` in `New/Filling/AmplitudeSums.lean`). □

20.6 Energy, pairs of congruent terms, and removal of lower cells

The following notation allows the estimates for the moment to be stated with their actual normalizations. For a subset $\Omega \subseteq \mathfrak{G}$ define

$$\mathcal{E}(A_T;\Omega) = \frac{1}{\nu^2}\sum_{u,u'\in\mathcal{C}}\sum_{s\in\Omega}\|A_T(u,s,u')\|_{\mathrm{HS}}^2. \tag{20.14}$$

The norm here is the Euclidean norm on the $I \times I$ tensor. Put

$$D = \frac{c^2}{d_\infty \operatorname{mass}_r(E)}, \quad D_1 = |\bar{S}_B|c, \quad \tau = b^{kf/4}, \quad \mathfrak{c} = 2\tau^2(b/g)^{kf},$$

where $\bar{S}_B$ is the reduction of the integral Bessel subgroup. These are `mainEnergy`, `besselEnergy`, `tailScale`, and `codimFactor`, respectively; the formulas and inequalities in the following proof use this normalization. In particular, the admissibility assumptions give

$$\tau^{44}D_1 \le D, \qquad \tau^{18}\mathfrak{c}D \le D_1, \qquad \mathfrak{c} \le 1. \tag{20.15}$$

This pair of bounds is central: the Bessel-subgroup energy, despite its large number of translations, is extraordinarily small on the scale of the full expected moment. The first bound means it is inexpensive to use coarser estimates on the lower cells; the second expresses the strong saving from nontrivial finite congruence codimension. They are part of `Filling.IsAdmissible` and are verified for the data in `New/Filling/FillingAdmissible.lean`; the verification is a *Formal-source input* using the mass and unipotent measure formulas and degree bounds. The first equality in (20.2) shows $D/|\mathfrak{G}| = c^2/n_0$.

For two compatible collections of amplitudes A, B, any $\delta > 0$ gives

$$\mathcal{E}(A+B;\Omega) \le (1+\delta)\mathcal{E}(A;\Omega) + (1+\delta^{-1})\mathcal{E}(B;\Omega). \tag{20.16}$$

This follows from $2\Re\langle a,b\rangle \le \delta\|a\|^2 + \delta^{-1}\|b\|^2$ in each summand. By absolute convergence the amplitude for a set decomposes into amplitudes for disjoint subsets. The global truncation error follows from Proposition 20.8 by applying the triangle inequality to each tensor, squaring, and summing:

$$\mathcal{E}(A_T - A_T^{\flat};\Omega) \le |\mathcal{C}|\,|\mathfrak{G}|\,b^{-2k^3f}/4 \le b^{-30}D_1.$$

For the last inequality use $|\mathfrak{G}| \le (g_0g_1)^{r^2f} \le b^{22r^2f}$, $c \ge |\mathcal{C}|$, $|\bar{S}_B| \ge 1$, and k much larger than h, hence $2k^3f - 22r^2f \ge 30$. This is `energy_sub_cut_le` in `New/Filling/SchurTest.lean`. Thus there is substantial freedom to cut off the Bessel functions. No comparable freedom is allowed in the constant of the main term.

The expansion of the squared amplitude in (20.14) gives pairs of rational elements *with equal finite reductions*: for each (u,s,u'), their relative positions belong to the same coset of $K(g)$ in K_f. These pairs are stratified by degeneracy of their flags and by the links between their middle spaces. We describe the stratification and record its precise resulting estimate.

Character orthogonality. Call a set of terms stable if it is invariant under left and right multiplication by rational unipotents. Write ψ_U for the additive Bessel character on rational unipotents as read in finite adelic coordinates; equivalently it is the inverse product of the infinite-place characters. For a term γ, a necessary condition to contribute is that

$$\psi_U(u) = \psi_U(\gamma^{-1}u\gamma) \quad (u \in U(E^+) \cap \gamma U(E^+)\gamma^{-1}). \tag{20.17}$$

Indeed the product $W(y) \otimes W(\gamma_\infty^{-1}y)^*$ transforms, on replacing y by $u_\infty y$, by the quotient of these characters. To see why cutoffs cause no obstruction, group the sum over $T \cap \mathcal{E}(u, s, u')$ into $U_u(g)$–$U_{u'}(g)$ double cosets. On a double coset, the product of the cutoffs can be replaced by any measurable weight of total one over the double action: this follows by Fubini from summing the two cutoff identities (20.8). The resulting integral is an orbit integral, and the mismatch character on the stabilizer makes it vanish. For a nontrivial character choose an element at which it is not one; the integral equals that nontrivial scalar times itself. In the formal proof the rational stabilizers relevant to the finite cosets and the full unipotent intersection give the same vanishing condition by the approximation theorem for the unipotent group, and the exact weight-change identity (*Formal-source inputs*: `hasSum_amplitudeSum_change_of_weight` in `New/Filling/AmplitudeSums.lean`, `amplitudeSum_eq_zero_of_not_charsAgree` in `New/Filling/CrudeEnergy.lean`, and `exists_rat_approx_unipotentChar_eq` in `NearTerms.lean`). The character must be retained until this step; replacing it by its modulus would lose the exact vanishing.

Normal forms and links. A term with degenerate discriminant $\mathfrak{d}(\gamma) = 0$ but compatible characters has a shared vector in the frame flags. The proof uses a descending family of subgroups G_j fixing the first j hyperbolic pairs ($0 \le j \le h$). For such a shared vector, compatibility of characters forces equality of the Hermitian norms of the two leading coefficients and propagates the shared-vector relation to subsequent levels. To see the mechanism, let v be a vector orthogonal to the relevant preceding flag, and w an isotropic vector of the flag lying in that orthogonal complement. The unitary Eichler transformation is

$$x \longmapsto x + w\langle v, x\rangle - (v + \eta\langle v, v\rangle w)\langle w, x\rangle, \qquad \eta + \bar{\eta} = 1.$$

Expanding the two pairings using $\langle w, w\rangle = \langle w, v\rangle = 0$ and the relation on η shows that this transformation preserves the Hermitian form. Its simple-root coordinates against the flag are linear in the coefficients of w and in the pairings of v with the successive flag vectors. The Bessel character of the transformation tests this linear combination. Applying the transformation with the corresponding common vector on each side of the conjugation, equality of characters for all rational scalings, and nondegeneracy of the additive pairing, forces equality of the linear functionals. Elementary linear algebra then propagates the shared-vector relation up the flag. In the source these steps are `exists_mem_besselU_eq_eichlerMap` in `EichlerFlag.lean`, `hasCommonFlagVector_of_le` and `CharsAgree.exists_hasLevel` in `CrudeEnergy.lean`. The latter supplies the exact assertion used below: a term satisfying (20.17) with discriminant zero has a level $j \le h$ in the sense of `Filling.HasLevel`. We use this implication as a *Formal-source input*, retaining its character hypothesis.

The link of a term is a linear subspace of $X_0 \oplus X_0$. In the split basis, let $M(\gamma)$ have as columns $e_0, \dots, e_{h-1}, \gamma e_0, \dots, \gamma e_{h-1}$ and let p_{X_0} denote projection to the rank-k positive middle coordinates. The link is the image of the kernel of $M(\gamma)^* J_Y$ under the map

$$z \longmapsto (p_{X_0} z, p_{X_0}\gamma^{-1} z).$$

It describes the relation between the two positive middle coordinates after imposing orthogonality to both isotropic frames. This definition is `link` and `linkMatrix` in `New/Filling/`

`EichlerFlag.lean`. When two terms have different links, their common part has smaller dimension; the resulting loss is reflected in the congruence codimension factor $\mathfrak{c}$. For the top cell (nonzero discriminant) the primary surviving energy is the sum of squared norms of blocks for a fixed link, not the sum of squares of individual terms. For a set $\mathcal{L}$ of links put

$$\mathcal{E}_{\mathrm{link}}(T,\Omega,\mathcal{L}) = \sum_{u,u',s\in\Omega}\sum_{\Lambda\in\mathcal{L}} \|A^{\flat}_{T\cap\{\mathrm{link}(\gamma)=\Lambda\}}(u,s,u')\|^2.$$

Unlike (20.14), this definition does not divide by ν^2; we keep the source normalization.

To estimate the discarded terms one first replaces the double cutoff by a partition in frame boxes of size $b^{2s_0^2}$, using a cutoff modulo the frame stabilizer U^0. Within such boxes all frame-product entries have bounded archimedean norm. Their finite congruence constraints and the product formula give lower bounds on the nonzero discriminants and on relevant lattice indices. The proof further selects near representatives of the U^0 double cosets (`IsNearRep` of `BoxPartition.lean`). For each pair of near representatives with matching finite reductions and level j, one bounds its contribution by products of explicit weights, introducing the lattice of common link vectors. The weights are inverse powers of: the discriminant of the lattice of link vectors, the index of the frame-value lattice, the saturation index of the paired frame lattice, and the factor $g^{2f\min(j,h)}$. In particular the pair invariant of `pairInvariant` is the product of the discriminant, frame-value index, and saturation index. The weight `termWeight` includes the powers $-(k+1), -k, -k$ of the three corresponding invariants (with a count normalization depending on the codimension).

The point of these inverse powers is that the total weight can be bounded by estimates for sublattices rather than a direct count of pairs. The geometry-of-numbers input (*Formal-source input*: `Filling.tsum_sublatticeDisc_inv_le` in `NearTerms.lean`) bounds sums of inverse discriminant powers using short generators obtained by a relative Siegel lemma. Bounds on the congruence indices and lattice intersections followed by these sums give the explicit theorem `sum_pairWeight_le` in `BlockBound.lean`: the total pair weight is at most $2\tau^2$. The class amplitude size prior to summation is the function `classSize` of `SchurTest.lean`. With M_0 the normalization `countWeight0` and z the relevant index parameter it has the exact form

$$a_j(z) = \tau^2\left(\frac{D_j}{M_0 g^{2\min(h,j)kf} z^{k+1}}\right)^{1/2}, \quad D_j = \begin{cases} \tau^{13-32j}D_1 & j \le h, \\ D & j = h+1. \end{cases}$$

At the last level the estimate for $a_j(z)$ is obtained from (20.7) and the archimedean and finite discriminants. At earlier levels the weighted Schur estimate bounds the retained common-vector blocks. The exact comparisons with class sizes and the total weights are `IsNear.le_classSize_and_cellMass_le` in `SchurTest.lean`, and `enorm_amplitudeSum_le_classSize` and `enorm_amplitudeSum_le_classSize_top` in `BlockBound.lean`. We take these as formal-source inputs. These estimates are uniform in the rank and in the field; qualitative finiteness of the double cosets or a fixed-rank Siegel lemma estimate alone would not give the stated bound.

Proposition 20.11 (Formal-source input: reduction to coherent top-cell pairs)**.** *Retain the notation D, D_1, τ. Suppose T is stable, Ω is a set of reductions with the common fixed-vector norm bound $X \ge 1$ of `Filling.FixesNormAtLeast`, and $C_0 \ge D_1$ is a constant satisfying the following extra restrictions. If a term in T over Ω is in normal form zero, require $C_0 \ge \tau^{20}D_1$; if an omitted link exists, require $C_0 \ge \tau^{12}D/X^k$. For $w \ge 0$, suppose*

$$\mathcal{E}_{\mathrm{link}}(T\cap\{\mathfrak{d}\neq 0\},\Omega,\mathcal{L}) \le w\nu^2.$$

Then

$$\mathcal{E}(A_T;\Omega) \le (1+b^{-4})^2 w + b^{-12}C_0. \tag{20.18}$$

This is `Filling.energy_le_of_linkEnergy_le` in `New/Filling/CrudeEnergy.lean`; the two subordinate assertions are `energy_le_of_level_pairs` and `energy_le_of_level_lt`. Their quantitative errors are $C_0/(b^{30}4^{h+1-j})$ at level j and the accumulated error below level n' is at most $C_0/(b^{30}4^{h+1-n'})$. To explain the accumulation, split off the next level by (20.16) (taking $\delta = 1$ here); its own error and the preceding one together cost at most four times the preceding-level budget. Thus a geometric factor 4 suffices from one level to the next. If a term has discriminant zero, the character obstruction or a lower normal form just discussed applies, so it contributes only to this error. At the last level, separate the coherent equal-link blocks, of cost $w\nu^2$, from the unequal-link blocks estimated by the common link and its codimension. Returning to the full amplitudes with (20.16), now using $\delta = b^{-4}$ and the extremely small truncation error, gives (20.18). Both the precise coefficient of w and the estimate on the omitted terms matter: choosing a fixed loss at the final step instead of $1 + b^{-4}$ would not yield a nearly spanning family. The nontrivial block estimates above, rather than their algebraic assembly, are the imported part of this proposition.

A first consequence concerns translations that locally lie in the image of the Bessel subgroup. For $i = 0, 1$, set

$$\Omega_i = \{s : \text{there exists } t \in \bar{S}_B \text{ with } s_i = t_i\}.$$

Proposition 20.12 (Formal-source input: Bessel-coset error)**.** *For each i one has*

$$\mathcal{E}(A_{G(E^+)\backslash S_B}; \Omega_i) \le b^{-12} D_1.$$

This is `energy_compl_besselS_le` in `New/Filling/Moments.lean`. For these reductions the frames satisfy the exact Bessel constraints on an entire prime's chosen factors. The finite congruences and the character condition imply that a remaining normal-form-zero term would actually be in the Bessel subgroup, so such a term does not contribute to the complement. The common constraints supply the norm bound X for applying Proposition 20.11 with $\mathcal{L} = \varnothing$, $w = 0$, $C_0 = D_1$; the scale inequality (20.15) guarantees the requisite domination of the full-cell error. The exact group-theoretic constraint and norm calculation are part of the cited source proposition. This restrictive error estimate isolates the diagonal term and is used again to bound the non-Bessel part of a central projection.

20.7 The sharp coherent main term

The preceding estimates deliberately allow coarse constants, because they are used only for errors. For the coherent open-cell pairs a different argument must retain enough positivity to give the leading constant one. We record the form of this argument and isolate its three substantive inputs: local orbital kernels, rational-to-local block comparison, and the global sieve.

Kernel systems and their products. Consider a measured commutative ring A with an additive character $\psi : A \to \mathbb{C}^\times$ of modulus one, and let $A^\times$ be its group of units. For a general semilocal ring put $u(z) = z$ if z is a unit and $u(z) = 1$ otherwise. Let p, q denote sequences of units; $p[i : c]$ means the sequence obtained by replacing entry i by $u(c)$. A sieve system is a family of forms $J_i(p, q)$, $0 \le i \le h + 1$, with the following relations:

$$\int_A J_{i+1}(p[i : c], q[i : c])\, dc = J_i(p, q), \qquad |J_0(p, q)| \le \mathfrak{e},$$

$$\left| \int_A \psi(\xi c) J_{i+1}(p[i : c], q[i : c])\, dc \right| \le \beta(\xi)\Lambda_i(p)\Lambda_i(q). \tag{20.19}$$

The Fourier inequality is required for unit frequencies, and all the integrals in these relations are absolutely convergent. For $i < h$ one also requires the unequal-entry bound

$$|J_{i+1}(p,q)| \le \beta(\iota(p_i) - \iota(q_i))\Lambda_{i+1}(p)\Lambda_{i+1}(q)$$

when the difference of inverse-head coordinates in this expression is a unit; the map ι is inverse of a unit, with the additional factor $1/g$ at the finite places. At the top the kernel is continuous and every matrix $\big(J_{h+1}(p_a,p_b)\big)_{a,b}$ with common last entry is positive semidefinite. This is an abstraction of `SieveSystem` in `New/Filling/EichlerFlag.lean`; we have omitted the infinite tails of the sequences from the notation.

Here is a useful consequence. If p_a is any finite family of heads, for $i \le h$ define the function

$$F_i(c) = \sum_{a,b} J_{i+1}(p_a[i:c], p_b[i:c]).$$

This is nonnegative, because the top forms are positive semidefinite and lower forms are obtained by integrating a common coordinate, successively. It is lower semicontinuous wherever the head maps are continuous, by Fatou's lemma after integration over subsequent entries. Fubini and (20.19) imply

$$\int F_i = \sum_{a,b} J_i(p_a,p_b), \qquad |\widehat{F}_i(\xi)| \le \beta(\xi)\left(\sum_a \Lambda_i(p_a)\right)^2.$$

The same scheme applies to the product of systems at finitely many places; the top positive matrices multiply entrywise and remain positive semidefinite by the Schur product theorem, the integration compatibility follows from Fubini, and the constants and majorants multiply. This is the elementary content of `SieveSystem.pi`, `SieveSystem.step_sum`, and `SieveSystem.nonneg_and_lowerSemicontinuous`. Positivity is retained in the sum over both heads; the individual summands need not be nonnegative, and no absolute value may be inserted before the sum without changing the argument.

Finite orbital kernels. Let Σ be a set of chosen places above g_0 and fix, for each $v \in \Sigma$, a nonzero frame vector at the place or its conjugate. Define $\Omega^\circ(\Sigma)$ to consist of the level reductions fixing the specified vectors, and for which, unless Σ is the full set, there exists at least one selected place above g_0 at which all required leading frame minors are units. This is the set denoted `fixingSet` in the formal proof. This is the exact source condition; imposing simultaneous unit-minor tests at all places gives a smaller subset and is sufficient for the later positivity lower bound. At an arbitrary finite conjugation-stable set of places containing the level primes and all denominators of a finite initial selection of rational terms, the argument works over the product of the corresponding local fields. Choose an intermediate compact open subgroup between $K(g)$ and K_f whose local factor at each place contains the appropriate fixed-vector subgroup. The ratio of the bottom orbital integral to the group measure is diminished by a factor b^{-6k} at each specified place. The precise construction is `exists_intermediateGroup_one_prime` and the product version `exists_intermediateGroup_lines` in `New/Filling/FixedLines.lean`; it uses counting on split matrices and a bound on the volume of nonunits (at all the level places this is at most b^{-9}).

The finite system incorporates the Bessel character of conductor given by $1/g$ and the different. If $p = (p_i)$ is a head over the local product of fields, let $[D_j]$ denote the generalized ring index associated to the partial product $D_j = p_0 \cdots p_j$. Write $\mathrm{den}(\xi)$ for its denominator index. The finite head majorant is

$$\Lambda_i^f(p) = \prod_{j<i} [D_j]^{s_0^6/2} \quad \text{if all } D_j \text{ are integral, and 0 otherwise.}$$

The finite frequency majorant is

$$\beta_f(\xi) = \frac{[\xi]^{(k-s_0^7)/2}}{\mathrm{den}(\xi)^6} b^{s_0^9 f},$$

where $[\xi]$ is the generalized index, not necessarily an integer. These definitions are `finHeadWeight` and `frequencyGain` in `EichlerFlag.lean` and `GlobalSieve.lean`. After dividing the top orbital kernel by the volume of a compact open additive neighborhood of zero and by the prescribed mass ratio, the bottom constant of the finite sieve system is at most

$$(1 + b^{-6}) b^{-6k|\Sigma|}. \tag{20.20}$$

These are the conclusions of the *Formal-source input* `Filling.IsFillingLevel.exists_finite_sieveSystem` in `New/Filling/FiniteSystem.lean`. We emphasize that the normalization of the orbital kernel and the bottom constant include the measure of the fixed-vector subgroup, rather than only a bound on its order. The proof passes down the tower of groups fixing successive hyperbolic pairs, integrating one entry of the frame Gram matrix at each step. Its key step is a local integral comparison (`Level.integral_succ_floor_eq` and `Gauge.integral_floorForm_succ`, `New/RealPlace/LocalTheorem.lean`). In the nonzero-frequency bound one uses the local character along a simple-root direction of an Eichler transformation, and averaging gives a power of the inverse module of the frequency. A parallel unequal-head calculation uses frequency $(p_i^{-1} - q_i^{-1})/g$; this is the origin of the unequal-entry bound in the definition of a sieve system. These local comparisons are substantive statements for indefinite unitary groups of growing rank, not applications of Plancherel alone.

The archimedean and finite systems together. At the infinite places use the systems of Section 20.4, and at the finite ones the systems of (20.20). The diagonal embedding of the rational frame entry into the local products scales it at each v by l_{vi}. Its covolume factor in Poisson summation is

$$c_i = \frac{\mathrm{vol}(\widehat{\mathcal{O}}_E)}{\prod_{v\in\Phi} l_{vi}^2} \qquad (0 \le i \le h).$$

The square in the denominator is the real Jacobian of multiplication by l_{vi} on a complex coordinate. Let A_0 be a finite set of rational units integral away from the auxiliary finite set, along with their inverses, and let $V_0 \subseteq A_0$ index the last rational entry. With the product system write $p[t]$ for the $h+1$ entries obtained from $p \in A_0^h$ and $t \in V_0$, read diagonally in the local coordinates. The following inequality is the output of the global sieve.

Proposition 20.13 (Formal-source input: the global sieve)**.** *If a finite sieve system has bottom bound $\mathfrak{e}_f$, head and frequency weights as above, then*

$$\Re \sum_{t\in V_0} \sum_{p,q\in A_0^h} J_{h+1}^f(p[t], q[t]) \prod_{v\in\Phi} J_{h+1}^v(p[t], q[t]) \le \prod_{i=0}^{h} c_i \left(\mathfrak{e}_f \prod_{v\in\Phi} \frac{1 + b^{-10}}{d_v} + \frac{b^{-6kf-10}}{d_\infty} \right).$$

The exact theorem is `sieve` in `New/Filling/GlobalSieve.lean`. The proof uses `one_sided_poisson` in that file; we give its mechanism because it explains the small additive error. For the function F_i above, one-sided Poisson summation on the additive adele lattice gives a bound for the sum over a finite set of rational points by c_i times its zero-frequency integral plus c_i times an error estimated by the sum of its nonzero frequency coefficients. A one-sided version is needed for lower semicontinuous kernels; it follows by comparison with integrable smoothing approximants to a nonnegative function and the elementary finite-subset inequality. The Fourier bounds and the assumption $(\sum_p \Lambda_i(p))^2 \le \Gamma \prod_{a<i} c_a$ give an error at

most $\Gamma\varepsilon\prod_{a\le i}c_a$ at this step if the sum of nonzero-frequency majorants is at most ε. When undoing a common head coordinate to descend another floor, the unequal head contributions give a second error of this size, bounded using the off-diagonal condition. Consequently induction from $i=0$ gives a bound

$$\prod_{a\le i}c_a\bigl(\mathfrak{e}+(2i+1)\Gamma\varepsilon\bigr).$$

The source's `one_sided_poisson` states this explicitly, with rational frequencies that are integral away from the finite set. Products of the weights β_f and β_v are summable uniformly in the finite set: by the product formula the large powers of the norms cancel, leaving the damping power 6 and denominator factors. The needed elementary sum with exponent k' is the precise inequality `sum_frequencies_le`: over a finite set of nonzero T-integers y, if $|c_v|^{-1}\le|x_v|\le B_1$, $B_1\ge1$, then

$$\sum_y\frac{[y]_T^{k'/2}}{\mathrm{den}(y)^6}\prod_v\lambda_{k'}(|c_vv(y)|^{-1})\le(b\,\mathrm{rd}_E)^{3f}\prod_v\bigl(|x_v|^{k'}(bB_1^2)^3\bigr).$$

This is proved in the source using a sum of inverse discriminants of rank-one sublattices of a rank-two lattice (the vector $(1,y)$), including its denominator lattice. It is a formal-source input, together with the one-sided Poisson lemma in the stated level normalization. Taking $k'=k-s_0^7$ and the actual archimedean scaling the strong inequalities of `Filling.IsAdmissible` give $(2h+1)\Gamma\varepsilon\le b^{-6kf-10}/d_\infty$. This is substantially better than a sieve bound with a factor b^η, and is the reason the second moment has the requisite leading constant.

Comparison of blocks with the orbital kernel. On the nondegenerate cell the terms are put in Weyl normal form $w_p\sigma$ with common residual middle element σ, common top entry $t=\langle z,\sigma z\rangle$, and $|v(t)|>N$ at every infinite place. The terms supported on identical finite reductions and identical link are sorted into such packets. Finite sets of initial representatives suffice by the absolute convergence of the amplitudes; subsequently enlarge the finite set of places to contain all denominators and take limits of local compact-open approximations. The exact comparison is `sum_blocks_le_topKernel` in `New/Filling/FiniteSystem.lean`; here is its normalization. Let $\mathcal{U}=|\bar U|=g^{fh(2k+2h+1)}$ be the order of the reduced integral unipotent subgroup, let

$$c_*=(\pi^k/((k-1)!N^k))^f\,D_E^{1/4}\,v_f^k,\qquad A_*=|\bar S_B|\mathcal{U}|\mathcal{C}|c_*.$$

For a finite packet of distinct quadruples (u,u',s,σ) with the given t, the squared norm of the corresponding sum of coset amplitudes, multiplied by the common normalization, is bounded by

$$(1+b^{-6})A_*\,\Re\sum_{p,q}R_{pq}\frac{J^{\mathrm{top}}(p,q)}{\mathrm{vol}(L_0)\ \mathrm{massRatio}},$$

for sufficiently small compact open additive L_0. Here R is the Gram matrix of the product of normalized archimedean correlation tensors. The exact factors in R are those of (20.5), and the finite module factors in the packet comparison cancel against these factors. This is a *Formal-source input*: the proof uses a count of integral points on a quadric for the last coordinate (`siegel_sum_le_lintegral_quadric`), and a local block comparison with the positive phase kernel, then the global-to-local limit identity `exists_hasSum_cosetAmplitude` in `New/Filling/GlobalToLocal.lean`. The normalization A_* incorporates the unipotent order and the same-class condition. It should not be replaced by a crude number of rational points. The factor c_* is the constant of the rank-k definite sphere sum. The factor $(|v(t)|^2-N^2)^{k-1}$ in

this count is the one that cancels against the open-cell Jacobian in the normalized real-place correlations.

The size comparison used in the compact proof is the following hypothesis of the filling datum (with our notation):

$$A_* \le (1+b^{-9})D\, d_\infty c_G^f\, v_f^{-s_0} \left(\frac{\nu\mathcal{L}}{|I|}\right)^2.$$

This formula is `packetConst_mul_mass_le` in `Filling.IsAdmissible`. It accounts for the fact that the second moment is compared with a definite genus mass, not just a count of the rows in the particular target representation. Combining the block comparison, Proposition 20.13, the scale factors, and the crude stratification bound yields the following sharp estimate. In contrast with a fixed-group spectral equidistribution statement, the constants here are uniform for the auxiliary r, k, f of this proof.

Proposition 20.14 (Formal-source input: sharp energy on the open congruence set)**.** *For every set Σ of chosen places above g_0, with specified nonzero fixed vectors, one has*

$$\mathcal{E}(A_{\mathrm{all}};\Omega^\circ(\Sigma)) \le (1-b^{-4})(1+9b^{-4})D\, b^{-6k|\Sigma|}.$$

This is `energy_fixingSet_le` in `New/Filling/Moments.lean`. To obtain exactly this factor the coherent bound is taken with slack $(1+b^{-4})Db^{-6k|\Sigma|}$ before applying the two small error steps of Proposition 20.11. The resulting expression is at most $(1-b^{-4})(1+9b^{-4})Db^{-6k|\Sigma|}$ by the size bounds and $\tau^{20}D_1 \le Db^{-6kf}$, which follows from (20.15). The main analytical content of this proposition is precisely the uniform packet and local sieve inputs specified above. We have not replaced those inputs by a claimed classical trace formula theorem without quantitative estimates.

20.8 A finite-field positivity argument for singular minors

The open-set estimate of Proposition 20.14 does not yet imply a total energy estimate, since it leaves out the locus of singular minors. The next finite group argument shows that a nonnegative function of positive type cannot concentrate too heavily on such a locus unless it concentrates on fixed-vector subgroups. The distinction between the energy itself and a general positive-type function is useful because multiplying by a fixed-vector indicator preserves positive type, allowing induction.

Let G be a finite group, $\pi : G \to \mathrm{GL}_R(\mathbb{F}_q)$ a representation, and let A, A' be $R \times j$ matrices. Let $\Xi : G \to \mathbb{R}_{\ge 0}$ be of positive type; choose vectors u_g in a complex Hilbert space with $\langle u_a, u_b\rangle = \Xi(a^{-1}b)$ (the Gram representation follows directly from positive semidefiniteness). Write $S_\Xi = \sum_g \Xi(g)$. For any nonzero $x, y \in \mathbb{F}_q^j$, suppose

$$\sum_{g:,\pi(g)Ax=Ax} \Xi(g) \le Y, \qquad \sum_{g:,\pi(g)^{-T}A'y=A'y} \Xi(g) \le Y.$$

Lemma 20.15 (Finite-field determinant estimate)**.** *Under these assumptions,*

$$q \sum_{g:,\det(A'^T\pi(g)A)=0} \Xi(g) \le 2S_\Xi + q^{j+R/2}Y.$$

Proof. Fix a nontrivial additive character ψ of $\mathbb{F}_q$. For nonzero y, x consider the weighted exponential sum

$$S(y,x) = \sum_g \psi(y^T A'^T \pi(g) A x)\Xi(g).$$

In $\ell^2(\mathbb{F}_q^R)$ let e_z be the delta basis and f_z its normalized Fourier basis with entries $q^{-R/2}\psi(z^T z')$. These are orthonormal bases by additive character orthogonality. For the tensors $U_1 = \sum_a f_{\pi(a)^{-T}A'y}\otimes u_a$ and $U_2 = \sum_b e_{\pi(b)Ax}\otimes u_b$, the squared norms are at most $|G|Y$: on expanding the inner products of the tensor sums, the basis inner products give exactly the indicators of the associated fixed-vector subgroups. Their mixed inner product is $|G|q^{-R/2}S(y,x)$, or its conjugate according to the Fourier convention, by reindexing $a^{-1}b$. Cauchy–Schwarz thus gives $|S(y,x)| \le q^{R/2}Y$. If at least one of y, x vanishes then $S(y,x) = S_\Xi$.

Put $B_g = A'^T\pi(g)A$. Additive orthogonality gives $\sum_y q^{-j}\psi(y^T B_g x) = 1_{B_g x=0}$. Summing in x and weighting by $\Xi(g)$ shows

$$\sum_g \Xi(g)|\ker B_g| = q^{-j}\sum_{y,x} S(y,x) \le 2S_\Xi + q^{j+R/2}Y.$$

On the singular locus, $|\ker B_g| \ge q$, and all summands are nonnegative. This proves the inequality. It reproduces `card_mul_sum_det_le` in `New/Heights/OneDeterminant.lean`. □

In the application $R = r$ and j is at most the size s_0 of the frame. A vector fixer indicator is of positive type, because it is the matrix coefficient of the permutation representation on the orbit of that vector. Multiplying it by Ξ gives another nonnegative function of positive type, by taking a tensor product of Gram representations. Thus one can successively impose new fixed-vector conditions. This is exactly the feature needed to derive the following result from Proposition 20.14.

Proposition 20.16 (Removing the unit-minor condition)**.** *Let $\Xi \ge 0$ be of positive type on $\mathfrak{G}$ and satisfy*

$$\sum_{s\in\Omega}\Xi(s) \le \mathcal{E}(A_{\mathrm{all}};\Omega)$$

for every set Ω of translations. Suppose its support consists of reductions fixing specified nonzero frame vectors over every place in a set Σ above g_0. Then

$$\sum_{s\in\mathfrak{G}}\Xi(s) \le (1+9b^{-4})D\,b^{-6k|\Sigma|}.$$

Proof. Use descending induction in $|\Sigma|$. If all places above g_0 belong to Σ, there is no additional minor condition, and the assertion follows from Proposition 20.14. In general suppose the assertion fails for Ξ at Σ. At each new place v outside Σ, intersect its support with the prescribed fixer of any nonzero frame vector x in the appropriate standard or dual frame. The restricted function remains of positive type, is dominated by the energy, and is supported on the extended set of fixers. By the induction hypothesis its total mass is at most $(1+9b^{-4})Db^{-6k(|\Sigma|+1)} < b^{-6k}S_\Xi$. Lemma 20.15 bounds the mass of each singular-minor locus by

$$\frac{2+q^{j+r/2}b^{-6k}}{q}S_\Xi \le 3b^{-10}S_\Xi,$$

where $q = g_0$, and the last inequality follows from $q \le b^{11}$, $q \ge b^{10}$, and $j + r/2 \le k/2 + 2h + 2$, which with the scale bounds gives $11(j+r/2) < 6k$ (in particular $11(2h+2) < k/2$). The exact singular-minor test of `fixingSet` involves the standard and contragredient frame matrices, with one test for each of the leading minors at a chosen place; this identification is the finite matrix calculation in `sum_le_of_positive_type` of `New/Filling/Moments.lean`. Even bounding the number of tests by $2f(h+1)$, the excluded mass is at most $6f(h+1)b^{-10}S_\Xi \le b^{-4}S_\Xi$, by $f \le b/1000$, $h+1 \le b$. Consequently at least $(1-b^{-4})S_\Xi$ lies in $\Omega^\circ(\Sigma)$. Domination by energy and Proposition 20.14 imply $(1-b^{-4})S_\Xi \le (1-b^{-4})(1+9b^{-4})Db^{-6k|\Sigma|}$, contradicting the supposition. This proves the claimed induction. □

Here we have used only the imported sharp estimate on the open set and finite group positivity; we have not assumed that the geometric energy is automatically of positive type. In fact the function

$$\Xi(s) = \|P^*\rho(s)P\|_{\mathrm{HS}}^2$$

is of positive type. To see this, with $T_0 = PP^*$ put $u_s = \rho(s)T_0\rho(s)^*$ in the Hilbert space of Hilbert–Schmidt operators. Then $\langle u_a, u_b\rangle = \mathrm{Tr}(T_0\rho(a^{-1}b)T_0\rho(b^{-1}a)) = \Xi(a^{-1}b)$. It equals the energy per reduction by Lemma 20.9. With $\Sigma = \varnothing$ Proposition 20.16 therefore gives

$$\frac{1}{|\mathfrak{G}|}\sum_s \|P^*\rho(s)P\|_{\mathrm{HS}}^2 \le (1+9b^{-4})\frac{c^2}{n_0}. \tag{20.21}$$

This is where both local and global sharpness are indispensable.

20.9 First moment and bad finite-type loss

The estimates needed in addition to (20.21) are not general consequences of that second moment. They use the diagonal and central projections at the two different level primes. In fact

$$\sum_{i=1}^{c}\|(P^*P-I)e_i\| \le b^{-4}c, \qquad \|(I-R)P\|_{\mathrm{HS}}^2 \le 4b^{-4}c. \tag{20.22}$$

We explain the derivation and specify the further inputs.

For a central self-adjoint idempotent $z \in \mathbb{C}[\mathfrak{G}_i]$ and the inclusion $\jmath_i : \mathfrak{G}_i \to \mathfrak{G}$, let $M_z(s)$ be the $c \times c$ matrix with block from u to u' equal to

$$\nu^{-1}\sum_{a\in\mathfrak{G}_i} z(a)A_{G(E^+)\setminus S_B}(u, \jmath_i(a)^{-1}s, u').$$

The complement-coset energy controls this matrix at 1 after summing its row norms. Indeed convolution by the coefficients of z is an orthogonal projection on the regular representation and hence contracts the squared sum over s. To restrict to the relevant set Ω_{1-i}, note that it is invariant under left translation by the embedded factor $\mathfrak{G}_i$: the other component is unchanged. It contains the Bessel reductions. The convolution estimate and the covariance under the Bessel subgroup give

$$|\bar{S}_B|\|M_z(1)\|_{\mathrm{HS}}^2 = \sum_{s\in\bar{S}_B}\|M_z(s)\|_{\mathrm{HS}}^2 \le \mathcal{E}(A_{G(E^+)\setminus S_B}; \Omega_{1-i}).$$

The equality uses the following exact covariance calculation, a *Formal-source input* in the proof of `sum_norm_amplitudeSum_le` in `New/Filling/Moments.lean`. Each $s \in \bar{S}_B$ induces a permutation of the column classes by the rational Bessel element over (u, s, u'). The left and right covariance of the Bessel functions replaces the corresponding block by a unitary matrix acting on its coordinate space, times a scalar of modulus one. Norms are unchanged because the finite-dimensional compact types are unitary. This is derived from `amplitudeSum_move` in `New/Filling/AmplitudeSums.lean` and `exists_besselS_mem_termsOver` in `NearTerms.lean`. The convolution estimate is applied after keeping the terms whose factor-$(1-i)$ reduction lies in the image of the Bessel subgroup; the exact group-module contraction, together with this covariance, is the assertion `sum_norm_amplitudeSum_le` cited above. In particular the resulting sum of row norms is at most $b^{-4}c$ (using $\sqrt{c}\|M_z(1)\|_{\mathrm{HS}}$ and Proposition 20.12). Taking $z = 1$ and using the identity term gives the first inequality of (20.22); bounding row or column norms is equivalent for the self-adjoint $P^*P - I$.

The remaining statement is the trace of a small central projection on the Bessel-subgroup packet. If z is central and $\Re(z(1)) \le b^{-8}/2$, then

$$\Re \sum_{a\in\mathfrak{G}_i} z(a) \sum_{u,j} A_{S_B}(u, \jmath_i(a)^{-1}, u)_{j,j}/\nu \le b^{-4}c. \tag{20.23}$$

This is the *Formal-source input* `re_trace_besselS_le` in `Moments.lean`. It is not a trace formula on arbitrary cusp forms. Its proof realizes the class sum on the finite orbital integrals over the Bessel subgroup, using invariance of z under conjugation and the local character. The *ell*^{2} norm of the coefficients of z is exactly $z(1)$ since $z = z^* = z^2$ in the regular group algebra. The remaining orbital coefficient has squared norm at most $2c^2$ after the measure normalizations (the lower-floor estimate in `New/Filling/FixedLines.lean`); hence Cauchy–Schwarz bounds the trace by $c\sqrt{2z(1)} \le b^{-4}c$. The crucial nontrivial part is the bound on the orbital coefficient, incorporated in (20.23). Applying this with $z = 1 - e_i$ and Lemma 20.2, the trace of $(1-\rho(e_i))PP^*$ is at most $2b^{-4}c$, since the non-Bessel part contributes at most the same bound $b^{-4}c$ from the previous paragraph. The two projections commute, so $1 - R \le (1-\rho(e_0)) + (1-\rho(e_1))$ as positive operators. Taking their traces against PP^* proves the second inequality in (20.22).

The notation of the formal proof for these two inputs is deliberately separate: `re_trace_besselS_le` controls the Bessel term, whereas `sum_norm_amplitudeSum_le` controls the complement for any central projection, including the identity. Both are needed even though the discarded relative Plancherel mass itself is tiny.

20.10 From the moment bounds to the covering theorem

We now give the final argument without any further automorphic estimate; it is a general Hilbert-space inequality. First, W is nonzero. If $R = 0$ then $\|P\|_{\mathrm{HS}}^2 \le 4b^{-4}c$ by (20.22), but the first bound in (20.22) gives $\mathrm{Tr}(P^*P) \ge (1-b^{-4})c$. These two inequalities contradict $c > 0$ and $b \ge 1000$. Thus $n > 0$.

For $s \in \mathfrak{G}$ put $T_s = \rho(s)PP^*\rho(s)^*$ and $\overline{T} = |\mathfrak{G}|^{-1}\sum_s T_s$. Unitarity and the trace identity show

$$\begin{aligned}\|\overline{T}\|_{\mathrm{HS}}^2 &= |\mathfrak{G}|^{-2}\sum_{s,t}\mathrm{Tr}(T_sT_t)\\ &= |\mathfrak{G}|^{-1}\sum_s \|P^*\rho(s)P\|_{\mathrm{HS}}^2.\end{aligned}$$

If $J \subseteq W$ and p_J is its orthogonal projection in V, set $U_s = (R\rho(s)P)^{-1}J$. For each unit vector $u \in U_s$, the two orthogonal components of $\rho(s)Pu$ lie respectively in J and $\ker R$. Summing over an orthonormal basis of U_s and enlarging the sums for these components gives

$$\mathrm{Tr}(p_{U_s}P^*P) \le \mathrm{Tr}(p_JT_s) + \|(I-R)P\|_{\mathrm{HS}}^2.$$

In this computation p_{U_s} is the orthogonal projection within C and R commutes with $\rho(s)$. For any orthogonal projection p_U in C, one has

$$|\,\mathrm{Tr}(p_U(P^*P - I))| \le \sum_i \|(P^*P-I)e_i\|,$$

by expanding in the coordinate basis, applying Cauchy–Schwarz to each inner product, and using $\|p_Ue_i\| \le 1$. Applying this with $U = U_s$ gives

$$\dim U_s \le 5b^{-4}c + \mathrm{Tr}(p_JT_s).$$

Averaging and using Hilbert–Schmidt Cauchy–Schwarz, $\|p_J\|_{\mathrm{HS}} = \sqrt{\dim J}$, and (20.21), we obtain

$$\begin{aligned}\frac{1}{|\mathfrak{G}|}\sum_s \dim U_s &\le 5b^{-4}c + \sqrt{\dim J}\sqrt{(1+9b^{-4})c^2/n_0}\\ &\le \left(5b^{-4} + \sqrt{(1+9b^{-4})\alpha}\right)c\\ &\le \tfrac{1}{2}(1+\alpha+19b^{-4})c,\end{aligned}$$

where $\alpha = \dim J/n$. The second line uses $n \le n_0$, and the third uses $2\sqrt{uv} \le u+v$. Thus there exists s attaining the indicated bound. This is `Fits.exists_translate_finrank_comap_le` in `New/Filling/Moments.lean`, with the underlying Hilbert–Schmidt variance argument `finrank_sub_le_re_inner_dyadSum` in `New/Heights/OneDeterminant.lean`. The constant 19 comes from the projected and identity estimates ($2 \cdot 5$) and from the second moment (9).

Finally, ramify the columns by applying $D_N = \prod_{w|\pi}(1 - z_w)$, with z_w the generator of the finite central-unit action at a selected place above N. Its image is C^{ram}, since in a cyclic finite group the image of $1 - z_w$ is the kernel of the average over the group, and these operators at different places commute. In the source the estimate `ramifyingOp_finrank_comap_and_short` in `New/Columns/ChainValuation.lean` shows that for any subspace $U \subset C$ the dimension of its inverse image under D_N is at most $\dim U + fc/(N-1)$: the central-unit action permutes classes freely in cycles of $N-1$, so the kernel has relative dimension at most $1/(N-1)$ at each place. The freeness and compatibility with the column type are the formal-source input used here; the dimension inequality also follows immediately from rank-nullity for D_N from its kernel bound. Admissibility gives $19b^{-4}/2 + f/(N-1) \le b^{-2}$, since $N - 1 \ge b^9$ and $f \le b/1000$. Inserting this loss and removing the common nonzero normalization scalar proves (20.1). This completes the derivation of the spread property appearing in `Data.ramifiedColumns_fill_rows` in `New/Filling/FillingTheorem.lean`.

It remains to convert spread into a small number of translates. Put $\eta = b^{-2}$ and start with $J = 0$. If the current codimension $q_* = n - \dim J$ satisfies $q_* > 4\eta n$, choose s by (20.1). By rank-nullity applied to the quotient map from C to W/J, the increase Δ in dimension obtained by adjoining $P_s(C^{\mathrm{ram}})$ satisfies

$$\Delta \ge c\left(\frac{q_*}{2n} - \eta\right) \ge \frac{cq_*}{4n} > 0.$$

In particular this is a genuinely new translate. Define a potential on the remaining codimension by

$$\Psi(q_*) = 5n \log\left(1 + \frac{q_*}{\eta n}\right).$$

When $q_* > 4\eta n$, adjoining the new image decreases the potential by at least

$$\Psi(q_*) - \Psi(q_* - \Delta) = -5n \log\left(1 - \frac{\Delta}{q_* + \eta n}\right) \ge \frac{5n\Delta}{q_* + \eta n} \ge c,$$

since $q_* + \eta n \le 5q_*/4$. The process terminates because dimension increases by a positive integer at every step. The potential remains nonnegative; hence the number of selected translates times c is at most $\Psi(n) = 5n\log(1+b^2)$. The final codimension is at most $4b^{-2}n$. This proves Theorem 20.1. The greedy covering is the result `exists_few_translates` of `New/Heights/OneDeterminant.lean` (the preceding derivation is a complete mathematical proof of that deduction).

For the application, take a spanning set of short integral columns as in Theorem A, and for the above finite set T select independent vectors from its projected images after the rational

rescaling. The span has the dimension supplied by Theorem 20.1, and the accumulated logarithmic index bound is at most $5n\log(1+b^2)B$. Apply the individual squared-length estimates to these independent vectors and Hadamard's inequality to their lifted Gram matrix. The loss from the small missing fraction of dimensions is controlled in the final comparison by the crude bound for the heights. No further spanning theorem is needed. The substantive assumptions for B have thus been reduced from the theorem itself to the precise uniform Bessel system, lattice block, trace transfer and local orbital/Poisson estimates; the finite-field counting, second-moment positivity, variance, and greedy selection are explicit deductions from those estimates.

Chapter 21

The analytic and arithmetic mechanism of the Bessel sieve

In Chapter 20 we isolated the quantitative covering property of the projected Poincaré map and its second-moment proof. We now give a more detailed account of the estimates behind that property. It is useful to organize the material by the three changes of measure that occur. First one goes from Haar measure on a real unitary group to a Bessel quotient. Next one introduces a tower of measures on successive open cells, replacing one pivot coordinate at a time by a Fourier integral. Finally one compares the rational orbit count with the finite-place measure and uses Poisson summation. The last two steps must preserve the leading constant. A conventional argument that merely bounded the integrals up to a factor depending on the ranks or the fields would not produce a sufficiently large subspace of rows.

We retain the notation $h = m/2 - 1$, $s_0 = h + 1$, $r = k + 2h + 1$ of Section 20.3. The letter d_v is a real formal degree (not the quadratic prime d). Unless the measure is explicitly specified otherwise, matrix norms in this section are operator norms. We use $\|\cdot\|_{\mathrm{HS}}$ for the Hilbert–Schmidt norm and write J_* for a Hermitian matrix with $J_*^2 = 1$. Thus $U(J_*)$ is embedded in $\mathrm{GL}_r(\mathbb{C})$ and

$$g^{-1} = J_* g^* J_*, \qquad \|g^{-1}\| = \|g\|, \qquad \|g\| \geq 1.$$

The symbols $\mathbb{A}$ and $\mathcal{O}$ refer to the adeles and integers of the CM field E. When using an assertion with "formal-source input" below, we mean the precise result at the indicated Lean declaration. In several instances we derive its consequences, or a substantial part of its proof, but the remaining uniform coordinate estimate is explicitly retained as an input. The purpose is to separate those inputs from elementary arguments of harmonic analysis and linear algebra, and to make their quantitative roles intelligible; we do not claim to have independently established every assertion.

21.1 Norm balls in the local groups

A useful feature of coefficient estimates on $U(J_*)$ is that a fixed inverse power of the operator norm is sufficient to dominate all integrals on the Bessel subgroup, and the exponent required depends only polynomially on the rank. We give the argument because it is repeatedly used both to justify rearrangement of integrals and to prevent the normalization of a Bessel function from destroying the tail bound.

Lemma 21.1 (Norm-ball covering)**.** *Let S be a locally compact group, $j : S \longrightarrow U(J_*)$ a continuous homomorphism, and ν_S a right Haar measure on S. Suppose that $D, R \geq 1$,*

$y \in U(J_*)$, and $K \geq 3DR^2$. Then

$$\nu_S\{s : \|j(s)y\| \leq R\} \leq K^{2r^2}\nu_S\{s : \|j(s) - 1\| \leq D^{-1}\}.$$

If j is proper, the set on the right has finite measure; if the Haar measure is nonzero it has positive measure.

Proof. In a real normed vector space of dimension $2r^2$, a set of points in the radius-R ball with mutual distances at least $\delta > 0$ has cardinality at most $(1 + 2R/\delta)^{2r^2}$. Indeed disjoint open balls of radius $\delta/2$ about them lie in the radius-$R + \delta/2$ ball; comparison of Lebesgue volumes proves the bound. Take $\delta = (DR)^{-1}$ and choose a maximal such separated family among the matrices $j(s)y$ with norm at most R. There are at most $(1 + 2DR^2)^{2r^2} \leq K^{2r^2}$ of them, with lifts $s_i \in S$. Every s in the set under consideration satisfies $\|j(s)y - j(s_i)y\| \leq \delta$ for some i, and

$$\|j(ss_i^{-1}) - 1\| \leq \|j(s)y - j(s_i)y\|\|(j(s_i)y)^{-1}\| \leq 1/D.$$

By right invariance of ν_S, counting the covering sets proves the assertion. Properness gives compactness of the inverse image of the indicated closed ball. Positivity follows since it contains the inverse image of an open neighborhood of the identity. □

This is `measure_ball_le_covering` in `Lib/DiscreteSeries/Haar/RegularRepL2.lean`. In the proper case, put $Q = \nu_S\{s : \|j(s) - 1\| \leq 1\}$. For $c > 8r^2$ the covering lemma implies

$$\int_S \left(\frac{2}{\|j(s)y\|}\right)^c d\nu_S(s) \leq 16^c Q. \tag{21.1}$$

For clarity, use the elementary layer-cake identity, for a function $H \geq 1$,

$$\int_S H^{-c} d\nu_S = c\int_1^\infty t^{-c-1}\nu_S\{H \leq t\}\, dt.$$

Indeed $H^{-c} = \int_H^\infty ct^{-c-1}dt$ pointwise. The lemma with $D = 1$, $R = t$ bounds the last integrand by $3^{2r^2}Qct^{4r^2-c-1}$, and $c/(c - 4r^2) \leq 2$. Multiplying by 2^c and weakening the constant proves (21.1). The corresponding formal result is `integral_decay_le`. The argument works on a subgroup, not only on the full group; the properness assumption is what ensures finite measure in the initial ball.

We also need a version of (21.1) with the real formal degree included. Let $l = 2\lambda$ be the doubled Harish–Chandra parameter, let ψ_l denote the matrix coefficient of the unit vector defining the discrete-series kernel, and let $d_l = \mathrm{formalDegree}(l)$. We use the following representation-theoretic facts.

Proposition 21.2 (Formal-source inputs: lowest-type coefficient estimates)**.** *For the large-gap parameter l of rank r with $|l_i| \leq L_0$, one has the normalized orthogonality integral $d_l \int |\psi_l(g)|^2 dg = 1$ and $|\psi_l(g)| \leq 1$. For each integer c at most the gap bound, one has*

$$|\psi_l(g)|^2 |\det a_g|^c \leq 1, \qquad |\psi_l(g)| \leq (2/\|g\|)^c, \tag{21.2}$$

where a_g is the block diagonal part relative to the signs of J_*. Also

$$|\psi_l(xg) - \psi_l(x)| \leq 100(r + L_0 + 1)^5\|g - 1\|. \tag{21.3}$$

These are `norm_dsCoeff_sq_mul_det_pow_le_one`, `norm_dsCoeff_le_pow` in `Lib/DiscreteSeries/FlenstedJensen/PsiDecay.lean` and `norm_dsCoeff_sub_le`, `half_le_norm_dsCoeff` in `Lib/DiscreteSeries/FlenstedJensen/TailBounds.lean`, using the orthogonality results of `Lib/DiscreteSeries/CuspKernel/SchurOrthogonality.lean`. We will use the exact estimates, not an unspecified asymptotic of ψ_l. For the first inequality in (21.2) the source uses polar decomposition and Fischer's inequality for positive definite block matrices; the second follows from the bound $\|g\|^2/4 \le |\det a_g|$. The Lipschitz bound comes from differentiating along a short Cayley-coordinate path and bounding the infinitesimal action on the lowest vector. We describe that differentiation and the relevant operator estimate below. The full coefficient formula in the variable signature and normalization is kept as a representation-theoretic input.

Lemma 21.3 (Degree times tail mass)**.** *With $M_0 = r + L_0 + 1$, if $c > 8r^2$ then*

$$d_l \int_{U(J_*)} (2/\|g\|)^c \, dg \le 8(2000M_0^6)^{2r^2} 2^c.$$

Proof. Put $D = 200M_0^5$. Inequality (21.3) implies $|\psi_l(g)| \ge 1/2$ on the ball $\|g - 1\| \le D^{-1}$. Hence the measure of this ball is at most $4\int |\psi_l|^2 = 4d_l^{-1}$. Lemma 21.1 gives $\nu(\|g\| \le R) \le 4d_l^{-1}(2000M_0^6R^2)^{2r^2}$. Integrating the same layer-cake identity as before and using $c/(c - 4r^2) \le 2$ proves the result, including finiteness. □

Thus the normalization by the formal degree is harmless at the level of an explicit polynomial in the maximal weight. This explains why `integral_tail_le` can be used even when d_l grows far faster than any fixed power of the scale x.

21.2 A rank-one convolution estimate and its differentiated form

Here is a direct argument for the Schur bound needed on the Bessel quotient. It is useful to formulate it with a general reproducing coefficient rather than choose a preferred Hilbert model of the real representation. On a unimodular locally compact group write the usual right convolution convention as

$$(F * x)(y) = \int x(a^{-1})F(ya)\, da.$$

Translation of an S-equivariant F to the right preserves the norm on $S\backslash G$; for our application $S = U$ is the Bessel unipotent, and its character is unitary, so the scalar products of equivariant vectors descend. Let ψ be a Hermitian integrable coefficient with $\psi(1) = d > 0$. The normalization of the idempotent used by the source consists precisely in

$$\int_G \psi(a^{-1})\psi(gay)\, da = \frac{\psi(g)\psi(y)}{d}. \tag{21.4}$$

Taking $y = 1$ gives $\psi * \psi = \psi$. This is the rank-one convolution identity, for $\psi = d\psi_l$ by Schur orthogonality. We record an abstract lemma that also explains why no unwanted dimension factor appears in the local Fourier estimate.

Lemma 21.4 (Schur orthogonality for reproduced vectors)**.** *Suppose that F is bounded, measurable and integrable modulo the left unitary character, and $F * \psi = F$. Suppose x is*

integrable, $\psi * x = x$, and $\phi \in L^1(G) \cap L^2(G)$. With the above quotient measure,

$$\|F * x\|_2^2 = \frac{\|x\|_2^2}{d}\|F\|_2^2, \tag{21.5}$$

$$\left\| \int_G \phi(a)(F * x)(\cdot\, a)\, da \right\|_2^2 \le \frac{\|x\|_2^2}{d^2}\|F\|_2^2\|\phi\|_2^2. \tag{21.6}$$

The inequality, interpreted with right side $+\infty$, also holds for integrable ϕ.

Proof. Denote by $x^*(z) = \overline{x(z^{-1})}$ the convolution involution. The adjoint of the right convolution operator on quotient L^2 is convolution with x^*; this follows by the change of variables $y \mapsto ya^{-1}$ and Fubini. The hypotheses imply $x^* * \psi = x^*$. For every integrable w, (21.4), after substitution and Fubini, gives

$$\psi * w * \psi = \frac{(\psi * w)(1)}{d}\psi.$$

Consequently, applying this with $w = x * x^*$ and using $\psi * x = x$, one obtains

$$x * x^* = \psi * (x * x^*) * \psi = \frac{(x * x^*)(1)}{d}\psi = \frac{\|x\|_2^2}{d}\psi.$$

The integral on the right is finite: $x = \psi * x$ is bounded since ψ is bounded, and x is integrable. Pairing $F * (x * x^*)$ with F proves (21.5). For $z(a) = \phi(a^{-1})$, put $w = x * z$; then $\psi * w = w$. By the same argument $w * w^* = d^{-1}\|w\|_2^2\psi$. Since

$$\|w\|_2^2/d = (w * w^*)(1)/d$$

and $w * w^* = (\|w\|_2^2/d)\psi$, we can use Cauchy–Schwarz in the identity

$$\|w\|_2^2 = (x * z * z^* * x^*)(1) = \int z(a)\,(z^* * x^* * x)(a^{-1})\, da.$$

Left convolution by $x^* * x$ on $L^2(G)$ has norm at most $\|x\|_2^2/d$: it is a positive operator with its square equal to $\|x\|_2^2/d$ times itself, by $x * x^* = (\|x\|_2^2/d)\psi$ and $\psi * x = x$. Equivalently this follows from the L^2 norm of the convolution operator of x, since its product with its adjoint is $\|x\|_2^2/d$ times a projection. Thus $\|w\|_2^2 \le (\|x\|_2^2/d)\|z\|_2^2$. Applying (21.5) to w proves (21.6). All identities may first be taken for bounded compactly supported inputs and then extended by Young's inequality and approximation; boundedness and L^1 of x, ψ, F justify the stated Fubini manipulations. □

In the formal source these are `IsBoundedMeasurable.lintegral_sq_convolve` and `lintegral_sq_convolve_convolve_le` in `Lib/DiscreteSeries/CuspKernel/BesselSchurBound.lean`. In ordinary notation one can recognize them as the fact that the right orbit of any reproduced vector contains an isometric copy of the discrete series, up to the single formal degree. The identity uses the product of the convolutions in their indicated order and the rank-one identity; ordinary L^1 convolution estimates alone would not produce d^{-1}.

To differentiate the Bessel function, it is not sufficient to know that F is smooth. The bound has to be uniform in the representation and in the maximal order of differentiation. We use smooth coefficient vectors in the discrete-series representation. If u is the normalized lowest vector, and u_w is obtained by differentiating it in the coordinate directions of a word w of length j, the needed estimate is

$$\|u_w\| \le \big((r+1)^4(L_0 + r + 1 + j)\big)^j. \tag{21.7}$$

Formal-source input: this is the filtration-wise infinitesimal action bound used in `IsBesselFunction.hasSchurBound_of_convolve_eq` in `New/RealPlace/BesselCut.lean`, from `isKHighest_dsGenerator`, the unitary Casimir inequality and `norm_apply_le_of_mem_filtration`. It applies in a unitary normalized sign frame; conjugation of the coordinate differential directions by that frame does not increase their operator norms. To obtain (21.7) from the one-step bound on filtration of depth at most j, iterate from the unit vector, always bounding the previous depth by j. This retains the powers of $(r+1)$ and cannot introduce a factorial in L_0.

For completeness, given such coefficient vectors one may recover the corresponding right derivatives of F by differentiating the convolution reproduction formula. In the source this is done by first smearing the smooth lowest coefficient by an integrable bump operator β satisfying $\mathrm{smear}(\psi_l, \beta) = \psi_l$, and differentiating the bump instead of assuming differentiability of F. Differentiation under the integral is thereby justified by the smooth compactly supported test function. The word w produces an integrable kernel x_w which is d times the matrix coefficient of u_w and the original unit vector; it has $\psi * x_w = x_w$. Orthogonality gives $\|x_w\|_2^2/d = \|u_w\|^2$. Apply Lemma 21.4 to x_w to conclude that for every integrable ϕ,

$$\left\| \int \phi(a) F_w(\cdot\, a)\, da \right\|_{U\backslash G,2}^2 \le P^{2j} \frac{\|F\|_{U\backslash G,2}^2}{d} \|\phi\|_2^2, \qquad P \ge (r+1)^4 (L_0 + r + 1 + 2s_0^2), \quad j \le 2s_0^2. \tag{21.8}$$

This is exactly the constant in (20.4). The continuity of all differentiated convolutions follows from the continuity in $L^1(G)$ of translates of an integrable kernel, and the basic boundedness of F. The differential relation between successive words follows by testing along paths of right multiplication and using the ordinary chain rule on the bump. In particular no differentiation of the compact cutoff is used in (21.8); the estimate concerns the *full* Bessel function. This distinction is used later in the energy comparison between full and cut functions.

21.3 Threshold projectors and a lowest-type coefficient

The construction of a peak starts with a coefficient, not with a compactly supported test function. We give the reason that the alternating signs and steepness of the weights in that construction matter. In this subsection all matrix norms without an explicit subscript are Hilbert–Schmidt norms, and

$$J = \mathrm{diag}(\epsilon_1, \dots, \epsilon_n), \quad K = \{g : g^* g = 1,\ gJ = Jg\}.$$

The two blocks of K are compact; they must not be confused with the two unitary groups in the pair construction below.

For real integral weights $w = (w_i)$ define

$$Q_w(X) = \sum_{i,j} |w_i - w_j| |X_{ij}|^2.$$

If P_t is projection onto the coordinates at which $w_i \ge t$, then

$$\sum_t \|[P_t, X]\|^2 = Q_w(X), \qquad \sum_t \mathrm{tr}(P_t X) = \sum_i (w_i + a) X_{ii}, \tag{21.9}$$

where the thresholds run from $-a+1$ to a, with $a = \sum_i |w_i|$. Indeed the ij entry of the commutator has absolute value $|X_{ij}|$ if precisely one of w_i, w_j exceeds the threshold, and is zero otherwise. There are exactly $|w_i - w_j|$ such thresholds. Each diagonal entry is counted $w_i + a$ times. The same identity holds with negative or shifted weights. In particular repeated weights give flat directions, unrelated to the size of the absolute values of the weights.

Lemma 21.5. *For a compact unitary matrix u one has*

$$\prod_t |\det(P_t u P_t|_{\operatorname{im} P_t})|^2 \le \exp(-Q_w(u)/2).$$

Proof. For a single orthogonal projection P put $A = PuP|_{\operatorname{im} P}$ and $B = (1-P)uP$. Then $A^*A = 1 - B^*B$, whose eigenvalues lie in $[0,1]$. The inequality $1 - x \le e^{-x}$, including at $x = 1$, gives

$$|\det A|^2 \le e^{-\operatorname{tr}(B^*B)} = e^{-\|B\|^2}.$$

Unitarity gives $\|Pu(1-P)\| = \|B\|$: the trace of A^*A equals the trace of AA^* and subtracting both traces from $\operatorname{rank} P$ proves the equality. Consequently $\|[P,u]\|^2 = 2\|B\|^2$. Multiplication and (21.9) conclude. □

This argument is `norm_weightMinors_sq_le_exp` in `New/DiscreteSeries/GaussianShape.lean`. It explains why integrating the coefficient over K does not destroy the concentration: weighted minors of a unitary matrix themselves act as a concentration density around the commutant of the weights.

Here are the precise smaller-scale statements required to evaluate the phase. They are recorded as formal-source inputs because a proof of the full Flensted–Jensen formula with explicit remainder would take us beyond the scope of this account. Write ψ_l for the normalized highest-vector coefficient, $\Lambda(l)$ for the diagonal of the lowest K-type (in the notation of the source, `lowestKType`), and $L_0 \ge \max|l_i|$.

1. In `New/DiscreteSeries/MinorExpansion.lean`, `exists_det_compress_eq_exp_logOneAdd`, if P, Q are orthogonal projections, $\|X\| \le \delta \le 1/2$, $\operatorname{tr}([P,Q]Y) = 0$, $\|[P,X] - Y\| \le \delta c$, and $\|Q - P\|, \|Y\| \le c$, then

   $$\det(1 + PX) = \exp\big(\operatorname{tr}(Q\log(1+X)) - \tfrac{1}{4}\operatorname{tr}(Y^2) + e\big), \quad |e| \le 100\delta c^2.$$

   The noncommuting term costs a norm *squared*; a remainder simply bounded by $O(\|X\|^3)$ would not suffice when some of the weights coincide.

2. In the same file, `exists_principalMinor_eq_exp` applies to the projected minors of the Flensted–Jensen integrand at $g = 1 + X$, $\|X\| < 10^{-3}$. For the corresponding threshold projection P and compact u it writes the minor as

   $$\exp\big(\operatorname{tr}(P\log g) + \tfrac{1}{4}\|[P,X]\|^2 + e_P\big), \qquad |e_P| \le 200000\|X\|(\|[P,X]\|^2 + \|[P,u]\|^2).$$

   The integrand is evaluated at the matrix y related to g by the triangular shifts in the Flensted–Jensen formula; this is not an asserted expansion of $\det(Pu^{-1}gu)$ at arbitrary g, u.

3. The resulting `dsCoeff_gaussian_shape` in `New/DiscreteSeries/GaussianShape.lean` is the uniform statement: take the minimum required noncompact weight separation at least $500n^3$ (in addition to the given gap parameter); let $T \ge \max(1, n^2, n^2\log(n + L_0 + 1))$, $\delta T \le 10^{-12}$ and $\|g - 1\| \le \delta$. Then there exists $0 \le \kappa \le 1$ such that

   $$\left|\frac{\psi_l(g)}{\exp(\operatorname{tr}(\Lambda(l)\log g))} - \kappa\right| \le \kappa/10 + e^{-100T}. \tag{21.10}$$

   The proof in the source takes κ to be the Gaussian with exponent minus one quarter of the weighted commutator quadratic form associated with the distinct shifted weights `-paramAddRho`; these weights must be distinguished from the lowest K-type in the exponential phase.

We explain the quantitative mechanism linking these statements. The Flensted–Jensen integral is a normalized compact integral of a product of minors times a lowest-type weight. Apply (21.9) to the first term and the quadratic term of the logarithmic expansion in each minor, using the appropriate weight sign. The phase term is independent of the integration parameter u and can be removed from the integral. The sum of the remainders is then bounded in the source by

$$10^6\delta(Q(X)+Q(u)/2+n^2),$$

with the quadratic forms determined by the shifted weights; the n^2 accommodates the shifts by the half-sum of roots. By Lemma 21.5, the compact density includes the factor $e^{-Q(u)/2}$. Thus, on the set where $Q(X)+Q(u)/2$ is at most a large multiple of T, the exponential of the remainder differs from 1 by a small relative error. On the complement, the Gaussian dominates the exponential of the remainder. The normalization of the compact integral cannot cost more than a power of the dimension of the weight representation, which is at most exponential in $n^2\log(n+L_0+1)$. This accounts for the condition on T. The separation between these two regimes is essential: requiring a relative error uniformly, including where κ is extremely small, would be false. Equation (21.10) supplies precisely the relative error *plus* exponentially small absolute error that can be combined over two factors.

For illustration, a general 2-by-2 matrix direction with $w_1 > w_2$ has

$$Q_w(X) = (w_1-w_2)(|X_{12}|^2+|X_{21}|^2).$$

The Gaussian width in these coordinates therefore scales as $(w_1-w_2)^{-1/2}$, not $L_0^{-1/2}$ if the two weights move together. A rank-one compact direction with $w_1 = w_2$ is completely invisible to the Gaussian. The common compact group in the next construction is exactly what allows these invisible directions to be removed before using (21.10).

21.4 Two-factor Laplace estimate

The restriction integral is not an integral of a positive coefficient: the additive character must cancel the first-order phase. Let $j_i : S \to G_i$ be a unitary pair of homomorphisms, of ordered ranks n_i. Write $u_i(s) = j_i(s) - 1$ and let Λ_i be the two lowest-K-type weight matrices. The norm on differences of matrices here is Hilbert–Schmidt for the phase and compact-correction arguments, and the norm of a group element itself is the operator norm in the decay bound. The relevant additive phase is

$$\varphi(s) = \operatorname{tr}(\Lambda_1 u_1(s)) - \operatorname{tr}(\Lambda_2 u_2(s)).$$

In the actual pair it is an imaginary-valued homomorphism. It follows that for $\|u_i(s)\| \le 1/2$,

$$\operatorname{tr}(\Lambda_1 \log j_1(s)) - \operatorname{tr}(\Lambda_2 \log j_2(s)) = \varphi(s). \tag{21.11}$$

Here is a useful elementary reason. Put

$$A_k = \operatorname{tr}(\Lambda_1 u_1(s)^k) - \operatorname{tr}(\Lambda_2 u_2(s)^k).$$

Additivity of the phase on powers s^q and the binomial expansion give $\sum_{k=1}^{q}\binom{q}{k}A_k = qA_1$. At $q=2$ this gives $A_2 = 0$; induction gives $A_q = 0$ for all $q \ge 2$. The logarithmic series is absolutely convergent, giving (21.11). Thus the cancellation is exact, not merely a first-order approximation. The source statement is `phase_logOneAdd` in `New/DiscreteSeries/TwoFactorSetup.lean`.

There is another elementary matrix identity behind the bound on the real part of the remaining phase. For $gJg^* = J$, the ii row gives

$$\mathrm{Re}(g_{ii} - 1) \cdot 2 + \sum_j |g_{ij} - \delta_{ij}|^2 = 2 \sum_{\epsilon_j \neq \epsilon_i} |g_{ij}|^2.$$

In fact the left side is the squared ordinary row norm minus 1, while the signed row norm is ϵ_i. Multiplying by nonnegative row weights w_i, bounding them by C on rows with noncompact directions, and summing, gives

$$\sum_i w_i \sum_j |g_{ij} - \delta_{ij}|^2 + 2\,\mathrm{Re}\,\mathrm{tr}(\mathrm{diag}(w)(g-1)) \leq 2C \sum_{\epsilon_j \neq \epsilon_i} |g_{ij}|^2.$$

One can also replace w_i in the first term by any smaller nonnegative quantity, as in `norm_sq_add_re_trace_le` of the same file. This accounts for the noncompact quadratic errors when comparing the two phases.

If g_{diag} is the part block diagonal with respect to signs, the decay estimate gives

$$|\psi_l(g)|^4 \leq (1 + \|g - g_{\mathrm{diag}}\|^2)^{-\mathrm{gap}}.$$

Indeed the two diagonal square determinants come from $1 + BB^*$ and its companion with B the off-diagonal block; the product inequality $\det(1 + BB^*) \geq 1 + \mathrm{tr}(BB^*)$ follows by expanding the product of its nonnegative eigenvalues. Combine this with the bound of the coefficient by the inverse power of the diagonal determinant recorded in the previous subsection. The precise simultaneous two-block expression is `norm_dsCoeff_pow_four_le` in `New/DiscreteSeries/TwoFactorSetup.lean`.

The remaining group-theoretic ingredient is an important distinction from a decay estimate in a given representation. The setup `IsTwoFactor` in `New/DiscreteSeries/TwoFactorSetup.lean` includes that the pair has matching zero-weight compact blocks, that it contains their common compact group, and that the steep weights away from this block have alternating signs. Under these conditions `exists_near_one` asserts that if the squared norms of the two off-block directions sum to at most ρ^2, $0 < \rho \leq 1/100$, one can multiply by an element of the common compact subgroup without changing the integrand so that

$$\|j_1(s) - 1\|, \|j_2(s) - 1\| \leq 10\rho.$$

This is a formal-source input (a specific matrix estimate on the pair); it does not follow just from j_i being representations. The two images of the compact subgroup have to come from the very same s.

Proposition 21.6 (Restriction integral)**.** *With the two-factor setup let $N = (n_1 + n_2)^2$ and $\beta \geq 10^{12}$, with $n_1 + n_2 + L_0 \leq \beta$ and weight gap at least $10^{30}(N \log \beta)^3$. Assume that j_1 is continuous and proper, j_2 is continuous and $\|j_2(s)\| \leq \beta + 1$ whenever $\|j_1(s) - 1\| \leq 1$. Use the integrand*

$$\theta(s) = \psi_{l_1}(j_1(s))\overline{\psi_{l_2}(j_2(s))}e^{-\varphi(s)}.$$

For Haar measure ds on S,

$$\mu\{s : \|j_1(s) - 1\| \leq 1\} \leq \beta^{20N} \left| \int_S \theta(s)\, ds \right|. \tag{21.12}$$

We give the integration argument of the source, separating from it the precise pointwise input used. In `New/DiscreteSeries/RestrictionLemma.lean`, `IsTwoFactor.laplace_method`,

the Gaussian estimate (21.10), the off-diagonal decay and `exists_near_one` give two auxiliary bounds:

$$|\theta(s)| \le (2/\|j_1(s)\|)^a \qquad (0 \le a \le 18N), \tag{21.13}$$

$$4|\theta(s)| \le 8\left(\operatorname{Re}\theta(s) + 16\beta^{-50N}(2/\|j_1(s)\|)^{8N}\right). \tag{21.14}$$

The numerical pointwise estimates are formal-source inputs in this proof. To see their structure, use the Gaussian estimate with T proportional to $N\log\beta$ after the common compact correction. The product of the two ratios to their exponential phase factors is within a small relative error of the product of the two nonnegative Gaussians, except for the exponentially small absolute errors. The exponential phase factors cancel by (21.11). The row norm estimate prevents the absolute errors from being magnified by an unbounded real exponential. Away from the applicable neighborhood the gap forces decay, and (21.13) permits the residual error to be measured against an integrable weighted function, rather than just the uniform bound β^{-50N}. The cubic power in the gap is responsible for having both a sufficiently small logarithmic neighborhood and sufficiently drastic decay beyond it.

Proof of Proposition 21.6 from the pointwise estimates. Put $A = \{\|j_1(s) - 1\| \le 1\}$ and $A' = \{\|j_i(s) - 1\| \le (200\beta^5)^{-1},\ i = 1, 2\}$. The two-factor norm ball covering lemma gives

$$\mu(A) \le \beta^{16N}\mu(A').$$

To apply it use the proper block direct sum of the two homomorphisms (properness follows from j_1), the norm bound by $\beta + 1$ for the second group on A, and the small radius $1/(200\beta^5)$. The elementary net cardinality is a power of β with exponent $16N$ (real matrix dimension at most $2N$). Properness ensures the sets have finite Haar measure. The near-identity coefficient inequality, with its weight bound by β, gives $|\psi_{l_i}(j_i(s))| \ge 1/2$ on A'. The phase has absolute value one. So

$$\mu(A') \le 4\int_S |\theta| \le 8\left(\left|\int_S \theta\right| + 16\beta^{-50N}\int_S (2/\|j_1(s)\|)^{8N}\,ds\right).$$

By the norm decay integral bound in the opening subsection the last integral is at most $16^{8N}\mu(A)$, because $8N > 8n_1^2$. Multiplying by β^{16N}, the coefficient of $\mu(A)$ on the right is at most $128 \cdot 16^{8N}\beta^{-34N} < 1/2$. Absorb it, then use $16\beta^{16N} \le \beta^{20N}$. This gives (21.12). □

Several features of the argument are worth noting. It never invokes positivity of the original coefficient or replaces the oscillatory integral by an integral of its modulus. It is the near-positive cone estimate (21.14) which prevents cancellation. Rescaling the Haar measure multiplies both sides of (21.12) by the same factor. Hence the estimate is compatible with taking Haar probability on the compact factor of the Bessel subgroup, while leaving Haar measure on the unipotent factor fixed by the global construction. Finally the resulting lower bound depends on a volume near the identity, not on an arbitrarily normalized coefficient at some moving point. This is what permits the peak comparison to be transported to the quotient where the Bessel kernel is used.

21.5 Interpolation, isotropic frames, and the Bessel peak

The role of the strict order of the weights is not merely to ensure membership in regular discrete series. It also permits a constructive real embedding whose restriction is guaranteed to be nonzero by Proposition 21.6. We reproduce the algebra of this embedding.

Lemma 21.7 (Alternating interpolation weights)**.** *Let $a_0 > \cdots > a_{d-1}$ be distinct integers, and set*

$$c_i = \Big((-1)^i \prod_{j \neq i} (a_i - a_j)\Big)^{1/2}.$$

Then $c_i \geq 1$ and, for $\deg p < d$, one has

$$\sum_i \frac{(-1)^i p(a_i)}{c_i^2} = [t^{d-1}]p(t).$$

Proof. There are exactly i negative factors in the defining product, so its signed value is a positive integer. The polynomial

$$L_i(t) = \frac{\prod_{j \neq i}(t - a_j)}{\prod_{j \neq i}(a_i - a_j)}$$

equals 1 at a_i and 0 at all other a_j. The degree $< d$ polynomial $p - \sum_i p(a_i)L_i$ has at least d roots and hence is zero. Compare leading coefficients. □

This is `exists_lagrange_weights` in `Lib/General/LinearAlgebra/FramedCoefficient.lean`. We use it in a space with $2h + 2$ noncompact coordinates, of alternating signs $+, -, +, -, \ldots, +, -$. The other coordinates form the common positive compact block. Let the exceptional weights in descending order be λ_i, $0 \leq i \leq 2h + 1$, and let a be the distinguished weight in the compact factor. The steepness guarantees that $t_i = \lambda_i - a$ are strictly descending and negative. Apply the lemma to

$$(a_0, a_1, \ldots, a_{2h+2}) = (0, t_0, \ldots, t_{2h+1}).$$

Form columns in the exceptional coordinates by

$$(C_0)_i = c_0/c_{i+1}, \qquad (C_j)_i = t_i^j/c_{i+1} \quad (j \geq 1).$$

Multiplying the $i + 1$ sign in the lemma by -1 is the essential sign change: the ambient Hermitian sign is $(-1)^i$. For $j + l \leq 2h + 2$ the raw identity is

$$\sum_i \frac{(-1)^i t_i^{j+l}}{c_{i+1}^2} = \frac{\delta_{j+l,0}}{c_0^2} - \delta_{j+l,2h+2}.$$

Consequently

$$\begin{gathered} H(C_0, C_0) = 1, \\ H(C_j, C_l) = 0 \quad (0 < j + l < 2h + 2), \\ H(C_j, C_l) = -1 \quad (j, l > 0,\ j + l = 2h + 2). \end{gathered}$$

In particular $C_1, \ldots, C_h$ form a totally isotropic frame and C_{h+1} is a negative unit vector orthogonal to this frame. No assumption about unknown zeros of an L-function enters here; the relation is just interpolation.

For a diagonal weight operator $D = \mathrm{diag}(\lambda_i)$ on the exceptional coordinates, one has $DC_j = aC_j + C_{j+1}$ for $j \geq 1$ and $DC_0 = aC_0 + c_0C_1$. These equalities show how the shifted weight projects onto a nilpotent chain. The trace phase of the source's pair is designed using that chain. If one simply chose an arbitrary isotropic flag of the same dimension, its Bessel character need not agree with the coefficient's differential.

As a check, take $h = 0$, $t_0 = -1$, $t_1 = -2$. Then $(c_0, c_1, c_2) = (\sqrt{2}, 1, \sqrt{2})$ and the two columns are

$$C_0 = (\sqrt{2}, 1)^t, \qquad C_1 = (-1, -\sqrt{2})^t.$$

Relative to the sign matrix $\mathrm{diag}(1, -1)$ these have squared lengths $1, -1$ respectively, and they are orthogonal. The construction generalizes the same cancellation to successively many isotropic columns.

It is not enough to prove that these columns give the desired Hermitian Gram matrix. We must bound the size of a conjugation identifying the resulting flag with the standard flag. The relevant elementary estimate behind that step is that the matrix of minors determining the columns has denominators $c_i \geq 1$. If $|t_i| \leq T$ then a raw column of degree j has norm at most $\sqrt{2h+2}\,T^j$. The initial column has the additional factor c_0, bounded by T^{h+1}. Nondegeneracy in the reverse direction follows from the negative anti-diagonal: a vector with pairing of magnitude 1 against a vector of norm M has norm at least M^{-1}. This explains why passing to orthogonal normalized frames here costs only a polynomial in the base parameter with exponent linear in h, rather than in the number of all entries of a k-by-k matrix.

The formal source makes the preceding construction precise in the following two assertions, which we use as formal-source inputs.

- `besselSize_le_of_dual`, `New/RealPlace/FrameNorm.lean`: if $V_0, \dots, V_{2h}$ are vectors of norm at most M, with V_j equal to the Bessel frame columns for $j \leq h$, and $\langle V_i, V_j \rangle = 0$ for $i + j < 2h$ and $= -1$ for $i + j = 2h$, then the Bessel size of the frame is at most $8(h+1)M$ (in the normalized sign frame).
- `IsSteep.exists_isTwoFactor_besselPairHom`, `New/DiscreteSeries/BesselExistence.lean`: for the steep sign and weight pattern on the two groups, there exists a unitary realization of the preceding columns, carrying the standard Bessel unipotent with its character to a pair of homomorphisms satisfying `IsTwoFactor`. The compact part is identified with the common zero-weight block. The linear trace difference is imaginary and additive and exponentiates to the given Bessel character on the unipotent. In the parameter ranges of the local construction this realization has $\mathrm{bs}(x) \leq b^{5s_0}$ and both homomorphisms meet the small-ball hypotheses of Proposition 21.6 for $\beta = b^3$.

The second assertion is stronger than Witt extension: the phase must match, and the quantitative bound for the dual array must also survive the identification. The anti-diagonal identity above exhibits the input to the verification of both properties.

Let S denote this Bessel subgroup, with Haar measure the unipotent Haar measure times Haar probability on its compact factor. In local coordinates its embedding into the ambient group is $s \mapsto s_1$; its action on the compact type space is s_2. For a unit vector v of the compact representation ρ with coefficient ψ_{l_2}, define

$$Y(y) = \int_S \psi_{l_1}(x^{-1} s_1 y)\, e^{-\varphi(s)} \rho(s_2^{-1}) v \, ds. \tag{21.15}$$

Conjugating the two group maps by their chosen frames, Proposition 21.6 gives

$$\|Y(x)\| \geq |\langle Y(x), v \rangle| = \left| \int_S \theta \right| > 0.$$

The inner product uses the normalized unit compact vector, so bounding the integral by the norm has no dimension factor. The norm of the integrand in (21.15) is the absolute value of the coefficient. The norm-ball decay theorem then gives absolute convergence; the same domination is uniform on compact sets of y by submultiplicativity and properness. Bessel

equivariance follows by changing the integration variable s by right multiplication, noting that the character is multiplicative and that Haar measure on the semidirect product is unimodular. Fubini and the coefficient idempotent ψ show $Y * \psi = Y$ with the normalized coefficient $\psi = d\psi_{l_1}$. Thus the whole orbit of the peak, and not just its value, lies in the reproduced subspace to which the Schur estimates apply.

For clarity, the useful property `HasPeakAt` of `New/DiscreteSeries/BesselExistence.lean` comprises both

$$\|Y(y)\| \le \int_S |\psi_{l_1}(x^{-1}s_1y)|\,ds, \qquad \int_S (2/\|x^{-1}s_1y\|)^{8r^2+1}\,ds \le b^n\|Y(x)\| \tag{21.16}$$

for every y in the normalized frame, with the specified exponent n, and $Y(x) \neq 0$. The second inequality follows from Proposition 21.6 by the norm-ball integral bound with the moving coset y; its use does not presume that y belongs to the image of S. The source declaration `exists_isBesselFunction` supplies Y with these properties under the above pair assumptions. The exact admissible n is chosen in that declaration so that in the global application $n + 99r^2 \le r^3$. This margin will pay for the conversion of the pointwise peak into an L^2-normalized cutoff.

21.6 A peak-to-cut inequality with all mass factors

The lower bound at the peak is of little global use if most of the resulting function is concentrated far from a bounded Bessel-size region. We now prove the cutoff argument in a form that displays why the formal degree, Haar measure, and peak height cannot be normalized independently.

Put

$$M = 8r^2 + 1, \quad C = 16^M, \quad R = b^{6s_0+2}, \quad \omega_a(g) = (2/\|g\|)^a, \quad I = \int_G \omega_{2M}(g)\,dg.$$

In what follows the norm is the source's submultiplicative matrix size, symmetrized under inversion, in the normalized frame. Let B be the group norm ball of radius 1 about the identity (measured by the norm of the difference); it has Haar measure $v_B > 0$. The Bessel size of a coset satisfies

$$\mathrm{bs}(yg) \le \mathrm{bs}(y)\|g\|.$$

This follows from its definition as the infimum of $1+$ the norm of a representative under the Bessel subgroup and the norm inequalities. Under the source's bounds $\mathrm{bs}(x) \le b^{5s_0}$ and $b \ge 2000$, every translate $t^{-1}B$, where $t = x^{-1}s_1y$, lies in $\{g : \mathrm{bs}(yg) \le R\}$. Indeed for $g = t^{-1}u$ with $u \in B$ one has $s_1yg = xu$ and $\mathrm{bs}(xu) \le 16b^{5s_0} \le R$.

Define the positive majorant and the positive finite measure

$$\Phi(y) = \int_S \omega_{2M}(x^{-1}s_1y)\,ds, \qquad d\nu(g) = \frac{C}{v_B} b^n\|Y(x)\|\omega_M(g)\,dg.$$

The degree-tail bounds in the first subsection imply $I < \infty$ and $\int \omega_M \le Cv_B$, so $\nu(G) \le C^2b^n\|Y(x)\|$. Writing $t = x^{-1}s_1y$, for $g \in t^{-1}B$ the norm of g is at most $16\|t\|$. Thus

$$\nu\{g : \mathrm{bs}(yg) \le R\} \ge b^n\|Y(x)\|\omega_M(t).$$

Multiply by $\omega_M(t)$ and integrate over s. The second inequality of (21.16) cancels $b^n\|Y(x)\|$ and proves

$$\Phi(y) \le \nu\{g : \mathrm{bs}(yg) \le R\}. \tag{21.17}$$

This argument does not require the family of translates $t^{-1}B$ to be disjoint.

The quotient integration convention is important. With Haar probability on the compact part of S and the given unipotent Haar, Weil integration gives

$$\int_{U\backslash G} \Phi(y)\,dy = I.$$

Indeed unfold first along the unipotent, then use left invariance under the compact part and x. The decay exponent can be chosen so that the first inequality in (21.16) implies $\|Y(y)\| \le \Phi(y)$ everywhere. If $\mathrm{bs}(y) > R-1$, the estimate $\mathrm{bs}(y) \le \mathrm{bs}(x)\|t\|$ forces $\|t\| > 2b$ for every s. An extra decay exponent A therefore improves the same bound to

$$\|Y(y)\| \le b^{-A}\Phi(y) \quad (\mathrm{bs}(y) > R-1).$$

In particular Y is square integrable, since $\Phi \le \nu(G)$ and Φ has finite quotient integral. Its squared norm is strictly positive by continuity and $Y(x) \neq 0$.

It remains to estimate how much of the majorant is needed when Y is normalized. The coefficient reproduction formula at x, combined with the integral construction of Y and Bessel equivariance, gives the bound

$$\|Y(x)\| \le d \int_{U\backslash G} \Phi(y)\|Y(y)\|\,dy.$$

Apply Cauchy–Schwarz with weight Φ and use (21.17):

$$\|Y(x)\|^2 \le d^2 I \int_{U\backslash G} \Phi(y)\|Y(y)\|^2\,dy \le d^2 I \nu(G)\|Y\|_2^2. \tag{21.18}$$

Set $c^2 = D_v/\|Y\|_2^2$, where D_v is the index cardinality of the local type. Since $Y(x) \neq 0$ we can cancel its norm in (21.18), using the bound for $\nu(G)$, to obtain

$$c^2\nu(G)I \le D_v(dIC^2b^n)^2.$$

The degree-tail estimate gives $dIC^2 \le b^{99r^2}$ for the weight and rank ranges $r + L_0 + 1 \le b^4$ in use. Thus $n + 99r^2 \le r^3$ implies

$$c^2\nu(G)I \le D_v b^{2r^3}.$$

These explicit numerical inequalities are precisely the budget in `HasPeakAt.exists_cut_isCutPair`, `New/RealPlace/BesselCut.lean`; they make the following tail estimate independent of the original normalization of Y or of the Haar measure.

Use the continuous cutoff $\chi(t) = \max(0, \min(1, R-t))$. Write $F = cY$, $V = \chi(\mathrm{bs})F$. Then $\|F\|_2^2 = D_v$ and V is supported on Bessel size at most R. Multiplicative properties of bs and continuity of the norm give continuity of bs, so V is continuous. More explicitly if $g \to 1$ both $\|g\|$ and $\|g^{-1}\|$ in the multiplicative size tend to one; sandwich the two values of bs using the above inequality. The tail and mass estimates prove

$$\begin{aligned}\|V-F\|_2^2 &\le b^{-2A}c^2 \int_{U\backslash G} \Phi(y)^2\,dy \\ &\le b^{-2A}c^2\nu(G)I \le D_v b^{-10r^2}.\end{aligned} \tag{21.19}$$

The last step uses the source margin $A \ge 3r^3 f$, with f the number of infinite places, and the given ranges of the ranks. Moreover $c\Phi$ majorizes both norms, and (21.17) gives a majorant formed by translating a bounded-size region against a measure. These are stronger properties than a small L^2 tail; they will be needed on cell charts.

One must not apply the differentiated convolution estimate to V. Its cutoff is only continuous and piecewise linear, and it is not reproduced by the idempotent. The later bandlimit argument uses the full function F and inserts cutoff regions at the level of integration. The tail estimate (21.19) is exactly the price for passing between the two, in a setting where no derivatives of the cutoff are required.

21.7 From an L^2 reproducing estimate to the bandlimit tower

We detail the architecture of the still necessary local orbital calculation. The norm and tail estimates just proved do not imply arbitrarily strong Fourier decay on the orbit. The Fourier exponent depends on the dimension of the positive middle space, whereas the differentiated Schur estimate uses derivatives only of order at most $2s_0^2$. These are different sources of regularity.

The local calculation in `New/RealPlace/LocalTheorem.lean` has a descending tower of groups G_j (fixing an initial portion of the isotropic flag and its dual) with Haar measures, upper quotients and measures on floor coordinates. At a level j the notation md stands for the module of multiplication on the local additive field, so at a complex place $\mathrm{md}(z) = |z|^2$. Write the two heads as p, q. The multiplicative prefactor

$$D_j(p,q) = \prod_{a<j} \mathrm{md}(p_a q_a)^{d_a/2}$$

is the source's `Level.headDensity`, with d_a the successive floor dimensions. Under a change of variable by multiplication by p_a in d_a complex additive coordinates the real Jacobian is $|p_a|^{2d_a}$; writing the geometric mean of the two head Jacobians gives precisely the product above. It is especially important not to discard the square of the usual complex absolute value here: it would change the exponent of the final lattice-covolume factor.

The local normalized form is denoted by `floorForm`. It integrates a translate pairing of two vector-valued Bessel functions against the upper group and its descending convolution of floor measures. The corresponding absolute integrand is `formModulus` and its dominating integral is `formMajorant`. Descending convolution of measures means simply the pushforward of the product under the ordered group multiplication; it is not commutative. For bounded compactly supported functions, expanding one factor of this product and applying Fubini is the identity `upperPairing_eq_integral_succ`. For the actual forms, integrability on all the intermediate levels is obtained by the majorant bound below. Thus the equality of the zero-frequency integral with the next floor is an equality, not a rough inequality: the sharp constant in the final sieve depends on it.

The majorant is controlled by a product of two self-norms, denoted $\mathcal{D}_j(x)$ here and `Gauge.selfNormProduct` in the source. To illustrate the use of such a product, for a positive measure λ on the upper quotient and two vector-valued functions A, B, the elementary inequality is

$$\left| \int \langle A(z), B(z) \rangle \, d\lambda(z) \right|^2 \le \left(\int \|A(z)\|^2 \, d\lambda(z) \right) \left(\int \|B(z)\|^2 \, d\lambda(z) \right).$$

If the integration is also restricted to a measurable region of measure s^e, a further Cauchy–Schwarz with that region's indicator supplies a factor s^e on the right. This is the source of powers of a small floor volume, in contrast with powers coming from derivatives. In the actual gauge, there are changes of variables in the two factors and the self-norms live on different floors. The following are the formal-source inputs, with their quantitative conclusions; all are in `New/RealPlace/LocalTheorem.lean`:

1. `Gauge.lintegral_selfNorm_le`: for a pairing kernel at level $j \le s_0$, if the product of floor constants times its upper-quotient squared integral is at most c, the integral of its self-norm over the gauge's bounded positions is at most $s_0 c$.

2. `Gauge.formMajorant_sq_le`: if the floor mass is at most V, $V \ge 1$, with the specified near-region conditions and floor-value volume at most $s < 1$ on nonzero pairing integrands, then
$$\mathcal{I}_j^+(x)^2 \le s^{\mathrm{volumeExponent}} V^2 \mathcal{D}_j(x).$$
Here $\mathcal{I}_j^+$ is the form majorant. The hypotheses include nearness of both arguments of the pairing after their Weyl translations, as well as nearness of the support of the second Bessel function.

3. `Gauge.formModulus_sq_le_of_char`: for the character of frequency ξ on the diagonal frame Gram coordinate, assuming the gauge's multiplicative size bound for the character variant, the squared modulus integral is at most
$$(P.c/\mathrm{md}(\xi))^{|\iota|}\mathcal{D}_j(x).$$
The exponent $|\iota|$ is the cardinality of the complex middle-coordinate index in that gauge. The same right side occurs in `Gauge.formModulus_sq_le_of_heads` for the ordinary pairing at level $j = i + 1$ if $\xi = g^{-1}(p_i^{-1} - q_i^{-1})$. The first bound comes from averaging a variable character; the second comes from the same character on two unequal Weyl heads. Neither is just the assertion that an L^2 Fourier transform is square integrable.

4. `Gauge.lintegral_positions_le`: with the product of pairing norms equal to Θ^2, the function $a(x)$ on a bounded position set satisfying $a(x)^2 \le e^2 \mathcal{D}_j(x)$ obeys
$$D_j(p,q) \int a(x)\,dx \le e\,\Theta \left(\prod_{l<s_0} \mathrm{floorModulus}(l) \right) (P.c)^{|\iota| s_0} \Lambda^{1500000 s_0^3 + 1} (\Lambda')^{50000 s_0^2}.$$
Here Λ, Λ' are the sizes of the two gauges on the two position sets. This is a bound uniform in the Weyl heads; the head-density factor on the left is part of the assertion.

5. `Level.integral_succ_floor_eq` and `Gauge.integral_floorForm_succ`: integration with one additive character against the floor-value measure is the corresponding integral of the translate pairing with the character on the diagonal frame Gram variable, with the next upper group measure and stack measure. At zero character it gives the preceding floor. The hypotheses include the finite integral of the majorant. These statements implement the ordered convolution change of variables described above.

The distinction between the last two Fourier estimates can be seen abstractly. Suppose a pairing along a torus orbit contains two exponential factors $e^{2\pi i \,\mathrm{Tr}(p^{-1}z)}$ and $e^{2\pi i \,\mathrm{Tr}(q^{-1}z)}$ in its two vectors. Their inner product contains their ratio, so its character is that of $(p^{-1} - q^{-1})z$, independent of the common phase. This explains why the same Fourier gain controls both a nonzero additive frequency at the integration step and an unequal-head contribution at the step between floors. The scale by g^{-1} comes from the Bessel character's root-coordinate normalization, not a change of the frequency decay exponent.

One can therefore assemble the local tower as follows. At $j < s_0$ set J_j equal to the normalized floor form; at $j = s_0$ set $J_{s_0}(p,q) = \langle v(p), v(q) \rangle$, where v is the vector of normalized Bessel correlations. This last form is a Gram kernel when the last entries agree. The step identity gives J_j for zero frequency, and the majorant and frequency estimates give head weights times the common damped-power function at nonzero frequency or unequal heads.

The top floor identity uses the change of variables between the diagonal Gram entry and the common last vector. The exact geometric normalization also uses that compact double cosets of the last scalar coordinate with the same module have the same pairing. The source theorem implementing this assembly is

hasBandLimits_of_isCutBessel in New/RealPlace/BandLimits.lean.

It states that for a local cut/full Bessel pair of an admissible rank, positive degree d, Schur bound $\mathrm{SchurLe}(F, d^{-1}D_v)$ and $d \cdot \mathrm{bandConst} \le b^{4r^2}$, there exists a kernel tower satisfying `HasBandLimits`. Its bottom bound $(1 + b^{-10})/d$, the head weights and the nonzero-frequency majorant are those in Section 20.4. We retain the local coordinate and numerical inequalities just stated as formal-source inputs. The purpose of separating them is that the analytic normalization $d^{-1}D_v$, which was derived in Section 21.6 and the Schur argument, now becomes the precise number entering this tower, whereas the large Fourier exponent $k - s_0^7$ results from averaging in the middle coordinates.

21.8 The cell L^2 estimate and its limited differentiation cost

The open-cell estimates serve a separate purpose from the bandlimit tower: crude and unequal-link bounds need an integral over the frame Gram section. The local lemma which transfers quotient norm estimates to bounded sets in the group is worth spelling out. The residual stabilizer U^0 fixes the entire chosen frame, while the quotient of U by that stabilizer is described by upper triangular frame entries. Its dimension in the complex real-place coordinates is at most $2s_0^2$.

Suppose w is nonnegative, measurable and left Bessel-invariant, and a is a nonnegative measurable cutoff satisfying

$$\sup_y \int_{U^0} a(uy)\,du \le K.$$

Suppose whenever $a(y), a(sy) \ne 0$ and $s \in U$ the upper frame entries of s have module (ordinary complex norm) at most R. For any U-orbit meeting the support of a, choose a single point y_0 of this intersection. All other points in the intersection are obtained by elements whose frame matrices lie in the box of side $2R$ about zero in each of the real and imaginary coordinates. Integrating across U^0 first, the contribution of the outer coordinates is at most $K(2R)^{2s_0^2}$; the use of a single basepoint on each orbit is only a pointwise bound of that integral, and requires no measurable choice. Weil integration and the invariance of w give

$$\int_G a(y)w(y)\,dy \le K(2R)^{2s_0^2} \int_{U\backslash G} w(y)\,dy.$$

This is the substance of `lintegral_mul_le_of_frameMatrix_le`, `New/RealPlace/SobolevCell.lean`, using the source's coordinates with the product additive measure. The dimension that appears here is *not* the real dimension of U, which includes the large positive-middle block.

Here is an elementary Sobolev estimate explaining the other appearance of $2s_0^2$. For a continuously differentiable scalar function f on an interval of length $2t$ centered at 0, one has

$$|f(0)| \le (2t)^{-1} \int_{-t}^{t} |f(x)|\,dx + \int_{-t}^{t} |f'(x)|\,dx.$$

This follows from $f(0) = f(x) - \int_0^x f'(u)\,du$ and averaging. The same inequality holds with the norm of a vector-valued function. Iterate the corresponding two operators in each of d_0

coordinates of a box of side $2t$; by Cauchy–Schwarz and the inequality $(\sum_I a_I)^2 \le 2^{d_0}\sum_I a_I^2$, obtain

$$\|f(0)\|^2 \le 2^{d_0}\sum_{I\subseteq\{1,\dots,d_0\}} (2t)^{-d_0+2|I|}\int_{[-t,t]^{d_0}} \|\partial_I f(z)\|^2\,dz.$$

Only derivatives of order at most d_0 appear, one per coordinate. Apply this to the functions obtained by smoothing F with an L^2 test function; the derivatives are controlled by (21.8). Choosing the small coordinate box comparable in size to P^{-6}, as in the source, costs a power of P of order s_0^2. The ambient rank affects derivative sizes but not the number of derivatives.

What converts these inequalities into the actual open-cell bound is the source's change of variables through the frame Gram map. Specifically `le_lintegral_of_gramSection` of the real-place cell calculation supplies the Jacobian proportional to $c_G\,\mathrm{disc}(Z)^{k-1}$, with Haar normalization c_G fixed by the Gram normalization. Together with the chart and near-coordinate estimates, `lintegral_gramSection_integral_le` in `New/RealPlace/SobolevCell.lean` is the formal-source input with the following conclusion: for a full Bessel function satisfying Schur bound with parameter $P \ge 8r$, frame Gram discriminant at least $c > 0$, and the specified chart bounds on the P^{-6} box, the integral of

$$\left\|\int \phi(y)F(\gamma_Z(uh)^{-1}y)\,dy\right\|^2$$

over the compact-factor variable h and the indicated unipotent cell variables u ($\|u\| \le P^{20}$), multiplied by $c_G c^{k-1}$, is bounded by $P^{2000s_0^2}D\int|\phi|^2$ when ϕ is supported in the P^{20} size region and D is the Schur mass bound. This is the chart-dependent inequality used by the `BesselSystem.open_cell` clause of Section 20.5. Our box estimate explains the required exponent scale and the order of differentiation; it does not on its own identify c_G or verify the boundedness of every Gram section.

21.9 A finite-subset version of adelic Poisson summation

We give a direct proof of the inequality actually used in the global sieve. Its hypotheses and conclusion differ from classical Poisson equality for a Schwartz function. For a number field E let $\mathbb{A}_E$ carry the self-dual measure relative to the usual trace character, so that the quotient $\mathbb{A}_E/E$ has measure one. The diagonal copy of E is a discrete cocompact subgroup and its annihilator is E. These structural properties are the background `WeilData` in the formal source. For a nonnegative real-valued L^1 function F on $\mathbb{A}_E$, define $\widehat{F}(\xi) = \int F(x)\overline{\psi(\xi x)}dx$.

Lemma 21.8 (One-sided Poisson)**.** *Assume F is lower semicontinuous at each member of a finite subset V of E. Then*

$$\sum_{t\in V} F(t) \le \sum_{\xi\in E} |\widehat{F}(\xi)|.$$

More precisely, if the nonzero coefficients have summable bounds $B(\xi)$, the right side may be replaced by $\int F + \sum_{\xi\neq 0} B(\xi)$.

Proof. Use the quotient Haar probability and the nonnegative periodization $P(x) = \sum_{t\in E} F(x+t)$, which is defined almost everywhere and belongs to $L^1(\mathbb{A}_E/E)$ by Tonelli. Its Fourier coefficients are $\widehat{F}(\xi)$, by Fubini and the annihilator identification. If the sum of their absolute values diverges the claim is immediate. Otherwise the Fourier series converges absolutely and uniformly to a continuous function P_0 and equals P almost everywhere. This last assertion follows from uniqueness of Fourier coefficients on a compact abelian group: characters span a uniformly dense subalgebra of the continuous functions, by conjugation

stability and separation of points, so the measure with density $P - P_0$ is zero. Given $\delta > 0$, lower semicontinuity gives a neighborhood O of zero on which $F(t + x) \geq F(t) - \delta$ for all $t \in V$. For almost every x in this neighborhood,

$$\sum_{t \in V} (F(t) - \delta) \leq P(x) = P_0(x) \leq \sum_{\xi} |\widehat{F}(\xi)|.$$

Nonempty open subsets of the quotient have positive measure. Now let $\delta \to 0$; the same argument with arbitrarily large bounds applies if some $F(t) = \infty$. □

The source supplies this inequality in `Lib/Adelic/Adeles/OneSidedPoisson.lean`. Finite subsets and lower semicontinuity permit the original orbital functions to be used without assuming differentiability at zero or at nonunit elements. For example a single upward spike at a rational point invalidates the inequality without lower semicontinuity: it contributes to the left side but is invisible to the Fourier transform. A downward spike is harmless for a one-sided bound.

We verify the Jacobian and the frequency support in the setting of the sieve. Let T be the finite auxiliary set of finite places, A_T the product of E_w at $w \in T$, and

$$U_T = \{x \in \mathbb{A}_E : x_w \in \mathcal{O}_w \ (w \notin T)\}.$$

This is open in the restricted product topology. For a function H on $\mathbb{C}^f \times A_T$ and positive or nonzero real scales l_{vi}, form

$$F(z) = \mathbf{1}_{U_T}(z) H((l_{vi} z_v)_{v|\infty}, (z_w)_{w \in T}).$$

Multiplication by l_{vi} on $\mathbb{C}$ has absolute real determinant l_{vi}^2. Off T, integrating the compact indicator introduces its Haar volume. Thus for the normalized measures on the local factors in the source,

$$\int F(z)\, dz = c_i \int H(y)\, dy, \qquad c_i = \frac{\mathrm{vol}(\widehat{\mathcal{O}})}{\prod_{v|\infty} l_{vi}^2}.$$

The finite-source convention is that the local integral volumes of the unit rings at T are one; equivalently one may absorb them into the normalization of H. In a nonzero-frequency integral the same change of variable replaces the infinite frequency by its quotient by l_{vi} (since l_{vi} is real). The off-T Fourier integral of an indicator of a compact open additive subgroup is zero outside its annihilator: translating by an element of the subgroup that has a nontrivial character multiplies the integral by a phase different from one. With the self-dual character divided by the source's different-normalizing element, the annihilator of the off-T integral lattice is the integral lattice. Thus the rational frequencies that remain are T-integers. The constants κ_v and g in the source convert the usual trace character to the local Bessel root characters; those constants occur in the frequency arguments, not in the absolute Jacobian factor. This is the measure and frequency content of `measurePreserving_scale_evalPlaces`, `extendToAdeles` and `one_sided_poisson` in `New/Filling/GlobalSieve.lean`.

21.10 An explicit induction for the sieve system

For the precise kernels defined in Section 20.13, we work with a finite alphabet A of admissible rational units (the elements and the relevant inverses integral away from T). The head maps $H_i(p, t)$, for $p \in A^i$ and next entry t, place these entries into the local product scaled by l_{va}, filling later entries with the chosen units. They are continuous on the unit locus. The bottom term is bounded by E_0; to simplify notation let $\lambda_i(p)$ be the nonnegative weight of a head of

length i, in the normalized coordinates. The scaling and filling maps are understood at all levels. We derive the finite-set induction used in `one_sided_poisson`; abstracting from those coordinate choices, it has two distinct error terms.

First, the positivity hypothesis at the top of a sieve system means that, for any finite family p_a with a common last entry, $\sum_{a,b} \overline{z}_a J_{s_0}(p_a, p_b) z_b \geq 0$. It is preserved by taking products over places: represent each matrix as the Gram matrix of a family e_a of vectors, and similarly f_a for a second matrix. The new Gram vectors $e_a \otimes f_a$ have pairings equal to the product of the original entries. Integration at zero frequency preserves the property for any fixed coefficients z_a, by linearity and integrability. It also preserves lower semicontinuity of these nonnegative sums, by Fatou's lemma. More explicitly, if the continuity hypothesis at the top is restricted to the open unit locus, a path of common head entries approaching a unit can be taken within this open set; the other integrated coordinates are units almost everywhere or replaced by the map `unitOrOne`. Iterating Fatou gives the required inequality even when the lower kernels have not been shown continuous separately. This proves the positivity and lower-semicontinuity content of `SieveSystem.pi` and `SieveSystem.nonneg_and_lowerSemicontinuous` in `New/Filling/EichlerFlag.lean`.

For a finite $V \subseteq A$ put

$$S_i(V) = \operatorname{Re} \sum_{t \in V} \sum_{p,q \in A^i} J_{i+1}(H_i(p,t), H_i(q,t)).$$

The summand after summation over heads is nonnegative, as just proved. We take the uniform nonzero-frequency assumption of the source in the form

$$\sum_{y \in X} \beta(\operatorname{ratPoint}(c_v / l_{vi}, y)) \leq \varepsilon, \tag{21.20}$$

for every finite set X of nonzero rational T-integers, for each i in the range, and for each family of admissible scale constants c_v (in `one_sided_poisson`, $|c_v|^{-1} \leq |\kappa_v|/g$). Here β is the product frequency majorant. The head sum assumption is

$$\left(\sum_{p \in A^i} \lambda_i(p) \right)^2 \leq \Gamma \prod_{a<i} c_a. \tag{21.21}$$

The zero-frequency step integral descends to J_i; at each nonzero frequency the sum over heads is at most the majorant times $(\sum \lambda_i)^2$, by triangle inequality. Fubini justifies this identity and the bound for a finite sum of head pairs: no convergence in the number of heads is being used. Lemma 21.8 and the Jacobian computation give

$$S_i(V) \leq c_i \operatorname{Re} \sum_{p,q \in A^i} J_i(p,q) + c_i \varepsilon \Gamma \prod_{a<i} c_a. \tag{21.22}$$

At $i = 0$ the sum on the right has one entry and the bound is $c_0(E_0 + \Gamma\varepsilon)$. For $i \geq 1$, split the right-hand sum according to the last entries of p and q. The equal-entry part is exactly an $S_{i-1}(A)$, not a sum of norms of individual head vectors. For the unequal-entry part the difference frequency at level i is $g^{-1}(t^{-1} - u^{-1})$. For fixed t and variable $u \neq t$, the differences are nonzero and distinct. They are rational T-integers before the stipulated normalization of the character. With

$$L_t = \sum_{p \in A^{i-1}} \lambda_i(p,t),$$

the absolute value of the unequal-entry contribution is bounded by

$$\sum_{t \neq u} \beta(g^{-1}(t^{-1} - u^{-1})) L_t L_u.$$

One can estimate this by the row and column bounds for the nonnegative frequency matrix: each row sum is at most ε by (21.20), and so is each column sum, because the differences there too are nonzero and distinct. Since $2L_tL_u \le L_t^2 + L_u^2$, the bound is $\varepsilon \sum_t L_t^2 \le \varepsilon(\sum_t L_t)^2$. Combining this with the Poisson error gives

$$S_i(V) \le c_i S_{i-1}(A) + 2\varepsilon\Gamma \prod_{a\le i} c_a.$$

Together with the base case this yields

$$S_i(V) \le \left(\prod_{a\le i} c_a\right)(E_0 + (2i+1)\Gamma\varepsilon)\,.$$

This is the induction and the coefficient $2i+1$ in `one_sided_poisson` of `New/Filling/GlobalSieve.lean`. The complex forms $J_i(p,q)$ in the source include the head padding and unit replacements; the displayed shorthand drops those redundant arguments. The term $2i+1$ accounts for one Poisson error at every floor and a different error each time the previous floor is split by its last entries. Bounding each Gram entry in absolute value at the top would lose the exact identity between floors, which supplies the leading term E_0.

21.11 Frequency summation and the product formula

The bound (21.20) would be useless if it contained the size of A or the sizes of denominators of its members. The damping exponent 6 is designed to avoid either dependence after product-formula cancellation.

Let $y \in E^\times$ be integral outside the conjugation-stable set T. In a local field write $|y|_w$ for the module normalized by the cardinality of the residue field, and $[y]_T = \prod_{w\in T} |y|_w^{-1}$ for the generalized index. For a complex place v, $|v(y)|$ denotes the ordinary absolute value. The product formula gives

$$\prod_{v|\infty} |v(y)|^2 = \prod_{w \text{ finite}} |y|_w^{-1} \ge [y]_T.$$

The inequality uses integrality away from T, not integrality at T. In particular the potentially small or large quantity $[y]_T$ can be nonintegral. Write $L_K(u) = u^K \min(1,u)^6$ for the damped power. At scales satisfying $|c_v|^{-1} \le |x_v| \le B$, $B \ge 1$, we get

$$\prod_v L_K(|c_v v(y)|^{-1})\,[y]_T^{K/2} \le \prod_v |x_v|^K \min\left(1, \frac{B}{|v(y)|}\right)^6.$$

The remaining local factor satisfies

$$\min(1, B/t)^6 \le \frac{(1+B^2)^3}{(1+t^2)^3} \quad (t>0).$$

If $t \le B$, the right side is at least one. If $t > B$, use $B^2(1+t^2) \le (1+B^2)t^2$ and cube. Thus each local term is controlled by the inverse cube of the natural two-dimensional vector norm $1+|v(y)|^2$, with the factor B^6 separated.

The denominator index $\mathrm{den}(y)$ is the index of the ideal $\{z \in \mathcal{O}_E : zy \in \mathcal{O}_E\}$, with the finite contribution outside the stipulated places taken according to the lattice convention. The point y determines a rational line in E^2 via $(1,y)$, and the lattice on that line is formed by multiplying $(1,y)$ by the denominator lattice. If $y = a/b$ is rational in lowest terms, this is simply the primitive vector (b,a), whose length squared is b^2+a^2; indeed $b^6(1+|a/b|^2)^3 = (a^2+b^2)^3$.

The corresponding sum over distinct lines converges: in concentric integer shells of maximum coordinate n, the number of pairs is at most $8n$ and each summand is at most n^{-6}. This illustrates why the same denominator power and the same infinite-place damping are used, rather than unrelated tapering functions at the two kinds of places.

The general CM assertion also needs the lattice covolumes and ideal classes. We retain as formal-source input the rank-one sublattice discriminant sum `tsum_sublatticeDisc_inv_le` and discriminant comparison `sublatticeDisc_mul_index_sq_le` used in the proof of `sum_frequencies_le` in `New/Filling/GlobalSieve.lean`. In that declaration, under WeilData and conjugation stability of T, the precise consequence is

$$\sum_{\substack{y\in X\\ y\neq 0}} \frac{[y]_T^{K/2}}{\operatorname{den}(y)^6} \prod_v L_K(|c_v v(y)|^{-1}) \le \left(\prod_v |x_v|^K (bB^2)^3\right) (b\,\mathrm{rd}_E)^{3f}$$

for any finite subset of nonzero T-integers. The explanation above proves the product-formula and damping portions of the inequality; the full sublattice discriminant sum is more than ordinary Euclidean lattice-point counting, because one must count rational lines with nonprincipal denominator ideal.

In the source $K = k - s_0^7$, and the finite multiplier defining `frequencyGain` is exactly $[\xi]_T^{K/2} \operatorname{den}(\xi)^{-6} b^{s_0^9 f}$. Thus the large power K disappears from the summation at the cost of the explicit powers of the scales x_v. The other weight bound (21.21) comes from summing over the successive head products in `sq_sum_globalWeight_le` (same file). Those products can be used as summation coordinates because knowing all successive products and the leading unit recovers each entry by division; denominators are again accounted for by the integral-lattice condition. The numerical comparison with the parameters of `Filling.IsAdmissible`, including these head bounds and frequency sums, is the remaining formal-source input `sieve`. It gives the error $(2h+1)\Gamma\varepsilon \le b^{-6kf-10}/d_\infty$ and the leading constant of Proposition 20.13. The part now proved directly is the one-sided Poisson argument and its induction, with the correct covolume factor; it requires the uniform frequency and head sums just described, not merely absolute summability for each fixed rank.

21.12 Finite packets, their normalization, and the global comparison

We expand the finite-place mechanism in the same notation as Section 20.13. At a finite product of local fields, a compact-open level permits a locally constant vector function to be integrated by counting cosets. Here is the elementary step used in constructing the top kernel. If f is continuous and locally constant on a compact open subset D of a finite product of nonarchimedean fields, cover D by finitely many translates of compact open additive subgroups on each of which f is constant; intersect those subgroups. The result is a compact open subgroup L with D a finite union of its cosets, on each of which f is constant. Consequently

$$\int_D f(t)\,dt = \operatorname{vol}(L) \sum_{t+L\subset D} f(t).$$

For a finite family of functions the same subgroup works, by further finite intersection. If the functions are indexed by finitely many cosets of a compact packet group, the same argument applies simultaneously to their pairings. This is the basic step-function mechanism in `New/Filling/FiniteSystem.lean`. It identifies the normalized top floor as an averaged Gram matrix, not just the indicator of an algebraic equation over a residue field.

There is an independent counting problem for the mass at the bottom of the tower. In the lattice model the quadratic condition for a last-vector packet is of the form

$$Ny^{*}y = t\bar{t} - N^{2}.$$

This follows by pairing the last frame vector with itself in the changed basis and substituting its coordinates in the positive middle space. Thus the number of choices for the k-component vector in the packet is subject to a Hermitian norm condition. For a finite field of square cardinality q^2, the equation $\sum_{j=1}^{k} x_j\overline{x_j} = a \neq 0$ has $q^{2k-1} - (-1)^k q^{k-1}$ solutions: for $k = 1$, each nonzero a is the norm of exactly $q + 1$ elements. If Z_k and N_k denote the counts at zero and at a fixed nonzero value, respectively, then $Z_k + (q-1)N_k = q^{2k}$ and adding one coordinate gives

$$N_{k+1} = N_k + (q+1)(Z_k + (q-2)N_k) = (q+1)q^{2k} - qN_k.$$

This recursion proves the formula by induction, and subtraction gives the corresponding zero count. It demonstrates the high-dimensional equidistribution in the packet counting step, but it does not prove the particular local mass comparison when congruences, ramified factors and higher floors are imposed. The latter uses the weighted quadric integral and the passage between successive floors in the source.

Precisely, `Filling.IsFillingLevel.exists_finite_sieveSystem` of `New/Filling/FiniteSystem.lean` constructs an intermediate packet group Z containing all level stabilizer elements whose reductions belong to the prescribed fixing set, and a compact open additive subgroup L of the finite-local product. Its kernels have the form

$$J_j^{\mathrm{fin}}(p,q) = \begin{cases} \mathrm{floorForm}_j(p,q;Z)/\mathrm{massRatio}(Z), & j < s_0, \\ \mathrm{topKernel}(L,Z;p,q)/(\mathrm{vol}(L)\mathrm{massRatio}(Z)), & j = s_0. \end{cases}$$

The upper kernel is a Gram kernel at a common top entry. The bottom bound in that declaration is

$$(1 + b^{-6})/(b^{6k})^{|S_g|}.$$

The head weights are `finHeadWeight`, and the nonzero frequency majorant is `frequencyGain` in Section 21.11. The exact floor and mass comparison, including `sum_blocks_le_topKernel`, is a formal-source input. The factor $1/\mathrm{vol}(L)$ compensates for the additive coset volume in the integration identity. The factor $1/\mathrm{massRatio}(Z)$ accounts for the measure of the fixed-vector subgroup; an arbitrary upper bound for its cardinality could not be used instead, because the subsequent product with the archimedean leading terms must give the stated main-energy normalization.

To compare the local orbital integrals with the rational amplitude, one first removes the compact unipotent class cutoffs by unfolding, then collects all rational terms lying in the prescribed finite cosets. For a convergent sum, partitioning the rational orbits into those cosets is just reindexing the sum. The height restrictions and quotient cutoff estimates ensure convergence for the actual amplitude. The identity `IsFillingLevel.correlation_smul_eq_cosetAmplitude` in `New/Filling/GlobalToLocal.lean` gives the exact conversion. For a term s at the last floor, local head q obtained by reading a rational head p at T, and packet cosets a, b satisfying the level-congruence relation, it equates

$$\mathrm{correlation}_{Z,q}(a,b) \otimes \mathrm{besselCorrelation}_{\infty}(\gamma) = D_{s_0}(q,1)\ \mathrm{vol}(K(g))\,\mathrm{cosetAmplitude}_T(\gamma).$$

The left arguments in the source include the inverse of the local term position of s, and γ denotes the rational Weyl-head translate of s. This is a formal-source input: proving it

requires the changes of variables in the exact integral normalization at all places. It specifies which mass factors survive the global comparison. Taking the product sieve system and applying Section 21.10 to the common-cell entries of that amplitude is therefore a comparison of normalized Gram matrices with the prescribed bottom constant, rather than a pointwise comparison of positive scalar functions.

21.13 Degenerate terms and the accumulation of energy errors

For completeness we supply two small calculations that govern the nonmain cell estimates. The first is character compatibility. Consider an unfolded integral on a double orbit with left and right Bessel characters. Translating within their common stabilizer by s leaves the cell and measure unchanged, but changes the integrand by the quotient $\eta(s)$ of the two characters. If $\eta(s) \neq 1$, the integral I must satisfy $I = \eta(s)I$, hence $I = 0$. For rational orbit sums with unipotent class cutoffs the same argument applies after averaging over the stabilizer, because the sum of the translates of the cutoff equals one. The precise source identity, using stable sets of rational terms and the hypotheses of admissibility, is `Filling.amplitudeSum_eq_zero_of_not_charsAgree` in `New/Filling/CrudeEnergy.lean`. We use the cutoff-invariance statement there; pointwise invariance of an arbitrarily chosen cutoff is not assumed.

To check the Hermitian geometry behind the compatible degenerate terms, use the convention that H is conjugate-linear in its first variable. For $H(x,x) = H(x,v) = 0$ and $\alpha + \overline{\alpha} = 1$ the transformation

$$T(y) = y + xH(v,y) - (v + \alpha H(v,v)x)H(x,y)$$

preserves H. Indeed write $l(y) = H(x,y)$, $m(y) = H(v,y)$ and $a = H(v,v)$ real. Then $T(y) = y + x(\overline{m(y)} - \alpha a\overline{l(y)}) - vl(y)$. Expanding $H(T(y),T(z)) - H(y,z)$, the cross terms $\overline{l(y)}m(z)$ and $\overline{m(y)}l(z)$ cancel, leaving $a(1-\alpha-\overline{\alpha})\overline{l(y)}l(z) = 0$. Its difference from the identity is nilpotent, since its image is contained in the span of x, v, sends x to zero and sends v to a multiple of x. It is thus an invertible unitary transformation. Choosing x in the appropriate shared flag gives the source's elementary Eichler map. The precise placement in the Bessel group is `exists_mem_besselU_eq_eichlerMap` in `New/Filling/EichlerFlag.lean`; the paired-coordinate propagation with compatible characters is `hasCommonFlagVector_of_le` in `New/Filling/CrudeEnergy.lean`. Their flag-dependent consequence `Filling.CharsAgree.exists_hasLevel` is a formal-source input: a rational term with compatible characters and discriminant zero has a level below the top open cell in the sense of `Filling.HasLevel`. The explicit calculation shows why preservation of the indefinite form in those test transformations does not itself impose any restriction on their Bessel characters. It is the additional character compatibility that supplies the relation between the two flags.

The second calculation is the handling of many error strata. The amplitude spaces in the energy definition are Hilbert spaces (with the common positive scalar unipotent-covolume normalization). For $\delta > 0$,

$$\|A + B\|^2 \leq (1+\delta)\|A\|^2 + (1+\delta^{-1})\|B\|^2. \tag{21.23}$$

This is obtained from $2|\langle A, B\rangle| \leq \delta\|A\|^2 + \delta^{-1}\|B\|^2$. It applies to integrable families by taking their Hilbert space norms; it does not require orthogonality of the strata.

We recall the actual pair-error bound from the source. Let C satisfy the prescribed size inequalities (at least the Bessel energy, with stronger bounds τ^{20} times that energy at level zero and τ^{12} times the main energy divided by X^k if the retained set of links omits one). The

input `Filling.energy_le_of_level_pairs` in `New/Filling/CrudeEnergy.lean` says that for a stable set of a common level j, the cut energy is at most

$$w + \frac{C}{b^{30}4^{j^{\mathrm{rev}}}},$$

if the retained coherent link energy divided by the square of the unipotent covolume is at most w. The reversed level index is the distance from the top of the finite tower. This inequality uses the summable pair weights, the exact assertion `sum_pairWeight_le` in `New/Filling/BlockBound.lean` being their total mass at most $2\tau^2$. Incoherent links contribute the arithmetic error rather than being discarded as orthogonal. Those lattice and pair estimates are retained as formal-source inputs.

To sum all levels strictly below a floor, begin with zero energy below the first level (the incompatible terms vanish by the character identity). If the error budget at one floor is E, the previous floor budget is $E/4$. The new single-level budget is likewise $E/4$ by the pair bound with no retained coherent link energies. Apply (21.23) with $\delta = 1$ to get $2(E/4)+2(E/4) = E$. This proves the induction of `Filling.energy_le_of_level_lt` with the geometric factor $4^{j^{\mathrm{rev}}}$; there is no factor proportional to the number of levels.

One still needs a bound for replacing the locally cut functions by the full ones after global unfolding. The formal-source estimate `energy_sub_cut_le` in `New/Filling/SchurTest.lean` bounds this difference energy by b^{-30} times the Bessel energy. It uses the quotient tail and transported majorants, not just the local inequality (21.19); sums of translates at distinct rational points would prevent the latter alone from proving the former. Now the cut top-cell energy is at most $w + Cb^{-30}$ and the lower-cell cut energy at most Cb^{-30} by the preceding induction. A first use of (21.23) with $\delta = b^{-4}$ bounds the total cut energy by

$$(1 + b^{-4})w + (2 + b^4 + b^{-4})Cb^{-30}.$$

A second use with the same δ, for full minus cut, shows that the full energy is at most

$$(1 + b^{-4})^2 w + \Big((1 + b^{-4})(2 + b^4 + b^{-4}) + 1 + b^4\Big)\, Cb^{-30}.$$

Since $b \geq 1000$, the error coefficient is less than b^{-12}. This gives precisely the combination in `Filling.energy_le_of_linkEnergy_le` of `New/Filling/CrudeEnergy.lean`. The retained pair estimate, cutoff transfer, flag-level theorem and link/coherent comparison are substantial formal-source inputs. The geometric summation and energy combination are elementary deductions. In particular the local Fourier sieve supplies the only non-negligible w; the other cells are controlled separately with large exponents. Substituting the coherent bound into this deduction gives the sharp energy bound of Section 20.11 and the subsequent `energy_fixingSet_le` calculation, with the arithmetic numerical comparisons of the specialist chapter left unchanged.

Chapter 22

Quantitative Bessel construction and the arithmetic of the filling datum

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 25, 26, 27 and the final audit in Chapter 57.

The filling argument is unusually sensitive to the normalization of local integrals and the exact exponent in their bounds. The proof of the first and second moments in Chapter 20 and the local mechanism in Chapter 21 left a number of these statements as formal-source inputs. In this chapter we reconstruct further parts of those arguments: the explicit Gaussian approximation of the discrete-series coefficient, the simultaneous removal of compact directions and the phase-cancelling realization of the Bessel subgroup, the arithmetic level and its unipotent covolume, the elimination of vanishing-discriminant terms by a flag argument, the height summation in the global sieve, and the finite-group covariance required for the moments. We maintain all quantitative bounds needed by those arguments. There remain local Fourier-floor estimates, some uniform arithmetic pair-counting estimates, and the exact global packet comparison; these are separately identified rather than inferred from the results proved here. Standard representation-theoretic results used here include the identification of the normalized lowest-type coefficient with the Flensted–Jensen integral, its square-integrable representation, the Schur orthogonality and Harish-Chandra degree formulas. These are logically different from the particular quantitative estimates proved below.

Throughout the chapter the hyperbolic rank is $h = \varrho$, $n = h + 1$, the dimension of the definite part is k, the full rank is $r = k + 2h + 1$, the number of complex places is f, $g = q_0 q_1$ is the product of the two split rational level primes, and N is the separate split prime used for the marked lattice. In the discrete-series calculations, where we use matrices of varying sizes, we use d for a matrix rank, not for the formal degree, which we denote by d_λ or d_∞ for the product of degrees. For a matrix A let A^* denote its conjugate transpose, $\|A\|_2$ its Frobenius norm, and $\|A\|_{\mathrm{op}}$ the Euclidean operator norm. This distinction is used throughout the estimates.

22.1 The compact integral and the change of splitting

Let $J = \mathrm{diag}(\epsilon_i)$, where each $\epsilon_i = \pm 1$, and let $G_J = \{g : g^* J g = J\}$ and $K = G_J \cap U(d)$. The doubled Harish-Chandra parameter ℓ has distinct entries, with $\ell_i + d - 1$ even. Put $\lambda = \ell/2$,

let ρ be the half sum of roots positive for the order of λ, and ρ_c the half sum of the compact positive roots (the roots $e_i - e_j$ with $\epsilon_i = \epsilon_j$). Thus

$$\mu = \lambda + \rho, \qquad \Lambda = \mu - 2\rho_c$$

are integral. In particular μ has the same strict order as ℓ. The weight Λ is dominant in each compact block for this order. The parity and distinctness here are essential: the compact type with highest weight Λ is an honest representation, not merely a character on a cover.

For an integral weight w and square matrix A write $m_w(A)$ for the product of principal minors used in the source:

$$m_w(A) = \det(A)^{-a} \prod_{t=-a+1}^{a} \det A_{\{i:w_i \geq t\}}, \qquad a = \sum_i |w_i|.$$

Empty principal minors have determinant 1. This is independent of enlarging a. If all the principal minors in the product are invertible it obeys

$$m_{-w}(A) = m_w(A^{-1}), \qquad m_{w+c}(A) = m_w(A)\det(A)^c. \tag{22.1}$$

For the first identity apply Jacobi's complementary-minor identity at each threshold, pairing threshold t with the complementary threshold $1 - t$; the determinant factors compensate because $(A^{-1})_S$ has determinant $\det A_{S^c}/\det A$. The second follows by shifting the thresholds.

Let a be invertible block diagonal for J and z have strictly positive Hermitian part, so that every principal minor of $u^* z u$ is nonzero for $u \in K$. Put

$$F_\ell(a, z) = \frac{1}{Z_\Lambda} \int_K \frac{m_\Lambda(au)\overline{m_\Lambda(u)}}{m_{-\mu}(u^* z u)}\, du, \qquad Z_\Lambda = \int_K |m_\Lambda(u)|^2\, du.$$

Here the Haar measure on K is probability. This is `HCParam.flenstedJensen` in `Lib/DiscreteSeries/FlenstedJensen/RepresentationDefs.lean`. If $g \in G_J$, write a_g for its block diagonal part. The representation-theoretic identification used in the source defines its coefficient by

$$\psi_\lambda(g) = F_\ell(a_g, a_g^{-1} g).$$

We rely on the lowest-type coefficient theorem in this identification, not on an analytic continuation assertion about arbitrary highest-weight functions.

We prove directly the splitting property of the integral. This proof is useful because the pointwise Gaussian calculation uses a different splitting in which the large weights occur only in the denominator.

Lemma 22.1 (Change of splitting)**.** *Let g be a fixed complex invertible matrix. On the set of block diagonal matrices a such that $g^{-1}a$ has strictly positive Hermitian part, the value $F_\ell(a, a^{-1}g)$ is constant.*

Proof. The domain is convex: positivity of the Hermitian part of $g^{-1}a$ is a convex condition and implies that a is invertible. Positivity is equivalent for an invertible matrix and its inverse, by conjugating the Hermitian part with the inverse. Hence both arguments of the integrand are valid on the domain. Rewrite it, by (22.1), as

$$\Phi(a, u) = \frac{m_\Lambda(au) m_\Lambda(u^{-1})}{m_\mu(u^{-1} g^{-1} a u)}.$$

The integral of this function is locally holomorphic in the block diagonal matrix a: all denominators are nonzero, and on the compact set K boundedness of the derivatives on a

sufficiently small neighborhood permits differentiation under the integral. For the same reason the integrand extends holomorphically in u to a neighborhood of K in its complexified group.

Let X be a block diagonal variation of a and write $Y = u^{-1}a^{-1}Xu$. At fixed u the derivative in direction X of Φ is the sum of the derivatives in directions $auE_{ij}u^{-1}$ multiplied by Y_{ij}, over pairs with $\epsilon_i = \epsilon_j$. If $\ell_i \geq \ell_j$ then right multiplication by $1 + tE_{ij}$ is an upper triangular elementary matrix (ordering the weights in descending order), and the principal-minor transformation law gives the derivative

$$D_{auE_{ij}u^{-1}}\Phi(a,u) = \delta_{ij}(\Lambda_i - \mu_i)\Phi(a,u).$$

Indeed for an upper triangular elementary right factor each high-weight principal submatrix is multiplied on the right by its restricted triangular factor; the diagonal case multiplies by $(1+t)^{\Lambda_i - \mu_i}$.

Suppose instead that $\ell_i < \ell_j$. A lower triangular unipotent left factor leaves the high-weight principal minors unchanged, since the principal submatrix of each initial segment depends only on the corresponding rows of the left factor. Under $u \mapsto u(1+tE_{ij})$, the left factor $(1+tE_{ij})^{-1}$ therefore drops out of both $m_\Lambda(u^{-1})$ and the denominator. The derivative of Φ in that u-direction equals $D_{auE_{ij}u^{-1}}\Phi$. In the same variation, Y is conjugated by $1+tE_{ij}$, so the derivative of its ij-entry is $Y_{ii} - Y_{jj}$. Haar right invariance implies

$$\int_K \frac{d}{dt}\Big|_{t=0} \big(\Phi(a, u(1+tE_{ij}))Y_{ij}(u(1+tE_{ij}))\big)\, du = 0.$$

This follows first for real Lie algebra directions and hence for E_{ij} by complex linearity of the holomorphic derivative (the Lie algebra of each complexified compact block is the complexification of that of the compact block). We conclude

$$\int_K Y_{ij} D_{auE_{ij}u^{-1}}\Phi\, du = -\int_K \Phi(Y_{ii} - Y_{jj})\, du \quad (\ell_i < \ell_j).$$

Sum over the indicated opposite triangular directions. For each i the coefficient of Y_{ii} in $\sum_{\ell_i<\ell_j,\epsilon_i=\epsilon_j}(Y_{ii} - Y_{jj})$ is $-2(\rho_c)_i$. Thus their contribution to the derivative is $+\int_K \Phi \sum_i 2(\rho_c)_i Y_{ii}\, du$. The diagonal contribution from the first formula is $\int_K \Phi \sum_i (\Lambda_i - \mu_i) Y_{ii}\, du$, precisely its negative. The total derivative vanishes along every variation X. Constancy follows from convexity. □

This is the argument of `flenstedJensen_eq_of_accretive` and `dsCoeff_eq_flenstedJensen` in `Lib/DiscreteSeries/FlenstedJensen/FlenstedJensenInvariance.lean`. In particular the half sum of the compact positive roots accounts exactly for the derivative of the conjugating matrix in the compact integral. Replacing Λ with λ would not give splitting invariance.

The following useful consequence isolates the triangular shifts. If L is strictly lower triangular in the order of descending ℓ and block diagonal for J, with X and $(1-L)X$ accretive, then

$$F_\ell(1, (1-L)X) = F_\ell(1, X). \tag{22.2}$$

Indeed apply Lemma 22.1 to $(1-L)X$, using the splitting $a = 1 - L$, and use $m_\Lambda((1-L)u) = m_\Lambda(u)$. Taking conjugate transpose shows the analogous invariance under a right factor $1 - L^*$, since $F_\ell(1, X^*) = \overline{F_\ell(1, X)}$. The adjoint statement is `flenstedJensen_shift_eq_conj` in `Lib/General/LinearAlgebra/FramedCoefficient.lean`.

22.2 Logarithmic principal-minor estimates by resolvents

The delicate feature of the next calculation is that the error must be proportional to the *squared commutator* rather than to a uniform cubic matrix error. This is what permits repeated or nearly equal compact weights. We give a proof using the derivative of the compressed logarithm.

Lemma 22.2 (Compressed logarithm with an absolute error)**.** *Let P, Q be orthogonal projections, $\|X\|_2 \le \delta \le 1/2$, and suppose*

$$\operatorname{tr}([P,Q]Y) = 0, \qquad \|[P,X] - Y\|_2 \le \delta c, \qquad \|Y\|_2, \|Q-P\|_2 \le c.$$

Then

$$\log\det(1+PX) = \operatorname{tr}(Q\log(1+X)) - \tfrac{1}{4}\operatorname{tr}(Y^2) + e, \qquad |e| \le 100\delta c^2,$$

where the logarithm of the determinant denotes the branch obtained by continuation along $1+tPX$, $0 \le t \le 1$.

Proof. The trace logarithm is valid because the operator norm of X is at most δ. Write X in blocks relative to P: its blocks are A, B, C, D. For $0 \le t \le 1$, put $U = (1+tA)^{-1}$ and $W = (1+tX)^{-1}$. The PP-block of the inverse satisfies

$$W_{PP} - U = t^2 U B W_{P^\perp P^\perp} C U.$$

This follows by multiplying the inverse equations, or twice using the equations for the off-diagonal inverse blocks. Differentiate the trace logarithms of $1+tPX$ and $1+tX$, compressed by P in the second case. The difference of their derivatives is $t^{-1}\operatorname{tr}(W_{PP} - U)$ (at $t = 0$ use the continuous extension). Hence

$$\operatorname{tr}\log(1+PX) - \operatorname{tr}(P\log(1+X)) = \int_0^1 t\,\operatorname{tr}(UBW_{P^\perp P^\perp}CU)\,dt. \tag{22.3}$$

The inverses have operator norm at most 2; their difference from the identity has operator norm at most $2t\delta$. Replace the three inverse factors in the integrand successively by 1 and use $|\operatorname{tr}(B'C')| \le \|B'\|_2\|C'\|_2$. The error is at most $14t\delta\|B\|_2\|C\|_2$ before multiplication by t. Thus the right side of (22.3) differs from $\frac{1}{2}\operatorname{tr}(BC) = -\frac{1}{4}\operatorname{tr}([P,X]^2)$ by at most

$$\tfrac{14}{3}\delta\|B\|_2\|C\|_2 \le 3\delta\|[P,X]\|_2^2.$$

We next estimate the trace incurred by replacing P with Q. Let $L = Q - P$, and separate its diagonal and off-diagonal blocks relative to P. The diagonal blocks are respectively negative and positive semidefinite. The sum of their trace norms is

$$\operatorname{tr}(P - PQP) + \operatorname{tr}(P^\perp Q P^\perp) = \|P - Q\|_2^2.$$

Their trace against $\log(1+X)$ has absolute value at most $2\delta c^2$. For the off-diagonal block part L_{off} one has

$$\operatorname{tr}(L_{\mathrm{off}}X) = -\operatorname{tr}([P,Q][P,X]).$$

The hypothesis on Y and $\|[P,Q]\|_2 \le \|L\|_2$ give an absolute bound δc^2. The logarithm series also shows

$$\|[P, \log(1+X) - X]\|_2 \le \sum_{m\ge 2} \delta^{m-1}\|[P,X]\|_2 \le 2\delta\|[P,X]\|_2.$$

Here the commutator of X^m is a sum of m terms with a single commutator of X. The off-diagonal part of a matrix has the same Frobenius norm as its commutator with P, so the remaining trace is at most $2\delta c\|[P,X]\|_2$.

Finally $\|[P,X]\|_2 \le (1+\delta)c \le 3c/2$. Replacing its trace square by $\operatorname{tr}(Y^2)$ costs at most $\|[P,X]-Y\|_2(\|[P,X]\|_2+\|Y\|_2) \le (5/2)\delta c^2$. Adding these estimates, the factor 100 is more than sufficient. □

In the source this is `exists_det_compress_eq_exp_logOneAdd`, `New/DiscreteSeries/MinorExpansion.lean`. In the proof the resolvent identity avoids a factor exponential in the matrix rank. All estimates use a Hilbert–Schmidt product only twice; other factors are measured in operator norm.

Proposition 22.3 (Triangularly shifted principal minors)**.** *Let $g = 1+X \in G_J$ with $\delta = \|X\|_2 < 10^{-3}$. There is an accretive y such that $\psi_\lambda(g) = F_\ell(1,y)$ and, for each projection P onto an initial segment of the* ascending *ℓ-order, each $u \in K$,*

$$\det((u^*yu)|_{\operatorname{im} P}) = \exp\left(\operatorname{tr}(P\log g) + \tfrac14\|[P,X]\|_2^2 + e_P\right),$$

with

$$|e_P| \le 200000\delta(\|[P,X]\|_2^2 + \|[P,u]\|_2^2).$$

Here the compressed matrix in the determinant is $Pu^*yuP|_{\operatorname{im} P}$.

Proof. Order the indices by descending ℓ. Let L and R be the strictly lower and upper parts, respectively, of the same-sign block part of X. Put

$$y = (1-L)(1+X)(1-R).$$

Since $\|L\|_2, \|R\|_2 \le \delta$, $\|y-1\|_2 \le 4\delta$, and both intermediate shifted matrices are accretive. Lemma 22.1 first yields $\psi_\lambda(g) = F_\ell(1,g)$, and (22.2) and its right-hand version give $F_\ell(1,y) = F_\ell(1,g)$.

Write $A=[P,X]$, $B=[P,L]$, $C=[P,R]$ and $A_o = A-B-C$. Thus A_o is the opposite-sign part of A because P is diagonal. The product rule for the commutator shows

$$\|[P,y-1]-A_o\|_2 \le 8\delta\|A\|_2.$$

Explicitly, $[P,y] = -B(1+X)(1-R) + (1-L)A(1-R) - (1-L)(1+X)C$; comparing the three terms to $-B, A, -C$ incurs respectively at most $(2\delta+\delta^2)\|B\|_2$, $(2\delta+\delta^2)\|A\|_2$, and $(2\delta+\delta^2)\|C\|_2$. Each commutator of a strictly triangular mask is a submatrix mask of A. The same estimate holds with P replaced by $1-P$ and A_o negated.

Put $P' = uPu^*$. The norm $\|P'-P\|_2 = \|[P,u]\|_2$. Consequently

$$\|[P',y-1]-A_o\|_2 \le 8\delta(\|A\|_2 + \|[P,u]\|_2). \tag{22.4}$$

Moreover P', P are same-sign block diagonal, whereas A_o is opposite-sign block off diagonal, so $\operatorname{tr}([P',P]A_o) = 0$. Lemma 22.2 applies with first projection P', second P, matrix $y-1$, radius 4δ, and $c = 4(\|A\|_2 + \|[P,u]\|_2)$. It yields

$$\log\det(1+P'(y-1)) = \operatorname{tr}(P\log y) - \tfrac14\operatorname{tr}(A_o^2) + e_0,$$

with $|e_0| \le 12800\delta(\|A\|_2^2 + \|[P,u]\|_2^2)$.

The complementary projection $P^\perp$ corresponds to a high-weight initial segment in the descending order. Its principal determinant is not changed by the lower triangular left or upper triangular right unit factors. The determinant of the full matrix is likewise unchanged. These equalities hold for the corresponding analytic logarithms: use the path $(1-tL)(1+tX)(1-tR)$, for $0 \le t \le 1$, to fix the branches. Apply the same compression lemma twice with both

projections $P^{\perp}$, first to X with $Y=-A$ and then to $y-1$ with $Y=-A_o$. With error bounds $100\delta\|A\|_2^2$ and $6400\delta\|A\|_2^2$ respectively this gives

$$\operatorname{tr}(P\log y)=\operatorname{tr}(P\log g)+\tfrac{1}{4}\operatorname{tr}(A^2-A_o^2)+e_1,\qquad |e_1|\le 6500\delta\|A\|_2^2.$$

To evaluate the quadratic term, put $W=(1+X)^{-1}$. The relation $g^*=Jg^{-1}J$ and the commutator rule for inverses imply $A^*=JWAWJ$. Since $\|W-1\|_{\mathrm{op}}\le\delta/(1-\delta)$, this gives

$$\|A^*-JAJ\|_2\le 3\delta\|A\|_2.$$

On the same-sign part $JAJ=A$ and on the opposite-sign part it equals $-A$. Taking traces against A and A_o respectively therefore proves

$$\operatorname{tr}(A^2)-2\operatorname{tr}(A_o^2)=\|A\|_2^2+\varepsilon,\qquad |\varepsilon|\le 9\delta\|A\|_2^2.$$

Combining the three formulas and using $\det(1+P'(y-1))=\det(Pu^*yuP|_{\operatorname{im}P})$ proves the assertion with the stated constant. □

This is `exists_principalMinor_eq_exp` in `New/DiscreteSeries/MinorExpansion.lean`. The exact sign of the quadratic term was determined by the signed adjoint identity. Without the triangular shifts, it would not in general have this sign.

22.3 Gaussian shape of the coefficient

For a real weight vector w set $Q_w(A)=\sum_{ij}|w_i-w_j||A_{ij}|^2$. As shown in Section 21.3, summing commutator squares over threshold projections gives Q_w. We now apply Proposition 22.3 to the weight $w=-\mu$, whose upper threshold sets are the required lower ℓ-sets.

Theorem 22.4 (Uniform near-identity Gaussian)**.** *Let $|\ell_i|\le L$, and let*

$$T\ge\max(1,d^2,d^2\log(d+L+1)),\quad \delta T\le 10^{-12},\quad \|g-1\|_2\le\delta.$$

Under the discrete-series hypotheses,

$$\left|\frac{\psi_\lambda(g)}{\exp(\operatorname{tr}(\operatorname{diag}(\Lambda)\log g))}-\exp(-Q_{-\mu}(g-1)/4)\right|\le\tfrac{1}{10}\exp(-Q_{-\mu}(g-1)/4)+e^{-100T}.$$

Proof. Write $X=g-1$, $q=Q_{-\mu}(X)/4$ and $p(u)=Q_\Lambda(u)/2$. Sum the logarithmic estimates for the shifted principal minors, using the determinant correction in $m_{-\mu}$. The trace of $\log y$ equals that of $\log g$. By the threshold identity and the formula for Λ,

$$\frac{\psi_\lambda(g)}{\exp(\operatorname{tr}(\operatorname{diag}(\Lambda)\log g))}=\frac{e^{-q}}{Z_\Lambda}\int_K|m_\Lambda(u)|^2e^{-\mathcal{R}(u)}\,du,\tag{22.5}$$

where $|\mathcal{R}(u)|\le 10^6\delta(q+p(u)+d^2)$. To verify the constant, the sum of the errors in Proposition 22.3 is bounded by $200000\delta(Q_{-\mu}(X)+Q_{-\mu}(u))$. Now $|\mu_i-\Lambda_i|\le d$, which implies $Q_{-\mu}(u)\le Q_\Lambda(u)+2d\|u\|_2^2=2p(u)+2d^2$. The trace incurred by replacing μ by Λ is at most $2d^2\delta$, using $\|\log g\|_2\le 2\delta$. These inequalities give the displayed bound on $\mathcal{R}$.

On the compact group the minor bound of Lemma 21.5 gives $|m_\Lambda(u)|^2\le e^{-p(u)}$. Schur orthogonality for the compact highest-weight coefficient gives $Z_\Lambda^{-1}=\dim V_\Lambda$, the dimension of the compact irreducible of highest weight Λ. For completeness, the function $m_\Lambda(u)$ is precisely the unit highest-vector coefficient: write the dominant weight as a sum of successive fundamental weights and a determinant weight on each sign block; take the tensor product

of the corresponding unit exterior highest vectors. The tensor highest vector generates the desired irreducible, and its coefficient is the product of the principal minors. The Weyl dimension formula, $\max|\Lambda_i| \le (L+3d)/2$, and the fact that there are at most $d(d-1)/2$ compact positive roots give

$$\dim V_\Lambda \le (L+4d+1)^{d(d-1)/2} \le e^{2T}.$$

If $|\mathcal{R}(u)| \le 1/20$, then $|e^{-\mathcal{R}(u)}-1| \le e^{1/20}/20 < 1/10$. Its contribution to the difference between (22.5) and e^{-q} is at most $e^{-q}/10$, using total probability of the normalized density. Otherwise the bound on $\mathcal{R}$ gives $q+p(u) > 49999T$. On this complementary set the absolute value of the integrand in the difference, including $Z_\Lambda^{-1}e^{-q}$, is at most

$$2e^{2T}\exp\big(-(1-10^{-6})(q+p(u))+10^{-6}\big) \le e^{-100T}.$$

Integrate against Haar probability. This proves the claimed estimate. □

This proves `dsCoeff_gaussian_shape` of `New/DiscreteSeries/GaussianShape.lean` using the Flensted–Jensen identification and ordinary compact representation theory. The absolute error is independent of the size of the Gaussian; this is what is needed in the restriction integral. The estimates concern the varying rank and weights, not merely the limit of a fixed representation at the identity.

22.4 Determinant decay and a small-ball bound

We give a further direct quantitative argument for the bounding functions used in the restriction calculation. Let γ be an integer such that $0 \le \gamma \le h_{\mathrm{gap}}$, where h_{gap} is the minimum allowed separation of λ_i, λ_j for indices of opposite sign. Thus any interval of the shifted weights μ of length γ, half open at one end, contains only indices of one sign. Let a_z denote the same-sign block part of a matrix z.

Lemma 22.5 (Decay from complementary and Fischer minors)**.** *For all $g \in G_J$,*

$$|\psi_\lambda(g)|^2|\det a_g|^\gamma \le 1, \qquad |\psi_\lambda(g)| \le (2/\|g\|_{\mathrm{op}})^\gamma, \qquad |\psi_\lambda(g)|^4 \le (1+\|g-a_g\|_2^2)^{-\gamma}.$$

Proof. First let $z \in G_J$ be positive Hermitian. Then $\det z = 1$ and $z^{-1} = JzJ$, so Jacobi's identity gives $\det z_S = \det z_{S^c}$ for every principal minor. Define $S = \{i : \mu_i + \gamma \ge t\}$ and $R = \{i : \mu_i \ge t\}$ for an integral threshold t. The difference $S \setminus R$ lies in one sign block. In that block Fischer's inequality gives

$$\det(a_z)_S \le \det(a_z)_R \det z_{S\setminus R}.$$

By the complementary-minor identity and then Fischer's inequality on the complementary set,

$$\det z_{S\setminus R} = \det z_{R\cup S^c} \le \det z_R \det z_{S^c} = \det z_R \det z_S.$$

Multiply these inequalities over a range of thresholds containing all weights, and include the determinant corrections. Since $\det z = 1$ and $z^{-1} = JzJ$, the result is

$$(\det a_z)^\gamma = \frac{m_{\mu+\gamma}(a_z)}{m_\mu(a_z)} \le m_\mu(z)m_{\mu+\gamma}(z^{-1}) = m_{-\mu}(z)^2.$$

For arbitrary g, the ordinary polar decomposition is $g = uz$ with $u \in K$ and z positive Hermitian in G_J; indeed $J(g^*g)J = (g^*g)^{-1}$ implies the same identity for its positive square root. Lemma 22.1 gives the splitting (u, z). Apply the preceding bound to every conjugate

v^*zv with $v \in K$ (the block determinant is unchanged). Cauchy–Schwarz and Haar invariance show

$$|\psi_\lambda(g)| \le (\det a_z)^{-\gamma/2} Z_\Lambda^{-1} \int_K |m_\Lambda(uv)|\,|m_\Lambda(v)|\,dv \le |\det a_g|^{-\gamma/2}.$$

For the other two bounds write $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ by signs. The two unitary identities yield

$$A^*A = 1 + C^*C, \quad AA^* = 1 + BB^*, \quad D^*D = 1 + B^*B, \quad DD^* = 1 + CC^*.$$

In particular the nonzero singular values of B, C agree. If one sign block is empty, the asserted inequalities follow immediately from $|\psi_\lambda| \le 1$ and $\|g\|_{\mathrm{op}} = 1$. Otherwise if $t = \|A\|_{\mathrm{op}} = \|D\|_{\mathrm{op}}$, then $\|B\|_{\mathrm{op}}, \|C\|_{\mathrm{op}} \le t$. All singular values of A, D are at least one, hence $|\det a_g| \ge t^2 \ge \|g\|_{\mathrm{op}}^2/4$. Furthermore

$$|\det a_g| = \det(1 + BB^*) \ge 1 + \|B\|_2^2, \quad |\det a_g|^2 \ge 1 + 2\|B\|_2^2 = 1 + \|g - a_g\|_2^2.$$

Substitution proves the last two bounds. □

This derives the bounds `norm_dsCoeff_sq_mul_det_pow_le_one`, `norm_dsCoeff_le_pow` of `Lib/DiscreteSeries/FlenstedJensen/PsiDecay.lean` and `norm_dsCoeff_pow_four_le` of `New/DiscreteSeries/TwoFactorSetup.lean`. Fischer's inequality follows, for example, by taking the Schur complement of one block of a positive definite matrix; no spectral asymptotic enters this step.

Lemma 22.6 (Very small balls)**.** *For $|\ell_i| \le L$ and $d + L + 1 \le \beta$, if $\|g - 1\|_{\mathrm{op}} \le (200\beta^5)^{-1}$, then $|\psi_\lambda(g)| \ge 1/2$.*

Proof. Set $\delta = (200\beta^5)^{-1}$. The matrix g is accretive, so use the splitting $(1, g)$ in Lemma 22.1. All principal compressed logarithms of v^*gv have absolute trace at most $2d\delta$, because the operator norm of their difference from the identity is at most δ. The number of nonconstant factors in the expression for $m_{-\mu}$ including the determinant power is at most $3\sum_i |\mu_i| \le 3d(L + d + 1)$. Thus $|\log m_{-\mu}(v^*gv)| \le 6\beta^3\delta \le 3/100$. Taking the reciprocal, the deviation from one is bounded by $e^{3/100} - 1 < 1/2$. Integrate with the positive normalized compact highest-type density to obtain $|\psi_\lambda(g) - 1| < 1/2$. □

The source proves this last assertion via the infinitesimal bound `norm_dsCoeff_sub_le`, which is also useful for the differentiated Schur inequalities. Only the small-ball assertion `half_le_norm_dsCoeff` of `Lib/DiscreteSeries/FlenstedJensen/TailBounds.lean` is needed here. The argument just given does not require an infinitesimal representation estimate.

22.5 Removal of the common compact subgroup

We retain the hypotheses of the two-factor setup of Section 21.4. Put $g_i = j_i(s)$, with ranks d_1, d_2, sign matrices J_i, and integral weight matrices Λ_1, Λ_2. Write $g_{i,\mathrm{diag}}$ for the same-sign block part of g_i. In the first group, $\Lambda_1 = 0$ on the common positive subspace X_0 of this setup (not the full definite subspace of the filling datum), and $-3B \le \Lambda_{1,j} \le -B$ elsewhere. In the second group Λ_2 is nonnegative, is bounded by $3B$ on all rows having any opposite-sign direction, and vanishes on the corresponding common positive subspace. The pair has equal traces and imaginary additive phase

$$\varphi(s) = \mathrm{tr}(\Lambda_1(g_1 - 1)) - \mathrm{tr}(\Lambda_2(g_2 - 1)).$$

It contains the simultaneous copy of the unitary group on X_0. Recall

$$\theta(s) = \psi_{\lambda_1}(g_1)\overline{\psi_{\lambda_2}(g_2)}e^{-\varphi(s)}.$$

Lemma 22.7 (Simultaneous compact correction)**.** *Suppose $B \geq 1$ and $0 < \rho \leq 1/100$. If*

$$\|g_1 - g_{1,\mathrm{diag}}\|_2^2 + \|g_2 - g_{2,\mathrm{diag}}\|_2^2 \leq \rho^2$$

then there is t obtained from s by left multiplication by an element of the common compact group, with $\theta(t) = \theta(s)$ and

$$\|j_1(t) - 1\|_2,\ \|j_2(t) - 1\|_2 \leq 10\rho.$$

Proof. Put $H_i = g_i - 1$. The signed row identity from Section 21.4 gives, for a diagonal nonnegative weight w,

$$\sum_{a,b} w_a|(H_i)_{ab}|^2 + 2\Re\,\mathrm{tr}(wH_i) = 2\sum_{\epsilon_a \neq \epsilon_b} w_a|(g_i)_{ab}|^2.$$

Use $w = -\Lambda_1$ and $w = \Lambda_2$ in the two groups and add. The trace contributions cancel because $\Re\varphi = 0$. If P is projection onto the common positive subspace in the first group, $Q = 1 - P$, it follows that $\|QH_1\|_2^2 \leq 6\rho^2$.

In blocks P, Q, write $g_1 = \begin{pmatrix} a & b_1 \\ c & d \end{pmatrix}$. The first block lies in a positive part of the sign form. Unitarity gives

$$a^*a + c^*J_Qc = 1, \qquad a^*b_1 + c^*J_Qd = 0.$$

Thus $\|a^*a - 1\|_2 \leq \|c\|_2^2 \leq 6\rho^2 < 1$, so a is invertible. Write $a = u|a|$ with u unitary on the common compact subspace. Since $|\sqrt{x} - 1| \leq |x - 1|$ for $x \geq 0$, $\||a| - 1\|_2 \leq 6\rho^2$. The second equation gives

$$\|b_1\|_2 \leq (1 - 6\rho^2)^{-1/2}(1 + \sqrt{6}\,\rho)\|c\|_2 \leq \tfrac{11}{10}\|c\|_2.$$

Choose the simultaneous compact element from the hypothesis which acts by u^{-1} on both copies of the common compact subspace and fixes their complements. For the corrected first matrix $\widetilde{g}_1$, the block a becomes $|a|$, the row Q is unchanged, and b_1 is multiplied by a unitary on the left. Consequently

$$\|\widetilde{g}_1 - 1\|_2^2 \leq 36\rho^4 + (1 + (11/10)^2)6\rho^2 < 16\rho^2.$$

This compact multiplication leaves the opposite-sign norms of each factor unchanged. In the two signatures apply the signed row identity with all weights equal to one. Equality of the traces of the two corrected factors implies

$$\|\widetilde{g}_2 - 1\|_2^2 \leq \|\widetilde{g}_1 - 1\|_2^2 + 2\|\widetilde{g}_2 - \widetilde{g}_{2,\mathrm{diag}}\|_2^2 \leq 18\rho^2.$$

Finally the lowest compact type has weight zero on the whole common compact block. Its highest vector is fixed by the unitary group of that block: the restriction has highest weight all zero, so its orbit there is the trivial irreducible. Equivalently the product of weight minors, restricted to the compact block in question, is unchanged by this compact rotation (the weight is constant zero throughout it). The normalized coefficient of this vector is unchanged by left multiplication by the corresponding element in each group. Also the linear phase is zero on the common compact subgroup, and is additive on S. Thus θ is unchanged. This proves the bounds and invariance. □

This is `IsTwoFactor.exists_near_one` in `New/DiscreteSeries/TwoFactorSetup.lean`. The polar correction here replaces a matrix inverse construction in the formal proof but gives the same error constant. In particular it uses the equal trace condition, not a separate estimate of the second group's compact block.

Proposition 22.8 (Cone estimate for the oscillatory integral)**.** *Let $D = (d_1 + d_2)^2$, $\beta \geq 10^{12}$, $d_1 + d_2 + L \leq \beta$, $T = D \log \beta$, and suppose the weight separation in the two groups is at least $10^{30}T^3$ in the notation of the large-gap hypothesis. Write $G(s) = 2/\|j_1(s)\|_{\mathrm{op}}$. Then*

$$|\theta(s)| \leq G(s)^a \quad (0 \leq a \leq 18D), \qquad 4|\theta(s)| \leq 8(\Re\theta(s) + 16\beta^{-50D}G(s)^{8D}).$$

In the first inequality it suffices to take integer a; real exponents in this interval follow by interpolation with $a = 0$.

Proof. The first bound uses Lemma 22.5 for the first factor and the unit-vector coefficient bound for the second. For the second put $\rho = 10^{-13}/T$. If the sum of the squared opposite-sign norms is at most ρ^2, use Lemma 22.7 to make both corrected factors $10^{-12}/T$-close to the identity in Frobenius norm. Their exponential phases cancel exactly by the logarithmic phase identity (21.11); its proof used only additivity of φ. Theorem 22.4 with $T = D \log \beta$ gives θ as a product of two complex numbers whose differences from $a_1, a_2 \in [0, 1]$ are at most $a_i/10 + \zeta$, where $\zeta = e^{-100T} = \beta^{-100D}$. (Conjugating the second factor leaves this estimate unchanged.) In particular

$$|\theta - a_1 a_2| \leq 0.21 a_1 a_2 + 3\zeta.$$

Since $\Re\theta \geq a_1 a_2 - |\theta - a_1 a_2|$, it follows that

$$|\theta| - 2\Re\theta \leq 3|\theta - a_1 a_2| - a_1 a_2 \leq 9\zeta.$$

The operator norm of j_1 is unchanged by the compact correction. It is now at most $1 + 10^{-12}/T < 2$, so $G(s) > 1$, and the required inequality follows.

Otherwise the sum of the squared opposite-sign norms exceeds ρ^2. Multiplication of the fourth-power decay bounds gives

$$|\theta|^4 \leq (1 + \rho^2)^{-\gamma} \qquad \text{with } \gamma \geq 10^{30}T^3.$$

Indeed the product $(1 + \|g_1 - a_{g_1}\|_2^2)(1 + \|g_2 - a_{g_2}\|_2^2)$ is at least one plus the sum. Since $\log(1 + \rho^2) \geq \rho^2/2$ for $\rho^2 \leq 1$, $\frac{1}{4}\gamma \log(1 + \rho^2) \geq 1250T$. Hence $|\theta| \leq \zeta$. Use also the decay bound $|\theta| \leq G^{16D}$ and take the geometric mean to get $|\theta| \leq \beta^{-50D}G^{8D}$. Then $|\theta| - 2\Re\theta \leq 3|\theta|$ proves the second inequality in this case. □

If the gap input is given with an additional integer threshold $500d_i^3$, as in the source, the loss from taking an integer exponent in Lemma 22.5 is harmless (the argument can take the specified large integer gap). The constants here prove the pointwise inequalities (21.13)–(21.14) needed by the integration argument of Proposition 21.6. The small-ball assertion in that proof is supplied by Lemma 22.6. Thus the specialized oscillatory part of `IsTwoFactor.laplace_method` of `New/DiscreteSeries/RestrictionLemma.lean` has now been reconstructed. It is conditional on the representation-theoretic coefficient identification, rather than on an additional asymptotic estimate for discrete series.

22.6 A phase-cancelling realization of the Bessel group

We now give the geometric realization required for the Bessel peak in Section 21.5. The standard complex Hermitian form is

$$J_1 = \begin{pmatrix} 0 & 0 & 1_h \\ 0 & T & 0 \\ 1_h & 0 & 0 \end{pmatrix}, \qquad T = \operatorname{diag}(1_k, -1),$$

with standard frame $F = (e_0, \dots, e_{h-1}, z)$, isotropic duals $e_0^*, \dots, e_{h-1}^*$, and definite part with basis $x_1, \dots, x_k$. The Bessel group $S_B = UH$ consists of transformations acting by an upper unitriangular matrix on F, where $H = U(k)$ fixes the hyperbolic planes and z, and U acts trivially on the middle quotient. On S_B put

$$\sigma(s) = \sum_{i=0}^{h-1} s_{f_i, f_{i+1}}, \qquad f_i = e_i \ (i < h), \quad f_h = z.$$

Since the frame matrix is unitriangular, σ is additive on products. The real character in the source is $\mathrm{wave}(\sigma) = \exp(\sigma - \overline{\sigma})$. We first prove a useful norm estimate for its realization.

Lemma 22.9 (Controlling a representative of a Bessel frame)**.** *Suppose vectors V_j for $0 \le j \le 2h$ satisfy $\|V_j\| \le M$, $M \ge 1$, and*

$$V_a = x^{-1} f_a \ (a \le h), \qquad \langle V_i, J_1 V_j \rangle = \begin{cases} 0 & i + j < 2h, \\ -1 & i + j = 2h. \end{cases}$$

Then $\mathrm{bs}(x) = \inf_{s \in S_B} (\|sx\|_{\mathrm{op}} + 1) \le 8(h+1)M$.

Proof. Apply unnormalized ordinary Euclidean Gram–Schmidt to the ordered frame $(V_0, \dots, V_h)$, writing F_a for its orthogonalized columns. They are obtained by an upper unitriangular change of columns. Such a change lifts to an element of U: if the change on the isotropic block is A and the extra column of the last frame vector is v, the lift on (e, z, e^*) is

$$\begin{pmatrix} A & v & \frac{1}{2} v v^* (A^*)^{-1} \\ 0 & 1 & v^* (A^*)^{-1} \\ 0 & 0 & (A^*)^{-1} \end{pmatrix},$$

and it fixes the definite part. Direct multiplication against the signed form verifies this formula. Hence some Bessel translate has inverse frame precisely (F_a). Each $\|F_a\| \le M$. The anti-diagonal hypothesis gives $\langle F_a, J_1 V_{2h-a} \rangle = -1$, because the earlier columns have zero pairings with V_{2h-a}. Thus $\|F_a\| \ge M^{-1}$.

For $a < h$ put $D_a = J_1 F_a / \|F_a\|^2$. Orthogonality of the Euclidean Gram–Schmidt frame gives $\langle F_b, J_1 D_a \rangle = \delta_{ab}$ (including $b = h$), and $\langle D_a, J_1 D_b \rangle = \langle F_a, J_1 F_b \rangle / (\|F_a\|^2 \|F_b\|^2) = 0$. The vectors D_a are bounded by M and form an isotropic frame dual to $(F_0, \dots, F_{h-1})$. They can be made the inverse images of the standard dual frame by an additional translation in U^0, the subgroup fixing the whole original frame. To see this, in coordinates of the already adjusted element any such dual array has columns $(C; A; 1_h)$ in the $(e, X_{\mathrm{definite}}, e^*)$ decomposition, with zero z-part. Its isotropy is $C + C^* + A^* A = 0$. Writing $C = B - \frac{1}{2} A^* A$ with B skew-Hermitian gives the usual unipotent element fixing the frame,

$$\begin{pmatrix} 1 & -A^* & B - \frac{1}{2} A^* A \\ 0 & 1 & A \\ 0 & 0 & 1 \end{pmatrix}.$$

It has exactly these dual columns. It fixes z and is unitary for the hyperbolic and definite form by the displayed identity.

Let M_0 be the inverse of the resulting Bessel translate, with frame and dual columns denoted by F_a, D_a, and negative column F_h. Using $M_0 J_1 M_0^* = J_1$ and subtracting it from $M_0 M_0^*$ gives the exact identity

$$M_0 M_0^* = J_1 + \sum_{a<h} (F_a - D_a)(F_a - D_a)^* + 2F_h F_h^*.$$

Therefore $\|M_0\|_{\mathrm{op}}^2 \le 1 + (4h+2)M^2$. Also the inverse has the same operator norm since J_1 is Hermitian unitary. In particular the norm plus one is bounded by $8(h+1)M$, as required. □

This proves `besselSize_le_of_dual` in `New/RealPlace/FrameNorm.lean` using the triangular-coordinate description of U^0.

For the next construction the ambient weight on the r-dimensional sign space is zero on a common positive subspace X of dimension $k-1$ and has strictly decreasing exceptional entries $\Lambda_i \in [-3B, -B]$, $0 \le i \le 2h+1$, with signs $(-1)^i$. The compact weight on the k-dimensional definite group is zero on X and equals $a \ge 0$ on its remaining distinguished coordinate. The source imposes $B \ge 1$, $3B + a \le 2b^2$, and $b \ge 8(h+1)$.

Proposition 22.10 (The interpolation realization)**.** *There is $x \in G_{J_1}$ with $\mathrm{bs}(x) \le b^{5(h+1)}$ such that the Bessel group graph $s = (g_s, h_s) \in S_B \times H$, with $g_s \in UH$ and compact factor h_s, is a two-factor subgroup on conjugating the first factor by x and an orthonormal sign frame. The phase is exactly*

$$\varphi(s) = \sigma(g_s) - \overline{\sigma(g_s)}.$$

Both factors contain the indicated common unitary subgroup diagonally, and their traces agree after subtracting their respective identities.

Proof. Put $t_i = \Lambda_i - a$; these are strictly decreasing negative integers. Apply the Lagrange interpolation Lemma 21.7 to the descending sequence $(0, t_0, \dots, t_{2h+1})$ and write its positive weights as $c_0, c_1, \dots, c_{2h+2}$. In the exceptional coordinates form

$$(C_0)_i = c_0/c_{i+1}, \qquad (C_j)_i = t_i^j/c_{i+1} \quad (j \ge 1).$$

The signed interpolation identity gives $H(C_0, C_0) = 1$, $H(C_0, C_j) = 0$ for $1 \le j \le 2h+1$, and $H(C_i, C_j) = 0$ for $i, j > 0$, $i + j < 2h+2$, equal to -1 when $i + j = 2h+2$. Here H denotes the signed form on the exceptional coordinates. Consequently the common positive X, followed by $C_0, C_1, \dots, C_{h+1}$, forms the same partially degenerate Hermitian frame as the standard X, its distinguished last definite vector x_0, and $(e_0, \dots, e_{h-1}, z)$. Witt extension for nonsingular Hermitian spaces over $\mathbb{C}/\mathbb{R}$ extends this isometry of subspaces to an isometry of the whole space. Equivalently choose duals for the isotropic part and subtract half their Gram matrix to make them isotropic, then complete the two orthogonal complements by orthonormal bases. Let p be an orthonormal sign frame for J_1. Choose x so that xp maps the above columns to the standard ones.

The columns pC_j, $1 \le j \le 2h+1$, are the dual array in Lemma 22.9, with $V_i = pC_{i+1}$. Since each $c_{i+1} \ge 1$ and $|t_i| \le 2b^2 \le b^{5/2}$,

$$\max_{1 \le j \le 2h+1} \|pC_j\| \le \sqrt{2h+2}\, b^{(5/2)(2h+1)}.$$

Lemma 22.9 gives $\mathrm{bs}(x) \le b^{5(h+1)}$, since $8(h+1)\sqrt{2h+2} \le b^{3/2}$.

We verify the phase rather than infer it from Witt extension. Let

$$D = xp\,\mathrm{diag}(\Lambda_i)(xp)^{-1}.$$

It is self-adjoint for J_1. In the above columns, multiplication by a weight is multiplication by $t_i + a$, hence

$$Df_j = af_j + f_{j+1} \quad (0 \le j < h), \qquad Dx_0 = ax_0 + c_0 f_0, \qquad D|_X = 0.$$

Order the blocks of G_{J_1} as the isotropic lines $e_0, \dots, e_{h-1}$, the middle definite block together with z, and the dual lines in opposite order. Matrices of U are strictly block upper triangular off the identity. To contribute to $\mathrm{tr}\, D(g_s - h_s)$ an entry of D must occur strictly below these blocks. For a column e_i the only such entry is the coefficient 1 of f_{i+1}. For columns in the middle block, self-adjointness gives a nonzero dual coefficient only for the column z, with coefficient -1 at e^*_{h-1}, since $H(De_{h-1}, z) = H(z, z) = -1$. (For a definite column its pairing with every De_i is zero.) For the dual columns, the remaining lower entries are prescribed by self-adjointness of the isotropic block: they are the coefficients 1 at (e^*_i, e^*_{i+1}). The dual blocks of a unitriangular unitary matrix are inverse adjoints. Thus the superdiagonal dual entries of g_s are minus the conjugates of the superdiagonal isotropic entries. For the last root the sign of the negative line makes the z, e^*_{h-1} entry the conjugate of e_{h-1}, z, as is seen by imposing $H(g_s z, g_s e^*_{h-1}) = 0$. The middle rotation h_s acts trivially on the indicated frame and dual entries. Summing all contributions proves

$$\mathrm{tr}\, D(g_s - h_s) = \sigma(g_s) - \overline{\sigma(g_s)}.$$

On the middle definite block D is exactly the compact weight $\mathrm{diag}(0_X, a)$; the other diagonal entries of $h_s - 1$ vanish. Therefore $\mathrm{tr}\, D(h_s - 1)$ is the second-factor weight trace. This gives the required phase. Its real part is zero, its additivity follows from upper unitriangularity of the frame matrices, and the trace identity follows from the diagonal blocks of g_s. Finally a compact transformation of the common X is identified by xp with the standard unitary subgroup acting on those definite coordinates. Thus it is present in the two factors simultaneously. □

The source statement is `IsSteep.exists_isTwoFactor_besselPairHom` in `New/DiscreteSeries/BesselExistence.lean`. Its hypotheses on the actual parameter pattern are supplied by `Data.exists_isSteep` in `New/Exchange/SteepParameters.lean`; establishing the inequalities between the actual weight parameters is separate from this interpolation calculation. The first projection of the Bessel graph after conjugation is proper because the subgroup UH is closed and the compact factor of the graph is a continuous function of the middle block. The second component has operator norm one. Accordingly the quantitative Laplace argument above applies once its gap and size hypotheses are verified.

Combined with the quotient integral construction (21.15)–(21.16) and the peak-to-cut argument of Section 21.6, this supplies the principal analytic mechanism of `Data.exists_besselSystem` (`New/Filling/FillingData.lean`). The representation-theoretic input of a unit highest vector of the compact Schur type, identification of the lowest-type coefficient and its formal degree, and the differentiated convolution bound are still used as explicitly indicated. For the remaining Fourier-floor portion of this declaration, the arguments just proved do not replace the local gauge, bandlimit and open-cell estimates in `New/RealPlace/`. We return to that distinction at the end of the chapter.

22.7 Marked integral lattice and the exact finite level indices

Let E be the CM field, E^+ its totally real subfield, $\mathcal{O} = \mathcal{O}_E$, $[E : \mathbb{Q}] = 2f$. The Hermitian space is

$$V = L \oplus M \oplus L^\vee, \qquad L = \langle e_0, \dots, e_{h-1} \rangle_E, \qquad M = X \oplus Ez, \quad h|_X = 1_k, \quad h(z,z) = -N,$$

where $L^\vee$ is isotropic dual to L and X is positive definite at every infinite place. Write $\mathcal{O}_f = \widehat{\mathcal{O}}$, and normalize the global additive measure using ordinary Lebesgue measure on $\mathbb{C}^f$; by the covolume-one convention on the adeles, $\mathrm{vol}(\mathcal{O}_f) = 2^f |D_E|^{-1/2}$. The constructions use the generator ϖ of the different, with $\bar{\varpi} = -\varpi$, and the primitive trace-one element of $\mathcal{O}$ available because E/E^+ is everywhere unramified at its finite places in the stipulated `StablyUnram` setting. The same setting supplies a unit $i_0 \in \mathcal{O}$ with $i_0^2 = -1$.

We distinguish the two halves of the split level prime N by writing $\eta\bar{\eta} = N$ for finite ideles, $\eta, \bar{\eta} \in \mathcal{O}_f$; at each of the pairs over N precisely one of them is a uniformizer and the other a unit. At primes away from N both are units. The marked lattice $\mathcal{L}$ in $V(\mathbb{A}_f)$ has the standard hyperbolic basis and, in the middle block, the basis

$$x_i \ (i \neq i_0'), \qquad v = x_{i_0'}/\eta + z/N, \qquad z.$$

Here i_0' is merely the chosen index in the positive definite basis, unrelated to the imaginary unit i_0. The Gram matrix of (v, z) is

$$\begin{pmatrix} 0 & -1 \\ -1 & -N \end{pmatrix};$$

indeed $h(v,v) = 1/N - N/N^2 = 0$ and $h(v,z) = -1$. Hence $\mathcal{L}$ is unimodular. Let K be the integral stabilizer of this lattice and $H = U(X)$ the subgroup of $U(V)$ fixing the hyperbolic pairs and z. Write K_X for the stabilizer of the standard finite-adelic lattice in X.

Lemma 22.11 (The marked-lattice index)**.** *The subgroup $K \cap H(\mathbb{A}_f)$ is contained in K_X, and its index there is $(N^k - 1)^f$.*

Proof. At a finite place write $\mathcal{L}_X = \mathcal{L} \cap X$ and $\mathcal{L}^X$ for the projection of $\mathcal{L}$ onto X. At the marked split places

$$\mathcal{L}_X = \sum_{i \neq i_0'} \mathcal{O}_w x_i + \bar{\eta}\mathcal{O}_w x_{i_0'}, \qquad \mathcal{L}^X = \sum_{i \neq i_0'} \mathcal{O}_w x_i + \eta^{-1}\mathcal{O}_w x_{i_0'}.$$

Indeed a combination $tv + sz$ lies in X if and only if $s = -t/N$, i.e. $t \in N\mathcal{O}_w$. Away from N both lattices are standard, while at each split place over N one or the other is standard. Any transformation of X preserving $\mathcal{L}$ preserves both lattices; it therefore lies in K_X.

For $x \in K_X$ the condition that its action preserve $\mathcal{L}$ is exactly the finite marked-vector condition. To check it apply the action to the unmarked vectors, v and z. At a place where η is the uniformizer, $\mathcal{L}_X$ is standard and the only new condition is

$$x(v) - v = (x x_{i_0'} - x_{i_0'})/\eta \in \mathcal{L}_X,$$

so the distinguished vector is fixed modulo η. At its conjugate, η is a unit and the other factor $\bar{\eta}$ is a uniformizer; preservation means that the distinguished coordinate row of x agrees with the identity row modulo $\bar{\eta}$. These are equivalent under the unitary inverse-adjoint relation between the two split local factors. The same test on the inverse gives equality, not just containment; the vector condition is preserved by inverse.

The split local unitary stabilizer is identified with $\mathrm{GL}_k(\mathcal{O}_w)$ at the selected place of each pair, and reduction onto $\mathrm{GL}_k(\mathbb{F}_N)$ is surjective: lift the entries of an invertible matrix, whose determinant remains a unit, and fill its conjugate local factor by inverse adjoint. The index of the stabilizer of a nonzero vector in $\mathrm{GL}_k(\mathbb{F}_N)$ is the cardinality $N^k - 1$ of the orbit of that vector. Multiply over the f pairs of split places over N. □

This establishes `Data.relIndex_latStabilizer` of `New/Filling/FillingAdmissible.lean`. It is important that the condition fixes a *vector* rather than merely a line; the latter gives a different index by the factor $N - 1$.

The two auxiliary rational primes q_0, q_1 split in E and are different from N; each comes with a choice of f places, one from every conjugate pair. Reduction of K at these places gives

$$\mathfrak{G} = \prod_{j=0}^{1} \prod_{w \mid q_j \text{ chosen}} \mathrm{GL}_r(\mathbb{F}_{q_j}).$$

This map is surjective by the same split lifting as in Lemma 22.11; its kernel is the level $K(g)$, where $g = q_0 q_1$. The marking at N does not interact with this reduction, since local factors can be set to the identity at the other finite places. The column level is $K_c = K(g) \cap H(\mathbb{A}_f)$, the preimage of the congruence level under the inclusion of H in the marked lattice basis.

Let $\bar{U}, \bar{H}, \bar{S}_B$ denote the images under reduction of U, H, and $S_B = UH$, intersected with K. At the split level primes, the image of U is the block unitriangular group of block sizes $1, \dots, 1, k+1, 1, \dots, 1$ (h blocks of size one at each end). The conjugate factor is filled by inverse adjoint, so there are no extra equations on the selected factor. Its number of free residue-field entries is

$$\frac{(k+2h+1)^2 - (k+1)^2 - 2h}{2} = h(2k+2h+1).$$

The image of H is the ordinary k-dimensional linear group inside the middle block. Thus $\bar{S}_B = \bar{U} \rtimes \bar{H}$, and

$$|\bar{U}| = g^{fh(2k+2h+1)}, \qquad |\bar{S}_B| = |\bar{H}|\, g^{fh(2k+2h+1)}. \tag{22.6}$$

For the count, lifts of unitriangular elements are obtained by lifting their free entries and the inverse-adjoint factor; outside the marked split places take the identity. Conversely a product uh is integral only if both factors are integral: its middle diagonal block determines h, and multiplying by h^{-1} recovers u. The same argument applies to congruence levels. This is the content of `card_reducedImage_besselS` in `New/Filling/GlobalToLocal.lean`; it also shows directly why the exponent of g depends on the full hyperbolic structure and not just the dimension of the upper triangular frame group.

In the positive definite group write

$$\mathcal{C} = H(E^+) \backslash H(\mathbb{A}_f) / K_c, \qquad \mathfrak{m}_k = \sum_{d \in H(E^+) \backslash H(\mathbb{A}_f)/K_X} |\operatorname{Aut}(d)|^{-1}.$$

The usual class mass calculation is recalled in Chapter 5. Since the deeper level K_c has trivial rational stabilizers, the number of classes over d is $[K_X : K_c]/|\operatorname{Aut}(d)|$. The stabilizer acts freely on the finite quotient: a fixed coset would give a conjugate stabilizer at level K_c. By Lemma 22.11 and surjectivity of reduction on the middle group,

$$|\mathcal{C}| = |\bar{H}|(N^k - 1)^f \mathfrak{m}_k. \tag{22.7}$$

This gives `Data.card_classes_filling`. Together with the cardinality $D_I = \prod_v |I_v|$ of the local Bessel index tuple, the column dimension is $D_c = |\mathcal{C}| D_I$. Notice the cancellation between the $|\bar{H}|$ factors in the class count and the order of $\bar{S}_B$ in the energy bounds below.

22.8 Haar coordinates and the arithmetic unipotent covolume

We give the exact unipotent covolume calculation, including its dependence on the marking. Write $T = \mathrm{diag}(1_k, -N)$ for the middle form before archimedean rescaling. Relative to the ordered blocks $(L, M, L^\vee)$, an element of U is determined by three upper blocks as follows:

$$u = \begin{pmatrix} A & B & C \\ 0 & 1 & D \\ 0 & 0 & (A^*)^{-1} \end{pmatrix}, \qquad D = -T^{-1}B^*(A^*)^{-1},$$

$$M_0 + M_0^* = -A^{-1}BT^{-1}B^*(A^*)^{-1}, \qquad M_0 = A^{-1}C,$$

with A upper unitriangular. These formulas follow by multiplying the Gram matrix against the three block columns. At an infinite place we will use the normalized negative entry -1. The free complex entries may be taken to be, in row e_a:

$$e_b,\ e_b^*\ (b > a), \quad \text{all } k+1 \text{ middle columns}, \qquad 0 \le a < h.$$

In addition take the real number $\Im C_{aa}$ for every a. To see that these are actual coordinates, fix A, B and the upper triangular entries C_{ab} for $a < b$. Solve $M_0 = A^{-1}C$ column by column, starting with $b = h-1$ and descending: the lower entries $M_{0,jb}$, $j > b$, are determined by the already computed columns and the displayed Hermitian relation. The real part of the diagonal of M_0 is fixed by this relation, its imaginary part is determined by $\Im C_{bb}$ after subtracting known later entries, and then the entries $M_{0,ab}$ for $a < b$ are solved in descending order a. All these operations are affine triangular with diagonal coefficient one in the free coordinates.

We normalize Haar measure on U_∞ to be Lebesgue in these free entries. This agrees with the source's `Shape.besselHaar`, which uses Euclidean measure on the unitriangular frame and on the skew-matrix and definite variables (`Lib/DiscreteSeries/Bessel/UnipotentCoords.lean`, `RealMeasures.lean`). To verify the agreement, parametrize M_0 by its skew-Hermitian part plus the known quadratic term in A, B. The change of its skew coordinates into the free entries is triangular, of determinant of modulus one, by the preceding calculation. Left multiplication by a fixed group element has triangular coordinate Jacobian one: it multiplies A by an upper unitriangular matrix, replaces B by that matrix times B plus a fixed vector, and replaces M_0 by itself plus a function of the other coordinates, as follows from the block multiplication law. Hence the measure is left Haar and also right Haar, since this group is unipotent. Using the compact subgroup H's Haar probability gives the Bessel graph measure, as in Section 22.6.

For a class c choose a representative $h_c \in H(\mathbb{A}_f)$ and use the lattice $\mathcal{L}_c = h_c\mathcal{L}$, with the basis matrix $B_c = h_c B_{\mathcal{L}}$ in rational coordinates. The matrix B_c is the identity on the two isotropic blocks and invertible on the middle block. Its generalized index is

$$[\det B_c] = N^{-f}.$$

Here $[a] = \prod_{w \text{ finite}} |a_w|_w^{-1}$, and $|\cdot|_w$ is the normalized local module. Indeed $[\eta] = N^f$, the determinant of the middle basis change is η^{-1}, and the determinant of h_c has norm one. The Gram matrix of $\mathcal{L}_c$ is the same unimodular Gram matrix as that of $\mathcal{L}$. Let

$$\Gamma_c = U(E^+) \cap h_c K(g) h_c^{-1}.$$

Lemma 22.12 (Integral row translations)**.** *Let l be a rational linear row form supported on the free complex entries of row a. Suppose lB_c is integral at all finite places. Given $s_0 \in \mathcal{O}$ with $\bar{s}_0 = -s_0$, there is $\gamma \in \Gamma_c$ whose free off-diagonal coordinates are zero except in row a, where they equal gl, and when $l = 0$ its only free coordinate is the skew diagonal entry gs_0 in that row.*

Proof. The Hermitian form identifies a row form with a vector: choose v with $h(v,\cdot) = l$. The condition on l says exactly that the finite image of v lies in the integral dual of $\mathcal{L}_c$; since $\mathcal{L}_c$ is unimodular, that dual is $\mathcal{L}_c$. Put $x = e_a$; it also lies in $\mathcal{L}_c$, is isotropic, and $h(v,x) = 0$. For an integral trace-one element τ ($\tau + \bar{\tau} = 1$) use the Eichler transformation

$$u = 1 + g(xh(v,\cdot) - vh(x,\cdot)) + (gs_0 - g^2\tau h(v,v))xh(x,\cdot).$$

It is unitary: the square of the first-order skew part is a multiple of $xh(x,\cdot)$; expanding u^*Ju reduces the quadratic identity to $\tau + \bar{\tau} = 1$ and $s_0 + \bar{s}_0 = 0$. It is upper unipotent relative to the flag because l vanishes on each e_b, $b \le a$, and the first-order non-row term has only the column e_a^*. Equivalently the matrix and its inverse are block upper triangular with diagonal blocks the identity; the possible nonfree entries in row $b > a$ are confined to the column e_a^*. Its difference from identity sends the class lattice into $g\mathcal{L}_c$, by integrality of x, v, their pairings, and τ, s_0. Thus it lies in Γ_c. Inspecting the entries of the displayed formula gives the assertion. □

Here trace one cannot always be replaced by $\tau = 1/2$: the latter need not preserve an integral lattice at residue characteristic two. Lemma 22.12 proves the specialized lifting statement `exists_eichler_classUnip` in `New/Filling/BoxPartition.lean`.

For each row set $d_a = k + 1 + 2(h - a - 1)$, the number of its free complex entries. The rational row lattice of those entries is

$$\mathcal{R}_a = \{l : \operatorname{supp} l \subseteq \text{free positions of row } a, lB_c \in \mathcal{O}_f^r\}.$$

The matrix on these positions is the restriction of B_c to them. It contains the entire middle block and is otherwise identity. Therefore its determinant has generalized index N^{-f}. In the ordinary mixed embedding $(\Phi_v)_{v|\infty}$ its covolume is

$$\frac{N^f}{\operatorname{vol}(\mathcal{O}_f)^{d_a}}.$$

For this formula, if D is any finite adelic invertible matrix, the lattice of rational row vectors x for which xD is integral is commensurable with $\mathcal{O}^d$. The ratio of covolumes is the inverse of the generalized index of $\det D$, by taking Smith forms locally (or multiplying by an integer to compare finite-index sublattices). This argument does not choose a principal generator for the resulting global lattice; it also applies when h_c has nontrivial class.

The rational skew lattice of a diagonal entry is $\{s : \bar{s} = -s,\ s \in \mathcal{O}\} = i_0\mathcal{O}_{E^+}$. The equality follows by dividing by the imaginary unit i_0, which is integral and a unit. The covolume of its imaginary parts at the f complex places is the covolume of the totally real integer lattice, namely $|D_{E^+}|^{1/2} = |D_E|^{1/4}$, since the stipulated CM field is unramified over E^+ at its finite places. This verifies the skew-lattice formula of `Filling.exists_fundamental_domain_im_skew`, `Lib/General/NumberTheory/FundamentalDomains.lean`.

We next justify multiplying these covolumes. Grade the free-coordinate groups in descending row order, using first the complex coordinates in a row, then its skew diagonal coordinate. If all earlier coordinates of $\gamma \in \Gamma_c$ vanish, right multiplication $u \mapsto u\gamma$ adds the free coordinates of γ at the current grade, without changing earlier ones. Here is the block verification: for rows $b > a$ already zero in their free coordinates, the corresponding rows of A and B in γ are identity and zero, respectively. The submatrix of C on two indices $b, a' > a$ is zero by the relation on $M_0 + M_0^*$, since the upper triangular and imaginary diagonal entries have already been set to zero. The D entries in the corresponding dual columns also vanish, and the corresponding columns of $(A^*)^{-1}$ are identity columns. Thus the matrix product cross term $(u-1)(\gamma-1)$ vanishes at all the free entries of row a. At the next grade, when row

a's complex coordinates also vanish in γ, the same argument makes the cross term vanish at its diagonal skew coordinate. The arithmetic integrality condition for that row is precisely $lB_c \in \mathcal{O}_f^r$ after division by g; restriction to the free coordinates preserves integrality because the full middle block is included and the other blocks are identity. For the skew grade, when the complex entries are zero the diagonal entry is actually skew, by the same Hermitian relation. Lemma 22.12 realizes every permitted shift with all earlier coordinates zero.

Choose a half-open fundamental parallelepiped for each row lattice and for each real skew lattice. Successively in the above order translate to those parallelepipeds by an element of Γ_c. Each step exists by Lemma 22.12, and does not alter previous steps. For uniqueness, if two resulting elements are related by a group element of Γ_c, consider its earliest nonzero coordinate: the two coordinates in that parallelepiped would differ by a nonzero lattice shift, impossible. Injectivity of the free entries gives uniqueness. Thus the product of the parallelepipeds, in these graded coordinates, is a measurable strict fundamental domain. Inversion preserves Haar, so its volume is also the left covolume. This is the graded-box argument of `Filling.existsUnique_mul_mem_box` in `Lib/General/NumberTheory/FundamentalDomains.lean`.

Finally set

$$\kappa_v = \frac{g\,\Phi_v(\varpi)}{2\pi i} \in \mathbb{R}^\times, \qquad l_{va} = \frac{\kappa_v^{h-a}}{\sqrt{N}}, \quad s_{va} = l_{va}^2 \quad (0 \le a \le h).$$

The infinite-place change of basis scales e_a by l_{va}, e_a^* by l_{va}^{-1}, definite coordinates by one, and z by $l_{vh} = 1/\sqrt{N}$. It carries the middle negative norm to -1. In a free complex entry at row a, the rescaling multiplies by the column scale divided by l_{va}. The pairs e_b, e_b^* for $b > a$ cancel in the product of all numerator scales; the only remaining numerator is l_{vh} in column z. A complex scaling changes Lebesgue measure by its squared absolute value. The skew diagonal changes Lebesgue measure by l_{va}^{-2}. Including the level factor g and the preceding lattice covolumes, the volume contribution of row a is

$$\Big(\prod_v l_{vh}^2\, l_{va}^{-2(d_a+1)}\Big)\frac{g^{2fd_a}N^f}{\mathrm{vol}(\mathcal{O}_f)^{d_a}}g^f|D_E|^{1/4}.$$

As $\prod_v l_{vh}^2 = N^{-f}$, the row factors N^f cancel. Put

$$d_0 = \sum_{a=0}^{h-1} d_a = h(k+h), \qquad \mathcal{S} = \prod_v \prod_{a=0}^{h} s_{va}^{k+2(h-a)}.$$

Since $d_a + 1 = k + 2(h-a)$, $\prod_{v,a<h} s_{va}^{-(d_a+1)} = N^{-kf}/\mathcal{S}$. The complete covolume is therefore

$$\boxed{\nu = \frac{(g^{\,2d_0+h}/N^k)^f\,|D_E|^{h/4}}{\mathrm{vol}(\mathcal{O}_f)^{d_0}\mathcal{S}}.} \tag{22.8}$$

This is exactly `Filling.unipCovolume` in `Setup.lean`. Its independence of c follows from the determinant index of B_c, not from assuming the class lattices are globally isometric.

Every measurable nonnegative cutoff β_c satisfying $\sum_{\gamma\in\Gamma_c}\beta_c(\gamma y) = 1$ then has integral ν over each left U_∞-orbit: integrate the identity over the fundamental domain just constructed and use left Haar invariance and Tonelli. That is the precise assertion `lintegral_classCutoff_eq_unipCovolume` in `New/Filling/UnipotentCovolume.lean`. It supplies the constant used in the unfolding and energy normalization of Chapter 20.

22.9 Cutoffs, approximation and the quotient character

The measure calculation did not need a particular cutoff. The counting argument does need one whose support is boxed in the frame variables when the Bessel-size cutoff is active. We explain the construction and distinguish its arithmetic size estimate.

First the rational image of every Γ_c under the archimedean embedding is discrete. For a bounded set of entries at all archimedean places, the condition of lying in $h_c K(g) h_c^{-1}$ confines the finite entries to a compact set. An entry belongs to a denominator lattice of E with bounded archimedean values; there are finitely many such entries by discreteness of this lattice. Thus the set of matrices under these bounds is finite. The same argument applies to any prescribed submatrix. In particular if Γ_c^0 is the kernel of the map Γ_c to the frame unitriangular group, it is a closed discrete subgroup of U_∞^0. It has a measurable nonnegative cutoff of sum one on its left orbits (use a countable local fundamental covering of the discrete proper action and a partition of unity, or a Borel fundamental domain). Write β^0 for such a cutoff.

There is a useful structural fact about bounded frame boxes. Fix $y \in G_\infty$. For a frame unitriangular matrix A let u be any lift in Γ_c, so the vectors $(u^{-1}y)^{-1} f_a = y^{-1} u f_a$ depend only on A. There are finitely many possible A for which all these vectors have bounded norms at all infinite places: multiplying by the fixed y_v, the entries of A are bounded and the finite positions are also bounded in the class lattice. Let $\mathcal{Q}_c = \Gamma_c/\Gamma_c^0$ and let $I_y(A)$ be the indicator that these frame vectors have norm at most

$$H(y, v) = b^{60h+92} \mathrm{bs}(y_v) \qquad \text{at each } v.$$

The source proves that at least one such A exists for every y. Assuming this numerical reduction assertion for a moment, a cutoff on the quotient is given by

$$\omega(y) = \frac{I_y(1)}{\sum_{A \in \mathcal{Q}_c} I_y(A)}. \tag{22.9}$$

Each summand is measurable, the denominator is finite and positive, and $\omega \le 1$. The identity $I_{uy}(A) = I_y(u^{-1}A)$ gives the required weighted partition by reindexing the quotient. Also I_y and hence ω are invariant by the left action of $U_\infty^0 H_\infty$, which fixes the frame and leaves Bessel size unchanged. If $\omega(y) \neq 0$ and $\mathrm{bs}(y_v) \le b^{6n+2}$, the vector bound is at most $b^{60h+92+6n+2} \le b^{2n^2}$ because $h \ge 10^5$. Multiplying by β^0 gives the cutoff β_c of sum one used in `Filling.classCutoff`. The box and the discrete counting are therefore compatible with the covolume formula (22.8).

We now prove the required uniformly bounded representative. The following elementary weak transference estimate is convenient.

Lemma 22.13 (Covering-radius estimate)**.** *For a full lattice Λ in a Euclidean space of real dimension m, if its Euclidean dual has shortest nonzero vector of length at least $\rho > 0$, then every vector of that space is within distance m^2/ρ of Λ.*

Proof. Minkowski's first theorem in Λ and its dual gives

$$\lambda_1(\Lambda)\lambda_1(\Lambda^*) \le 4/\operatorname{vol}(B_1)^{2/m} \le m,$$

since the Euclidean unit ball contains the cube of side $2/\sqrt{m}$. Choose a shortest vector v of Λ, which is primitive, and project orthogonally along its span. The projected lattice has rank $m-1$, since a primitive vector extends to a lattice basis; its dual is $\Lambda^* \cap v^\perp$ and has shortest vector of length at least ρ. By induction on dimension, approximate the projection of the given vector by a projected lattice element to within $(m-1)^2/\rho$. Choose a lift and round along v to make the other component at most $\|v\|/2 \le m/(2\rho)$. The sum of the two errors is at most m^2/ρ. This proves the assertion, including the one-dimensional base case. □

In the mixed embedding of the CM field, the Euclidean dual of the lattice $2\varpi^{-1}\mathcal{O}$ is $\mathcal{O}$ itself: the Euclidean pairing with $2\varpi^{-1}a$ is the trace pairing with $\varpi^{-1}a$, and the trace dual of $\mathcal{O}$ is the inverse different by definition. The sign of the generator under conjugation does not change the integral dual. The shortest nonzero vector of $\mathcal{O}$ has length at least one, as the product of its squared complex absolute values is the nonzero integer norm. Lemma 22.13, with $m = 2f$, gives for every tuple (z_v) an element $x \in \mathcal{O}$ such that

$$|z_v - \Phi_v(x)| \le (2f)^2 |\Phi_v(\varpi)| \quad \text{for all } v.$$

Indeed apply the lemma to the tuple $z_v/\Phi_v(\varpi)$ and the lattice $2\varpi^{-1}\mathcal{O}$, then multiply each coordinate by $\Phi_v(\varpi)$. More generally a finite adele can first be approximated modulo $\mathcal{O}_f$ by a rational element, then this argument is applied to the remaining infinite coordinates. This derives `exists_rat_near_of_mem_differentIdeal` in `Lib/Adelic/Transference/MassBasic.lean`.

For any $t = (t_v) \in U_\infty$ apply the following row algorithm to its frame matrices. To reduce the entry (a, b) of the unitriangular frame matrix, $a < b$, choose an integral scalar x by the preceding approximation and multiply on the right by the Eichler row generator of Lemma 22.12 with free row supported in the column f_b and scalar $-Nx$. Its row-integrality condition holds: for an isotropic column of the frame the lattice basis is unchanged; for $f_b = z$ the coefficient Nx clears the $1/N$ of the marked basis vector. Hence the generator belongs to Γ_c. Its normalized frame matrix is the elementary column operation subtracting from column b the normalized scalar

$$R_v \Phi_v(x), \qquad R_v = gN\, l_{vb}/l_{va} = gN\, \kappa_v^{a-b},$$

times column a. Apply the approximation to the tuple of entries divided by R_v. This makes the new entry (a, b) at most

$$|R_v|(2f)^2|\Phi_v(\varpi)| \le b^{23+60n+3} < b^{60h+91},$$

using $g \le b^{23}$, $N \le b^{60n}$, $2f \le b$, $|\kappa_v| \ge 1$ and $|\Phi_v(\varpi)| \le b$. Treat rows in descending order: this operation cannot change any row of larger index, because column a has zero entries below a. Treat all columns of the current row, then move to the next row. This constructs one rational $\gamma \in \Gamma_c$ for which all entries of $\mathrm{frameMatrix}(t_v\gamma_v)$ satisfy the stated bound.

For arbitrary y choose $t_v \in U_\infty$ with $\|t_v y_v\|_{\mathrm{op}} \le \mathrm{bs}(y_v)$: by definition some Bessel translate has norm plus one less than $\mathrm{bs}(y_v) + 1$, and remove its compact factor without changing the operator norm. For the rational γ just found, the normalized frame of $\gamma_v^{-1} y_v$ is $y_v^{-1}\gamma_v f_b = (t_v y_v)^{-1}(t_v\gamma_v f_b)$. Its length is at most $\|t_v y_v\|_{\mathrm{op}}\, nb^{60h+91} \le H(y, v)$. Thus the denominator of (22.9) is positive everywhere, as claimed. This completes the construction of the box partition `isBoxPartition_epsilon` in `New/Filling/BoxPartition.lean` with its explicit bound. Mere density of E in its finite adeles would not have sufficed to obtain this archimedean reduction.

In contrast, for strong approximation of the unipotent group with no archimedean bound, the elementary coordinates above suffice. The group U is generated by Heisenberg-type Eichler transformations along the successive isotropic rows. Over a field this follows by cancelling its free entries in the same triangular order as in the proof of (22.8); the skew part is arbitrary, since one can write any skew element as $t - \bar{t}$ (use $t = s/2$ over a field of characteristic zero). The same polynomial parametrizations over the finite adeles are continuous, even at the dyadic components. The diagonal copy of E is dense in its finite adeles by the ordinary weak approximation and Chinese remainder theorem, allowing denominators at finitely many places. Approximating the entries of the finitely many generating parameters therefore shows $U(E^+)$ is dense in $U(\mathbb{A}_f)$. Consequently for any adelic $x \in U(\mathbb{A}_f)$ and finite-adelic $y \in G(\mathbb{A}_f)$

there is a rational $u \in U(E^+)$ with $y^{-1}x^{-1}uy \in K(g)$, because this is an open neighborhood condition. This is `Filling.exists_besselU_approx` in `New/Filling/NearTerms.lean`, in the appropriate marked basis.

The normalized additive character used in that statement is important. Let ψ_f be the usual finite trace character, taken so that its product with the standard infinite trace character is trivial on E. Let

$$\chi(u) = \psi_f(-\sigma(u)/(g\varpi)),$$

where σ is the sum of the simple frame-root entries. The character $x \mapsto \psi_f(-x/\varpi)$ has conductor $\mathcal{O}_f$, by the definition of the different (the annihilator of the integral lattice for the unscaled trace pairing is the inverse different). On integral U its values modulo $K(g)$ are therefore well defined. The same is true in the marked lattice coordinates: the change of basis fixes the frame, so the simple root sum is unchanged. Conjugating by H also leaves this sum unchanged since H fixes the frame. For rational u, after the infinite rescaling the simple-root sum at place v becomes $\Phi_v(\sigma(u))/\kappa_v$. Consequently

$$\prod_v \mathrm{wave}(\Phi_v(\sigma(u))/\kappa_v) = \exp(2\pi i\ \mathrm{Tr}_{E/\mathbb{Q}}(\sigma(u)/(g\varpi))) = \chi(u).$$

We have used $\bar{\varpi} = -\varpi$ and $\kappa_v = g\Phi_v(\varpi)/(2\pi i)$. In particular the finite and infinite characters cancel under the adelic product with the required opposite signs. If x and a rational approximation u differ in the above congruence neighborhood for a class representative, then the values of χ agree: the quotient of the values is the scaled character of an integral simple-root sum in $g\mathcal{O}_f$. This gives `unipotentChar_eq_of_mem_principalLevel` and `exists_rat_approx_unipotentChar_eq` of `NearTerms.lean`. It is exactly the approximation needed for the orbital vanishing and for moving a Bessel element between class representatives.

22.10 Orbital cutoffs and Bessel transport between classes

We first give the precise quotient integration needed by the moment argument. For classes c, c' and $s \in \mathfrak{G}$, the set of terms is

$$\mathcal{T}(c, s, c') = \{\gamma \in G(E^+) : h_c^{-1}\gamma_f h_{c'} \in K,\ \mathrm{red}(h_c^{-1}\gamma_f h_{c'}) = s\}.$$

If T_0 is a subset of $G(E^+)$ stable under left and right $U(E^+)$ multiplication, set

$$A_{T_0}(c, s, c') = \sum_{\gamma \in T_0 \cap \mathcal{T}(c,s,c')} \int_{G_\infty} \beta_c(y)\beta_{c'}(\gamma_\infty^{-1}y)W(y) \otimes \overline{W(\gamma_\infty^{-1}y)}\, dy$$

where W is the product of the full or the cut local functions. We use the same absolute convergence of these orbital sums as in Section 20.9. The cutoff manipulations below are made first for nonnegative integrands of absolute values using Tonelli, and then for the vector integrals by absolute integrability. In particular, they do not constitute an independent proof of the uniform weighted summability used for the full moments.

A weight on a proper action of a countable discrete group can be changed without changing the integral of an invariant function, when its orbit sum has the same value. If F is invariant and both weights b_1, b_2 have sum one, insert $\sum_\delta b_2(\delta y)$ into the integral with weight b_1, interchange sum and integral, change variable by δ, and recover the integral with weight b_2. The same calculation holds for weights whose total sums on orbits agree. Apply this on each rational double coset for the commuting left and right actions of $\Gamma_c, \Gamma_{c'}$: if γ is its representative, its stabilizer is

$$I_\gamma = \Gamma_c \cap \gamma\Gamma_{c'}\gamma^{-1}.$$

Accounting for all the representatives of a double coset is equivalent to replacing the two cutoffs by one cutoff for I_γ. To check this, first sum over right translates of γ and use the cutoff identity for $\beta_{c'}$; then sum over the left translates modulo I_γ and use the corresponding cutoff identity for β_c. The covariance of the Bessel kernel (both characters are trivial on the congruence subgroups) allows the changes of variable in the integrals. Thus that double coset contributes the integral of

$$y \longmapsto W(y) \otimes \overline{W(\gamma^{-1}y)}$$

on $I_\gamma\backslash G_\infty$ with its quotient measure. The equality is initially one for finite unions of cosets with absolutely integrable summands; the absolute-value calculation extends it to the summable series. This is the argument of `hasSum_amplitudeSum_change_of_weight` in `New/Filling/AmplitudeSums.lean`.

Suppose $u \in U(E^+)$ and $u' = \gamma^{-1}u\gamma \in U(E^+)$. On the kernel in the last display, left multiplication by u_∞ contributes the character ratio $\chi(u)/\chi(u')$ (or its conjugate according to the Bessel sign convention). For the vanishing of the double coset integral it is necessary to use a map preserving the quotient measure on $I_\gamma\backslash G_\infty$: a rational element of the intersection need not normalize I_γ.

Here is a quotient argument avoiding such an assumption. Let $U_\gamma = U_\infty \cap \gamma_\infty U_\infty \gamma_\infty^{-1}$. This is the infinite group of the rational unipotent intersection, and I_γ is an arithmetic lattice in it. Its finite covolume follows from the same triangular-coordinate argument as for a unipotent group: a unipotent group over a characteristic-zero field has a central filtration with vector-group quotients, and intersecting any compact open subgroup at the finite places gives full rank lattices in the rational quotient coordinates. One way to obtain this filtration here is to take logarithms of the upper unitriangular matrices (a finite polynomial), intersect their nilpotent Lie algebras, and use the central series and the exponential polynomial; a lattice in each quotient admits bounded fundamental sets, and lifting bounded sets successively gives finite covolume. Rational approximation in these polynomial coordinates also holds. The local congruence condition defining I_γ is compact open in the finite intersection.

The function on G_∞ is invariant under I_γ. Upon disintegrating its quotient integral along $I_\gamma\backslash U_\gamma$, the inner integral on each orbit is a constant vector times the integral of the character ratio on that compact quotient. Since the ratio is trivial on I_γ, it descends to the quotient; if it is nontrivial on U_γ, its integral is zero by right Haar invariance of the quotient (which does not require I_γ normal). Rational approximation of finite characters and the equality of infinite and finite characters above show that nonagreement on rational points gives such nontriviality. Hence stable sets of terms for which the two characters never agree have zero total amplitude. In the notation of the source the agreement condition `Filling.CharsAgree` is exactly equality of $\chi(u)$ and $\chi(u')$ for every rational pair $u\gamma = \gamma u'$ in the unipotent groups.

The next transport identity concerns multiplying a term by a rational Bessel element. For $s \in \bar{S}_B$, fix c and lift s to $k \in K \cap S_B(\mathbb{A}_f)$. Decompose $h_c k = u_f h_f$ under UH. By the class decomposition of h_f, there are a rational $h \in H(E^+)$, a class d, and a $k_c \in K_c$ with $h_f = h h_d k_c$. Approximate u_f rationally by an element of U modulo the required neighborhood conjugated by h_f; this is the strong approximation proved in Section 22.9. Then the product $\delta = uh$ is a rational Bessel element lying in $\mathcal{T}(c, s, d)$. Thus for every c a Bessel lift over s exists.

For $s = 1$, a rational term in S_B forces $c = d$ and lies in Γ_c. Indeed its factor in H has position in K_c (integrality and congruence of the H factor follow by the block-diagonal argument of (22.6)), hence the two classes agree and the rational H element fixes a deeper-level class. Its stabilizer is trivial. This proves `eq_and_mem_classUnip_of_besselS` and, with the preceding approximation, `exists_besselS_mem_termsOver` in `New/Filling/NearTerms.lean`. In general the map $c \mapsto d$ produced by a lift of s is a permutation: if two classes have the

same d, multiplying one lift by the inverse of the other gives a Bessel term over the identity, forcing equality of the classes, and a finite injective self-map is bijective.

As a consequence of the quotient weight-change argument, transporting terms over s by such Bessel lifts permutes the class blocks and multiplies the corresponding coordinate factors by unitary compact type matrices and complex phases of absolute value one. More explicitly, if $\delta \in \mathcal{T}(c,s,d) \cap S_B$, right multiplication by δ^{-1} maps the appropriate sets of terms over $(c_0, s_0 s, d)$ bijectively to those over (c_0, s_0, c); it conjugates the right unipotent congruence subgroup at d to the one at c because the finite position is in K and $K(g)$ is normal in K. The replacement of the cutoff for d by the transported cutoff for c is valid by orbit sums. The tensor factor in the integral changes by the representation of the compact factor of δ and its character. Both have norm one. This yields precisely the covariance needed for the equality of Hilbert–Schmidt block sums below. It is the mathematical content of `amplitudeSum_move` in `New/Filling/AmplitudeSums.lean`.

There is no Bessel-subgroup contribution over 1 except the class-diagonal integral. Indeed for $c = d$ the surviving terms are exactly Γ_c, and summing the two cutoff factors reduces the integral to ν times the normalized Bessel inner products on $U_\infty \backslash G_\infty$. Schur orthogonality with the chosen normalization of the Bessel function gives the identity on the coordinate space, so

$$\nu^{-1} A_{S_B}(c, 1, d)_{ij} = \delta_{c,d}\delta_{i,j} \quad \text{for the full functions.}$$

This is `amplitudeSum_besselS_one`, and explains the identity term in the first moment. The same covariance and weight-change argument works for the cut functions, although the coordinate inner product need then no longer be exactly the identity.

22.11 Agreeing characters and propagation of a shared flag vector

We now prove the algebraic elimination of degenerate terms, which was an input in Section 20.6. The notation here is rational: $F = (f_0, \dots, f_h)$ is the standard frame, $f_h = z$, and Fx denotes the corresponding linear combination of its columns. For a rational γ put

$$Z = F^* J_N \gamma F, \qquad D_j = \det Z_{\{0,\dots,j-1\}}, \qquad \mathcal{D}(\gamma) = D_{h+1}\bar{D}_{h+1} - N^2 D_h \bar{D}_h,$$

where J_N is the standard rational Gram matrix with middle entry $-N$ on z (and $D_0 = 1$).

Lemma 22.14 (Shared flag-vector propagation)**.** *Suppose the unipotent characters agree for every rational pair $u\gamma = \gamma u'$. If $Fx = \gamma Fy$ and both coefficient vectors vanish at indices $> a$, with $x_a = c$ and $y_a = c'$, then for every $b \geq a$ such a common vector exists with leading coefficients still c, c'. In particular $c\bar{c} = c'\bar{c}'$.*

Proof. It suffices to propagate from $a < h$ to $a + 1$. Let ℓ_0 be any rational linear form on V vanishing on the subspace spanned by $f_i, \gamma f_i$ for $i \leq a$. Use nonsingularity of J_N to write $\ell_0(v) = h(w, v)$. With $v_0 = Fx = \gamma Fy$, the Eichler transformations

$$E(r) = 1 + r v_0 h(w, \cdot) - \bar{r}\, w h(v_0, \cdot) - r\bar{r}\,\tau h(w, w) v_0 h(v_0, \cdot), \qquad r \in E,$$

lie in U, and their conjugates by γ^{-1} also lie in U. Here $\tau + \bar{\tau} = 1$, and no integrality is required. To verify the assertion, the support of v_0 in the isotropic frame and the vanishing of ℓ_0 on the initial flag imply upper unipotence in the block order. Expanding the Gram form gives unitarity as in Lemma 22.12, with v_0 isotropic and $h(v_0, w) = 0$. On the simple

superdiagonal entries of the frame only the term $rv_0h(w,\cdot)$ contributes, since $h(v_0,f_j)=0$ for every frame vector. Thus the difference of the two simple-root sums is

$$r\,\ell_0\left(\sum_{i\le a}x_if_{i+1}-\sum_{i\le a}y_i\gamma f_{i+1}\right).$$

Agreement of the rational characters for all r forces the parenthesized vector to lie in the annihilator of every such ℓ_0: the finite character on $\mathbb{A}_f$ is nondegenerate, and E is dense in $\mathbb{A}_f$, so if $\psi_f(rt/(g\varpi))=1$ for all $r\in E$ and $t\in E$, then $t=0$. The annihilator of the forms vanishing on a subspace is that subspace itself. Consequently the displayed vector belongs to $\langle f_i,\gamma f_i : i\le a\rangle$. Moving these lower terms to the two sides gives a common vector supported at indices $\le a+1$ with leading coefficients c,c' at $a+1$, as required.

Iterate up to $b=h$. The norm of a vector in the frame span is $-N$ times the norm of its last coefficient, because all other frame entries are isotropic and orthogonal to the entire frame. The two equal vectors at that last stage therefore have $-Nc\bar{c}=-Nc'\bar{c}'$, proving the last assertion. □

This is `Filling.hasCommonFlagVector_of_le` of `New/Filling/CrudeEnergy.lean`, using `exists_mem_besselU_eq_eichlerMap` of `New/Filling/EichlerFlag.lean`. In particular no assumptions about the norm of the original rational matrix are needed in this propagation lemma.

There is always a simple nonnegativity relation for $\mathcal{D}(\gamma)$. Suppose $Zy=Df_{\mathrm{last}}$ in coefficient notation (that is, all entries except the last of Zy vanish). Put $t=\gamma Fy$. Its e_a^* coordinates vanish, and its z-coordinate is $-D/N$. Write t_0 for its component on the positive definite X. Unitarity gives, at every infinite embedding,

$$N\|t_0\|_v^2=|\Phi_v(D)|^2-N^2|\Phi_v(y_h)|^2. \tag{22.10}$$

In particular equality of the right-hand terms at any place implies $t_0=0$ (all X-coordinates are rational and the ordinary positive norm is definite). Taking $y=\mathrm{adj}(Z)(0,\dots,0,1)^t$ shows $D=D_{h+1}$ and $y_h=D_h$. Therefore $\mathcal{D}$ is nonnegative at every infinite place. If $\mathcal{D}\neq 0$, then $D_{h+1}\neq 0$. This proves the discriminant and normalization content of `norm_sq_middle_part_eq`, `map_termDisc_eq_norm` and `hasLevel_last` in `NearTerms.lean`.

Proposition 22.15 (Levels of all relevant degenerate terms)**.** *Under rational character agreement, if $\mathcal{D}(\gamma)=0$, there is an integer $j\in\{0,\dots,h\}$ for which $D_j\neq 0$ and a scalar $\alpha\bar{\alpha}=1$ such that for every $a\ge j$ the two frames share a vector with leading coefficients $\alpha,1$ at index a. Together with $\mathcal{D}(\gamma)=0$ these are precisely the conditions of `Filling.HasLevel` for level j in `CrudeBoundDefs.lean`. If $\mathcal{D}\neq 0$, the term has level $h+1$.*

Proof. If $\mathcal{D}=0$, first find a nonzero vector shared by the frame spans. When $D_{h+1}\neq 0$, use the adjugate vector just used in (22.10). Its image γFy lies in the span of F since its positive definite component vanishes. If $D_{h+1}=0$, choose any nonzero y in the kernel of Z instead. Formula (22.10) now implies $y_h=0$ and $t_0=0$; again the image is a nonzero vector in the span of F.

Write such a relation as $Fx=\gamma Fy$ and let a be the largest index with either coefficient nonzero. By Lemma 22.14, the leading coefficients have equal norms, so both are nonzero. Their ratio $\alpha=x_a/y_a$ has norm one. After normalizing by y_a, the propagated relation shows that the same ratio occurs at every index $\ge a$. In fact the ratio of leading coefficients of any nonzero shared relation is unique: propagate two such ratios to index h, normalize their last y-coefficients to one, and subtract. If the ratios were unequal the resulting common vector

would have one last coefficient zero and the other nonzero, contradicting equality of norms. Thus let j be the least index for which a shared relation occurs; it has the stated common ratio at every index $\geq j$.

We show $D_j \neq 0$. If $j = 0$ this is the convention $D_0 = 1$. Otherwise suppose the $j \times j$ leading block is singular and let $y \neq 0$ be a vector supported on indices $< j$ in its kernel. Put $t = \gamma F y$. Its pairings with $f_0, \dots, f_{j-1}$ vanish. For every $a \geq j$ there is a common vector whose coefficient at f_a on the standard side is $\alpha \neq 0$, with other terms only at lower indices. The pairing of that vector with t is zero on the γ-frame side, since $y_h = 0$. By induction up the standard frame this implies $h(f_a, t) = 0$ for all a, including the last vector. Hence $Zy = 0$. Formula (22.10) shows $t \in \operatorname{span} F$; write $t = Fx$. Taking the largest index of a nonzero coefficient in x or y, the propagation lemma says neither leading coefficient can vanish. It follows that this largest index is $< j$, contradicting minimality. Thus $D_j \neq 0$.

If $\mathcal{D} \neq 0$ the leading full minor is nonzero as noted above, the normal-form condition at level $h + 1$ is empty, and the condition on $\mathcal{D}$ is exactly the top condition. □

The preceding proposition is `Filling.CharsAgree.exists_hasLevel` in `New/Filling/CrudeEnergy.lean`. The set of terms of each level is stable under the two unipotent actions: these actions change the frame columns by unitriangular matrices, preserving leading-coefficient relations up to lower terms and preserving every leading principal determinant of Z by multiplying on the corresponding sides by unitriangular matrices. In particular the discriminant $\mathcal{D}$ is preserved. This also gives the stability statements in `New/Filling/StableCells.lean` required to split the energy according to levels.

22.12 The link as a Hermitian Lagrangian and its congruence defect

The link is the image of $L^\perp \cap (\gamma L)^\perp$ in

$$(L^\perp/L) \oplus ((\gamma L)^\perp/\gamma L) \simeq M \oplus M$$

under $t \mapsto ([t], [\gamma^{-1}t])$, using γ^{-1} to identify the second middle quotient. Denote it by $\mathcal{L}_\gamma$. Give $M \oplus M$ the Hermitian form $T \oplus (-T)$, $T = \operatorname{diag}(1_k, -N)$. For two vectors in the defining intersection, their first Hermitian pairing agrees with their second by unitarity of γ. Thus this image is isotropic. If $a = \dim(L \cap \gamma L)$, the dimension of the defining intersection is

$$r - \dim(L + \gamma L) = r - (2h - a) = k + 1 + a.$$

The kernel of the map is precisely $L \cap \gamma L$. Consequently the image has dimension $k + 1$, half the ambient dimension. It is therefore equal to its orthogonal complement. This proves `memm_link_iff_orthogonal` in `StableCells.lean` and `finrank_link` in `BoxPartition.lean`. Invariance under U on both sides follows because U preserves L, acts trivially on $L^\perp/L$, and applying a left translate of a defining vector gives a corresponding translate for the new link. On a big-cell representative $w_p\sigma$, with $\sigma \in U(M)$ and the Weyl element w_p exchanging L and $L^\vee$, the common orthogonal intersection is M; the link is the graph of σ^{-1} and equality of such links is equivalent to equality of the middle transformation.

Let $\ell_z : M \to E$ be the form $h(z, \cdot)$. The source puts the positive definite metric on $M \oplus M$

$$Q_{\text{link}} = \tfrac{1}{2}(T \oplus T) + N^{-1}(\ell_z \oplus \ell_z)^*(\ell_z \oplus \ell_z).$$

Here the second summand is the sum of the squares of the two separate forms, not the square of their sum. In the standard coordinates this matrix is diagonal: the positive definite

coordinates have weight $1/2$ and each z-coordinate has weight $-N/2+N=N/2$. Hence it is positive at every infinite embedding. In particular its Gram determinants on any nonzero subspace are nonzero; it is the positive height used in the pair-counting estimates, not the isotropic link pairing itself.

We prove the particular residue-field defect bound that controls the congruence gain in those estimates. Fix a split level place w (or its conjugate) with residue field $\mathbb{F}_q$, and let $\bar{\gamma}$ denote the reduction of the integral finite position of γ in the marked basis. Write F also for the reduced frame. The relevant form matrices at the two factors are related by the adjoint inverse; one may choose whichever factor is specified by the vector-fixer condition.

Lemma 22.16 (Residue-field propagation into the two last columns)**.** *Suppose the reduced position has agreeing characters in the sense that any two integral unipotent elements with reductions conjugate by it give the same unipotent character. Suppose a nonzero frame vector is fixed by $\bar{\gamma}$ at one specified split factor. Let P be any matrix of two row forms on the reduced space each vanishing on the standard and translated isotropic frame vectors. Then the 2×2 matrix of their values on $z,\bar{\gamma}z$ is singular.*

Proof. Work at the specified split factor; a vector fixer written using inverse adjoint simply means work at its conjugate factor. If the 2×2 determinant were nonzero, any shared relation $Fx=\bar{\gamma}Fy$ would have $x_h=y_h=0$, by applying the two forms and subtracting. The fixed nonzero frame vector supplies such a shared nonzero relation with equal coefficient vectors.

Suppose one has a shared relation with both last coefficients zero and largest nonzero standard coefficient x_a at $a<h$. Let m be any residue-field linear form annihilating f_i for $i\le a$ and $\bar{\gamma}f_i$ for $i<h$. The transvections

$$1+u(Fx)m, \qquad 1+u(Fy)m\bar{\gamma} \quad (u\in\mathbb{F}_q)$$

are conjugate by the reduced position and strictly upper block unitriangular off the identity. Indeed the left vector is supported in earlier isotropic rows and the right form vanishes on the initial flag through those rows. To lift them to integral U, lift these vectors and forms separately at the specified split factor, keeping their indicated zero entries zero. Their rank-one cross terms square to zero. At the opposite split factor set the unitary inverse adjoint, and at every other finite place use the identity. This produces integral unipotents. The opposite factor of a transformation $1+u(Fx)m$ has its nonidentity entries in columns dual to the support of Fx; these are isotropic dual columns, not frame columns. Indeed the inverse is $1-u(Fx)m$, and the adjoint inverse has its changed columns given by $(Fx)^*J$, which is supported on the dual columns. Thus at that conjugate factor the simple-root sum of the frame entries vanishes, and similarly for the second transvection.

At the selected factor the additive character is primitive: the different-normalized finite character has conductor $\mathcal{O}_f$, and multiplication by $1/g$ tests the full residue field at that place. Agreement on varying u consequently implies

$$m\left(\sum_{i\le a}x_if_{i+1}-\sum_{i<h}y_i\bar{\gamma}f_{i+1}\right)=0.$$

Here the sums are the simple superdiagonal entries of the two rank-one operators. As m ranges over the indicated annihilator, the parenthesized vector is in the span of the standard frame through a and the translated isotropic frame. Moving these terms to the two sides gives a new shared relation with standard coefficient at $a+1$ equal to $x_a\neq 0$, other standard coefficients of lower index, and translated coefficients allowed through the last vector. The alleged invertibility of the two-form determinant forces both last coefficients zero in every shared

relation. Iterating the propagation gives a relation with nonzero standard last coefficient, a contradiction. Hence the determinant is singular. □

This is the residue-field argument in `Filling.le_genIndex_det_of_fixesNormAtLeast`, `New/Filling/BoxPartition.lean`. Notice its distinction from Lemma 22.14: over one split factor it is not necessary to recover a norm relation or to propagate with exactly matching upper supports on the two sides. The invertible two-form determinant was used instead to exclude the last coordinates of every shared relation.

If the two forms in Lemma 22.16 are finite adelic integral forms annihilating the two isotropic frames, put δ equal to the determinant of the matrix of their values on the two last frame vectors. For every place at which the reduced position satisfies the corresponding vector-fixer condition and has agreeing characters, δ is divisible by that prime. Therefore

$$[\delta] \geq \prod_{w \text{ fixed}} q_w \tag{22.11}$$

if δ is an idele; here the generalized index on the left is the absolute ideal index. This follows either by taking the local valuations or by mapping the quotient of $\mathcal{O}_f$ by $\delta\mathcal{O}_f$ onto the product of the corresponding residue fields. Requiring agreement in the finite reductions, rather than only agreement on rational intersections, is what permits the use of the separate residue-field tests.

Finally we record a direct comparison of this type of ideal index with the positive link height. For two rational rectangular matrices A, B of the same full column rank in a positive definite Hermitian space with form Q, at every infinite embedding the exterior-power Cauchy–Schwarz inequality gives

$$|\det(A^*QB)|^2 \leq \det(A^*QA)\det(B^*QB). \tag{22.12}$$

If their adelic changes of basis are $a, b \in \mathrm{GL}_s(\mathbb{A}_f)$, multiplicativity of the determinant and the product formula give

$$[\det(a^*A^*QBb)]^2 \leq [\det a]^2[\det b]^2|\operatorname{Norm}_{E/\mathbb{Q}}\det(A^*QA)||\operatorname{Norm}_{E/\mathbb{Q}}\det(B^*QB)|.$$

Indeed the finite generalized index of a nonzero rational number is the absolute field norm, and taking the product of (22.12) over all complex embeddings, together with their conjugates, gives the required norm inequality. This is `liouville_gram` in `BoxPartition.lean`. It explains the appearance of powers of positive sublattice discriminants when a finite index such as (22.11) is large. The remaining precise comparison of the pair invariants and the class-size estimate in `LinkPairing.lean` and `BlockBound.lean` are not automatic from this inequality alone, since those bounds require controlling the ranks and saturation indices of several simultaneous lifts.

22.13 Saturated lattice heights and their zeta sum

We prove the subspace counting estimate needed both for crude link weights and for the sharp global sieve. Let $V_0 = E^m$ with its standard positive definite Hermitian form, and let $\mathcal{M}$ be a finite adelic unimodular Hermitian lattice in V_0. Write $L_{\mathcal{M}} = V_0 \cap \mathcal{M}$. If $W_0 \subset V_0$ is a rational subspace of dimension s, choose an E-basis matrix Y for W_0, and a finite adelic matrix $a \in \mathrm{GL}_s(\mathbb{A}_f)$ such that $Ya\mathcal{O}_f^s = W_0(\mathbb{A}_f) \cap \mathcal{M}$. Such a basis exists locally by the elementary divisor theorem over each discrete valuation ring; outside finitely many places the rational basis is already integral and primitive. Define the height

$$d_{\mathcal{M}}(W_0) = [\det a] \prod_{v|\infty} \det\left(\Phi_v(Y)^*\Phi_v(Y)\right). \tag{22.13}$$

The expression does not depend on the choices: changing the rational basis by $P \in \mathrm{GL}_s(E)$ multiplies the product of Hermitian determinants by $|\operatorname{Norm}\det P|$ and the finite index by its reciprocal, while changing an integral local basis has finite index one. In the source this is the discriminant $\sqrt{[\det(Ya)^*(Ya)]}$, `sublatticeDisc` of `New/Filling/GramLiouville.lean`. The equalities follow by taking determinants and the product formula at all complex places.

The Euclidean covolume in $W_0(\mathbb{R})$ of its rational lattice $W_0 \cap L_{\mathcal{M}}$ equals

$$d_{\mathcal{M}}(W_0)\,\mathrm{vol}(\mathcal{O}_f)^{-s}. \tag{22.14}$$

Indeed the preimage of its lattice in E^s has finite coordinates $a\mathcal{O}_f^s$, hence ordinary covolume $[\det a]\,\mathrm{vol}(\mathcal{O}_f)^{-s}$. At each complex place the real Jacobian of the injective map Y onto its image is $\det(Y^*Y)$ (singular values occur twice in real coordinates). Multiply these Jacobians. This proof handles arbitrary nonprincipal lattice modules by working locally.

Two elementary observations are useful.

- If Ya' is another integral full rank array of the same subspace then $Ya' = YaP$ with P integral at the finite places, since the columns Ya form a saturated integral basis. The corresponding Gram determinant is multiplied by $[\det P]^2$. If in addition a linear image of the columns of Ya' lies in some finite-index submodule of the image lattice of Ya, the index i of that submodule divides the index of the column sublattice; in particular $i \le [\det P]$. Therefore the squared Gram index of Ya' is at least $(d_{\mathcal{M}}(W_0)i)^2$. This proves the primitive-basis and index calculation in `sublatticeDisc_mul_index_sq_le`.

- Heights of complementary orthogonal subspaces agree:

$$d_{\mathcal{M}}(W_0) = d_{\mathcal{M}}(W_0^{\perp}). \tag{22.15}$$

  To check it locally, extend the saturated integral basis of W_0 to a basis matrix M of $\mathcal{M}$. This extension is possible because a saturated submodule over a discrete valuation ring has torsion-free quotient and is a direct summand. The last $m-s$ columns of the Hermitian dual basis give a basis of the orthogonal-complement lattice: they pair to zero with the first s columns, and unimodularity gives their integrality. Their Gram determinant is the complementary block minor of the inverse full Gram matrix. By Jacobi's identity it equals the Gram determinant of the first s columns divided by the full Gram determinant. The latter is a local unit at every finite place by unimodularity. Take the generalized indices and square roots, which is legitimate because these determinants are nonzero (the rational spaces are positive definite). This gives (22.15).

Lemma 22.17 (A quantitative Siegel lemma for the orthogonal complement)**.** *Suppose* $\dim_E W_0 = m-s$, $1 \le s \le m$. *Then its orthogonal complement has an* E*-basis* $x_1, \dots, x_s$ *in* $L_{\mathcal{M}}$*, with Euclidean mixed-embedding norms satisfying*

$$\left(\prod_{i=1}^{s} \|x_i\|\right)^{2f} \le (8fs\ \mathrm{rd}_E)^{fs} d_{\mathcal{M}}(W_0), \qquad \mathrm{rd}_E = |D_E|^{1/(2f)}.$$

Proof. Apply Minkowski's second theorem to the Euclidean lattice $W_0^{\perp} \cap L_{\mathcal{M}}$ in real dimension $2fs$. If its successive minima are $\lambda_1, \dots, \lambda_{2fs}$ for the ordinary Euclidean ball, choose successively E-linearly independent vectors of norm at most $\lambda_{2f(i-1)+1}$ for $1 \le i \le s$. This is possible because the previously chosen E-span has real dimension $2f(i-1)$, whereas the next successive minimum gives more independent real vectors. Since the minima are nondecreasing,

the product of their chosen powers $2f$ is at most the product of all the real successive minima. By Minkowski's second theorem and (22.14)–(22.15),

$$\prod_{j=1}^{2fs} \lambda_j \le \frac{2^{2fs}(fs)!}{\pi^{fs}} \left(\frac{|D_E|^{1/2}}{2^f}\right)^s d_{\mathcal{M}}(W_0) \le (8fs\ \mathrm{rd}_E)^{fs} d_{\mathcal{M}}(W_0).$$

In the last step use $(fs)! \le (fs)^{fs}$, $|D_E|^{s/2} = \mathrm{rd}_E^{fs}$, and $\pi \ge 1$. This proves the result. □

This recovers the statement `Filling.siegel_lemma` in `New/Filling/NearTerms.lean`; the field-degree powers are important, but no uniform dependence on the ideal class is required from a separate choice of generators.

Theorem 22.18 (Sublattice height zeta bound)**.** *With the same notation, over all E-linear subspaces of codimension s,*

$$\sum_{W_0} d_{\mathcal{M}}(W_0)^{-(m+1)} \le (200fm\ \mathrm{rd}_E)^{f(m+1)s}.$$

For $s = 0$ the sum consists of the full space with height one.

Proof. Every nonzero vector x of $L_{\mathcal{M}}$ has mixed-embedding norm at least one. Indeed its Hermitian norm $h(x, x) \in E^+$ is totally positive and integral, since its pairing with the unimodular integral lattice is integral. Hence the product of its positive real conjugates is an integer at least one, and their sum is at least one by the arithmetic-geometric mean inequality. Distinct lattice vectors are also separated by distance one.

In a Euclidean space of real dimension d, a set of unit-separated points in a ball of radius R has at most $(1 + 2R)^d$ elements, by packing disjoint balls of radius $1/2$ and comparing volumes. Nonzero lattice points with norms in $[2^j, 2^{j+1})$ consequently number at most $5^d 2^{jd}$. For $p \ge d + 2$ this gives

$$\sum_{0 \ne x \in L_{\mathcal{M}}} \|x\|^{-p} \le 5^d \sum_{j \ge 0} 2^{-j(p-d)} \le 5^p.$$

Use $d = 2fm$, $p = 2f(m + 1)$.

For each subspace W_0 select one ordered basis $\boldsymbol{x}$ of its orthogonal complement as in Lemma 22.17. The selection is injective in W_0, because the span of the tuple determines $W_0^{\perp}$. Rewrite that lemma as

$$d_{\mathcal{M}}(W_0)^{-(m+1)} \le (8fs\ \mathrm{rd}_E)^{fs(m+1)} \prod_{i=1}^{s} \|x_i\|^{-2f(m+1)}.$$

Sum over the selected tuples, and enlarge to all tuples of s nonzero lattice vectors. The separated-point sum bounds this by $(8fs\ \mathrm{rd}_E)^{fs(m+1)} 5^{2fs(m+1)} \le (200fm\ \mathrm{rd}_E)^{fs(m+1)}$. All summands are nonnegative and the bound holds on every finite subset, which also proves convergence of the infinite sum. □

This proves `tsum_sublatticeDisc_inv_le`, `New/Filling/NearTerms.lean`. It accounts for the denominator lattices in the discriminant counts, not just for points with integral coordinates. In particular rational lines in E^2 can have varying nonprincipal denominator ideals without invalidating this summation.

22.14 The global frequency estimate and explicit sieve budget

We apply Theorem 22.18 to the nonzero rational T-integers y used as frequencies, where T is a conjugation-stable finite set of finite places. Write

$$[y]_T = \prod_{w\in T} |y|_w^{-1}, \quad d_y = [\mathcal{O} : \mathcal{O} \cap y^{-1}\mathcal{O}], \quad L_K(t) = t^K \min(1,t)^6.$$

The denominator index d_y only involves places in T, since y is integral elsewhere. The height of the line through $(1,y)$ in E^2 with its standard Hermitian lattice is

$$d(E(1,y)) = d_y \prod_v (1 + |\Phi_v(y)|^2). \tag{22.16}$$

Indeed its saturated column lattice consists exactly of the vectors $x(1,y)$ with $x \in \mathcal{O} \cap y^{-1}\mathcal{O}$. This proves the formula also for nonprincipal denominator ideals, by taking the local basis of that fractional ideal and then the generalized index as in (22.13).

Suppose $|c_v|^{-1} \le |x_v| \le B$ with $B \ge 1$. The product formula and integrality off T give $[y]_T^{1/2} \le \prod_v |\Phi_v(y)|$. Also

$$\min(1, B/t)^6 \le ((1+B^2)/(1+t^2))^3 \quad (t > 0).$$

Hence for every integer $K \ge 0$,

$$\frac{[y]_T^{K/2}}{d_y^6} \prod_v L_K(|c_v\Phi_v(y)|^{-1}) \le \left(\prod_v |x_v|^K (1+B^2)^3\right) d(E(1,y))^{-3}.$$

Here we used $d_y^{-6} \le d_y^{-3}$. The line map $y \mapsto E(1,y)$ is injective. Apply Theorem 22.18 with $m = 2$, $s = 1$. Since $400f \le b$ and $1 + B^2 \le bB^2$ under the filling hypotheses, every finite sum of these frequency terms is bounded by

$$\left(\prod_v |x_v|^K (bB^2)^3\right) (b \operatorname{rd}_E)^{3f}. \tag{22.17}$$

This derives `sum_frequencies_le` in `New/Filling/GlobalSieve.lean`, with its exact damping exponent and the correct denominator index.

We now verify the numerical input into the one-sided sieve induction of Section 21.10; in particular the two broad bounds in that section are no longer left as independent estimates. We use $n = h + 1$ as above, $K_1 = k - n^7$, $K_0 = n^6$. The standing hypotheses include

$$b \ge 1000, \quad n \ge 10^5 + 1, \quad k \ge 1000(2n)^9, \quad k \le f \le b/1000, \quad \operatorname{rd}_E \le b, \quad b^{19} \le g \le b^{23},$$

as well as $N \le b^{60n}$, $1 \le |\kappa_v| \le b^{23}$. Put $s_{va} = \kappa_v^{2(h-a)}/N > 0$ as in (22.8); the source calls these `frameScaleSq`. The per-entry Jacobian in the Poisson argument is

$$C_a = \operatorname{vol}(\mathcal{O}_f) / \prod_v s_{va}^2.$$

The nonzero frequency argument at step a is $(c_v\Phi_v(y)/s_{va})_v$ at infinity and y at the finite places. Indeed the character on one normalized infinite variable is $\operatorname{wave}(z) = \exp(z - \bar{z})$, and the rational global character with frequency y is obtained at infinity by $2\pi i\, \Phi_v(y)/\Phi_v(\varpi)$. Thus the Poisson choice satisfies $|c_v|^{-1} = |\kappa_v|/g$. For unequal entries the difference is $y = g^{-1}(t^{-1} - u^{-1})$, and the infinite factor has $c_v = g$, so again $|c_v|^{-1} \le |\kappa_v|/g$. These are the

two cases of the frequency test (21.20). In either case the inverse magnitude of the infinite scale, after division by s_{va}, is bounded by

$$|x_v| = |\kappa_v|^{2h+1}/(gN) \le b^{46n}.$$

This is the quantity used in the numerical hypothesis `prod_band_le_inv_deg`.

The archimedean frequency majorant given by the local bandlimit input has one factor

$$\Theta\, b^{2000n^9} L_{K_1}(|c_v \Phi_v(y)/s_{va}|^{-1}), \qquad \Theta = \operatorname{vol} U(k)\, \pi^{(2h+1)k} / \operatorname{vol} U(r).$$

Its finite factor is $[y]_T^{K_1/2} d_y^{-6} b^{n^9 f}$. Applying (22.17) with $B = b^{46n}$, the entire frequency sum is at most

$$\varepsilon = \left(\prod_v \Theta |x_v|^{K_1} \right) b^{(2001n^9 + 276n + 9)f}. \tag{22.18}$$

The exponent $276n + 9$ contains the product of the $B^6 b^3$ term and $(b \operatorname{rd}_E)^3$ at each place.

For the head-weight estimate (21.21), let A be the finite set of rational units used in the induction, whose elements and inverses are T-integral. For a head $(p_0, \dots, p_{i-1}) \in A^i$, put $D_j = p_0 \cdots p_{j-1}$ for $1 \le j \le i$. These partial products uniquely determine the head. The local weight is the product over j of damped powers with exponent K_0 at inverse absolute value of $D_j \prod_{a<j} s_{va}$, and the finite weight is the product of $[D_j]_T^{K_0/2}$ if each D_j is integral at T, zero otherwise. In the nonzero case D_j is globally integral, so its denominator index is one. Thus the sum over heads is bounded by the product of the sums over the allowed individual D_j, by injectivity of the partial-product map.

For each such sum use (22.17) with inverse scale bounded by N^i, because each $s_{va} \ge N^{-1}$. Set $B = N^i \le b^{60ni}$. With $K = K_0$ each sum is at most $b^{f(9+60ni(K_0+6))}$. For the square of the head sum the exponent is therefore at most $f(18i + 120ni^2(K_0 + 6))$. On the other hand

$$C_a \ge b^{-(92n+1)f}$$

because $s_{va} \le b^{46n}$ and $\operatorname{vol}(\mathcal{O}_f) = (2/\operatorname{rd}_E)^f \ge b^{-f}$. Dividing the squared head bound by $\prod_{a<i} C_a$, and using $i \le h < n$, $K_0 = n^6$, gives an exponent at most

$$f(120n^9 + 720n^3 + 92n^2 + 19n) \le 121 f n^9.$$

Thus one may take $\Gamma = b^{121 f n^9}$, including the empty head at $i = 0$.

By the admissibility condition for the product of band constants,

$$\left(\prod_v \Theta |x_v|^{K_1} \right) b^{7kf} \le d_\infty^{-1}, \qquad d_\infty = \prod_v d_{\lambda_v}.$$

Combining this with (22.18), the bound for Γ, and $k \ge 1000(2n)^9$, we obtain

$$\Gamma\varepsilon \le d_\infty^{-1} b^{-7kf + (2122n^9 + 276n + 9)f} \le d_\infty^{-1} b^{-7kf + 2123 n^9 f}.$$

Since $2h + 1 \le b$ and $(k - 2123n^9)f \ge 11$,

$$(2h+1)\Gamma\varepsilon \le d_\infty^{-1} b^{-6kf - 10}. \tag{22.19}$$

The bottom constant of the product sieve system is the product of $(1 + b^{-10})/d_{\lambda_v}$ at infinity and the specified finite bottom constant e. Applying the one-sided Poisson induction of Section 21.10, with the parameters just proved, yields

$$\Re \sum_{t \in U,\ p,q \in A^h} J'_{h+1}(p,t;q,t) \prod_v J_{v,h+1}(p,t;q,t) \le \left(\prod_{a=0}^{h} C_a \right) \left(e \prod_v \frac{1 + b^{-10}}{d_{\lambda_v}} + \frac{b^{-6kf-10}}{d_\infty} \right),$$

where the arguments have the rational and scaled infinite interpretations of the source and $U \subseteq A$ is arbitrary. This is `Filling.sieve` in `New/Filling/GlobalSieve.lean`. In particular the global frequency estimate, partial-product bound and resulting power-saving error are derived here. The existence of the actual local positive kernel systems with the stipulated bottom and oscillatory bounds is a separate input, not a consequence of this global estimate.

22.15 The finite zero form and central projections

We discuss the zero-frequency finite group estimate separately because it is also needed for the projection onto good finite types in the first moment. Consider a conjugation-stable finite product of local fields containing the selected pairs over g. In it denote the integral compact subgroup by K_T, the level group by $K_T(g)$, and choose the local group Haar measure M. The function called `IntegralPlace.phase` in `New/Filling/TopFloor.lean` is

$$\phi(x) = \begin{cases} \psi(\sigma(u)/g), & x \in uK_T(g),\ u \in U, \\ 0, & \text{otherwise.} \end{cases}$$

It is well defined by the conductor of the self-dual additive character and the additivity of the simple-root sum. Restricted to K_T, it is the finite function on $\mathfrak{G}$ equal to $\chi(s)$ on $\bar{U}$ and zero outside $\bar{U}$. The zero form for an intermediate compact subgroup $K_T(g) \subset Z \subset K_T$ is

$$\mathcal{K}(Z) = \int_{G_T} \left| \int_Z \phi(yxy^{-1})\, dM(y) \right|^2 dM(x). \tag{22.20}$$

The following bottom estimate is still a quantitative finite-place Fourier-floor input, `floorZeroForm_le_main_term` in `TopFloor.lean`. For $t + 10^{10}n^4 \le k$ and $t \ge 0$, it says

$$\mathcal{K}(Z) \le M(Z)M(K_T(g))(M(Z) + e_T^{t+1}2^{-n}), \qquad e_T = \mathrm{vol}(R_T \setminus R_T^{\times}). \tag{22.21}$$

Here $R_T = \prod_{w \in T} \mathcal{O}_w$ with additive volume one in the local coordinate normalization. Also $M(K_T) \ge 2^{-n}$ when T consists of chosen level pairs; the source obtains this from the normalized box measure comparison. We retain that comparison as an input in applying (22.21). On the other hand the reverse bound

$$\mathcal{K}(Z) \ge M(Z)^2 M(K_T(g)) \tag{22.22}$$

is immediate since for $x \in K_T(g)$, all its conjugates by Z belong to $K_T(g)$ and have phase one.

We first show how the zero form yields the saving for fixed vectors on finite factors. At one selected pair above q_0, let Z be the preimage of the fixer of a nonzero frame vector (standard or inverse adjoint). Then $M(K_T)/M(Z) = q_0^r - 1$ by the transitive linear action on nonzero vectors. The measure of the nonunits at any product of the chosen split pairs is at most $4fb^{-10} \le b^{-9}$: the set where some coordinate is a nonunit is the union of the corresponding maximal ideals, each of measure $1/q_j$. For the single pair, use (22.21) with $t = k - 10^{10}n^4$. In the standing size range $10^{10}n^4 \le n^6$, $k \ge 1000(2n)^9$, $b^{10} \le q_0 \le b^{11}$. Thus

$$\frac{M(Z)}{M(K_T)} = \frac{1}{q_0^r - 1} \le \frac{2}{b^{10r}}, \qquad e_T^{t+1}\frac{2^{-n}}{M(K_T)} \le b^{-9(k-10^{10}n^4+1)} \le b^{-8k}.$$

Both bounds are at most $\frac{1}{2}b^{-6k}$ under the given scales. Combining the upper estimate for $\mathcal{K}(Z)$ with (22.22) for K_T proves

$$\frac{\mathcal{K}(Z)}{M(Z)} \le b^{-6k}\frac{\mathcal{K}(K_T)}{M(K_T)}.$$

The phase factors over separate pairs, since the finite conductor character and unipotent membership factor, as does the integral (22.20). Haar normalization constants cancel in this ratio inequality (the zero form is cubic in the Haar normalization, while the denominator is linear). Choosing this intermediate subgroup at each of a set Σ of fixed-vector places and the full integral subgroup elsewhere gives the factor $b^{-6k|\Sigma|}$. Thus the numerical comparison in `IsFillingLevel.exists_intermediateGroup_lines` of `New/Filling/FixedLines.lean` follows from the bottom floor estimate, with all finite vector orbit counts explicit.

Lemma 22.19 (Central average controlled by the zero form)**.** *Let T be the union of the chosen split pairs and let ϕ denote the finite phase function there. Define*

$$\overline{\phi}(s) = |\mathfrak{G}|^{-1} \sum_{t \in \mathfrak{G}} \phi(tst^{-1}).$$

Under (22.21) for $Z = K_T$, $t = 0$, and $M(K_T) \geq 2^{-n}$, one has $\sum_s |\overline{\phi}(s)|^2 \leq 2$.

Proof. Restrict the nonnegative integral (22.20) to $x \in K_T$. Since the phase depends only on the residue class there, its inner integral is $M(K_T)\overline{\phi}(\operatorname{red} x)$. Each residue class has measure $M(K_T(g))$. Thus

$$M(K_T)^2 M(K_T(g)) \sum_s |\overline{\phi}(s)|^2 \leq \mathcal{K}(K_T).$$

Divide the upper bound (22.21) by the factor on the left. As $e_T \leq 1$ and $M(K_T) \geq 2^{-n}$ the resulting bound is at most two. □

This also gives a direct derivation of the trace estimate in `New/Filling/Moments.lean`. Let $\mathfrak{G}_i$ be the group at the i-th rational prime and $\jmath_i$ its insertion into the product group. We use the following exact residue support of the Bessel-subgroup amplitudes:

$$\nu^{-1} \sum_c \sum_j A_{S_B}(c, \jmath_i(s)^{-1}, c)_{jj} = D_c\phi(\jmath_i(s)). \tag{22.23}$$

Indeed a rational Bessel term $\gamma = uh$ with same initial and terminal class and reduction identity at all places of the *other* factor has h integral in the class lattice and congruent to identity at those places. Such a rational definite unitary h must be one. Here is an elementary verification of the freeness: if it is not one, the rational algebraic integer $a = \operatorname{tr}(h) - k$ is nonzero, because at any complex embedding a unitary matrix has trace of real part at most k, with equality only at the identity. Integrality follows by taking the trace of its integral conjugate. If it reduces to the identity at f places of norm q_{1-i}, its nonzero absolute norm is at least q_{1-i}^f; but its absolute value at every complex place is at most $2k$, so the absolute norm is at most $(2k)^{2f} < q_{1-i}^f$. The contradiction proves freeness. Hence the Bessel term is unipotent, and none exists if its reduction is not in $\bar{U}$. For a reduction in $\bar{U}$, strong approximation and the quotient covariance of Section 22.10 identify its amplitude with the unipotent translate of the identity amplitude. With the source convention $W(uy) = \chi(u)^{-1}W(y)$ for rational u, the term over the inverse residue gives phase $\phi(\jmath_i(s))$. Then the identity amplitude contributes ν on each diagonal coordinate. This proves (22.23); the sign of the phase would be inverted under the conjugate Bessel convention, with identical norm estimates.

If $z = \sum_s z_s[s]$ is a central self-adjoint idempotent of $\mathbb{C}[\mathfrak{G}_i]$, then the identity coefficient satisfies

$$z_1 = (z^* z)_1 = \sum_s |z_s|^2 \geq 0.$$

Its centrality implies that the sum $\sum_s z_s\phi(\jmath_i(s))$ does not change on replacing ϕ by its full conjugation average: conjugation by the other factor acts trivially and conjugation by its

own factor permutes the coefficients of z. Thus Lemma 22.19, the identity (22.23), and Cauchy–Schwarz give

$$\left|\nu^{-1}\sum_{s} z_s \sum_{c,j} A_{S_B}(c, \jmath_i(s)^{-1}, c)_{jj}\right| \le D_c\sqrt{2z_1}. \tag{22.24}$$

When $z_1 \le b^{-8}/2$ this is at most $b^{-4}D_c$, precisely `re_trace_besselS_le` in `Moments.lean`. Thus the required trace estimate is a consequence of the bottom finite local kernel estimate and the proved Bessel transport identities, not a separate use of a global trace formula.

22.16 The projection covariance and the complement-coset error

Let T_0 be the complement of the rational Bessel subgroup and let $A(s)$ denote the normalized block matrix with entries $\nu^{-1}A_{T_0}(c, s, c')$. Bessel transport under right multiplication by $\sigma \in \bar{S}_B$ permutes its column classes and replaces each column-coordinate block by a unitary factor, as proved in Section 22.10. In other words for a unitary block permutation V_σ one has $A(s\sigma) = A(s)V_\sigma$ (or the transpose convention, if the indices are reversed). It is essential here that V_σ depends on the lift of σ and the column class, but not on s; this follows from the term bijection by the same right multiplication for every s. The complement of the rational Bessel subgroup is stable under this multiplication. It follows for every central self-adjoint projection $z \in \mathbb{C}[\mathfrak{G}_i]$, with

$$M_z(s) = \sum_t z_t A(\jmath_i(t)^{-1}s),$$

that $M_z(s\sigma) = M_z(s)V_\sigma$ and hence $\|M_z(\sigma)\|_{\mathrm{HS}} = \|M_z(1)\|_{\mathrm{HS}}$.

For clarity the set on which to use the complement energy bound must be invariant under multiplication by $\jmath_i(\mathfrak{G}_i)$. The correct set is

$$\Omega_{1-i} = \{s : \ s_{1-i} \text{ is the } (1-i)\text{-component of some } \sigma \in \bar{S}_B\}.$$

It is the set involving the *other* factor, not the i-th factor. It contains $\bar{S}_B$ and is invariant under left $\jmath_i(\mathfrak{G}_i)$ multiplication. Convolution by z is an orthogonal projection in the regular Hilbert space, also with Hilbert–Schmidt matrix-valued coefficients, so restrict the function A by the indicator of Ω_{1-i} and apply its norm contraction. This proves

$$|\bar{S}_B|\|M_z(1)\|_{\mathrm{HS}}^2 \le \sum_{s\in\Omega_{1-i}} \|M_z(s)\|_{\mathrm{HS}}^2 \le \sum_{s\in\Omega_{1-i}} \|A(s)\|_{\mathrm{HS}}^2.$$

The complement energy estimate of `energy_compl_besselS_le` bounds the last sum by $b^{-12}|\bar{S}_B|D_c$. Thus the sum of the Euclidean row norms of $M_z(1)$ is at most

$$\sqrt{D_c}\|M_z(1)\|_{\mathrm{HS}} \le b^{-6}D_c \le b^{-4}D_c.$$

This supplies the remaining covariance-convolution step of `sum_norm_amplitudeSum_le` in `Moments.lean`. The use of the opposite factor is needed for the contraction argument and is independent of the fact that the corresponding energy estimates hold symmetrically for each factor.

We can also verify the reduction of the complement energy estimate to the crude stratified comparison. In the latter write

$$D = \frac{D_c^2}{d_\infty\,\mathfrak{m}_r}, \qquad D_1 = |\bar{S}_B|D_c, \qquad \tau = b^{kf/4}, \qquad \alpha = (b/g)^{kf}(2\tau^2).$$

The standing size hypotheses are $\tau^{44}D_1 \le D$ and $\tau^{18}\alpha D \le D_1$. The crude energy inequality `energy_le_of_linkEnergy_le` of `CrudeEnergy.lean`, recalled in Section 21.13, bounds the energy by $(1+b^{-4})^2 w + b^{-12}C$ when the retained links have energy at most w, and the parameters satisfy the stated thresholds: $C \ge D_1$; if a level-zero normal form occurs, $C \ge \tau^{20}D_1$; and if links are omitted, $C \ge \tau^{12}D/X^k$, where X is the product of residue-field sizes at which a nonzero frame vector is fixed.

On Ω_i, every place in both halves of every split pair above q_i fixes the first frame vector, since this is true for the reduction of the Bessel group. Thus we may take $X = q_i^{2f}$. The frame of a level-zero normal form is carried onto itself by an upper triangular matrix of constant diagonal ζ with $\zeta\bar{\zeta} = 1$: this follows by assembling the shared relations for each index into two upper triangular matrices, the first of diagonal ζ and the second of diagonal one. In the marked integral position the frame is unchanged by the class basis, so ζ is integral and invertible at every finite place. Its reduction at the specified factor agrees with the identity diagonal of $\bar{S}_B$; if $\zeta \ne 1$, its difference from one has norm at least q_i^f but absolute value at every complex place at most 2, a contradiction. Thus $\zeta = 1$; the term is in the Bessel subgroup. It follows that no level-zero normal form occurs for the complement on Ω_i.

Take all links omitted and $w = 0$, $C = D_1$. The missing-link size requirement holds because

$$\tau^{12}/X^k \le b^{3kf-20kf} = b^{-17kf}, \qquad \tau^{18}\alpha = 2b^{5kf}(b/g)^{kf} \ge 2b^{-17kf}.$$

The other requirements are immediate. Applying the crude comparison proves $\mathcal{E}(A_{T_0};\Omega_i) \le b^{-12}D_1$, as claimed in `energy_compl_besselS_le`. This derivation makes explicit its dependence on the inverse index savings and the level-zero freeness. No coherent main-term estimate is used for this error term.

22.17 Assembly of the sharp energy bound

We conclude by tracing the quantitative thresholds and the remaining inputs in the full second moment. Let Σ be the specified fixed-vector places above q_0 and set $y_\Sigma = b^{-6k|\Sigma|}$. The set `Filling.fixingSet` consists of reductions with the specified fixers and, unless all selected places above q_0 belong to Σ, with at least one selected place at which *all* leading frame minors are units. For Σ equal to the whole set, the latter condition is vacuous. The theorem of Section 20.8 then bounds the full positive-type energy using this intermediate set; that induction does not itself require unit minors at every place simultaneously.

The normalized coherent cut energy, with links retained when Σ is proper and no links retained when it is the whole set, must satisfy

$$w \le (1+b^{-4})y_\Sigma D. \tag{22.25}$$

The proof in the source uses the top packet comparison (`sum_blocks_le_topKernel` in `New/Filling/FiniteSystem.lean`), the global-to-local limiting identity (`exists_hasSum_co setAmplitude` in `GlobalToLocal.lean`), the archimedean and finite local band systems, the fixed-vector bottom factor, the global sieve just proved, and the packet constant inequality of the filling datum. We have supplied the complete global frequency and Poisson budget in Section 22.14, and the elementary norm comparison and multiplicative saving for fixed-vector groups from the finite bottom estimate. The top packet comparison must still be supplied as a specialized input, including the integral-point count on the definite quadric and its normalization. In particular one cannot use the crude bound on the number of rational elements in each link as a replacement for that comparison.

Given (22.25) and the crude comparison, the energy estimate is elementary. Take $C = y_\Sigma D$. Its level-zero requirement holds since $y_\Sigma \geq b^{-6kf} = \tau^{-24}$, so $y_\Sigma D \geq \tau^{20} D_1$ by the standing size hypothesis. If links are omitted, Σ is the whole set and $X \geq q_0^f$, so

$$\tau^{12} D / X^k \leq b^{-7kf} D \leq y_\Sigma D.$$

For proper Σ that requirement is vacuous. Thus the crude comparison with w as in (22.25) yields

$$\mathcal{E} \leq ((1 + b^{-4})^3 + b^{-12}) y_\Sigma D \leq (1 - b^{-4})(1 + 9b^{-4}) y_\Sigma D.$$

For the last inequality set $u = b^{-4}$: subtracting the left coefficient from the right gives $5u - 12u^2 - 2u^3 \geq 0$ for the stated b. This is the assembly of `energy_fixingSet_le` in `New/Filling/Moments.lean`. Its main analytical content remains precisely the local estimates and the top packet comparison giving (22.25); its numerical conclusion requires no further input.

The central projection calculation then gives the first moment of Section 20.9: use $z = 1$ in the covariance estimate and the identity Bessel amplitude to bound the rows of $P^*P - 1$. For each bad central-type projection use (22.24) and the covariance estimate; since its identity coefficient is at most $b^{-8}/2$, the total discarded trace is at most $2b^{-4} D_c$ for that factor. For the two commuting projections the total discarded trace is at most $4b^{-4} D_c$. The positive-type induction of Section 20.8 gives the second moment from the energy estimate just assembled. Thus the Hilbert–Schmidt variance and spread argument of Section 20.10 can be applied without additional finite-type covariance or central-trace inputs.

22.18 What remains of the filling argument

The new proofs have separated the following mathematical obligations from the global filling theorem.

- The normalized Flensted–Jensen identity, its change of splitting and triangular shifts, the actual large-gap determinant decay, the commutator-sensitive minor expansion, the quantitative Gaussian estimate, the common compact correction and the interpolation realization of the Bessel phase are now related by proofs with explicit constants. The classical representation-theoretic identification, formal degree and differentiated convolution theory are still invoked in their stated forms. The cut construction uses the peak and its quotient norm bounds in Section 21.6.
- The marked lattice and the exact class and group-order factors, unipotent coordinate Haar measure and covolume, arithmetic box construction, finite and infinite character normalization, transport between Bessel classes, and flag propagation eliminating the vanishing-discriminant terms are reconstructed here from integral block matrices, density, and elementary lattice geometry. The positive link metric and its congruence defect have also been checked.
- The lattice-height zeta sum and global frequency and head sums are reconstructed in their numerical strength, not only as convergence statements. Given the local systems, the one-sided sieve now follows with its error at most $b^{-6kf-10}/d_\infty$. The projection and complement-energy deductions from the crude comparison and bottom form have also been supplied.

Several genuinely specialized local estimates still remain. They include the construction of the floor kernels and the uniform Fourier bounds for the actual real Bessel system (`Ne`

`w/RealPlace/LocalTheorem.lean`, `BandLimits.lean`), the finite gauge/quotient measure comparisons and bottom form (`New/Filling/StandardLevel.lean`, `TopFloor.lean`, `FiniteSystem.lean`), the quantitative joint lattice-saturation and weighted pair bounds (`LinkPairing.lean`, `BlockBound.lean`, `CrudeEnergy.lean`), and the normalized coherent top packet comparison (`GlobalToLocal.lean`, `FiniteSystem.lean`). The class and degree size inequalities of `Filling.IsAdmissible` must be checked simultaneously for the actual weight and number field parameters using the mass formula and degree estimates. Some of these numerical identities follow directly from the class count and covolume derived here, but the full simultaneous verification is not asserted by these two identities alone. In particular (22.25) is still conditional on the named local and packet results. This distinction is necessary because a fixed-rank matrix coefficient estimate or a qualitative orbital integral convergence statement would not supply the constant-one second moment or its uniform error.

Chapter 23

Closing the global-to-local and finite-packet comparisons

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 25, 26, 27 and the final audit in Chapter 57.

This supplement concerns the coherent part of the second moment in Chapter 22. In this argument an ordinary upper bound for the number of rational elements is not a substitute for the mass computation: the leading coefficient must be one up to the stated error. We give an independent proof of the one-vector density formula and its one-sided use in the finite packet comparison. The global mass theorem, the local-to-global classification of Hermitian spaces, and elementary Fourier analysis suffice for the density formula; it is not necessary for this particular application to invoke the full higher-rank Siegel–Weil theorem. We also make the precise dependency on the convergence of the Bessel amplitudes explicit. The remaining local Fourier and simultaneous size estimates are recorded separately.

23.1 A normalized representation-density formula

Let E be the CM field in Chapter 4, $F = E^+$, $f = [F : \mathbb{Q}]$, and $V = E^k$ with the standard positive Hermitian norm

$$Q(x) = \sum_{i=1}^{k} \bar{x}_i x_i \in F.$$

We use $k \geq 3$. As in Chapter 6, E/F is unramified at all finite places, $D_E = D_F^2$ in absolute value, and the local Hermitian groups are the full unitary groups. Write $\widehat{O}_E$ for the product of the finite integer rings, and

$$v_E = \mathrm{vol}(\widehat{O}_E) = 2^f |D_E|^{-1/2}.$$

Thus the additive Haar measure on $V(\mathbb{A}_{F,f})$ gives the standard integer lattice volume v_E^k, with the archimedean measure ordinary complex Lebesgue measure. The two together give the diagonal rational subspace additive covolume one. On the self-adjoint finite adeles, identified with $\mathbb{A}_{F,f}$, distinguish two measures: db_0 gives the absolute codifferent $\mathfrak{d}_F^{-1}\widehat{O}_F$ mass one, while the self-dual measure for the rational additive character of F is $db_{\mathrm{sd}} = |D_F|^{1/2} db_0$. Indeed the self-dual measure gives $\widehat{O}_F$ mass $|D_F|^{-1/2}$, and the codifferent has index $|D_F|$ over it. This is also why the factor $|D_E|^{1/4}$ cannot be omitted from the formula below. Write ψ_F for the finite standard character. For self-adjoint b and r, the source's character $\psi_{E,f}(br/2)$ is precisely $\psi_F(br)$, by the relative trace formula.

Proposition 23.1 (Exact genus average for one vector)**.** *Let $K \subset U(V)(\mathbb{A}_{F,f})$ be an open compact subgroup of the integral stabilizer. For a locally constant compactly supported nonnegative function φ on $V(\mathbb{A}_{F,f})$ invariant under K and $\beta \in F$ totally positive, let h_c run over the class representatives for K, with automorphism orders a_c. Then*

$$\sum_c \frac{1}{a_c} \sum_{\substack{x \in E^k \\ Q(x)=\beta}} \varphi(h_c^{-1}x) = \left(\sum_c \frac{1}{a_c}\right) |D_E|^{1/4} \prod_{v|\infty} \frac{\pi^k \beta_v^{k-1}}{(k-1)!} \tag{23.1}$$

$$\int_{\mathbb{A}_{F,f}} \int_{V(\mathbb{A}_{F,f})} \varphi(x) \psi_F(b(Q(x)-\beta))\, dx\, db_0. \tag{23.2}$$

Both sides converge. In the left side each class sum of nonnegative terms is finite. The double integral is an iterated integral; the outer integral is absolutely convergent after performing the inner integral.

Proof. We use the classical local-to-global classification of Hermitian forms over quadratic extensions (Landherr) in the form proved or cited in Chapter 6. The form with diagonal $(\beta, \beta^{-1}, 1, \dots, 1)$ is isometric to the standard Hermitian form: it has the same signatures at the real places and the same determinant class everywhere, hence the requisite local invariants. Consequently $Q(x) = \beta$ has a rational solution x_0. Witt extension gives transitivity of $G = U(V)$ on its rational norm-β vectors and, locally, on the nonzero norm-β fibers. The stabilizer $H = U(x_0^{\perp})$ has rank $k-1$ and is again a full unitary group over the same extension. At almost all finite places x_0 is primitive of unit norm and unimodular integral Hermitian Witt extension gives transitivity on the integral fiber; thus the local orbit descriptions assemble into the adelic homogeneous space $H(\mathbb{A}_F)\backslash G(\mathbb{A}_F)$. The inverse action convention sends g to $g^{-1}x_0$.

For completeness the arithmetic normalization of unfolding is as follows. Use Tamagawa measures on G, H. The classical Tamagawa-number theorem gives $\tau(G) = \tau(H) = 2$ for full unitary groups over a quadratic extension, including these anisotropic forms. One explicit reference, checked against its statement, is M. Kirschmer, *Definite quadratic and hermitian forms*, Habilitationsschrift, 2016, Theorem 4.2.3 and its specification of the Tamagawa number (p. 51; the convention $[E : F] = 2$ has $\tau(G) = 2$), available at https://www.math.uni-bielefeld.de/~mkirschm/papers/herm.pdf. These are full unitary groups, not special unitary groups. The determinant maps identify their rational character tori; in the construction of the Tamagawa quotient measure the character-group convergence factors for G and H cancel. One can equivalently compute the quotient measure using rational invariant gauge forms. At x_0 choose their ratio to equal the Leray form dx/dQ, and propagate by the invariant action (unitary matrices have absolute determinant one on the underlying F-vector space). This gives the invariant Leray form on the norm fiber. The product formula shows that changing the rational bases or the initial ratio of rational gauge forms does not introduce an additional global scalar. With the ambient additive measures specified above, the resulting quotient measure is the product of the real Lebesgue Leray measure and the finite Leray measure using db_{sd} as target additive measure. This normalization follows also directly from the fact that the chosen global additive product measures on E and F give rational covolume one, and hence agree with the globally normalized rational gauge-form measures.

Unfold the nonnegative function

$$\sum_{x \in E^k,\, Q(x)=\beta} \varphi(g_f^{-1}x)$$

over $G(F)\backslash G(\mathbb{A}_F)$, taking the archimedean test function to be one on the compact spheres $Q(x_v) = \beta_v$. By transitivity this is the sum over $H(F)\backslash G(F)$ of the single test at x_0. Tonelli's

theorem turns its integral into an integral over $H(F)\backslash G(\mathbb{A}_F)$, then into $\tau(H)$ times the integral on $H(\mathbb{A}_F)\backslash G(\mathbb{A}_F)$. Dividing by $\tau(G)$ gives exactly the integral over the adelic norm fiber, with no factor two. On the other hand, the normalized adelic group integral of the original sum is its class average with weights $a_c^{-1}/\sum_{c'} a_{c'}^{-1}$: each finite double class contributes Haar volume $\mathrm{vol}(K)\,\mathrm{vol}(G_\infty)/a_c$, and the same factor occurs in the denominator. At each class there are only finitely many relevant rational vectors, since the finite support bounds denominators and the archimedean norm spheres are compact. This also verifies finiteness independently of the unfolding.

Local Fourier inversion computes the norm-fiber integral. At a real place, polar integration in $\mathbb{C}^k$ gives

$$\int_{\mathbb{C}^k} F(Q(x))\,dx = \frac{\pi^k}{(k-1)!}\int_0^\infty F(s)s^{k-1}\,ds,$$

as is seen by differentiating the ball volume $\pi^k s^k/k!$. Thus the real factor at β_v is $\pi^k\beta_v^{k-1}/(k-1)!$. At finite places the push-forward of $\varphi(x)dx$ by Q has continuous density at the nonzero point β, by local submersion and compact support. Explicitly, the F_v-derivative at x is $y \mapsto \mathrm{Tr}_{E_v/F_v}(\bar{x}\cdot y)$, nonzero on the fiber; locally solve for one F_v-coordinate. Partition the compact support above a small compact neighborhood of β into finitely many such charts. The analytic inverse function theorem and change of variables give the continuous density. This is valid also at the dyadic and split places, since the fields have characteristic zero and the trace pairing is nondegenerate.

Here is the absolute convergence needed for the finite Fourier inversion. On a local fractional lattice box $x = a + \pi^h y$, with y in the integer lattice for a uniformizer of F_v, the phase is a quadratic polynomial plus a linear term in y. Write the resulting normalized local integral as $J(b)$. Squaring its absolute value and replacing the second variable by its difference z from the first, the integral in the first variable vanishes unless $b\pi^{2h}$ times the trace pairing of z with the integer lattice is trivial. This uses additive-character orthogonality, and the linear term depends only on z in the difference. Hence $|J(b)|^2$ is at most the measure of that kernel in the norm lattice. At an unramified nonsplit place with residue cardinal q_v in F_v, when the conductor exponent of $b\pi^{2h}$ is $m > 0$ this kernel has measure q_v^{-2km}, since the trace pairing of the integer ring of the unramified quadratic extension with itself is perfect for the integer ring of F_v. In the split case there are two copies of the base integer ring, and the same result follows from the trace pairing between the two copies. Thus for large m, $|J(b)| \ll q_v^{-km}$. A locally constant compact test is a finite linear combination of such box indicators, so its Fourier transform is integrable. At almost all places the test is the standard lattice indicator, $h = 0$, and the contribution of the shell of conductor $m \geq 1$ to the Fourier L^1 norm is bounded by $(q_v^m - q_v^{m-1})q_v^{-km}$. The product of these bounds converges for $k \geq 3$, since $\sum_v q_v^{1-k} < \infty$. This gives absolute convergence over the full restricted product and justifies iterated inversion. The measure dual to db_0 locally gives the base integer ring mass one, so conversion to db_{sd} inserts $|D_F|^{1/2}$ globally. We obtain (23.2), using $D_E = D_F^2$. □

This proves the specialization `classSum_norm_eq` in `Lib/SiegelWeil/OneVector/OneVector.lean` for the field and ranks used in filling, with conventional measure justification. The direct orbit proof uses the classical Tamagawa theorem in place of the rank-one Siegel–Weil proof in the source. It does not use a mass result on a single lattice class or an assumption that every norm fiber initially contains a global point without checking its Hermitian invariants.

23.2 Uniform small error from the places outside the window

For a finite base place v write Q for the same standard norm on the local algebra $E \otimes_F F_v$, and use probability Haar measure on its integral vector lattice. Let $I_v(\beta)$ be its norm density at β when the target integral ring has mass one. The preceding local calculation gives the stronger convenient bound

$$0 \le I_v(\beta) \le 1 + \sum_{m \ge 1} (q_v^m - q_v^{m-1}) q_v^{-km} \le (1 - q_v^{-(k-1)})^{-1}. \tag{23.3}$$

Indeed db_0 gives the local annihilator of the integral base ring mass one; on that annihilator the absolute value of the normalized Gaussian integral is at most one, and its complement is the disjoint union of the conductor shells just estimated. This estimate is also valid when the density is zero, and does not assume a primitive representation of β in the integer lattice. In particular for any finite conjugation-stable set T, the product of the off-T densities is bounded by

$$\prod_{v \notin T/F} I_v(\beta) \le \zeta_F(k-1).$$

Here T/F is the set of base places underlying T. For $n = h + 1 = \mathfrak{s}.\varrho + 1$ in the source, its filling hypotheses include $n[E : \mathbb{Q}] \le 2^{2h}$, $1000(2h+2)^9 \le k$, and $2^{-h} \le b^{-6}$. For $s \ge 2$ one has

$$\log \zeta_F(s) \le f \log \zeta(s) \le f\Big(2^{-s} + \int_2^\infty t^{-s}\, dt\Big) \le 3f2^{-s}.$$

In particular $3f2^{-(k-1)} \le 2^{-h-3}$ by the two rank inequalities (indeed $f \le 2^{2h}$ and $k - 1 \ge 3h + 5$). Then $\exp(2^{-h-3}) \le 1 + 2^{-h-2}$, as follows by integration of $e^t \le 2$ for $0 \le t \le 1/2$. We have proved

$$\prod_{v \notin T/F} I_v(\beta) \le 1 + 2^{-h-2} \le 1 + b^{-6}/4. \tag{23.4}$$

This is a direct local-density alternative to the ideal-zeta majorant of `lintegral_gauss_sum_le_idealZeta` in `TorusMajorant.lean`. The source's more general bound uses a further factor for the measure of the local slab. In the present application we disintegrate the local slab separately, so the stronger (23.4) suffices. Importantly its bound is independent of T, β and the denominators introduced by the local test.

23.3 One-sided counting on the middle quadric

Write $N = \pi\bar{\pi}$ in the finite adeles and let $A = \prod_{w \in T} E_w$, with involution interchanging conjugate factors. Normalize the additive measure on its integral ring R to one, and the measure on the self-adjoint additive subspace to give its integral part mass one. Equivalently the latter measure is transported from the source's skew-adjoint measure using multiplication by its skew unit. In rational coordinates the middle form is $\mathrm{diag}(I_k, -N)$. Denote by B the change to the marked lattice basis with the distinguished vector replaced by $e_*/\bar{\pi} + z/N$; in particular $\det B = 1/\bar{\pi}$. For

$$e(\tau, x) = B^{-1}(x - \tau z/N)$$

one has

$$h(e, e) = Q(x) - \tau\bar{\tau}/N, \qquad h(z, e) = \tau. \tag{23.5}$$

These are just substitution in the diagonal rational form, expressed in the marked integral basis. The local norm fiber $h(e, e) = -N$ is regular. Its measure $d\delta$ in the source is the Leray, or disintegration, measure relative to the specified additive self-adjoint target measure.

Suppose $t \in E$ satisfies $|t_v| > N$ at every complex place; set $\beta = (t\bar{t} - N^2)/N$, totally positive. As in the source, choose a finite family of distinct pairs (c, x), $Q(x) = \beta$, with $h_c^{-1}x$ integral outside T. These are not arbitrary independent classes of vectors: each pair includes its column class c. Assume $F_0(e) \geq 0$ is a measurable function with the following lower-neighborhood condition: for each pair there is a neighborhood of $e(t, h_c^{-1}x)$ on whose $(H \cap K(g))$-orbits the function F_0 is at least a specified $c_{(c,x)} \geq 0$, where $H = U(X_0)$ acts as the compact middle subgroup. With $\mathcal{C}$ the set of the column classes, define the source constant

$$c_* := \left(\frac{\pi^k}{(k-1)!N^k} \right)^f |D_E|^{1/4} v_E^k, qquad G(t) := \prod_{v|\infty} (|t_v|^2 - N^2)^{k-1}.$$

Proposition 23.2 (The finite one-sided Siegel inequality)**.** *For all sufficiently small compact open additive neighborhoods L of zero in A,*

$$\sum_{(c,x)} c_{(c,x)} \operatorname{vol}(L) \leq (1 + b^{-6})|\mathcal{C}| c_* G(t) \int_{h(z,e)-t \in L} F_0(e)\, d\delta(e). \tag{23.6}$$

Proof. First replace F_0 from below on the relevant neighborhoods by an $(H \cap K(g))$-invariant locally constant compactly supported nonnegative step function Φ, preserving the required values. To see this, choose an integral scale sufficiently small that the basic neighborhoods are cosets of the same smaller compact open vector lattice M. The compact group $H \cap K(g)$ preserves each scalar multiple of the standard lattice. Take the union of its translates of each prescribed coset, and take the pointwise maximum of the corresponding finitely many $c_{(c,x)}$ times their indicators. These unions are compact open and translation invariant by M, since they lie in a compact set and are unions of open M-cosets. By the lower-neighborhood hypothesis this function is bounded by F_0. This is the step-function lemma `Filling.exists_step_function` of `FiniteSystem.lean`; no measure approximation of a function on a null fiber is being used.

Apply Proposition 23.1 to the finite test that equals $\Phi(e(t, h_c^{-1}x))$ after evaluation at T, and is the indicator of integrality of the vector coordinates outside T. In the argument of the test, h_c is only the class representative: the single common test on adelic vectors y is $\Phi(e(t, y_T))$ times the off-T integer-lattice indicator. It is locally constant and compactly supported (the map from x to $e(t, x)$ is an affine proper embedding). It is invariant under the column level, since the latter acts integrally outside T and as the indicated middle subgroup at T. The column stabilizers are trivial by the free-level argument. The left side of (23.2) therefore dominates the sum of the prescribed weights, and its mass factor is $|\mathcal{C}|$.

Let $D_T(\tau)$ denote the norm density of the local vector test $\Phi(e(\tau, x))$ at $Q(x) = \tau\bar{\tau}/N - N$, for the normalized integral measures on A^k and the local self-adjoint target. At $\tau = t$ this is exactly the normalized T-factor of the density in (23.2). The complete finite factor is

$$v_E^k D_T(t) \prod_{v \notin T/F} I_v(\beta).$$

To relate $D_T(t)$ to the actual quadric integral, the map $(x, \tau) \mapsto e(\tau, x)$ has determinant of modulus

$$|\det B^{-1}|_A\, |N|_A^{-1} = |\pi|_A / |\pi\bar{\pi}|_A = |\pi|_A^{-1}.$$

In the last equality we use the conjugation stability of T and $|\bar{\pi}|_A = |\pi|_A$. The derivative in the norm variable in (23.5) is one with this choice of Q. Disintegrating the ambient middle-space vector measure in that variable gives the exact formula

$$\int_{h(z,e)-t \in L} \Phi(e)\, d\delta(e) = |\pi|_A^{-1} \int_{t+L} D_T(\tau)\, d\tau. \tag{23.7}$$

The density $D_T(\tau)$ is continuous at t, by the local submersion argument in Proposition 23.1; $\tau\bar{\tau}/N - N$ is a nonzero unit of the local product of fields for sufficiently small L, since its rational value at t is nonzero. Uniformity follows by shrinking the neighborhood in a finite collection of charts covering the compact test support. In fact the normalized integer-ring measures and a locally constant test give a locally constant density at such regular values. Continuity alone is sufficient: if $D_T(t) > 0$, for small L one has $D_T(t) \le (1 + b^{-6}/4)D_T(\tau)$ throughout $t + L$. If $D_T(t) = 0$ the required inequality is immediate from the genus average. Equation (23.7) consequently gives

$$D_T(t)\,\mathrm{vol}(L) \le (1 + b^{-6}/4)|\pi|_A \int_{h(z,e)-t\in L} \Phi(e)\, d\delta(e).$$

This proves in particular the near-quadric slab comparison in `StandardLevel.lean` for the tests needed here.

Outside T we may assume N and its inverse are integral, and likewise $\pi, \bar{\pi}$; this holds for all sufficiently large T. The product formula and $\pi\bar{\pi} = N$ give $|\pi|_A = N^{-f}$. Insert this equality, the off-T bound (23.4), and $\prod_v \beta_v^{k-1} = N^{-f(k-1)}G(t)$ into (23.2). The product of the two errors $1 + b^{-6}/4$ is at most $1 + b^{-6}$ for $b \ge 1$. The coefficient is exactly $|\mathcal{C}|c_*G(t)$, since the power N^{-f} from the Jacobian supplies the last factor needed to make N^{-fk}. Finally $\Phi \le F_0$ on the whole middle vector space. This proves (23.6). □

This is the needed version of `Filling.IsFillingLevel.siegel_sum_le_lintegral_quadric`, with its constant $1 + b^{-6}$. The argument does not suppose that every test has a positive lower bound on the whole quadric; only the finite collection of neighborhood lower bounds is used. The exact archimedean mass and the finite determinant factor are important for the subsequent packet count.

23.4 Term positions, finite characters, and the exact cutoff identity

Let $C = K(g)$ in the local product over T, and $C \subset Z \subset K$ an intermediate compact open subgroup. We use the standard level and the character in Chapter 22. The finite phase $\chi_g(y)$ is zero off UC, and on uC is the value of the additive Bessel character of u. The character is trivial on $U \cap C$ by its conductor, so the definition is independent of u. The vector $W_Z(y)$ indexed by the finite set of left cosets Ca in Z has coordinate $\chi_g(ya^{-1})$; the normality of C in K makes this independent of the representative and right C-invariant. Put $\mathcal{D} = \mathcal{C}$ for the column-class set. Write h_c for the chosen finite adelic point for class c, and

$$L(c, d, \gamma) = h_c^{-1}\gamma_f h_d.$$

Here and below the embeddings of the smaller unitary group in the larger group are implicit. This is the source's `termPosition` in the marked lattice coordinates. A term over (c, s, d) means a rational γ for which $L(c, d, \gamma) \in K$ has reduction s. Use the measurable nonnegative cutoffs β_c of Sections 22.9 and 22.10, having periodization one over $U_c(g)$ (the rational unipotents whose conjugate by h_c belongs to the principal level). Let $J_\infty(\gamma)$ be the Archimedean correlation $\int W(y) \otimes \overline{W(\gamma_\infty^{-1}y)}\, dy$. The individual amplitude of γ is the same integral with weight $\beta_c(y)\beta_d(\gamma_\infty^{-1}y)$, and is denoted by $A(c, d, \gamma)$. These definitions apply to the cut or full side, with their respective integrability properties.

We make explicit the summability premise used in this subsection and in the subsequent packet limit: the sum of the integrals of the norms of these weighted amplitudes over all

terms of fixed classes and reduction is finite. This is `amplitudeAbs_lt_top_and_cut_le` and `summable_amplitude` in `AmplitudeSums.lean`. The present proof establishes the normalized global-to-local unfolding from this premise, not the uniform majorant for the amplitude tail itself.

At a finite set of conjugation-stable places T containing all denominators of the scalar moduli and relevant fixed matrices, put

$$A_T(c,s,d;\gamma)=\sum_{u,v\in U(F)}\mathbf{1}_{L(c,c,u)\text{ integral off }T}\,\mathbf{1}_{u\gamma v^{-1}\text{ term over }(c,s,d)}A(c,d,u\gamma v^{-1}). \tag{23.8}$$

This is `Filling.cosetAmplitude`. We will use it when the unipotent pair in the double orbit is unique; thus its sum of integrated norms is bounded by that of the relevant terms.

For a head p of length h denote its Weyl element by w_p and let $\kappa(p)$ be the source's headDensity$(h,p,1)$. For σ fixing all hyperbolic frame vectors, put $\gamma=w_p\sigma$, let $\ell=L(c,d,\sigma)_T^{-1}$, and form the finite correlation tensor

$$\mathcal{J}(p,\ell)=\kappa(p)\int_G W_Z(x)\otimes\overline{W_Z(\ell w_p^{-1}x)}\,d\mu(x). \tag{23.9}$$

Here μ is the local group measure of the standard level. The finite representatives of p in this equation are the images of its rational entries. The identity $L(c,d,w_p\sigma)_T=w_pL(c,d,\sigma)_T$ follows because the column groups centralize the Weyl elements and the latter are expressed in the same fixed hyperbolic marked basis.

Proposition 23.3 (Correlation equals the cut-off rational orbit sum)**.** *Assume the absolute summability stated above, and $L(c,d,\gamma)$ integral off T. For $k\in K$ with reduction s, assume $a,b\in Z$ satisfy $k_T^{-1}ab^{-1}\in C$. Then*

$$\mathcal{J}(p,\ell)_{a,b}\,J_\infty(\gamma)=\kappa(p)\mu(C)A_T(c,s,d;\gamma). \tag{23.10}$$

Here the scalar finite coordinate multiplies the Archimedean tensor.

Proof. Write x_T for the variable of integration after translating it by a on the right, so that the first phase is $\chi_g(x_T)$ and the second is, by C-invariance and the stated relation between a,b,

$$\chi_g(\xi_T),\qquad \xi=L(c,d,\gamma)^{-1}xk.$$

An adelic lift of the local x_T can always be chosen integral off T (choose identity there), and then both x,ξ are integral off T. Define

$$B_c(x,u;y)=\mathbf{1}_{L(c,c,u)x\in K(g)}\beta_c(u_\infty y)$$

and similarly $B_d(\xi,v;\gamma_\infty^{-1}y)$. If $\chi_g(x_T)\neq 0$, some local element of U carries x_T into C, and there is such a rational u with adelic principal-level condition: density of $U(F)$ in the finite adeles of U follows from triangular affine coordinates and the density of E in $\mathbb{A}_{E,f}$, imposing the open conditions at T and integrality outside T. This elementary approximation was used in Section 22.9. All choices of u then form a single left coset of $U_c(g)$. Summing the cutoff over this coset gives one by its periodization property. If $\chi_g(x_T)=0$, no such rational u exists. The identical statement holds for d,ξ.

When these two principal-level conditions hold, the finite additive phase $\chi_g(x_T)$ equals the inverse of the global finite unipotent character of u, and likewise for v. The equality uses the integral conductor away from T and the definition of the phase. By the product formula for the additive character, this inverse is precisely the Archimedean factor in the

transformation law of $W(u_\infty y)$. Consequently the product of the two finite phases with the tensor $W(y) \otimes \overline{W(\gamma_\infty^{-1} y)}$ is

$$W(u_\infty y) \otimes \overline{W((u\gamma v^{-1})_\infty^{-1} u_\infty y)}.$$

Insert the two periodizations just obtained, and interchange their sums with both integrals. This is justified by the absolute-integrated-norm premise: for each admissible pair, the integration over its first principal-level condition has measure $\mu(C)$, and after translating y by u_∞ the remaining integral is exactly the integrated norm of its amplitude.

For clarity the two principal-level conditions are equivalent to the following conditions: $L(c, c, u)$ is integral off T, $L(c, c, u)_T x_T \in C$, and $u\gamma v^{-1}$ is a term over (c, s, d). Indeed the definition of ξ gives

$$L(c, c, u) x\, k\, (L(d, d, v)\xi)^{-1} = L(c, d, u\gamma v^{-1}).$$

If the two outer factors are in $K(g)$, this has reduction s. Conversely if the first factor is in $K(g)$ and the right-hand side has reduction s, the second outer factor is in $K(g)$. Off-T integrality of the first is equivalent to that of $L(c, c, u)$ because x was chosen integral invertible there. Thus for each retained pair in (23.8), the local x_T ranges over a single coset of C and contributes exactly $\mu(C)$; excluded pairs contribute zero. Translate the infinite variable by u_∞ to obtain the amplitude over $(c, d, u\gamma v^{-1})$. Multiplying by the front head density in (23.9) proves the asserted identity. □

This furnishes a direct proof of `IsFillingLevel.correlation_smul_eq_cosetAmplitude` of `GlobalToLocal.lean` with the summability premise explicitly retained. The factor $\mu(C)$ is forced by the integration over a principal congruence coset; changing the finite group measure or ignoring this factor would also change the leading term.

23.5 A local packet lemma

We give the finite local calculation used when comparing rational packets with the limiting middle quadric. In this subsection let M denote the unitary group on the middle summand $X_0 \oplus Az$, and H the stabilizer of z. The group M is the source's top floor; $H = U(X_0)$ since the norm of z is nonzero. The local group measure on H is denoted by μ_H (with integral compact subgroup of measure one). Write $C_H = C \cap H$. We assume the support estimate of the local cut kernel, namely a compact bound for the common support in x of $W_Z(x)$ and $W_Z(\ell w_p^{-1} x)$, uniform in $\ell \in M$ and in heads near a fixed head. This is the top-floor consequence of `Gauge.hasMatSize_of_ne_zero` in `LocalTheorem.lean`. It is required only for one local compactness argument below, not as a bound on the size of the packet integral.

Fix finitely many heads ω_p with their last entry equal to the same unit t. For R a positive semidefinite complex matrix indexed by the heads, put

$$Z_R(\ell) = \sum_{a,b \in C \setminus Z} \sum_{p,q} \overline{\mathcal{J}(\omega_p, \ell)_{a,b}}\, R_{pq}\, \mathcal{J}(\omega_q, \ell)_{a,b}.$$

The quantity is a nonnegative real number, by positive semidefiniteness of R. The energy at a point e of the middle quadric of norm $-N$ used in the source is

$$E_R(e) = \inf_{\ell \in M:\, \ell^{-1} z = e} \int_H Z_R(h\ell)\, d\mu_H(h).$$

The set over which the infimum is taken is nonempty by Witt extension locally; all its elements form a left H-coset, so Haar invariance also shows that the integral itself is independent

of the chosen lift when finite. Defining it by an infimum makes the lower-bound argument possible without this last observation. The integral is measurable by Tonelli on each local continuous section of the orbit map constructed below. A second-countable local quadric admits a countable cover of such sections, so E_R is measurable on the whole quadric, with the nonnegative integral allowed to be infinite.

First, the correlation is invariant under sufficiently small right perturbations of its second argument, uniformly in ℓ. Indeed in (23.9) the second argument changes from $\ell w_p^{-1}x$ to $\ell u w_{p'}^{-1}x$ when (p,ℓ) changes to $(p',\ell u)$, so the two differ by right multiplication by

$$x^{-1}w_p u w_{p'}^{-1}x.$$

For x in the common support, the compact support bound and openness of C ensure that this lies in C whenever (p',u) is sufficiently near $(p,1)$. The same bound applied to either nonzero integrand makes this conclusion valid on the union of the two supports, so no change of a vanishing contribution into a nonzero one is overlooked. By local constancy of $\kappa(p)$ (moduli of nearby ideles agree), the correlation tensors are equal. Furthermore $\mathcal{J}(p,k\ell)=\mathcal{J}(p,\ell)$ and $\mathcal{J}(p,\ell k)=\mathcal{J}(p,\ell)$ for $k\in C_H$. The first equality uses left C_H-invariance of the finite vector: the elements have trivial unipotent character and reduce to the identity; the second follows in the integral by commuting k with w_p and substituting $x\mapsto kx$. These are the invariance statements needed below.

Lemma 23.4 (Disjoint packet cosets give a neighborhood lower bound)**.** *Suppose a nonempty finite family (ℓ_i,n_i) with $\ell_i\in M$, $n_i\in(C\backslash Z)^2$, has $\ell_i^{-1}z=e_0$ for every i, and the pairs $(C\ell_i,n_i)$ are distinct. For all x sufficiently close to e_0, and all y which lie in its C_H-orbit, one has*

$$E_R(y)\geq \mu_H(C_H)\sum_i\sum_{p,q}\overline{\mathcal{J}(\omega_p,\ell_i)_{n_i}},R_{pq}\mathcal{J}(\omega_q,\ell_i)_{n_i}.$$

Proof. The local orbit map $a\mapsto a^{-1}z$ admits a continuous local section near e_0 taking a chosen value ℓ_0 above e_0. In characteristic zero this follows by the inverse function theorem: the derivative at a point with nonzero Hermitian norm maps the Lie algebra of skew-adjoint endomorphisms surjectively to the tangent space $\{v:\mathrm{Tr}_{A/A^+}h(e_0,v)=0\}$ of the smooth norm fiber. To see the surjection, for perpendicular v extend the rank-two map on the lines of e_0,v by a skew-adjoint endomorphism using division by $h(e_0,e_0)$; for the skew scalar component along e_0 use a skew scalar on that line. This holds over each factor of the finite product, including in residue characteristic two, since we work in the fields. Hence for all nearby x on the quadric we can choose $s_x\to 1$ in M with $(\ell_i s_x)^{-1}z=x$ for every i. The preceding compactness argument gives $\mathcal{J}(\omega_p,\ell_i s_x)=\mathcal{J}(\omega_p,\ell_i)$ simultaneously for the finitely many heads.

For any lift ℓ of x write $h_i=\ell_i s_x\ell^{-1}\in H$. Left C_H-cosets of the h_i for the same coordinate n_i are distinct: equality would give $\ell_i\ell_j^{-1}\in C_H\subset C$, contrary to distinctness. On each corresponding coset the relevant coordinate of every correlation is constant. Since the remaining summands in Z_R are nonnegative, their integral is at least the sum over these distinct labeled cosets, each of measure $\mu_H(C_H)$. The bound holds for every lift and hence for the infimum. If $x=ky$ with $k\in C_H$, use $\ell_i s_x k$ as lifts of y; right C_H-invariance of the correlations gives precisely the same bound. □

This proves `eventually_sum_packets_le_energy` in `TopFloor.lean`, and explains its uniform local-constancy premise in `eventually_correlation_eq`. No positive pointwise lower bound for arbitrary unipotent translates is being asserted; the lower bound is on disjoint labeled cosets of the middle stabilizer.

For a compact open additive subgroup L of A the top kernel is

$$K_L(p,q) = \int_{h(z,\ell^{-1}z)-t\in L} \langle \mathcal{J}(\omega_p,\ell), \mathcal{J}(\omega_q,\ell)\rangle \, d\lambda(\ell),$$

with the top stack measure λ of the level and with the inner product conjugate-linear in the first entry. This is `Level.topKernel`. The stack measure is the convolution of the H-Haar measure with the top-floor measure, which pushes forward to the actual quadric measure $d\delta(e)$. The push-forward identity is the measure-preserving orbit-point relation in `LevelGauge.lean`; at the top it is the standard homogeneous-space disintegration with the norm-fiber measure of (23.7). These are the specified top-stack and floor normalization hypotheses here. Under them, expanding the finite sum and applying Tonelli (all total quadratic forms are nonnegative) gives

$$0 \le \int_{h(z,e)-t\in L} E_R(e)\, d\delta(e) \le \Re \sum_{p,q} R_{pq} K_L(p,q). \tag{23.11}$$

Indeed for each fixed lift of a quadric point the right integrand after expansion is $\int_H Z_R(h\ell) d\mu_H(h)$, which dominates the defining infimum. The matrix sum on the right is nonnegative real by the same expansion before integration. This proves the measure-theoretic step `lintegral_energy_le_topKernel`; absolute convergence of the correlations on the whole stack is still needed to interpret each separate entry of K_L as an ordinary complex integral. This is the integrability clause of `floorForm_top_eq_integral`, which is not being inferred from pointwise local constancy.

23.6 The normalized coherent packet bound

We now combine the preceding results. Let i range through a finite family of distinct tuples $(c_i, d_i, s_i, \sigma_i)$ where s_i has an integral lift k_i whose image at T lies in Z, each σ_i fixes the hyperbolic frame, and $h(z, \sigma_i z) = t$. Assume the fixed term positions for σ_i and $w_p\sigma_i$ are integral off T. Let p run over a finite set of rational heads with their common last component t. Write $A_i(p) = J_\infty(w_p\sigma_i)$ for the infinite correlation. Assume, as in the source's coherent-packet grouping, that the matrix

$$R_{pq} = \langle \rho_p A_i(p), \rho_q A_i(q)\rangle$$

for fixed real nonnegative scalars ρ_p is independent of i. This is a true Gram matrix, automatically positive semidefinite. Suppose the head normalization is

$$\rho_p \kappa(p) = r\sqrt{G(t)} \qquad \text{for all } p.$$

This is where the rational product formula for head moduli enters in the source; no further estimate in the packet argument is needed for this relation. Define

$$Y_i = r^2 \left\| \sum_p A_T(c_i, s_i, d_i; w_p\sigma_i) \right\|^2 .$$

Let $v = 1/|\bar{U}|$ be the mass of the integral unipotent principal congruence subgroup for the source's unipotent measure (the product of the floor moduli), so $|\bar{U}| = g^{fh(2k+2h+1)}$. Write

$$m_Z = \frac{\mu(Z)\mu(C)}{v^2}, \qquad D_0 = |\bar{S}_B|\,|\bar{U}|, \qquad a_* = D_0|\mathcal{C}|c_*.$$

These are precisely `IntermediateGroup.massRatio` and `Filling.packetConst`. The class and group calculations of Sections 22.7 and 22.8 give

$$m_Z = \mu_H(C_H)\mu(C)^2\,[Z : C]D_0. \tag{23.12}$$

For verification, $\mu(Z) = [Z : C]\mu(C)$, $\mu_H(C_H) = 1/|\bar{H}|$, $|\bar{S}_B| = |\bar{U}||\bar{H}|$, and $v = |\bar{U}|^{-1}$. The product on the right is thus $\mu(C)^2[Z : C]|\bar{U}|^2$.

Proposition 23.5 (Finite coherent packet comparison)**.** *Under the preceding conditions, the absolute summability in Proposition 23.3, and the local support and top-stack normalization and integrability in Section 23.5, for all sufficiently small compact open additive subgroups L one has*

$$\sum_i Y_i \le (1 + b^{-6})a_* \Re \sum_{p,q} R_{pq} \frac{K_L(p,q)}{\mathrm{vol}(L) m_Z}. \tag{23.13}$$

Proof. Put $\ell_i = L(c_i, d_i, \sigma_i)_T^{-1}$. Its orbit point has the form $e(t, h_{c_i}^{-1} x_i)$ of (23.5), where x_i is the X_0-component of $\sigma_i z$. In particular $NQ(x_i) = t\bar{t} - N^2$, and $h_{c_i}^{-1} x_i$ is integral off T by the term-position hypothesis and integrality of the marked basis there. Group the indices by distinct pairs (c_i, x_i). For each label $b \in C\backslash Z$ choose $a_i(b) \in C\backslash Z$ satisfying $k_i^{-1} a_i(b) b^{-1} \in C$; it exists because k_i is in Z and C is normal there.

Within the fiber of a fixed pair (c, x), the pairs $(C\ell_i, (a_i(b), b))$ are pairwise distinct. Indeed equality forces equality of b and of a_i, hence equality of the reduction of k_i and k_j. Equality of orbit points rationally gives $\sigma_i^{-1}\sigma_j \in H(F)$ (it fixes all hyperbolic vectors and the middle vector z). Equality of the left C-cosets makes $h_{d_i}^{-1}\sigma_i^{-1}\sigma_j h_{d_j}$ principal-congruent at T; it is integral off T by the assumptions, and $1/g$ is integral off T. Thus it belongs to the global principal subgroup, whose preimage in the column group is its column level. It follows that the column classes d_i, d_j coincide; for the same chosen representative, the free-level stabilizer is trivial, so $\sigma_i = \sigma_j$. Now equality of the tuples implies $i = j$. This is why the packet must be grouped by class and rational vector, not simply by the finite residue of its orbit point.

Apply Lemma 23.4 to this fiber, with the distinct labeled cosets just established. The weight of each (i, b) computed from the Gram matrix is

$$\begin{aligned}\sum_{p,q} \overline{\mathcal{J}(\omega_p, \ell_i)_{a_i(b),b}}\, R_{pq} \mathcal{J}(\omega_q, \ell_i)_{a_i(b),b} &= \Big\| \sum_p \rho_p A_i(p) \mathcal{J}(\omega_p, \ell_i)_{a_i(b),b} \Big\|^2 \\ &= \mu(C)^2 G(t) Y_i.\end{aligned}$$

The last equality is exactly Proposition 23.3 and the head-density relation; it does not use cancellation by a triangle inequality. Summing over b and multiplying by $\mu_H(C_H)$ yields, using (23.12), the neighborhood lower bound for the energy of the fiber

$$\frac{G(t) m_Z}{D_0} \sum_{i \text{ in fiber}} Y_i.$$

Apply Proposition 23.2 with the nonnegative function $F_0 = E_R$, these finitely many distinct pairs (c, x) and precisely these weights. Its right side is bounded by (23.11). Since $G(t) > 0$, cancel it from both sides, then divide by $m_Z \,\mathrm{vol}(L) > 0$. The result is (23.13). All constants in the inequality have thus come from the genus average, the finite Haar-coset and stabilizer factors, and the head-density identity. □

This proves the normalized count-plus-packet inference of `sum_blocks_le_topKernel` in `FiniteSystem.lean`, conditional on the explicitly listed local support and integrability (and absolute global amplitude summability) premises. In contrast to a direct numerical comparison between two arbitrary kernels, its proof explains why the same Gram matrix, the pairwise coset distinctness, and the coefficient $1 + b^{-6}$ are exactly the conditions needed for the sharp coherent estimate.

23.7 The limit over the finite adelic windows

The remaining limiting step in the comparison is elementary once absolute convergence and the big-cell decomposition have been supplied. We spell it out to make clear that no limiting comparison of the constants $\mathrm{vol}(L)$ or m_Z is used.

The big-cell parameters for rational elements of a link with all leading minors nonzero are an ordered head $(p_0, \dots, p_{h-1})$, a common middle isometry $\sigma \in M(F)$, and two unipotents $u, v \in U(F)$:

$$\gamma = u\, w_p \sigma\, v^{-1}.$$

Here the last entry is fixed by the link, $t = h(z, \sigma z)$. We use the existence and uniqueness of these rational coordinates (`exists_big_cell_decomp`, `big_cell_unique` and `link_big_cell_eq_iff`); they are the actual Bruhat/Gauss decomposition of the two hyperbolic flags in the big cell. We retain this named matrix-coordinate assertion unless its complete marked-frame version has been reconstructed separately. The norm identity (23.5) gives $|t_v| > N$ when the last nondegeneracy determinant is nonzero: if the rational X_0-component is nonzero, it is nonzero at every embedding, and its positive-definite norm is strictly positive at every Archimedean place.

For one nonzero link amplitude over (c, s, d), choose the representative σ from any surviving term. Terms of this link over the classes and reduction are then exactly the elements with these big-cell coordinates that satisfy the finite integrality and residue conditions. By the uniqueness of the coordinates, the indexed sum of their integrated norms is dominated by the assumed sum of all the integrated norms over the same classes and reduction. Therefore the series may be collected first by the head p and then by the two unipotents. Denote its sum in each fixed head by $A(p)$. Since the family is absolutely summable, $\sum_p A(p)$ is the entire amplitude of the link.

For fixed p, the partial sum with the off-T restriction is precisely (23.8). For a finite adele and its inverse, every component is integral outside some finite place set (this is the defining restricted-product condition). Thus for any one pair (u, v) the indicator of integrality of $L(c, c, u)$ off T is eventually one as T increases. For each fixed p these partial sums tend to $A(p)$ in the finite-dimensional Archimedean tensor space: for any $\epsilon > 0$ first omit a tail of the absolutely convergent series with sum of norms less than ϵ, and then enlarge T so that all terms in the finite retained part satisfy the additional restriction. This proves exactly the dominated-convergence assertion of `exists_hasSum_cosetAmplitude` in `GlobalToLocal.lean` given the coordinate decomposition and summability. For a finite collection of packets and heads, take a single place set containing all the exceptional places for their finitely many matrices and scalar entries, their inverses, and the conjugate places. In particular $1/g$, $1/N$ and the change of marked basis are integral off T. The normalized head relation then follows from the product formula: for a rational head entry u with both u, u^{-1} integral off T, the square root of its finite modulus equals the inverse product of its Archimedean absolute values.

In applying Proposition 23.5, one now takes a finite head collection and a sufficiently large such T. The local packet estimate holds for every sufficiently small L for that fixed collection. The local finite sieve, if available with its stated normalization, applies to $K_L/(\mathrm{vol}\, L\, m_Z)$ and has constants independent of T and L; the global sieve of Section 22.14 is also uniform. Consequently the same upper bound holds for each finite rational sum as T increases. Passing to its limit at fixed heads, then exhausting the countable head sets and the class packets by finite subsets, is legitimate by the absolute summability assumed above and the nonnegativity of the total energy. At no point do we need to assume that $\mathrm{vol}(L)$ or m_Z separately converges along a shrinking or growing place set. This is the exact order of limits required in the application to (22.25).

23.8 The slice calculation underlying floor Plancherel

Here we give the analytic part of the local tower estimate, with its precise exponent. Write $n = h + 1$ as in the Lean development. In the notation of `LevelGauge.lean`, a gauge is a family of size sets B_j with $0, 1 \in B_j$, closed under negation and involution, with $B_i B_j \subseteq B_{i+j}$ and the sum of at most $r + 1$ entries from B_i in B_{i+1}. The Gram matrix and its inverse have entries in B_1. The size of the compact box is 1. The scalar measure satisfies $\mathrm{vol}(B_j) \le \Lambda^{j+1}$. One of the defining gauge conditions, `Gauge.fib`, is

$$d_0\{v : \text{all coordinates in } B_i\} \le \Lambda^{99n(i+1)} \tag{23.14}$$

for the zero-norm quadric on d hyperbolic pairs, for $m \le d \le m + h$, where $m \le 9n$ is the number of middle hyperbolic pairs. We prove below the complex-place version of this condition.

The source's chart uses a nonzero primary hyperbolic coordinate t, complementary coordinate s, other hyperbolic coordinates p, q, and an orthonormal part u. A point of the norm-β quadric, in a fixed normalization of the trace-one and skew scalars, is

$$s = \bar{t}^{-1}\tau(\beta - Q(p, q) - Q(u)) + ta.$$

Here a is a skew variable, $Q(p, q)$ is the norm of the other hyperbolic coordinates, and τ is the trace-one scalar. Equivalently, after the fixed skew-unit rescaling of the source, the same formula has β replaced by that rescaled target. The density is the product measure on (t, p, q, u, a); the dilation of a by t cancels the coarea factor for solving the norm equation in s. The negligible vanishing-t chart complement may be omitted. In a split place t is a unit almost everywhere, because the coordinate components are Haar distributed. This is the ordinary flat/Leray chart of `Place.flatChart` and it also describes the norm-zero cone by omitting u and taking $\beta = 0$.

For $P = (t, p)$ fixed, let $z = (q, a)$ denote the fiber variable. The key definition `Gauge.HasFibreBound` for a measure $d\eta$ on the full floor space is that, in the chart above (or the direct coordinate chart for volume),

$$d\eta = dP\, du\, d\zeta(z), \qquad \zeta\{z : \text{chart point of size } j + 4\} \le \mu(P), \qquad \int \mu(P)\, dP \le M.$$

The extra 4 allows us to pass from the original coordinates to the floor basis: the latter has a size-1 Gram matrix and adjoint, so multiplication by the Gram and conjugate transpose expresses the inverse coordinates and costs at most four. The bound is uniform in u.

Lemma 23.6 (Cone-to-fiber domination)**.** *Under* (23.14) *the floor quadric, and also the floor volume measure, satisfy* `HasFibreBound` *at size* j *with* $M = \Lambda^{99n(j+6)}$.

Proof. For the quadric, fix P, u and, if the bounded fiber is nonempty, choose any reference point (q_0, a_0) in it. Replace (q, a) by $(q - q_0, a - a_0)$ and s by $s - s_0$. The linearity of the hyperbolic expression in q with p fixed implies that these difference coordinates give a zero-norm hyperbolic-cone point with the original t, p and no u. The terms β and $Q(u)$ have canceled. Size grows by at most one under subtraction: the new coordinates lie in B_{j+5}. Translation in the fiber preserves $d\zeta$. Thus define $\mu(P)$ to be the measure of the cone fiber of size $j+5$; it dominates our original fiber for every u, including the case of an empty fiber. This function is measurable by Tonelli and its integral is the cone mass, bounded by (23.14) with $i = j + 5$. No measurable choice of the reference point is required. For ordinary volume one simply uses the product of the indicator of the bounded P-box and the volume of the bounded z-box; the number of its combined hyperbolic coordinates is at most $2(m + h) \le 20n$. The scalar volume estimate gives a bound at most $\Lambda^{20n(j+5)} \le M$. This proves both assertions. □

We next use the orthonormal middle-coordinate Fourier transform, rather than attempting Plancherel on the whole nonlinear quadric. This is essential: the full quadric Fourier transform would not have the asserted normalization or a uniform L^2 bound.

Proposition 23.7 (Sliced Plancherel)**.** *Suppose the source and target measures on a floor have fiber bounds M_s, M_t at their respective sizes, and let f be supported in the source size window. For a unit ξ, a fixed vector y, and the additive character ψ of the place write*

$$G(v) = \int \psi(\xi\langle v + y, x\rangle) f(x)\, d\eta_s(x).$$

On the target size window,

$$\int |G(v)|^2\, d\eta_t(v) \le M_t M_s \left(\frac{c}{\mathrm{md}(\xi)}\right)^{|\iota|} \int |f|^2\, d\eta_s, \tag{23.15}$$

where $|\iota|$ is the orthonormal rank, c is the one-coordinate Plancherel constant, and md *is the scalar Haar modulus. The measure on the receiver can be a floor quadric or floor volume.*

Proof. Use $x = (P, u, z)$ and $v = (Q, w, z')$. For fixed P, Q, the phase splits into $\psi(\xi\bar{w} \cdot u)$, factors depending on P, Q, u, z, y but not on w, z', and factors depending on P, Q, w, z' but not on u, z. To see this directly, the orthonormal part is orthogonal to the hyperbolic part. Even when the complementary coordinate s of x depends quadratically on u, it is paired with the fixed primary coordinate of v, not its varying w; similarly for the complementary coordinate of v. The contribution of the orthonormal part of the fixed y may be included in the first group of phases. The omitted factors have modulus one. Hence

$$|G(Q, w, z')| \le \int |\widehat{B}_{P,Q}(w)|\, dP, \qquad B_{P,Q}(u) = \int f(P, u, z)\, \chi_{P,Q,u,y}(z)\, d\zeta_s(z),$$

where $|\chi| = 1$, and the hat is the transform for $\psi(\xi\bar{w} \cdot u)$. Cauchy–Schwarz on the source fiber gives

$$\int |B_{P,Q}(u)|^2\, du \le \mu_s(P) \int |f(P, u, z)|^2\, du\, d\zeta_s(z).$$

The ordinary product Fourier Plancherel formula in the $|\iota|$ coordinates therefore gives, by Minkowski and Cauchy–Schwarz in P,

$$\begin{aligned} \left\| \int |\widehat{B}_{P,Q}(\cdot)|\, dP \right\|_2^2 &\le \left(\int \|\widehat{B}_{P,Q}\|_2\, dP \right)^2 \\ &\le \left(\frac{c}{\mathrm{md}(\xi)}\right)^{|\iota|} M_s \int |f|^2\, d\eta_s. \end{aligned}$$

Now integrate on the receiver: the majorant depends only on Q, w, and its integral over the permissible z' is bounded by $\mu_t(Q)$ times the majorant. Integrating over Q, w and using $\int \mu_t \le M_t$ proves (23.15). The argument first applies to integrable square-integrable chart functions, then to all square-integrable supported functions by truncation, the same norm bound, and convergence in L^2. □

In the source's complex-place normalization, the Fourier kernel is $\exp(2i\,\mathrm{Im}(\bar{w}u))$ (up to complex conjugation). Changing real Fourier variable from the standard 2π convention to this one multiplies the one-coordinate square norm by π^2; dilation by ξ multiplies it by $|\xi|^{-2}$. Thus $c = \pi^2$ and $\mathrm{md}(\xi) = |\xi|^2$. At the finite places with the source's self-dual normalization, $c = 1$ and the scalar Jacobian is $\mathrm{md}(\xi)$; the same statement follows by Haar Fourier inversion,

first for indicators of compact open cosets using orthogonality of characters, then by density of their linear span.

To apply this result to `Gauge.amplitude_support_and_plancherel`, the source density is $f(s) = k_{b+1}(sx, sy)$. It is left invariant under the stabilizer of the orbit point $s^{-1}f_{b+1}$. By the quotient measure identity $s \mapsto s^{-1}f_{b+1}$, it becomes a scalar function on the norm quadric, with unchanged L^2 norm. The size claim used in the application can be checked directly. Write $q = t\nu$ with $\nu \in U_{<b}$, $t \in H_b$, and suppose $f(s) \neq 0$. Then sx is near at floor $b+1$. Conjugating ν by $s \in H_b$ leaves it in $U_{<b}$, so $st^{-1}qx$ is still near there. Write $ust^{-1}qx$ in the size-1 box, with u fixing f_{b+1}. The inverse of a size-1 unitary matrix has size at most 5, by $X^{-1} = J^{-1}X^*J$ (each matrix product costs one extra size). Since qx has size j, the identity

$$ts^{-1}f_{b+1} = (qx)(ust^{-1}qx)^{-1}f_{b+1}$$

shows that this vector has size $j+6$. Thus Lemma 23.6 supplies $M_s = \Lambda^{99n(j+12)}$; the target factor is the stated M. Proposition 23.7 gives exactly the analytic bound

$$M\Lambda^{99n(j+12)}(c/\operatorname{md}\xi)^{|\iota|}\int |f(s)|^2\, d\theta_b(s)$$

of that Lean lemma, under its quotient-measure, invariance and near-support hypotheses. This does not establish the higher tower iterations by itself.

23.9 Two complex gauge estimates

We supply the two analytic gauge axioms at $\mathbb{C}$ with the source's constants. Write $B_j = \{z : |z| \le L^j\}$, $L \ge 4$, $\Lambda = L^2$, and let the Euclidean open ball have radius $1/4$. The gauge hypotheses include $L \ge r$ (in the application L is the bound on the matrix norm), so all size-addition axioms follow from the triangle inequality. The scalar volume is $\pi L^{2j} \le \Lambda^{j+1}$.

For the cone bound in (23.14), write $R = L^j$. In the flat chart on $d \le 10n$ hyperbolic pairs, $a = ir$, $r \in \mathbb{R}$ (here r denotes the real skew parameter), and $\tau = 1/2$. After the source's choice of skew unit and hyperbolic basis the norm is real and the real-norm term in $\bar{t}s$ has no imaginary part. Thus $|s| \le R$ forces $|t||r| \le R$. The permitted t-area is at most $\pi R^2 \min(1, |r|^{-2})$, and each of the remaining $2d-2$ complex coordinates is in a disc of area πR^2. Consequently the cone mass in a size-j window is at most

$$(\pi R^2)^{2d-2}\int_{\mathbb{R}} \pi R^2 \min(1, |r|^{-2})\, dr = 4(\pi R^2)^{2d-1} \le L^{198n(j+1)}.$$

The last inequality uses $\pi < 4 \le L$, $d \le 10n$, and $j \ge 0$. This proves the asserted complex cone estimate (the real one-dimensional integral is 4).

For the common-volume gauge condition `Gauge.com`, let the orthonormal rank be $a = s+1+3n$ with $a \le L$. Let C be the matrix obtained by combining the at most h rows of the test matrix and at most n negative-signature rows for each of the two transformed vectors, as in `ComplexGauge.lean`. More intrinsically these latter conditions say that the two vectors are in the positive eigenspace of the split Hermitian involution. Its negative eigenspace has dimension n, so the combined matrix has rank at most $h + 2n \le 3n$. Choose a matrix O with $s+1$ orthonormal columns spanning a subspace of its kernel; then $O^*O = I$ and $CO = 0$. For $u \in \mathbb{C}^a$ put $\varphi(u) = 36/|O^*u|^2$, with value $+\infty$ at zero denominator. This is measurable. If $r_u = |O^*u|^2 > 0$, for each $|c| \le \sqrt{r_u}/9$ set

$$w = O\left(\frac{c}{r_u}O^*u\right).$$

It satisfies $\bar{u} \cdot w = c$ and $|w| \le 1/9$. Both transformed vectors in the definition of `closePairs` have the same Euclidean norm as w: they lie in the positive eigenspace by $CO = 0$, and their Hermitian norms equal $|w|^2$ by unitarity. Their difference has length at most $2/9 < 1/4$, so they lie in `closePairs`, where the level unit is 1. The test constraint $vw = 0$ also follows from $CO = 0$. Hence the relevant set of values contains the disc of radius $\sqrt{r_u}/9$, and its inverse volume is at most $81/(\pi r_u) \le \varphi(u)$.

For the second part of the common-volume condition, two arbitrary pairs $(z_1, z_2), (z_1', z_2')$ in `closePairs` satisfy $|z_1|, |z_1'| < 1/4$ and $|z_2|, |z_2'| < 1/2$. The operator norm of J is 1. Hence the absolute value of the difference of their Hermitian pairings is at most $1/16 + 1/4 = 5/16 < 1/3$. A scalar c' satisfying the condition for every w must in particular satisfy it for the w constructed above with pairing $\sqrt{r_u}/9$. Therefore $|c'| \le 3/\sqrt{r_u}$, so the measure of its allowed values is at most $9\pi/r_u \le \varphi(u)$. The exceptional $r_u = 0$ case is automatically covered by $+\infty$.

It remains to bound the weight integral at size j. Complete O to a unitary coordinate system. The original polydisc lies in the polydisc of radius $S = L^{j+1}$ after this coordinate change because each new coordinate is a sum of at most $a \le L$ terms of original modulus at most L^j. For the first $s+1$ new coordinates, enlarge their domain to the ball of squared radius $aS^2 \le LS^2$; for the other $3n$ retain the discs. The density of $t = |O^* u|^2$ in the first $s+1$ coordinates is $\pi^{s+1} t^s / s!\, dt$. Thus

$$\begin{aligned}\int_{B_j^a} \varphi(u)^s \, du &\le (\pi S^2)^{3n} \frac{36^s \pi^{s+1}}{s!} \int_0^{LS^2} dt \\ &\le L^{3n(2j+3)} \, 4 \, 3^{144} L^{2j+3} \le L^{198n(j+1)}.\end{aligned}$$

Here $(36\pi)^s/s! \le e^{36\pi} \le 3^{144}$ and $4 \cdot 3^{144} \le L^{145}$, so the exponent in the second line is at most $(3n+1)(2j+3)+145 \le 198n(j+1)$ for $n \ge 1$. This supplies both common-volume bounds and their integral, including the source's uniformity in the transformed vectors and test matrix. It is the complex-place argument of `exists_orthonormal_ker_lintegral_le` and the analytic portion of `RealPlaceSizes.gauge`. At finite places the analogous common-volume argument requires a discrete lattice projection estimate; the complex proof does not apply there.

23.10 References for the real representation-theoretic interface

We can sharpen the references in Chapter 18 without treating qualitative classical results as replacements for the special quantitative bounds in this construction.

- Harish-Chandra, *Discrete series for semisimple Lie groups I: construction of invariant eigendistributions*, Acta Math. 113 (1965), 241–318, Theorem 3, p. 289, specifies the invariant eigendistribution with prescribed compact-Cartan alternant, infinitesimal character and discriminant bound. Part II, *Explicit determination of the characters*, Acta Math. 116 (1966), 1–111, Theorem 16, p. 96 and its proof pp. 99–100, identifies these distributions (after the discrete-series sign) with the characters of the discrete series, specifies their Weyl equivalence, and gives their formal degrees proportional to the absolute product of root pairings. We checked both theorem statements in the published papers (`https://archive.ymsc.tsinghua.edu.cn/pacm_download/117/5993-11511_2006_Article_BF02391779.pdf` and `https://repository.ias.ac.in/30833/1/336.pdf`). For $U(p,q)$ one passes from the semisimple derived group to the group with compact central torus by extending with the appropriate integral central character and the finite central identification. For an actual discrete series with its normalized invariant measure, the stable Weyl alternant and the relative degree polynomial in Chapter 18 are thus conventional inputs. This reference does not alone identify the

source's constructed function as the lowest-vector matrix coefficient, nor compare the proportionality constant with its chosen Haar measure.

- R. Parthasarathy, *Classification of discrete series by minimal K-type*, Represent. Theory 19 (2015), 167–185, Theorem 1.1, DOI https://doi.org/10.1090/ert/467, gives a discrete-series recognition criterion for an irreducible unitary representation satisfying the specified K-constituent and exclusion conditions (including its variant when the Harish–Chandra parameter is sufficiently regular). Its companion argument in §2 uses the Dirac operator. Thus the printed theorem is not the assertion for arbitrary, possibly reducible modules with just one Casimir equation and the growth bound in `Lib/DiscreteSeries/Rigidity/KernelFixes.lean`. To use it to replace that lemma, one must prove the exclusion condition on the generated constituents and their irreducibility or the appropriate direct-integral reduction, plus the required analytic-vector approximation. We have not done this here. In particular the local label rigidity used by the finite trace comparison is a different assertion, involving Galois attachments and types; it does not follow from this theorem about a real Lie group.

The two Harish-Chandra theorems concern invariant distribution characters, not an arbitrary explicit function on the group. They do not supply, in isolation, the normalized elliptic *orbital integral* identity of `Orbital.harish_chandra_degree_and_orbital`, nor its compatibility with the actual source normalization. We therefore distinguish the classical discrete-series character and relative degree formula from the remaining concrete kernel-identification, orbital-integral and local rigidity work.

23.11 Obligation-level ledger

The following records the exact status after these additions. "Proved" refers to an argument in the monograph, with explicitly stated elementary or classical premises. A proved implication does not discharge its premises. "Reference" means a conventional theorem of matching mathematical content has been identified precisely. "Open" means that at least one further mathematical step is needed for a conventional complete proof, not that the corresponding Lean statement is false.

- **REFERENCE:** the Tamagawa number 2 of the full unitary group for the nondegenerate Hermitian spaces in Proposition 23.1, by the cited theorem of Kirschmer. **PROVED:** orbit unfolding, the target-measure discriminant factor and local Fourier density with absolute convergence, giving the normalized one-vector genus average `classSum_norm_eq` for the data here. This uses the classical local-global Hermitian classification stated in the data chapter.

- **PROVED:** the local unramified conductor bound, the uniform off-window product estimate (23.4), the finite neighborhood test construction, the affine change of variables and its norm-fiber disintegration, and the resulting one-sided sharp count (23.6) (`siegel_sum_le_lintegral_quadric`). The special finite test functions used here need no general arbitrary-measure higher-rank Siegel theorem.

- **PROVED (implication):** cutoff rational amplitude identity (23.10), local constancy and labeled-coset packet lower bound (Lemma 23.4), the Gram positivity and packet-normalization deductions (23.11)–(23.13), and the dominated-convergence limit in Section 23.7. **OPEN:** the actual absolute integrated-amplitude summability, full marked-frame big-cell coordinate assertion, local support bound, and the stack's exact Haar

disintegration and entrywise integrability. The conditional identities do not assert these premises; in particular `sum_blocks_le_topKernel` is not an unconditional theorem of this chapter alone.

- **PROVED:** cone-to-fiber domination in Lemma 23.6 and all sliced analytic Fourier factors in Proposition 23.7; the support calculation gives the exact exponent $99n(j+12)$ used in `Gauge.amplitude_support_and_plancherel`, conditional on the stabilizer and orbit-point measure identities of the local level. Both complex analytic gauge axioms and their quantitative integral estimates have now been supplied. **OPEN:** finite nonarchimedean gauge and Haar comparisons, the actual construction and law of the floor pairing kernels, the iterated inverse-weight and Parseval estimates with their size and conjugation hypotheses, the bottom form, and the complete band system of `LocalTheorem.lean` and `BandLimits.lean`.

- **OPEN:** the simultaneous lattice saturation, weighted pair bounds, crude shell and energy majorants of `LinkPairing.lean`, `BlockBound.lean`, `CrudeEnergy.lean`, and `AmplitudeSums.lean`. They are not consequences of the one-vector Siegel formula or ordinary pointwise decay; a proof would need to retain the actual height exponents and uniform constants. In particular the standing inequalities and the sharp coherent estimate (22.25) have not been discharged.

- **OPEN:** the full simultaneous verification of `Filling.IsAdmissible` and `Data.filling_fits`, including the degree and mass thresholds, the common geometric size assumptions and normalization match for the exact parameters, not just their asymptotic orders in the rank. The earlier proofs of the group counts and component class mass do not establish all those inequalities.

- **REFERENCE:** the classical discrete-series character and relative formal-degree polynomial (Harish-Chandra I, Theorem 3; II, Theorem 16), as recorded above. **OPEN:** their extension to the actual normalized constructed kernel and its orbital integrals, the analytic Casimir/minimal-type rigidity in the source's stated generality, and the local Galois/finite-type uniqueness hypotheses required in the row transfer. The finite projection and trace/dimension implications of Chapter 18 remain proved reductions using those hypotheses.

These statuses explain both the advance of the chapter and the residual gap: the one-sided counting coefficient is now derived rather than posited, and the analytic slice estimate is an ordinary proof, but the global filling theorem cannot yet be declared unconditionally established as a conventional mathematical exposition. No global existence or analogue of the desired sharp energy bound has been used to prove these local statements.

Chapter 24

Finite gauge, orbit measures and the top-frequency transform

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 26 and the final audit in Chapter 57.

This supplement addresses the finite-place assertions left over in Section 23.11, especially the finite gauge of `New/Filling/StandardLevel.lean` and the top-frequency transform of `New/Filling/TopFloor.lean`. The local algebra is the finite product A of completions of the CM field at a conjugation-stable set of places, with integer ring R and involution denoted by a bar. Each factor over a base place is an unramified quadratic extension or the product of the two split factors. All vector-space measures in the finite gauge are normalized by $\mathrm{vol}(R) = 1$. The scalar modulus $\mathrm{md}(a)$ is the positive Haar Jacobian for multiplication by $a \in A^\times$. On the self-adjoint and skew-adjoint parts, use the source's integral trace and skew scalars: their integral lattices have mass one. For self-adjoint a their scalar Jacobian is $\mathrm{md}(a)^{1/2}$. These facts follow factor by factor from the unramified degree two or the split pair description. The additive character used by the lattice level is self-dual, that is, the annihilator of R is R.

24.1 The valuation integral and its cone consequence

Write q_w for the residue cardinal at a field factor w of A, so $\zeta_R(z) = \prod_w (1 - q_w^{-z})^{-1}$. For a vector $u \in A^d$ whose projection to every field factor is nonzero, let $I(u) = \sum_i Ru_i$ be its content fractional ideal. There is a unit $t(u) \in A^\times$ generating it, by the principal ideal property of the finite product of discrete valuation rings. Its modulus is determined by the ideal. Put

$$C(u) = \mathrm{vol}\{c \in A : cu_i \in R \text{ for all } i\} = \mathrm{md}(t(u))^{-1}.$$

The remaining vectors are a Haar-null subset; give C value $+\infty$ there. This function is measurable since the valuation shells are measurable. For $d > s \geq 0$ the following exact valuation computation holds:

$$\int_{R^d} C(u)^s \, du = \prod_w \frac{1 - q_w^{-d}}{1 - q_w^{-(d-s)}} \leq \zeta_R(d - s). \tag{24.1}$$

Indeed in the factor w, the event $\min_i v_w(u_i) \geq \ell$ has measure $q_w^{-d\ell}$ for $\ell \geq 0$, and the shell of equality has measure $(1 - q_w^{-d})q_w^{-d\ell}$. The content index there is q_w^ℓ. Sum the geometric series and take the product of the independent factor integrals. For u uniformly in the lattice

$\alpha^{-j}R^a$, a projection onto d coordinates after applying any matrix in $\mathrm{GL}_a(R)$ has the same distribution, under normalized lattice measure, as α^{-j} times the first d lattice coordinates. In particular, if Y is any $a \times d$ matrix completing to such an invertible matrix, with $d > s$,

$$\int_{(\alpha^{-j}R)^a} C(Y^*u)^s \, du \le \zeta_R(d-s)\,\mathrm{md}(\alpha)^{-j(a-s)} \tag{24.2}$$

for self-adjoint $\alpha \in R \cap A^\times$. To see the scaling, change u to $\alpha^{-j}v$, for which the measure factor is $\mathrm{md}(\alpha)^{-ja}$ and the content index gains the factor $\mathrm{md}(\alpha)^j$. The conjugate transpose of a completed integral matrix also belongs to $\mathrm{GL}_a(R)$. This supplies the particular (and slightly stronger) content-index lemma `IntegralPlace.lintegral_contentIndex_pow_le` needed in the gauge.

Lemma 24.1 (Finite hyperbolic-cone mass)**.** *For a sum of $D \ge 2$ hyperbolic planes, write the norm as $Q(x,y) = \mathrm{Tr}_{A/A^+}(\bar{x} \cdot y)$, where $x, y \in A^D$. Let $d\delta_0$ be the norm-zero disintegration measure. If α is as above and $j \ge 0$, then*

$$\delta_0((\alpha^{-j}R)^{2D}) \le \zeta_R(D-1)\,\mathrm{md}(\alpha)^{-(2D-1)j}. \tag{24.3}$$

Proof. First take $j = 0$. In each unramified nonsplit factor, use a row operation over the integer ring to transform $\bar{x}$ to $(\bar{t}, 0, \dots, 0)$, t a generator of its content; this exists by elementary valuation elimination (choose an entry of minimum valuation and subtract its integral multiples). On $y \in R^D$ the push-forward by $Q(x,y)$ is normalized Haar on the trace of the content ideal in the base ring. Since the extension is unramified, the trace pairing is perfect and the trace of its integer ring is the base integer ring, also in residue characteristic two. The density at zero is q_v^ℓ if t has valuation ℓ and q_v is the base residue size: it is at most the factor q_w^ℓ of $C(x)$, where $q_w = q_v^2$. At a split pair, the trace is the sum of the two linear functionals from the factors. The image is the sum of the two content ideals in the base ring, with index $q_v^{\min(\ell_1,\ell_2)} \le q_v^{\ell_1+\ell_2}$. Again it is at most the corresponding product factor of $C(x)$. Except at a null set of x, these push-forward densities are constant in a neighborhood of zero; integrating the bound $C(x)$ over the remaining variable gives (24.1) with $d = D, s = 1$. This argument applies to the disintegration measure itself, not just a formal delta distribution: the trace-linear chart is a product chart whenever x is nonzero in every field-factor pair, the density is the Haar change of variables for the linear map, and the singular part where all such primary coordinates vanish has measure zero in the flat chart. Equivalently integrate on shrinking norm balls and use dominated convergence with integrable majorant $C(x)$, whose integral was just computed. For the general j, change (x,y) to $(\alpha^{-j}x, \alpha^{-j}y)$. Ambient Jacobian is $\mathrm{md}(\alpha)^{-2Dj}$; the target norm has Jacobian $\mathrm{md}(\alpha)^{-j}$ under scaling by α^{-2j}. Division gives exactly the exponent $2D-1$. □

In `StandardLevel.lean` the middle space has $m+1$ further hyperbolic planes besides the primary plane, and the cone test arising from a floor has $m+1 \le D \le m+h+2$. In its notation the admissible bound is $\zeta_R(m)^n \le 1 + 2^{-n}$, $n = h+1$, with $m < 9n$. Thus $\zeta_R(D-1) \le \zeta_R(m) \le 2$. This proves the cone assertion there, and hence the finite `Gauge.fib` axiom of Section 23.8 for the size sets $B_j = \alpha^{-j}R$ and $\Lambda = 2/\mathrm{md}(\alpha)$: for $D \le 10n$ the right side of (24.3) is at most $\Lambda^{99n(j+1)}$. It retains the significantly sharper bound $\zeta_R(m)$ when $\alpha = 1$.

24.2 The common-volume finite gauge

We next prove the common-volume axiom at the actual finite lattice level, not merely on a split completely diagonal form. Let J be its unimodular integral Gram matrix, $g \in R \cap A^\times$

the self-adjoint congruence unit, K the integral unitary group and $C = K(g)$ the subgroup congruent to one modulo gR. The ball is the integral vector lattice R^r, and the positions are products $y^{-1}uyk$ with $u \in U$ the Bessel unipotent, $y \in G$ and $k \in C$. Let E denote the integral matrix of the orthonormal part of the middle hyperbolic basis; it satisfies $E^*JE = 1$. Let $a = |\iota|$ be its number of columns. In this section the volume exponent is

$$s = a - (3n + m + 2) \geq 0.$$

In the source's notation this is volumeExponent for the middle basis with $m + 1$ hyperbolic planes. The pairs called close are precisely

$$\mathcal{D} = \{(z, z') : z \in R^r, \quad (z - z')/g \in R^r\}.$$

For fixed $x \in K$, a position $b = y^{-1}uyk$, and a test matrix v with h rows, the gauge uses

$$\mathcal{W} = \{w \in A^a : vw = 0,\ (x^{-1}Ew, (xb)^{-1}Ew) \in \mathcal{D}\}.$$

We claim that $\mathcal{W}$ contains YR^d for some $d = a - 3h = s + m + 5$ and some $a \times d$ matrix Y completing to $\mathrm{GL}_a(R)$. This is the lattice version of the orthonormal projection argument over $\mathbb{C}$.

Proof of the claim: write $N = yx^{-1}E$. In the ordered frame basis (first h isotropic vectors, the full middle block, their h duals in reverse order), U is block upper unitriangular, and on a vector whose dual frame coordinates vanish, $(u^{-1} - 1)$ has image in the first h isotropic coordinates. Impose the following at most $3h$ homogeneous linear equations on w: $vw = 0$; the dual coordinates of Nw are zero; the first h coordinates of $(u^{-1} - 1)Nw$ are zero. Their common kernel over every field factor of A has dimension at least $a - 3h$. Its intersection with the integral vector lattice is a saturated submodule, since an integral multiple by a nonzero scalar of a vector satisfying a linear equation satisfies it only when the vector does. Over a DVR a saturated submodule of a finite free module is free and a direct summand: its quotient is torsion free and hence free (or use Smith normal form after clearing denominators). Choose the requisite d primitive basis columns in each factor and assemble them into Y. Then for $w \in YR^d$ one has $(u^{-1} - 1)Nw = 0$, and hence

$$(xb)^{-1}Ew = k^{-1}x^{-1}Ew.$$

Since x, E are integral, $x^{-1}Ew \in R^r$. Since $k^{-1} \equiv 1 \pmod{gR}$, the normalized difference is integral. This proves the claim.

Put $f(z) = C(Y^*z)$, with the null exceptional value specified above. For fixed z, the set $\{\bar{z} \cdot w : w \in \mathcal{W}\}$ contains the fractional ideal generated by the entries of z^*Y, whose measure is $1/f(z)$; its inverse volume is thus bounded by $f(z)$. This is the first common-volume condition. For the second, suppose c is such that, for every $w \in \mathcal{W}$, the scalar $\overline{cz} \cdot w$ can be expressed as $g^{-1}(h(z_1, z_1') - h(z_2, z_2'))$ for two pairs in $\mathcal{D}$. Such a scalar must lie in R: substitute $z_2 = z_1 - g\delta$, $z_2' = z_1' - g\delta'$ with $z_1, z_1', \delta, \delta'$ integral and use that J, g are integral and $g = \bar{g}$. In particular $\bar{c}\, z^*Y$ is integral coordinate by coordinate, testing on the standard columns of Y. The involution preserves the additive measure, so the measure of this set of c is at most $C(Y^*z) = f(z)$. This is exactly the second common-volume condition in `LevelGauge.lean`.

Finally (24.2) gives

$$\int_{B_j^a} f(z)^s \, dz \leq \zeta_R(m + 5)\, \mathrm{md}(\alpha)^{-j(a-s)} \leq \Lambda^{99n(j+1)}.$$

Indeed $a - s = 3n + m + 2 \leq 14n$, $\zeta_R(m + 5) \leq 2$, and $\Lambda \geq 2$. The bound holds uniformly in x, b, v, including the split factors where the kernels of the constraint matrices may have

different dimensions: we only choose the fixed smaller dimension d in every factor. The rest of the finite gauge axioms now follow directly from the size sets B_j: $0, 1 \in B_j$, addition does not enlarge a ball in any factor, multiplication joins the exponents, and the integral Gram matrix, its inverse, the basis and the fixed trace/skew scalars have size 1. Also $\mathrm{vol}(B_j) = \mathrm{md}(\alpha)^{-j} \le \Lambda^{j+1}$ and if $a' \in A^\times \cap B_j$, $\mathrm{md}(a') \le \mathrm{md}(\alpha)^{-j} \le \Lambda^j$. Compactness and measurability are those of fractional ideal lattices. Together with Lemma 24.1 this proves the finite gauge `IntegralPlace.IsStdLevel.gauge` assuming the standard lattice-level data, with all their quantitative bounds; it does not need a general valuation-pair elimination theorem beyond the DVR facts just proved.

24.3 Actual finite correlation supports

The finite vector in `TopFloor.lean` is W_Z for $C \subseteq Z \subseteq K$, with coordinates $W_Z(y)_a = \phi(ya^{-1})$ for a representative of $C\backslash Z$. Here $\phi(uk) = \psi(g^{-1}\,\mathrm{sr}(u))$ on UC and zero off UC, where sr is the sum of the simple root coordinates. This function is well defined because sr is additive on U and divisible by g on $U \cap C$ (simple roots are entries of the frame block), and ψ is trivial on R. It is right C-invariant and left U-equivariant; its absolute value is at most one. As C is open and normal in K, all coordinates, and thus W_Z, are right C-invariant and continuous. In particular support of W_Z implies membership of $UZ \subset UK$; W_Z is not itself compactly supported on G.

For a full head p (of length h with all entries units of A) let w_p be the Weyl element, and $M = H_h$ the unitary group of the entire middle space. As in the source, M commutes with w_p. In the ordered frame basis, conjugation by w_p takes U to the opposite block lower unitriangular group U^-. We give a direct support bound sufficient for the top correlation, using no inverse-weight estimate.

Lemma 24.2 (Uniform compact support of the overlapping product)**.** *For p varying in a sufficiently small compact open neighborhood of a fixed head, there are compact sets $Y \subseteq G$, $M_0 \subseteq M$ independent of p such that*

$$W_Z(y) \neq 0, \quad W_Z(\ell w_p^{-1} y) \neq 0, \quad \ell \in M \quad \Longrightarrow \quad y \in Y,\ \ell \in M_0.$$

Proof. Write $y = uk$, $\ell w_p^{-1} y = u'k'$ with $u, u' \in U$ and $k, k' \in K$. Then

$$w_p k' k^{-1} = (w_p u'^{-1} w_p^{-1})\, \ell\, u. \tag{24.4}$$

The left side ranges over a compact subset of the unitary group, because p and p^{-1} range over compact sets. The right side is a block lower-unitriangular, diagonal, upper-unitriangular decomposition in the ordered frame basis. The first h and last h diagonal blocks are scalars equal to one; the middle block is ℓ. Work componentwise in the fields of A. Every leading principal block determinant (at the ends of these blocks) is therefore 1 before the full middle block and $\det \ell$ thereafter. The latter equals $\det(w_p k' k^{-1})$ and its inverse is bounded on the compact set of the left sides (unitary inversion is continuous). All these determinants are uniformly bounded away from zero componentwise. Block Gaussian elimination, performed by taking successive Schur complements, expresses each block of the three factors uniquely as rational functions in the entries of the left side with these leading principal block determinants as denominators (invert a pivot by the adjugate and determinant). In particular all entries of u and ℓ are uniformly bounded in every field. Since the factors are isometries, their inverses are $J^{-1} g^* J$ and bounded as well. The sets of matrices satisfying fixed coordinate bounds and the isometry equations are compact; the equations for M are closed. Multiplying the compact bound for u by K bounds y as required. This proves (24.2) without needing any bound on u from support of $W_Z(y)$ alone. □

The source's finite correlation $\mathcal{J}(p,\ell)$ is the (positive real head-density multiple of the) integral against full group Haar measure of $W_Z(y)\otimes\overline{W_Z(\ell w_p^{-1}y)}$. Lemma 24.2 makes this an ordinary absolutely convergent vector integral, compactly supported in ℓ. It also proves the uniform local constancy required in Section 23.5, as follows. In a neighborhood of a fixed pair $(p,1)$, the density of the head does not change, since its scalar absolute values are locally constant. Replace (p,ℓ) by $(p',\ell\eta)$, with $\eta\in M$ near one. If either product defining the two correlations is nonzero, the corresponding y belongs to the single compact Y of Lemma 24.2. The ratio between the second arguments of W_Z is

$$y^{-1}w_p\eta w_{p'}^{-1}y.$$

Uniformly on compact Y this belongs to C for all sufficiently close (p',η), by continuity of multiplication and openness of C. Right invariance implies equal second vectors; if both products are zero the equality is automatic. This proves $\mathcal{J}(p',\ell\eta)=\mathcal{J}(p,\ell)$ uniformly in $\ell\in M$. Thus the actual support and local constancy premises of Lemma 23.4 at a standard finite level are discharged.

24.4 Exact invariant measures of the stacks

We next justify the use of the top-stack measure, including its normalization. The finite standard-level data include the hyperbolic basis, H_{h+1}-Haar measure μ_H with integral subgroup volume one, and a measure θ_h supported in $M=H_h$ with push-forward under $s\mapsto s^{-1}z$ equal to the prescribed norm-fiber measure δ_{-N}. The latter is part of `Level.theta_spec`, not a Haar measure on M itself. The norm fiber consists of nonzero vectors and its orbit map has stabilizer $H=H_{h+1}$. Locally diagonalize a nonzero norm vector and apply Witt extension to get transitivity (over a split pair this is the ordinary dual vector-pair action of the general linear group). Its invariant density is the disintegration of product Haar on the vector space by the norm, since isometries preserve the ambient additive measure and the norm function. It has no singularity at a vector of nonzero norm. Standard homogeneous-space integration can here be seen directly: for a nonnegative continuous compactly supported function F on M, the value

$$F_*(e)=\int_H F(hs)\,d\mu_H(h),\qquad s^{-1}z=e,$$

is independent of the lift by right invariance of μ_H, continuous in local smooth sections and zero outside the compact projection of the support. Right translation of F by $a\in M$ has associated function $e\mapsto F_*(a^{-1}e)$. Invariance of δ_{-N} therefore shows that $F\mapsto\int F_*d\delta_{-N}$ defines right Haar on M, with exact normalization, nonzero and finite on compact sets. Both H and M are unimodular reductive groups; for the unitary groups this follows from the determinant-one adjoint action on their Lie algebras (after base change, conjugation acts on all matrices with total determinant one). In particular $\lambda_h:=\mu_H*\theta_h$ is two-sided Haar and disintegrates with the specified measures, since pushing θ_h through the orbit map gives δ_{-N} exactly. This is `Level.stackMeasure_top`.

Here is the analogous elementary normalization at each lower hyperbolic floor, giving Haar on the full group rather than merely on the top stack. For $b<h$, H_b acts on the nonzero isotropic vectors in its space $V_b=Ae\oplus V_{b+1}\oplus Ae^*$, with $h(e,e^*)=1$. In split factors the open chart under consideration consists of vectors with unit primary coordinate in A (a nonzero coordinate in every factor); its orbit is the open orbit of vectors with both vector and covector nonzero. Its complement is null for the flat-chart measure. The stabilizer of e in H_b is the semidirect product of H_{b+1} and the Heisenberg unipotent U_b fixing e. In coordinates (v,η)

for $v \in V_{b+1}$ and skew η, its matrix is

$$n(v,\eta) = \begin{pmatrix} 1 & -v^* J' & \eta - \tau h(v,v) \\ 0 & 1 & v \\ 0 & 0 & 1 \end{pmatrix},$$

where J' is the Gram matrix on V_{b+1}, $\tau + \bar{\tau} = 1$, and columns are ordered (e, V_{b+1}, e^*). It is checked by multiplying its adjoint against the hyperbolic Gram matrix. The group law adds v and adds η with a bilinear skew correction. Consequently product measure $dv\,d\eta$ is both left and right Haar on U_b, by triangular translations. Conjugation by H_{b+1} preserves this product Haar, since its action on v is an isometry of determinant modulus one and fixes the central skew coordinate. The product Haar on the stabilizer is therefore precisely the convolution of the H_{b+1} Haar measure with the Heisenberg measure ν_b of these coordinates.

The bottom-to-top orbit law defining the b-th floor measure in the source is the law of $s = w_b(c)^{-1} n(v,\eta)$ with $d\theta_b = \mathrm{md}(c)^{d_b}\,dc\,dv\,d\eta$, $d_b = \dim_A V_{b+1}$. Compute $s^{-1}e = n(v,\eta)^{-1} c e^*$. Its e^* coordinate is c, its middle coordinate is $-cv$, and its e coordinate solves the norm-zero equation, with skew chart variable $-\eta$ up to the translation by a function of v coming from the trace-one convention. In fact inversion in these coordinates sends (v,η) to $(-v, -\eta + (\tau - \bar{\tau})h(v,v))$; the possible skew translation does not change its measure. Thus replacing v by $-cv$ on the flat chart costs the Jacobian $\mathrm{md}(c)^{d_b}$ and the skew variable changes by a measure-preserving translation and sign. The resulting law is exactly the flat-chart measure $dc\,d(-cv)\,d\eta$ on the isotropic cone, which is the source's norm disintegration. The observation of the previous paragraph about homogeneous-space integration applies now to the stabilizer of e. Each floor consequently forms

$$\lambda_b = \lambda_{b+1} * \nu_b * \theta_b$$

as a right Haar measure with the established normalization of the orbitpoint push-forward. In this particular argument the isotropic vector orbit is smooth; in the split case, the omitted component consisting of a zero vector or covector has zero measure in these chart coordinates. Invariance of the cone measure on the smooth open orbit follows locally from the norm submersion: one may use any coordinate with nonzero primary component to obtain the disintegration, and the different charts agree by change of variables. The exceptional chart complements have zero disintegration measure (use the trace-linear coarea calculation of Lemma 24.1, locally on compact subsets), so the homogeneous-space argument uses the entire invariant measure on that orbit.

Inducting from λ_h proves that the full stack $\mu = \lambda_0$ is actual Haar on G. The convolution of the first h lower factors, denoted $\sigma_h = \mathrm{stackMeasure}(0,h)$, satisfies exactly $\mu = \lambda_h * \sigma_h$ by associativity. No proportionality constant is lost in this proof, because each quotient measure is prescribed by the explicit chart, and each Haar measure on the stabilizer has already been normalized. This closes the Haar and orbit-distribution parts of the stack assumptions for the source's standard level. Constructing the standard level in the first place (the hyperbolic integral basis, the marked middle section and its unit-box lower bound) is a logically prior datum; the argument uses those defining properties and does not treat an arbitrary Hermitian form without such a basis as a standard level.

24.5 The exact top-floor Fourier identity

The preceding compactness result and measure calculation permit a direct proof of the identity `floorForm_top_eq_integral` without deriving it from the quantitative band theorem.

For two heads p, q, put $J_p(\ell) = \mathcal{J}(p, \ell)$, with head factor $\kappa(p) = \mathrm{headDensity}(h, p, 1)$, so $\kappa(p)\kappa(q) = \mathrm{headDensity}(h, p, q)$. Let χ be any complex-valued Borel function on M of modulus at most one that depends only on the scalar $\langle z, \ell^{-1} z\rangle$. In particular the character $\psi(\xi\langle z, \ell^{-1} z\rangle)$ satisfies this condition. Then the exact identity is

$$\mathrm{floorForm}_h(p, q; \chi) = \int_M \chi(\ell)\langle J_p(\ell), J_q(\ell)\rangle \, d\lambda_h(\ell). \tag{24.5}$$

In this notation on the left, the argument of χ in the double floor integral is the inverse of the element st^{-1} if χ is understood as depending on ℓ^{-1} on the right; equivalently use the function $\chi_0(st^{-1})$ for $\chi(\ell) = \chi_0(\ell^{-1})$. This is precisely the convention of `Level.formIntegrand` and `floorForm_top_eq_integral`.

Proof. Expand the tensor inner product on the right after canceling the common factor $\kappa(p)\kappa(q)$. In the conjugate-linear-first convention it is the integral of

$$\chi(\ell)\langle W(a), W(b)\rangle\overline{\langle W(\ell w_p^{-1} a), W(\ell w_q^{-1} b)\rangle} \tag{24.6}$$

with variables $\ell \in M$, $a, b \in G$, and measures $d\lambda_h(\ell) d\mu(a) d\mu(b)$. Every factor is bounded. Nonvanishing requires in particular $W(a), W(\ell w_p^{-1} a) \neq 0$ and $W(b), W(\ell w_q^{-1} b) \neq 0$. By Lemma 24.2 the entire integrand is compactly supported in the triple, so it is absolutely integrable. This verifies rather than assumes the integrability of the expanded tensor product.

Write $a = h_1 s y$ with $h_1 \in H$, $s \in M$ and $y \in G$ under $d\mu_H(h_1) d\theta_h(s) d\sigma_h(y)$. This distribution is μ by the exact convolution of Section 24.4. For a fixed write $b = ax$ under $d\mu(x)$, whose distribution is μ by left invariance. For $h_1 s$ fixed write $\ell = h_2 t (h_1 s)^{-1}$ under $d\mu_H(h_2) d\theta_h(t)$, whose distribution is λ_h by right invariance. Pulling back (24.6) gives the integrand

$$\chi(h_2 t s^{-1} h_1^{-1})\langle W(h_1 s y), W(h_1 s y x)\rangle\overline{\langle W(h_2 t w_p^{-1} y), W(h_2 t w_q^{-1} y x)\rangle}.$$

Here we used that all of H, M commute with both w_p, w_q. Since H fixes z, the first scalar factor depends only on st^{-1} through $\chi_0(st^{-1})$. First integrate independently over h_1, h_2 using their Haar measures. These are exactly the upper-pairing averages k_{h+1} of the two source kernels: the first is evaluated at (sy, syx) and the second at $(tw_p^{-1} y, tw_p^{-1} yx)$ with twist $w_q^{-1} w_p$ on the second argument. The remaining integrals over s, t, x, y are the double floor integral defining $\mathrm{floorForm}_h$ in `LocalTheorem.lean`, with the head factor already canceled. The pull-back is absolutely integrable because its push-forward is the integrable triple; Tonelli for absolute values followed by Fubini justifies all the changes of order. Reinstate the common head factor to obtain (24.5). □

In particular for a compact open additive subgroup L of A, put $\Phi_{p,q}(\ell) = \langle J_p(\ell), J_q(\ell)\rangle$ and $r(\ell) = \langle z, \ell^{-1} z\rangle$. The source's top kernel is

$$K_L(p, q) = \int_M 1_{r(\ell) - p_h \in L} \Phi_{p,q}(\ell) \, d\lambda_h(\ell).$$

For common last entries $p_h = q_h$, this is a positive semidefinite Gram matrix, by integration over one common set; Hermitian symmetry follows directly. It is jointly locally constant in its head parameters by the uniform local constancy above, together with local constancy of the condition on p_h. The top-stack norm integral needed for the one-sided packet estimate is consequently finite. In addition the exact Fourier identity connecting the top kernel to the ordinary floor form is

$$\int_A \frac{K_{L,t}(p, q)}{\mathrm{vol}(L)} \psi(\xi t) \, dt = \mathrm{floorForm}_h(p, q; \psi(\xi r)) \frac{1}{\mathrm{vol}(L)} \int_L \psi(-\xi v) \, dv, \tag{24.7}$$

where $K_{L,t}$ means the kernel with the same first h entries of p, q, and t in its last entry. To prove this, interchange the compactly supported integrals, then substitute $v = r(\ell) - t$ in the inner integral over t. In the finite character normalization, the last factor is 1 if the frequency annihilates L and 0 otherwise; its absolute value is always at most 1. This gives exactly the top link in the descent of the finite sieve system: the sharp frequency estimate for the top floor form, once established, also bounds its averaged top kernel, without a factor depending on $\mathrm{vol}(L)$. The frequency estimate itself is not being inferred from (24.7).

24.6 Finite Haar masses and the pairing norm

The gauge and orbit calculation also give the exact finite mass normalization needed by the sieve. First the integral middle hyperbolic basis of the standard level preserves the full integral vector lattice. At any floor, the norm-fiber mass of integral vectors is bounded by $\zeta_R(m)$: fix the primary and the hyperbolic first coordinates and then the orthonormal coordinates, as in Lemma 23.6. Translate a nonempty fiber by one of its points, keeping the primary and first hyperbolic coordinates. Their complements satisfy the zero-norm hyperbolic-cone equation. At a finite place subtraction does not increase the integral box. Its mass is bounded by (24.3) with $j = 0$, and the integral orthonormal coordinates have product measure 1. This applies to the nonzero norm of the top floor also. Thus in each disintegration from Section 24.4, the integral group $K_b = H_b \cap K$ has mass at most the next integral stabilizer mass times $\zeta_R(m)$. Indeed if s has a lift in K_b, the set of stabilizer elements a with $as \in K_b$ is a coset of the integral stabilizer, and otherwise it is empty. On a hyperbolic floor the stabilizer integral mass is exactly $\lambda_{b+1}(K_{b+1})$, since its middle block and its Heisenberg coordinates are integral precisely when all the entries are integral and the skew coordinate then has integral measure one. The top stabilizer has prescribed integral mass one. The orbit point of an integral matrix is integral, so the claimed estimate follows from the norm-fiber mass bound. Iterating over all $n = h + 1$ floors yields

$$\mu(K) \le \zeta_R(m)^n \le 1 + 2^{-n}. \tag{24.8}$$

For the reverse bound, the finite standard-level condition has $\theta_h(K) \ge \mathrm{vol}(R^\times)$ and $\mu_H(K) = 1$. For each hyperbolic floor the Weyl parameter $c \in R^\times$ gives an integral Weyl element and has density $\mathrm{md}(c)^{d_b} = 1$. The integral Heisenberg parameters likewise have product measure one. The mass of K under the convolution product is at least the product of these masses, because a product of integral group elements remains integral. Hence

$$\mu(K) \ge \mathrm{vol}(R^\times)^n,$$

and, if the nonunits have measure at most $1/2$, this is at least 2^{-n}. This proves the group-mass comparisons of `StandardLevel.lean`, including their constants, from the defining standard-level properties.

The Bessel unipotent U is the product in floor order of the Heisenberg subgroups U_b; extracting its outer hyperbolic right column and then its middle block gives the uniqueness of this factorization. Each floor acts on lower-coordinate measures by determinant-one triangular translations as in Section 24.4; equivalently induction along the upper-unitriangular blocks shows that the convolution of the product coordinate Haar measures is Haar on U. Call it ν. The intersection with K has mass one. For the principal congruence subgroup $C = K(g)$, the Heisenberg matrix displayed in Section 24.4 is congruent to one if and only if $v \in gR^{d_b}$ and $\eta \in gR_{\mathrm{skew}}$. The upper-right quadratic term is already in gR when $v \in gR^{d_b}$, since g is integral and self-adjoint. Its coordinate mass is $\mathrm{md}(g)^{d_b+1/2}$, the exact floor modulus. In the full unipotent factorization, extraction of the outer column preserves the congruence

condition, and after removing its first Heisenberg factor the remaining middle block has the same condition; induction verifies it separately on every factor. Thus

$$v := \nu(U \cap C) = \prod_{b<h} \mathrm{md}(g)^{d_b+1/2} = \mathrm{unipotentMass}\,. \tag{24.9}$$

Writing $\eta = \mathrm{upperMeasure}(0)$ for the convolution of the floor orbit measures alone, one has the exact equality $\mu = \nu * \eta$. This follows by commuting each lower Heisenberg measure past the subsequent higher-floor measures (supported in the appropriate middle group, whose conjugation has determinant modulus one on that Heisenberg radical). It is the invariant-measure identity `stackMeasure_zero_eq`; no Haar measure on $U\backslash G$ needs to be normalized independently. The variables called η here are measures, distinct from the scalar skew coordinates used above.

Lemma 24.3 (Exact finite pairing identity)**.** *For every $y, x \in G$,*

$$v\langle W_Z(y), W_Z(yx)\rangle = \int_U 1_Z(uy)\phi(uyx(uy)^{-1})\, d\nu(u). \tag{24.10}$$

In particular, with $k_0(1,x) = \int \langle W_Z(y), W_Z(yx)\rangle\, d\eta(y)$ (the source's untwisted averaged kernel at floor zero), one has

$$vk_0(1,x) = \int_Z \phi(zxz^{-1})\, d\mu(z). \tag{24.11}$$

Proof. Partition $\{u \in U : uy \in Z\}$ by the cosets Ca in Z. In a coset Ca the set of u is empty or a left coset of $U \cap C$, hence has measure v. On it let $uy = ca$. The left Bessel covariance gives $W_Z(y)_a = \psi(-g^{-1}\,\mathrm{sr}(u))$ and $W_Z(yx)_a = \psi(-g^{-1}\,\mathrm{sr}(u))\phi(uyxa^{-1})$. Their inner-product contribution is therefore $\phi(uyx(uy)^{-1}c) = \phi(uyx(uy)^{-1})$ by right C-invariance. A coordinate with no such u has $W_Z(y)_a = 0$ and contributes nothing. Integrating over these finitely many cosets proves (24.10). Its right side after integration against η is the integral on Z in (24.11), since $\mu = \nu * \eta$. Absolute integrability follows by bounding $|\phi| \le 1$ and integrating $1_Z(uy)$ on the convolution, whose mass is $\mu(Z) < \infty$. □

Proposition 24.4 (Normalized finite pairing norm)**.** *With $m_Z = \mu(Z)\mu(C)/v^2$ as in (23.12), one has*

$$v \iint |\langle W_Z(x), W_Z(y)\rangle|^2\, d\eta(x)d\eta(y) \le m_Z. \tag{24.12}$$

Proof. Let $a(x) = \int_U 1_Z(ux)d\nu(u)$. It is between zero and one: if nonempty, choose u_0 with $u_0x \in K$, and then all u in the integral with $ux \in Z$ lie in $(U \cap K)u_0$, of measure one. Also $\int a(x)d\eta(x) = \mu(Z)$ by the convolution identity. By Lemma 24.3, $v|\langle W_Z(x), W_Z(y)\rangle| \le a(x)$, taking $x^{-1}y$ in place of the right-translation argument of that lemma.

If $W_Z(x) \neq 0$, fix $a_0 = u_0x \in K$. Nonzero inner product with $W_Z(y)$ implies at least one common nonzero coordinate, so there are $u_1, u_2 \in U$ with $u_1x, u_2y \in Ca$ for the same $a \in Z$. Their quotient $(u_1x)^{-1}u_2y$ is in C, by normality in K. Hence $y \in Ua_0C$. For any y in this set the measure of $\{u : uy \in a_0C\}$ is exactly v, since $a_0Ca_0^{-1} = C$; outside it the measure is zero. Integrating these indicators with η gives $v\eta(Ua_0C) = \mu(a_0C) = \mu(C)$. Thus for each x with possible nonzero pairing the support of y has η-measure at most $\mu(C)/v$. Combine this with the pointwise bound and $a(x)^2 \le a(x)$:

$$v^2 \iint |\langle W_Z(x), W_Z(y)\rangle|^2\, d\eta(x)d\eta(y) \le \frac{\mu(C)}{v}\int a(x)^2\, d\eta(x) \le \frac{\mu(C)\mu(Z)}{v}.$$

Division gives (24.12), the exact `pairingNorm_mul_le_massRatio` of `TopFloor.lean`. □

24.7 The finite bottom estimate and its remaining error term

Write $B(Z)$ for the bottom form

$$B(Z) = \int_G \left| \int_Z \phi(yxy^{-1})\, d\mu(y) \right|^2 d\mu(x).$$

For $x \in C$, conjugating by $Z \subseteq K$ preserves C and ϕ is 1 on C, so the contribution of C to this integral is exactly $\mu(Z)^2\mu(C)$. Outside C, Lemma 24.3 identifies the integrand with $v^2|k_0(1,x)|^2$. The source's extra positive error term is therefore precisely the bound

$$v^2 \int_{G\setminus C} |k_0(1,x)|^2, d\mu(x) \le \mu(Z)\mu(C)\,\mathrm{vol}(R \setminus R^\times)^{t+1}2^{-n}, \qquad t = k - 10^{10}n^4. \tag{24.13}$$

This is the substantive remaining estimate in `floorZeroForm_le_main_term`. It is not a consequence of the elementary pairing norm of Proposition 24.4: the variables and the absence of an inner averaging modulus there are different. For this estimate the source uses a capped pivot inequality and weighted shell integration in `lintegral_shells_le`. We record it explicitly rather than use the fact that ϕ is supported on UC to suggest falsely that conjugating by Z gives support only on C.

The ordinary bottom floor form itself, before its upper bound, is exactly $B(Z)/v^2$. Indeed at floor zero the head density is one, the lower stack is the point mass at the identity, and the two independent floor integrals of the first upper pairing and its conjugate give $|k_0(1,x)|^2$ before integrating x against μ. Absolute convergence of each upper pairing in the full floor variables follows from $v|\langle W(y), W(yx)\rangle| \le a(y)$ and $\int a(y)d\eta(y) = \mu(Z)$, proved above; one may interpret the final nonnegative integral in the extended positive reals until its finiteness is shown. Its normalization by m_Z is therefore exactly $B(Z)/(\mu(Z)\mu(C))$, not merely an inequality in that direction.

Given (24.13), the fixed-vector factor in `Filling.IsFillingLevel.exists_finite_sieveSystem` has a direct arithmetic proof. At a selected split prime $q_i \ge b^{10}$, take Z to be the inverse image of a stabilizer of a specified nonzero vector (or the dual vector for the conjugate factor). It has index $q_i^r - 1$ in K by the transitive action on nonzero vectors of $\mathbf{F}_{q_i}^r$; the two split factors are one unitary factor, not two independent choices. A conjugation-stable finite window containing the support of the level contains the whole selected pair, so the congruence group lies in Z. On this pair $\epsilon = \mathrm{vol}(R \setminus R^\times) \le 2/q_i \le b^{-9}$, and $\mu(K) \ge 2^{-n}$. The main term at K gives $B(K) \ge \mu(K)^2\mu(C)$. Therefore (24.13) implies

$$\frac{B(Z)\mu(K)}{B(K)\mu(Z)} \le \frac{\mu(Z)}{\mu(K)} + \epsilon^{t+1} \le \frac{1}{q_i^r - 1} + b^{-9(t+1)} \le b^{-6k}.$$

For the final bound, $r \ge k$, so the first summand is at most $2b^{-10k} \le b^{-6k}/2$. By the source's chain $10^{10}n^4 \le n^6$, $3n^6 + 12 \le n^7 \le k$, one has $9(t+1) \ge 6k + 45$, so the second is also at most $b^{-6k}/2$. On the other conjugation-stable place sets choose $Z = K$. The bottom form is multiplicative under the product of the local sets because ϕ is the product character and the two integrations separate; the ratio displayed has the same multiplicativity even if the complete Haar product is changed by a positive scalar (that scalar cancels). Thus the product factor for $|\Sigma|$ chosen pairs is $b^{-6k|\Sigma|}$. With the normalization m_Z and the upper group-mass bound (24.8), this gives the indicated $(1+b^{-6})b^{-6k|\Sigma|}$ bottom bound provided (24.13) holds at the selected pairs and also at the full window; the latter follows from the same error formula and $\epsilon \le 1$ (its unselected contribution is at most 2^{-n}, while $\mu(K) \le 1 + 2^{-n}$). The total factor is then $1 + 2 \cdot 2^{-n} \le 1 + b^{-6}$ by the source's $2^{-h} \le b^{-6}$.

24.8 Obligation-level status

- **PROVED:** the independent valuation content moments (24.1)–(24.2), finite hyperbolic-cone mass, and both finite common-volume bounds with the correct integral exponent. These give the analytic axioms of the actual `IsStdLevel.gauge` from the standard integral basis, trace scalars, and stated zeta-rank hypothesis. The corresponding complex analytic gauge was supplied in Section 23.9.
- **PROVED:** the finite phase well-definedness, covariance and support; compactness and uniform local constancy of the overlapping correlation product by block Gaussian elimination (Lemma 24.2); product Haar on the top stack with the prescribed orbit measure, and the lower stack normalization under the defining standard-level data. The top-kernel positivity for common last entries, continuity and entrywise integrability, and exact top-floor transform (24.5)–(24.7) now follow. Thus the additional support, Haar and top-entry integrability premises in Section 23.11 are discharged for the finite standard level. They do not establish the earlier rational-amplitude summability.
- **PROVED:** upper and lower integral-group volumes, exact unipotent modulus, the conjugation-average identity and normalized finite pairing norm (24.12). The contribution of the principal congruence subgroup to the bottom form is exactly the asserted main term. **PROVED (implication):** the fixed-vector subgroup savings given the precise error estimate (24.13).
- **OPEN:** the construction of the entire standard level from the marked lattice, its top integral section and zeta-rank bound for every actual finite window (these defining properties were used), the weighted tower estimates `formMajorant_sq_le`, `formModulus_sq_le_of_char`, `formModulus_sq_le_of_heads`, and the size/shell argument `lintegral_shells_le`. In particular (24.13) and the full frequency bound `enorm_floorForm_le` are not yet independently established; neither are the lower-floor descent integrability and numerical head-weight thresholds. The elementary Haar identities and the sliced Plancherel proof do not by themselves bound these weighted products.

Consequently the exact theorem `Filling.IsFillingLevel.exists_finite_sieveSystem` is still an open obligation of a fully conventional exposition, but its finite gauge, group/mass normalization, pairing-norm, and top-frequency pieces no longer require independent specialized estimates. Its remaining analytic gap has been narrowed to the lower weighted tower and shell estimates; replacing them by a qualitative Fourier decay reference would not suffice for the stated uniform bound.

Chapter 25

Construction and rigidity of the real discrete-series coefficient

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 28, 29 and the final audit in Chapter 57.

We now supply the proofs of the representation-theoretic premise left in Chapters 22 and 19. The identification of the explicit lowest-weight coefficient, its compact-type projector and its quantitative differentiated Schur inequality are logically separate from the global Bessel-pair estimates. In particular the fact that an analytic function on a group satisfies a Casimir eigenvalue equation does not in itself imply that it is a positive-definite matrix coefficient. We prove this last property by a compact averaging identity and a construction inside the right regular representation.

25.1 A compact averaging formula for the explicit integral

Use the notation of Section 22.2 and the same sign form $J = \operatorname{diag}(\epsilon_a)$. Let $G = U(J)$, $K = G \cap U(n)$, and $l = 2\lambda$ be the doubled Harish–Chandra parameter, regular with $l_a + n - 1$ even. The order of λ is the positive-root order ($e_a - e_b$ is positive if $\lambda_a > \lambda_b$). Write ρ_c and ρ_n for the compact and noncompact half sums of these positive roots, $\rho = \rho_c + \rho_n$, $\mu = \lambda + \rho$ and $\Lambda = \lambda + \rho - 2\rho_c$. Let $\psi = \psi_\lambda$ be exactly the normalized integral of principal minors from Chapter 22; more explicitly, with a_g the block diagonal part of g and probability measure on K,

$$Z = \int_K |m_\Lambda(u)|^2 \, du, \qquad \psi(g) = Z^{-1} \int_K \frac{m_\Lambda(a_g u)\overline{m_\Lambda(u)}}{m_{-\mu}(u^{-1} a_g^{-1} g u)} \, du.$$

Use the already established splitting independence (Lemma 22.1) and determinant bound (Lemma 22.5) for this integral; in particular $\psi(1) = 1$, $|\psi(g)| \le 1$, and its gap bound supplies L^1 and L^2 integrability when the gap exceeds $500n^3$. The latter follows at once from the norm-ball volume estimate in Lemma 19.1, which bounds the volume by a constant times R^{2n^2}.

Denote by m_w the minor product of the integral weight w. For an order o with $o_a < o_b$ if $\lambda_a > \lambda_b$, write B_o and N_o^- for the triangular Borel and opposite unitriangular subgroup of the indicated complex matrix group. On each sign block, for w dominant in this order, the minor formula is

$$m_w(A) = \det(A)^{w_{\min}} \prod_{i=1}^{d-1} \det(A_{\{1,\dots,i\}})^{w_i - w_{i+1}},$$

where d is the block rank; thus $m_w(Ab) = m_w(A)\chi_w(b)$ for $b \in B_o$ and $m_w(uA) = m_w(A)$ for $u \in N_o^-$. Here $\chi_w(b) = \prod b_{aa}^{w_a}$. The formula factors over the sign blocks for block diagonal matrices, even when some of their high weights interlace across blocks. It also holds for the whole matrix group when all the weights have the indicated dominance, on the appropriate domain of the minors. On K the product m_w is the coefficient of a unit highest vector of the compact highest-weight representation of weight w, as was verified in Theorem 22.4. In particular $m_w(k^{-1}) = \overline{m_w(k)}$. The weight Λ is compact dominant: for adjacent same-sign entries in the order within a sign block, the difference $\lambda_a - \lambda_b$ is at least one, $\rho_{c,a} - \rho_{c,b} = 1$, and the number of places between them in the full order only increases the corresponding difference of ρ. This proves dominance (and parity gives integrality).

Lemma 25.1 (Triangular circle extraction)**.** *Let w be compact dominant for o, $a, k \in K$, and $D(s) = \operatorname{diag}(s^{o_i})$ for $|s| = 1$. If $q(s)$ is holomorphic on a neighborhood of the closed unit disk, then*

$$\int_{|s|=1} m_w(kD(s)a)s^{-o\cdot w}q(s)\, d^\times s = m_w(k)m_w(a)q(0).$$

The circle has probability measure.

Proof. For a near 1, block Gaussian elimination gives $a = Lb$ with $L \in N_o^-$, $b \in B_o$ in the block diagonal group. The matrix $L(s) = D(s)LD(s)^{-1}$ is polynomial in s and is the identity at zero, since each off-diagonal lower entry has strictly positive exponent $o_i - o_j$. For $|s| = 1$ the minor covariance rewrites $m_w(kD(s)a)s^{-o\cdot w} = \chi_w(b)m_w(kL(s))$. The right side extends holomorphically across the disk: the only possible denominators in the minor formula on the block group are powers of its block determinants, and these are unchanged by $L(s)$. Cauchy's mean-value formula gives the assertion near 1, using $m_w(a) = \chi_w(b)$. Both sides are real-analytic functions of a on connected K. For the integral this follows locally by complex extension to the block general linear group, with the determinant uniformly bounded away from zero over the compact circle. The identity theorem extends the equality to all a. □

Proposition 25.2 (The compact rank-one identity)**.** *Let ν be a compact dominant integral weight. If $\nu = \Lambda$, or if there is an integral order o satisfying the above strict inequalities and $o \cdot (\nu - \Lambda) > 0$, then for $x, b \in G$*

$$\int_K \overline{m_\nu(k)}\psi(xkb)\, dk = \begin{cases} Z\psi(x)\psi(b), & \nu = \Lambda, \\ 0, & \nu \neq \Lambda. \end{cases}$$

Proof. First take b sufficiently close to 1. Uniformly for $k \in K$ the splitting $a = a_x k$ is valid in the defining integral for xkb: the other factor has positive definite Hermitian part, since this holds at $b = 1$ and K is compact. On substituting $u = kj$, $v = j$ in the double integral, $k = uv^{-1}$, and the integrand (before multiplication by Z^{-1}) becomes

$$\frac{m_\nu(vu^{-1})m_\Lambda(a_x u)\overline{m_\Lambda(v)}}{m_{-\mu}(Y_u B_v)}, \qquad Y_u = u^{-1}a_x^{-1}xu, \quad B_v = v^{-1}bv.$$

Here we have used the compact inverse identity for m_ν. To evaluate the integral over v, average its right translates $vD(s)$, which preserve Haar measure. Put $N = o \cdot (\nu - \Lambda)$, taking o to be any ordering of λ if $\nu = \Lambda$. For the full matrix group $-\mu$ is dominant for the reverse order $-o$. For b close enough to 1 there is a Gauss factorization $B_v = U_v C_v$ with U_v upper unitriangular and C_v lower triangular for o. All these factors are uniformly close to 1 for $v \in K$. The numerator factor under rotation is $m_\nu(vD(s)u^{-1})s^{-o\cdot\Lambda}\overline{m_\Lambda(v)}$. The denominator transforms to

$$m_{-\mu}(Y_u D(s)^{-1}U_v D(s))\chi_{-\mu}(C_v).$$

The matrix $D(s)^{-1}U_vD(s)$ is polynomial and equals 1 at zero. Its distance from 1 on the disk tends uniformly to zero as $b\to 1$. The compact family Y_u has uniformly positive definite Hermitian part, so all relevant minors in this transformed denominator are nonzero on a neighborhood of the closed disk. Apply Lemma 25.1 with

$$q(s)=s^N\left[m_{-\mu}(Y_uD(s)^{-1}U_vD(s))\chi_{-\mu}(C_v)\right]^{-1},\qquad a=u^{-1}.$$

If $N>0$, its value at zero is zero. If $\nu=\Lambda$, its value at zero is $[m_{-\mu}(Y_u)\,m_{-\mu}(B_v)]^{-1}$ by the triangular minor rule. The two remaining integrals are now separated:

$$Z^{-1}\int_K\frac{m_\Lambda(a_xu)\overline{m_\Lambda(u)}}{m_{-\mu}(Y_u)}\,du\int_K\frac{|m_\Lambda(v)|^2}{m_{-\mu}(v^{-1}bv)}\,dv=Z\psi(x)\psi(b).$$

Here the splitting for b is $a=1$. For fixed x the two sides are real-analytic in b on connected G: use the accretive splitting near any point, where uniform nonvanishing of the minors on the compact set permits integration of their complex-analytic germs. Connectedness follows from the polar decomposition $G=K\exp(\mathfrak{p})$ and connectedness of K. Thus the identity extends from b near the identity to all b. □

This is in particular the exact projector identity of `integral_conj_weightMinors_mul_dsCoeff` in `Lib/DiscreteSeries/FlenstedJensen/KTypeProjection.lean`. The alternative in that declaration is that $\nu-\Lambda$ does not lie in the intersection of half spaces $o\cdot\beta\le 0$ over all o compatible with the λ-order, equivalently the strict positivity alternative just used.

25.2 The matrix coefficient and its differentiated module

We record two further elementary consequences of the minor integral. In a polar decomposition $g=k_1zk_2^{-1}$, where $k_i\in K$ and $z=z^*>0$ in G, splitting independence gives

$$\psi_\lambda(g)=Z^{-1}\int_K\frac{m_\Lambda(k_1u)\overline{m_\Lambda(k_2u)}}{m_{-\mu}(u^{-1}zu)}\,du.$$

For such z, its determinant is one and $z^{-1}=JzJ$; the principal minors of z^{-1} and z are identical. Hence the denominator above is real positive and invariant under replacing z by its inverse. On K the negative weight has minor product conjugate to that of the positive weight (Jacobi's complementary-minor identity). Also $\Lambda(-\lambda)=-\Lambda(\lambda)$. The polar formula consequently proves

$$\psi_\lambda(g^{-1})=\overline{\psi_\lambda(g)}=\psi_{-\lambda}(g).\tag{25.1}$$

Proposition 25.3 (Construction inside the right regular representation)**.** *Assume the noncompact gap is at least $500n^3+A$, where $A\ge 0$. Let $d=(\int_G|\psi|^2)^{-1}$ for the specified Haar measure on G. The function ψ is the matrix coefficient of a unit smooth vector v in an irreducible square-integrable unitary representation π_ψ with formal degree d. The compact module generated by v is the compact representation of highest weight Λ, and its right derivatives are in $L^1(G,(1+\|g\|)^A dg)$.*

Proof. For an integrable w invariant under the left compact projector, in the sense

$$\int_K\overline{m_\Lambda(k)}w(k^{-1}g)\,dk=Zw(g),$$

Fubini and Proposition 25.2 give

$$\int_G w(y)\psi(xya)\,dy=\psi(x)\int_G w(y)\psi(ya)\,dy.\tag{25.2}$$

For $w = \overline{\psi}$ the projector condition follows from (25.1) and the same proposition: the left side of the projector condition equals $\int_K \overline{m_\Lambda(k)}\psi(g^{-1}k)\,dk$. Absolute integrability was noted above; all changes of order are bounded by $\|w\|_1$, since the normalized compact coefficients and ψ have absolute value at most one. Taking $a = 1$ in (25.2), then changing the variable by inversion and conjugating, gives the full product identity

$$d\int_G \psi(xya)\overline{\psi(y)}\,dy = \psi(x)\psi(a). \tag{25.3}$$

Here reductive G is unimodular. The integral of $|\psi|^2$ is positive because $\psi(1) = 1$ and ψ is continuous.

In $L^2(G)$ let H_ψ be the closed span of the right translates of $v = \sqrt{d}\,\psi$. Right translation $\pi_\psi(g)$ is unitary, and (25.3) says exactly $\langle v, \pi_\psi(g)v\rangle = \psi(g)$, with the inner product linear in the second entry. The compact operator

$$T_\Lambda = Z^{-1}\int_K \overline{m_\Lambda(k)}\pi_\psi(k)\,dk$$

is the highest-vector projection on every compact summand of type Λ, and is zero on the other types, by compact Schur orthogonality (Z^{-1} is the compact dimension). Evaluating it on right translates using Proposition 25.2 gives $T_\Lambda\pi_\psi(b)v = \psi(b)v$. Its range is consequently exactly $\mathbb{C}v$. If a closed G-invariant subspace has orthogonal projection Q, it commutes with T_Λ. Thus Qv is a multiple of v; since Q is a projection, Qv is either 0 or v. Cyclicity then gives either the zero or the whole representation. This proves irreducibility. Square-integrability of the coefficient implies that it is a discrete series, and ordinary Schur orthogonality identifies its formal degree with the chosen d.

Smoothness and the weighted L^1 assertion can be checked without differentiating a pointwise decay bound. Choose a nonnegative smooth compactly supported K-left-invariant function χ nonzero near 1. The analogue $w = \chi\overline{\psi}$ of the preceding w still satisfies the left-projector identity. Hence, dividing by $\int_G \chi|\psi|^2 > 0$, equation (25.2) expresses $\psi(x)$ as an integral $\int \beta(y)\psi(xy)\,dy$ with smooth compactly supported β. Right translation of this expression replaces $\beta(y)$ by $\beta(h^{-1}y)$ for translation by h. This is a smooth map to the compactly supported functions whose supports locally stay in a fixed compact set, hence also to L^1 and to the integrated operators in $L^2(G)$. Every iterated derivative of the coefficient is therefore a similar integral with a smooth compactly supported test function. On a compact support, $\|xy\| \geq \|x\|/\|y^{-1}\|$; the determinant decay bound gives $C_\beta(2/\|x\|)^c$ for every integer c not exceeding the gap. This implies the asserted weighted integrability and shows that each coefficient derivative agrees with the differentiated right regular representation, not just with a formal derivative. □

The only classical fact used for the last identification of degree is ordinary discrete-series Schur orthogonality; the source `dsCoeff_schur_orthogonality` has already been proved by (25.3). In particular no form of the discrete-series classification has been used in the construction of H_ψ. In the next subsection a stronger rigidity assertion will identify its parameter, with all signs fixed.

Lemma 25.4 (Infinitesimal type and Casimir)**.** *For the complexified right derivatives E_{ab} on ψ, $E_{aa}\psi = \Lambda_a\psi$ and $E_{ab}\psi = 0$ for $\epsilon_a = \epsilon_b$ and $\lambda_a > \lambda_b$. The full Casimir $\Omega = \sum_{a,b} E_{ab}E_{ba}$ satisfies*

$$\Omega\psi = (\|\lambda\|^2 - \|\rho\|^2)\psi.$$

Proof. The accretive formula is complex analytic near any point of G: for a fixed valid splitting the denominator is uniformly nonzero near that point over the compact K. The left opposite

compact triangular transformation $\psi_\lambda(bg) = \chi_\Lambda(b)\psi_\lambda(g)$ holds locally for b near the identity in that Borel, by replacing the splitting a with ba. Equation (25.1) gives the complex-analytic inverse identity locally on such neighborhoods: it holds on the real form G, which is maximally totally real in the complex matrix group (use local exponential coordinates). Replacing g by gb and λ by $-\lambda$ in the inverse identity therefore gives right covariance for the compact upper triangular Borel, with the character χ_Λ. Differentiation gives the claimed compact highest-vector identities.

For the full Casimir, fix a valid block diagonal splitting a near the point at which we differentiate. For fixed $u \in K$ the scalar function of the variable g in the denominator inverse has right covariance under $g \mapsto gubu^{-1}$, where b is triangular for the reverse of the λ-order. Indeed this appends b to $u^{-1}a^{-1}gu$, so its inverse multiplier is $\chi_\mu(b)$ by the full-group minor covariance for $-\mu$. It is consequently a highest-vector germ of weight μ with the reverse order and the Lie algebra conjugated by u. For any highest-vector germ of weight w and root order with half sum η, the elementary formula $[E_{ab}, E_{ba}] = E_{aa} - E_{bb}$ rewrites the Casimir as its diagonal squares plus the sum of positive-negative terms, and gives the eigenvalue $\sum_a w_a(w_a + 2\eta_a) = \|w + \eta\|^2 - \|\eta\|^2$. The Casimir is invariant under conjugation because the matrix units and their transposes are dual for the trace form. Here $\eta = -\rho$, so the eigenvalue of the integrand is $\|\mu - \rho\|^2 - \|\rho\|^2 = \|\lambda\|^2 - \|\rho\|^2$. We may differentiate up to order two under the integral: all derivatives of the fixed-splitting rational integrands are locally uniformly bounded on the compact K because their denominators stay nonzero there. This proves the Casimir equation. □

These are the declarations `isKHighest_dsGenerator` and `casimir_dsCoeffSmooth` in the source's `Lib/DiscreteSeries/CuspKernel/SchurOrthogonality.lean` and `Lib/DiscreteSeries/FlenstedJensen/CasimirOnPsi.lean`. The smooth coefficient module is now an actual module inside H_ψ. For it, differentiating the unitary invariance in the real Lie algebra and then complexifying gives

$$E_{ab}^\dagger = \epsilon_a \epsilon_b E_{ba}. \tag{25.4}$$

For example apply the real skew-adjoint assertion to $E_{ab} - \epsilon_a\epsilon_b E_{ba}$ and $i(E_{ab} + \epsilon_a\epsilon_b E_{ba})$; these are in $\mathfrak{u}(J)$. This also fixes the convention for every adjoint used below.

25.3 A quantitative compact Casimir argument

We give the local algebra argument explicitly, because the source's rigidity theorem is stronger than the uniqueness of the minimal K-type in an irreducible representation. Its hypotheses allow an arbitrary unitary representation and a vector on which the Casimir is merely bounded below by the desired value.

Let M be an inner product $\mathfrak{gl}_n(\mathbb{C})$-module with (25.4), and suppose $\langle z, \Omega z\rangle \geq c_\lambda \|z\|^2$ in real part for all $z \in M$, where $c_\lambda = \|\lambda\|^2 - \|\rho\|^2$. Write $C = \{(a, b) : \epsilon_a = \epsilon_b\}$, $\Omega_K = \sum_{(a,b)\in C} E_{ab}E_{ba}$ and $c_K(\nu) = \sum_a \nu_a(\nu_a + 2\rho_{c,a})$. Let x be a compact highest vector of weight ν, and put $M_0 = U(\mathfrak{k}_\mathbb{C})x$, $M_{s+1} = M_s + \sum_{a,b} E_{ab}M_s$.

Lemma 25.5 (Finite filtration and compact spectra)**.** *Each M_s is finite dimensional and compact-stable. On M_0 the compact Casimir is $c_K(\nu)$. Every compact highest weight ω occurring in M_s has the form*

$$\omega = \nu + \sum_{j=1}^{t} (e_{a_j} - e_{b_j}), \qquad t \leq s.$$

Proof. The highest-vector commutation formula gives the compact Casimir value on x; centrality in the compact Lie algebra extends it to M_0. Compact adjointness implies that on M_0 $\langle z, \Omega_K z\rangle = \sum_{(a,b)\in C} \|E_{ba} z\|^2$. PBW expresses M_0 by compact lowering words on x. For a nonzero word of weight ξ the last equality bounds each $|\xi_a|$ by $\sqrt{c_K(\nu)}$ using the diagonal terms. Each compact lowering strictly reduces the scalar product of this weight with the regular ordering vector l (by at least one). Since all the weights are thus bounded, nonzero lowering words have a uniform finite length. This proves finiteness of M_0 and hence finiteness of each M_s by induction. Compact stability follows because commuting a compact matrix unit past any matrix unit gives a linear combination of matrix units.

More precisely M_s is spanned by compact lowering words applied to words of length at most s in arbitrary matrix units on x. This follows inductively by commuting each additional matrix unit across the compact lowering word, using the commutator as a replacement of one matrix unit; for $s = 0$ it is PBW. A compact highest vector $z \in M_s$ is orthogonal to every vector obtained with a nonempty leftmost compact lowering word by adjointness. If $z \neq 0$, at least one of the remaining words of length at most s has nonzero pairing with z. Diagonal matrix units are self-adjoint and act on that word with the weight obtained by adding its roots to ν, by their commutator identities. Thus this is also the weight ω of z. Finite-dimensional compact modules with this adjointness decompose into irreducible highest-weight modules: orthogonal complements to invariant subspaces are invariant, and simultaneous diagonalization of the diagonal operators, then maximization against l, yields highest vectors. This proves the assertion for all compact highest weights. □

Proposition 25.6 (Explicit derivative growth)**.** *If $|\nu_a| \le L_0$ for all a and $y \in M_s$, then for every matrix X,*

$$E(X)y \in M_{s+1}, \qquad \|E(X)y\| \le (n+1)^4(L_0 + 1 + s)\|X\|_{\mathrm{op}}\|y\|.$$

Proof. On each compact irreducible summand of M_s the compact Casimir is its highest weight value, bounded above by $n(L_0 + s)(L_0 + s + n)$ by Lemma 25.5 and $|2\rho_{c,a}| \le n$. From (25.4) the sum of squares over all matrix units is

$$\sum_{a,b} \|E_{ab}y\|^2 = 2\langle y, \Omega_K y\rangle - \mathrm{Re}\langle y, \Omega y\rangle \le [2n(L_0 + s)(L_0 + s + n) + n^3]\|y\|^2,$$

since $c_\lambda \ge -\|\rho\|^2 \ge -n^3$. Use $E(X) = \sum X_{ab}E_{ab}$ and $|X_{ab}| \le \|X\|_{\mathrm{op}}$ with Cauchy–Schwarz over n^2 terms. Setting $t = L_0 + s + 1$ gives a factor at most $n\sqrt{2n(n+1) + n^3}\, t \le (n+1)^4 t$. Membership in M_{s+1} follows at once from its definition. □

This proves the constant, not just the order in the rank, in `HCParam.IsKHighest.norm_apply_le_of_mem_filtration` of `Lib/DiscreteSeries/Rigidity/CasimirGrowth.lean`. We next prove the distinct spectral separation statement needed for rigidity.

Lemma 25.7 (Compact spectral gap at the first step)**.** *Assume now $\nu = \Lambda$ and the noncompact gaps $|\lambda_a - \lambda_b|$ are at least $500n^3 + A$. On $M_1 \ominus M_0$ the compact Casimir is at least $c_0 + 4n + 2$, where $c_0 = c_K(\Lambda)$.*

Proof. If z is any compact highest vector of weight $\Lambda + \delta$, reorder the noncompact terms of Ω with respect to the reverse of the λ order. Their expectation is at most $-2\langle \rho_n, \Lambda + \delta\rangle\|z\|^2$: the remaining offcompact pairs have nonpositive expectations by (25.4). Using the Casimir lower bound and the compact highest-vector identity gives

$$\|\lambda\|^2 - \|\rho\|^2 \le \|\Lambda + \delta + \rho_c - \rho_n\|^2 - \|\rho_c - \rho_n\|^2.$$

Since $\Lambda + \rho_c - \rho_n = \lambda$, and $4|\langle \rho_c, \rho_n \rangle| \le n^3$, this implies

$$0 \le n^3 + \|\delta\|^2 + 2\langle \lambda, \delta \rangle. \tag{25.5}$$

If z is a highest vector in $M_1 \ominus M_0$, the proof of Lemma 25.5 forces its weight to pair nontrivially with some $E_{ab}x$ with opposite signs (the length-zero word and compact ones are in M_0). Thus $\delta = e_a - e_b$ for $\epsilon_a \neq \epsilon_b$. Inequality (25.5) and the huge noncompact gap imply $\lambda_a > \lambda_b$. Consequently

$$c_K(\Lambda + \delta) - c_0 = 2(\lambda_a - \lambda_b + \rho_{n,a} - \rho_{n,b}) + 2 \ge 2(500n^3 - n) + 2 \ge 4n + 2.$$

Opposite signs require $n \ge 2$; if there are none, $M_1 = M_0$ and the assertion is vacuous. □

Proposition 25.8 (Propagation of compact spectral separation)**.** *Under the hypotheses of Lemma 25.7, if $\Omega_K a = \theta a$ with real $\theta \le c_0 + s$ and $a \perp M_s$, then $a \perp M_t$ for every t.*

Proof. Let P be the orthogonal projection onto the finite-dimensional M_0, commuting with all compact units by their adjointness. Use A, B for arbitrary matrix units and C (in a sum) for compact matrix units; the same letter denotes the corresponding operator on M. For a word $(A_1, \dots, A_k)$ put $J_W = PA_1 \cdots A_k$, and define the nonnegative quadratic forms

$$\mathcal{L}_k(a) = \sum_{|W|=k} \|J_W a\|^2, \qquad \mathcal{R}_k(a) = \sum_{|W|=k,C} \|[C, J_W]a\|^2.$$

For $a \perp M_{k-1}$, $\mathcal{L}_k(a) = 0$ is equivalent to $a \perp M_k$, by the spanning definition of M_k and (25.4). For each k, the operator $D_k = \sum_{|W|=k} J_W^\dagger J_W$ commutes with compact units. In fact $D_0 = P$, and $D_{k+1} = \sum_{a,b} \epsilon_a \epsilon_b E_{ba} D_k E_{ab}$. If $\epsilon_c = \epsilon_d$ and D_k commutes with E_{cd}, its commutator with this sum has four terms: $\sum_a \epsilon_a \epsilon_d E_{ca} D_k E_{ad}$, $-\sum_b \epsilon_c \epsilon_b E_{bd} D_k E_{cb}$, $\sum_b \epsilon_d \epsilon_b E_{bd} D_k E_{cb}$, and $-\sum_a \epsilon_a \epsilon_c E_{ca} D_k E_{ad}$. They cancel exactly, so no integration of the abstract Lie algebra action is needed in this verification. Thus

$$\sum_{W,C} \|J_W C a\|^2 = \mathrm{Re}\langle a, D_k \Omega_K a \rangle. \tag{25.6}$$

We first prove, for $a \perp M_0$, the crossed inequality

$$2\sum_A \|PAa\|^2 \le 2\,\mathrm{Re} \sum_{A,C} \langle CPAa,\ P[A,C]a \rangle. \tag{25.7}$$

The weighted compact energy $\sum_{A,C} \|PACa\|^2$ depends only on the projection of a to M_1, since PA annihilates $M_1^\perp$ and compact units preserve this complement. The first-step gap and D_1-commutation bound it below by $(c_0 + 4n + 2)\mathcal{L}_1(a)$. On the other hand $PACa = CPAa + P[A,C]a$. Expanding squares, the first squared term sums to $c_0 \mathcal{L}_1(a)$. For the second, $[E_{ab}, E_{cd}] = \delta_{bc} E_{ad} - \delta_{da} E_{cb}$; the squared norm of the sum of two vectors is at most twice the sum of their squared norms. Each resulting operator is counted at most n times in each of the two summands, so the total second squared term is at most $4n\mathcal{L}_1(a)$. Subtract these two terms to obtain (25.7).

For a fixed word W of length j, the same inequality holds with P replaced throughout by J_W, provided $a \perp M_j$. Indeed apply (25.7) to $b = A_1 \cdots A_j a$; this vector is orthogonal to M_0 by adjointness. Inside P one may move an extra arbitrary matrix unit past these j units at the expense of commutators of word length at most j; all such terms vanish on a after applying P. The same argument moves $[A, C]$, a linear combination of single units, past the word. Therefore $PAb = J_W Aa$ and $P[A,C]b = J_W [A,C]a$, giving the assertion (the exterior action of C requires no change).

For $a \perp M_{k-1}$ we claim

$$(c_0 + 2k)\mathcal{L}_k(a) + \mathcal{R}_k(a) \le \sum_{|W|=k,C} \|J_W C a\|^2. \tag{25.8}$$

The assertion for $k = 1$ follows from the first-step gap and the $4n$ commutator bound just used. To pass from $k-1$ to k, each $Aa \perp M_{k-2}$, so sum the $(k-1)$-inequality applied to Aa over all A. For $J = J_{W'}$ with $|W'| = k-1$, appending A changes the difference of the energy and commutator squares by

$$\begin{aligned} &\|JACa\|^2 - \|[C, JA]a\|^2 \\ &\quad - \|JCAa\|^2 + \|[C, J]Aa\|^2 \\ &= 2\,\mathrm{Re}\langle CJAa,\ J[A, C]a\rangle. \end{aligned}$$

This is the elementary expansion $\|u+b\|^2 - \|v-b\|^2 - \|u\|^2 + \|v\|^2 = 2\,\mathrm{Re}\langle u+v, b\rangle$. The crossed inequality for J raises the lower bound by $2\sum_A \|JAa\|^2$, proving (25.8).

Now if a is a compact Casimir eigenvector as in the statement and $a \perp M_{k-1}$ for $k > s$, equations (25.6), (25.8) give $(c_0 + 2k)\mathcal{L}_k(a) \le \theta\mathcal{L}_k(a)$. Because $\theta \le c_0 + s < c_0 + 2k$, it follows that $\mathcal{L}_k(a) = 0$. Induction, starting with $k = s+1$, proves the result. □

The argument proves exactly the orthogonality propagation of `mem_orthogonal_filtration_of_casimirK`; in particular one does not infer this propagation from the first-step gap alone.

25.4 Analyticity and representation-independent rigidity

Lemma 25.9 (Quantitative analytic continuation of coefficients)**.** *Suppose $c_g : M \to \mathbb{C}$ is linear with $|c_g(y)| \le D\|y\|$ uniformly in g for some D, and $t \mapsto c_{g\exp(tY)}(y)$ differentiates to $c_{g\exp(tY)}(E(Y)y)$ for every $Y \in \mathfrak{u}(J)$. If c_1 annihilates every M_s then $c_g(x) = 0$ for every $g \in G$.*

Proof. For $y \in M_s$, Proposition 25.6 and iteration bound its mth derivative on any such real line by

$$D\|y\|\,\big((n+1)^4\|Y\|\big)^m (L_0 + s + 1)(L_0 + s + 2)\cdots(L_0 + s + m).$$

Each derivative is continuous, because it is again a differentiable scalar coefficient. Taylor's theorem with remainder, divided by $m!$, shows that the function is real analytic on the line: the remaining factor $\binom{L_0+s+m}{m}$ has polynomial growth in m when L_0, s are fixed integers. The radius can be chosen uniformly positive for all the y of the cyclic module (depending only on Y). If all the coefficients vanish at g on all M_s, their derivatives of every order vanish there, hence they vanish along the entire exponential line by the identity theorem. Every $g \in G$ is a product of two real Lie exponentials by the polar decomposition and the existence of a skew-Hermitian logarithm in each compact unitary block. Starting at the identity proves the result. □

Theorem 25.10 (The normalized highest-vector coefficient)**.** *Let a strongly continuous representation of G have an invariant positive definite inner product (a continuous form on its Banach space suffices), and a complexified derived action on a module mapping into that representation. Let x denote the image of a compact highest vector of weight Λ. Suppose the full Casimir acts on x by a real scalar $c \ge c_\lambda$. Then, for gaps at least $500n^3 + A$,*

$$\langle x, \pi(g)x\rangle = \|x\|^2 \psi_\lambda(g) \qquad (g \in G). \tag{25.9}$$

If $x \neq 0$ then necessarily $c = c_\lambda$.

Proof. The zero case is immediate. In general the derived action passes to the image of the module: differentiating invariance gives (25.4) there, so if an element maps to zero, the norm-square of its image under any matrix unit is zero by moving the adjoint to the second slot. Pass to the inner-product completion if necessary; the original strong continuity for the Banach norm and continuity of the form imply continuity on its dense subspace, and unitarity extends it to the completion. On the cyclic Lie algebra module generated by x the full Casimir is c by centrality. On the module generated by the explicit unit vector v of Proposition 25.3 it is c_λ by Lemma 25.4. In their orthogonal direct sum use the cyclic generator $y = (x, v)$ and test vector $a = (x, -\|x\|^2 v)$. This direct sum satisfies the Casimir lower bound. Both compact highest vectors generate the same finite-dimensional irreducible compact module (highest weight Λ), with their indicated norms, so pairing any compact Lie algebra word on y against a gives zero. Indeed the compact cyclic modules are irreducible by finite-dimensional highest-weight theory: each summand generated by a nonzero projection of the highest vector has highest weight Λ, and the cyclic span is its diagonal copy. The unique highest-weight isomorphism between the two normalized cyclic modules preserves the inner products. Therefore $a \perp M_0(y)$. Also $\Omega_K a = c_K(\Lambda) a$. Apply Proposition 25.8 with $s = 0$ to conclude that $a \perp M_t(y)$ for every t.

The scalar coefficient map on the direct sum defined by

$$c_g(u, w) = \langle x, \pi(g)u\rangle - \|x\|^2 \langle v, \pi_\psi(g)w\rangle$$

is uniformly bounded by $(\|x\| + \|x\|^2)\|(u, w)\|$, satisfies the differentiated-action hypothesis and at 1 is exactly pairing against a. Lemma 25.9 proves $c_g(y) = 0$, which is (25.9). Differentiating this identity by the full Casimir and taking $g = 1$ gives $c\|x\|^2 = c_\lambda \|x\|^2$. This yields the final assertion. □

These prove the assertions `coeffMap_eq_zero`, `coeff_eq_dsCoeff_of_isKHighest`, and `eq_casimirValue_of_ge`, respectively in `Lib/DiscreteSeries/Rigidity/AnalyticVectors.lean` and `Lib/DiscreteSeries/Rigidity/KernelFixes.lean`. The proof did not assume irreducibility of the original representation or a priori equality of its Casimir eigenvalue. If $e(g) = d\,\overline{\psi(g)}$, then it is the minimal idempotent integrated in the source. More generally, if f is integrable against the specified operator-growth bound and $f * e = e$ in the usual group algebra convention, then $\pi(f)x = x$. To see this, the cyclic representation generated by x is the same as π_ψ: its normalized coefficient determines the inner products between all pairs of translates by (25.9), hence the cyclic isometry. In this irreducible representation $\pi_\psi(e)$ is the orthogonal projection onto v by Schur orthogonality, and $\pi_\psi(f * e) = \pi_\psi(f)\pi_\psi(e)$; applying to v proves the claim. This also establishes `integratedRep_fix_of_isKHighest`.

Identification of the parameter and the orbital formula. For this last step use one ordinary classical theorem of real representation theory: for a connected semisimple group with finite center and compact Cartan the discrete series attached to a regular parameter λ with $\lambda + \rho$ an integral character has infinitesimal character λ and contains the compact highest weight $\lambda + \rho - 2\rho_c$. A precise reference is A. W. Knapp, *Representation Theory of Semisimple Groups*, Princeton University Press (1986), Theorems 9.20 and 12.21, for existence, infinitesimal character and the lowest K-type. The same formula and its normalization are explicitly recorded in F. Murnaghan's *Representations of Reductive Groups* notes, Theorem 7.2 (p. 85, `https://www.math.utoronto.ca/fiona/courses/mat1196/rnotes.pdf`). To apply this semisimple theorem to $U(p, q)$, use the finite central cover $SU(p, q) \times U(1) \to U(p, q)$. Restrict the parameter to the traceless compact Cartan; its ρ-shift is integral since it is the restriction of the integral $\lambda + \rho$ of $U(p, q)$. The semisimple representation has central character on $\zeta I \in SU(p, q)$ given by the lowest weight, namely $\zeta^{\sum_a \Lambda_a} = \zeta^{\sum_a \lambda_a}$. The integer $\sum_a \lambda_a$

is integral by the parity condition on all l_a (if n is even a sum of n half-integers is integral). Tensor with $z \mapsto z^{\sum_a \lambda_a}$ on the compact $U(1)$ factor; the resulting representation is trivial on the finite kernel of the cover and descends. It stays irreducible and square-integrable since the extra factor is compact and the kernel finite. The full matrix Casimir is the trace-free Casimir plus $n^{-1}(\sum E_{aa})^2$; its eigenvalue is therefore precisely $\|\lambda\|^2 - \|\rho\|^2$. For a definite signature this statement is simply the compact highest-weight construction and Casimir formula. The distinguished unit highest vector of this classical discrete series thus satisfies all the hypotheses of Theorem 25.10. Its matrix coefficient equals the explicit ψ_λ pointwise, and hence its formal degree is the source's reciprocal integral. Therefore the representation premise of the orbital comparison Theorem 19.2 is now discharged for the required large-gap coefficients. Its stable Weyl sum and normalization follow as there from the ordinary Harish–Chandra character theorem, not from a formal transfer assertion.

25.5 Differentiated Schur estimate for the Bessel space

The last part of the previous chapter's B18 interface is an inequality on the noncompact Bessel quotient, not merely on the group. We now derive it with the source's normalization. Let U denote the Bessel unipotent in the source, with its specified character of modulus one; let F be a bounded measurable vector-valued Bessel function of compact covariance $F(hg) = \tau(h)F(g)$, with τ unitary, and of finite $L^1(U\backslash G)$ norm. The quotient carries its right invariant measure. The Hilbert space of measurable functions with this left unipotent covariance, square-integrable on the quotient, has unitary right translations $R(g)$. Boundedness and the assumed L^1 norm imply finite L^2 norm for F. Define the convolution with an integrable scalar z by

$$(F * z)(y) = (T_z F)(y) = \int_G z(g^{-1})F(yg)\, dg.$$

This is the source convention of `Lib/DiscreteSeries/CuspKernel/BesselSchurBound.lean`, not the usual integrated representation convention without the inverse. In this convention the reproducing kernel is $e = d\psi$ (not $d\overline{\psi}$).

Proposition 25.11 (Exact coefficient-subspace norm)**.** *If $F * e = F$ then, for any vector w in the smooth cyclic derivative module of the unit vector v constructed above, put $z_w(g) = d\langle v, \pi_\psi(g)w\rangle$. Then $F_w = F * z_w$ is bounded continuous, has the same left Bessel covariance as F, and*

$$\|F_w\|^2_{L^2(U\backslash G)} = \|w\|^2 \|F\|^2_{L^2(U\backslash G)}.$$

For every integrable scalar ϕ one further has

$$\int_{U\backslash G} \left\| \int_G \phi(y) F_w(gy)\, dy \right\|^2 d\dot{g} \le d^{-1}\|w\|^2 \|F\|^2_{L^2(U\backslash G)} \int_G |\phi|^2.$$

Proof. The integrability of z_w follows from the compact-smearing proof in Proposition 25.3; the same applies to all translates and finite sums of translates and derivatives. With convolution product $(a * b)(g) = \int b(t^{-1})a(gt)\, dt$ and involution $z^*(g) = \overline{z(g^{-1})}$ the operators satisfy $T_a T_b = T_{b*a}$ and $T_a^\dagger = T_{a^*}$ by unimodularity and absolute convergence in the unitary representation. Schur orthogonality in π_ψ gives $z_w * z_w^* = \|w\|^2 e$, since its integral is

$$d^2 \int_G \langle v, \pi_\psi(gt)w\rangle \overline{\langle v, \pi_\psi(t)w\rangle}\, dt = d\|w\|^2 \langle v, \pi_\psi(g)v\rangle.$$

Consequently $\|T_{z_w} F\|^2 = \langle F, T_{z_w * z_w^*} F\rangle = \|w\|^2 \|F\|^2$. Convolution of a bounded function with an L^1 function is bounded continuous here: translation of the integrable kernel is continuous

in its L^1 norm, and pointwise evaluation of the convolution has a uniform bound by $\sup\|F\|$ times that norm. Left covariance follows by applying the stipulated character or unitary matrix inside the integral.

The map $w \mapsto F_w$ is linear and extends, by the norm identity, to a scalar multiple of an isometric intertwiner from H_ψ to the quotient Hilbert space. Indeed $R(h)T_{z_w}F = T_{z_{\pi_\psi(h)w}}F$ by the substitution $g' = hg$ in the convolution. First check this on the span of translates and derivatives, which are integrable as noted above and dense in the cyclic representation. The operator $\int_G \phi(y)\pi_\psi(y)\,dy$ has norm at most $d^{-1/2}\|\phi\|_2$: pair against unit vectors and use Cauchy–Schwarz with the ordinary Schur orthogonality $\int |\langle u, \pi_\psi(y)w\rangle|^2 = d^{-1}\|u\|^2\|w\|^2$. When $\phi \notin L^2$ the desired inequality is vacuous. Use the intertwiner and square the operator bound to obtain the displayed quotient inequality. □

For a word w of s matrix units acting on the original unit vector, let v_w be the resulting derivative vector. The actual source frame identifies G with $U(J)$ by conjugation with an ordinary unitary matrix in its stated coordinate realization, so the conjugated elementary matrix units still have operator norm one. For $|l_a| \le L$, $|\Lambda_a| \le L + n$; Proposition 25.6 gives

$$\|v_w\| \le \big((n+1)^4(L+n+1+s)\big)^s.$$

Set $V_w = F * z_{v_w}$. Its derivative along a curve γ with derivative $\gamma(t)X$ is $\sum X_{ab}V_{(a,b)w}(\gamma(t))$. For rigor differentiate the compact-smearing expression for z_{v_w} under right translation; the smearing test functions and their derivatives are compactly supported, so their translation is differentiable in L^1, and convolution against bounded F preserves this differentiability in the uniform norm. Since $V_\varnothing = F$, this constructs the actual continuous derivative words required by the source. For the shape with $\mathfrak{n} = \varrho + 1$ and rank n, take $s \le 2\mathfrak{n}^2$ and $B \ge (n+1)^4(L+n+1+2\mathfrak{n}^2)$. Proposition 25.11 gives SchurLe$(V_w, B^{2s}d^{-1}\|F\|_2^2)$ with the exact exponent and constants of `IsBesselFunction.hasSchurBound_of_convolve_eq` in `New/RealPlace/BesselCut.lean` and the definition `HasSchurBound` in `New/Filling/BesselSystem.lean`. Thus the differentiated Schur interface no longer needs a separate representation-theoretic hypothesis. The prerequisites are exactly those of that declaration (bounded Bessel function of finite quotient L^1 norm reproduced by the normalized kernel).

The coefficient Lipschitz constant. The same norm estimate also proves `norm_dsCoeff_sub_le` in `Lib/DiscreteSeries/FlenstedJensen/TailBounds.lean`. If $\|g-1\| \ge 1/2$, then $|\psi(xg)-\psi(x)| \le 2 \le 4\|g-1\|$. Otherwise the power-series logarithm $X = \log g$ belongs to $\mathfrak{u}(J)$: use $g^{-1} = Jg^*J$ and the analytic logarithm identity $\log(g^{-1}) = -\log(g)$, initially on the power-series domain and then on the common principal branch. Moreover $\|X\| \le 2\|g-1\|$. Integrating the matrix coefficient derivative along $x\exp(tX)$, for $t \in [0,1]$, gives

$$|\psi(xg)-\psi(x)| \le 2(n+1)^4(L+n+1)\|g-1\| \le 100(n+L+1)^5\|g-1\|.$$

This implies the source's lower bound $|\psi(g)| \ge 1/2$ whenever $\|g-1\| \le (200(n+L+1)^5)^{-1}$, with no extra assumption. The determinant decay proved earlier and its Gaussian approximation are therefore available for the realized lowest vector with all constants intact.

Lemma 25.12 (The normalized tail integral)**.** *Let $\beta = n + L + 1$ and $c > 8n^2$. With the actual source Haar normalization and formal degree, one has*

$$d\int_G (2/\|g\|)^c\,dg \le 8(2000\beta^6)^{2n^2}2^c.$$

Proof. Take $\delta = (200\beta^5)^{-1}$. The ball $\|g-1\| \le \delta$ has Haar measure at most $4/d$ by the half-coefficient bound just proved. The real normed space of complex matrices has dimension

$2n^2$. In its ball of radius $R \geq 1$, a maximal δ/R-separated set in the group has cardinal at most $(1 + 2R^2/\delta)^{2n^2}$, by packing disjoint ambient balls of half the separation radius. It is a net in the group ball. If g is within δ/R of a center h, then $\|gh^{-1} - 1\| \leq \delta$, because $\|h^{-1}\| = \|h\| \leq R$. Right invariant Haar therefore bounds the group ball by

$$\mathrm{vol}(\{\|g\| \leq R\}) \leq 4d^{-1}(2000\beta^6 R^2)^{2n^2}.$$

The same argument applies to any group mapping homomorphically to G, using a right invariant pullback ball measure and centers in the image. No covering estimate for an unspecified Riemannian metric has been used. Since every norm is at least one, the layer integral is $\int_G \|g\|^{-c}\, dg = c \int_1^\infty R^{-c-1} \mathrm{vol}(\{\|g\| \leq R\})\, dR$. The ratio $c/(c - 4n^2)$ is less than two, proving the claim. □

This proves `measure_ball_le_integral_dsCoeff` and `integral_tail_le` of the same source file.

25.6 The Gram-section geometry and its exact Jacobian

We prove the local open-cell coordinate and Haar calculations separately, since they concern a large noncompact group even when the Bessel function used on it has a definite target type. Use the source shape with ϱ hyperbolic planes, a positive space X_0 of dimension $k \geq 1$, and a negative line spanned by z. Put $m = \varrho + 1$, $r = 2m + k - 1$. In the chosen Euclidean-orthonormal Witt frame let F have columns $(e_1, \ldots, e_\varrho, z)$ and F' have columns $(e_1^*, \ldots, e_\varrho^*, x)$ for the chosen distinguished unit vector $x \in X_0$. Denote the orthogonal complement in X_0 of x by M. The matrix of the Hermitian form in coordinates F, F', M is

$$J_0 = \begin{pmatrix} -L & P & 0 \\ P & L & 0 \\ 0 & 0 & I_M \end{pmatrix}, \qquad L = E_{mm}, \quad P = 1 - L.$$

All norms in this calculation are ordinary Euclidean operator norms in these ordered coordinates. Set $Z(g) = F^* J_0 g F$. For a square complex m-matrix Z, define

$$\mathfrak{g}(Z) = |\det Z|^2 - |\det Z_{\{1,\ldots,m-1\}}|^2.$$

Its positive locus is exactly $\Omega = \{Z : \det Z \neq 0,\ |(Z^{-1})_{mm}| < 1\}$ by the adjugate formula. Let $T = Z^{-1}$, $\xi = T_{mm}$, $s = \sqrt{1 - |\xi|^2}$, $D = P + sL$, $\Delta = D^{-1}$. The two matrices

$$Y_0 = -L - \tfrac{1}{2} P T^* L T (P + 2L), \qquad Y_1 = -L - \tfrac{1}{2} P T L T^* (P + 2L)$$

are precisely the `sectionCoeff` matrices with arguments (Z, Z^*) and (Z^*, Z) in `New/RealPlace/GramSectionBounds.lean`.

Lemma 25.13 (The actual section and double-coset parameter)**.** *For $Z \in \Omega$ the source Gram section acts on F, F' by*

$$\gamma_Z = \begin{pmatrix} Y_0 & T^*\Delta \\ D & 0 \end{pmatrix} \begin{pmatrix} T & Y_1\Delta \\ 0 & 1 \end{pmatrix}^{-1}$$

and is the identity on M. It is unitary and satisfies $Z(\gamma_Z) = Z$. If $Z(g) = Z$ then $g = s_L \gamma_Z s_R^{-1}$ with s_L, s_R in the pointwise stabilizer S_0 of F.

Proof. The displayed product has first block column $\binom{Y_0}{D}Z$, and second column $\binom{T^*}{0}\Delta - \binom{Y_0}{D}ZY_1\Delta$, which are exactly the source's two columns `sectionFrame` and `sectionMatrix`. For the verification of unitarity write X and Y for the right and left factors before the inverse. Directly from the above formulas,

$$LY_0 = LY_1 = -L, \quad PY_0 + Y_0^*P = -T^*LT + |\xi|^2 L, \quad PY_1 + Y_1^*P = -TLT^* + |\xi|^2 L.$$

Using $DLD = s^2L$, $\Delta P = P$, the upper-left block of Y^*J_0Y is $-T^*LT$: it is $-L + (Y_0^*P + PY_0) + s^2L$. The upper-right block is $(-Y_0^*L + DP)T^*\Delta = T^*\Delta$. The lower-right block is $-\Delta TLT^*\Delta$. Those of X^*J_0X are respectively $-T^*LT$, $T^*(-LY_1\Delta + P) = T^*\Delta$, and

$$-s^{-2}L + \Delta(Y_1^*P + PY_1)\Delta + L = -\Delta TLT^*\Delta.$$

The last cancellation is $1 + (-1 + |\xi|^2)s^{-2} = 0$. Thus $Y^*J_0Y = X^*J_0X$, proving unitarity. Furthermore $F^*J_0\gamma_Z F = (-LY_0 + PD)Z = Z$.

The Gram matrix of the two frames (F, gF) is $\left(\begin{smallmatrix} -L & Z \\ Z^* & -L \end{smallmatrix}\right)$. It is nondegenerate: if a pair (a, b) is in its kernel, $b = TLa$, $a = T^*Lb$, then $La = |\xi|^2 La$, whence both vanish. The isometry of the two nondegenerate frame spans $(F, \gamma_Z F)$ and (F, gF) respecting their ordered bases extends to the ambient space by choosing an isometry of their orthogonal complements, of identical signature. This extension s_L fixes F, and so does $s_R = g^{-1}s_L\gamma_Z$, proving the assertion. □

In particular the source's `coe_gramSection`, `realFrameGram_gramSection`, and `exists_eq_mul_gramSection_mul` are proved by the actual displayed section, not by an unspecified measurable choice.

Lemma 25.14 (Quantitative section bounds)**.** *Suppose $P_0 \geq 8$ and at every Y within operator distance $t > 0$ of Z one has $\|Y\|, \|Y^{-1}\|, (1 - |(Y^{-1})_{mm}|^2)^{-1/2} \leq P_0$ with $Y \in \Omega$. Then*

$$\|\gamma_Z\| \leq P_0^{12}, \qquad \|D\gamma_Z\| \leq 2P_0^{12}/t.$$

Moreover, if $M_0 \geq 8m$, $\mathfrak{g}(Z) \geq M_0^{-1}$, $|Z_{ab}| \leq M_0^2$ and $P_0 \geq M_0^{9m}$, every Z' with $\|Z' - Z\| < P_0^{-6}$ satisfies

$$\mathfrak{g}(Z') \geq \tfrac{1}{2}\mathfrak{g}(Z), \quad Z' \in \Omega, \quad \|Z'\|, \|(Z')^{-1}\|, (1 - |((Z')^{-1})_{mm}|^2)^{-1/2} \leq P_0.$$

Proof. For independent complex variables Z_1, W_1 in the product balls centered at Z, Z^* of radius t, use the defining formulas of `sectionMatrix` with Z^* replaced by W_1. The square root is of $1 - (Z_1^{-1})_{mm}(W_1^{-1})_{mm}$, in the right half-plane; its absolute value is at most $\sqrt{2}$ and its inverse at most P_0, by the assumed corner-defect bounds on both variables. The two section coefficients have norm at most $1 + P_0^2 \leq 2P_0^2$. The first block-column matrix $S = (FY_0 + F'D)Z_1$ has norm at most $3P_0^3$. In the source formula

$$1 + (S - F)F^* + ((FW_1^{-1} - SY_1)\Delta - F')F'^*$$

these bounds give norm at most $3 + 3P_0^3 + P_0^2 + 6P_0^6 \leq P_0^{12}$. The formula is holomorphic in the two independent complex variables. Cauchy's estimate on complex lines in the product ball bounds its derivative by P_0^{12}/t in the maximum norm. Restricting to the real slice (Y, Y^*) gives the claimed, slightly weaker, derivative bound.

For the second part, $\|Z\| \leq mM_0^2 \leq M_0^3$, and along a segment of length at most 1 from Z the operator norm is at most M_0^4. For a j-minor determinant the derivative is bounded in a unit operator direction by $jM_0^{4(j-1)}$: use the column multilinear formula and Hadamard's

inequality. Thus on such a segment the derivative of $\mathfrak{g}$ has norm at most $4mM_0^{8m}$. Multiplying by $P_0^{-6} \le M_0^{-54m}$ bounds its variation by $M_0^{-1}/2$. This proves the discriminant bound and positivity. The adjugate identity gives $Z' \in \Omega$, $|\det Z'| \ge (2M_0)^{-1/2}$, and

$$\|(Z')^{-1}\| \le M_0^{4(m-1)}\sqrt{2M_0}, \qquad (1-|((Z')^{-1})_{mm}|^2)^{-1/2} \le M_0^{4m}\sqrt{2M_0}.$$

Both are at most $P_0 \ge M_0^{9m}$; so is $\|Z'\|$. □

These are the precise bounds in `norm_sectionMatrix_le` and `mem_openCell_and_bounds_of_near`.

Theorem 25.15 (Open-cell Haar density)**.** *Let U^0 be the source frame-fixing Bessel unipotent, with its parameter Lebesgue measure, and $H = U(X_0)$ with probability Haar. For a local smooth section $h(c)$ of the map $H \to S^{2k-1}$, $h \mapsto hx$, the injective coordinate map from $\Omega \times S^{2k-1} \times U^0 \times U^0 \times H$ (using a section on a measurable partition of the sphere) is*

$$(Z, c, u, v, h_R) \longmapsto uh(c)\gamma_Z(vh_R)^{-1}.$$

Its Haar density is

$$\frac{\operatorname{vol} U(k)}{\operatorname{vol} U(r)}\mathfrak{g}(Z)^{k-1}\, dZ\, d\sigma(c)\, du\, dv\, dh_R. \tag{25.10}$$

Here $\operatorname{vol} U(j) = \prod_{i=1}^{j} 2\pi^i/(i-1)!$ is the source's compact dual normalization and $d\sigma$ is ordinary (unnormalized) sphere area.

Proof. First, the pointwise stabilizer of F is $U^0 \rtimes H$. In coordinates (e, X_0, e^*) with the negative vector fixed, elements fixing the e's must act identically on the quotient dual to them; the Hermitian relations force the induced action on X_0 to be unitary. After removing this action the matrix has the unique form

$$u(a,B) = \begin{pmatrix} 1 & a & B - aa^*/2 \\ 0 & 1 & -a^* \\ 0 & 0 & 1 \end{pmatrix}, \qquad B^* = -B.$$

This is the source's `unipotentOf`. Its parameters are the complex entries of a and independent real and imaginary entries of the skew matrix B (diagonal imaginary entries). The expression is a product-group parametrization with this Lebesgue Haar measure (triangular additive changes of variables under multiplication).

The two frames $F, \gamma_Z F$ span $F \oplus F'$ since the bottom block DZ of the latter is invertible. Consequently $S_0 \cap \gamma_Z S_0 \gamma_Z^{-1}$ is exactly the compact unitary group of M, acting identically on F, F'. If two of the displayed chart points are equal, their Gram matrices Z are equal, and the relative left and right stabilizer factors belong to this intersection. By the unique semidirect-product decomposition on the left, the two sphere columns hx then coincide; with the chosen section this forces equality of the left factors. The right factors are then equal too. This proves injectivity. The image is the open cell, by Lemma 25.13 and absorption of the remaining compact $U(M)$ into the right stabilizer (it commutes with γ_Z).

We calculate the Jacobian in the invariant skew coordinates $J_0 g^{-1} dg$; Lebesgue measure on skew matrices means the independent entries just specified. The compact-dual Haar measure in these coordinates is divided by $\operatorname{vol} U(r)$: at the identity the source's Cayley chart has derivative the identity in skew coordinates, so its density there has exactly this value. Left or right invariance extends the normalization to any chart. For the product $g = s_L \gamma_Z s_R^{-1}$,

discard the outer congruence by s_R^{-1}, of absolute determinant one since s_R is unitary. The remaining differential is the sum of

$$\gamma_Z^* B_L \gamma_Z, \qquad \gamma_Z^* J_0\, d\gamma_Z, \qquad -B_R,$$

where $B_L = J_0 s_L^{-1} ds_L$, $B_R = J_0 s_R^{-1} ds_R$. Both stabilizer skew matrices have zero rows and columns on F. Introduce the invertible block matrix on F, F', M

$$K_Z = \begin{pmatrix} T\Delta & Y_1\Delta & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

By the exact section formula,

$$\gamma_Z K_Z = \begin{pmatrix} Y_0\Delta & T^*\Delta & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

After congruence by K_Z, the left tangent of the stabilizer is therefore supported on the skew block $F \oplus M$, and the right tangent on $F' \oplus M$. Their only overlap is the skew M, M block; that block may be assigned to the right. On the left the independent coordinates are precisely the skew block with the M, M entries omitted, corresponding to the unipotent and the sphere tangent. On the right they are the full stabilizer skew block, corresponding to the unipotent and all of H. The section differential supplies the remaining complex cross block F', F:

$$F'^* K_Z^* \gamma_Z^* J_0 (d\gamma_Z) K_Z F = \Delta T(dZ) T \Delta.$$

Indeed $(\gamma_Z K_Z)F' = FT^*\Delta$, $K_Z F = FT\Delta$, and $F^* J_0 \gamma_Z F = Z$. The coordinates of the full skew matrix now split, by a permutation and conjugations of independent entries of absolute determinant one, into these three sets of blocks. The overall map is block triangular; the section differential may also contribute to the stabilizer blocks but not conversely. The determinant of $dZ \mapsto \Delta T(dZ)T\Delta$ as a real linear map on complex m-matrices is $|\det(\Delta T)|^{4m} = \mathfrak{g}(Z)^{-2m}$. The stabilizer-coordinate map has determinant one before the compact coordinates are converted to their invariant measures. For the unipotent this follows explicitly from its logarithmic derivative: $u^{-1}du$ has off-diagonal blocks $da, -da^*$, and skew central block $dB + (a\, da^* - da\, a^*)/2$. The change is triangular with unit determinant. Conjugating this by the compact factor rotates the da block with absolute determinant one. On the left the compact skew entries are the column of $h(c)^{-1}dh(c)$ at x, and its imaginary diagonal entry: together they are an isometry of the real tangent of the unit sphere into the independent column coordinates, since $dh(c)x = dc$ and $h(c)$ is unitary. Thus its volume element is exactly $d\sigma(c)$. On the right all compact independent skew entries have volume element $\operatorname{vol} U(k)\, dh_R$ by the source's probability normalization.

Finally skew congruence $B \mapsto K_Z^* B K_Z$ has real determinant of absolute value $|\det K_Z|^{2r}$: by singular-value decomposition it suffices to check positive diagonal entries, where each diagonal skew coordinate is multiplied by the square of one entry and each complex off-diagonal coordinate by the product of two entries. Here $|\det K_Z|^2 = |\det T|^2 s^{-2} = \mathfrak{g}(Z)^{-1}$. Removing this congruence converts the determinant to $\mathfrak{g}(Z)^{r-2m} = \mathfrak{g}(Z)^{k-1}$, as required. Ordinary change of variables on local smooth sections of the sphere, partitioned measurably to avoid overlap, gives (25.10). □

The theorem gives the exact constant

$$c_G = \frac{\operatorname{vol} U(k)}{\operatorname{vol} U(r)} \frac{2\pi^k}{(k-1)!}$$

of `Shape.openCellConst`. Integrating the sphere with probability area shows the precise Haar lower bound `le_lintegral_of_gramSection` in `New/RealPlace/SobolevCell.lean`: if a nonnegative test function has lower bound L_1 on the integral over Z, u, v, h_R with density $c_G \mathfrak{g}(Z)^{k-1}$ for every fixed left compact factor, then its full group integral is at least L_1. The argument proves this directly, without restricting to a particular sphere chart. In the source's explicit chart, the sphere column of its rank-one Cayley coordinates and the full right Cayley coordinates supply exactly the two ordinary Cayley densities. For completeness, this can be checked in the sphere coordinates without any undisclosed volume factor. In the order x, M, let $S = \left(\begin{smallmatrix} it & -w^* \\ w & 0 \end{smallmatrix}\right)$ and $D_c = 1 - S/2$. Its Cayley column is $(2\alpha - 1, \alpha w)$, where $\alpha = (1 - it/2 + \|w\|^2/4)^{-1}$. It is injective and covers the sphere except $-x$: recover α from the first entry, then w from the remaining entries, and $t = -2\,\mathrm{Im}(1/\alpha)$; the real part relation follows from the unit norm. The logarithmic Cayley derivative in skew matrices is $D_c^{-*}(dS)D_c^{-1}$ (differentiate $2D_c^{-1} - 1$). Let $\mathcal{V}$ be the space of skew matrices supported on the column and row of x, and $\mathcal{M}$ the complementary skew block on M. The determinant of the projected column derivative is the determinant on the full skew space of the map

$$(V, Q) \longmapsto D_c^{-*} V D_c^{-1} + Q, \qquad V \in \mathcal{V},\ Q \in \mathcal{M}.$$

Congruence by D_c converts this map into $V + D_c^* Q D_c$. Its $\mathcal{M}$ block is precisely Q, since D_c has identity lower-right block, so its determinant has absolute value one. The original map has determinant $|\det D_c|^{-2k}$, the exact `cayleyDensity` of the partial compact chart. The full right Cayley derivative is the same congruence formula in dimension k and likewise has determinant $|\det D_c|^{-2k}$ with its full skew argument. In the source's `skewOfReal` the independent off-diagonal real coordinate pairs are multiplied by the orthogonal matrix taking them to $((y_{ab} - y_{ba})/\sqrt{2}, (y_{ab} + y_{ba})/\sqrt{2})$; hence the Lebesgue normalization used in these computations is exactly that of the source. This also supplies the specific `openCellChart_jacobian` and injectivity assertion of `New/RealPlace/CellNeighbourhood.lean`.

25.7 A quantitative tube and the Sobolev restriction

Two analytic bounds complete the use of the Gram section for restriction at a fixed Gram matrix.

Lemma 25.16 (Tube of controlled fibre measure)**.** *For $T_0 \geq 8r$ there are a compact A in the group and a measurable $\Omega_0 \subset U^0$ of positive finite Haar measure such that*

$$\|g\| \leq T_0^{20m} \quad (g \in A), \quad \Omega_0\{\|g\| \leq T_0\} \subset A, \quad \int_{U^0} 1_A(ug)\, du \leq T_0 \,\mathrm{vol}(\Omega_0) \quad (g \in G).$$

Proof. Conjugate by the inverse of $d_0 = \mathrm{diag}(T_0^6 I_e, I_{X_0,z}, T_0^{-6} I_{e^*})$; this automorphism j of U^0 sends a to $T_0^{-6}a$ and B to $T_0^{-12}B$. For $R \geq 2$ let $A_R = \{u : \|j(u)\| \leq R\}$. These sets are compact of positive finite Haar measure. The matrix packing proof of Lemma 25.12, with $\delta = 1$ and center group ball A_2, shows

$$\mathrm{vol}(A_{T_0^2}) \leq (1 + 2T_0^4)^{2r^2}\,\mathrm{vol}(A_2) \leq T_0^{10r^2}\,\mathrm{vol}(A_2).$$

The same proof works for the scaled Haar on the image of j, without any degree factor. Since $10r^2 + 2 \leq T_0^2$, some integer R between 2 and $10r^2 + 1$ satisfies $\mathrm{vol}(A_{R+1}) \leq T_0\,\mathrm{vol}(A_R)$. Otherwise multiply the strictly opposite inequalities. Take $\Omega_0 = A_R$ and $A = A_R\{\|g\| \leq T_0\}$. Its elements have norm at most $T_0^{12} R T_0 \leq T_0^{15} \leq T_0^{20m}$. For a nonempty fibre $\{u : ug \in A\}$ choose $u_0 g = s_0 z_0$ with $s_0 \in A_R$, $\|z_0\| \leq T_0$. If $ug = sz$, also with $s \in A_R$, $\|z\| \leq T_0$, then $u u_0^{-1} s_0 = s(z z_0^{-1})$, and $v = z z_0^{-1}$ belongs to U^0 and has norm at most T_0^2. All three

nonidentity blocks of $j(v)-1$ are scaled by at most T_0^{-6}, so $\|j(v)-1\| \le 3T_0^{-6}(T_0^2+1) \le T_0^{-2}$. Hence $\|j(uu_0^{-1}s_0)\| \le R(1+T_0^{-2}) \le R+1$. By right invariance the whole fibre has measure at most $\mathrm{vol}(A_{R+1})$, proving the claim. □

This is the `exists_tube` of `Lib/DiscreteSeries/Bessel/RealMeasures.lean`.

Lemma 25.17 (Bounded frame fibres)**.** *Let $w \ge 0$ be measurable and invariant under the source left Bessel group including U^0H. Suppose $\eta \ge 0$ is measurable, and $\int_{U^0} \eta(uy)\,du \le K_0$ for every y. Assume that whenever $\eta(y)\eta(sy) \ne 0$ for $s \in U$, where U is the full Bessel unipotent, all the entries of its frame matrix are at most $R_0 \ge 1$ in absolute value. Then*

$$\int_G \eta(y)w(y)\,dy \le K_0(2R_0)^{2m^2} \int_{U\backslash G} w(y)\,d\dot{y}.$$

Proof. Disintegrate the Haar integral over U and the quotient, using the free-entry parameter measure of the source; the invariant factor is constant along each orbit. In a nonempty fibre, translate by a chosen point of its support. The assumed frame bound puts its image in the upper unitriangular frame coordinates inside complex disks of radius R_0. There are at most m^2 such coordinates. Each disk has area $\pi R_0^2 \le (2R_0)^2$. Disintegrating over these free coordinates and the kernel U^0 of the frame projection gives the remaining bound K_0 on each fibre, proving the formula. Empty fibres contribute zero. □

This is `lintegral_mul_le_of_frameMatrix_le`; the quotient and additive parameter normalization already proved in the Bessel-measure chapter is essential.

Proposition 25.18 (Sobolev restriction at one Gram matrix)**.** *Let $P_0 \ge 8r$ and let every Y with $\|Y-Z\| < P_0^{-6}$ lie in the bounded-cell region of Lemma 25.14 with bound P_0 and $\mathfrak{g}(Y) \ge c_0 > 0$. Suppose F has the source's differentiated Schur bounds with constants P_0, D_0 (words of length at most $2m^2$). For an integrable scalar ϕ supported where $\|y\| \le P_0^{20}$,*

$$c_G c_0^{k-1} \int_H \int_{\substack{u\in U^0\\ \|u\|\le P_0^{20}}} \left\| \int_G \phi(y) F(\gamma_Z(uh)^{-1}y)\,dy \right\|^2 du\,dh \le P_0^{2000m^2} D_0 \int_G |\phi|^2.$$

Proof. Let p run over the real cube $[0, P_0^{-8}]^{2m^2}$ in the independent real and imaginary entries of a complex matrix; write $E(p)$ for that matrix. Its norm is at most $2mP_0^{-8} < P_0^{-6}/2$. For $g(p,u,h) = \gamma_{Z+E(p)}(uh)^{-1}$ and a coordinate derivative of the cube, write $\partial g = gA_0$. The norm of the section derivative on this smaller ball is at most $4P_0^{18}$ by Lemma 25.14, and $\|\gamma\|, \|\gamma^{-1}\| \le P_0^{12}$. Since $\|u\|, \|u^{-1}\| \le P_0^{20}$ and h is Euclidean unitary, one obtains

$$\|A_0\| \le P_0^{40} P_0^{12}\, 4P_0^{18} \le P_0^{71}.$$

This holds for each coordinate direction. Differentiating $F(g(p,u,h)y)$ replaces the infinitesimal direction by $y^{-1}A_0y$. Expand this into matrix units. To keep the coefficient test function independent of p, u, h, for each four-index tuple use the scalar multiplier $y_{ac}^{-1}y_{db}/P_0^{40}$ on ϕ. Its absolute value on the support is at most one. After a word of such operations denote the new multiplier by ϕ_i and the Bessel derivative word by F_i. Then $|\phi_i| \le |\phi|$, all have the same support, and the coordinate derivative of $f_i(p,u,h) = \int \phi_i(y)F_i(g(p,u,h)y)\,dy$ has norm bounded by $P_0^{112}\sum_j \|f_{ji}(p,u,h)\|$, with at most $r^4 \le P_0^4$ summands. Differentiation under the integral is legitimate: each Bessel derivative is bounded continuous by the definition of the differentiated Schur bounds, and the scalar multiplier is bounded on the support of ϕ.

The elementary one-dimensional estimate, for a differentiable Hilbert-valued function f on $[0,\varepsilon]$, is $\|f(0)\|^2 \le (2/\varepsilon)\int_0^\varepsilon(\|f\|^2+\|f'\|^2)$ if $0<\varepsilon\le 1$: write $f(0)=f(t)-\int_0^t f'(s)ds$, square by Cauchy–Schwarz and average over t. Iterating this along the $2m^2$ independent coordinates and applying Cauchy–Schwarz to the at most r^4 summands of the derivative just estimated, gives a factor at most

$$[2P_0^8(1+P_0^{232})]^{2m^2} \le P_0^{484m^2}$$

times the maximum integrated squared norm of f_i on the cube for $|i|\le 2m^2$.

To bound the latter integrals use the Haar formula (25.10) with the Gram matrices restricted to the ball $\mathcal{Q}$ of radius $P_0^{-6}/2$ about Z. Since the norm of $f_i(g)=\|\int\phi_i(y)F_i(gy)dy\|$ is invariant under the left stabilizer U^0H, the left compact average and left unipotent factor in the chart can equivalently be disintegrated using a nonnegative measurable cutoff ζ for $U^0\backslash G$ of U^0-orbit integral one. Such a cutoff is obtained from quotient Haar disintegration, or from a measurable local cross-section and any nonnegative integrable function of mass one on U^0. With the Haar density lower bound $c_G c_0^{k-1}$ on $\mathcal{Q}$, the integral over the cube and the restricted right variables is bounded by $\int_G \zeta(g)1_{\{Z(g)\in\mathcal{Q}\}}|f_i(g)|^2\,dg$. Indeed in the chart the left unipotent average of ζ is one, so this last integral is the corresponding unrestricted right-variable integral with its exact Jacobian. Apply Lemma 25.17 to $\eta(g)=\zeta(g)1_{\{Z(g)\in\mathcal{Q}\}}$. For any $s\in U$ with both $Z(g),Z(sg)\in\mathcal{Q}$, the frame matrix A_s satisfies $Z(g)=A_s^*Z(sg)$ by unitary invariance. Both matrix norms and the inverse norm are bounded by P_0, so entries of A_s are bounded by P_0^2. The U^0 fibre integral is at most one. Thus the integral is bounded by $(2P_0^2)^{2m^2}\int_{U\backslash G}|f_i|^2$, which is at most $(2P_0^2)^{2m^2}P_0^{2|i|}D_0\int|\phi|^2$ by the assumed Schur bound. Combine this with the cube trace factor above. All these factors together are at most $P_0^{2000m^2}$, proving the assertion with the stated constant. □

This supplies `lintegral_gramSection_integral_le` of `New/RealPlace/SobolevCell.lean`, using the actual section and Jacobian just proved.

25.8 The randomized open-cell bound

We can retain the exact mass and support conditions in the source by combining the preceding arguments. Write $d_I=|I|$ and $d=\deg$ for a Bessel system W with its full-function L^2 norm-squared d_I, and denote the full function by W in this subsection. Let b be the system parameter, with $b\ge 8r$, $\rho_0\ge b$. Suppose that there is a point in the intersection of $\mathrm{Box}(\rho_0)$ and $\gamma\,\mathrm{Box}(\rho_0)$, where the box means that all frame columns $y^{-1}f_a$ have length at most ρ_0. Let $Z=Z(\gamma)$, and suppose $\mathfrak{g}(Z)>0$.

Theorem 25.19 (Open-cell bound with the exact cell mass)**.** *There are a compact set A, a probability space bearing a measurable random translation $t\in U^0H$, and a finite measure n_0 on the group with*

$$n_0(G)\le\left(\frac{\rho_0^{9m}}{\mathfrak{g}(Z)}\right)^{10^5m^4}\frac{d_I^2}{d\,c_G\,(\mathfrak{g}(Z)/2)^{k-1}},$$

such that

$$\mathbb{E}\left\|\int_A\phi(y)W(\gamma ty)\otimes\overline{W(ty)}\,dy\right\|^2\le\int|\phi|^2\,dn_0.$$

For every translation in the probability support, A contains every $y\in\mathrm{Box}(\rho_0)$ with $\mathrm{besselSize}(\gamma ty)\le\rho_0$.

Proof. The intersection of the two boxes implies $|Z_{ab}| \le \rho_0^2$ by invariance of the Hermitian pairing: if $y_0, \gamma^{-1}y_0$ lie in the box, pair their inverse frame columns. Put

$$M_0 = \rho_0^{9m}/\mathfrak{g}(Z), \quad P = M_0^{9m}, \quad T_0 = P^5, \quad P_1 = T_0^m.$$

Since $\mathfrak{g}(Z) \le |\det Z|^2 \le (m\rho_0^2)^{2m} \le \rho_0^{6m}$, one has $M_0 \ge \rho_0 \ge 8r$ and $\mathfrak{g}(Z) \ge M_0^{-1}$. Lemma 25.14 supplies the bounded-cell region with parameter P, and hence with the larger parameter P_1, and lower discriminant $c_0 = \mathfrak{g}(Z)/2$. Also $P_1 \ge b^5$, so the Bessel system's differentiated Schur bound applies at P_1 with $D_0 = d_I/d$.

Choose the tube A, Ω_0 of Lemma 25.16 with parameter T_0, and use the double-coset factorization $\gamma = s_L \gamma_Z s_R^{-1}$. Take $h \in H$ with probability Haar, $u \in \Omega_0$ with relative probability Haar, and put $t = s_R(uh)^{-1}$. Left Bessel covariance implies that the norm of the tensor integral in the statement equals that of $\int_A \phi(y) W(\gamma_Z(uh)^{-1}y) \otimes \bar{W}(y)\, dy$: the same fixed unitary factors on the two tensor entries can be extracted outside the integral. Apply Proposition 25.18 to each coordinate test function $1_A \phi \overline{W_j}$, and add the inequalities. The range of $u \in \Omega_0$ is allowed by that proposition because $\|u\| \le T_0^{14} \le P_1^{20}$ by the tube proof, and A has norm bound $T_0^{20m} = P_1^{20}$. Thus the random-translation inequality holds with

$$dn_0(y) = \frac{P_1^{2000m^2} d_I}{d\, c_G c_0^{k-1} \operatorname{vol}(\Omega_0)} 1_A(y) \|W(y)\|^2\, dy.$$

For $y, vy \in A$, the frame entries of $v \in U$ have absolute value at most $\|v\| \le T_0^{40m}$. Lemma 25.17, the tube fibre bound, and the full-function norm give

$$\int_A \|W\|^2 \le T_0 \operatorname{vol}(\Omega_0)(2T_0^{40m})^{2m^2} d_I.$$

The measure is therefore finite. The power multiplying $d_I^2/(d\, c_G c_0^{k-1})$ is at most $M_0^{10^5 m^4}$: $P_1^{2000m^2} = M_0^{90000m^4}$, the tube factor contributes $M_0^{45m+3600m^4}$, and $2^{2m^2} \le M_0^{2m^2}$. This proves the exact mass constraint. All integrals and random-variable inequalities first hold for bounded measurable tests on compact A, the source's `IsL2Contraction` tests, and extend by completion to the indicated finite-measure L^2 space.

For the support condition put $q = (uh)^{-1}y$ so that q lies in the same frame box as y. From the Bessel-size assumption and its definition as an infimum of norms plus one, choose $v \in U$ with $\|v\gamma_Z q\| \le \rho_0$. Indeed the strict one-unit slack realizes the infimum within one by an element of UH, and its compact factor may be removed without changing the Euclidean norm. Set $g = v\gamma_Z q$ and let A_v be the frame matrix of v, an upper unitriangular matrix. The frame Gram of $v\gamma_Z$ is $Z' = A_v^{-*} Z$ because $v^{-1}F = FA_v^{-1}$. Its entries have absolute value at most ρ_0^2: they pair the columns $g^{-1}F$ and $q^{-1}F$. Its discriminant is $\mathfrak{g}(Z)$, since multiplication by a lower unitriangular factor preserves the determinant and the leading principal minor.

For any such Z' one has $\|Z'\| \le M_0^3$ and, using $\mathfrak{g}(Z') \ge M_0^{-1}$, $\|(Z')^{-1}\| \le M_0^{3m}$ by the singular-value bound. The same bounds hold for Z. Thus $\|A_v\| \le M_0^{3+3m}$ from $A_v^* = Z(Z')^{-1}$, and

$$\|q^{-1}\gamma_Z^{-1}F\| = \|g^{-1}FA_v\| \le M_0^{3m+5}, \qquad \|q^{-1}F\| \le M_0^2.$$

The inverse of the explicit section satisfies $\gamma_Z^{-1}F = (FY_1 + F'D)Z^*$ by multiplying the two block factors of Lemma 25.13. As $\|\Delta\| \le M_0^{3m+1}$ and $\|Y_1\| \le 2M_0^{6m}$, solve for $q^{-1}F'$:

$$\|q^{-1}F'\| \le (M_0^{3m+5}M_0^{3m} + 2M_0^{2+6m})M_0^{3m+1} \le 3M_0^{9m+6}.$$

Control of these frame and dual columns gives an operator norm bound independent of k. In fact let P_- be the Euclidean orthogonal projection onto the negative subspace, spanned by z and $(e_a - e_a^*)/\sqrt{2}$. Since $q^* J_0 q = J_0 = 1 - 2P_-$ and $q^* = J_0 q^{-1} J_0$,

$$\|q\|^2 \le 1 + 2\|q^* P_- q\| \le 1 + 2 \sum_{w \text{ negative orthonormal basis}} \|q^{-1}w\|^2.$$

The frame bounds make this at most $1 + 4m(3M_0^{9m+6})^2$, so $\|q\| \le T_0 = M_0^{45m}$. Finally $y = uhq$, and $\|hq\| = \|q\| \le T_0$. The tube inclusion $\Omega_0\{\|g\| \le T_0\} \subset A$ proves the required support claim. □

This gives the whole `BesselSystem.open_cell` in `New/RealPlace/OpenCellBound.lean` with its `cellMass`, for a Bessel system satisfying the stipulated Schur bounds and full-function norm. It does not require the lower-floor bandlimit theorem to prove this implication.

Current scope. The coefficient construction, Casimir rigidity, normalized orbital comparison and differentiated Schur bound have now been derived. The compact averaging argument applies to a genuine noncompact unitary group rather than only a definite one. These results do not themselves establish the existence of a simultaneous marked arithmetic Bessel datum, all the lower-floor weighted towers, or global packet coherence. The exact local Jacobian, tube, section derivative, Sobolev estimate and randomized open-cell bound have now been proved; the remaining uniform and arithmetic arguments stay separate. In the obligation ledger this provides the written proofs for B01 (the exact coefficient premise of normalized orbital comparison), B17 (lowest discrete-series coefficient and tail estimates), B18 (differentiated Schur), B20 (Gram-section restriction), and B06 (real open-cell uniform estimate). The Bessel coefficient's representation-theoretic identification needed by A06 is thereby supplied, but the actual arithmetic marked choices and good-datum realization in A06 do not yet follow. In particular no inference of global filling from local estimates alone is intended.

Chapter 26

Green labels, quantitative finite mass, and further finite-place comparisons

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 29, 30 and the final audit in Chapter 57.

We continue the proof of the finite interfaces of Chapter 18 and Chapter 24. In particular the degree and regular-representation bound previously marked as B03 are proved here; no asymptotic character estimate is needed. We identify the class functions used by the actual source, including the sign, character naming, and coefficient of the identity, rather than assuming an arbitrary parametrization of the irreducible representations.

26.1 The class functions actually used for the labels

Let $k = \mathbb{F}_q$, and let $G_n = \mathrm{GL}_n(k)$. Fix an injective multiplicative character $T : \bar{k}^\times \to \mathbb{C}^\times$ as in `faithfulChar`. Such a character exists by choosing compatible generators in the direct limits of prime-to-char(k) roots of unity. For an element $x \in \mathbb{F}_{q^d}^\times$, take $a \geq 0$ with $T(x) = \exp(2\pi i a/(q^d - 1))$ and write $\theta_x = T^a$ on $\mathbb{F}_{q^d}^\times$. The restriction is independent of a modulo $q^d - 1$. This is precisely the restriction of `thetaCharT(q^d)x`. The naming map is an isomorphism to the multiplicative character group and satisfies

$$\theta_{x^q}(z) = \theta_x(z^q), \qquad \theta^{(d)}_{a_0}(z) = \theta^{(1)}_{a_0}(N_{\mathbb{F}_{q^d}/k} z) \quad (a_0 \in k^\times).$$

Indeed the exponent naming a_0 in degree d is the one naming it in degree one multiplied by $(q^d - 1)/(q - 1)$. In particular if P is a monic irreducible polynomial of degree d with $P(0) \neq 0$ and x is a root, θ_x has Frobenius orbit of length d.

Let $\mathcal{H}$ denote the graded algebra of class functions on the groups G_n, completed in the degree. Its multiplication in degree n is parabolic induction: for homogeneous F, H of degrees u, v it is

$$(F * H)(g) = \sum_{\substack{W \subset k^{u+v} \\ gW = W,\ \dim W = u}} F(g|_W) H(g|_{k^{u+v}/W}).$$

This follows from induction from the parabolic stabilizer of a u-dimensional subspace, summing over its cosets. We can restrict the source's Hall algebra of similarity classes of all

endomorphisms to invertible endomorphisms, since these stay invertible on invariant subspaces and their quotients. The constant class function of value one in every degree is $\mathbf{1}$; the unit for $*$, denoted 1, instead is supported on degree zero. Let $\mu = \mathbf{1}^{-1}$ for this product. Its value at g is the Möbius number of the lattice of g-invariant subspaces: it satisfies

$$\sum_{gW=W} \mu(g|_W) = \begin{cases} 1, & \dim g = 0, \\ 0, & \dim g > 0. \end{cases}$$

If $\eta = T^a$, write $\beta_\eta(g) = \sum_\xi \eta(\xi)$, with roots of the characteristic polynomial counted with multiplicity. This is `Hall.brauerLift`. The source defines

$$C_\eta = (\beta_\eta \mu) * \mathbf{1} = -\mu * \beta_\eta, \tag{26.1}$$

where juxtaposition $\beta_\eta\mu$ is pointwise multiplication. For the second equality, use the additivity of β_η on invariant subspaces and quotients: the sum $C_\eta + \mu * \beta_\eta$ at g is $\beta_\eta(g) \sum_{gW=W} \mu(g|_W) = 0$ in positive degree and also vanishes in degree zero. The formulas are those of `Lib/LocalGL/Green/HallAlgebra.lean` in the unabridged source and `greenCuspidal_mul_constOne` in `CuspidalFunction.lean`.

Lemma 26.1 (Exact Möbius and cuspidal values)**.** *Consider g as a $k[X]$-module with X acting as g. If it is semisimple with primary multiplicities t_P over $k[X]/P$, then*

$$\mu(g) = \prod_P (-1)^{t_P} (q^{\deg P})^{\binom{t_P}{2}}.$$

If it is not semisimple, $\mu(g) = 0$. The function $C_\eta(g)$ vanishes if the characteristic polynomial has more than one distinct irreducible factor. If that polynomial is P^m, where $e = \deg P$ and $t = \dim_{k[X]/P} \ker P(g)$, then

$$C_\eta(g) = -\sum_{\xi:\, P(\xi)=0} \eta(\xi) \prod_{j=1}^{t-1} (1 - q^{ej}). \tag{26.2}$$

Here t is the number of primary Jordan blocks, not the multiplicity m in the characteristic polynomial.

Proof. For semisimple g its invariant subspace lattice is a product of ordinary subspace lattices over $k[X]/P$. For a space of dimension t over $\mathbb{F}_Q$ the Möbius formula follows from the Gaussian binomial identity

$$\sum_{j=0}^{t} \begin{bmatrix} t \\ j \end{bmatrix}_Q (-1)^j Q^{\binom{j}{2}} z^j = \prod_{i=0}^{t-1} (1 - Q^i z)$$

with $z = 1$ (the identity follows by induction from the Gaussian binomial recursion). If a nonzero module is not semisimple, every proper submodule with possible nonzero Möbius number is semisimple by induction on dimension. All lie in the proper nonzero socle $S = \bigoplus_P \ker P(g)$, the kernels taken in their primary components. The sum of their Möbius numbers is the sum over all submodules of S, namely zero. The recursive definition gives $\mu(g) = 0$.

By the first expression for C_η in (26.1), only submodules of the socle contribute. The summands factor over the primary components, except that $\beta_\eta(g|_W)$ is additive over them. If at least two components are nonzero, each term of this additive decomposition contains the unweighted Möbius sum over another nonzero component, so the total is zero. For a single component of socle dimension t over $\mathbb{F}_{q^e}$, a submodule of dimension j has $\beta_\eta = j \sum_{P(\xi)=0} \eta(\xi)$, giving (26.2) by differentiating the Gaussian identity at $z = 1$. □

Let $\tau_P = (-1)^d (C_{\theta_x})_d$ (subscript means homogeneous component of degree d). This is exactly `greenTauTP`. Formula (26.2) shows independence of the chosen root x: replacing x by x^q permutes the roots ξ in the sum. Taking $g = 1$ of dimension d gives

$$\tau_P(1) = \prod_{j=1}^{d-1} (q^j - 1) > 0. \tag{26.3}$$

26.2 Irreducibility, projections, and finite twists

The only non-elementary finite-group character input needed in what follows is Green's theorem on symmetric functions of lifted eigenvalues: J. A. Green, *The characters of the finite general linear groups*, Trans. Amer. Math. Soc. **80** (1955), 402–447, Theorem 1 (statement in the introduction, proof in Section 3, pp. 413–415), DOI `https://doi.org/10.1090/S0002-9947-1955-0072878-2`. In its notation the chosen injection of the multiplicative group of a sufficiently large finite field into $\mathbb{C}^\times$ can be any such injection. An integral symmetric polynomial evaluated at the lifted eigenvalues is a virtual ordinary character. We apply it only to the power sum $\sum z_i^a$ with $a \geq 0$, using a finite field containing every eigenvalue in the rank in question, and the restriction of our T. This gives β_η as a virtual character in every degree, not merely an identification of some functions by their dimensions. We derive the remaining representation identities below.

For $u + v = n$, let $\Delta_{u,v} F$ be the mean of F on block triangular matrices with diagonal (g_u, g_v) over the off-diagonal block. On characters this is the character of the unipotent-radical invariants, as a representation of $G_u \times G_v$. Hence it is adjoint to parabolic induction for the probability-normalized finite group inner products. The bialgebra identity $\Delta(F * H) = \Delta(F) * \Delta(H)$ follows by the following Mackey calculation, with tensor factors and degrees kept separate. Write $V = A \oplus B$ for the outer block decomposition and P_A, P_W for the outer and inner subspace stabilizers. The P_A-orbits on subspaces W of a given dimension are indexed by $\dim(W \cap A)$, since P_A acts independently on $A, V/A$, and on the graph map into $A/(W \cap A)$. Choose the representative $W = A' \oplus B'$ with $A' \subset A$, $B' \subset B$. Restrict the induction from P_W to P_A using Mackey's decomposition over these orbits. Passing to invariants under the unipotent radical U_A of P_A, a summand is induced in the quotient P_A/U_A from the image of $P_A \cap P_W$; its inducing module is the invariants under $U_A \cap P_W$ of the original inducing module (this follows by writing induced functions constant along U_A-orbits). That image is the product of parabolics stabilizing A', B' in $\mathrm{GL}(A), \mathrm{GL}(B)$. On W and V/W, the intersection $U_A \cap P_W$ gives independently the two block unipotent radicals, arising from the arbitrary maps $B' \to A'$ and $B/B' \to A/A'$. The invariant inducing module therefore has the two corresponding block averages of F, H. Summing the resulting products of the two lower-rank parabolic inductions over $\dim A'$ proves the identity on characters, hence on all class functions by their linear span.

Now $\Delta \mathbf{1} = \mathbf{1} \otimes \mathbf{1}$, so $\Delta \mu = \mu \otimes \mu$. On any block triangular matrix β_η is the sum of its diagonal values. Using the first expression in (26.1) yields

$$\Delta C_\eta = C_\eta \otimes 1 + 1 \otimes C_\eta.$$

Thus every positive-degree component has zero block averages at nontrivial splits, the assertion `isCuspidal_greenTau`. Also μ is virtual in every rank by its inverse-series expansion in $1 - \mathbf{1}$, which is finite in each rank. By (26.1) every C_η is virtual.

We compute the ordinary inner product in rank d. Put $T_d = \mathbb{F}_{q^d}^\times$ embedded in G_d by multiplication on its underlying k-space. There is an exact pointwise identity

$$(C_\eta)_d \, \mu_d = \mathrm{Ind}_{T_d}^{G_d}(\eta|_{T_d}). \tag{26.4}$$

Both functions vanish off semisimple elements with just one primary factor: on the left by Lemma 26.1, on the right since elements in T_d are of that form. If g has characteristic polynomial P^m, $\deg P = e$, $d = em$, and is semisimple, then $t = m$ and its centralizer is $\mathrm{GL}_m(\mathbb{F}_{q^e})$. The formula for induced class functions gives

$$\mathrm{Ind}_{T_d}^{G_d}\eta(g) = \frac{|\,\mathrm{GL}_m(\mathbb{F}_{q^e})|}{q^d-1}\sum_{P(\xi)=0}\eta(\xi).$$

On the left of (26.4), the signs from C_η and from μ cancel, leaving $(q^e)^{\binom{m}{2}}\prod_{j=1}^{m-1}(q^{ej}-1)$ times the same root sum. This is the centralizer quotient by the elementary formula for the order of the general linear group. Hence (26.4) holds on every conjugacy class and gives the identity underlying `rankInner_greenCuspidal_mulMu`.

Let $\langle F,H\rangle = |G_d|^{-1}\sum_g \overline{F(g)}H(g)$. By adjointness and cuspidality only the rank d part of $\beta_{\eta'}\mu$ in $C_{\eta'} = (\beta_{\eta'}\mu) * \mathbf{1}$ contributes. Since μ is real, (26.4) and Frobenius reciprocity give

$$\begin{aligned}\langle C_{\eta,d}, C_{\eta',d}\rangle &= \langle C_{\eta,d}\mu_d, \beta_{\eta',d}\rangle \\ &= \langle \eta, \beta_{\eta'}|_{T_d}\rangle_{T_d} = \sum_{i=0}^{d-1}\langle \eta, \eta'\circ \mathrm{Frob}^i\rangle_{T_d}.\end{aligned}$$

For $\eta = \theta_x, \eta' = \theta_y$ with x, y of degree d the last sum is one if y is a conjugate of x, otherwise zero. Thus the τ_P are orthonormal virtual characters, with positive degree by (26.3). A virtual character of norm one is plus or minus a single irreducible character, and positive degree eliminates minus. For distinct irreducible polynomials P_i, the product $\Xi_c = \tau_{P_1} * \cdots * \tau_{P_s}$ for $c = \prod_i P_i$ has norm one: iterate adjointness and the bialgebra identity, assigning each primitive factor to a tensor slot. Only terms assigning exactly one factor to each slot can have nonzero inner product with the positive-degree cuspidal character there, and orthonormality gives a unique nonzero term when the unordered lists coincide. This also proves orthogonality for distinct squarefree labels (the same computation works when numbers of factors differ). The products are actual inductions. Thus the squarefree labels give distinct irreducible characters and establish the source's `labelChar`, `IsLabel.isIrrChar_labelChar`, and `rankInner_labelChar` for these ranks. Neither a classification of repeated labels nor the other parts of Green's complete character table are needed.

For a character χ of $k^\times$ let a_χ be named by $\theta^{(1)}_{a_\chi} = \chi$. Then

$$\tau_P(g)\chi(\det g) = \tau_{P^{(a_\chi)}}(g), \qquad \Xi_c(g)\chi(\det g) = \Xi_{c^{(a_\chi)}}(g), \tag{26.5}$$

where superscript (a) means multiply every root by a and make the polynomial monic. Indeed if $\mathrm{charpoly}(g) = Q^m$, $\deg Q = e$, the determinant is the norm from $\mathbb{F}_{q^d}$ of any root ξ of Q (here $d = em$); multiply the root sum in (26.2) by $\chi(N_{\mathbb{F}_{q^d}/k}\xi)$ and use the norm identity for names. This gives the equation on the nonzero classes, and it holds trivially elsewhere. For the product use multiplicativity of determinants in subspaces and quotients. This proves `labelChar_mul_det`. By finite Schur orthogonality,

$$e_c = \frac{\Xi_c(1)}{|G_r|}\sum_{g\in G_r}\overline{\Xi_c(g)}\,g \tag{26.6}$$

is the self-adjoint central projector onto this isotypic component, precisely `labelIdem`. Invalid labels are assigned zero. Distinct projectors are orthogonal. Tracing left multiplication on the regular representation shows $[1]e_c = \Xi_c(1)^2/|G_r|$, where $[1]$ denotes coefficient at the identity (each $g \neq 1$ permutes the group basis without fixed points). The same construction on a product of groups gives `tupleIdem`; their coefficients at the identity are the products of the separate coefficients.

Reflection annihilation. Let $l \neq 0$ be a linear functional on k^r and

$$N_l = \{1 + ul : u \in k^r,\ 1 + l(u) \neq 0\}.$$

This is the group fixing $\ker l$ pointwise, acting on its one-dimensional quotient by the determinant. The sum $\sum_{h \in N_l} \chi(\det h)^{-1} h$, divided by $|N_l|$, projects onto the $\chi \circ \det$ isotypic part under N_l. In the Jacquet space for blocks $(r-1, 1)$ of Ξ_c, the one-dimensional block has character χ only if the Hall factor $\tau_{X - a_\chi}$ occurs: apply the bialgebra identity and cuspidality of the factors, so only a single factor of degree one can enter this block. In degree one the formula above gives $\tau_{X - a_\chi} = \chi$. Thus if $c(a_\chi) \neq 0$ this isotypic part is zero. Multiplication by the reflection sum kills e_c in the group algebra. This proves `reflectionSum_mul_labelIdem` and, by tensoring, `Data.reflectionSum_mul_e_0` (where $\chi^{M_\omega} = 1$ makes the name an excluded root). It holds as well for $r = 1$ with the rank-zero block the unit.

26.3 Exact character degree and quantitative bad share

Proposition 26.2 (The coefficient bound for every label)**.** *For a squarefree label c of degree r write $d_1, \dots, d_s$ for the degrees of its monic irreducible factors. Then*

$$\Xi_c(1) = \frac{\prod_{j=1}^{r} (q^j - 1)}{\prod_{i=1}^{s} (q^{d_i} - 1)}, \qquad [1]e_c \ \geq\ \frac{1 - r/q}{q^r}.$$

Here $[1]$ *is the ordinary group-algebra coefficient at the identity, with no probability factor omitted.*

Proof. Counting ordered bases gives $|G_r| = q^{\binom{r}{2}} \prod_{j=1}^{r} (q^j - 1)$. The parabolic for block sizes d_i has order $q^{\sum_{i<j} d_i d_j} \prod_i |G_{d_i}|$. Their quotient is precisely $\prod_{j=1}^{r} (q^j - 1) / \prod_i \prod_{j=1}^{d_i} (q^j - 1)$, since the powers of q cancel. Multiply by the degrees $\tau_{P_i}(1)$ already computed to obtain the first identity. It follows from (26.6) that

$$\begin{aligned} [1]e_c &= \frac{\prod_{j=1}^{r} (q^j - 1)}{q^{\binom{r}{2}} \prod_i (q^{d_i} - 1)^2} = q^{-r} \frac{\prod_{j=1}^{r} (1 - q^{-j})}{\prod_i (1 - q^{-d_i})^2} \\ &\geq q^{-r} (1 - r/q). \end{aligned}$$

For the inequality discard the denominator (at most one) and use $\prod_{j=1}^{r} (1 - q^{-j}) \geq 1 - \sum_{j=1}^{r} q^{-j} \geq 1 - r/q$. This proves `labelChar_degree_bounds`, including nonnegativity of the real degree. □

Primitive is stronger than irreducible here: it means a root is a multiplicative generator of $\mathbb{F}_{q^r}^{\times}$, not merely that its character has a full Frobenius orbit. For any integer $N \geq 1$, writing its s distinct primes p_j in increasing order, $p_j \geq j + 1$ gives

$$\frac{\phi(N)}{N} = \prod_{j=1}^{s} (1 - 1/p_j) \geq \frac{1}{s+1}, \qquad s \leq \frac{\log N}{\log 2} \leq 2 \log N.$$

In particular $\phi(N) \geq N / (1 + 2 \log N)$, proving `le_totient_mul_log`. There are exactly $\phi(q^r - 1)/r$ primitive labels: generators lie in no proper subfield (their order would then divide $q^d - 1 < q^r - 1$), and each Frobenius orbit has r members.

Theorem 26.3 (Exact bad-share bound)**.** *Let f finite fields have common cardinality q; fix $1 < r < q$ and $M \geq 1$. A tuple of degree r monic polynomials is good if every factor is squarefree, has nonzero constant term and has no root $\xi \in k$ with $\xi^M = 1$, and at least one factor is primitive. If $e = \sum_{\mathbf{c}\ \mathrm{good}} \otimes_w e_{c_w}$, then for*

$$(1 + M + 2r)/q \leq \epsilon \leq 1, \qquad 2r(2r \log q + 1)L \leq f$$

we have

$$[1](1 - e) \leq f\epsilon + \exp(-L). \tag{26.7}$$

Proof. Use the uniform lower mass $m = (1 - r/q)q^{-r} > 0$ on every allowed label by Proposition 26.2. There are q^r degree r monics. The excluded roots are zero and at most M nonzero roots of $X^M - 1$; each excludes q^{r-1} monics. A non-squarefree monic is divisible by the square of a monic polynomial of degree j for some $1 \leq j \leq \lfloor r/2 \rfloor$. Counting all such factorizations (which overcounts) gives at most $\sum_{j=1}^{\lfloor r/2 \rfloor} q^j q^{r-2j} \leq rq^{r-1}$ bad monics. Thus the number A of allowed monics satisfies

$$Am \geq 1 - (1 + M + 2r)/q \geq 1 - \epsilon, \qquad Am \leq 1.$$

There are equally many in each field since finite fields of the same order are isomorphic. If B is the number of allowed but nonprimitive monics, then $B \leq q^r - \phi(q^r - 1)/r$. Using $m \leq q^{-r}$ and $q^r - 1 \geq q^r/2$ gives

$$Bm \leq 1 - \frac{\phi(q^r - 1)}{rq^r} \leq 1 - \frac{1}{2r(2r \log q + 1)}.$$

There are $A^f - B^f$ good tuples. By multiplying the coefficient bounds, $[1]e \geq (Am)^f - (Bm)^f$. If $f > 0$ and $L > 0$, the condition on L gives $Bm \leq 1 - L/f$, whence

$$\begin{aligned}[1](1 - e) &\leq 1 - (1 - \epsilon)^f + (1 - L/f)^f \\ &\leq f\epsilon + \exp(-L).\end{aligned}$$

If $L \leq 0$, use $[1](1 - e) \leq 1 \leq \exp(-L)$; if $f = 0$ the hypotheses force $L \leq 0$. This covers every case. The coefficient on the left is real because the sum consists of distinct orthogonal projectors. □

This proves `bad_share_le` of `New/Trace/BadShare.lean`, with exact constants. In the global data there are two families of chosen places. Applying it separately to their respective products of groups and using $1 - ab \leq (1 - a) + (1 - b)$ for $a, b \in [0, 1]$ gives the bound for the intersection of the two good conditions. The equality between this coefficient and the regular-representation share follows by tracing the regular action as above. It does not assert an analogous bound on an arbitrary nonregular representation.

26.4 Primitive labels and cubic avoidance

We verify the other arithmetic and character assertions of `BadShare.lean` which were previously deduced conditionally in Lemma 18.4. Put $N = q^r - 1$. If $M^2 \mid q - 1$ and $a_0^M = 1$, then scaling a primitive root ξ by a_0 leaves it primitive. Choose a generator ζ of $\mathbb{F}_{q^r}^\times$, write $\xi = \zeta^h$ with $(h, N) = 1$ and $a_0 = \zeta^{kN/M}$. Every prime dividing N divides N/M: for primes dividing M this holds by $M^2 \mid N$, and for the others automatically. Thus $h + kN/M$ is coprime to N. The norm of a primitive root is primitive in $k^\times$, since it equals $\xi^{(q^r-1)/(q-1)}$

and has order $q-1$. This is the product of all its Frobenius conjugates and equals that same power of every conjugate.

Suppose $r = 3^{a+1}$ and $9 \mid q-1$. If for some prime ℓ, exponent $d \geq 0$ and $0 < i < r$ one had $(\xi^{\ell^d})^{q^i} = \xi^{\ell^d}$, then $N \mid \ell^d(q^i-1)$. The gcd identity $\gcd(q^r-1, q^i-1) = q^{\gcd(r,i)}-1$ implies

$$\frac{q^r-1}{q^{r/3}-1} \mid \ell^d.$$

But the quotient is $1 + q^{r/3} + q^{2r/3} > 3$, congruent to 3 mod 9; it has both the prime divisor 3 and another prime divisor. This is impossible. Thus ξ^{ℓ^d} has full Frobenius orbit, proving `frobenius_conj_pow_injOn`. In particular scaling ξ by the three cubic roots of unity gives three distinct Frobenius orbits (raise a purported pair of conjugates to the third power). Each is primitive by the preceding paragraph with $M = 3$. Given two sets S, T each of cardinality at most one, some scaled label is outside $S \cup T$.

If charpoly(g) is reducible and $\tau_P(g)$ is nonzero for a primitive label P, it is a power Q^m with $m > 1$ by (26.2), so $3 \mid m$. The determinant is $((-1)^{\deg Q} Q(0))^m$, a cube in $k^\times$. Consequently every cubic twist of the primitive label has the same character value at g by (26.5), including when that value vanishes. It also has the same degree; hence its coefficient (26.6) at g is unchanged. For an original nonprimitive label take it unchanged (also if it is not a valid label). We have proved `exists_labelIdem_eq_off_irreducible` in the exact formulation allowing arbitrary polynomials, not only good ones. Together these calculations make the finite substitution step in Section 18.8 unconditional as a finite-group argument. The geometric transfer and the local–global hypothesis `Data.HasRigidLabels` are distinct assertions; this proof does not infer them from finite polynomial arithmetic.

26.5 An integral pivot calculation for the actual finite kernel

We now address the remaining error in the bottom form of Section 24.7. Work with the finite standard level of that chapter, over the semilocal algebra A with its product integer ring R. Write $h = \varrho$, $n = h+1$, K for the integral group, $C = K(g_*)$ with $g_* = \bar{g}_* \in R \cap A^\times$ for the principal level, U for the full Bessel unipotent, and $U_{<l}$ for the product of its first l Heisenberg factors N_a $(0 \leq a < l)$. We assume the standard-level basis, quadric, and Haar data of Chapter 24, not yet their assembly from all the global parameters. The first h positive frame coordinates are $e_0, \dots, e_{h-1}$, the middle block is V_h, and the remaining coordinates are the duals $e^*_{h-1}, \dots, e^*_0$. Let μ be the group Haar measure, ν the Haar measure of U, θ_l the successive floor measures, and $v = \nu(U \cap C) = \prod_{l<h} \mathrm{fl}_l(g_*)$, with the exact factors of (24.9). Here $\mathrm{fl}_l(b) = \mathrm{md}(b)^{\dim_A V_{l+1}+1/2}$ for a self-adjoint unit b. The finite character is self-dual for the Haar measure with R of mass one, so its one-coordinate Plancherel factor is one.

For a pair (x, xy) with nonzero $k(x,xy) = \langle W_Z(x), W_Z(xy)\rangle$, there are $u, v_0 \in U$ such that $a = ux$, $b = v_0 xy$ belong to K and $a^{-1}b \in C$. This is the common-coordinate support used in Proposition 24.4. Put $T_0 = v_0 u^{-1} \in U$. The pivot set at the l-th floor, denoted $I_l(x,y)$, is precisely the set `floorValues`:

$$\{w(e_l^*) : w(e_j^*) = 0 \ (j < l), x^{-1}w, \ g_*^{-1}(x^{-1}w - (xy)^{-1}w) \in R^r\}.$$

Since u is triangular, multiplying w by u preserves the earlier-zero conditions and the l-th coordinate on that subspace. The first integrality condition is then equivalent to $uw \in R^r$ because $a \in K$, and the second to $(T_0 - 1)uw \in g_* R^r$ because $a, b \in K$ and $b \equiv a \pmod{g_*}$. Hence $I_l(x,y)$ is the coordinate image of the lattice

$$\{z \in R^r : z(e_j^*) = 0 \ (j < l), \ (T_0 - 1)z \in g_* R^r\}.$$

It is a nonzero open ideal of R (it contains all sufficiently divisible elements times e_l^*). Each factor is a DVR, so write $I_l = c_l R$ with $c_l \in A^\times \cap R$. Thus $\mathrm{vol}(I_l) = \mathrm{md}(c_l)$, proving `exists_unit_mem_floorValues` in fact with equality of the volumes.

Here is a direct finite-field-factor substitute for the size-back-substitution argument of `FlatCharts.lean`. Let $\delta = \prod_{l<h} c_l$. Choose $z_l \in R^r$ in the indicated lattice with pivot c_l and preceding dual coordinates zero. Its dual coordinate block forms the l-th column of a triangular matrix P of determinant δ. Write the upper-unitriangular block form of T_0 (the duals can be ordered opposite to the positives):

$$T_0 = \begin{pmatrix} A_0 & B_0 & D_0 \\ 0 & 1 & D_1 \\ 0 & 0 & A_1 \end{pmatrix}.$$

The first and last blocks A_0, A_1 are unitriangular in opposite orders, and are inverse adjoints under the pairing of positive and dual blocks. The middle Gram matrix J_M is integral unimodular. The last two block rows of $(T_0 - 1)z_l$ imply

$$(A_1 - 1)P,\ D_1 P \in g_* M(R), \qquad \delta(A_1 - 1),\ \delta D_1 \in g_* M(R)$$

by adjugates of P. To avoid conjugation of δ in the bounds, put $\Delta = \delta\bar{\delta}$. This only improves integrality. Then $A_0^{-1} - 1$ has entries in $g_*\Delta^{-1}R$, by the inverse-adjoint relation. Expanding the inverse of the unitriangular block as the finite geometric series in its strictly triangular nilpotent part (nilpotence index $\le h$) gives $A_0 - 1 \in g_*\Delta^{-h}M(R)$. Orthogonality of the middle and last block columns gives $B_0 = -A_0 D_1^* J_M$ in matched dual coordinates, hence $B_0 \in g_*\Delta^{-(h+1)}M(R)$. The first row of $(T_0 - 1)z_l$, followed by the adjugate of P, now shows

$$T_0 - 1 \in g_*\Delta^{-(h+2)}M(R). \tag{26.8}$$

The inverse is the adjoint with the unimodular integral Gram matrix, so the same bound holds for $T_0^{-1} - 1$. Since $y = a^{-1}T_0^{-1}b$ and $b - a \in g_* M(R)$, it also holds with $y - 1$ in place of $T_0 - 1$. This proof is factorwise over R and hence applies to split pairs as well as unramified quadratic factors; no assumption that c_l itself is self-adjoint is needed.

Let $\mathfrak{a}_y = \{d \in R : d(y-1)/g_* \in M(R)\}$ and let $d_y = \mathrm{vol}(\mathfrak{a}_y)$. This is a nonzero integral principal ideal with $0 < d_y \le 1$. By (26.8), $\Delta^{h+2}R \subseteq \mathfrak{a}_y$, so

$$\min_{l<h} \mathrm{vol}\, I_l(x, y) \le d_y^{1/(2h(h+2))}. \tag{26.9}$$

If $y \notin C$, at least one I_l is a proper ideal: otherwise one can use δ a unit in (26.8) to get $(y-1)/g_* \in M(R)$, and unimodularity makes y an integral isometry congruent to one. Every proper ideal of R has mass at most $\epsilon = \mathrm{vol}(R \setminus R^\times)$ and at most $1/2$. Thus in addition the minimum in (26.9) is at most $e = \min(\epsilon, 1/2)$. These caps hold for every nonzero elementary pair, uniformly in x. The case $h = 0$ needs no cap: $U = 1$ and $\phi = 1_C$, so the bottom integral has no error outside C.

26.6 Weighted square estimates at a finite standard level

We give the estimates needed for the bottom error, including the shell integrability, separately from the general unequal-head theorem. This permits using the ultrametric size rule throughout; no real-place inverse-weight assertion is needed for this particular application. Put $a = |\iota|$ and $s = \mathrm{volumeExponent} = a - (3n + m + 2)$ as in Section 24.2; the standard rank relation

$k+1 = a+2(m+1)+2$ and $m < 9n$ give $s \geq k - 40n$. For a self-adjoint $\alpha \in R \cap A^\times$ let $B_i = \alpha^{-i}R$, $\Lambda_\alpha = 2/\operatorname{md}(\alpha)$, and

$$\Omega_\alpha = \{y \in \text{positions} : \alpha(y-1)/g_* \in M(R)\}.$$

Use $P_l(x,y) = \prod_{a'<l} \operatorname{vol}(I_{a'}(x,y))^s$ with $P_0 = 1$. Even without a nonzero pairing the pivot sets are images of intersections of lattices, hence have positive finite Haar mass; all are measurable. For a fixed y set

$$k_l(x,xy) = \int k(\sigma x, \sigma xy)\, d(\theta_{h+1} * \cdots * \theta_l)(\sigma)$$

and denote by $A_l(x) = \int |k_{l+1}(\sigma x, \sigma xy)|\, d\theta_l(\sigma)$ the pivot modulus. These are the actual `upperPairing` and `pivotModulus`, not kernels on newly normalized quotients. We have $|k_l(x,xy)| \leq A_l(x)$. Their bounded absolute integrals at each fixed pair follow by the same floor mass induction as the group-mass argument: each nonzero $W_Z(z)$ has its U-coset meeting K, and the mass of a floor with σz near at the next stage is bounded by the integral norm-quadric mass $\zeta_R(m) \leq 2$. At the final stabilizer the mass is one by the standard-level property. Thus all uses of successive absolute integrals are legitimate, as are Tonelli integrals of their nonnegative majorants.

For $l \leq h$ let

$$B_l(x) = \int \left(P_l(\sigma x, y)|k_{l+1}(\sigma x, \sigma xy)|^2 + B_{l+1}(\sigma x)\right) d\theta_l(\sigma), \qquad B_{h+1} = 0.$$

The source's self-norm $Z_0(y) = B_0(1)$ is the sum over $l = 0, \ldots, h$ of the integrals of $P_l(x,y)|k_{l+1}(x,xy)|^2$ against $\lambda_{l+1} = \theta_l * \cdots * \theta_0$. The floor values are invariant under simultaneous left multiplication of both vectors by the flag group $Q_{a'+1}$: change the defining vector by that matrix, which preserves the preceding-zero constraints and the a'-th coordinate. In particular for $\sigma \in H_l$ the factors of $P_l(\sigma x, y)$ equal those of $P_l(x,y)$. Also the absolute value of $k_{l+1}(x,z)$ is invariant under independent left multiplication on its two arguments by $U_{<l}$, since conjugation by its integration group H_{l+1} preserves the earlier simple-root sums; the full phase is the quotient of the two respective Bessel characters. All these invariances follow from the upper-triangular block description and the simple-root character of W_Z. They also imply that k_{l+1} is zero unless both arguments are near at $l+1$, i.e. their respective Q_{l+1}-cosets meet K.

One-floor inverse volume. If x is near at l and σx is near at $l+1$, $l < h$, then after putting x in K by its Q_l-action, the orbit point $\sigma^{-1}e_l$ is integral: write σx near using a matrix in Q_{l+1}, take the inverse frame column and discard the earlier positive coordinates, which are fixed by H_l. The movement by $Q_l = U_{<l} \rtimes H_l$ preserves the floor measure on its left quotient and the preceding pivot flags. For the reference $x \in K$ and position y, choose the space $\mathcal{W}$ of Section 24.2, with test matrix zero. The projected vectors $\sigma E w$ for $w \in \mathcal{W}$ are eligible vectors in the definition of $I_l(\sigma x, y)$, since $(\sigma x)^{-1}\sigma E w = x^{-1}Ew$ and the preceding dual coordinates vanish. Their e_l^* coordinates are $h(\sigma^{-1}e_l, Ew)$. Hence

$$\operatorname{vol}(I_l(\sigma x, y))^{-1} \leq f(E^* J \sigma^{-1} e_l)$$

for the common-volume majorant f proved in that section, with $\int_{R^a} f^s \leq \zeta_R(m+5) \leq 2$. Integrating over the integral norm quadric on the floor uses the orbit measure of Section 24.4; fixing the orthonormal coordinates and bounding the hyperbolic fiber gives a further factor at most $\zeta_R(m) \leq 2$ by its cone computation. If x is not near at l, the near subset of the l-th floor is empty. We thus have uniformly

$$\int_{\sigma x \text{ near at } l+1} \operatorname{vol}(I_l(\sigma x, y))^{-s}\, d\theta_l(\sigma) \leq 4. \tag{26.10}$$

This is a finite version of `floorMassLe_floorWeight_div`, with constant no larger than the source's $\Lambda^{10^4 n}$.

Capped pivot. Suppose that, for every elementary nonzero pair $k(\sigma x, \sigma x y)$ with $\sigma \in H_l$, at least one $a' \in [l, h)$ has $\mathrm{vol}(I_{a'}(\sigma x, y)) \le b$. Then

$$A_l(x)^2 \le b^s 4^{h-l}\, B_l(x)/P_l(x,y). \tag{26.11}$$

Indeed for $l = h$ the hypothesis says no such pair is nonzero and the left side vanishes. Otherwise use weighted Cauchy–Schwarz and (26.10):

$$A_l(x)^2 \le 4 \int \mathrm{vol}(I_l(\sigma x, y))^s |k_{l+1}(\sigma x, \sigma x y)|^2 \, d\theta_l(\sigma).$$

For σ where the pivot mass is $\le b$, the integrand is at most $b^s |k_{l+1}|^2$. Else the original hypothesis forces a cap at a deeper floor for every nonzero pair in the upper integration: I_l is invariant under H_{l+1}. Apply the induction hypothesis to $|k_{l+1}| \le A_{l+1}$ and cancel $\mathrm{vol}(I_l)^s$ against the relation $P_{l+1} = P_l \,\mathrm{vol}(I_l)^s$. Combining both contributions proves (26.11), precisely the weighted recurrence of `pivotModulus_sq_le_of_capped` at a finite standard level.

Integrating the direct weights on the finite Heisenberg tower. For any x near at $j \le h$ and any y_0, set $y(u) = x^{-1} u y_0$ for $u \in U_{<j}$. With $d\nu_{<j} = d\nu_0 * \cdots * d\nu_{j-1}$ we claim

$$\int_{U_{<j}} \mathbf{1}_{\Omega_\alpha}(y(u)) P_j(x, y(u))\, d\nu_{<j}(u) \le \prod_{l<j} \mathrm{fl}_l(g_*)\, \Lambda_\alpha^{4600nj}. \tag{26.12}$$

Putting x in K by the Q_j-action is harmless: $Q_j = U_{<j} \rtimes H_j$ normalizes $U_{<j}$ and preserves its measure, and the weights preceding j are invariant. For the measure statement, conjugation by the middle isometry has modulus one on each vector block and fixes each skew coordinate; conjugation by $U_{<j}$ is inner in a unimodular unipotent group. If the integrand is nonzero somewhere, translate y_0 by that chosen $u_0 \in U_{<j}$, and replace u by $u u_0^{-1}$. We can assume $x_0 = x^{-1} y_0 \in \Omega_\alpha$. Whenever $y(u) \in \Omega_\alpha$,

$$\frac{u-1}{g_*} = x \,\frac{y(u) - x_0}{g_*}\, x_0^{-1} x^{-1} \;\in\; \alpha^{-2} M(R).$$

Here x and its inverse are integral and x_0^{-1} has entries in $\alpha^{-1}R$ by the integral adjoint formula (using $\alpha = \bar\alpha$).

Prove (26.12) by induction, writing the last factor as $u = u' N_l(v_1, \beta)$ with $l = j-1$, $u' \in U_{<l}$, $v_1 \in V_{l+1}$ and $\bar\beta = -\beta$. The Heisenberg matrix on the l-th hyperbolic pair and its middle has middle-to-dual column v_1, positive-to-middle row $-v_1^* J_{l+1}$, and positive-to-dual entry $\beta - \tau h(v_1, v_1)$ with $\tau + \bar\tau = 1$. Its product with u' has this same V_l block (all earlier factors act as identity there on the quotient by the earlier positive frame). Consequently v_1/g_* has entries in $\alpha^{-2}R$, and for each fixed v_1 the possible β lie in a translate of $g_*\alpha^{-2}R$ intersected with the skew scalars. The latter has mass at most $\mathrm{md}(g_*)^{1/2}\Lambda_\alpha$.

To bound the last weight independently of u', take $\mathcal{W}$ from the gauge common-volume condition for (x, x_0) with test matrix zero. For $c \in I_l(x, y(u))$ choose the defining vector z with earlier dual coordinates zero and l-th coordinate c. For $w \in \mathcal{W}$ the two pairs

$$(x^{-1}z, (xy(u))^{-1} z), \qquad (x^{-1}Ew, y_0^{-1}Ew)$$

are close. The difference of their two Hermitian pairings, divided by g_*, is

$$g_*^{-1}\big(h(z, Ew) - h(N_l(v_1,\beta)^{-1} u'^{-1} z, Ew)\big) = g_*^{-1} \bar c\, h(v_1, Ew).$$

Indeed the earlier dual coordinates of z vanish, so u'^{-1} leaves its projection to V_l unchanged; the displayed Heisenberg block makes the change in the middle orthonormal coordinates exactly $-cv_1$ (projected to those coordinates). The second common-volume inequality of Section 24.2, applied with $E^*J(v_1/g_*)$ as the test vector, now gives

$$\operatorname{vol} I_l(x, y(u)) \le f(E^*J(v_1/g_*)).$$

Perform the u' integral using the induction hypothesis with $N_l(v_1, \beta)y_0$ in place of y_0, because this last factor bound is independent of u'. Integrate the remaining bound over $v_1/g_* \in (\alpha^{-2}R)^{\dim V_{l+1}}$. The orthonormal part of the integral basis has coordinate map E^*J; the complementary rank is at most $2h + 2(m+1) + 2 \le 24n$. Its coordinate measures on this box cost at most Λ_α^{48n}. The integral of f^s in the orthonormal coordinates costs at most $\Lambda_\alpha^{99n(2+1)}$ by the common-volume computation; scaling the vector back by g_* gains $\mathrm{md}(g_*)^{\dim V_{l+1}}$. Together with the skew mass the total is at most

$$\mathrm{fl}_l(g_*)\Lambda_\alpha^{48n+297n+1} \le \mathrm{fl}_l(g_*)\Lambda_\alpha^{4600n}.$$

This proves (26.12), the finite direct-weight integration assertion `lintegral_floorWeight_le`. It does not bound arbitrary inverse weights on unequal heads.

The integrated self-norm and its one-step Parseval estimate. We prove the finite estimate

$$\int_{\Omega_\alpha} Z_0(y)\, d\mu(y) \le n\mathcal{N} \prod_{l<h} \mathrm{fl}_l(g_*)\, \Lambda_\alpha^{4600nh}, \tag{26.13}$$

where $\mathcal{N} = \iint |k(x,z)|^2 d\eta(x) d\eta(z)$ and $\eta = \theta_{h+1} * \cdots * \theta_0$ as in Proposition 24.4. The proof works with nonnegative majorants, so before the resulting bound all square-integrals may take value infinity. Set

$$\rho_l = (\nu_l * \cdots * \nu_{h-1}) * \eta, \qquad J_l(x,z) = \int \mathbf{1}_{\Omega_\alpha}(x^{-1}uz)\, d\nu_{<l}(u),$$
$$\Phi_l(x,z) = \mathbf{1}_{\{J_l>0\}} |k_{l+1}(x,z)|^2.$$

The empty convolution $\nu_{<0}$ is unit mass. The Haar factorization of Section 24.4 gives $\mu = \nu_{<l} * \rho_l$ and $\rho_l = \nu_l * \rho_{l+1}$ for $l < h$. The push-forward of ρ_l to the left quotient by $U_{<l}$ is the quotient measure for μ with Haar measure $\nu_{<l}$. Indeed the factorization equality tested against compact functions is its defining quotient integral. It is invariant under left multiplication by H_l, which normalizes $U_{<l}$ with modulus one (same coordinate determinant calculation as above). Thus for any $U_{<l}$-left-invariant nonnegative function of z we can transform the z-variable under H_l in its ρ_l integral. Both the condition $J_l > 0$ and the kernel absolute value have that invariance under $U_{<l}$ in each argument.

Changing y to $x^{-1}z$ and using $\mu = \nu_{<l} * \rho_l$, the l-th contribution to $\int_{\Omega_\alpha} Z_0$ is bounded by

$$\prod_{a'<l} \left(\mathrm{fl}_{a'}(g_*)\Lambda_\alpha^{4600n} \right) \iint \Phi_l(x,z)\, d\rho_l(z)\, d\lambda_{l+1}(x), \tag{26.14}$$

by (26.12). We claim the one-step estimate, for $l < h$,

$$\int \Phi_l(x, Nz)\, d\nu_l(N) \le \mathrm{fl}_l(g_*)\Lambda_\alpha^{4600n} \int \Phi_{l+1}(\sigma x, \sigma z)\, d\theta_{l+1}(\sigma). \tag{26.15}$$

Here are the explicit support and Fourier details of this step. Let S consist of N on the left with nonzero integrand. If it is empty the inequality holds. Otherwise choose a center

$N_0 \in S$, and for each $N \in S$ choose $u \in U_{<l}$ with $y_N = x^{-1}uNz \in \Omega_\alpha$ (and a reference u_0 for N_0). The nonzero upper pairing implies x is near at $l+1$, so choose $\xi = tq \in Q_{l+1}$ with $\xi x \in K$, $t \in H_{l+1}$, $q \in U_{<l+1}$. Then

$$(\xi x) y_{N_0} y_N^{-1} (\xi x)^{-1} = \xi u_0 N_0 N^{-1} u^{-1} \xi^{-1} \in 1 + g_* \alpha^{-2} M(R).$$

On the V_l quotient block, the earlier factors u_0, u disappear and q acts as a Heisenberg element of N_l. Conjugation by that element changes only the central skew coordinate in the Heisenberg differences. Writing v_N, v_0 for the middle coordinates of N, N_0, the middle-to-dual column gives $t(v_N - v_0)/g_* \in \alpha^{-2} R^{\dim V_{l+1}}$. For fixed v_N the skew coordinates lie in a translate of $g_* \alpha^{-2} R \cap A^{\mathrm{skew}}$ (difference of two positive-to-dual entries); their mass is at most $\mathrm{md}(g_*)^{1/2} \Lambda_\alpha$. Both statements hold regardless of the size of the center itself.

Conjugation of $N_l(v_N, \beta)$ by $\sigma \in H_{l+1}$ sends v_N to σv_N and does not change β. Only its simple root in the l-th position contributes to the Bessel character, with entry $-h(v_N, \sigma^{-1} f_{l+2})$ (the formula `simpleRootSum_conj_heisenbergElem` follows directly from the row $-v_N^* J_{l+1}$). The covariance of the upper pairing therefore gives

$$k_{l+1}(x, Nz) = \int \psi(-g_*^{-1} h(v_N, \sigma^{-1} f_{l+2})) k_{l+2}(\sigma x, \sigma z)\, d\theta_{l+1}(\sigma).$$

Here f_{l+2} is the next frame vector in one-based notation, possibly the last norm vector. The phase is independent of the skew variable. The source amplitude is a function of the orbit point $\sigma^{-1} f_{l+2}$. Indeed the stabilizer inside H_{l+1} is the next Heisenberg radical and H_{l+2} (at the last norm floor just the norm-vector stabilizer). Simultaneous multiplication by the former cancels the two Bessel phases, by the conjugation formula for the preceding root entry. For the latter, the upper measures $\theta_{h+1} * \cdots * \theta_{l+2}$ have precisely the quotient normalization by the remaining unipotent in H_{l+2}, by the same Haar factorization as in Section 24.4 starting at floor $l+2$. The elementary pairing is invariant under simultaneous left multiplication by that unipotent, and right invariance of quotient integration thus makes its upper average invariant under simultaneous multiplication by H_{l+2}. This proves the factorization with unchanged square integral. Its supported orbit point after multiplication by t is integral. In detail choose $\zeta \in Q_{l+2}$ with $\zeta \sigma x \in K$ whenever the amplitude is nonzero. The vector $(\xi x)(\zeta \sigma x)^{-1} f_{l+2}$ is integral. Its V_{l+1} projection is $t\sigma^{-1} f_{l+2}$: ζ^{-1} adds only earlier positive frame columns to f_{l+2}, σ fixes them, and the earlier unipotent q leaves the middle quotient unchanged. This proves the requisite support in the source's size 7 (in fact integral).

After the change $v = t(v_N - v_0)/g_*$ the Fourier frequency is a unit of modulus one, because g_* is self-adjoint and t is an isometry. The center gives a unit-modulus factor multiplying the source amplitude only. Apply Proposition 23.7 on this floor, with receiver volume and source quadric. For a size i receiver or source the fiber constant is at most $\Lambda_\alpha^{99n(i+6)}$ by the hyperbolic-cone gauge bound and the integral basis coordinate change; for the present source size ≤ 7 and receiver size ≤ 14 their product is at most $\Lambda_\alpha^{99n(13+20)}$. The Fourier coefficient contributes no extra scalar since the one-coordinate Plancherel constant is one. Returning from v to v_N multiplies by $\mathrm{md}(g_*)^{\dim V_{l+1}}$; the skew fiber contributes at most $\mathrm{md}(g_*)^{1/2} \Lambda_\alpha$. Thus the required factor is at most $\mathrm{fl}_l(g_*) \Lambda_\alpha^{4600n}$. Finally for every $\sigma \in H_{l+1}$ the indicator in $\Phi_{l+1}(\sigma x, \sigma z)$ is one: some N_0 and u_0 as above witness it via $\sigma u_0 N_0 \sigma^{-1} \in U_{<l+1}$. Nonemptiness implies positive Haar mass because the positions are a union of right cosets of the open group C, hence open, and the integral displacement condition is open as well. Consequently inserting the indicator does not alter the right-side square integral. This proves (26.15); its proof is the self-pairing instance of `parseval_heisenbergMeasure_of_box`, rather than an appeal to an unweighted Fourier transform of a noncompact group.

By the quotient invariance above, integrating (26.15) against $\rho_{l+1}(z) \lambda_{l+1}(x)$ bounds the double integral in (26.14) by the same integral at $l+1$ times $\mathrm{fl}_l(g_*) \Lambda_\alpha^{4600n}$: absorb the

simultaneous left multiplication by σ in the z quotient, and use $\lambda_{l+2} = \theta_{l+1} * \lambda_{l+1}$ in x. At $l = h$ the remaining average is over the last stabilizer H_{h+1} with integral mass one on each relevant near set (the defining μ_H property). Cauchy–Schwarz on that support gives $|k_{h+1}(x,z)|^2 \le \int |k(ax,az)|^2 d\mu_H(a)$. Using its quotient invariance and $\eta = \mu_H * \lambda_{h+1}$ bounds the terminal double integral by $\mathcal{N}$. Thus each of the n summands (26.14) has the same bound; summing proves (26.13). By Proposition 24.4, its right side is further bounded by

$$n\, m_Z\, \Lambda_\alpha^{4600nh}, \qquad m_Z = \mu(Z)\mu(C)/v^2. \tag{26.16}$$

26.7 Closing the finite bottom error

Theorem 26.4 (Bottom form at a finite standard level)**.** *In the notation of Section 24.7, for $t \ge 0$, $t + 10^{10}n^4 \le k$ one has*

$$B(Z) \le \mu(Z)\mu(C)\big(\mu(Z) + \epsilon^{t+1}2^{-n}\big), \qquad \epsilon = \mathrm{vol}(R \setminus R^\times).$$

Proof. The contribution of C is exactly $\mu(Z)^2\mu(C)$ by (24.11). If $h = 0$, $\phi = \mathbf{1}_C$ and there is no contribution outside C. Suppose $h \ge 1$. The support of $k_0(1,y)$ is contained in the positions, since every nonzero elementary pair gives $y = a^{-1}T_0^{-1}b$ with $T_0 \in U$, $a^{-1}b \in C$. By (26.9) and (26.11), at every position outside C,

$$|k_0(1,y)|^2 \le 4^h \min\big(e, d_y^{1/(2h(h+2))}\big)^s Z_0(y), \quad e = \min(\epsilon, 1/2).$$

The finite-window algebra with no factors, if allowed, is trivial and gives zero error; otherwise $\epsilon > 0$. Partition the positions by their denominator ideals $\mathfrak{a}_y = \mathfrak{a}$ with volume $d \in (0,1]$. Each integral ideal is principal in the product integer ring, choose a generator $c_\mathfrak{a}$ and put $\alpha = c_\mathfrak{a}\bar{c}_\mathfrak{a}$. Every position in the shell belongs to Ω_α because $\alpha \in \mathfrak{a}$, and $\Lambda_\alpha = 2d^{-2}$. Put $K_0 = 4600nh$ and $D = s - (t+1)$, so $D \ge 10^{10}n^4 - 40n - 1$. Choosing $R_0 = K_0 + 4n + 10$, we have $D > R_0$ and

$$\min(e, d^{1/(2h(h+2))})^s \le e^{t+1+R_0}\, d^{(D-R_0)/(2h(h+2))}.$$

Thus (26.16) gives

$$v^2 \int_{G\setminus C} |k_0(1,y)|^2\, d\mu(y) \le v^2 m_Z\, n\, 4^h 2^{K_0}\, e^{t+1+R_0} \sum_\mathfrak{a} \mathrm{vol}(\mathfrak{a})^{(D-R_0)/(2h(h+2))-2K_0}.$$

This sum includes even the ideal R harmlessly. Its exponent is at least m: use $2h(h+2) \le 2n^2$, $K_0 \le 4600n^2$, $m < 9n$, and the displayed bound on D. The sum over all integral ideals is the elementary geometric product ζ_R at that exponent, at most $\zeta_R(m) \le 2$ by the standard-level zeta-rank condition. Since $v^2 m_Z = \mu(Z)\mu(C)$ and $e^{R_0} \le 2^{-R_0}$, the error is at most

$$\mu(Z)\mu(C)\,\epsilon^{t+1} \cdot n\, 2^{2h+K_0+1-R_0} \le \mu(Z)\mu(C)\,\epsilon^{t+1}2^{-n}.$$

The last inequality follows from $h = n - 1$ and $n \le 2^n$. The identity $vk_0(1,y) = \int_Z \phi(zyz^{-1})\, d\mu(z)$ from Lemma 24.3 gives the assertion. □

This is the exact estimate `floorZeroForm_le_main_term` in `TopFloor.lean`, including the error (24.13). In particular the fixed-vector subgroup bottom savings already deduced from that error in Section 24.7 now hold for the finite standard level. This proof used the cap argument and the direct self-pairing Plancherel at finite places. It did not use the general `lintegral_shells_le` assertion on unequal-head form majorants and their oscillatory kernels.

26.8 Status of the continued finite interfaces

- **B03 — proved.** The exact source `labelChar_degree_bounds`, `bad_share_le`, primitive scaling, cubic avoidance and reflection annihilation are derived in Sections 26.1–26.4. The only classical character input used is the precise Green Theorem 1 on virtual symmetric-power-sum characters; the actual identification, degree, and constants have proofs here.
- **B02 — still conditional.** The label arithmetic, representation and correction operations are now justified, but the common attached Galois representation, exact local killing polynomials and determinant, and parahoric transfer in `New/Trace/LabelRigidity.lean` must be established as actual global input. No assertion of arbitrary common spectra follows from our finite proof.
- **B05 / B13 — finite bottom and self-pairing estimates proved for the standard level.** Sections 26.5–26.7 give the exact finite pivot ideals, a quantitative denominator bound, inverse one-floor volume, capped pivot, direct-weight integration, self-pairing Parseval and integrated self-norm, and the previously missing `floorZeroForm_le_main_term`. The actual global construction of every finite standard-level datum, the full arbitrary-head Fourier estimates (including inverse-weight integration and unequal-head Parseval) and the rest of `exists_finite_sieveSystem` are not inferred from these calculations. The broader B05 and B13 interfaces remain conditional on those.
- **B19 — conditional.** Some abstract recurrence and weight mechanisms are now transparent, but the ultrametric direct-weight and denominator proofs here are finite-only. The real-place `lintegral_selfNorm_le`, inverse-weight, character and unequal-head band bounds are not established in this supplement.

Chapter 27

Rational frames, amplitude convergence, and pair discriminants

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 30, 29, 28 and the final audit in Chapter 57.

The normalization of the rational amplitude sums is useful only if the sums are absolutely convergent with a tail bound *uniform in the classes and the reduction.* Chapter 22 constructed the arithmetic cutoffs, while Chapter 23 retained absolute integrated-amplitude summability as a hypothesis of the packet limit. We now derive that summability, with the exact loss prescribed in `New/Filling/AmplitudeSums.lean`, from the local Bessel majorant and the elementary standing arithmetic size assumptions. The key is a packing bound on rational frame images, not a count of all elements of the unitary group. We keep the dependence on the actual analytic Bessel system explicit.

27.1 Packing the arithmetic frame boxes

Use the notation of Chapter 22: E is a CM field, $f = [E : \mathbb{Q}]/2$, h is the hyperbolic rank, $n = h + 1$, $k = \dim_E X_0$, $r = k + 2h + 1$, and the rational form is J_N with hyperbolic pairs e_a, e_a^* $(0 \le a < h)$, positive space X_0, and vector z with norm $-N$. Write

$$F = (e_0, \dots, e_{h-1}, z) = (f_0, \dots, f_h).$$

The normalized complex form is J_1; it satisfies $J_1^2 = 1$. For the embedding $\Phi_v : E \to \mathbb{C}$ chosen at the infinite place v, the diagonal change of basis from the normalized complex coordinates to the original ones has entries

$$s_{v,a} = \kappa_v^{h-a}/\sqrt{N} \quad (0 \le a \le h), \qquad s_{v,e_a^*} = s_{v,a}^{-1}, \qquad s_{v,X_0} = 1.$$

Here $s_{v,h} = 1/\sqrt{N}$ is the scale of z. Denote the corresponding map by $\gamma \mapsto \gamma_\infty$. Thus the frame Gram matrix in normalized coordinates is $Z_v(\gamma)_{ab} = s_{v,a} s_{v,b} \Phi_v(Z(\gamma)_{ab})$ with $Z(\gamma) = F^* J_N \gamma F$. The scales are real and may have signs, which do not affect the estimates.

We use the explicit hypotheses

$$b \ge 1000, \quad n \ge 10^5 + 1, \quad |\kappa_v| \in [1, b^{23}], \quad N \le b^{60n}. \tag{27.1}$$

Let $R = b^{2n^2}$, $C = b^{6n+2}$ and $\delta = 1/(NR)$. The box condition on $y \in G_\infty$ is

$$\max_{v,a} \|y_v^{-1} f_a\| \le R, \qquad \mathrm{bs}(y_v) := \inf_{\sigma \in S_B(\mathbb{C})} (\|\sigma y_v\|_{\mathrm{op}} + 1) \le C.$$

For clarity the norm of an array of frame vectors will always mean the maximum of the Euclidean norms of its individual vectors. The Bessel group $S_B = UH$ is the subgroup preserving the frame flag and acting by an upper unitriangular matrix on F; H acts only in X_0 and is compact at infinity. By removing the compact factor from an approximant to the defining infimum, the Bessel-size condition implies that some $u_v \in U(\mathbb{C})$ satisfies

$$\|u_v y_v\|_{\mathrm{op}} \le C. \tag{27.2}$$

Indeed the infimum of the norms plus one is at most C, so an approximant has norm plus one strictly less than $C+1$; left multiplication by the inverse of its H-factor is norm preserving.

Fix c, s, a class and a finite residue of the filling data. Recall the term sets $\mathcal{T}(c,s,d)$ and the rational congruence subgroup $\Gamma_d = U(E^+) \cap h_d K(g) h_d^{-1}$; its subgroup Γ_d^0 fixes every vector of F. The finite integral basis of a class is $B_d = \mathrm{diag}(1, B_{d,M}, 1)$ in the hyperbolic–middle–dual block order. In its middle block the marked change of basis replaces only the distinguished positive vector by $x_{k_0}/\bar{\pi} + z/N$, leaving z fixed; the definite-class part also fixes z. Consequently

$$B_d F = F, \quad B_d^{-1} F = F, \qquad (B_d^{-1} t)_{e_a} = t_{e_a}, \quad (B_d^{-1} t)_{e_a^*} = t_{e_a^*}. \tag{27.3}$$

Each finite class Gram matrix $B_d^* J_N B_d$ is integral and unimodular as recalled in Chapter 22. Term position in integral coordinates means $B_c^{-1} \gamma B_d \in K$ of reduction s. These identities are all in finite adeles. In particular, if $\eta \in \mathcal{T}(c,s,d)$ and $\eta' \in \mathcal{T}(c,s,d')$, then

$$B_d^{-1} \theta B_{d'} \in K(g), \qquad \theta = \eta^{-1}\eta'. \tag{27.4}$$

By (27.3), each entry of $Z(\theta) = F^* B_d^* J_N B_d (B_d^{-1}\theta B_{d'}) F$ is integral at every finite place. The rational top rows of θF are also integral by the first-row identity in (27.3). This observation is important: neither full rational matrix η nor the full matrix of θ need have coefficients in $\mathcal{O}_E$.

Lemma 27.1 (Quantitative separation of boxed frames)**.** *Fix an arbitrary $y \in G_\infty$ and terms η, η' as in (27.4). Suppose that $\eta_\infty^{-1} y$ and $\eta_\infty'^{-1} y$ are both boxed. If their frame arrays differ by norm strictly less than δ, then $d = d'$ and $\eta^{-1}\eta' \in \Gamma_d^0$.*

Proof. Put $Y = \eta_\infty^{-1} y$, so the two arrays at each place are $Y_v^{-1} F$ and $Y_v^{-1}\theta_v F$. The difference of their pairings with the first array under J_1 has each entry of absolute value less than $R\delta$ because $\|J_1\|_{\mathrm{op}} = 1$. The Gram is $Z_v(\theta) - Z_v(1)$; all $|s_{v,a} s_{v,b}| \ge 1/N$. Undoing the diagonal normalization proves

$$|\Phi_v((Z(\theta) - Z(1))_{ab})| < NR\delta = 1 \quad \text{for every } a, b, v.$$

This is an integral rational element by (27.4); it vanishes because otherwise its absolute field norm is an integer at least one, whereas the product of all conjugate absolute values is strictly less than one. Thus $Z(\theta) = Z(1)$.

Fix a column θf_b. Its pairings with F determine the dual hyperbolic coordinates and its z-coordinate. The equality of Gram matrices says that those coordinates equal those of f_b. Hence

$$\theta f_b - f_b = \sum_{a<h} t_{ab} e_a + x_b, \qquad x_b \in X_0.$$

Since θ is unitary, comparison of the norms of θf_b and f_b in the rational form gives $\langle x_b, x_b \rangle = 0$. The form on X_0 is positive definite at each infinite place, so $x_b = 0$. Also each $t_{ab} \in \mathcal{O}_E$ by the top-row assertion above.

Work in descending order of a and suppose all $t_{ib} = 0$ for $i > a$. For every place choose u_v as in (27.2). It is upper unitriangular on the isotropic frame; the e_a coordinate of

$$u_v(\theta_v f_b - f_b) = (u_v Y_v)(Y_v^{-1}(\theta_v f_b - f_b))$$

is exactly $\Phi_v(t_{ab}) s_{v,b}/s_{v,a}$: the higher rows are absent by induction. Therefore

$$|\Phi_v(t_{ab})| < C\delta\,|\kappa_v|^{b-a} \le b^{6n+2+23h}\delta < 1.$$

In the exponent of κ_v the column index in the exponent of κ_v lies in $\{0, \dots, h\}$; in particular its difference from a has absolute value at most h. The final inequality follows from $6n+2+23h < 2n^2$ and $N \ge 1$. The same norm argument gives $t_{ab} = 0$ at this step. This proves $\theta F = F$.

A unitary map fixing F lies in $U^0 H$ (use the block form of the frame stabilizer). In particular it belongs to S_B. By (27.4) it is a Bessel term over residue 1 from d to d'. The free-column-level argument of Section 22.10 implies that $d = d'$ and $\theta \in \Gamma_d$. Since $\theta F = F$, in fact $\theta \in \Gamma_d^0$. This is the implication `termsOver.eq_and_mem_classUnip0` in `New/Filling/NearTerms.lean`. □

Proposition 27.2 (Weighted frame count)**.** *Let P be a countable label set, d_p a class and γ_p a rational element, with $p \mapsto (d_p, \gamma_p)$ injective. Let $x \ge 0$ and $y, y' \in G_\infty$. Consider nonnegative weights w_p such that, whenever $w_p > 0$, there is a term $\eta_p \in \mathcal{T}(c, s, d_p)$ with $\gamma_p^{-1}\eta_p \in \Gamma_{d_p}$ and $\eta_{p,\infty}^{-1} y$ boxed, and*

$$w_p = x\beta_{d_p}(\gamma_{p,\infty}^{-1} y').$$

Here β_d is the arithmetic cutoff for Γ_d with periodization one. Suppose the differences of the array $\mathcal{F}_p = (y_v^{-1}\eta_{p,v} f_a)_{v,a}$ lie in a real subspace $\mathcal{V}$ of dimension D. Then

$$\sum_p w_p \le b^{6n^2 D}\, x. \tag{27.5}$$

Proof. Restrict to the support of the weights. Group the labels by equality of the frame arrays. By Lemma 27.1, different arrays are δ-separated. All arrays lie in an affine translate of $\mathcal{V}$ and have norm at most R. If the set is nonempty, choose one array as center. All points then lie in a ball of radius $2R$ in a normed real D-dimensional space. Open balls of radius $\delta/2$ about distinct points are disjoint. Lebesgue volume comparison for a finite collection bounds their number by

$$\left(\frac{2R + \delta/2}{\delta/2}\right)^D = (1 + 4NR^2)^D \le b^{6n^2 D},$$

since $1 + 4NR^2 \le 5b^{60n+4n^2} \le b^{6n^2}$. Taking the supremum over finite collections proves finiteness with this same bound; when $D = 0$ the number is at most one.

All labels in a fiber have the same class d and their terms differ by right multiplication by Γ_d^0 by the lemma. Take a representative term η_0 in the fiber. Since $\gamma_p^{-1}\eta_p \in \Gamma_d$ for every p, it follows that $\gamma_p = \eta_0 u_p$ with $u_p \in \Gamma_d$. The u_p are distinct by injectivity of the labels. Hence the total weight in this fiber is bounded by

$$x \sum_{u \in \Gamma_d} \beta_d(u_\infty^{-1}\eta_{0,\infty}^{-1} y') = x.$$

Multiply by the number of fibers. This is the exact exponent in `tsum_classCutoff_box_le` of `New/Filling/BoxPartition.lean`. In particular it counts possible *frames*, with total weight at most one for each, not all translations in an arithmetic unipotent subgroup. □

27.2 Uniform summability and the cut error

Let $W_v^{\mathrm{full}}, W_v^{\mathrm{cut}}$ be the local Bessel functions with Euclidean coordinate space of dimension d_v. The following is precisely the local analytic majorant property stipulated by `IsCutPair.exists_majorant` in `New/Filling/BesselSystem.lean`. We do not infer this property from the square-integrability of the individual functions. It is an actual quantitative input from the construction of the local Bessel system.

For every v, let $A_v \geq 0$ be measurable, $U(\mathbb{C})$-left invariant, and m_v a finite nonnegative measure on the normalized group, satisfying, for either side t,

$$\begin{aligned} \lambda_v(y)\|W_v^t(y)\| &\leq A_v(y), \\ A_v(y) &\leq m_v\{x : \mathrm{bs}(yx) \leq b^{6n+2}\}, \\ m_v(G_v)\int_{U(\mathbb{C})\backslash G_v} A_v\, d\dot{y} &\leq b^{2r^3} d_v, \end{aligned} \tag{27.6}$$

where $\lambda_v(y) = 1$ if $W_v^{\mathrm{full}}(y) = W_v^{\mathrm{cut}}(y)$ and $\lambda_v(y) = b^{3r^3 f}$ otherwise. The quotient measure is the one used for the Bessel functions in Chapter 22; ν is the covolume of Γ_c for the compatible unipotent Haar measure, independent of c. In addition to (27.1), use

$$k \geq 1000(2n)^9, \qquad d := \prod_v d_v \leq b^{2kf}. \tag{27.7}$$

The latter bound is the Bessel-coordinate cardinality in `Filling.IsAdmissible`; no assumption on the number of definite classes is needed in the argument below.

Put $A(y) = \prod_v A_v(y_v)$, $m = \bigotimes_v m_v$, and $M = m(G_\infty)$. The norms of the tensor products $W^t(y) = \bigotimes_v W_v^t(y_v)$ are the products of the local norms. Define

$$K_t(\gamma, y) = W^t(y) \otimes \overline{W^t(\gamma_\infty^{-1} y)}.$$

For arbitrary T_0 (it need not be a stable set), write $\mathcal{T}_d = T_0 \cap \mathcal{T}(c, s, d)$, using d as a class index in sums, and β_c for the normalized cutoff of Section 22.9. For $T_0 = \varnothing$ all assertions below are immediate.

Theorem 27.3 (Absolute convergence with explicit cut error)**.** *Under* (27.1), (27.6) *and* (27.7), *for every c, s, c' and either side t there is a finite constant, independent of y, bounding*

$$\sum_{\gamma \in \mathcal{T}_{c'}} \beta_{c'}(\gamma_\infty^{-1} y)\|W^t(\gamma_\infty^{-1} y)\|.$$

Furthermore the sum of absolute integrated amplitudes, even over all terminal classes, is finite:

$$\sum_{c'} \sum_{\gamma \in \mathcal{T}_{c'}} \int_{G_\infty} \beta_c(y)\beta_{c'}(\gamma_\infty^{-1} y)\|K_t(\gamma, y)\|\, dy \leq \nu b^{(12rn^3+2r^3)f} \prod_v d_v. \tag{27.8}$$

The exact cut loss needed for the filling-moment estimate is

$$\sum_{c'} \sum_{\gamma \in \mathcal{T}_{c'}} \int_{G_\infty} \beta_c(y)\beta_{c'}(\gamma_\infty^{-1} y)\|K_{\mathrm{full}}(\gamma, y) - K_{\mathrm{cut}}(\gamma, y)\|\, dy \leq \frac{\nu}{2} b^{-k^3 f}. \tag{27.9}$$

Proof. Fix y and $x \in G_\infty$. Consider the label pairs (c', γ) with $\gamma \in \mathcal{T}_{c'}$ and weights

$$w_{c',\gamma} = \beta_{c'}(\gamma_\infty^{-1} y)\, \mathbf{1}_{\{\forall v : \mathrm{bs}((\gamma_\infty^{-1} yx)_v) \leq b^{6n+2}\}}.$$

If the weight is nonzero, the periodization of $\beta_{c'}$ at $\gamma_\infty^{-1} yx$ has a nonzero term $\beta_{c'}(u_\infty \gamma_\infty^{-1} yx)$ with $u \in \Gamma_{c'}$. The Bessel size is invariant under left translation by $U(\mathbb{C})$; the cutoff support

property from Section 22.9 means that this translate is boxed. The term $\eta = \gamma u^{-1}$ is still in $\mathcal{T}(c, s, c')$, and $\gamma^{-1}\eta \in \Gamma_{c'}$. Here η need not be in T_0, since the weighted count requires only term membership, not membership of its witnesses in the original set.

Apply Proposition 27.2 with boxing variable yx, cutoff variable y, and scalar 1. Use the ambient real space of all frame arrays, of dimension $2frn$. It gives, uniformly in x, y,

$$\sum_{c',\gamma} w_{c',\gamma} \le b^{12frn^3} =: B.$$

Integrating against the product measure m and using the second inequality in (27.6), Tonelli gives

$$\sum_{c',\gamma} \beta_{c'}(\gamma_\infty^{-1} y) A(\gamma_\infty^{-1} y) \le MB. \tag{27.10}$$

In particular the first assertion follows from $\|W^t(z)\| \le A(z)$. If $M = 0$, all majorants vanish and the remaining assertions follow; otherwise each $m_v(G_v) > 0$.

Multiply (27.10) by $\beta_c(y)A(y)$ and integrate. Because A is invariant under the full real unipotent group, its quotient integral with the cutoff is

$$\int_{G_\infty} \beta_c(y)A(y)\,dy = \nu \int_{U_\infty \backslash G_\infty} A(y)\,d\dot{y}.$$

This follows by disintegrating along left U_∞-orbits: on each orbit A is constant and the integral of the cutoff is ν, since its periodization along the lattice Γ_c is one. Both sides are nonnegative, so no prior integrability is needed. The quotient measures and product factors yield

$$\begin{aligned}\sum_{c',\gamma} \int \beta_c(y)\beta_{c'}(\gamma_\infty^{-1} y)A(y)A(\gamma_\infty^{-1} y)\,dy &\le \nu B \prod_v \left(m_v(G_v) \int A_v\,d\dot{y}_v \right) \\ &\le \nu B\, b^{2r^3 f} \prod_v d_v.\end{aligned}$$

Since $\|K_t\| \le A(y)A(\gamma_\infty^{-1} y)$, this proves (27.8).

If the two kernels differ, some local full and cut functions fail to agree at either y_v or $(\gamma_\infty^{-1} y)_v$. At this argument the first inequality in (27.6) saves the factor $b^{-3r^3 f}$ for *both* functions. For each tensor kernel use this saving at that factor; bound the other factors by their majorants. The triangle inequality gives

$$\|K_{\mathrm{full}}(\gamma, y) - K_{\mathrm{cut}}(\gamma, y)\| \le 2b^{-3r^3 f} A(y)A(\gamma_\infty^{-1} y),$$

including the case when the left side vanishes. Integrating as above gives

$$2\nu b^{(12rn^3 - r^3)f} \prod_v d_v \le 2\nu b^{(12rn^3 + 2k - r^3)f}.$$

As $r = k + 2n - 1 \le 2k$, we have $r^3 - k^3 \ge 3k^2 n$. Now $k \ge 1000(2n)^9$ implies $k \ge 9n^2$, so

$$3k^2 n \ge 27kn^3 \ge 24kn^3 + 2k + 2 \ge 12rn^3 + 2k + 2.$$

Thus the last integral bound is $\le 2\nu b^{-k^3 f - 2f} \le \nu b^{-k^3 f}/2$, as asserted. □

The theorem proves the full global conclusion of `amplitudeAbs_lt_top_and_cut_le` and `summable_amplitude` in `New/Filling/AmplitudeSums.lean`, conditional on the explicitly stated local analytical majorant and arithmetic size hypotheses. These are not additional global convergence assumptions. In particular the packet integrability and limiting arguments

of Chapter 23 may now invoke Theorem 27.3 rather than an assumed summability of rational amplitudes, once the local Bessel system itself has been constructed.

Here is also the useful exact weight-change version. For any stable T_0, on the discrete-times-continuous space $\mathcal{T}_{c'} \times G_\infty$ the action of $\Gamma_c \times \Gamma_{c'}$ is

$$(u, v)(\gamma, y) = (u\gamma v^{-1}, u_\infty y).$$

It preserves counting measure times Haar measure, and each kernel K_t is invariant, since the Bessel character is trivial on the two principal congruence subgroups. Both the original nonnegative weight $b_0(\gamma, y) = \beta_c(y)\beta_{c'}(\gamma_\infty^{-1} y)$ and any other normalized nonnegative weight $b_1(\gamma, y)$ with orbit sum one give the same integral of a nonnegative invariant function F:

$$\int b_1 F = \int b_1 F \sum_{g \in \Gamma_c \times \Gamma_{c'}} b_0(g(\gamma, y)) = \int b_0 F \sum_g b_1(g^{-1}(\gamma, y)) = \int b_0 F.$$

The middle equality follows by Tonelli and the measure-preserving change of variables, with no assumption of a free action. Apply this to the norms of K_t and $K_{\mathrm{full}} - K_{\mathrm{cut}}$. The preceding estimates then imply absolute integrability and equality for the corresponding vector-valued integrals; individual amplitudes are summable and the cut error retains exactly (27.9). This is `hasSum_amplitudeSum_change_of_weight`, including its quantitative assertion. The orbital cancellation and transport identities of Section 22.10 are justified by this argument for the stipulated local systems.

27.3 The determinant of a regularized orthogonal complement

We require an index identity valid for an indefinite or even degenerate subspace of a unimodular lattice. It is stronger than the positive-definite complementary-height formula of Section 22.13. We write $A = \mathbb{A}_{E,f}$, $\mathcal{O}_f = \prod_{w<\infty} \mathcal{O}_{E_w}$, and

$$[x] = \prod_{w<\infty} |x_w|_w^{-1} \quad (x \in A^\times).$$

In particular, $[x] = |\operatorname{Norm}_{E/\mathbb{Q}}(x)|$ on nonzero rational scalars. For an adelic invertible matrix a, $[\det a]$ is the factor by which it multiplies the covolume of an $\mathcal{O}_f$-lattice. We use $[x^*] = [x]$, since conjugation permutes the places, and $[\det a] = 1$ for $a \in \mathrm{GL}_q(\mathcal{O}_f)$.

An adelic matrix of full column rank can be made primitive by choosing a basis for the lattice of coefficients which it maps into $\mathcal{O}_f^m$: at every finite place use the elementary divisor theorem over the discrete valuation ring. Here primitive means that its columns are an integral basis for the intersection of their adelic span with $\mathcal{O}_f^m$, or equivalently that they extend to an integral invertible basis. At almost all places every given rational matrix of full rank is already primitive, so all changes of basis may be chosen to be identities there. We also need simultaneous primitive bases adapted to the first block of a matrix. If $(U_0 \,|\, U_1)$ has full column rank, the coefficient lattice of its integral points has the direct-summand sublattice lying in the first coordinate block. Its intersection with that block is saturated, since the quotient injects into the last coordinates. Choose a basis of the first part and lift a basis of the projected lattice in the last coordinates. This constructs a change of basis

$$a = \begin{pmatrix} a_0 & z \\ 0 & a_1 \end{pmatrix}, \qquad a_i \in \mathrm{GL}(A),$$

such that $(U_0 \,|\, U_1)a$ is primitive. The construction also proves that $U_0 a_0$ is primitive in its own span. These statements give precisely the block-preserving primitive-basis assertion `exists_isPrimitiveMatrix_mul_fromBlocks` in `Lib/General/Algebra/LatticeComplement.lean`.

Lemma 27.4 (Regularized complementary determinant)**.** *Let J be an invertible Hermitian m-by-m matrix over $\mathcal{O}_f$ with integral inverse. Let F_0 be an integral matrix with s columns of full rank, and L a primitive matrix of $m-s$ columns with $F_0^* J L = 0$. Let P be any matrix with q columns and $c \in A^\times$. Define*

$$Q = J + c^{-1} J P P^* J, \qquad M = \begin{pmatrix} F_0^* J F_0 & F_0^* J P \\ P^* J F_0 & P^* J P + c I_q \end{pmatrix}.$$

Assume $\det M$ *is invertible in A. If i denotes the index of $F_0 \mathcal{O}_f^s$ in its saturation in $\mathcal{O}_f^m$, then*

$$[\det(L^* Q L)]\, i^2 [c]^q = [\det M]. \tag{27.11}$$

Moreover the orthogonal complement of $\operatorname{span}_A(L)$ *equals* $\operatorname{span}_A(F_0)$*.*

Proof. Choose $a \in \mathrm{GL}_s(A)$ such that $F_1 = F_0 a$ is primitive. Then $i = [\det a]^{-1}$ and the corresponding regularized matrix M_1 is obtained from M by multiplying its first block of columns by a and its first block of rows by a^*. So $[\det M_1] = [\det a]^2 [\det M]$.

Because F_1 extends to an integral basis, $F_1^* J : \mathcal{O}_f^m \to \mathcal{O}_f^s$ is surjective. Choose integral columns B with $F_1^* J B = I_s$. The kernel is exactly $L\mathcal{O}_f^{m-s}$: their adelic spans agree by rank, and L is primitive. Therefore $(B \mid L)$ is integral invertible. This rank argument also proves the last assertion of the lemma.

For the determinant compare the two eliminations of the block matrix

$$\mathcal{M} = \begin{pmatrix} J^{-1} & F_1 & P \\ F_1^* & 0 & 0 \\ P^* & 0 & -c I_q \end{pmatrix}.$$

Eliminating its first block by the Schur complement gives $\det \mathcal{M} = (-1)^{s+q} \det(J)^{-1} \det M_1$. Eliminating the bottom block instead gives $(-1)^q c^q \det \mathcal{B}$, where

$$\mathcal{B} = \begin{pmatrix} J^{-1} + c^{-1} P P^* & F_1 \\ F_1^* & 0 \end{pmatrix}.$$

Conjugate the latter matrix (in the sense of congruence) first by $\operatorname{diag}(J, 1)$, then by $\operatorname{diag}((B \mid L), 1)$. Neither operation changes the finite determinant index, because the matrices and their inverses are integral. The result has the block form

$$\begin{pmatrix} * & * & I_s \\ * & L^* Q L & 0 \\ I_s & 0 & 0 \end{pmatrix}.$$

Expanding on its last s rows and columns gives determinant $(-1)^s \det(L^* Q L)$. Comparing the two formulas for $\det \mathcal{M}$ yields $[\det M_1] = [c]^q [\det(L^* Q L)]$. This also shows the last determinant is invertible. Substitute the earlier expression for $[\det M_1]$, and obtain (27.11). □

This is the exact determinant identity `Filling.jacobi_perp_discriminant` in `LatticeComplement.lean`, with the full-rank condition already available for the frame matrices below. The proof uses only elementary determinant operations and the existence of a primitive basis; neither definiteness nor nondegeneracy of the form restricted to the initial columns is needed.

27.4 The upper bound for integral link lifts

Now fix $\gamma \in \mathcal{T}(c, s, c')$ of level $j \in \{0, \dots, h+1\}$, in the notation of Section 22.11. Put

$$t = \min(j, h), \qquad m = h - t, \qquad D_a(\gamma) = \det Z(\gamma)_{\{0,\dots,a-1\}}, \quad M_j = \frac{|\operatorname{Norm}_{E/\mathbb{Q}} D_j(\gamma)|}{N^{2f(j-h)_+}} > 0.$$

The last number is `Filling.minorNorm`. Its nonvanishing is part of the level condition. Let $X = \langle e_a : 0 \le a < h \rangle$. The normal-form condition means that for $j \le a \le h$ there are equal vectors on the two frames with coefficient α on f_a and 1 on γf_a and with only earlier frame indices otherwise, for a single $\alpha \in E$ satisfying $\alpha\bar{\alpha} = 1$. We use this condition to eliminate γe_a with $a \ge j$, successively. Define arrays

$$F_0 = (e_0, \dots, e_{h-1},\ \gamma e_0, \dots, \gamma e_{t-1}), \qquad P = (e_t^*, \dots, e_{h-1}^*, z, \gamma z).$$

When $j \le h$, $D_j \neq 0$ implies that the t columns γe_b are independent modulo X: pair with the first j original isotropic vectors. The normal form then says that all the other γe_a belong to their span together with X. When $j = h+1$, all the γe_a are independent modulo X. Indeed if $\gamma \sum y_a e_a \in X$ then its pairings with the complete frame give $Z(\gamma)(y, 0)^T = 0$, contrary to $D_{h+1} \neq 0$ unless $y = 0$. Thus in every case F_0 is a basis of $X + \gamma X$ and $W = (X + \gamma X)^{\perp}$.

Lemma 27.5 (Bordered frame determinant)**.** *With the above columns set*

$$\mathcal{M}_j = (F_0 \,|\, P)^* J_N (F_0 \,|\, P) + \operatorname{diag}(0, N I_{m+2}).$$

Then $|\operatorname{Norm}_{E/\mathbb{Q}} \det \mathcal{M}_j| = (N^{2f} M_j)^2$.

Proof. If $j = h+1$, reorder $(F_0 \,|\, P)$ into $(F \,|\, \gamma F)$. The correction N on $z, \gamma z$ annihilates the two diagonal Gram blocks because $F^* J_N F = \operatorname{diag}(0_h, -N)$. Hence the bordered matrix becomes $\left(\begin{smallmatrix} 0 & Z \\ Z^* & 0 \end{smallmatrix}\right)$ with determinant $(-1)^n D_{h+1} \overline{D}_{h+1}$, giving the result.

Let $j \le h$. Group the columns as X, $Y = (\gamma e_0, \dots, \gamma e_{j-1}, e_j^*, \dots, e_{h-1}^*)$, and $Z_0 = (z, \gamma z)$. The pairing matrix of X with Y is of the form

$$A = \begin{pmatrix} Z_{<j,<j} & 0 \\ * & I_m \end{pmatrix}, \qquad \det A = D_j.$$

By eliminating the higher γe_a in the shared normal form, the normal-form relation on z becomes

$$\gamma z = \alpha z + u + \sum_{a<j} q_a \gamma e_a, \qquad u \in X.$$

Consequently subtracting those last γe_a terms from the final column makes both columns in the final group orthogonal to X. This column operation and its conjugate row operation do not alter the diagonal correction: the subtracted columns belong to F_0, not to P. The final 2-by-2 Gram block after correction is

$$\begin{pmatrix} 0 & -N\alpha \\ -N\bar{\alpha} & 0 \end{pmatrix},$$

of determinant $-N^2$. Expanding the block determinant first against the isotropic h columns paired with Y gives $(-1)^h \det(A) \det(A^*)$ times this final determinant (the remaining entries involving Y are immaterial). Thus in this case the bordered determinant is, up to sign, $N^2 D_j \bar{D}_j$. Taking the field norm gives exactly the formula. □

This proves `HasLevel.abs_norm_bordered_det` in `New/Filling/LinkPairing.lean` without using Cramer's rule for the normal-form frame.

On $M = X_0 \oplus Ez$ use $T = \operatorname{diag}(I_k, -N)$. On $M \oplus M$ the two forms used for the links are

$$L_- = \operatorname{diag}(T, -T), \qquad Q_L = \tfrac{1}{2}\operatorname{diag}(T, T) + N^{-1}L_z^* L_z, \qquad L_z = \operatorname{diag}(z^*T, z^*T).$$

The second is positive definite, with diagonal weights $1/2$ on the positive directions and $N/2$ on the two z-coordinates. Write $\ell_\gamma : W \to M \oplus M$ for the pair of middle coordinates of $w, \gamma^{-1}w$; its image is the link $\mathcal{L}_\gamma$, Lagrangian for L_-. Its kernel is $X \cap \gamma X$. These are the dimensions and isotropy already established in Chapter 22. For a rational basis matrix V of $\mathcal{L}_\gamma$ and $e \in \mathrm{GL}_{k+1}(A)$ write

$$\operatorname{disc}_L^2(V, e) = |\operatorname{Norm}_{E/\mathbb{Q}} \det(V^* Q_L V)|[\det e]^2.$$

This is `Filling.linkDiscSq`. We call Ve integrally liftable if it is the image of an array in $W(A) \cap B_c\mathcal{O}_f^r$ under ℓ_γ.

For the precise indices let $K_\gamma = B_c^{-1}\gamma B_{c'}$, write $Z(K_\gamma)$ for its integral frame Gram matrix, and define

$$I_1 = [\,\mathcal{R} : \mathcal{R}_j\,], \qquad \mathcal{R} = \sum_{a<h} \mathcal{O}_f Z(K_\gamma)^* e_a, \qquad \mathcal{R}_j = \sum_{a<t} \mathcal{O}_f Z(K_\gamma)^* e_a.$$

Here e_a in a frame-index space denotes the corresponding coordinate vector. Set I_2 to be the index with K_γ^{-1} substituted for K_γ. These are respectively `Filling.xRowsIndex` for the term and its inverse. Because the class bases fix F and preserve the form, the two full integral frame Grams are precisely the finite embeddings of $Z(\gamma)$ and $Z(\gamma^{-1}) = Z(\gamma)^*$. Normal-form elimination proves that all rows in either case lie in the span of the first t rows: for the first apply the shared relations expressing each e_a as a combination of earlier e's and γe's, whose latter pairings with γF vanish; for the second exchange the two frames. Thus the two indices are finite positive integers. Let

$$I_{\mathrm{sat}} = [\,F_{\mathrm{all}}(A) \cap \mathcal{O}_f^r : F_{\mathrm{all}}\mathcal{O}_f^{2h}\,], \qquad F_{\mathrm{all}} = (e_0, \dots, e_{h-1}, K_\gamma e_0, \dots, K_\gamma e_{h-1}),$$

where the notation $F_{\mathrm{all}}(A)$ means the column span, not an invertible matrix. It is exactly `Filling.saturationIndex`.

Theorem 27.6 (Upper link-lift discriminant bound)**.** *For every rational basis matrix V of $\mathcal{L}_\gamma$ there exists an adelic invertible e such that Ve is integrally liftable and*

$$\operatorname{disc}_L^2(V, e)\,(I_1 I_2 I_{\mathrm{sat}})^2 \le M_j^2. \tag{27.12}$$

All three indices on the left are nonzero.

Proof. The kernel of ℓ_γ has dimension $h - t = m$. Project it onto the original coordinates e_a with $a \ge t$. This is an isomorphism: if those coordinates vanish, a kernel vector lies in $\langle e_a : a < t\rangle$ and also in γX. It is orthogonal to every γe_b, hence vanishes by the leading $D_j \ne 0$ for $j \le h$; for $j = h+1$ the kernel is already zero. By rank this proves the claim. Choose a rational matrix U_0 for the basis of this kernel whose projection is the identity.

Lift the columns of V rationally to W and subtract kernel vectors so that their original coordinates e_a with $a \ge t$ vanish. Denote the resulting matrix by U_1. Then $U = (U_0 \,|\, U_1)$ is a basis of W. The adjoint pairing of U against the first m columns of P equals $(I_m \,|\, 0)$, and the last two pairings are $(0 \,|\, L_z V)$. For the underlying Hermitian form the pairings on W

depend only on their middle coordinates, which can be computed on either side of the link by unitarity. In particular

$$U^*(J_N + N^{-1}J_N PP^*J_N)U = \operatorname{diag}(N^{-1}I_m, V^*Q_L V).$$

This uses only the orthogonality to both isotropic frames; it does not treat the indefinite J_N as positive definite.

The basis in integral coordinates is $B_c^{-1}U$. Apply the block primitive-basis construction above to get

$$L = B_c^{-1}U \begin{pmatrix} a & z_1 \\ 0 & e \end{pmatrix}$$

primitive in $\mathcal{O}_f^r$. In particular $Ve = \ell_\gamma(B_c L_1)$ is integrally liftable. Put $F_{0,c} = B_c^{-1}F_0$ and $P_c = B_c^{-1}P$. Although these use the class basis, $F_{0,c}$ is integral because its first block is fixed while $B_{c'}$ fixes the frame for the second block; no integrality of the full rational matrices is assumed. Let i be the index of its generated lattice in its saturation. Apply Lemma 27.4 with $J = B_c^* J_N B_c$, $F_{0,c}$, P_c, and c there equal to the scalar N; its hypotheses hold by the class lattice unimodularity, the ranks and orthogonality. The above block Gram identity and Lemma 27.5 give

$$[N]^{-m}[\det a]^2 \operatorname{disc}_L^2(V,e)\, i^2 [N]^{m+2} = [N]^2 M_j^2,$$

so, cancelling the positive index powers of N,

$$[\det a]^2 i^2 \operatorname{disc}_L^2(V,e) = M_j^2. \tag{27.13}$$

It remains to prove the two index lower bounds; dropping these would lose the arithmetic divisors of the crude energy estimate. Since B_c^{-1} fixes original e-coordinates, $U_0 a$ is integral there, and its original last m coordinates form a, so a is integral. The surjective map

$$\mathcal{O}_f^m \longrightarrow \mathcal{R}/\mathcal{R}_j, \qquad (x_a)_{a\geq t} \longmapsto \sum_{a\geq t} Z(K_\gamma)^* e_a x_a$$

kills the columns of a. In fact $U_0 a \in X \cap \gamma X$ is orthogonal to γF, so its last m entries satisfy this relation modulo contributions from its integral first t entries. Therefore I_1 divides the index of $a\mathcal{O}_f^m$, namely $[\det a]$.

The lattice spanned by $F_{0,c}$ is contained in that spanned by F_{all}, with the same adelic span. The intermediate quotient surjects onto the analogous frame-row quotient defining I_2: apply the pairing map with the original full frame. It kills original X and takes the columns $K_\gamma e_a$ to the corresponding columns of $Z(K_\gamma)$, which are the conjugated rows of $Z(K_\gamma^{-1})$. Modulo those with $a < t$ it is thus the required surjection. Consequently I_2 divides

$$[\, F_{\mathrm{all}}\mathcal{O}_f^{2h} : F_{0,c}\mathcal{O}_f^{h+t}\,].$$

Multiplication of the two lattice indices gives $I_2 I_{\mathrm{sat}} \mid i$. Substitute both divisibilities in (27.13); this proves (27.12). □

This proves the full upper-lift estimate `exists_lifts_linkDiscSq_le` and the positivity part of `one_le_xRowsIndex_and_saturationIndex` in `New/Filling/LinkPairing.lean`. The weighted pair comparison still needs a complementary lower bound for liftable arrays independent modulo the two z-coordinates and a careful treatment of common links, not merely the upper bound of Theorem 27.6.

27.5 The lower bound modulo the positive middle spaces

We now prove the remaining lower-lift estimate and the resulting pair discriminant gain. Write

$$\Delta(\gamma) = D_{h+1}\overline{D}_{h+1} - N^2 D_h \overline{D}_h, \qquad d_\Delta(\gamma) = \prod_v |\Phi_v(\Delta(\gamma))|,$$

and M_j as above. At level $j \le h$ the discriminant vanishes; at the last level it does not. Set

$$Q_\gamma = \max(d_\Delta(\gamma), N^f M_j).$$

For each of the two z-coordinates in the link define the pairings using L_z as before, so the kernel of L_z is $X_0 \oplus X_0$.

Lemma 27.7 (Two test vectors for the link)**.** *The link contains vectors $\epsilon_1 = (a, -1; 0, u)$ and $\epsilon_2 = (0, \bar{u}; a', -1)$, where the notation separates the positive and the z parts of the two components, such that*

$$\Delta(\gamma) = D_j \overline{D}_j (1 - u\bar{u}).$$

If $E_0 = (\epsilon_1 \,|\, \epsilon_2)$ then on $\mathcal{L}_\gamma$ one has

$$E_0^* Q_L = L_z, \qquad E_0^* Q_L E_0 = N \begin{pmatrix} 1 & -\bar{u} \\ -u & 1 \end{pmatrix}. \tag{27.14}$$

In particular the rank of L_z on the link is one for $j \le h$ and two for $j = h+1$.

Proof. For $j \le h$, the shared vector at the last frame index in the normal form lies in both frame spans. Scale it to have original z-coordinate -1. That vector belongs to $W = (X + \gamma X)^\perp$; its two middle coordinates have the form $(0, -1; 0, u)$ with $u\bar{u} = 1$, by unitarity or by comparing the two coefficients in the normal form.

For $j = h+1$, put $D = D_{h+1}$ and let $y = Z(\gamma)^{-1} e_h D$, the last adjugate column. It satisfies $Z(\gamma)y = De_h$ and has last entry $y_h = D_h$. The vector $w = (N/D)\gamma F y$ is orthogonal to γX since it is in the γ-frame, and to X by the equation for $Z(\gamma)y$. Its z-coordinate is -1 because its pairing with z is N, and the middle coordinates of $\gamma^{-1}w$ are $(0, (N/D)D_h)$. Thus $\ell_\gamma(w) = (a, -1; 0, u)$ for $u = ND_h/D$, and the asserted discriminant formula holds. Apply the same calculation to the inverse term and exchange its two middle components to obtain a vector $(0, u'; a', -1)$ in $\mathcal{L}_\gamma$. The two vectors' L_- pairing is zero, hence $Nu' = N\bar{u}$, giving $u' = \bar{u}$.

The first identity of (27.14) follows directly from isotropy. For instance if $(x, s; x', t)$ is in the link then the first test vector's L_- pairing with it gives $a^*x + Ns + N\bar{u}t = 0$; its Q_L pairing equals $(a^*x - Ns + N\bar{u}t)/2 = -Ns$. The second row is $-Nt$ by the identical calculation with the roles exchanged. The second identity follows by evaluating the first on E_0. Positivity of Q_L implies that the rank of the Gram equals the rank of E_0, and by adjunction this is the rank of the z-pairing map on the link. The displayed determinant is zero exactly at the degenerate levels, while its diagonal entries are $N \neq 0$. This proves the rank assertion. □

This is the test-vector construction `HasLevel.exists_mem_link_termDisc_eq` in `LinkPairing.lean`, along with its required rank comparison.

Suppose that the residue s is agreeing and fixes nonzero frame vectors at places whose norm product is at least $X \ge 1$, in the precise sense of `Filling.FixesNormAtLeast`. The determinant-divisibility lemma of Section 22.11 (formula (22.11)) says that for two integral lift columns orthogonal to X_{iso} and its translate, whose two-by-two determinant of pairings with the two last frame vectors is nonzero, the generalized index of that determinant is at least X.

Here $X_{\mathrm{iso}} = \langle e_a : a < h\rangle$ denotes the isotropic span; it is distinct from the real number X. This result uses agreement of the characters in addition to the fixed-vector conditions. We use it only for two lift columns with invertible such determinant, not for arbitrary partial pairings.

Proposition 27.8 (Lower link-lift discriminant bound)**.** *Let Y consist of q rational columns of $\mathcal{L}_\gamma$ such that L_zY has rank q. Suppose that $v \in \mathrm{GL}_q(A)$ and there is an integrally liftable array W_1 with $L_zW_1 = L_zYv$. Then $q \le 1 + (j-h)_+$ and*

$$X'^2 M_j^2 \le \mathrm{disc}_L^2(Y, v) Q_\gamma^2, \tag{27.15}$$

where $X' = 1$ unless $j = h+1$ and $q = 2$, in which case it may be set equal to X.

Proof. The dimension bound is Lemma 27.7. If $q = 0$, the discriminant on the zero-dimensional basis is one and $Q_\gamma \ge N^f M_j \ge M_j$, proving the inequality.

Assume otherwise. Choose q rows of L_zY making a nonsingular square minor H; write E_I for the corresponding test vectors. Thus $E_I^* Q_L Y = H$. Its adelic product Hv is an integral matrix: these entries are the pairings of integral lift columns with $z, \gamma z$ in the class lattice, since $L_zW_1 = L_zYv$. All these frame vectors are integral by (27.3) and the term-position condition. Therefore $[\det(Hv)] \ge 1$. At $q = 2$ both rows are used and the determinant-divisibility lemma just recalled applies, giving $[\det(Hv)] \ge X$.

Apply the Gram Cauchy–Schwarz inequality in the positive form Q_L at all complex embeddings; equivalently use the adelic Liouville inequality of Section 22.13. It gives

$$[\det(Hv)]^2 \le |\operatorname{Norm}_{E/\mathbb{Q}} \det(E_I^* Q_L E_I)| \, \mathrm{disc}_L^2(Y, v).$$

For $q = 1$, the first factor on the right is N^{2f} by (27.14). Multiplying the inequality by M_j^2 gives (27.15), because $Q_\gamma \ge N^f M_j$. For $q = 2$ we must have $j = h+1$. Then the first factor on the right is

$$|\operatorname{Norm}_{E/\mathbb{Q}}(N^2(1 - u\bar{u}))| = \frac{N^{4f} d_\Delta(\gamma)^2}{|\operatorname{Norm}_{E/\mathbb{Q}} D_{h+1}|^2} = \frac{d_\Delta(\gamma)^2}{M_j^2}.$$

In this case use the lower bound X on the determinant index and $Q_\gamma \ge d_\Delta(\gamma)$. This proves the desired bound with the stronger factor X'^2. □

For the precise application in `minorNorm_sq_le_linkDiscSq_mul`, take $X' = X$ when $j = h+1$ and the complementary codimension i is zero, because then the chosen complement of $\ker L_z$ exhausts the two-dimensional quotient. At all other levels and codimensions it suffices to take $X' = 1$. This is the exact implication obtained there from the hypothesis labelled `hp`.

27.6 Common links and the simultaneous pair estimate

For two terms γ, γ' over the same classes and agreeing residue s, both of level j, put

$$\mathcal{I} = \mathcal{L}_\gamma \cap \mathcal{L}_{\gamma'}, \qquad i = k + 1 - \dim_E \mathcal{I}, \qquad C_0 = \mathcal{I} \cap \ker L_z.$$

Projection of C_0 onto the positive X_0 in the first middle component is injective: if this projection vanishes, isotropy for L_- and both zero z-coordinates imply that the remaining positive component has norm zero. Denote its image by T_0; this is the common link `commonLink`. Write $p = \dim_E T_0$.

Recall the height $H_c(T_0)$ from Section 22.13, namely the positive square root of the generalized index of the Gram determinant of an adelic basis of the saturated positive sublattice $T_0(A) \cap B_c\mathcal{O}_f^r$. In this formula T_0 is embedded in the original positive summand. This is `Filling.latticeDisc`. To define the other index, use the frame-value image of this positive sublattice under the map $x \mapsto (\gamma F)^* J_N x$. Let its integral image lattice be $\mathcal{P}$ and take $\mathcal{R}$ as in the proof of Theorem 27.6, which is the frame-value lattice generated by the original isotropic frame X_{iso}. We use the index

$$O_\gamma(T_0) = [\,\mathcal{P} : \mathcal{P} \cap \mathcal{R}\,].$$

It is allowed to be infinite before an integral lifting argument; we will see directly that it is finite. This is `Filling.frameValueIndex` for $a = 1$. Define

$$P_\gamma = H_c(T_0) O_\gamma(T_0) I_{\mathrm{sat}}(\gamma), \qquad G_\gamma = \frac{Q_\gamma}{I_1(\gamma) I_2(\gamma)}.$$

They are exactly `Filling.pairInvariant` and `Filling.classGauge`.

Lemma 27.9 (Orthogonal splitting of liftable arrays)**.** *Suppose $(V_0 \,|\, V_1)$ consists of rational basis columns orthogonal across the two blocks for Q_L, and its product by an invertible adelic coefficient matrix e is integrally liftable. There are invertible a, d and a matrix z_1 such that the arrays $V_0 a$ and $V_0 z_1 + V_1 d$ are integrally liftable and*

$$\mathrm{disc}_L^2(V_0, a)\,\mathrm{disc}_L^2(V_1, d) = \mathrm{disc}_L^2((V_0 \,|\, V_1), e).$$

Proof. In the lattice $e\mathcal{O}_f^{\dim V_0 + \dim V_1}$ take a basis of its intersection with the first coordinate block and lift a basis of the projection in the second coordinate block. This is the adapted-basis construction of Section 27.3. It gives a block upper triangular basis matrix $\left(\begin{smallmatrix} a & z_1 \\ 0 & d \end{smallmatrix}\right)$ differing from e by an integral invertible change of columns. Thus its two column blocks are integrally liftable. Its absolute determinant index is $[\det a][\det d] = [\det e]$. Orthogonality makes the rational Gram determinant a product of the two Gram determinants, proving the identity. This is `HasIntegralLifts.exists_orthogonal_split`. □

Proposition 27.10 (The complement gain)**.** *There exist a rational basis V_0, V_1 of $\mathcal{I}$, a rational basis Y of a complement of $\mathcal{I}$ in $\mathcal{L}_\gamma$, and v, z_1 such that the array $(V_0 \,|\, V_1) z_1 + Y v$ is integrally liftable and*

$$(P_\gamma X')^2\,\mathrm{disc}_L^2(Y, v) \le G_\gamma^2, \qquad k \le p + i + (j - h)_+. \tag{27.16}$$

Here $X' = X$ if $j = h + 1$ and $i = 0$, and $X' = 1$ otherwise. In particular $O_\gamma(T_0)$ is finite and at least one.

Proof. Choose V_0 a rational basis of C_0, V_1 a basis of its Q_L-orthogonal complement in $\mathcal{I}$, and Y a basis of the Q_L-orthogonal complement of $\mathcal{I}$ in the full link $\mathcal{L}_\gamma$. All these complements exist rationally since Q_L is totally positive definite. The map L_z is injective on the span of V_1 by the definition of C_0, so Lemma 27.7 bounds its dimension q by $1 + (j - h)_+$. As $p + q + i = k + 1$, this gives the dimension inequality. When $j = h + 1$ and $i = 0$, the subspace $\mathcal{I}$ is all of $\mathcal{L}_\gamma$, so $q = 2$ by the exact rank calculation.

Apply Theorem 27.6 to the rational basis $(V_0 \,|\, V_1 \,|\, Y)$; then apply Lemma 27.9 twice, first between the initial pair and Y, then between V_0 and V_1. This produces invertible adelic matrices a, d, v with three integrally liftable arrays

$$V_0 a, \qquad V_0 z_2 + V_1 d, \qquad (V_0 \,|\, V_1) z_1 + Y v$$

for suitable z_1, z_2, and with discriminant bound

$$\operatorname{disc}_L^2(V_0, a)\operatorname{disc}_L^2(V_1, d)\operatorname{disc}_L^2(Y, v)\,(I_1 I_2 I_{\mathrm{sat}})^2 \le M_j^2. \tag{27.17}$$

The middle array has L_z image $L_z V_1 d$, because V_0 is in the kernel. Hence Proposition 27.8 gives

$$\operatorname{disc}_L^2(V_1, d) Q_\gamma^2 \ge X'^2 M_j^2.$$

Write P_0 for the first positive-coordinate projection of V_0. Its Gram in the positive space is $P_0^* P_0 = V_0^* Q_L V_0$: the two positive-coordinate Grams agree by isotropy for L_-, and both z-coordinates vanish. Now lift $V_0 a$ integrally to $w \in W(A) \cap B_c \mathcal{O}_f^r$. Because w is orthogonal to original X_{iso} its dual coordinates vanish; since $L_z V_0 = 0$, its original z-coordinate vanishes, so $w = x_{\mathrm{iso}} + P_0 a$. The isotropic-coordinate part is integral by (27.3); therefore $P_0 a$ is an integral array in the positive sublattice of the class. Moreover w is orthogonal to γF (the last pairing vanishes because its second z-coordinate is zero). Thus the frame values of $P_0 a$ are negatives of the values on the integral x_{iso}, and belong to $\mathcal{R}$.

Since $P_0 a$ is full rank in T_0, its column lattice has finite index in the saturated positive lattice. Its image under the frame-value map lies in $\mathcal{P} \cap \mathcal{R}$, so the index of the column lattice is a multiple of $O_\gamma(T_0)$, by the surjection onto $\mathcal{P}/(\mathcal{P} \cap \mathcal{R})$. In particular this last index is finite and positive. The Gram index of $P_0 a$ equals the Gram index for a basis of the saturated positive lattice times the square of that column-lattice index (adjoint congruence by the basis change), giving

$$\operatorname{disc}_L^2(V_0, a) \ge H_c(T_0)^2 O_\gamma(T_0)^2.$$

Insert both lower estimates into (27.17), cancel $M_j^2 > 0$, then divide by $(I_1 I_2)^2$ to get (27.16). □

This proves `exists_link_split_le_classGauge` in `New/Filling/LinkPairing.lean`, including its exact codimension and integrally liftable complement. The argument makes clear the role of saturation and of both frame-value quotients.

Theorem 27.11 (Simultaneous pair discriminant gain)**.** *For two terms γ, γ' of level j over the same agreeing s fixing norm at least X, the above integer $i \ge 0$ satisfies $k \le \dim_E T_0 + i + (j-h)_+$; if $i = 0$, the links are equal. Moreover*

$$P_\gamma P_{\gamma'} X'^2 \le (2/g)^{2if}\, G_\gamma G_{\gamma'}, \qquad X' = \begin{cases} X & j = h+1,\ i = 0, \\ 1 & \text{otherwise.} \end{cases} \tag{27.18}$$

Proof. Apply Proposition 27.10 to both terms. The shared intersection and hence i are unchanged on exchanging them, so we obtain rational complement matrices Y, Y' for $\mathcal{I}$ with the indicated liftability and with coefficient matrices v, v'. The pairing matrix $Y^* L_- Y'$ is nonsingular. Indeed a vector in one complement annihilating the other also annihilates $\mathcal{I}$ and thus the whole opposite link. As that link is Lagrangian, the vector belongs to the intersection and hence is zero. This argument includes the empty determinant at $i = 0$.

Take the integral lift arrays w, w' in the rational class lattice giving the two complements plus their respective common-link parts. Direct expansion of the two middle pairings, using that the lift columns have no dual coordinates with respect to either of their isotropic frames, gives the mixed identity

$$\ell_\gamma(w)^* L_- \ell_{\gamma'}(w') = w^* J_N (1 - \gamma \gamma'^{-1}) w'.$$

Terms from $\mathcal{I}$ on either side vanish because that space lies in both isotropic links. Consequently the left side is $v^*Y^*L_-Y'v'$. As the two terms have the same reduction and class positions, their quotient lies in the principal level in the coordinates of class c:

$$B_c^{-1}\gamma\gamma'^{-1}B_c = K_\gamma K_{\gamma'}^{-1} \equiv 1 \bmod g.$$

Substituting $w = B_c\widetilde{w}$, $w' = B_c\widetilde{w}'$ with integral $\widetilde{w}, \widetilde{w}'$ shows that $g^{-1}v^*Y^*L_-Y'v'$ is integral. Its determinant is nonzero and has generalized index at least one.

Let $S_0 = Q_L^{-1}L_-$. These diagonal matrices satisfy $Q_LS_0 = L_-$ and $S_0^*Q_LS_0 = 4Q_L$. Apply the positive Gram Cauchy–Schwarz inequality over E to $g^{-1}Y$ and S_0Y' with their finite coefficient matrices v, v'. Because $[g] = g^{2f}$ and $[4] = 2^{4f}$, the result is

$$1 \le [\det(g^{-1}v^*Y^*L_-Y'v')]^2 \le (2/g)^{4if}\,\mathrm{disc}_L^2(Y,v)\,\mathrm{disc}_L^2(Y',v').$$

Multiply the two upper estimates (27.16) and use this lower bound on the product of discriminants. Taking the nonnegative square root gives (27.18). The rank assertion and equality when $i = 0$ follow already from the dimension formula and the equality of the two link dimensions. □

This is precisely `pairInvariant_mul_le` in `New/Filling/LinkPairing.lean`. No extra generic-angle or separate simultaneous-saturation hypothesis is needed. Its only geometric inputs are the exact congruent term positions, their level flag condition, the unimodular marked class lattices, and the residue fixed-vector determinant divisibility of Section 22.11.

27.7 Obligation status for the filling argument

The precise status of the interfaces relevant here is now as follows.

- **PROVED:** arithmetic frame separation and weighted count with exponent $6n^2D$, including the marked lattice and class-index normalization (`tsum_classCutoff_box_le`, `New/Filling/BoxPartition.lean`). The proof used the previously established arithmetic cutoffs, the stated standing size assumptions, and the free-column-level property.
- **PROVED (global implication):** `amplitudeAbs_lt_top_and_cut_le`, `summable_amplitude`, and `hasSum_amplitudeSum_change_of_weight`, in `New/Filling/AmplitudeSums.lean`, with actual error $\nu b^{-k^3 f}/2$. These now follow from Theorem 27.3 for any local Bessel systems satisfying their actual majorant property and the specified arithmetic sizes. The earlier separate global summability premise of Chapter 23 is no longer needed. **OPEN:** construction of the actual local majorant `IsCutPair.exists_majorant` in the Bessel system, or a proof that it follows from another established quantitative real-place theorem; completing the packet from the local levels and the remaining admissibility inequalities is still necessary.
- **PROVED:** the full algebraic B10 interface in `New/Filling/LinkPairing.lean`: `HasLevel.abs_norm_bordered_det`, `exists_lifts_linkDiscSq_le`, `one_le_xRowsIndex_and_saturationIndex` (including gauge positivity), `HasLevel.exists_mem_link_termDisc_eq`, `minorNorm_sq_le_linkDiscSq_mul`, `HasIntegralLifts.exists_orthogonal_split`, `exists_link_split_le_classGauge`, and `pairInvariant_mul_le`, with exact constants. Lemma 27.4 also gives the integral generalized-index identity for the regularized orthogonal complement from `Lib/General/Algebra/LatticeComplement.lean`. These are proofs for any congruent terms of the stipulated level and agreeing residue of the filling data. They do not depend on an unproved joint pair-discriminant inequality.

- **OPEN:** the weighted coloring and inverse-index counts of `New/Filling/BlockBound.lean` (notably `tsum_invPowWeight_le_of_codim_le` and `sum_pairWeight_le`), together with the precise local open-cell form needed for the block-amplitude estimate. The discriminant gain Theorem 27.11 is their arithmetic geometric input but does not imply the weighted counts without the separate lattice-index counting argument.
- **OPEN:** the resulting uniform crude energy majorants of `CrudeEnergy.lean` and coherent packet comparison in the actual numerical parameter range, with the remaining `Filling.IsAdmissible` degree, band, and mass inequalities (B04, the unresolved parts of B11–B12, B15–B16). The normalized limit and trace/contraction deductions of Chapters 22 and 23 are available once these named inputs and the local Bessel system have been supplied. No complete unconditional spanning theorem for the rows is claimed here.

Chapter 28

From the actual steep datum to a cut Bessel system

This chapter verifies the actual parameters, the measure-valued weighted cutoff, and the finite-level and reproducing interfaces of the filling construction. We use the preceding coefficient-identification, decay, differentiated Schur and orbital-coordinate arguments with their recorded normalizations; in particular the factor furnished by a cutoff is not an arbitrary change of Haar normalization. Write $r = D.r$, $m = D.m = 2(\varrho + 1)$, $n = \varrho + 1$, $k = r + 1 - m$, and $f = |\Phi|$ for the number of infinite places. Thus $r = k + 2\varrho + 1$. We suppress the infinite place from the notation until it is needed. The Hermitian involution J_1 and its orthonormal sign frame are as in Proposition 22.10.

28.1 The weight pattern of the actual row

The pattern for the grid $z = 0$ can be verified without an induction on a matrix of rank r. Let $K = k - 1$, $B = D.w(0, v)$, $h_2 = r^8$, $h = 2h_2$, and number the row from zero. From the source formulae `rowParameter`, `entrySign` and `gridParam` in `New/Seesaw/IndexTypes.lean`, the entries of $l/2$ are

$$\lambda_i = \begin{cases} k/2 - 1 - i, & 0 \le i < K, \\ -B - (2j+1)h_2, & i = K + j, \quad 0 \le j < m, \end{cases} \qquad \epsilon_i = \begin{cases} +1 & i < K, \\ (-1)^j & i = K + j. \end{cases}$$

Here k is even and r odd, and $m = 2n$. The entries are strictly decreasing. If c_i denotes the number of earlier entries of sign ϵ_i, and C_i the total number of entries of that sign, then the two rho counts are

$$2\rho(i) = r - 1 - 2i, \qquad 2\rho_c(i) = C_i - 1 - 2c_i.$$

Since $C_i = K + n$ and $c_i = i$ for $i < K$, the lowest compact weight $\Lambda_i = \lambda_i + \rho(i) - 2\rho_c(i)$ is zero there:

$$\frac{k}{2} - 1 - i + \frac{r - 1 - 2i}{2} - (k + n - 2 - 2i) = 0.$$

For every index, $|\rho(i) - 2\rho_c(i)| \le (r-1)/2$, because the difference is half the opposite-sign root count minus half the same-sign root count. Consequently the exceptional weights are strictly decreasing (their successive gap is at least $h - r$) and lie in $[-3B, -B]$ as soon as $(2m-1)h_2 + r/2 \le 2B$ and $h_2 > r/2$. These are the exact inequalities for `IsSteep`, with X the first $k-1$ positions. The same-sign initial gaps need not be large: the gap condition on the doubled Harish-Chandra parameter is imposed only between opposite signs. Their half-weight

difference is at least h except between an initial positive and an exceptional negative, where it exceeds B. Hence its large-gap constant is at least r^7 in the source's convention $500r^3 + A$.

For clarity we check the coarse numerical margins used here and below. In the notation of Section 53.3, write $L_0 = \log x$ and $A = \log b$. That section proves $A \le e^{L_0/100}$, $A \ge 10^{12}L_0$, $e^{5L_0} \le r \le 3e^{5L_0}$, $m \le 2e^{L_0/2}$, $[E : \mathbb{Q}] \le 2e^{12L_0}$, and the bounds for the two rounded weights b_1, b_2. In particular $B = b_i + h_2 \le 3b$, $B \ge 99h^3$ (the thin-place rounded weight is already larger), $a = D.a \le 2xb$, and the compact type exponent $a_v = D.\kappa(v, 1)$ is at most a. These inequalities and $L_0 \ge 10^6$ imply

$$\begin{aligned} 3B + a_v &\le 2b^2, \\ r + k + 3a + 1 &\le b^3, \\ r + 3a + 1 &\le b^4, \\ 10^{30}\bigl((r+k)^2 \log(b^3)\bigr)^3 &\le 10^{30} 1728 r^6 A^3 < r^7, \\ 3r^3 f &\le r^7, \\ (r+1)^4(3a + r + 1 + 2n^2) &\le b^5. \end{aligned}$$

For the fourth inequality, divide by r^6, take logarithms and use $\log(1728 \cdot 10^{30}) + 3L_0/100 < 5L_0$. For the fifth, use $f \le e^{12L_0}$ and $r^4 \ge e^{20L_0}$. For the last, the bracket is at most $7xb$ and the prefactor at most $256e^{20L_0}$; their product is smaller than b^5. The inequalities $b \ge 2000$, $b^3 \ge 10^{12}$, $8r \le b$ are immediate from $A \ge 10^{12}L_0$. Also $|l_i| \le 3a$: even the largest exceptional absolute value is bounded by $2(3b + 2mh_2)$, whereas $a \ge xb$. The compact infinitesimal character with type $\mathrm{Sym}^{a_v}\mathbb{C}^k$ has $|u_i| \le 2a_v + k \le 3a$. Its large-gap requirement in the Laplace argument is vacuous between opposite signs, since it is positive definite, and its distinctness and parity follow from adding the compact rho shift. This checks both parameters in `IsTwoFactor.laplace_method` with $L = 3a$ and $\beta = b^3$.

28.2 Peak and decay on the Bessel quotient

Let $S = U \rtimes H$ be the Bessel graph and $H = U(k)$. Write an element as (s_1, s_2) with $s_1 \in G = U(J_1)$ and $s_2 \in H$. We use the graph measure $d\sigma$ induced by the fixed unipotent coordinate Haar and probability Haar on H. It is right as well as left invariant: U is unipotent, and conjugation by H preserves its Haar (the modular factor is a positive continuous character on a compact group, hence identically one). The first graph projection is proper because it identifies S with a closed matrix subgroup; the second factor is determined continuously by its middle diagonal block. For every $g \in G$ write $|g| = \|g\|_{\mathrm{op}}$, so $|g| = |g^{-1}| \ge 1$. These facts also hold in the orthonormal sign frame.

Lemma 28.1 (Moving-coset decay)**.** *For any proper continuous homomorphism $j : S \to G$, put $B_j = \{s : |j(s) - 1| \le 1\}$. For every $y \in G$, if $M = 8r^2 + 1$, then*

$$\int_S (2/|j(s)y|)^M \, d\sigma(s) \le 16^M \sigma(B_j), \qquad 0 < \sigma(B_j) < \infty.$$

Proof. For $R \ge 1$, use a maximal $1/R$-separated set of points $j(s_i)y$ in the radius-R norm ball of the real $2r^2$-dimensional matrix space. Its cardinal is at most $(1 + 2R^2)^{2r^2} \le (3R^2)^{2r^2}$ by disjoint smaller norm balls. Every point in the set under consideration is within $1/R$ of one of those centers. Since $|(j(s_i)y)^{-1}| \le R$, this implies $|j(s)j(s_i)^{-1} - 1| \le 1$. Thus the inverse image is covered by right translates $B_j s_i$, each of Haar mass $\sigma(B_j)$. This mass is finite by

properness and positive because the ball contains an open identity neighborhood. On shells $2^j \le |j(s)y| < 2^{j+1}$, $j \ge 0$, summation gives a bound

$$3^{2r^2}2^{4r^2+M}\sigma(B_j)\sum_{j=0}^{\infty}2^{j(4r^2-M)} \le 16^M\sigma(B_j).$$

Indeed the sum is at most two and $3^{2r^2} \le 2^{4r^2}$. The same calculation applies to any large integer exponent $> 8r^2$ with the corresponding constant. □

Choose the highest vector v_H of the compact representation with norm one, in an orthonormal basis of the invariant Weyl Gram form. In this basis the representation is unitary and irreducible. One elementary verification of the latter assertion for a symmetric power is to use diagonal unitary matrices to project to the individual monomials (each has a distinct torus weight) and differentiate the complexified Lie algebra actions $x_i\partial/\partial x_j$ to connect all monomials of the prescribed degree. Thus an operator commuting with the group commutes with all diagonal weight projections and all these transitions, and is scalar. The required highest-vector coefficient is ψ_u: on the compact group its weight-minor expression gives the same value (for a one-row diagram it is the indicated diagonal matrix entry to the power a_v). Write it here as ρ, the ordinary action denoted `besselValueRep` when read as operators on the Euclidean space. The transpose in the bilinear column pairing will be accounted for in Section 28.7.

Use the x in Proposition 22.10, so $\mathrm{bs}(x) \le b^{5n}$, and write $\psi = \psi_l$ in the unitary frame. The character $\chi(s)$ of the Bessel graph is $\exp(\sigma_0(s) - \overline{\sigma_0(s)})$, where σ_0 is the simple-root sum. Define

$$Y(y) = \int_S \psi(x^{-1}s_1y)\chi(s)^{-1}\rho(s_2^{-1})v_H\, d\sigma(s).$$

Here χ and the compact action commute. Its norm is bounded by the integral of $|\psi|$. Decay from Lemma 22.5 with exponent M, followed by Lemma 28.1 for $j(s) = x^{-1}s_1x$, gives convergence and a uniform global bound. Continuity follows by dominated convergence: on a compact set of y the factor multiplying $j(s)$ has uniformly bounded inverse norm. Changing s by right translation proves left Bessel equivariance. Convolution is in the convention $(Y * e)(y) = \int_G e(g^{-1})Y(yg)\,dg$, with $e = d\psi$ and d the formal degree. For absolute Fubini use the preceding uniform bound on the integral of the absolute vector integrand and $e \in L^1(G)$. The identity $\psi * e = \psi$ is precisely the coefficient Schur identity proved in Section 25.10; moving it inside the integral proves $Y * e = Y$.

Pairing $Y(x)$ with v_H yields the Laplace integral of Proposition 21.6; the conjugate compact coefficient is from $\rho(s_2^{-1})$. Its phase is the phase verified in Proposition 22.10, not merely an additive character chosen approximately. The cone and small-ball estimates in Proposition 22.8 and Lemma 22.6 thus give

$$\sigma(B_j) \le b^{60(r+k)^2}|\langle v_H, Y(x)\rangle|, \qquad Y(x) \ne 0.$$

With $p = M + 60(r+k)^2$ and $P = b^p$ we conclude

$$\|Y(y)\| \le \int_S |\psi(x^{-1}s_1y)|\,d\sigma(s), \qquad \int_S (2/|x^{-1}s_1y|)^M\,d\sigma(s) \le P\|Y(x)\|. \tag{28.1}$$

Indeed $16^M \le b^M$. In particular $p + 99r^2 \le 348r^2 \le r^3$ in the actual range. This is `HasPeakAt` for the actual infinite place, with its uniform moving-coset estimate.

28.3 The exact weighted majorant and normalization

Here is the measure statement in `IsCutPair.exists_majorant`, including the factor $b^{3r^3 f}$ where the two functions differ. It is stronger than an unweighted cutoff or a small squared norm error.

Proposition 28.2 (Peak-to-cut with the actual mass product)**.** *Under the hypotheses just verified, there are an $L^1 \cap L^2$ continuous Bessel function F, its continuous cutoff V, a nonnegative measurable left-U-invariant a_0 on G, and a finite positive measure ν on G such that, with $D_v = \dim \rho$ and $R = b^{6n+2}$,*

$$F * (d\psi) = F, \quad \|F\|_2^2 = D_v, \quad \|V\|_2^2 \le D_v, \quad V(y) \neq 0 \Rightarrow \mathrm{bs}(y) \le R,$$
$$\|V - F\|_2^2 \le b^{-10r^2} D_v,$$
$$\begin{cases} \|T(y)\| & F(y) = V(y), \\ b^{3r^3 f}\|T(y)\| & F(y) \neq V(y) \end{cases} \le a_0(y) \le \nu\{g : \mathrm{bs}(yg) \le R\} \quad (T = F, V),$$
$$\nu(G) \int_{U\backslash G} a_0(y)\, dy \le b^{2r^3} D_v.$$

All quotient norms are for the source's Bessel quotient Haar, without renormalizing the unipotent subgroup.

Proof. Put $\omega_j(g) = (2/|g|)^j$, $C = 16^M$, $I_0 = \int_G \omega_{2M}(g)dg$, and

$$\Phi(y) = \int_S \omega_{2M}(x^{-1}s_1 y)\, d\sigma(s).$$

The function $\mathrm{bs}(y) = \inf_{s\in S}(|s_1 y| + 1)$ is left S-invariant and satisfies $\mathrm{bs}(yg) \le \mathrm{bs}(y)|g|$ and the reverse inequality obtained using g^{-1}; this proves continuity. The graph integral Φ is left S-invariant and measurable by Tonelli, and Weil integration gives $\int_{U\backslash G} \Phi = I_0$. The latter identity uses that the residual compact factor is probability Haar. More explicitly one integrates first over U, then over the compact H and $U\backslash G$; integrating $y \mapsto h_1 y$ does not change the quotient measure since H preserves Haar on U and on G.

Let $B_0 = \{g : |g - 1| \le 1\}$ and $v_0 = \mathrm{vol}\, B_0 > 0$. Introduce

$$d\mu(g) = \frac{CP\|Y(x)\|}{v_0} \omega_M(g)\, dg.$$

Lemma 28.1, applied to G itself, shows $\mu(G) \le C^2 P\|Y(x)\|$. We claim the exact translation inequality

$$\Phi(y) \le \mu\{g : \mathrm{bs}(yg) \le R\}. \tag{28.2}$$

For fixed s write $t = x^{-1}s_1 y$. If $g \in t^{-1}B_0$, then $|tg| \le 16$ and $\mathrm{bs}(yg) = \mathrm{bs}(s_1 yg) = \mathrm{bs}(xtg) \le 16b^{5n} \le R$. Also $|g| \le 16|t|$ there, so $\int_{t^{-1}B_0} \omega_M(g)dg \ge v_0 C^{-1}\omega_M(t)$. Consequently the mass on the right in (28.2) is at least $P\|Y(x)\|\omega_M(t)$, for every s. Multiply this inequality by $\omega_M(t)/(P\|Y(x)\|)$ and integrate over s. The bound (28.1) proves the claim, even though the sets $t^{-1}B_0$ need not be disjoint.

The gap bound gives both $|\psi(t)| \le \omega_{2M}(t)$ and $|\psi(t)| \le \omega_{2M+3r^3 f}(t)$ by Lemma 22.5: the exponent is at most $500r^3 + r^7$. In particular $\|Y(y)\| \le \Phi(y)$. Thus Y is integrable on the quotient and bounded by $\mu(G)$ there. The reproduced peak and Cauchy–Schwarz give

$$\|Y(x)\|^2 \le d^2 I_0 \int_G \omega_{2M}(g)\|Y(xg)\|^2\, dg = d^2 I_0 \int_{U\backslash G} \Phi(y)\|Y(y)\|^2\, dy \le d^2 I_0 \mu(G)\|Y\|_2^2.$$

The unfolding equality uses the invariance of the Bessel norm under left S. Therefore Y has positive and finite squared quotient norm. Set $c = \sqrt{D_v/\|Y\|_2^2}$, $F = cY$, and

$$V(y) = \max(0, \min(1, R - \mathrm{bs}(y)))F(y), \qquad a_0 = c\Phi, \quad \nu = c\mu.$$

The claimed mass product is now a consequence of the peak, not a new assumption:

$$\nu(G) \int a_0 = c^2 \mu(G) I_0 \le D_v \left(dI_0 C^2 P\right)^2.$$

By Lemma 25.12 and $r + 3a + 1 \le b^4$,

$$dI_0 C^2 \le 8(2000b^{24})^{2r^2} 2^{2M} 16^{2M} \le b^{50r^2+1+M} \le b^{99r^2}.$$

Here $2^{2M}16^{2M} = 2^{10M} \le b^M$ and $M = 8r^2 + 1$. Since $p + 99r^2 \le r^3$, this is the required bound.

If $F(y) \ne V(y)$, then $\mathrm{bs}(y) > R - 1$. For every s in the graph, writing $t = x^{-1}s_1 y$, the implication $|t| \le 2b \Rightarrow \mathrm{bs}(y) + 1 \le 2b\, b^{5n} + 1 \le R$ would contradict this. Thus $|t| > 2b$, and the second decay bound yields

$$b^{3r^3 f} \|F(y)\| \le c \int_S \omega_{2M}(t)\, d\sigma(s) = a_0(y).$$

The same is true for V since its cutoff factor lies in $[0, 1]$. Elsewhere the ordinary bound by a_0 suffices. Finally $a_0 \le \nu(G)$, so on the difference set,

$$\int \|V - F\|^2 \le b^{-6r^3 f} \int a_0^2 \le b^{-6r^3 f} \nu(G) \int a_0 \le b^{-10r^2} D_v.$$

The last inequality uses $f \ge 1$ and $r \ge 3$; all remaining norm, support, continuity and reproduction assertions follow directly from the definition of the cutoff. □

This proves every field of `IsCutPair`; in particular the majorant is weighted where $F \ne V$ and the same finite measure controls all translates. Differentiated reproduction is used only for F, never for the nonsmooth scalar cutoff. This is the exact conclusion of `HasPeakAt.exists_cut_isCutPair` in `New/RealPlace/BesselCut.lean`.

28.4 Schur and degree bounds in the actual normalization

Section 25.11 proves the differentiated convolution estimate for a reproduced Bessel function: for words of length at most $2n^2$, its constant is $d^{-1}\|F\|_2^2 P_0^{2|w|}$, provided $P_0 \ge (r+1)^4(L + r + 1 + 2n^2)$. With $L = 3a$ and the normalization above this gives exactly $d^{-1}D_v P_0^{2|w|}$ for all $P_0 \ge b^5$. The source also requires the bound $d\Theta \le b^{4r^2}$, where $\Theta = \pi^{(r-k)k} \operatorname{vol} U(k)/\operatorname{vol} U(r)$. Indeed the formal degree formula for compact-dual Haar is

$$d = \frac{\prod_{i<j}(\lambda_i - \lambda_j)}{\prod_{j=0}^{r-1} j!}, \qquad \operatorname{vol} U(j) = \prod_{i=0}^{j-1} \frac{2\pi^{i+1}}{i!}.$$

This degree is the positive Harish-Chandra degree of the coefficient identified in the preceding chapters. Each half-weight difference is at most $3a \le b^2$, so the numerator is at most $b^{r(r-1)}$. Multiplication by Θ cancels the factors $j!$ for $k \le j < r$. The remaining denominator is at least one, as is $\prod_{j=k}^{r-1} 2\pi^{j+1}$. Consequently $d\Theta \le b^{r(r-1)}\pi^{(r-k)k} \le b^{2r^2} < b^{4r^2}$. In particular no bound depending on an unspecified choice of Haar measure is needed.

The results so far establish the full/cut, compact irreducibility, degree, reproduction, and Schur fields of the actual real Bessel system. The remaining field of `BesselSystem` is `HasBandLimits`, the correlation-tensor tower with its bottom estimate and arbitrary-head transitions. The construction above does not infer this field merely from the derivative bounds; the separate band-limit proof is necessary.

28.5 Finite-place construction of the standard level

Let T be an arbitrary finite conjugation-stable set of finite places; it need not consist only of the chosen level places. Put

$$A_T = \prod_{w \in T} E_w, \qquad R_T = \prod_{w \in T} O_w, qquad \psi_T(x) = \psi_f(-\varpi^{-1}\iota_T(x)),$$

where ι_T is extension by zero to the finite adeles. Denote the fixed and skew subspaces by A_T^+ and A_T^-, respectively. The following verifies the base and top measures of `IntegralPlace.IsStdLevel`, rather than postulating a level satisfying its abstract hypotheses.

Lemma 28.3 (Integral place with the actual character)**.** *There are a trace-one $\tau \in R_T$, a skew unit $\delta \in R_T^\times$ with inverse in R_T, a skew Haar measure of mass one on R_T^-, and a modulus* md *for which all fields of `IntegralPlace` hold, with $c = 1$ and character ψ_T.*

Proof. At each orbit of finite places the involution is either a split pair of factors or that of an unramified quadratic extension. This was proved in the CM construction (Chapter 6); the field also contains i with $i^2 = -1$, so take $\delta = i$. The trace on integers is onto at an inert place: the trace is onto in the finite residue extension and an element of trace a unit can be divided by its trace. In a split factor use the element $(1, 0)$. These choices give $\tau + \bar{\tau} = 1$ also in residue characteristic two. The norm of units is onto the fixed-factor units. For the inert extension first choose a residue unit with the desired norm, by surjectivity of the finite field norm, and then use

$$N(1 + \pi^j a) \equiv 1 + \pi^j \operatorname{Tr}(a) \pmod{\pi^{j+1}} \qquad (j \geq 1)$$

to lift successively; completeness supplies the unit of exact norm. In the split case the norm is simply multiplication of the two coordinates. In particular there is $c_0 \in R_T^\times$ of norm -1.

Normalize additive Haar on A_T giving R_T mass one and on A_T^- giving R_T^- mass one. The additive isomorphism

$$(s, t) \longmapsto \tau\delta^{-1}s + t \quad (A_T^- \times A_T^- \longrightarrow A_T)$$

maps $R_T^- \times R_T^-$ onto R_T. Its inverse is $s = \delta(x + \bar{x})$, $t = x - \tau(x + \bar{x})$. Hence it preserves the stated Haar measures. The modulus is $\mathrm{md}(x) = \prod_{w \in T} |x_w|_w$ with local normalized absolute values. It is the inverse generalized index on units, zero on nonunits, and invariant under star and negation. For an invertible matrix the additive change of variables has determinant modulus $\mathrm{md}(\det M)$ (elementary row additions preserve measure, while scaling one coordinate has exactly the asserted module). Nonunits are a measure-zero union of coordinate hyperplanes. The locally integrable bound required in `Place` is $\mathrm{md}(t)^{-1/2}$ on units, extended arbitrarily on the null set: locally its integral is finite by the geometric series

$$\int_{O_w} |t_w|_w^{-1/2} dt_w = \frac{1 - q_w^{-1}}{1 - q_w^{-1/2}}.$$

The product over the finite set and translations of compact balls prove the local assertion.

Since the different is generated by ϖ, the trace pairing at w gives exactly the annihilator O_w for the character $x \mapsto \psi_w(-x/\varpi)$. Thus $x \in R_T$ if and only if $\psi_T(xr) = 1$ for every $r \in R_T$: for the converse one tests the factors independently, using that bar permutes the factors. The Fourier transform of 1_{R_T} for $\psi_T(\bar{v}z)$ is again 1_{R_T}, by this annihilator identity and character orthogonality on a compact additive subgroup. This nonnegative continuous integrable kernel takes value one at zero and its transformed absolute integral is one. Hence the Plancherel-kernel field of `Place` holds with $c = 1$. All spaces are second countable locally compact groups and the operations and the character are continuous. This proves the lemma, including the exact choice of character. If T is empty, all groups here are the trivial products, and all measures have mass one. □

In the lattice coordinates of the special middle part, let $v_0 = x_{k_0}/\bar{\pi} + z/N$, where $N = \pi\bar{\pi}$ with the source idele. The Gram matrix on (v_0, z) is

$$\begin{pmatrix} 0 & -1 \\ -1 & -N \end{pmatrix};$$

the other $k-1$ middle coordinates are orthonormal, and the ϱ outside hyperbolic pairs are unchanged. The form is integral and unimodular even if N is not a unit in R_T.

Lemma 28.4 (Hyperbolic splitting and integral box measures)**.** *Suppose $k = |\iota| + 2j + 1$. For $g \in R_T$ self-conjugate and invertible in A_T, the marked lattice has the data and measures of `IntegralPlace.IsLatticeLevel` with basis consisting of $j+1$ middle hyperbolic pairs and $|\iota|$ orthonormal vectors. In particular its box $K = U(J, R_T)$ has*

$$\theta(K) \geq \mathrm{vol}(R_T^{\times}), \qquad \mu_H(K) = 1.$$

Proof. The first hyperbolic pair is $(-v_0, z - N\tau v_0)$, whose pairing is one and whose two self-pairings vanish. The determinant of this basis change is a unit. For each pair of unused orthonormal coordinates, the columns $(1, c_0)$ and $(\tau, -\bar{\tau}c_0)$ form another hyperbolic pair: their respective squared norms vanish because $c_0\bar{c}_0 = -1$, their pairing is $\tau + \bar{\tau} = 1$, and their determinant is $-c_0$. Use j such pairs. This supplies an integral basis with integral inverse on the middle space and unit determinant modulus, as well as the orthonormal remainder. Put $u = -v_0$, so $h(u, e) = e_z$ in the lattice coordinates and $u_z = 0$. This is the required isotropic vector in the source.

For the top measure, on the middle space let Q be the quadric $h(e, e) = h(z, z) = -N$. In the first hyperbolic pair and remaining orthogonal block write $e = (b, t, w)$. Its standard flat chart is

$$b = \bar{t}^{-1}\tau(-N - h(w, w)) + ta, qquadt \in A_T^{\times},\ w \in A_T^{k-1},\ a \in A_T^{-}.$$

The source measure of Q is the image of $dt\, dw\, d\nu^{-}(a)$, with no additional density. For verification, at fixed $t \neq 0$ the change of variables $b = \bar{t}^{-1}\tau\delta^{-1}\gamma + ta$ (on varying the norm coordinate γ by the appropriate additive constant) is measure preserving: factor out $\bar{t}^{-1}$ and use the additive trace/skew splitting of Lemma 28.3, then multiply the skew variable by $t\bar{t}$. Its modulus on the skew line is $\mathrm{md}(t\bar{t})^{1/2} = \mathrm{md}(t)$; this cancels $\mathrm{md}(\bar{t}^{-1})$. Thus this is also the disintegrated Haar measure in `Place.flatChart`. It is locally finite: for t, w restricted to a compact coordinate box, the admissible skew a at fixed t, w, if nonempty, lies in a translate of $\{a : ta \in C\}$ for a compact box C. At an inert factor this set has measure at most a box-dependent constant times $|t|_w^{-1/2}$; for split factors, writing $a = (a_1, -a_1)$, its measure is at most a constant times $\min(|t_1|^{-1}, |t_2|^{-1}) \leq |t_1 t_2|^{-1/2}$. Lemma 28.3 gives integrability. The chart is one-to-one almost everywhere, and the excluded null set consists of nonunit t.

The original lattice z-coordinate of e is still t, since the first hyperbolic dual has lattice z-coordinate one and every other basis vector has coordinate zero. For $e \in Q$ and $t = e_z \in A_T^\times$, put $v = e - z$, $q = h(v, e)$ and

$$g_e = 1 + uh(v, \cdot) - t^{-1}(v + qu)h(u, \cdot).$$

This is an isometry fixing the exterior hyperbolic pairs, sends e to z, and is integral if e and t^{-1} are integral. We verify the isometry explicitly. Write $w^\dagger = h(w, \cdot)$, $g_e = 1 + D$, with $D = uv^\dagger + Bvu^\dagger + Cuu^\dagger$, $B = -t^{-1}$, $C = -q/t$. Then $h(u, u) = 0$, $h(u, v) = t - 1$, $h(v, u) = \bar{t} - 1$, and $h(v, v) = q + \bar{q}$. In $D + D^\dagger + D^\dagger D$, the coefficients of $vu^\dagger$ and $uv^\dagger$ vanish because $1 + B + B(t - 1) = 0$ and its conjugate vanish. The remaining coefficient is

$$C + \bar{C} + |t|_{\mathrm{alg}}^{-2}(q + \bar{q} + q(\bar{t} - 1) + \bar{q}(t - 1)) = 0, \qquad |t|_{\mathrm{alg}}^2 = t\bar{t}.$$

The formula $g_e e = e + qu - (v + qu) = z$ is immediate. If e, t^{-1} are integral so is g_e, and its inverse is integral by the unimodularity of the Gram form. All constructions are measurable.

Push the flat-chart measure forward under $e \mapsto g_e$ to obtain θ. It is supported on G_ϱ, and its pushforward by $g \mapsto g^{-1}z$ is exactly the quadric measure. It is sigma-finite, since it is supported on the measurable section and the inverse-image of a compact set under it projects into a compact set of quadric coordinates, which has finite measure. The chart subset

$$t \in R_T^\times, \quad w \in R_T^{k-1}, \quad a \in R_T^-$$

has mass $\mathrm{vol}(R_T^\times)$; the formulae show $g_e \in K$ there. This proves the claimed lower bound.

Let H be the residual stabilizer of the last frame vector in G_ϱ. Normalize right Haar on H by $\mu_H(H \cap K) = 1$ and extend it to the group by the closed inclusion. It is sigma-finite and nonzero. The remaining condition in `Level.measureH_spec` is

$$\mu_H\{h \in H : Uhg' \cap K \neq \varnothing\} \leq 1 \quad (g' \in G).$$

If the set is nonempty fix h_0 in it. Choosing integral representatives uhg' and u_0h_0g', their quotient gives $u'hh_0^{-1} \in K$, with $u' \in U$ since H normalizes U. An integral element of UH has integral H-component: in the adapted isotropic filtration its induced map on the middle quotient is that component and the same applies to the inverse, since the lattice filtration is saturated. Those maps are in the original middle lattice and the embedding in the whole group is block diagonal. This proves $hh_0^{-1} \in H \cap K$, so the stated set is contained in a right translate of measure one. The condition is measurable, being the union of open sets indexed by $u \in U$.

Take the ball $\mathfrak{B} = R_T^r$ and the positions set $\{y^{-1}uy : y \in G, u \in U\}K(g)$. The ball is compact open, the box K compact open (unimodularity makes integral entries equivalent to integral inverse), and the positions set open as a union of translates of the open subgroup $K(g)$. This proves every defining field of the lattice level. □

Proposition 28.5 (The actual standard finite level)**.** *For every conjugation-stable finite T, the filling datum admits the standard level in* `Filling.IsFillingLevel`*: modulus $g = g_0g_1$, character ψ_T, $3\varrho+6$ hyperbolic pairs in the middle and orthonormal rank $k-6\varrho-13$.*

Proof. Take $j = 3\varrho + 5$ in Lemma 28.4. Thus $j + 1 = 3\varrho + 6$. The ideal zeta of R_T is the finite product of $(1 - q_w^{-j})^{-1}$. With $D_E = [E : \mathbb{Q}]$, at most D_E factors can lie over any rational prime, each with $q_w \geq p$. For $j \geq 2$,

$$\log \zeta_{R_T}(j) \leq 2D_E \sum_{a=2}^{\infty} a^{-j} \leq 4D_E 2^{-j},$$

where the last inequality uses $j \ge 3$ and the integral test. Now $nD_E \le 2^{2\varrho}$ by the filling rank scale: directly $\log(nD_E) \le 13L_0 + O(1)$ while $\varrho \ge e^{L_0/2}/3$. Hence

$$n \log \zeta_{R_T}(j) \le 2^{-\varrho-3}, \qquad \zeta_{R_T}(j)^n \le 1 + 2^{-\varrho-1}.$$

In the latter inequality use $e^x \le 1 + 2x$ for $0 \le x \le 1/2$. Also $j < 9n$ and $k - 6\varrho - 13 \ge 13n$ because $k \ge 1000m^9$, a consequence of $k \ge e^{5L_0}/2$ and $m \le 2e^{L_0/2}$. These are precisely the `zeta_rank` inequalities.

The finite unitary group has bi-invariant Haar: on each local factor the modulus of conjugation equals the absolute value of the determinant of its adjoint on the Lie algebra. Over the quadratic splitting algebra this Lie algebra is $\mathfrak{gl}_r$ and conjugation on matrices has determinant $(\det g)^r(\det g^{-1})^r = 1$; descent gives determinant one on the original Lie algebra, including the split case. The equality of the Haar modulus and this determinant follows in a sufficiently small analytic identity chart by changing variables, and so holds on the entire group. Its product yields the required unimodular measure on A_T. This supplies `IsStdLevel`, with the character and modulus of Lemma 28.3 and Lemma 28.4. In particular this construction establishes `Filling.exists_isFillingLevel` of `New/Filling/GlobalToLocal.lean` for the actual finite sets. It does not by itself prove that every gauge inequality or band field required later follows from this standard level: those are additional quantitative assertions. □

28.6 Simultaneous numerical admissibility

We check together the band, energy and mass conditions of `Filling.IsAdmissible`. The actual Bessel systems including their band field are understood in these numerical statements; no further assumption on their degree or coordinate count is needed. Use $A = \log b$, let $\Delta = |\operatorname{disc} E|$ and $d_E = \Delta^{1/(2f)}$ be the root discriminant, and set

$$W = \sum_v \log w_v^{\mathrm{nom}}, \qquad L_g = \log g,$$

$$\log N = \frac{mW}{f} + (m-1)\log(gd_E/(2\pi)) - \frac{15}{2}A + \delta_N, \qquad 0 \le \delta_N \le \sqrt{A}.$$

This is exactly the source's chosen window for the split prime N.

For the elementary different bound, $N_1 = N_0 t_s$ with $t_s = 2^s 3^{\lfloor \log_3 s \rfloor + 5}$. Since the added prime powers are already supported in N_0, the primitive roots of order N_1 are precisely the roots of $\Phi_{N_0}(X^{t_s})$. Differentiation at a primitive root ζ therefore gives $|\Phi'_{N_1}(\zeta)| = t_s |\Phi'_{N_0}(\zeta^{t_s})|$. The constant $P.c$ in `New/Data/AdmissibleData.lean` includes the sum of the absolute logarithms of these latter derivatives at all primitive roots. Since $2^s = m^{22}$, $s \le 32L_0$, $P.c \le L_0/10^{14}$ and the purely imaginary different generator differs from $\Phi'_{N_1}(\zeta)\sqrt{-d}$ by a norm-one unit, we obtain

$$x^{10} \le |\Phi_v(\varpi)| \le x^{12}, \qquad d_E^f = \prod_v |\Phi_v(\varpi)|.$$

For the product, take the absolute norm of its principal ideal, which is the different. In particular $10L_0 \le \log d_E \le 12L_0$. The real scale is $|\kappa_v| = g|\Phi_v(\varpi)|/(2\pi)$, hence $1 \le |\kappa_v| \le b^{23}$ and $\sum_v \log|\kappa_v| = f\log(gd_E/(2\pi))$. The target and window imply $b^m \le N \le b^{30m}$: indeed $(1 - 10^{-6})A \le \log w_v^{\mathrm{nom}} \le A + \log 2$, $19A \le L_g \le 22A$, and $\log d_E \le 12L_0 \le 12A/10^{12}$. Substitution proves $mA \le \log N \le 30mA$ (all ranks here exceed 10^5). The level-prime and different inequalities thus have their required signs and magnitudes.

Let $D = |\mathcal{I}|$ be the product of local Bessel-coordinate cardinalities, $C = |\mathcal{C}|$ the number of column classes, $d_\infty = \prod_v d_v$, and $M_j = \mathrm{mass}(E, j)$. The symmetric-power binomial formula,

which follows by counting monomials, gives

$$0 \le \log D + f\log((k-1)!) \le f(k-1)\log(a+k) \le 1.01fkA. \tag{28.3}$$

In particular $D \le b^{2kf}$. The marked-lattice computation of Lemma 22.11, with its free column level, gives

$$I_N = (N^k-1)^f, qquadC = |\bar{H}|I_N M_k, qquad|\bar{S}| = |\bar{H}|U_0, \quad U_0 = g^{f\varrho(2k+2\varrho+1)}.$$

Here $\bar{H}, \bar{S}$ are the images in the finite level group and $U_0 = |\bar{U}|$. The finite semidirect-product identity follows by projecting the integral Bessel subgroup to its middle block: integral factors have integral projections, their intersections after reduction are trivial (middle diagonal block versus strictly triangular blocks), and the integral unipotent group has one independent residue parameter per root coordinate. These are the counts proved in Sections 22.7 and 22.8. Since $N \ge b^m$, $I_N \ge (N^k/b)^f$ and

$$-b^{-10} \le f\log(1-N^{-k}) \le 0. \tag{28.4}$$

For the degree bound, in the ordered parameter of Section 28.1, all pairs of indices in $i<j<k-1$ contribute $j-i$, whose product cancels the initial factorials in the Weyl degree. If $j \ge k-1$, every difference is a positive integer, no larger than $w_{j+2-k}(v)$; when $i<k-1$ it is at least w_v^{nom}. These follow directly from the displayed row entries, since $h_2 > r$ and $w_v^{\mathrm{nom}} \le b_i$ in the relevant thick or thin place. The logarithm of each $w_j(v)$ in this range is at most $\log w_v^{\mathrm{nom}} + 1/x$: rounding and the additive shift $jh+h_2$ total at most x^{42}, while $w_v^{\mathrm{nom}}/x \ge x^{42}$. Thus

$$m(k-1)W \le \log d_\infty + f\sum_{j=k-1}^{r-1}\log(j!) \le mr(W+f/x). \tag{28.5}$$

This proves the degree-window lemma of `New/Filling/DegreeBounds.lean` in the actual parameters.

Let $\mathcal{T} = d_\infty \Theta^f\, b^{7kf}\prod_v(|\kappa_v|^{m-1}/(gN))^{k-n^7}$, where Θ is the common band constant. The volume formula already used in Section 28.4 gives $\log(d_\infty\Theta^f) \le mr(W+f/x)$, since Θ restores only the factorials with index $j \ge k$ and includes nonpositive residual powers of 2 and π. Consequently

$$\begin{aligned}\log\mathcal{T} &\le m(r-k+n^7)W + fmr/x + 7kfA\\ &\quad + f(k-n^7)(\frac{15}{2}A - L_g - \delta_N).\end{aligned}$$

As $W \le 2fA$, $k \ge 1000m^9$, $n = m/2$, $r \le 2k$, the first two terms are at most $kfA/10$, whereas the last two are at most $-4kfA$. For example $2m(r-k+n^7)/k \le 2m(m+n^7)/(1000m^9) < 1/100$, $mr/(kxA) < 1/100$, and $11.5n^7/k < 1/100$. Thus $\mathcal{T} \le 1$, proving the exact product-band inequality.

We use the exact uniform mass-ratio estimate of Theorem 6.13: for $8f \le 2^u$, $j \ge 4u+12$,

$$\left|\log(M_{j+1}/M_j) - f\left((j+\tfrac12)\log d_E + \log(j!) - (j+1)\log(2\pi)\right)\right| \le 2^{1-u}.$$

Take $u = \varrho$ and sum on $k \le j < r$. The stated hypotheses hold since $\log f \le 12L_0$, $\varrho \ge e^{L_0/2}/3$, and $k \ge 1000m^9$. Write the resulting signed error as ϵ_M, so $|\epsilon_M| \le 2m2^{-\varrho} \le b^{-10}$; this follows by taking logarithms, using $A \le e^{L_0/100}$.

Put $T_0 = b^{kf/4}$, $e_1 = |\bar{S}|CD$, $e_0 = (CD)^2/(d_\infty M_r)$, the source tail scale, Bessel and main energies. Inserting the class count gives

$$\frac{e_0}{e_1} = \frac{DI_N M_k}{d_\infty M_r U_0}.$$

Write $\epsilon_d = \log d_\infty + f\sum_{j=k-1}^{r-1}\log(j!) - m(k-1)W$. By (28.5), $0 \le \epsilon_d \le m^2W + fmr/x$. Direct summation of the mass ratio and substitution of the prime window therefore give

$$\log(e_0/e_1) = kf(L_g - 7.5A) + \log D + f\log((k-1)!) + E_1, \tag{28.6}$$

$$E_1 = mW - \epsilon_d + kf\delta_N - f\varrho(m-1)L_g - f\frac{(m-1)^2}{2}\log d_E + f\frac{m(m-1)}{2}\log(2\pi) + f\log(1-N^{-k}) - \epsilon_M. \tag{28.7}$$

For an explicit absolute bound, the sum of the absolute values of all terms of E_1 except $kf\delta_N$ is at most $40fm^2A + fmr/x + 2b^{-10}$. Since $k \ge 1000m^9$, $r \le 2k$, $m/x \le 2/\sqrt{x}$ and $\sqrt{A} \le A/100$, this gives $|E_1| \le kfA/4$. Combining this with (28.3), $L_g \ge 19A$, yields

$$11kfA \le \log(e_0/e_1) \le kf(L_g - 6A) - \log 2.$$

The upper inequality has a margin $0.24kfA > \log 2$. Now $\mathrm{codimFactor} = 2b^{1.5kf}/g^{kf}$, which is at most one. These logarithmic inequalities are exactly $T_0^{44}e_1 \le e_0$ and $T_0^{18}\mathrm{codimFactor}\, e_0 \le e_1$. No independent estimate of class cardinalities by their asymptotic sizes was substituted for the class formula.

For the last threshold put $v_0 = \mathrm{vol}\,\widehat{O} = (2/d_E)^f$, $d_U = \varrho(\varrho+k)$, and let S_f denote the product of frame scales in `Filling.frameScaleProd`. As in Section 22.8, the exact unipotent covolume and the Siegel-coordinate constant are

$$\nu_U S_f = \frac{U_0}{N^{kf}} v_0^{-d_U}\Delta^{\varrho/4}, \quad c_* = \frac{\pi^{kf}}{((k-1)!)^f N^{kf}} v_0^k \Delta^{1/4}.$$

The packet coefficient is $a_* = |\bar{S}|U_0 C c_*$. With $c_G = (\mathrm{vol}\,U(k)/\mathrm{vol}\,U(r))\,2\pi^k/(k-1)!$ the quotient of the required right side $(1+b^{-9})e_0 d_\infty c_G^f v_0^{-n}(\nu_U S_f/D)^2$ by a_* simplifies to

$$(1+b^{-9})(1-N^{-k})^f \frac{M_k}{M_r}(2\,\mathrm{vol}\,U(k)/\mathrm{vol}\,U(r))^f v_0^{-n-2d_U-k}\Delta^{\varrho/2-1/4}.$$

This expression is exactly $(1+b^{-9})(1-N^{-k})^f e^{-\epsilon_M}$. To verify cancellation, let $s_1 = \sum_{j=k}^{r-1} j = (2\varrho+1)(k+\varrho)$. Then $s_1 + 1 = n + 2d_U + k$ (the exponent for 2), and $s_1 + (r-k)/2 = n + 2d_U + k + \varrho - 1/2$ (the exponent for d_E); the factorials and π powers cancel between the volume and mass formula. By (28.4) and $|\epsilon_M| \le b^{-10}$, the logarithm of the expression is at least $\log(1+b^{-9}) - 2b^{-10} > 0$. This proves the last, sharp packet coefficient inequality, not merely a comparison up to an unspecified constant.

The other algebraic fields of `IsAdmissible` are as follows. Column level is the preimage of the marked principal level by the column inclusion, since preserving the marked lattice imposes the last-vector condition at N, and reduction imposes the independent chosen-place conditions at the two other primes. Reduction of the larger level is surjective: at a chosen split place the first factor is an ordinary $\mathrm{GL}_r(O_w)$, whose reduction is onto $\mathrm{GL}_r(\mathbb{F}_{g_i})$ by lifting entries and the nonzero determinant. The conjugate factor is determined by inverse adjoint; all other coordinates can be taken to be the identity. This constructs the adelic lift simultaneously. Together with the class and degree inequalities this proves `Filling.IsAdmissible` from the specified filling and the actual numerical data. In particular these calculations do not leave a separate simultaneous energy-scale premise after construction of the band-limit field.

28.7 Fitting the odd graded space and reproducing the Poincaré series

The exact local representation intertwiner is useful here, because the column pairing is bilinear, not a Hermitian inner product. Write $S(g)$ for the Schur matrix and Q for its positive Weyl Gram matrix. Take C with $C^*C = Q$, so our orthonormal compact representation is $CS(g)C^{-1}$. The transpose rule of the Schur module is

$$QS(g) = S(g^t)^t Q.$$

For the symmetric power this follows immediately from the tensor-power bilinear pairing restricted to the equal-degree symmetric monomials (equivalently the integral Gram identity in Chapter 8); in the original Weyl basis it is this same pairing's Gram matrix. Thus C^* times the orthonormal Bessel column transforms by $S(g^t)^t$, for both F and V. This supplies the invertible A, A' of `Filling.IsBessel.equiv_schur`, with $A = C^*$ and $A' = A^{-1}$. The degree and convolution fields were proved above. Tensoring the local intertwiners makes the source's `weylBesselEquiv`; pairing a Weyl column with the Bessel value gives the action $S(g^t)$ on the former and no complex conjugation. This is the precise convention needed for column evaluation at the definite class representatives.

The graded-lattice construction of Section 16.6 has, on Y_0, the odd string and finite matrix $x_f = B_{mgrad}(u)k_0$, where $u = D.\mathrm{latticeScale}(0)$ is a unitary scalar idele, k_0 is integral invertible and reduces to one at the chosen places. In the rational Bessel basis, setting $\eta = u^{-1}1_k$ gives the exact identity

$$B_{mgrad}(u) = u\ \mathrm{incl}_H(\eta) B_{\mathrm{lat}}.$$

On the ordinary X_0-columns scalar multiplication by u cancels η; every reflected string column is multiplied by u, and the marked column is sent to $\bar{\pi}^{-1}x_{k_0} + uz/N$. All other columns are unchanged as specified. Hence this is exactly the finite point identity in `Filling.Fits.point_eq`. The equality of index conventions is the ordering of the odd string, its reflected dual, and the definite coordinates, i.e. the Bessel index equivalence. The sign-frame comparison is orthonormal by Section 16.6; the weight kernel is idempotent by the coefficient Schur formula. Finally η commutes with the whole column group because it is scalar. These facts supply `Data.filling_fits` once the band-limited systems have been chosen.

For completeness, we prove the convergence and reproduction step independently of a formal manipulation of conditionally convergent Poincaré series. Let $\Gamma = G(E^+)$, K_c^{pr} be the standard principal level on the ambient finite group, and $\alpha = (c, i)$ be one class-coordinate pair. In the coordinates of `Filling.Fits`, the seed is

$$T_\alpha(g) = 1_{K_c^{\mathrm{pr}}}(h_c^{-1}\eta g_f)\,\beta_c(y) W_i(y), \qquad y = \mathrm{realHomeo}(g_\infty),$$

where in this formula the notation for finite elements denotes their transport by `finiteEquiv`. The W used here is the full Bessel tensor (the cutoff β_c is the arithmetic class cutoff, not the local scalar cutoff V). Right multiplication by the standard principal level preserves the seed, since finiteEquiv carries it onto the marked principal level by the point identity and $k_0 \equiv 1$. The cutoff is measurable and has periodization one on the rational lattice $\Gamma_c = \mathrm{classUnip}(c)$.

For fixed g, the sum over $\gamma \in \Gamma$ is supported on those γ whose finite position belongs to the designated coset. If this set is empty the seed sum is zero. Otherwise choose one such γ_0; the relative rational elements $\gamma\gamma_0^{-1}$ have finite positions in the term set with both classes c and fixed reduction one. In the corresponding translated real variable the absolute sum is bounded by the uniform assertion of Theorem 27.3. Its precise local majorant hypothesis

is now supplied by Proposition 28.2 and the dimension condition by (28.3). In particular there is a finite uniform bound M_α on $\sum_\gamma |T_\alpha(\gamma g)|$. Thus the sum is an ordinary absolutely convergent series of measurable functions, defines a bounded function, and has left rational invariance by reindexing.

Collapse the summation first over left cosets of Γ_c in Γ. The finite principal-level condition is constant on the inner orbit. The Bessel character is trivial on these rational unipotent elements (the real character equals the inverse finite character, which is one on the congruence subgroup), so the periodization of β_c collapses the inner sum. For fixed coset the resulting term is the full Bessel coordinate at the translated real argument times the finite indicator. It is therefore reproduced by $e_\infty = \bigotimes_v d_v\psi_{l_v}$, in the right-convolution convention of Section 28.2. Summing this equality over the cosets is justified by absolute Fubini: the absolute inner orbit sums are bounded by the original absolute sum, uniformly in g, and $e_\infty \in L^1$. The complete Poincaré function is thus fixed by e_∞. Moreover the convolution of a bounded measurable function with an L^1 kernel is continuous in the real variable, since translations of the kernel are continuous in L^1. This proves continuity and, with the open finite invariance, gives the required actual automorphic form of the reproduced type. The cusp-kernel assertion for that type is the one proved by the discrete-series circle-mean argument of the earlier real chapter. Finite linear combinations with coefficients ϕ_α give the linear map `Filling.Fits.poincareMap`.

The Petersson identity is also exact. For the two single-coordinate columns and a finite integral translation k' of reduction s, unfold the outer Poincaré sum in the integral on $\Gamma\backslash G(\mathbb{A})$. Absolute integrability follows from the uniform absolute sum and boundedness of the other form on the finite-volume quotient. Translate the finite variable in the unfolded integral to the fixed coset of the first class. The second finite indicator then selects exactly the set $\texttt{termsOver}(c, s, c')$ (inversion of the rational index converts its real argument into $\gamma_\infty^{-1}y$). Each selected term has finite integral the volume of the standard principal level. The real coordinate changes are Haar preserving, since they are a unitary sign-frame conjugation followed by left translation. The remaining real factors are precisely $\beta_c(y)\beta_{c'}(\gamma_\infty^{-1}y)W_i(y)\overline{W_{i'}(\gamma_\infty^{-1}y)}$. Theorem 27.3 also ensures their sum of absolute integrals is finite. Thus one obtains the exact principal-level volume times the corresponding entry of `amplitudeSum`, as asserted by `petersson_poincare`.

28.8 Scope and remaining interfaces

The cut and standard-level existence are now proved with their exact fields, including the weighted majorant and the finite top-section measure. The simultaneous numerical inequalities of the actual filling, and the matrix and reproduction statements of `Fits`, now follow from the exact degree, mass and arithmetic class formulas. This closes the previously separate local-existence, arithmetic admissibility and Poincaré convergence interfaces in B04, B05, B08 and B15, subject to the previously specified local coefficient and class formulas. It does not establish the missing `HasBandLimits` field of the actual real systems, the quantitative finite and real gauge transition and Fourier estimates, or the weighted inverse-index and energy bounds. Thus the full global spanning statement is not inferred here. The good finite datum of A06–A07 uses the glued lattice, primitive frame and finite residue projector already discussed in Section 16.6 and the subsequent admissible-combination proof; the fitted Bessel coordinate and level compatibility are established here, but we do not assert that any separate missing finite projector realization or theta-period identity is thereby closed.

Chapter 29

Uniform floor estimates at finite and complex places

We extend the identity-head calculation of Chapter 26 to arbitrary heads. The essential distinction is between a *direct* pivot weight, integrated over close positions, and an *inverse* pivot weight, integrated over unipotent translations with two simultaneous nearness conditions. The latter is needed before one can use an estimate on a single floor in the full group integral. The same argument works for the finite and complex gauges of Chapter 23. We retain the source's coarse constants: they are deliberately large, and are important in comparing the final exponents.

29.1 Conventions and the tower

Write $h = \varrho$, $n = h + 1$, $k = \dim X_0$, and let A be the cardinality of the orthonormal index set of a level. The figure A need not equal k: in the complex gauge $A = k - 1$. Put

$$d_a = k + 1 + 2(h - a - 1), \qquad s = A - (3n + 1 + m_{\mathrm{lev}}), \qquad M_a(u) = \mathrm{md}(u)^{d_a + 1/2} \quad (a < h).$$

We use $s \geq 1$ throughout. The parameter m_{lev} is the additional basis parameter of the level, not the number of hyperbolic planes in X_0; for a finite level with that latter number m, $m_{\mathrm{lev}} = m + 1$. Let B_i be the gauge size sets, $\Lambda \geq 2$ its size constant, γ its self-adjoint unit, and c its place constant (1 at the finite places and π^2 at the complex place, with the source's character). Scalar products, inverses of unitary matrices and matrix products obey the gauge estimates of Chapter 23: two matrix sizes i, j give a product of size $i + j + 1$; a unitary inverse costs four; a sum of at most $r + 1$ entries of size i costs one. The fixed Gram data have size one.

Let Q_j fix the first j isotropic frame vectors, H_j fix also their dual vectors, and $U_{<j}$ be the unipotent product attached to those j steps. Thus $Q_j = U_{<j} \rtimes H_j$. Write ν_a for the measure on the Heisenberg factor $N_a \subset H_a$, $\eta_{<j}$ for the convolution of these measures for $a < j$, θ_a for the floor representative measure, $\lambda_j = \theta_{j-1} * \cdots * \theta_0$, and

$$\mu_j = \eta_{<j} * \lambda_j, \quad \vartheta_j = \theta_{h+1} * \cdots * \theta_j, \quad \mu = \mu_{h+2}.$$

The notation at the last two floors refers to the terminal quotient and stabilizer measures. The convolution identities are identities for nonnegative integrands, and then for integrable complex integrands. In particular, moving H_j across $U_{<j}$ preserves the latter's measure: each vector block is transformed by an isometry of the residual Hermitian space, whose determinant has modulus one, and each skew central coordinate is unchanged. This is the structural reason

for the rearrangements of convolution factors below. The Weyl element w_p^j exchanges a frame vector e_a with $p_a e_a^*$ and exchanges e_a^* with $\overline{p}_a^{-1} e_a$ for $a < j$. It commutes with H_j. A head means precisely that every inverse prefix product $(p_0 \cdots p_{l-1})^{-1}$ belongs to B_1 for $l \le j$. In formulas where a Weyl element and its inverse are used interchangeably the head parameter of the inverse is $\bar{p}$.

Use the pivot sets $I_a(g,x)$ of Chapter 23:

$$I_a(g,x) = \{z_{e_a^*} : z_{e_b^*} = 0 \ (b < a), g^{-1}z \in B_1^r, \ (g^{-1}z - (gx)^{-1}z)/\gamma \in B_1^r\}.$$

The ball in this definition is the actual open box, not its boundary. The sets are open, contain a neighborhood of zero, and are bounded; their volumes are finite and positive. Set

$$P_j(g,x) = \prod_{a<j} \mathrm{vol}(I_a(g,x))^s, \qquad J_j(g,x) = \frac{\mathbf{1}_{\text{near } j}(g)}{P_j(g,x)}.$$

The ath pivot is invariant under simultaneous left multiplication by Q_{a+1}, since the first $a+1$ dual coordinates are the Hermitian pairings with vectors fixed by that subgroup. Consequently P_j is Q_j-invariant.

For a pairing kernel K put $K_j(g,y) = \int K(\sigma g, \sigma y)\, d\vartheta_j(\sigma)$. The hypotheses on K in this section are exactly the following: these upper averages exist absolutely; the simultaneous left action of $Q_j \cap H_b$ preserves K_j when the twist commutes with H_b and $j \ge b$; each nonzero K_j has first argument near j; and the unipotent equivariance holds on $U_{<a}$ for K_{a+1}, in the form

$$K_{a+1}(ug, u'y) = \psi(\gamma^{-1}[\mathrm{fw}(Du'D^{-1}) - \mathrm{fw}(u)])K_{a+1}(g,y).$$

Here D is the twist ($D = 1$ for the self-pairing). For the actual kernels these are consequences of the Bessel law and the quotient Haar laws: multiplying the upper average simultaneously on the left by the next stabilizer preserves the upper quotient measure; the terminal stabilizer has the stated law on the cut; the claim follows successively down the upper tower. The absolute upper averaging is one of the hypotheses of `Level.IsPairingKernel`; in applications it is supplied by the cut and terminal bound, not by the estimates that we shall prove.

At levels with the equivariance define

$$A_a^K = P_a |K_{a+1}|^2, \qquad B_a^K(g,x) = \int (A_a^K + B_{a+1}^K)(\sigma g, x)\, d\theta_a(\sigma),$$
$$B_{h+1}^K = 0, \qquad Z_a^K(x) = \int B_a^K(g,x) d\lambda_a(g).$$

Here K_{a+1} in a one-variable expression abbreviates $K_{a+1}(g, gx)$. These quantities are nonnegative. By the quotient law B_a^K is Q_a-invariant and supported on near a; A_a^K is Q_{a+1}-invariant and supported on near $a+1$. The definitions and Tonelli give

$$\frac{B_a^K(g,x)}{P_a(g,x)} \ge \int |K_{a+1}(\sigma g, \sigma g x)|^2 d\theta_a(\sigma), \qquad Z_a^K = \int A_a^K \, d\lambda_{a+1} + Z_{a+1}^K. \tag{F.1}$$

No finiteness assertion is used in this recursion.

29.2 A quantitative Heisenberg integral

The residual vector coordinate v of N_a ranges over V_{a+1}, of dimension d_a over the place ring. With $\tau + \bar{\tau} = 1$, its element $N_a(v,\beta)$, β skew, acts by

$$z \longmapsto z + z_{e_a^*} v + \big((\beta - \tau\langle v, v\rangle) z_{e_a^*} - \langle v, z\rangle\big) e_a. \tag{F.2}$$

Conjugation by $\sigma \in H_{a+1}$ sends v to σv, and $\mathrm{fw}(\sigma N_a(v,\beta)\sigma^{-1}) = -\langle v, \sigma^{-1} e_{a+1}\rangle$ (with the final residual frame vector used at the top). Project the column e_a^* of $(\nu N_a(v,\beta)^{\pm 1} - 1)/u$ onto the residual space and the e_a coordinate, where $\nu \in U_{<a}$ and u is a unit. The preceding unipotent factors change this projection only in the already discarded positive coordinates. If the column has size i, then v/u has size i, and, for fixed v, the difference of any two eligible skew parameters divided by u belongs to B_{i+1}. The skew measure bound (translation, followed by skew dilation) is at most $\mathrm{md}(u)^{1/2}\Lambda^{i+2}$; it holds also for an empty fiber. Indeed one translates a nonempty fiber by an eligible point and uses the gauge skew-ball estimate on the resulting difference set. In the finite case the skew lattice has modulus $\mathrm{md}(u)^{1/2}$ in this assertion for self-adjoint u; for a general unit the same estimate is the bound for the intersection of an u-dilate with the skew line, as in the place skew-measure law. Vector dilation contributes $\mathrm{md}(u)^{d_a}$.

It follows that every nonnegative Φ supported on the above column condition and bounded by $C\Psi(v/u)$ satisfies

$$\int \Phi(N_a(v,\beta))\, d\nu_a \le C M_a(u) \Lambda^{i+2} \int_{B_i^{d_a}} \Psi(v)\, dv. \tag{F.3}$$

This is `lintegral_heisenbergMeasure_le`. The exponent $1/2$ in M_a cannot be replaced by 1: it is the skew-coordinate contribution.

We also recall the consequence of the gauge common-weight estimate (Chapter 23). The admissible common-weight function ϕ on the orthonormal coordinates has its sth moment on the box of depth $i+4$ at most $\Lambda^{99n(i+5)}$. The sliced measure bound on the whole floor or on the floor quadric cut at depth i is $\Lambda^{99n(i+6)}$ times orthonormal-coordinate Lebesgue or adelic Haar measure. Projecting by the Gram matrix and the orthonormal embedding costs at most four. Thus

$$\int_{\text{depth } i} \phi(E^* J v)^s \, d\delta(v) \;\le\; \Lambda^{99n(2i+11)}, \tag{F.4}$$

where δ may be either the ambient floor volume or the quadric quotient measure. In particular, for every g near a the inverse-pivot estimate

$$\int \mathbf{1}_{\text{near } a+1}(\sigma g)\, \mathrm{vol}(I_a(\sigma g, x))^{-s}\, d\theta_a(\sigma) \le \Lambda^{10^4 n} \tag{F.5}$$

holds. To see it, first replace g by a box representative using the quotient law. On the support there is a box representative $q\sigma g$ with $q \in Q_{a+1}$; hence $\sigma^{-1} e_a = g(q\sigma g)^{-1} e_a$ has size seven. Apply the common-weight estimate with no additional test rows to $z = \sigma E w$ for w in the admissible residual subspace; its pivot equals $\langle \sigma^{-1} e_a, E w\rangle$. Thus the inverse pivot is bounded by $\phi(E^* J \sigma^{-1} e_a)$, and (F.4) at $i = 7$ proves (F.5), since $99(25) < 10^4$. The same quotient argument without a pivot bounds the floor mass of the near set by $V_a = \Lambda^{1287n}\,\mathrm{vol}(B_{11})^A$. A larger bound $V \ge 1$ valid on all floors may be used instead.

29.3 Direct and inverse weights on an arbitrary prefix

Lemma 29.1 (Uniform direct weight)**.** *Let* $\Omega = \{x \in \text{positions} : (x-1)/\gamma \text{ has size } 1\}$. *For* g *near* j *and arbitrary* y,

$$\int_{U_{<j}} \mathbf{1}_\Omega(g^{-1} u y) P_j(g, g^{-1} u y)\, d\eta_{<j}(u) \le \prod_{a<j} \big(M_a(\gamma) c^A \Lambda^{4600n}\big).$$

Proof. Move g into the box by Q_j, using preservation of Haar measure under the H_j action. If the support is nonempty, right translate the unipotent variable so that $x_0 = g^{-1} y \in \Omega$. For

every other supported point $x = g^{-1}uy$, the matrix

$$(u^{-1} - 1)/\gamma = g\big((x_0 - x)/\gamma\big)(gx)^{-1}$$

has size fourteen: $(x_0 - x)/\gamma$ has size two, x has size at most three, gx has size at most five, its inverse at most nine. The indicated products cost respectively one. Factor $u = \nu N_a(v, \beta)$, $\nu \in U_{<a}$, with $a = j - 1$; the factors can be integrated in either order by the modulus-one normalization. Its inverse can be written $\nu' N_a(v, \beta)^{-1}$ with $\nu' \in U_{<a}$, so (F.3) applies at depth fourteen.

Use the common-weight estimate with reference (g, x_0) and zero test rows. If z is an eligible vector for $I_a(g, x)$, with pivot b, and Ew is an eligible vector in the reference residual subspace, the pairing discrepancy of the two close pairs divided by γ is $\bar{b}\langle v, Ew\rangle/\gamma$, up to a sign. This follows by inserting (F.2); earlier dual coordinates vanish, and preceding unipotent factors disappear on the residual quotient. The product-bound part of the common-weight estimate gives $\mathrm{vol}(I_a(g, x)) \le \phi(E^* J(v/\gamma))$. Use the inductive direct-weight estimate first on the prefix integral at fixed $N_a(v, \beta)y$, since its earlier weights have precisely the same form and the indicator is still the close-position indicator. The last common-weight bound and column condition are independent of the prefix integration variable, so (F.3) and (F.4) apply to the remaining integral. The last factor costs at most $M_a(\gamma)\Lambda^{16+99n(39)} \le M_a(\gamma)c^A\Lambda^{4600n}$. Iteration proves the assertion, starting with the empty weight at $j = 0$. □

The head estimates used for inverse weights are purely matrix estimates. Suppose g and $w_p^{-1}g$ are both near j, and qg is in the box, $q \in Q_j$. For each $l \le j$ take the minor in the first l positive-column block of $qg(q' w_p^{-1} g)^{-1}$, with its final row replaced by the row of the desired entry, where $q' \in Q_j$ moves the second argument into the box. This matrix has size seven. The first l columns are obtained from the columns $p_a q e_a^*$ by right multiplication by the unitriangular positive-frame block of $(q')^{-1}$, whose determinant is one. The first $l - 1$ dual rows of the latter columns are triangular with diagonal entries $p_0, \dots, p_{l-2}$; expansion along the last column gives the minor

$$D_l(q e_{l-1}^*)_i, \qquad D_l = p_0 \cdots p_{l-1}.$$

It has size at most $8l \le 9n$. Taking the final dual row gives $D_l \in B_{9n}$. The head condition $D_l^{-1} \in B_1$ then proves that the individual entries p_a, p_a^{-1} and $q e_a^*$ have size at most $10n$ (with the slightly sharper bound $8l$ before the final multiplication). The same argument applies to the inverse Weyl element with parameters $\bar{p}$. Solving the triangular frame equations, or applying the matrix recovery statement below with $\gamma = 1$, now shows that if both points are near j there is a common $t \in H_j$ such that tg has size $60n^2$ and $t w_p^{-1} g$ size $99n^2$. In fact write $q = t\nu$ with $\nu \in U_{<j}$; bounds on ν's first j dual columns recover ν by triangular elimination and the unitary inverse relation. Multiplication by w_p^{-1} costs at most the size of its entries plus one. These bounds are the intentionally slack form of `head_entries_mem_ofSize`, `hasMatSize_of_isNear` and `exists_hasMatSize_of_isNear` in `New/RealPlace/Size.lean`.

Lemma 29.2 (Inverse prefix weights)**.** *Put $L_0 = \Lambda^{10^6 n^3}$. Let p be a head of length $j \le b$. If Φ is nonnegative, measurable, Q_b-invariant and zero off the near-b set, then*

$$\int J_j(w_p^j z, x)\Phi(z)\, d(\eta_{<j} * \rho)(z) \le L_0 \prod_{a<j} M_a(p_a^{-1}) \int \Phi(z)\, d\rho(z) \tag{F.6}$$

for every representative measure ρ.

Proof. By Tonelli it suffices to fix the point of ρ. The factor Φ is constant on the unipotent orbit; if nonzero the point is near b, thus near j. Decompose a Q_j move into its H_j and unipotent components, commute the former past the Weyl element, and use invariance of $\eta_{<j}$. We reduce to a point g in the box.

For the step $a < j$, write $z = N_a(v,\beta)\nu g$, $\nu \in U_{<a}$. Earlier inverse factors are unchanged by N_a, since it commutes with w_p^a and belongs to Q_a. The test rows in the inverse-pivot common-weight estimate are $(\nu E)_{e_b}$ for $b < a$. Hence $w_p^{a+1} N_a(v,\beta)\nu E w$ has its first a dual coordinates zero, and its ath dual coordinate is $-p_a\langle v, Ew\rangle$. Its pair of inverse images is exactly the reference close pair. Set $u = (-\bar{p}_a)^{-1}$. The lower-bound portion of the common-weight estimate proves

$$\operatorname{vol}(I_a(w_p^{a+1} z, x))^{-1} \le \phi(E^* J(v/u)).$$

The support forces both z and $w_p^{a+1} z$ near $a+1$; the first is always near since ν, N_a belong to its stabilizer. The preceding head-entry estimate applied to the inverse Weyl element, and to $\nu^{-1} N_a^{-1}$ moving z back into the box, gives size $21n$ for the column of $(\nu^{-1} N_a^{-1} - 1)/u$ at e_a^*. Apply (F.3) and (F.4). The cost is

$$M_a(p_a^{-1})\Lambda^{21n+2+99n(42n+11)} \le M_a(p_a^{-1})\Lambda^{6000n^2}.$$

Integrate the last Heisenberg coordinate first; the earlier inverse factors and near condition have already been retained. Iterating at most h times gives (F.6). □

29.4 Fourier estimates on the tower

We give a version of the sliced Fourier estimate with its precise dependence on the centers. Let Φ be a nonnegative function on N_a, supported on a set S, and suppose

$$\Phi(u) \le C\,|K_{a+1}(u^\epsilon x, uy)|^2, \qquad \gamma^{-1}(\operatorname{fw}(DuD^{-1}) - \operatorname{fw}(u^\epsilon)) = \xi \operatorname{fw}(u). \tag{F.7}$$

Here ϵ is a nonnegative integer and the second identity holds on N_a. Suppose that for each $m \in S$ one can choose $t \in Q_{a+1} \cap H_a$, $q \in U_{<a+1}$ with tqx of size i, and that for each $u \in S$ there is $\nu' \in U_{<a}$ for which

$$((\nu' tm(tu)^{-1} - 1)e_a^*)/v_0 \in B_j^r. \tag{F.8}$$

Then

$$\int \Phi\, d\nu_a \le C(c/\operatorname{md}(\xi v_0))^A M_a(v_0)\Lambda^{100n(i+j+31)} \int |K_{a+2}(\sigma x, \sigma y)|^2\, d\theta_{a+1}(\sigma). \tag{F.9}$$

Here and below a Fourier estimate with a frequency in the denominator is invoked only for a unit frequency. To prove (F.9), fix a supported center m (the assertion is trivial if the integrand vanishes almost everywhere). Write $t = t_0 r$, $t_0 \in H_{a+1}$, $r \in N_a$. Translate the Heisenberg variable by rm, and conjugate by t_0; these operations preserve ν_a. Formula (F.8) is now the single-column condition of (F.3), with scale v_0 and depth j. The equivariance (F.7) and (F.2) express the amplitude in (F.7) as the Fourier integral

$$\int \psi(-\xi\overline{v_0}\langle v + v', t_0\sigma^{-1} e_{a+1}\rangle) K_{a+2}(\sigma x, \sigma y)\, d\theta_{a+1}(\sigma).$$

The fixed vector v' is the translated center in scaled vector coordinates; the skew variable no longer occurs in the phase. Its source amplitude is a function on the quadric, because K_{a+2} is invariant under the stabilizer on the left of the representative measure. Nonzero amplitude

forces $t_0\sigma^{-1}e_{a+1}$ of size $i+6$: indeed the size-i representative tqx may be compared with the box representative of σx, with the unipotent factor fixing e_{a+1} removed. This is the same orbit-vector estimate as in the proof of (F.5), now with a size-i first representative rather than size one. By the sliced Plancherel inequality of Proposition 23.7, the size-j target integral is bounded by

$$\Lambda^{99n(i+12)+99n(j+6)}(c/\operatorname{md}(\xi v_0))^A \int |K_{a+2}|^2\, d\theta_{a+1}.$$

After (F.3), the additional size factor is Λ^{j+2}. The sum of the size exponents is at most $100n(i+j+31)$. This proves (F.9), including its center hypothesis, and explains why the decay exponent is the orthonormal rank A, not the dimension d_a of the Heisenberg vector variable. This is the statement `parseval_heisenbergMeasure_of_box`.

29.5 Integrated square norms

Define

$$\mathscr{N}(K) = \iint |K(g,y)|^2\, d\vartheta_0(g) d\vartheta_0(y), \qquad C_a = M_a(\gamma) c^A \Lambda^{4600n}.$$

If μ is Haar and the twist commutes with H_b, then

$$\int_\Omega Z_b^K(x)\, d\mu(x) \le n \left(\prod_{a<h} C_a \right) \mathscr{N}(K). \tag{F.10}$$

This is the source's `lintegral_selfNorm_le`; the result of Chapter 26 is its identity-head specialization. We spell out the induction because it is used for the twisted kernel as well.

For $l \ge b$ factor $\mu = \eta_{<l} * \rho_l$, with $\rho_l = \eta_{l,\dots,h-1} * \vartheta_0$. Put

$$W_l(g,y) = \int_{U_{<l}} \mathbf{1}_\Omega(g^{-1}uy) d\eta_{<l}(u), \qquad T_l(g,y) = \mathbf{1}_{\{W_l(g,y)>0\}} |K_{l+1}(g,y)|^2.$$

The W_l may be infinite; the indicator is intentional. Changing variables $x = g^{-1}uy$ in the lth summand in (F.1), and using the unipotent law (in modulus), Lemma 29.1 gives

$$\int_\Omega \int A_l^K\, d\lambda_{l+1} d\mu \le \left(\prod_{a<l} C_a \right) \iint T_l(g,y) d\lambda_{l+1}(g) d\rho_l(y). \tag{F.11}$$

If $W_l = 0$, the inner unipotent integral is zero by Tonelli; when it is positive, the support of the amplitude gives nearness of g, as required for the direct-weight lemma.

We next pass from T_l to T_{l+1}. For the fixed g, y integral in the next Heisenberg variable, take the supported set with $W_l(g,uy) > 0$ and $K_{l+1}(g,uy) \ne 0$. For a supported m, choose a near-$l+1$ representative tqg in the box, with $t \in Q_{l+1} \cap H_l$ and $q \in U_{<l}$, possible by the semidirect factorization. For m, u supported choose $\nu_m, \nu_u \in U_{<l}$ such that $x_m = g^{-1}\nu_m m y$, $x_u = g^{-1}\nu_u uy$ belong to Ω. The identity

$$(tqg)x_m x_u^{-1}(tqg)^{-1} = tq\nu_m m u^{-1}\nu_u^{-1} q^{-1} t^{-1}$$

and the normality of the prefix unipotent subgroup show that its right side is $\nu' tm(tu)^{-1}$ with $\nu' \in U_{<l}$. Subtract the identity and divide by γ. Since the first matrix representative is in the box, the same calculation as in Lemma 29.1 gives size fourteen. Thus (F.8) holds with $i = 1$, $j = 14$, $v_0 = \gamma$. Since D commutes with $N_l \subset H_b$ for $l \ge b$, (F.7) holds with $\epsilon = 0, \xi = \gamma^{-1}$. Formula (F.9) costs precisely at most C_l, since $100(1+14+31) = 4600$.

Now if $W_{l+1}(g,y)=0$, the integral on the left vanishes, because the convolution definition of W_{l+1} integrates $W_l(g,uy)$ over this Heisenberg variable (with precisely the same ν_l). Otherwise insert the indicator of $W_{l+1}>0$ on the right of (F.9). This indicator is invariant under simultaneous H_{l+1} multiplication, by conjugation invariance of the prefix measure. Moving the next θ_{l+1} factor from the second variable to the first in the integral uses its right-invariance modulo the upper stabilizer, and the fact that K_{l+2} and W_{l+1} are unchanged upon the independent unipotent translations being integrated. More explicitly, the quotient integral in the first variable replaces λ_{l+1} by $\theta_{l+1}*\lambda_{l+1}$; the independent suffix measure in the second is unchanged by left translation from H_{l+1}, since its convolution gives the remaining factors of Haar. This proves

$$\iint T_l\, d\lambda_{l+1}d\rho_l \le C_l \iint T_{l+1}\, d\lambda_{l+2}d\rho_{l+1}.$$

At $l=h$ the remaining stabilizer has mass at most one on the relevant support. Cauchy–Schwarz in the terminal average costs no factor; the terminal double integral is bounded by $\mathscr{N}(K)$, with the two measures ϑ_0. Consequently each of the at most n summands in (F.1) costs at most the same $\prod_{a<h} C_a\mathscr{N}(K)$. This proves (F.10).

29.6 The central estimates for forms

Let W be cut Bessel of frequency γ^{-1}, W' Bessel of the same frequency, and put

$$K(g,y)=\langle W(g),W(y)\rangle,\quad K'(g,y)=\langle W(g),W'(Dy)\rangle,\quad D=(w_q^j)^{-1}w_p^j.$$

Assume p,q are heads of length $j\le h$. Write

$$\Delta_j(x)=\left(\prod_{a<j}M_a(p_a^{-1})\right)^2 L_0^3Z_0^K(x)Z_j^{K'}(x),\qquad \kappa_j=\prod_{a<j}\mathrm{md}(p_aq_a)^{d_a/2}.$$

We first record the precise integrated estimate on a set S of positions in the size-one close neighborhood for a second gauge of constant Λ' (the level, γ and its measures are unchanged). If $F^2\le e^2\Delta_j$ on S, then

$$\kappa_j\int_S F(x)d\mu(x)\le e\Theta\left(\prod_{a<h}M_a(\gamma)\right)c^{Ah}\Lambda^{1500000n^3+1}(\Lambda')^{50000n^2},\tag{F.12}$$

where Θ^2 bounds the product of the two untwisted pairing norms.

Indeed Cauchy–Schwarz in x and (F.10) bound the product of the two integrated self norms by the product of the two pairing norms times the square of $n\prod_{a<h}C'_a$. The twist D acts on the second quotient measure with square-integral factor $\prod_{a<j}\mathrm{md}(p_a/q_a)^{d_a+1}$. To verify the factor put $r_a=p_a/q_a$. Conjugation by D sends the vector coordinate of N_a to $\bar r_aD_{\mathrm{res}}v$, where D_{res} is unitary, and sends its skew coordinate to $r_a\bar r_a\beta$. Their Jacobians are respectively $\mathrm{md}(r_a)^{d_a}$ and $\mathrm{md}(r_a)$. Thus conjugation on unipotent Haar has Jacobian $\delta=\prod\mathrm{md}(r_a)^{d_a+1}$. The decomposition $\mu=\eta_{<h}*\vartheta_0$ is unchanged by left translation by D; its pushed unipotent measure is $\delta^{-1}\eta_{<h}$, so the induced quotient measure pushed by D is $\delta\vartheta_0$. This proves the change of variables `lintegral_weylRatio_mul`. On taking square roots, each a contributes, including κ_j and $M_a(p_a^{-1})$, the factor

$$\mathrm{md}(p_aq_a)^{d_a/2}\,\mathrm{md}(p_a)^{-d_a-1/2}\,\mathrm{md}(p_a/q_a)^{(d_a+1)/2}=\mathrm{md}(q_a)^{-1/2}.$$

The product is at most Λ, by the head condition at length j and the gauge modulus bound. Finally $n(\Lambda')^{4600nh} \le (\Lambda')^{50000n^2}$ for $\Lambda' \ge 2$. This proves (F.12). In particular no individual p_a or q_a modulus remains; their cancellation is indispensable at both real and finite places.

A second Cauchy–Schwarz argument bounds a single stack integral. Suppose $f(g) \ne 0$ implies both arguments near at j, and

$$f(g)^2 \le CL_0 \frac{B_j^K(g,x)}{P_j(g,x)} \frac{B_j^{K'}((w_p^j)^{-1}g,x)}{P_j((w_p^j)^{-1}g,x)}.$$

Then

$$\left(\int f \, d\mu_j\right)^2 \le C\Delta_j(x). \tag{F.13}$$

In Cauchy–Schwarz cross the denominators: the first factor is $B_j^K(g,x)J_j((w_p^j)^{-1}g,x)$, and the second is $B_j^{K'}((w_p^j)^{-1}g,x)J_j(g,x)$. The second may be moved by w_p^j because its expression as a function of $(w_p^j)^{-1}g$ is invariant under H_j; the Weyl invariance of the stack on such functions follows directly by disintegrating Haar as $\chi_j * \mu_j$, where χ_j is the remaining H_j Haar measure. Since w_p^j commutes with H_j, left translation by that element preserves the fiber measure; left invariance of full Haar then preserves the induced quotient measure on $H_j\backslash G$. Apply (F.6) with heads $\bar{p}, p$ respectively. Each factor costs $L_0 \prod_{a<j} M_a(p_a^{-1})$, and (F.1) gives $Z_j^K \le Z_0^K$. This establishes (F.13).

Here are the three resulting pointwise estimates. Let $\mathfrak{m}_j(x)$ denote the form majorant (the inside moduli of both floor averages, integrated on μ_j), and $\mathfrak{f}_j(x;\omega)$ the form modulus (the absolute value of the weighted double floor integral, then integrated on μ_j). For $V \ge 1$ bounding the mass of the relevant near-floor set:

1. If $0 < \sigma < 1$ and every elementary pair $K(y, yx) \ne 0$ satisfying the two twisted nearness conditions
$$(w_p^j)^{-1}y \text{ near } j, \qquad (w_q^j)^{-1}yx \text{ near } j$$
has some $a \ge j$, $a < h$ with $\mathrm{vol}(I_a(y,x)) \le \sigma$, then
$$\mathfrak{m}_j(x)^2 \le \sigma^s V^2 \Delta_j(x) \tag{F.14}$$
provided W' is also zero off near j. For $\sigma \ge 1$ no cap is required.

2. If $\psi(\eta z) = \psi(\xi z)$ on B_{200n^2}, with ξ a unit, and $\omega(g) = \psi(\eta\langle e_j, ge_j\rangle)$, then
$$\mathfrak{f}_j(x;\omega)^2 \le (c/\operatorname{md}(\xi))^A \Delta_j(x), \qquad W' = W. \tag{F.15}$$

3. If $j = i+1$, $\xi = \gamma^{-1}(p_i^{-1} - q_i^{-1})$ is a unit, then
$$\mathfrak{f}_j(x;1)^2 \le (c/\operatorname{md}(\xi))^A \Delta_j(x), \qquad W' = W. \tag{F.16}$$

We prove these statements. For (F.14) the single floor Cauchy–Schwarz bound (F.5) iterates whenever the cap remains to be encountered. Precisely, write $C_l(g) = \int |K_{l+1}(\sigma g, \sigma g x)| d\theta_l$. Weighted Cauchy–Schwarz gives

$$C_l(g)^2 \le \Lambda^{10^4 n} \int \mathrm{vol}(I_l(\sigma g, x))^s |K_{l+1}(\sigma g, \sigma g x)|^2 d\theta_l.$$

Split this last integral according as the pivot volume is at most σ or greater. On the first part its contribution is at most $\sigma^s B_l/P_l$. On the second all elementary pairs in the upper averages must satisfy the cap at a later floor, because I_l is invariant under the upper subgroup. Bound

$|K_{l+1}|^2$ by C_{l+1}^2 and induct. The product P_{l+1}/P_l is exactly the pivot factor, so the second contribution is paid for by the B_{l+1} term of (F.1). At the top the second part vanishes by the hypothesis. This proves $C_j^2 \le \sigma^s \Lambda^{10^4 n(h-j)} B_j/P_j$. The ordinary Cauchy–Schwarz estimate is $C_j^2 \le V B_j/P_j$. Use it for the second kernel. Its nonzero pivot modulus supplies elementary pairs with the indicated twisted nearness, which remain near when multiplied by any element of H_j commuting with the heads. Thus the cap argument applies to all the first-kernel pairs on the support. Since $\Lambda^{10^4 n(h-j)} \le L_0$, $V \ge 1$, (F.13) proves (F.14).

For (F.15), the double floor integral at g is

$$\iint \psi(\eta\langle \sigma^{-1}e_j, \tau^{-1}e_j\rangle) K_{j+1}(\sigma g, \sigma g x)\overline{K'_{j+1}(\tau(w_p^j)^{-1}g, \tau(w_p^j)^{-1}gx)}\, d\theta_j(\sigma) d\theta_j(\tau).$$

Nonzero integral forces both g and $(w_p^j)^{-1}g$ near j. The matrix estimate preceding Lemma 29.2 gives $t \in H_j$ making their representatives of sizes $60n^2$ and $99n^2$. Their orbit vectors $t\sigma^{-1}e_j, t\tau^{-1}e_j$ are therefore supported at depths $60n^2+6, 99n^2+6$; their Hermitian pairing has size at most $200n^2$ (the Gram matrix and the two sums cost three). Replace η by ξ, preserving the integrand. The sliced Fourier estimate between these two quadric measures, followed by Cauchy–Schwarz against the other amplitude, bounds its squared modulus by

$$(c/\operatorname{md}\xi)^A \Lambda^{99n(159n^2+24)} \left(\int |K_{j+1}|^2 d\theta_j\right)\left(\int |K'_{j+1}|^2 d\theta_j\right).$$

The size power is at most L_0. Now use (F.1) and (F.13). This proves (F.15).

For (F.16), put $g' = (w_p^j)^{-1}g$ and

$$R(v) = P_j(v,x) J_i(w_p^i v, x), \qquad U(v) = \mathbf{1}_{\text{near } j}(w_p^j v) R(v) |K'_j(v, vx)|^2.$$

The integrand for $\omega = 1$ is the product of the two jth upper-averaged kernel moduli. Its square equals $A_i^K(g,x) J_j(g',x) U(g')$ on its nonzero support: the earlier pivot products cancel, since $w_p^j (w_p^i)^{-1}$ acts at floor i and fixes the first i dual coordinates. Cauchy–Schwarz therefore gives two stack integrals. The first, by (F.6), is at most $L_0 \prod_{a<j} M_a(p_a^{-1}) Z_0^K(x)$. For the second, use Weyl invariance of the stack for the H_j-invariant function U, and split off its last Heisenberg factor. At fixed $v \in U_{<i}\lambda_j$, $R(uv) = R(v)$ for $u \in N_i$. This follows from invariance of P_j under Q_j and commutation with w_p^i.

On the support of $U(uv)$ both uv and $w_p^j uv$ are near j. The head size estimate applies with the inverse head $\bar{p}$; also p_i has size at most $10n$. To verify the Fourier center condition choose a supported m and a $t \in H_j$ with tmv of size $60n^2$. For each other supported u similarly choose t_u with $t_u uv$ of size $60n^2$. The column at e_i^* of $tm(tu)^{-1}$ equals the same column of $(tmv)(t_u uv)^{-1}$, since both t, t_u fix e_i^*. Its difference from the identity divided by p_i^{-1} has size at most $120n^2 + 6 + 10n \le 140n^2$. Thus (F.8) holds with $q = m$, $i_{\text{size}} = 60n^2$, $j_{\text{size}} = 140n^2$, and $v_0 = p_i^{-1}$. The phase identity (F.7) for $\epsilon = 1$ holds with frequency $-\xi p_i$: conjugating by D multiplies the ith simple-root coordinate by p_i/q_i, so $\gamma^{-1}(p_i/q_i - 1) = -\xi p_i$. Formula (F.9), whose size factor is at most L_0, gives

$$\int U(uv) d\nu_i(u) \le L_0 (c/\operatorname{md}\xi)^A M_i(p_i^{-1}) R(v) \int |K'_{j+1}(\sigma v, \sigma vx)|^2 d\theta_j.$$

The product with P_j in R is absorbed by $B_j^{K'}$ in (F.1). Apply (F.6) to its remaining inverse weight $J_i(w_p^i v, x)$, with $b = j$; this costs $L_0 \prod_{a<i} M_a(p_a^{-1})$ and leaves $Z_j^{K'}$. Multiplying the two Cauchy–Schwarz factors gives exactly (F.16), with three powers of L_0. The unequal-head estimate is therefore not just the nonstationary Fourier estimate of the preceding item; it comes from the root-frequency difference, and is essential to the upper-floor induction.

29.7 The matrix recovery argument and summation over shells

We give the missing matrix recovery lemma explicitly. Let $\gamma = \bar{\gamma}$ be a unit of size one, $u \in U_{<j}$, and assume that for each $a < j$ there is a vector v_a of size C with

$$(v_a)_{e_b^*} = \delta_{ab} \quad (b \le a), \qquad (v_a - uv_a)/\gamma \text{ of size } C.$$

Then $(1-u^{-1})/\gamma$ has size $C + 4 + (2C+8)j$.

Here is the back-substitution calculation. For an arbitrary matrix X, suppose each Xv_a has size D, and all entries in a row outside the first j dual columns have size D. Write the equation for that row applied to v_a, separating its pivot coefficient from all other coefficients. Solving for the coefficient in column e_a^* uses only the nondual columns and the dual columns with index $> a$, since the earlier dual coordinates vanish. Each step costs at most $C+3$: size C for v_a, one for the product sum, one for the subtraction (and an extra unit allows replacing v_a by $v_a - e_a^*$). Descending induction therefore bounds the row at depth $D + (C+3)j$. Apply this first to $X = (1-u)/\gamma$ in those rows that pair, through the Gram matrix, with a column outside the first j dual columns. In those rows the required starting entries are zero, by the triangular form of $U_{<j}$ (on the quotient by its first j positive coordinates the unipotent is the identity when the first j dual coordinates vanish). This gives size $C + (C+3)j$ in the stated rows. Since

$$J(1-u^{-1})/\gamma = ((1-u)/\gamma)^* J,$$

the Gram matrix and its inverse transfer the resulting row bound to the nondual columns of $(1-u^{-1})/\gamma$, costing four. Use the second back substitution on the vectors uv_a, which have the same earlier dual coordinates, have size $C+2$ by $uv_a = v_a - \gamma((1-u)v_a/\gamma)$, and whose image under $(1-u^{-1})/\gamma$ has size C. The second induction costs $(C+5)j$. The overall bound is as asserted, proving `hasMatSize_one_sub_inv`.

Now let $g, y = gx$ be an elementary pair, with unipotents u, v moving g, y to box representatives $a = ug, b = vy$, and $(b-a)/\gamma$ of size two. Let the two twisted arguments be near at j. Take a new gauge in which each old size-$10n$ element divided by γ has size one if $j \neq 0$. Assume also that every pivot $I_i(g,x)$ for $i \ge j$ contains a unit with inverse of new size one. For $i < j$ use the vector ue_i^*: its size is at most two and its difference under $T = vu^{-1}$, divided by γ, has size at most two by the head-entry bound. For $i \ge j$ choose an eligible vector z with unit pivot c_i, and take $c_i^{-1}uz$. Its size is at most four since $a(g^{-1}z)$ has size three. Its difference under T, divided by γ, has size at most six: expand

$$(uz - vz)/\gamma = (a-b)g^{-1}z/\gamma + b(g^{-1}z - y^{-1}z)/\gamma.$$

The two terms have sizes four and three respectively, so after their sum and multiplication by c_i^{-1} the depth is six. Our recovery lemma with $C = 6$ gives new size $10 + 20h$ for $(1-T^{-1})/\gamma$. Finally

$$(x-1)/\gamma = a^{-1}\big((b-a)/\gamma - ((1-T^{-1})/\gamma)b\big)$$

has size at most $99n$. This proves `hasMatSize_sub_one_of_floorValues`, the common mechanism behind both shell arguments.

29.7.1 The finite shells

Assume a standard finite level, R its product of local rings of integers, and Z an intermediate open compact group between $K(\gamma)$ and $K(1)$. Let W_Z be its finite Bessel vector and M_Z its mass ratio, normalized as in Chapter 24. Use a depth-α gauge, with $\bar{\alpha} = \alpha \in R_{\mathrm{frac}}^{\times}$ integral (the fraction-field unit condition is not the integral unit condition), $B_i = \alpha^{-i}R$, $\Lambda = 2/\operatorname{md}(\alpha)$.

We require $\gamma^{-1} \in B_1$ if $j \neq 0$. The estimates of Chapter 26 give, for every nonzero elementary pair, close box representatives and a pivot generator $d_i \in R \cap E_{\mathrm{frac}}^{\times}$ in I_i, with $\mathrm{vol}(I_i) \leq \mathrm{md}(d_i)$. They also give the bound

$$\sqrt{\mathscr{N}(K)\mathscr{N}(K')}_{\mathrm{untwisted}} \prod_{a<h} M_a(\gamma) \leq M_Z. \tag{F.17}$$

This last inequality is the finite orthogonality/mass normalization, not an additional frequency estimate. For this basic finite gauge the floor mass is at most 2, since the quadric mass is at most 2 and the integral box has volume one; a larger Λ^{9999n} is convenient. Write $\mathfrak{a}_x$ for the integral denominator ideal of all entries of $(x-1)/\gamma$: c generates it exactly when c times every entry is integral and every scalar with that property is a multiple of c. Its modulus $d = \mathrm{md}(c)$ belongs to $(0, 1]$; this partitions the group into countably many measurable ideal shells. The form majorant is zero off the positions: a nonzero elementary pair has box representatives whose ratio belongs to the principal congruence group, so their relative position has the required unipotent–congruence factorization, exactly as in the finite bottom calculation of Chapter 26. Put

$$\sigma^{400n^2} = d\Lambda^{(10n+1)99n}.$$

If $\sigma < 1$, any nonzero elementary pair on this shell satisfying the two twisted nearness conditions must have some remaining pivot volume at most σ^2. Otherwise take the product $C_0 = \prod_{i \geq j} d_i$ and the new depth $\alpha^{10n+1}\bar{C}_0 C_0$. It absorbs γ^{-1} times the old depth-$10n$ elements by the assumed head-depth condition; each d_i^{-1} also belongs to the new size-one set, because C_0/d_i is integral. The preceding recovery lemma proves

$$\alpha^{(10n+1)99n}(\bar{C}_0 C_0)^{99n} \in \mathfrak{a}_x,$$

whence

$$d \geq \mathrm{md}(\alpha)^{(10n+1)99n} \prod_{i \geq j} \mathrm{md}(d_i)^{198n}.$$

If there is at least one remaining pivot and all volumes exceed σ^2, the right side exceeds $\Lambda^{-(10n+1)99n}\sigma^{396n(h-j)} \geq \Lambda^{-(10n+1)99n}\sigma^{400n^2} = d$. If there is no remaining pivot, the non-strict inequality already gives $d\Lambda^{(10n+1)99n} \geq 1$, a contradiction. This proves the cap even at the top floor.

The gauge of depth $\bar{c}c$ contains the shell in its size-one close-position set; its constant is $2d^{-2}$. Suppose $0 \leq f \leq \mathfrak{m}_j$ and, when $e < 1$, $f^2 \leq e^{2s}\Delta_j$ everywhere. Combine (F.14) with (F.12), (F.17), and $c_{\mathrm{place}} = 1$ to obtain the shell bound

$$M_Z \min(1, e, \sigma)^s \Lambda^{1500000n^3+1+9999n}(2d^{-2})^{50000n^2}.$$

If $t + 10^8 n^4 \leq k$, the level's rank bounds give $s - t \geq 400n^2(200000n^2)$: in a standard finite level the subtracted ranks total less than $40n$. Use

$$\min(1, e, \sigma)^s \leq \min(1, e)^t \min(1, \sigma)^{s-t}, \quad \min(1, \sigma)^{s-t} \leq (d\Lambda^{(10n+1)99n})^{200000n^2}.$$

The resulting modulus factor is $d^{100000n^2}$; the resulting power of Λ, including 2^{50000n^2}, is less than $9 \cdot 10^8 n^4 + 10^5 n^2$. The standard-level ideal zeta hypothesis bounds the sum of the modulus powers by 2 (their exponent exceeds the ideal zeta rank). Absorbing this last two, we have proved

$$\boxed{\kappa_j \int f \, d\mu \leq M_Z \min(1, e)^t \Lambda^{10^9 n^4}}. \tag{F.18}$$

This is `lintegral_shells_le`. In particular the unrestricted form majorant is integrable, with $e = 1$, whenever the stated gauge exists.

To choose its depth, if the form majorant is nonzero take a nonzero elementary pair from its upper integrals. The minor estimate in the basic integral gauge shows that all prefix products $D_l(p), D_l(q)$ are integral; otherwise the form vanishes. Put $\beta_p = \prod_{1\le l\le j} D_l(p)$, and similarly β_q. Then $\beta_p/D_l(p)$ and β_p/p_a are integral, the latter because $\beta_p/D_{a+1}(p)$ and $D_a(p)$ are integral. Choose $\alpha = \gamma\,\overline{\beta_p\beta_q}\,\beta_p\beta_q$. Both heads, the inverse γ^{-1} and every difference $p_a^{-1} - q_a^{-1}$ (where applicable) have the required size one. For $4E \le K_0$,

$$\Lambda^E \le (2/\operatorname{md}\gamma)^{K_0} H_{K_0}(p) H_{K_0}(q), \quad H_{K_0}(p) = \prod_{l\le j,\ l\ge 1} \operatorname{md}(D_l(p))^{-K_0/2}. \tag{F.19}$$

This follows from $\Lambda = (2/\operatorname{md}\gamma)(\operatorname{md}(\beta_p)\operatorname{md}(\beta_q))^{-2}$. The head weight is set to zero if any D_l is nonintegral, in agreement with the vanishing case.

Finally let ξ be a unit frequency, set $\beta = \alpha^{200n^2}$, and consider either the character weight of (F.15) at frequency ξ, or the unequal-head weight of (F.16). In the character case choose an auxiliary unit η componentwise: use $\eta = \xi$ when ξ/β is nonintegral in that component, and $\eta = \beta$ otherwise. Then $(\eta - \xi)/\beta \in R$ and the two characters agree on B_{200n^2} by self-duality. Moreover $\xi/\eta \in R$ and $w = \beta/\eta \in R$, with $\xi w \in R$. In the unequal-head case take $\eta = \xi, w = \beta$: indeed $\xi = \gamma^{-1}(p_i^{-1} - q_i^{-1})$ lies in B_2, so $\xi\beta$ is integral. In either case

$$\min(1, \operatorname{md}\eta^{-1/2})^2 \le \operatorname{md}(w)\operatorname{md}(\beta)^{-1}, \qquad \operatorname{md}(\xi) \le \operatorname{md}(\eta).$$

For character weights the first follows even with equality when $\operatorname{md}\eta \ge 1$; in the unequal-head case its right side is one. Since $A \ge s$, the frequency estimates (F.15)–(F.16) permit $e = \operatorname{md}(\eta)^{-1/2}$ in (F.18). Use the latter with exponent $t + 12$, under $t + 10^{10}n^4 \le k$. Extract the factor $\min(1, e)^t \le \operatorname{md}(\xi)^{-t/2}$; the remaining factor is at most $(\operatorname{md}(w)\operatorname{md}(\beta)^{-1})^6 \le \operatorname{vol}(R \cap \xi^{-1}R)^6 \Lambda^{1200n^2}$. The total gauge exponent is at most $2 \cdot 10^9 n^4$, with four times this exponent still at most $10^{10}n^4$. By (F.19), for $K_0 \ge 10^{10}n^4$ we have proved the actual standard-level frequency bound

$$|\mathcal{F}_j(p, q; \omega)| \le M_Z \operatorname{md}(\xi)^{-t/2} \operatorname{vol}(R \cap \xi^{-1}R)^6 (2/\operatorname{md}\gamma)^{K_0} H_{K_0}(p) H_{K_0}(q). \tag{F.20}$$

The notation $\mathcal{F}_j$ includes κ_j. The factor involving the intersection is precisely the inverse sixth power of the denominator index, not an estimate only for integral frequencies. This closes the passage from the arbitrary-head local estimates to the finite-place shell theorem `IntegralPlace.IsStdLevel.enorm_floorForm_le`.

29.7.2 The complex shells and the band bound

Now $\gamma = 1$, $B_i = \{|z| < L^i\}$, $\Lambda = L^2$, $L \ge \rho \ge 10$. The level has $A = k - 1$, $s = k - 3n - 2$, $c_{\mathrm{place}} = \pi^2$. The complex Haar normalization is $\mu = \lambda\,\mu_{\mathrm{Haar}}$, where μ_{Haar} is the Haar measure used in the Bessel norm and $\lambda > 0$ is the ratio. As in `IsCutBessel.formConst`, put

$$N_{\mathrm{form}} = \frac{\operatorname{vol} U(k)}{\operatorname{vol} U(r)} (\lambda|I|)^{-2}, \qquad C_{\mathrm{band}} = \frac{\operatorname{vol} U(k)}{\operatorname{vol} U(r)} \pi^{(2h+1)k}.$$

The cut satisfies $\int \|W\|^2 d\mu_{\mathrm{Bessel}} \le |I|$, and the second Bessel function has square norm $e_0^2|I|$ in that measure. Thus in (F.12) the untwisted Θ may be bounded by $e_0(\lambda|I|)^2$, by the pointwise inner-product inequality and Tonelli, using the quotient measure relation.

Fix heads for L, and a real parameter $t \ge 1$. Suppose $f \le \mathfrak{m}_j$ and, if $t > 1$, $f^2 \le ((\pi\rho^{24})^k/t^s)^2\Delta_j$. Set $T_i = tL^{11ni}$. The close-position balls of depth T_{i+1}^{99n} exhaust the positions; their differences give disjoint shells indexed $i \ge 0$. For $i \ge 1$, suppose a nonzero elementary pair with the two twisted nearness conditions has all remaining pivot volumes greater than

π/T_i^2. Then each remaining pivot set contains a point of modulus greater than $1/T_i$ (otherwise monotonicity of planar area gives the opposite volume bound). These points are nonzero. Use the gauge of size T_i in the matrix recovery lemma with the original size-L gauge for head-entry bounds; $L^{10n} < T_i$ and $\gamma = 1$. The close box representatives of the cut automatically satisfy the size-two difference condition. The lemma puts $(x-1)$ in the previous shell, a contradiction. Thus (F.14) applies with cap π/T_i^2. If this cap is above one, the same bound follows from the ordinary Cauchy–Schwarz case of (F.14).

For the basic gauge at ρ, the near-floor mass estimate after (F.5) is at most $\rho^{2574n}(\pi\rho^{22})^A$. Since $\pi^{s/2} \le \rho^k$, the square-root bound furnished by (F.14) on the ith shell is at most

$$(\pi\rho^{24})^k \rho^{9999n}/T_i^s \quad \text{times } \sqrt{\Delta_j}.$$

It also holds on shell zero by the hypothesis if $t > 1$ and by the ordinary mass bound if $t = 1$. Apply (F.12) with $\Lambda' = T_{i+1}^{198n}$ and sum. In particular $(\Lambda')^{50000n^2} = T_{i+1}^{9900000n^3}$, so the power of T_i on each shell is $9900000n^3 - s$. The source assumes $s > 10^7 n^3$, hence the geometric series has sum at most $2t^{9900000n^3-s}$. All extra factors in L, including $L^{108900000n^4}$ from T_{i+1}/T_i, $L^{3000000n^3+2}$ from (F.12), and ρ^{9999n}, are absorbed by $L^{10^9 n^4}$. The product of the $M_a(1)$ is one, and $(\pi^2)^{Ah}\pi^k \le \pi^{(2h+1)k}$. We obtain

$$\boxed{N_{\mathrm{form}}\kappa_j \int f\, d\mu \le C_{\mathrm{band}} e_0 L^{10^9 n^4} \rho^{24k} t^{9900000n^3-s}.} \tag{F.21}$$

This is `IsCutBessel.formConst_mul_lintegral_le`, under the source's top-floor and cut hypotheses.

For a nonzero form majorant, the minor estimate in the basic gauge at ρ shows all prefix products of p, q have absolute value at most $R_0 = \rho^{9n}$. If this fails the form is zero. Choose L to be the maximum of ρ and all inverse prefix absolute values. For $K_0 = n^6$, define $\ell_{K_0}(r) = r^{K_0}\min(1, r)^6$. Since each inverse prefix $r \ge R_0^{-1}$, $\max(1, r)^{K_0} \le R_0^{K_0+6}\ell_{K_0}(r)$. Consequently

$$L^{K_0} \le R_0^{(K_0+6)(2j+1)} H_{K_0}^{\infty}(p) H_{K_0}^{\infty}(q), \quad H_{K_0}^{\infty}(p) = \prod_{1 \le l \le j} \ell_{K_0}(|D_l(p)|^{-1}). \tag{F.22}$$

For a unit frequency y of (F.15) or (F.16) take $t = \max(1, \rho^{24}|y|)$. If $t > 1$, the frequency square-root estimate $(\pi/|y|)^A = (\pi\rho^{24}/t)^A$ is at most $(\pi\rho^{24})^k/t^s$, since $s \le A \le k$. Let $K_1 = k - n^7$. The enormous numerical size assumptions imply $10^9 n^4 \le n^6$ and $9900000n^3 + K_1 + 6 \le s$. Then

$$\rho^{24k} t^{9900000n^3-s} \le \rho^{24n^7}(\rho^{24}/t)^{K_1} t^{-6} \le \rho^{24n^7}|y|^{-K_1}\min(1, |y|^{-1})^6.$$

Combining these inequalities with the stated radius relation ρ to b and numerical bounds of `RealPlaceSizes` gives

$$\boxed{|N_{\mathrm{form}}\mathcal{F}_j(p, q; \omega)| \le C_{\mathrm{band}} b^{2000n^9} \ell_{k-n^7}(|y|^{-1}) H_{n^6}^{\infty}(p) H_{n^6}^{\infty}(q).} \tag{F.23}$$

For transparency, the only radius power before this substitution is $24n^7 + 9n(n^6 + 6)(2j + 1)$, bounded by $50n^8$ for the source's n; with the source's radius relation $\rho = b^{6n+2}$ (whose exponent is at most $40n$) it is at most $2000n^9$. This proves the complex Fourier band estimate `norm_floorForm_le`. In contrast with the finite estimate, the small-frequency upper bound is allowed to grow as $|y|^{-K_1}$; the damping factor is specifically $\min(1, |y|^{-1})^6$. Confusing these two asymptotics changes the sieve.

29.8 Descent and the actual sieve systems

It remains to check that the forms with the preceding estimates really are the consecutive forms of the sieve, rather than unrelated bounds. We use the following measure identity, which also takes care of integrability.

Suppose θ_b is the law of $v(c)^{-1}u$ under $\rho(c)\,dc\,d\zeta(u)$, with ζ supported on $Q_{b+1}\cap H_b$. Assume the unrestricted form majorant has finite group integral. In the double floor integral with character $\omega(g)=\psi(\xi\langle e_b,ge_b\rangle)$, replace τ by $\tau'\sigma$. This is legitimate in the quotient measure: the integrand as a function of τ is invariant under left multiplication by $Q_{b+1}\cap H_b$, and right translation of such a function preserves its integral. The ω weight becomes $\omega(\tau'^{-1})$; because the heads commute with H_b, the two kernel arguments now involve σg and $\tau'\sigma w_p^{-1}g$. Write $\tau'=v(c)^{-1}u$. The character is now $\omega(u^{-1}v(c))=\omega(v(c))$ since u fixes e_b. Transfer the variable u from the second kernel to the first by the simultaneous stabilizer invariance of the first kernel. The combined variable $u\sigma g$ has measure $\zeta*\theta_b*\mu_b$. The other variable x retains Haar measure. We have proved the identity of the following integrals:

$$\mathcal{F}_b(p,q;\omega)=\kappa_b\int\rho(c)\,\omega(v(c))\iint\frac{K_{b+1}(g,gx)}{\cdot\overline{K'_{b+1}((w_p^bv(c))^{-1}g,(w_p^bv(c))^{-1}gx)}d(\zeta*\theta_b*\mu_b)(g)d\mu(x)\,dc.} \tag{F.24}$$

Both the real and imaginary parts may be integrated in this identity. To check Fubini, make the preceding substitutions first in the integral of the absolute value of the integrand, which is bounded by the unrestricted majorant integral times κ_b. The substitutions are identities of measures for nonnegative functions, hence the right side is absolutely integrable as well. Multiplying by a fixed complex scalar is therefore also permitted. This is the general identity `integral_succ_floor_eq`. It requires the actual measure $\rho(c)\,dc$, rather than replacing the next floor by a formal sum.

For $b<h$ the flat-chart law is

$$v(c)=w_b(c),\qquad \rho(c)=\mathrm{md}(c)^{d_b},\qquad \zeta=\nu_b,\qquad \langle e_b,v(c)e_b\rangle=c.$$

Indeed the chart at a unit c uses the middle vector cv in its isotropic-vector coordinates and one skew coordinate; the ambient volume contributes its vector dilation and skew scale, and taking the scalar norm constraint removes precisely the extra skew scale. This is the lower-floor Weyl law of Section 24.4, valid also with Lebesgue measure on $\mathbb{C}$. The formula for κ_{b+1} with common next entries is $\kappa_b\,\mathrm{md}(c)^{d_b}$. The new Weyl ratio equals the old one, and $\zeta*\theta_b*\mu_b=\mu_{b+1}$ by the tower identities. Thus

$$\int\psi(\xi c)\,\mathcal{F}_{b+1}((p,c),(q,c);1)\,dc=\mathcal{F}_b(p,q;\psi(\xi Z_{bb})). \tag{F.25}$$

Nonunits form a null set, so the replacement of a nonunit by one in the unit-valued head family is harmless. Equation (F.25), with the preceding integrability proof, proves `integral_floorForm_succ` in either place.

29.8.1 Finite forms with a fixed-vector subgroup

Take now the actual data of a filling and a finite set of places as in `Filling.IsFillingLevel.exists_finite_sieveSystem`. In particular we assume that the local parameters are those of a standard level, the integral reduction is surjective as in `Filling.IsAdmissible`, the finite set is conjugation stable, and $1/\gamma$ is integral off it. Let E_S be the set of selected reduction places at which a vector is to be fixed. We describe all fields of the sieve system.

At a single selected conjugate pair of places, use the inverse image Z_1 of the nonzero-vector stabilizer in the residual general linear group (the action on the conjugate factor is the contragredient action). The reduction kernel contains $K(\gamma)$ and the inverse image lies in the integral group. Surjectivity and transitivity on nonzero vectors give relative index $q^r - 1$, where q is the residual cardinal, $r = k + 2h + 1$. Hence $\mu(Z_1)/\mu(K) = 1/(q^r - 1)$. The source's lower bound $q \geq b^{10}$ implies $q^r - 1 \geq 2b^{6k}$. The mass of the nonunits in a pair is at most $2b^{-10} \leq b^{-9}$ by the union bound. Apply Theorem 26.4 with $t = k - 10^{10}n^4$. With $\mu(K) \geq 2^{-n}$ the error in the ratio

$$\frac{B(Z_1)\mu(K)}{B(K)\mu(Z_1)} \leq \frac{\mu(Z_1) + 2^{-n}\,\mathrm{vol}(R \setminus R^{\times})^{t+1}}{\mu(K)}$$

is at most $b^{-9(t+1)} \leq (2b^{6k})^{-1}$. In this inequality B is the bottom quantity of that theorem, and its lower bound at K, namely $B(K) \geq \mu(K)^2\mu(K(\gamma))$, follows by restricting to the principal congruence group. Since the main term has the same upper bound $(2b^{6k})^{-1}$, each specified vector contributes the saving b^{-6k}.

Partition the set of places into the selected pairs and the remaining conjugation-stable subset. Choose Z to be the product of these inverse images, with integral group on the remainder. All globally fixing elements map into Z. On a product of local rings the phase factors and the character are products. The group Haar measure is a positive scalar times the product measure; if this scalar is a, then $\mu(Z) = a\prod \mu_i(Z_i)$ and $B(Z) = a^3 \prod B_i(Z_i)$, directly from the inner phase integral and outer Haar integral. Consequently the scalar cancels in the relative bottom comparison:

$$B(Z)\mu(K)b^{6k|E_S|} \leq B(K)\mu(Z). \tag{F.26}$$

This is the fixed-vector saving of `exists_intermediateGroup_lines`. The lower bound on each residual cardinal and the saving comparison use the actual rank k, not the dimension of a Heisenberg orbit.

For each compact open additive subgroup L define the consecutive forms

$$J_i(p,q) = \begin{cases} \mathcal{F}_i(p,q;1)/M_Z, & 0 \leq i \leq h, \\ \mathrm{topKernel}_L(p,q)/(\mathrm{vol}(L)M_Z), & i = h+1. \end{cases}$$

Their measure and normalization are exactly those of `FiniteSystem.lean`. For $i < h$ the step field is (F.25), divided by M_Z. At $i = h$ use the finite top Fourier identity of Section 24.5. In detail, if $R(\ell) = \langle z, \ell^{-1}z\rangle$ and $\Phi_{p,q}(\ell)$ is the inner product of the unnormalized correlations, then the Fourier integral of the last form at common top entry c equals

$$\frac{\mathcal{F}_h(p,q;\psi(\xi Z_{hh}))}{M_Z} \cdot \frac{1}{\mathrm{vol}(L)} \int_L \psi(-\xi v)\,dv. \tag{F.27}$$

This follows by the substitution $v = R(\ell) - c$ in the additive Haar integral. The final factor is one at zero and has modulus at most one at every frequency. The absolute integrability and joint continuity of the top kernel, and positive semidefiniteness on common last entries, were proved in Section 24.5: the top integral is a Gram matrix over the common band. In particular the normalization by $\mathrm{vol}(L)$ neither enlarges the Fourier bound nor damages positivity.

For all step frequency and unequal-entry fields apply (F.20) with $t = k - n^7$ and $K_0 = n^6$. The standing chain of inequalities $10^{10}n^4 \leq n^6$, $3n^6 + 12 \leq n^7 \leq k$ gives its numerical premises. Write f_∞ for the number of infinite places of the CM field. Since γ is the level product at the places in question, $\mathrm{md}(\gamma)^{-1} \leq g^{2f_\infty} \leq b^{46f_\infty}$, and $2 \leq b$; hence $(2/\,\mathrm{md}\,\gamma)^{n^6} \leq b^{n^9 f_\infty}$. The normalized bound is therefore

$$\Gamma_T(\xi)\,H_{n^6}(p)H_{n^6}(q), \qquad \Gamma_T(\xi) = \frac{\mathrm{genIndex}(\xi)^{(k-n^7)/2}}{\mathrm{denomIndex}(\xi)^6} b^{n^9 f_\infty}.$$

The frequency for unequal entries is precisely $\gamma^{-1}(p_i^{-1} - q_i^{-1})$ under the definition of the character of the finite filling. The character at zero is one. Finally, the bottom form at $i = 0$ is $B(Z)/(\mu(Z)\mu(K(\gamma)))$, by the exact pairing identity and the normalization of M_Z. Equations (F.26) and Theorem 26.4 imply

$$|J_0| \le b^{-6k|E_S|}(\mu(K) + 2^{-n}\epsilon^{t+1}) \le (1 + b^{-6})b^{-6k|E_S|},$$

using $\epsilon \le 1$, $\mu(K) \le 1 + 2^{-n}$ and $2^{-h} \le b^{-6}$. This gives exactly the bottom field of `exists_finite_sieveSystem`. All the analytic fields of that theorem now follow from the stated standard-level and filling data, and the actual normalized forms have been identified. The argument does not construct the global hyperbolic basis and local level data from arbitrary filling parameters; that construction is logically upstream.

29.8.2 Complex forms, the top vector, and the energy field

For the actual complex cut pair of the source, suppose the full function W and its cut F satisfy `IsCutBessel`, with Schur constant $|I|/\deg$, and $\deg C_{\mathrm{band}} = b^{4r^2}$. The lower forms are $J_i = N_{\mathrm{form}}\mathcal{F}_i(p, q; 1)$ for $i \le h$, and the top is the Gram form of the normalized Bessel correlation vector. Here is the normalization and descent at the last step.

At the last floor $V_h = X_0 \oplus \mathbb{C}z$, $h(z, z) = -1$, put $H = U(X_0)$ with probability Haar measure. For $|c| > 1$ choose the boost $v(c)$ acting on the plane of the chosen unit vector of X_0 and z, with diagonal entries $\bar{c}, -c$ and off-diagonal entries $\pm\sqrt{|c|^2 - 1}$ in the source's basis. It is an isometry and $\langle z, v(c)z\rangle = c$. The last quotient density is

$$\rho(c) = \mathrm{sphereDensity}_k(1)(|c|^2 - 1)_+^{k-1}.$$

One sees this directly by imposing $\|x_0\|^2 = |c|^2 - 1$ on the residual vector using polar coordinates in $\mathbb{C}^k$; H moves its normalized direction with invariant probability. Thus the representative measure is the law of $v(c)^{-1}a$ with $a \in H$. On $|c| \le 1$ the density is zero, with the arbitrary source extension of $v(c)$ immaterial. The convolution in (F.24) is now Haar measure μ in the first variable, and the second also has measure μ. Expanding tensor coordinates identifies its double integral with the inner product of the Bessel correlations

$$\mathcal{C}_p(v) = \int_G F(y) \otimes \overline{F((w_p^h v)^{-1} y)}\, dy$$

against ordinary Haar dy, times λ^2. This change uses $x \mapsto gx$ and preservation of Haar. The product integrals are absolutely integrable: if either overlapping cut pair is nonzero, both of its points lie in some common bounded two-frame size set, by the matrix recovery estimates; locally in the heads these sets are uniformly compact. One can alternatively obtain the integrability from the unrestricted form majorant as above.

Put $C_G = (\mathrm{vol}\, U(k)/\mathrm{vol}\, U(r))\,\mathrm{sphereDensity}_k(1)$. Since the complex measure modulus is $|p|^2$, the square-root factor in the normalized vector is

$$\mathcal{V}(p) = \frac{\sqrt{C_G}\,(|p_h|^2 - 1)_+^{(k-1)/2}}{|I|} \prod_{a<h} |p_a|^{d_a}\, \mathcal{C}_p(v(p_h)).$$

This is `sieveVector`. Its Gram product at common last entry c supplies all factors $N_{\mathrm{form}}\kappa_h\rho(c)\lambda^2$ in (F.24); thus the top step, including every character, is (F.25) with $J_{h+1} = \langle \mathcal{V}(p), \mathcal{V}(q)\rangle$. Continuity of the vector follows by dominated convergence, locally in the heads: the same two-frame estimate gives a single compact support for the integrand, which is continuous there. The boost has a continuous extension on the unit domain used in $\mathcal{V}$, and its

prefactor vanishes when $|p_h| \le 1$ (here $k > 1$). Positivity of the top Gram matrix is immediate. The isometry field in `HasBandLimits` allows any $v \in H_h$ with this same top scalar. Indeed for $|c| > 1$ its action on z has a residual vector of exactly the same length as $v(c)z$. The transitive action of H on that sphere makes $a^{-1}vz = v(c)z$ for some $a \in H$, and therefore $v = av(c)a'$ with $a' \in H$ fixing z. The Bessel law on H is a unitary representation law; substituting by a in the Haar integral and using that law for a' transforms the correlation tensors by a common unitary operator. Their inner products are unchanged. For $|c| \le 1$ both sides vanish by the prefactor. This proves the isometry field.

The Fourier step and unequal-entry estimates are (F.23), with $e_0 = 1$. It remains to prove the bottom estimate, which is strictly stronger than an unrestricted form-majorant bound. Let $a(x) = \int \langle F(g), F(gx)\rangle d\vartheta_0(g)$, $d(x) = \int \langle F(g), W(gx)\rangle d\vartheta_0(g)$. Pointwise

$$|a|^2 \le |d|^2 + 2|a|\,|a - d|.$$

The last product is bounded by the form majorant with second function $F - W$ at identity heads and floor zero, by the triangle inequality for the upper averages. Its normalized group integral is bounded by (F.21) with $t = 1$, $L = \rho$, $e_0 = b^{-5r^2}$, since this is the square-root L^2 difference bound of the cut pair. With the stated degree relation, twice this bound is at most $b^{-10}/\deg$: indeed it is

$$\frac{2}{\deg}\, b^{-r^2+(6n+2)(10^9 n^4+24k)} \le b^{-10}/\deg\,.$$

The last inequality follows from $n \ge 10^8$, $k \ge n^7$, $r \ge k$ and $b \ge 2$.

To bound d, the product in its integral is invariant under the Bessel unipotent, so the measure may be written λ times the Bessel quotient measure. Take a finite-Haar-measure subset S of G and use $f = \mathbf{1}_S \overline{d}$ in Schur's bound for W (or its complex conjugate according to the inner-product convention). This is integrable because d is bounded by the absolute upper-average bound. Fubini for the integrable tests, the inner-product inequality, and $\|F\|_{L^2} \le \sqrt{|I|}$ give

$$\int_S |d(x)|^2 dx \le \lambda\sqrt{|I|}\sqrt{(|I|/\deg)\int_S |d(x)|^2 dx}.$$

If the left side is zero the bound holds; otherwise divide by its square root. Exhaustion by finite measure sets now gives $\int |d|^2 dx \le \lambda^2 |I|^2/\deg$. In the real measure normalization, $\lambda \operatorname{vol} U(k)/\operatorname{vol} U(r) = 1$, so $N_{\mathrm{form}} \int |d|^2 d\mu \le 1/\deg$. Combining with the preceding error gives

$$|J_0| \le (1 + b^{-10})/\deg\,.$$

Thus one obtains `hasBandLimits_of_isCutBessel`, including its energy field, not merely a bound for the oscillatory lower forms. This argument uses the exact real Haar ratio and the full-function Schur estimate; neither may be replaced by a bound on the cut function alone.

29.9 Scope of the closure

The calculations above supply the general inverse-weight integral, the two centered Heisenberg Plancherel applications (including unequal heads), the twisted integrated self-norm, crossed Cauchy–Schwarz, all three pointwise form bounds, matrix recovery, shell summation and character modification at finite places, complex shell and frequency summation, lower-floor descent, and both sieve constructions from their stated local data and cut/Schur hypotheses. The identity-head finite bottom and top are used with the proofs already in Chapters 26 and 24. These are the analytic and measure-theoretic implications missing in the previous finite and real local audits (B05, B13, B19 and the band-system portion of B14).

In particular these results do not prove the global construction `Filling.exists_isFillingLevel`, the surjectivity assumption of `Filling.IsAdmissible`, or the existence of a cut pair with the required tail, Schur and normalization identities if those have not been established separately. The complex top step uses the Haar ratio identity $\lambda \operatorname{vol} U(k)/\operatorname{vol} U(r) = 1$, whose source is `RealPlaceSizes.haarRatio_spec`; the proof of that normalization and the representation-theoretic cut hypotheses belong to the upstream real orbital and discrete-series chapters. Accordingly these estimates remove the arbitrary-head Fourier and sieve-system implications as independent proof obligations, but do not themselves certify all the geometric and global premises of those obligations.

Chapter 30

Weighted index counting and the crude energy mechanism

The estimates proved in Chapters 22 and 27 control the height of rational links and pairs, but a pair discriminant gain does not by itself bound the sum over terms. We now prove the intervening weighted counting argument, with its precise marked-lattice indices and with its actual uniform exponent. We also separate the finite algebraic part of the energy estimate from the local real-cell estimate that supplies its amplitude bound.

30.1 An inverse-index counting lemma

Write $\mathcal{O}_f = \prod_{w<\infty} \mathcal{O}_{E_w}$ for the integral finite adeles, and write $[a] = \prod_w |a_w|_w^{-1}$ for the finite generalized index of an idele a. If $a \in \mathcal{O}_f$, then $[\mathcal{O}_f^d : a\mathcal{O}_f^d] = [a]^d$. The indices here and below are indices of additive groups, and all lattices are $\mathcal{O}_f$-modules. For a matrix Y let $L(Y)$ denote the $\mathcal{O}_f$-span of its columns. For arbitrary submodules K, M, write $[M : M \cap K]$ with value zero if the index is infinite, as in the source. The weight of an index zero is defined to be zero.

Lemma 30.1 (Inverse-index coloring)**.** *Let a be an integral idele, $K \subseteq \mathbb{A}_f^d$ an $\mathcal{O}_f$-submodule containing $a\mathcal{O}_f^d$, and Y_p integral d-by-$(r+s)$ matrices indexed by a set B. Let m be a primitive integral $(r+s)$-by-r matrix, so its columns extend to an integral invertible basis. Suppose*

$$[L(Y_p m) : L(Y_p m) \cap K] \mid n_p \qquad (p \in B)$$

for nonnegative integers n_p. If every nonempty subset of B on which all matrices Y_p agree columnwise modulo K has cardinal at most x, then, for every integer $l \geq r$,

$$\sum_{p \in B} n_p^{-l} \leq [a]^{d(d+s)} x.$$

Proof. Complete m to a matrix $(m \,|\, m') \in \mathrm{GL}_{r+s}(\mathcal{O}_f)$. At each place, the column module $L(Y_p m)$ has at most d generators, by the elementary divisor theorem. Choose d such generators (padding with zero); their entries can be assembled at all finite places into integral adelic vectors. Color p by their residues modulo $a\mathcal{O}_f^d$, and by the residues of the s columns of $Y_p m'$ modulo the same submodule. There are at most $[a]^{d(d+s)}$ colors.

In any fixed color, the group $(K + L(Y_p m))/K$ is independent of p: it is generated by the common residues of the chosen d generators. Denote it by H and its order by q, which is finite since it is a subgroup of the image of $\mathcal{O}_f^d$ modulo K. Thus $q \mid n_p$ on this color. Color

again by the residues of the r columns of $Y_p m$ modulo K; all these residues lie in H, so there are at most q^r second colors. Within any second color both blocks of $Y_p(m \mid m')$ agree modulo K. Right multiplication by the integral inverse shows the required congruence of Y_p. The sum of weights in this second color is at most $q^{-l}x$ (and terms with $n_p = 0$ contribute zero). Summing over the second colors gives at most $q^{r-l}x \le x$. Summing over the first colors proves the lemma. □

This is precisely the abstract `tsum_inv_index_pow_le_of_congr` of `New/RealPlace/VolumeKernel.lean`. The proof is insensitive to nonprincipal rational lattices: all choices of generators are local integral choices in the finite adeles. The two-stage coloring is essential, since the order of the image group may vary with the first color but its inverse power pays for the second color.

30.2 The labeled near terms and their middle congruences

Continue with the notation of Chapter 27. Thus h is the hyperbolic rank, $n = h + 1$, $\dim X_0 = k$, $F = (e_0, \dots, e_{h-1}, z)$, and J_N is the rational Hermitian matrix in the order (X, M, X^*), $X = \langle e_0, \dots, e_{h-1}\rangle$, $M = X_0 \oplus Ez$. Let q_0, q_1 be the two level primes, $g = q_0 q_1$, and let

$$\tau = b^{kf/4}, \qquad t = \min(j, h).$$

All of the standing hypotheses of `Filling.IsAdmissible` are used when specified, notably

$$b^{10} \le q_0, q_1 \le b^{11}, \quad g \le b^{23}, \quad k \ge 1000(2n)^9, \quad N \le b^{60n}, \quad b \ge 1000, \quad n \le b.$$

Write B_c for the integral class coordinate basis of Section 27.1, $C_c = B_c^{-1}|_{X_0}$ for the $(k+1)$-by-k matrix putting X_0 in the class middle coordinates. Its Gram with the marked middle form is $C_c^* J_c C_c = 1$; the integral J_c is the unimodular middle Gram in the lattice basis.

For $p = (c', \gamma)$ a term over (c, s, c') put $K_p = B_c^{-1}\gamma B_{c'}$. Its frame Gram equals the finite embedding of $Z(\gamma) = F^* J_N \gamma F$, so put

$$P(\gamma) = Z(\gamma)_{<h,*}.$$

The leading padded matrix $Y(\gamma)$ is the block diagonal matrix with the leading j-by-j block of $Z(\gamma)$ and identity on the other coordinates. By the level condition $D_j = \det Y \ne 0$. Let P_j be the first t rows of P. By the level flag relations, all h rows of P are in the rational span of P_j. For a rational subspace $T \subseteq X_0$, let m_T be an adelic basis of the saturated lattice

$$T(\mathbb{A}_f) \cap \{x : C_c x \in \mathcal{O}_f^{k+1}\}.$$

The primitive matrix $J_c C_c m_T$ and the integral matrix $Y_p = (\mathrm{middle}(K_p F))^*$ give the module

$$\mathcal{M}(p, T) = L(Y_p J_c C_c m_T).$$

The frame-value index of the source is

$$I_a(p, T) = [\, \mathcal{M}(p, T) : \mathcal{M}(p, T) \cap aL(P(\gamma)^*)\,]. \tag{30.1}$$

A labeled near term is a chosen representative of its double coset by the two frame-fixing congruence groups $\Gamma_c^0, \Gamma_{c'}^0$, is near and has level j, and its residue in the finite group $\mathfrak{G}$ has fixed second component under a choice of coordinates $\mathfrak{G} \simeq \mathfrak{G}/\bar{S} \times \bar{S}$. The following elementary

injectivity of transformed frame images will be used. If $p = (c', \gamma)$ and $p' = (c'', \gamma')$ have the same such label and

$$u\gamma F = u'\gamma' F, \qquad u, u' \in \Gamma_c^0, \tag{30.2}$$

then $p = p'$. Indeed $(u'\gamma')^{-1}u\gamma$ fixes the frame, hence is in the rational Bessel subgroup S_B. Its adelic integral term position lies in S_B, so the two residues differ by right multiplication by an element of $\bar{S}$. Equality of the fixed label, as well as the right coset, forces equality of these residues. The free-column-level argument of Section 22.10 now gives $c' = c''$ and the quotient belongs to $\Gamma_{c'}^0$. This makes the chosen double-coset representatives equal. This proves `IsLabelledTerm.eq_of_frameImage_eq` in `VolumeKernel.lean`.

We give explicit congruence elimination for fixed Z. For $b \geq j$, the vector $\gamma F Y^{-1} e_b$ lies in $F(E^n)$ by the shared flag property of the level. For $b = h$ and $j = n$ (so $Y = Z$), its pairings with F are e_h. Its dual isotropic coordinates therefore vanish and its z-coordinate is $-1/N$. The norm equation for $\gamma F Y^{-1} e_h$ gives squared X_0-norm at most $1/N \leq 1/5$ at each complex embedding, since the rational F-span has nonpositive Hermitian norm. These arguments prove the assertion of `HasLevel.middle_shared_vector_le`: whenever b is outside $\{0, \ldots, t-1\}$, the dual coordinates vanish and the X_0-part has squared norm at most $1/5$. In the shared-vector case it is zero.

Lemma 30.2 (Middle-row elimination)**.** *Suppose two level-j terms over the same left class have a common Z and the columns of $Y_p - Y_{p'}$ lie in*

$$K = gL(P^*) + gY^*\mathcal{O}_f^n.$$

Then some $u \in \Gamma_c^0$ makes the middle coordinates of $u\gamma F$ agree exactly with those of $\gamma' F$. More generally, in this assertion one may replace $L(P^*)$ by the span of any chosen subset of its rows containing the first t rows.

Proof. The middle z-coordinates agree because these are determined by Z. Let x be the difference of the rational X_0-components, a k-by-n matrix, and let Q be the array of the stipulated rows of P (in the first assertion $Q = P$). Taking adjoints gives

$$C_c x = gAQ + gDY \quad \text{with } A, D \text{ integral adelic matrices.}$$

Choose an invertible adelic a so that $C_c a$ is primitive and extend it to an integral basis. In particular there exists an integral left inverse L_0 of $C_c a$. Multiplying the last equation by aL_0 shows it has a solution $x = gl_f Q + gL_f Y$ with both $C_c l_f, C_c L_f$ integral. By density of rational matrices in finite adelic matrices, choose a rational matrix l such that $C_c l$ and $C_c L$ are integral, where

$$L = (x - glQ)(gY)^{-1}.$$

The two integrality conditions are open in l, and the preceding adelic solution belongs to their intersection.

For a column b outside the selected row indices (and for $b = h$), $QY^{-1}e_b = 0$: in the range $b \geq j$ the vector is shared, and the other omitted case is the last column at the top level. Thus $L_b = g^{-1}xY^{-1}e_b$. Its squared norm at each complex embedding is at most $4/(5g^2) < 1$ by the preceding middle-component estimate. But $C_c L_b$ is integral and its Gram is $L_b^* L_b$. If L_b were nonzero, this positive integral rational scalar would have absolute field norm at least one; its absolute values at every embedding are strictly less than one. Therefore $L_b = 0$. It follows that $LY = L_Q Q$ with L_Q the corresponding array of columns of L (for columns $\geq j$ the only possible such rows of L are zero by the same estimate, since $PY^{-1}e_b = 0$). We obtain $x = g(l + L_Q)Q$ with integral class coordinates for $l + L_Q$.

An elementary unitary block unipotent fixing each frame vector adds any such X_0-matrix times the dual isotropic coordinates. Here those coordinates are precisely P. To check integrality including primes above 2, use an integral trace-one element θ with $\theta + \bar{\theta} = 1$ (available for the stably unramified fields of the data). In coordinates (X, M, X^*), the translation with middle block A is

$$\begin{pmatrix} 1 & -A^*J_M & -\theta A^*J_M A \\ 0 & 1 & A \\ 0 & 0 & 1 \end{pmatrix}.$$

Its unitarity follows by multiplication against the block Gram. Take $A = g(l + L_Q)$ embedded in $X_0 \subset M$, with zero slots for unselected rows. It is congruent to 1 modulo g in the class basis since its off-diagonal blocks are divisible by g. It fixes F since $A^*J_M z = 0$. With the appropriate sign it cancels x. This is the claimed u. □

The more general row-subset assertion, including the rational approximation step, gives `HasLevel.exists_frameImage_toRows2_eq_add` in `VolumeKernel.lean`. To see explicitly the last vanishing used there, even if a selected row is numbered $b \geq j$, $QY^{-1}e_b = 0$ for all those b by the shared-vector identity. After the zero columns have been removed, the corresponding rows of Y equal those of P.

30.3 Counting the remaining top rows

A shared vector has a much stronger size estimate before passage to residues. Suppose γ is near, of level j, and write $Y = Y(\gamma)$. Let $b_0 \geq j$ be a column index. Then $\gamma F Y^{-1} e_{b_0} = Fw$ for an E-vector w. At every embedding v, let $y_v, \gamma_v^{-1} y_v$ be the two boxed points witnessing nearness, with the scale conventions of Section 27.1. Their normalized frame vectors are bounded by $R = b^{2n^2}$, and some $a_v = u_v y_v$, $u_v \in U(\mathbb{C})$, has operator norm at most b^{6n+2}. Extract the rows at the frame indices from $a_v y_v^{-1} F$ and $a_v y_v^{-1} \gamma_v F$, undoing the frame scales for rows and columns. Call their n-by-n matrices G_1, G_2. The first is unitriangular with determinant 1; the second has entries of absolute value at most

$$b^{23h+6n+2+2n^2} \leq b^{5n^2}.$$

Both bounds follow from the ratios of the frame scales, which are at most b^{23h}, also for the last frame vector. The same bound holds for the entries of G_1. The equations for w and $w' = \Phi_v(Y^{-1} e_{b_0})$ are

$$\begin{pmatrix} G_1 & -G_2 \\ 0 & \Phi_v(Y) \end{pmatrix} \begin{pmatrix} \Phi_v(w) \\ w' \end{pmatrix} = \begin{pmatrix} 0 \\ e_{b_0} \end{pmatrix}.$$

Every entry of Y is bounded by b^{5n^2}, by `IsNear.norm_frameProducts_le` or the box Gram bound. Cramer's rule and the permutation expansion yield

$$2|\Phi_v(\det Y \, w_a)| \leq 2(2n)! \, b^{10n^3} \leq b^{10n^3+4n+1} < q_0^{n^4}. \tag{30.3}$$

Here $n \leq b$ and $b \geq 1000$ were used in $(2n)! \leq b^{4n}$; the strict final inequality follows from $n \geq 10^5 + 1$ and $q_0 \geq b^{10}$. A left frame-fixing translation leaves the vector Fw unchanged. Thus this estimate is exactly the bound `HasLevel.norm_shared_vector_lt` (with room in its numerical inequality) under every allowed left translate.

Fix a subset of labeled near terms whose Z is constant and whose middle columns agree modulo K in Lemma 30.2, and choose a base element of that subset. For each element use the lemma to choose a left translate in Γ_c^0 aligning its middle rows with the base. All their

dual rows and their z-coordinates already agree, as these are entries of Z with the indicated scale. Write $P = Z_{<h,*}$ and put

$$d = q_0^{n^4}\, g\, \det Y\, \overline{\det Y}.$$

This is a nonzero integral rational scalar. Consider only those translated frame images whose top h rows agree modulo $d\mathcal{O}_f$. We show that there is at most one labeled term in each such color.

For two of these translates write ξ for the difference of the top h-by-n arrays. The top rows of the class inverse basis are the identity, by the marked coordinate construction. Consequently $\xi = dX$ with X an integral rational matrix. Both translated frame arrays have the same self Gram $F^* J_N F$ and their other rows agree, so

$$X^*P + P^*X = 0.$$

For a shared column $b_0 \geq j$, the vector in Section 30.3 is the same before and after each left translation. Estimate (30.3) bounds the difference of the top coordinates of $\gamma F \operatorname{adj}(Y) e_{b_0}$ for the two terms by strictly less than $q_0^{n^4}$. Therefore

$$\left|\Phi_v\left(g \det Y\, \overline{\det Y}\,(X \operatorname{adj} Y)_{a,b_0}\right)\right| < 1 \quad \text{for every } v.$$

This is an integral rational entry and hence zero by the product formula. It follows that $X \operatorname{adj}(Y)$ has zero columns for every $b_0 \geq j$.

Let α be the first h columns of $\operatorname{adj}(Y)$ and put $V = X\alpha$. From the skew relation we have

$$V^*P = -(P\alpha)^*X, \qquad V^*P\alpha = -(P\alpha)^*V.$$

The integral rational matrix

$$\Sigma = q_0^{n^4}\left(\overline{\det Y}\, V - \det Y\, V^* - (P\alpha)^*V\right)$$

is skew-Hermitian by the second equation. We claim $g\Sigma P = \xi$. If $j = n$, $Y = Z$, so $P\alpha = \det Y\, 1_h$; the first and third summands cancel and the first skew identity establishes the claim. If $j \leq h$, the zero trailing columns established above imply

$$VP = X \operatorname{adj}(Y) Y = \det Y\, X :$$

in the product only rows of Y of index $< j$ enter, and those rows equal the corresponding rows of P. Now the second and third summands in ΣP cancel by the first skew identity, giving the same result. The central unipotent with top-right block $g\Sigma$ fixes the frame and belongs to Γ_c^0 by integrality and its congruence modulo g. It translates one of the two frame images into the other. Now use (30.2); the two terms are identical. This establishes the required top-row injection, including the case $j = 0$ (then all the columns of $X \operatorname{adj}(Y)$ vanish).

30.4 The uniform weighted count over all labeled terms

Theorem 30.3 (Kept near terms with inverse-index weight)**.** *Let A be any subset of the labeled near terms at a fixed left class, level j and finite label. Let $T \subseteq X_0$ have codimension at most s, and suppose $n(p)$ is a nonnegative integer for each term such that $I_g(p,T) \mid n(p)$, with (30.1). Then*

$$\sum_{p \in A} n(p)^{-k} \leq b^{2000 n^6 (s+1) f}.$$

Proof. First color by the integral rational matrix Z itself. Every entry is bounded at all embeddings by b^{5n^2}. Distinct integral rational matrices are separated by at least one in the maximum norm of their complex entries: otherwise their nonzero integral difference would have norm smaller than one. A disjoint-ball argument in the product of fn^2 complex discs bounds the number of these matrices by

$$(1+2b^{5n^2})^{2fn^2} \le b^{12fn^4}.$$

For a fixed Z, put $d = q_0^{n^4} g \det(YY^*)$ as before. From $|\Phi_v(D_j)|^2 \le b^{11n^3}$ (the leading minor bound from the boxed Gram estimate and the determinant permutation expansion), we get

$$[d] \le b^{2f(11n^4+23+11n^3)} \le b^{46fn^4}.$$

In the matrix congruence use $K = gL(P^*) + gY^*\mathcal{O}_f^n$. It contains $d\mathcal{O}_f^n$: multiply a vector by the integral adjugate of Y^* and then by $q_0^{n^4} \det Y$. Also replacing $gL(P^*)$ by the larger K can only divide the index in (30.1). Apply Lemma 30.1 with row dimension n, primitive matrix $m = J_c C_c m_T$, column dimension $k+1$, $r = \dim T$ and complementary column count $l = k+1-r \le s+1$, and inverse-power exponent $k \ge r$.

In this lemma the cardinal bound on a subset congruent columnwise modulo K is exactly $[d]^{hn}$. Indeed align its middle rows by Lemma 30.2. The top h rows of the integral class frame images can be colored modulo d, one color per residue of each of their hn entries, and Section 30.3 establishes singleton fibers of this coloring. Consequently the weight of each original Z color is at most

$$[d]^{n(n+l)+hn} \le b^{46fn^5(2n+s)}.$$

Combining with the bound on the first coloring yields the stated bound, since

$$12n^4 + 46n^5(2n+s) \le 2000n^6(s+1).$$

All three colorings are on actual integral adelic arrays or on integral rational matrices. No count of all rational unitary elements with bounded real operator norm is needed. □

This proves `tsum_invPowWeight_le_of_codim_le` of `New/Filling/BlockBound.lean`. In particular its bound is in terms of the codimension of the chosen test space, not of the total dimension k. This feature allows one to sum it alongside the discriminant height.

30.5 The improved marked-class discriminant sum

Theorem 30.4 (Marked class averaging)**.** *Let $\mathcal{C}$ be the finite class set of the filling datum, and $d_c(T)$ the sublattice discriminant in the marked class lattice L_c of the rational subspace $T \subseteq X_0$. Under the standing hypotheses, for every codimension i,*

$$\sum_{c\in\mathcal{C}} \sum_{\operatorname{codim} T = i} d_c(T)^{-(k+1)} \le \frac{|\mathcal{C}|}{N^{kf}} \tau^{i+1}.$$

Proof. We use the elementary lattice description of the mark. Let $H = U(X_0)$ and K_f be the usual unmarked integral compact subgroup in $H(\mathbb{A}_f)$; let L be its subgroup preserving the marked lattice, namely the subgroup of integral unitary matrices whose k_0-th row agrees with that of the identity modulo $\bar{\pi}\mathcal{O}_f$. To verify this description, in the middle coordinate basis the distinguished positive vector is replaced by $x_{k_0}/\bar{\pi} + z/N$. The coordinate change on X_0 is integral on a vector exactly when that vector is integral and its distinguished coordinate belongs to $\bar{\pi}\mathcal{O}_f$. For an integral unitary matrix the marked condition on the

distinguished row entails the conjugate-column condition by taking the inverse adjoint. At each finite place at least one of $\bar{\pi}$ and π is a unit (since N is split). Thus on the overlap of the distinguished row and column no extra denominator occurs. This proves the subgroup description, including integrality of the conjugated full middle matrix and its inverse. It is `inclHFin_mem_latStabilizer` in `LatticeChart.lean`. The deeper level K_c is contained in L. If $A = K_f/L$ denotes the right coset set, its order satisfies

$$|A| \geq (N^k/b)^f$$

by the index hypothesis in `IsAdmissible`. Moreover $[\bar{\pi}] = N^f$ because $\bar{\pi}\pi = N$.

For any $u \in H(\mathbb{A}_f)$, consider the marked lattices obtained by uy with y ranging over representatives of A. Their sublattices inside the ordinary unimodular lattice $u\mathcal{O}_f^k$ are obtained by the single row condition

$$(y^{-1}u^{-1}x)_{k_0} \in \bar{\pi}\mathcal{O}_f.$$

Let T be fixed, with codimension i, and choose a primitive basis of the ordinary saturated lattice of T in $u\mathcal{O}_f^k$. In the u coordinates let its matrix be m. The image of m under $r_y = e_{k_0}^* y^{-1}$ has index I_y relative to the condition $\bar{\pi}\mathcal{O}_f$. The marked lattice in T is exactly the kernel of that row reduction, so its index in the ordinary lattice is I_y. Each lattice has the same rational span; by taking Gram determinants and local index (Section 22.13), its discriminant is therefore the ordinary $d_u(T)$ multiplied by I_y.

The rows r_y are integral. Two coincide modulo $\bar{\pi}\mathcal{O}_f$ precisely when their representatives belong to the same right coset of L. Indeed $r_y - r_z \in \bar{\pi}\mathcal{O}_f^{1\times k}$ implies $e_{k_0}^*(y^{-1}z - 1)$ has this divisibility by right multiplication by the integral z, and the subgroup description gives $y^{-1}z \in L$. Apply Lemma 30.1 with row dimension 1, col dimension k, subspace dimension $k-i$, complementary dimension i, $a = \bar{\pi}$ and exponent $k+1$. Each final fiber has cardinal at most one, so

$$\sum_{y\in A} I_y^{-(k+1)} \leq N^{f(i+1)}.$$

Sum first over y, then over the rational subspaces, using the ordinary sublattice zeta bound of Theorem 22.18. This gives

$$\sum_{y\in A}\sum_{\operatorname{codim} T=i} d_{uy}(T)^{-(k+1)} \leq N^{f(i+1)}(200fk\ \mathrm{rd}_E)^{f(k+1)i}.$$

Here the subscript uy indicates the marked lattice obtained in that way. Right averaging over K_f is exactly the sum over A divided by $|A|$, since the function is right-L-invariant and all those cosets have equal Haar measure. Hence this average is at most

$$N^{-kf}\ b^{f+60nf(i+1)+f(k+1)i/1000} \leq N^{-kf}\tau^{i+1}.$$

Use $N \leq b^{60n}$ and $(200fk\ \mathrm{rd}_E)^{1000} \leq b$ for the first inequality, and $1 + 60n(i+1) + (k+1)i/1000 \leq k(i+1)/4$ for the second. The latter follows with a large margin from $k \geq 1000(2n)^9$.

Finally, integrate the function of u over the rational adelic quotient $H(E^+)\backslash H(\mathbb{A}_f)$. It is right-$K_c$-invariant and the class stabilizers at level K_c are trivial, so its integral is $\mathrm{vol}(K_c)$ times the ordinary sum over class representatives (nonnegative integrands, or apply Tonelli first for truncated sums). Right Haar invariance leaves the integral unchanged by averaging over K_f. The last displayed uniform average bound gives the required inequality after division by $\mathrm{vol}(K_c)$. The mass of the whole quotient is $|\mathcal{C}|\,\mathrm{vol}(K_c)$, justifying precisely the factor $|\mathcal{C}|$. □

This is `sum_latticeDisc_inv_le` of `BlockBound.lean`; the extra factor N^{-kf} comes from averaging the mark, not from applying the ordinary zeta bound to each marked lattice separately.

30.6 The pair-weight estimate

We write $\bar{S}$ for the image of the Bessel subgroup in the finite residue group, and set

$$M_i = |\bar{S}|\,|\mathcal{C}|\,N^{-kf}(2\tau^2)^i, \qquad I_{\mathrm{sat}}(p) = [\,(X + K_pX)(\mathbb{A}_f) \cap \mathcal{O}_f^r : L((X \mid K_pX))\,],$$

where the last formula denotes the lattice spanned by the two isotropic basis arrays in integral coordinates, as in Section 27.4. For two congruent terms let $T = T^\circ$ be the projection of the common link with the two z-coordinates zero, and $i = \operatorname{codim} T$. The positive weight of the source for two kept near terms p, q over the same classes and reduction, at level j, is

$$W_j(p,q) = M_i^{-1} d_c(T)^{-(k+1)} (g^{2tf} I_1(p,T))^{-k} I_{\mathrm{sat}}(q)^{-k}, \tag{30.4}$$

where I_1 is the frame-value index (30.1) with $a = 1$, not the row index of Section 27.4. For pairs not of this form the weight is zero.

We establish the two divisibilities that allow Theorem 30.3 to apply. First,

$$I_g(p,T) \mid g^{2tf} I_1(p,T). \tag{30.5}$$

Indeed the rank of $L(P^*)$ is t, so its quotient modulo g has order g^{2tf}. The index after intersection with $gL(P^*)$ is the product of the old index and an integer dividing that order, by the injection of $(\mathcal{M} \cap L(P^*))/(\mathcal{M} \cap gL(P^*))$ into $L(P^*)/gL(P^*)$.

For the second divisibility, fix $p = (c', \gamma)$, let the varying term be $q = (c', \gamma')$ congruent to p and with the same common link $T^\circ = T$, and put

$$T' = T \cap \ker\big((\gamma F)^* J_N|_{X_0}\big).$$

Then $\operatorname{codim} T' \le i + n$. We claim

$$I_g(q,T') \mid I_{\mathrm{sat}}(q). \tag{30.6}$$

For a rational $x \in T'$ put $P_x = (0, x, 0) \in X_0$ as vector of the whole space. It is orthogonal to F and γF. Since T is the projected common link with both line coordinates zero, there is a vector $x' \in X_0$ such that (x, x') belongs to both middle links. The vector $\gamma^{-1}P_x$ belongs to $X^\perp$ and has zero z-coordinate (it is orthogonal to z). Its X_0-component equals x': subtract the link vector with this component from (x, x') inside the isotropic link. The difference has zero first component and second component in the positive space X_0. Its norm for the difference form is the negative of the positive norm; isotropy forces it to vanish.

Choose a rational lift $w' \in (X + \gamma' X)^\perp$ of the common link vector through the second link. Then $w' = (a', x, 0)$ and $\gamma'^{-1}w' = (b', x', 0)$ in the isotropic–middle–dual coordinates; similarly $\gamma^{-1}P_x = (a, x', 0)$. Thus

$$\gamma'\gamma^{-1}P_x - P_x = \gamma'(a - b', 0, 0) + (w' - P_x) \in X + \gamma' X.$$

This holds for all vectors in T' and hence after finite adelic base change. For any vector P_x with integral class coordinates, $g^{-1}B_c^{-1}(\gamma'\gamma^{-1}P_x - P_x)$ is integral because $K_qK_p^{-1} \equiv 1 \bmod g$. Therefore it lies in the saturated lattice defining $I_{\mathrm{sat}}(q)$.

Write $\widetilde{J}_c = B_c^* J_N B_c$ for the full class Gram (as distinct from the middle Gram J_c), and apply $g(K_qF)^*\widetilde{J}_c$ to this vector. As P_x is orthogonal to γF, its image equals, up to sign, the unscaled frame-value vector $(K_qF)^*\widetilde{J}_c B_c^{-1} P_x$. On the unsaturated generating lattice $(X \mid K_qX)\mathcal{O}_f$ the image of that same map lies in $gL(P(\gamma')^*)$: pairing with the second isotropic array K_qX gives zero, and pairing with the first gives exactly the conjugate rows of $P(\gamma')$. The quotient of the saturation by this generating lattice thus maps onto a group containing the image of the frame-value lattice of T' modulo $gL(P(\gamma')^*)$. Its order is divisible by that image order. This is (30.6); it proves divisibility of actual finite indices, even when the primitive bases of the subspaces have adelic denominators.

Theorem 30.5 (Pair weight bound)**.** *With exactly the weights* (30.4)*,*

$$\sum_{c,c'}\sum_{\gamma,\gamma'} W_j((c',\gamma),(c',\gamma')) \le 2\tau^2.$$

Proof. Overcount by the common codimension i, the left class c, the finite label in $\bar{S}$, and the rational space T of that codimension. The first term runs over the labeled terms at (c,j) (now allowing all right classes); the second must remain congruent over the same right class and have the indicated common link, so it is a subset of labeled terms of the same label. For a fixed first term, by (30.6) and Theorem 30.3 the sum of the second-term weights I_{sat}^{-k} is at most $b^{2000n^6(i+n+1)f}$. By (30.5) the sum of first-term weights is at most $b^{2000n^6(i+1)f}$. Their product is bounded by τ^{i+1}, since

$$2000n^6(2i+n+2) \le k(i+1)/4,$$

using $k \ge 1000(2n)^9$. The height sum of Theorem 30.4, now over c and T, costs at most $|\mathcal{C}|N^{-kf}\tau^{i+1}$. Including the finite label and the factor M_i^{-1} proves the bound for codimension i:

$$M_i^{-1}|\bar{S}|\,|\mathcal{C}|N^{-kf}\tau^{2(i+1)} = 2^{-i}\tau^2.$$

Finally sum this geometric series over $i \ge 0$. Nonnegative summands justify all reorganizations of the sums without requiring a priori finiteness. □

This proves `submoduleIndex_gIdele_smul` in the instance needed here by rank and index multiplicativity, and `sum_pairWeight_le` with the exact $2\tau^2$ constant. It is worth stressing that the index of a line in the marked lattice and the saturation index of an isotropic frame pair occur in different steps: charging their worst-case values without the weighted colorings would not produce this exponent.

30.7 The energy estimate once the local class bound is available

We give the exact quantitative deduction in `New/Filling/CrudeEnergy.lean`. This is a conditional analytic deduction, not an assertion that the local class bound has already been established. The notation is $\nu = \mathrm{unipCovolume}$, $B = \mathrm{besselEnergy}$, $D = \mathrm{mainEnergy}$, $M_m = \mathrm{countWeight}(m)$ as above, $\tau = b^{kf/4}$, and

$$\rho = \left(\frac{b}{g}\right)^{kf}(2\tau^2).$$

The admissibility conditions in use here are $\tau^{44}B \le D$, $\tau^{18}\rho D \le B$, and $0 < \rho \le 1$. For each level define

$$D_j = \begin{cases} D & j = n, \\ \tau^{13-32j}B & j \le h, \end{cases} \qquad a_j(G) = \tau^2\sqrt{\frac{D_j}{M_0 g^{2\min(j,h)kf}G^{k+1}}}.$$

Thus a_j is precisely `classSize`. The local class amplitude estimate that we require reads as follows, for a two-sided unipotent-stable set U of level j:

$$\|A_U^{\mathrm{cut}}(c,s,c')\| \le \nu \sum_{\substack{\gamma\in U\cap\mathcal{T}(c,s,c') \\ \gamma \text{ chosen near representative},\ s \text{ agrees}}} a_j(G_\gamma). \tag{30.7}$$

Here G_γ is exactly the gauge of Section 27.6. The proof below uses (30.7) also for each single link, not merely for the full set U. The link condition is two-sided stable by the algebra of the

link, so this is an instance of the same hypothesis. The thin- and thick-class proofs of (30.7) are given in Sections 30.8–30.10 below, under the actual local Bessel-system hypotheses.

First, the cut-error bound of `SchurTest.lean` already follows from the previously proved summable amplitude-tail theorem under its local majorant hypothesis. Indeed, the absolute sum of the errors contributing to any (c, s, c') is at most $\nu b^{-k^3 f}/2$ by Theorem 27.3. For the energy

$$\mathcal{E}(A;\Omega) = \nu^{-2} \sum_{c,c',s\in\Omega} \|A(c,s,c')\|^2,$$

it follows, even with the coarser estimate that charges every triple separately, that

$$\mathcal{E}(A_U^{\mathrm{full}} - A_U^{\mathrm{cut}};\Omega) \le \tfrac{1}{4}|\mathcal{C}|^2|\mathfrak{G}|b^{-2k^3 f}.$$

One can avoid the second class factor by first summing over c' before squaring, since the bound in Theorem 27.3 is already on the sum over c' for each fixed c, s. In particular the right hand side may be replaced by $\frac{1}{4}|\mathcal{C}||\mathfrak{G}|b^{-2k^3 f}$. Now put $r = k + 2h + 1$. The finite group has order at most g^{fr^2}, by the elementary matrix count at its split level places. Since $h \le k$ and k satisfies the standing lower size bound, $23(3k+1)^2 + 30/f \le 2k^3$. Use $g \le b^{23}$ to conclude

$$\mathcal{E}(A_U^{\mathrm{full}} - A_U^{\mathrm{cut}};\Omega) \le b^{-30}B. \tag{30.8}$$

We used $B = |\bar{S}||\mathcal{C}|\#\mathcal{I} \ge |\mathcal{C}|$, where $\mathcal{I}$ is the finite product Bessel index set. This is `energy_sub_cut_le`, with the exact b^{-30} factor.

Theorem 30.6 (Single-level crude energy)**.** *Assume (30.7), and let U be stable of level j, with every $s \in \Omega$ fixing norm at least $X \ge 1$. Let $C, w \ge 0$ satisfy*

$$B \le C, \qquad \tau^{20}B \le C \quad \text{if a level-zero term occurs over } \Omega,$$

and suppose moreover that $\tau^{12}D/X^k \le C$ if $j = n$ and some link is omitted from a designated set R. If

$$\sum_{c,c',s\in\Omega,\Lambda\in R} \|A^{\mathrm{cut}}_{U\cap\{\mathrm{link}=\Lambda\}}(c,s,c')\|^2 \le \nu^2 w,$$

then

$$\mathcal{E}(A_U^{\mathrm{cut}};\Omega) \le w + Cb^{-30}4^{-(n-j)}. \tag{30.9}$$

Proof. Split the sum of amplitudes into link blocks. Absolute convergence from Theorem 27.3 justifies this reorganization. Expand the squared norm, keep each diagonal term for a link in R with its actual value, and majorize each other ordered link-pair by the product of the two norms. Apply (30.7) to these products; the individual summands now correspond to pairs $p = (c', \gamma), q = (c', \gamma')$ that are congruent over (c, s, c'), are chosen near representatives, and have agreeing s. Each fixed pair determines at most one s, since the integral double-coset reduction of a term position is unique.

For such a pair, let i be the codimension of its common intersection in either link, and let T be the common positive link in Section 27.6. Put $m = k - \dim T$, so Theorem 27.11 gives $m \le i + (j-h)_+$. Also $i = 0$ implies equality of links; such a pair with $j = n$ is being majorized only when its link is omitted from R. Put $X' = X$ if $j = n, i = 0$, and $X' = 1$ otherwise. Write $I_p = I_1(p, T)$ and $J_p = I_{\mathrm{sat}}(p)$, and similarly for q. They are positive integers by the integral-lift proof of Section 27.4. The pair gain gives

$$G_pG_q \ge (g/2)^{2if}X'^2 d_c(T)^2 I_pJ_pI_qJ_q.$$

As $g/2 \ge 1$ and $X' \ge 1$, and the four indices are all at least one, this yields

$$(G_pG_q)^{(k+1)/2} \ge (g/b)^{ifk}X'^k d_c(T)^{k+1}(I_pJ_pI_qJ_q)^{k/2}.$$

In this last inequality the factor $g/2$ was weakened to g/b on the first k powers; the residual power and the extra index powers were discarded only in directions that enlarge the desired upper bound.

The geometric mean of the two ordered weights in (30.4) is

$$\sqrt{W(p,q)W(q,p)} = M_m^{-1} d_c(T)^{-(k+1)} g^{-2tkf} (I_p J_p I_q J_q)^{-k/2}, \qquad t = \min(j,h).$$

Consequently

$$\begin{aligned} a_j(G_p)a_j(G_q) &\le \frac{\tau^4 D_j (2\tau^2)^m (b/g)^{ifk}}{X'^k} \sqrt{W(p,q)W(q,p)} \\ &\le \frac{\tau^4 D_j (2\tau^2)^{(j-h)_+} \rho^i}{X'^k} \frac{W(p,q)+W(q,p)}{2}. \end{aligned}$$

The exact transformations of exponents in this inequality explain why a weighted count, rather than a finite bound on numbers of near terms with large frame-value index, was needed. The coefficient of the arithmetic mean is at most $C\tau^{-3}$ in every remaining case:

- If $j = n, i = 0$, use the omitted-link hypothesis to get $2\tau^6 D/X^k \le 2\tau^{-6} C \le \tau^{-3} C$.
- If $j = n, i \ge 1$, use $\rho^i \le \rho$ and $\tau^{18}\rho D \le B$: the coefficient is at most $2\tau^{-12} B \le \tau^{-3} C$.
- If $j = 0$, use the level-zero assumption: $\tau^{17} B \le \tau^{-3} C$.
- If $1 \le j \le h$, use $\rho^i \le 1$, giving $\tau^{17-32j} B \le \tau^{-15} B \le \tau^{-3} C$.

Sum the ordered pair weights by Theorem 30.5. The error in normalized energy is at most $2C/\tau$ (or $4C/\tau$ without the factor $1/2$ in the arithmetic mean). The size bound on k gives $\tau \ge 4b^{30}4^n$, hence $4C/\tau \le Cb^{-30}4^{-(n-j)}$. The kept diagonal contributes at most w. This proves (30.9). □

The vanishing of character-disagreeing stable sets and the flag propagation propositions are proved in Chapter 22; in particular Proposition 22.15 is the exact level alternative, including the nondegenerate top level. These are the first three lemmas of `CrudeEnergy.lean`. For a stable set on which all character-agreeing terms have level below a, induct on a to get

$$\mathcal{E}(A_U^{\mathrm{cut}}; \Omega) \le Cb^{-30}4^{-(n-a)}. \tag{30.10}$$

At $a = 0$ the amplitude vanishes by character cancellation. At the successor step, split off the stable level-a set and apply Theorem 30.6 with $R = \varnothing$ to it. The elementary bound

$$\mathcal{E}(A + A'; \Omega) \le (1+e)\mathcal{E}(A; \Omega) + (1+e^{-1})\mathcal{E}(A'; \Omega) \quad (e > 0)$$

follows from the triangle inequality and $2xy \le ex^2 + e^{-1}y^2$. For $e = 1$ combine the induction bound and the single-level bound to obtain four times $Cb^{-30}4^{-(n-a)}$, which equals the successor expression. Thus (30.10) is precisely `energy_le_of_level_lt`.

Corollary 30.7 (Full crude energy)**.** *Let the stable set U meet the single-level hypotheses on C and X, interpreting the omitted-link condition at the nondegenerate (top) level. Suppose its nondegenerate cut link energy over R is at most $\nu^2 w$. Under (30.7) and the hypotheses of (30.8),*

$$\mathcal{E}(A_U^{\mathrm{full}}; \Omega) \le (1 + b^{-4})^2 w + b^{-12} C.$$

Proof. Split the full amplitude into its cut part and the difference. Split the cut part into its nondegenerate and degenerate parts. Their energies are bounded respectively by $w + b^{-30}C$, $b^{-30}C$, and $b^{-30}B$, by (30.9), (30.10), and (30.8). Apply the elementary two-term energy inequality twice with $e = b^{-4}$. The resulting coefficient of w is $(1 + b^{-4})^2$; the remaining coefficient of C is at most

$$b^{-30}\left((1 + b^{-4})^2 + (1 + b^{-4})(1 + b^4) + (1 + b^4)\right) \le b^{-12},$$

where $b \ge 1000$ suffices for the last inequality. This proves the exact final energy estimate `energy_le_of_linkEnergy_le`. □

30.8 The real volume kernel used in the thin-class Schur estimate

The following argument supplies the precise determinant majorant of `exists_kernel_framVolume_le`, rather than an unspecified integrable bound. All matrices in this subsection are complex. Use the real-place model in which the ambient Hermitian matrix J is a Hermitian involution: its hyperbolic blocks exchange X and X^*, its restriction to X_0 is positive identity, and $Jz = -z$. Write F for the r-by-n frame matrix, X_0 also for its ambient column inclusion, and

$$H = F^*(1 - J), \qquad \phi(y) = |\det(Hy^{-1}F)|.$$

The columns of $(1 - J)F$ span the negative eigenspace, of dimension n. If $Hy^{-1}Fx = 0$, the vector $y^{-1}Fx$ lies in the positive eigenspace, whereas its J-norm is $x^*F^*JFx \le 0$. Hence it is zero, and $x = 0$. This proves $\phi > 0$. For a frame-fixing unipotent u, $(uy)^{-1}F = y^{-1}F$, so $\phi(uy) = \phi(y)$.

Theorem 30.8 (Integrated volume kernel)**.** *Fix y of Bessel size at most η, and an agreeing normal form j. Write $Z = F^*J\gamma F$ and let $i : \mathbb{C}^j \hookrightarrow \mathbb{C}^n$ select the initial j columns; $z_0 = i^*Zi$ is assumed invertible. Fix an inclusion $e : \mathbb{C}^s \hookrightarrow \mathbb{C}^j$ of distinct coordinate axes, an invertible $R \in M_s(\mathbb{C})$, and $V \in M_{k,j}(\mathbb{C})$. For all γ having this normal form and this same Z, whose initial X_0 coordinates satisfy*

$$X_0^*\gamma Fi = V + ARe^*z_0$$

for some $A \in M_{k,s}(\mathbb{C})$, there is one nonnegative measurable majorant K depending only on the fixed data such that

$$\left(\frac{\phi(y)}{\phi(\gamma^{-1}y)}\right)^{k+n} \le K(A),$$

and, for standard Lebesgue measure on the complex entries,

$$\int_{M_{k,s}(\mathbb{C})} K(A)\,dA \le \left(\frac{(2\eta^2)^j}{|\det z_0|}\right)^{k+n}\left(\frac{\pi^s}{s!\,(\eta^s|\det R|)^2}\right)^k. \tag{30.11}$$

Proof. The real Bessel-size property supplies a unipotent u such that $\|uy\|_{\mathrm{op}} \le \eta$ in the Bessel upper-triangular frame gauge (the precise property is `exists_norm_le_of_besselSize_le`). The inverse has the same operator norm because J is a unitary involution. Let U denote the unitriangular j-by-j restriction defined by $uFi = FiU$, and put $T = \eta U$. The determinants of T and e^*Te are η^j and η^s respectively. Put

$$I = Fi, \quad B = (Hy^{-1}F)^{-1}Hy^{-1}, \quad p = Ziz_0^{-1}, \quad t = \gamma Iz_0^{-1}, \quad C = Tp^*B.$$

These satisfy $BF = 1$, $i^*p = 1$, $F^*Jt = p$, and $CF = Tp^*$. The following determinant identity keeps the extra shared columns exactly, rather than estimating their contribution:

$$|\det(B\gamma F)| = |\det(Ct)|\,|\det z_0|\eta^{-j}. \tag{30.12}$$

To prove it, put $M = Btz_0 - i$. For $b < j$, take $X_b = Y_b$ equal to the bth standard basis column of $\mathbb{C}^n$; then $B\gamma FY_b = (1 + Mp^*)X_b$ since $p^*i = 1$. For $b \geq j$, normal form supplies $FX_b = \gamma FY_b$ with both coefficient arrays supported in indices at most b, and with leading coefficients of modulus one. As $j \leq h$ when such columns exist, $I^*JF = 0$; thus $p^*X_b = t^*JFX_b = t^*J\gamma FY_b = 0$. The same matrix equation $B\gamma FY_b = (1 + Mp^*)X_b$ follows. The two coefficient matrices are triangular with determinants of modulus one. Taking determinants, using Sylvester's identity $\det(1 + Mp^*) = \det(1 + p^*M)$, and using $1 + p^*M = p^*Btz_0 = T^{-1}Ctz_0$ gives (30.12).

Now put

$$Y = JtT^* - C^*, \qquad W = X_0^*Y, \qquad D = Cy, \qquad Q = \eta^{-1}(uy)^{-1}I.$$

We have $F^*Y = 0$ by the formulas for F^*Jt and CF. Thus Y is supported on $X_0 \oplus X^*$, so $Y^*JY = W^*W$. Also $QT = y^{-1}I$ and $DQ = 1$. The bound $\|Q\|_{\mathrm{op}} \leq 1$ implies $DD^* \succeq 1$. Since $By = (Hy^{-1}F)^{-1}H$ has rows in the negative eigenspace, $CJC^* = -DD^* \preceq -1$. Expanding Y^*JY now gives

$$-CtT^* - Tt^*C^* = W^*W - Tt^*JtT^* - CJC^* \succeq 1 + W^*W;$$

we used $t^*Jt = z_0^{-*}I^*JIz_0^{-1} \preceq 0$.

For completeness, if a square complex matrix L has positive Hermitian part $P = (L+L^*)/2$, then $P^{-1/2}LP^{-1/2} = 1 + iS$ for a Hermitian S. Spectral decomposition shows $|\det(1+iS)| = \prod_\ell \sqrt{1+s_\ell^2} \geq 1$, so $|\det L| \geq \det P$. Apply this determinant inequality to $L = -CtT^*$; monotonicity of positive definite determinants follows likewise by conjugation with the smaller positive matrix's inverse square root. We obtain

$$|\det(Ct)|\eta^j \geq 2^{-j}\det(1 + W^*W).$$

The original volume ratio is $|\det(B\gamma F)|^{-1}$, and therefore (30.12) gives

$$\frac{\phi(y)}{\phi(\gamma^{-1}y)} \leq \frac{(2\eta^2)^j}{|\det z_0|\det(1 + W^*W)}.$$

Choose $S = Re^*T^*e$ and $c = (Vz_0^{-1}T^* - X_0^*C^*)e$. The prescribed X_0 coordinate identity implies $We = AS + c$. Since $ee^* \preceq 1$, Sylvester's determinant identity and positive definite monotonicity give

$$\det(1 + W^*W) \geq \det(1 + (AS + c)^*(AS + c)).$$

All dependence on the possible γ has now vanished except for A. Use

$$K(A) = \left(\frac{(2\eta^2)^j}{|\det z_0|}\right)^{k+n} \det(1 + (AS + c)^*(AS + c))^{-k-n}.$$

This is continuous, hence measurable. The affine substitution $A \mapsto AS + c$ changes Lebesgue measure by $|\det S|^{2k} = \eta^{2sk}|\det R|^{2k}$.

We include the integral identity needed for this substitution. With $p_0 > k + s - 1$,

$$\int_{\mathbb{C}^{k\times s}} \det(1 + A^*A)^{-p_0}\,dA = \pi^{ks}\prod_{\ell=0}^{s-1}\frac{\Gamma(p_0 - k - \ell)}{\Gamma(p_0 - \ell)}.$$

Indeed write $A = (A' \,|\, a)$, use the matrix determinant lemma on $1 + AA^*$, and set $x = (1 + A'A'^*)^{-1/2}a$. The Jacobian on $\mathbb{C}^k$ is $\det(1 + A'A'^*)$, leaving the same integral on A' with exponent $p_0 - 1$. Integrating the last variable gives $\int_{\mathbb{C}^k}(1 + \|x\|^2)^{-p_0}\,dx = \pi^k\Gamma(p_0 - k)/\Gamma(p_0)$, by polar coordinates and the beta integral (equivalently by the elementary positive Laplace representation of $(1 + \|x\|^2)^{-p_0}$). Iterate from s down to zero, with the empty integral equal to one. At $p_0 = k + n$ and $s \le n$ the gamma product satisfies

$$\prod_{\ell=0}^{s-1}\frac{\Gamma(n-\ell)}{\Gamma(k+n-\ell)} = \left(\prod_{a=0}^{k-1}\prod_{\ell=0}^{s-1}(n-\ell+a)\right)^{-1} \le (s!)^{-k}.$$

Combined with the affine Jacobian, this proves (30.11), exactly the integral constant in `VolumeKernel.lean`. □

Thus the real determinant/Hua-integral part of the Schur argument is proved. The only real-place hypothesis used in this lemma beyond the explicitly given J and normal form is the actual Bessel-size property stated above. This proof does not infer the stronger global thin-class estimate merely from integrability of K: the coefficient lattice, the residual frame cutoff and the weighting of averages must also be accounted for.

30.9 The exact thin-class amplitude bound

Write $d_j = \mathrm{minorNorm}(j, \gamma) = |N_{E/\mathbb{Q}}(D_j(\gamma))|/N^{2(j-h)+f}$ in this subsection only (to distinguish the minor from the energy D_j of the preceding subsection). Put $t = \min(j, h)$, and let L_t and L_P be the integral row lattices generated by the first t rows of $P(\gamma)$ and all its h rows, respectively, after taking adjoints. Their index $I = [L_P : L_t]$ is the source's `xRowsIndex`. It is positive and finite since the level property makes the first t rows span the full row space. The double coset

$$\mathcal{D} = \Gamma_c^0\gamma\Gamma_{c'}^0$$

consists of terms over the same (c, s, c'), of the same level and with the same Z, as follows by multiplying the frame matrices (both outer actions fix the frame). Moreover their conjugate middle rows differ by elements of gL_P: the block formula for $u \in \Gamma_c^0$, used in Lemma 30.2, shows this when multiplying on the left, and the right action preserves the frame image itself. This is `terms_of_class_spec`. Let $m_B = \#\mathcal{I}$ denote the size of the product Bessel index set.

Define $\beta_c^b(y) = \beta_c(y)\mathbf{1}_{\mathrm{boxed}}(y)$ and $W(y) = \prod_v \|W_v^{\mathrm{cut}}(y_v)\|^2$. For any $d \in \mathcal{D}$ write

$$F_d(y) = \beta_{c'}^b(d_\infty^{-1}y)\prod_v\left(\frac{\phi(y_v)}{\phi(d_v^{-1}y_v)}\right)^{k+n}, \qquad S_d = \int \beta_c^b(y)W(y)F_d(y)\,dy.$$

Here S_d is `schurIntegral`, not an absolute amplitude. The sum $F(y) = \sum_{d\in\mathcal{D}} F_d(y)$ is Γ_c^0-left invariant, by reindexing d and using the frame-volume invariance. The averaging step takes the following form. If $s(x)$ is a measurable family in U_∞^0 parametrized by a set B_0 of measure at least one, and

$$\int_{B_0} F(us(x)y)\,dx \le H_0 \quad (u \in U_\infty^0,\ y \text{ boxed}),$$

then

$$\sum_{d\in\mathcal{D}} S_d \le \nu m_B H_0. \tag{30.13}$$

Indeed use the factorization $\beta_c = \beta_c^0\omega_c$ from (22.9); the second factor is U_∞^0-invariant, as are W and the boxed condition. The first factor periodizes to one under Γ_c^0. Disintegrate along

the left U_∞^0 orbits and use the quotient measure of $\Gamma_c^0\backslash U_\infty^0$, substituting $u \mapsto us(x)$ on this quotient. Its measure is right invariant (the unipotent group is unimodular). The integrals of the original and the translated F coincide. Tonelli then averages over $x \in B_0$ and bounds the integral by H_0 times $\int \beta_c W$. The latter is at most νm_B by the covolume formula (22.8), the local L^2 bound of the cut functions, and their unipotent covariance. Dropping the boxed indicator here only enlarges this bound. This proves (30.13), the precise averaging estimate `lintegral_boxCutoff_le_of_average`.

Theorem 30.9 (Thin-class Schur sum)**.** *With the same hypotheses as above,*

$$\sum_{d\in\mathcal{D}} S_d \le \nu m_B \tau^2 (b^4 g)^{-tkf} \left(\frac{I}{\sqrt{d_j}}\right)^{k+1}. \tag{30.14}$$

Proof. For each place choose the real scale $s_{v,a}$ of the frame as in Section 27.1. Write $e : \mathbb{C}^t \hookrightarrow \mathbb{C}^j$ for the first t coordinate inclusion, and put

$$R_v = \operatorname{diag}_{a<t}\left(\frac{\pi g}{b s_{v,a}}\right).$$

Let $x = (x_v)_v$, with $x_v \in M_{k,t}(\mathbb{C})$, and let B_0 be the product of open unit discs for all its complex entries. It has Lebesgue measure $\pi^{ktf} \ge 1$. Choose $s(x) \in U_\infty^0$ by the off-diagonal unipotent formula so that the X_0-matrix of $s(x)_v^{-1} d_v$ equals that of d_v plus $x_v R_v$ times the first t rows of the normalized Z_v. This identity is just the middle row of the unitary unipotent block multiplication from the proof of Lemma 30.2; it is `frameX0Part_unipotentOf_inv_mul`. It holds for all d and does not require integrality of $s(x)$. In particular this produces a continuous family of shears.

Color $d \in \mathcal{D}$ by its middle rows modulo gL_t. There are at most I^{k+1} colors: the rows already differ by gL_P, and there are $k+1$ rows in the middle. Fix a color and a representative d_0. By the rational middle-elimination argument of Lemma 30.2 (with the first t rows alone, and without the additional padded error term), the difference of the rational X_0 matrices of d and d_0 is $gl_d P_j$, with $C_c l_d$ integral. Multiplying on the left by u_v^{-1} for any fixed $u \in U_\infty^0$ adds the same X_0-matrix to both terms, as their dual coordinates agree. Thus still the difference is the archimedean normalization of $gl_d P_j$. Put $L_d(v) = (b/\pi)\Phi_v(l_d)$. The scale factors give, for $i : \mathbb{C}^j \hookrightarrow \mathbb{C}^n$,

$$X_0^*(u_v s(x)_v)^{-1} d_v F i = V_v + (x_v + L_d(v)) R_v e^* z_v, \quad z_v = i^* Z_v i, \quad V_v = X_0^* u_v^{-1}(d_0)_v F i.$$

To see the scales explicitly, the difference of the rational rows contributes the factor $g\Phi_v(l_{d,a})/s_{v,a}$ multiplying row a of Z_v, which is $(b/\pi)\Phi_v(l_{d,a})R_{v,a}$.

The normal form and Z_v are preserved by the left frame-fixing elements $u_v s(x)_v$. For any boxed y, apply Theorem 30.8 at each place with $\eta = b^{6n+2}$, V_v, z_v, and R_v. Denote its kernel by K_v. The volume factor in $F_d(us(x)y)$ is at most $\prod_v K_v(x_v + L_d(v))$. Translate x to $a = x + L_d$ in each term and use Tonelli. We must bound, at each a, the sum of the remaining cutoff factors

$$\sum_{d \text{ in this color}} \mathbf{1}_{B_0}(a - L_d)\beta_{c'}^b(d_\infty^{-1} us(a - L_d)y).$$

Any two summands with $a - L_d, a - L_{d'} \in B_0$ have $|\Phi_v((l_d - l_{d'})_{ab})| < 2\pi/b$ entrywise. Since $k \le b$, the square X_0 norm of each difference column is less than $k(2\pi/b)^2 < 1$ at every place. Its Gram is integral because $C_c(l_d - l_{d'})$ is integral, so the norm argument of Lemma 30.2 forces $l_d = l_{d'}$. Thus all nonzero summands have the same L and are evaluated at the same point $y_* = us(a - L)y$. Their frame images have identical X_0, dual and z coordinates; differences

are supported only in the top h rows. Therefore the differences of their boxed frame arrays $y_*^{-1}d_\infty F$ lie in a real subspace of dimension at most $2fhn$. Apply the weighted frame-count proposition 27.2, with the terms themselves as labels and cutoff weights as above. It bounds this sum by $b^{12fhn^3} \le \tau$. This argument is what prevents an unweighted count of translations in the noncompact frame-fixing group.

Consequently the average of the sum over one color is at most $\tau \prod_v \int K_v$, uniformly in u and boxed y. It remains to check constants, not just convergence. Put

$$P_t = \prod_{v,a<t} |s_{v,a}|^2, qquadu_0 = (j-h)_+ = j - t.$$

By the frame normalization and the definition of d_j,

$$\prod_v |\det z_v| = \sqrt{d_j}\, N^{u_0 f} P_t N^{-u_0 f} = \sqrt{d_j}\, P_t.$$

Also $P_t \ge N^{-tf}$ since $|\kappa_v| \ge 1$. The determinant of R_v in absolute value is $(\pi g/b)^t / \prod_{a<t} |s_{v,a}|$. The kernel integral bound (30.11), dropping only factors $\pi^{-t}/t! \le 1$, now gives

$$\prod_v \int K_v \le d_j^{-(k+n)/2} P_t^{-n} (2\eta^2)^{jfn} \left(2^j \eta^{2(j-t)} (b/g)^{2t}\right)^{kf}.$$

The rational minor is a nonzero integral element, so $d_j \ge N^{-2u_0 f}$. Hence the extraneous prefactor, after retaining $d_j^{-(k+1)/2}$, is bounded by

$$N^{(tn+u_0(n-1))f} (2\eta^2)^{jfn} \le b^{(72n^3+65n^2)f} \le \tau$$

using $N \le b^{60n}$ and $k \ge 1000(2n)^9$. Finally

$$2^j \eta^{2(j-t)} \le (g/b^6)^t.$$

For $j = t$ this is immediate from $g \ge b^{19}$. For $j = n$, $t = n-1$ and $\eta = b^{6n+2}$, the left side is at most $b^{12n+4+n/2}$ since $2 \le \sqrt{b}$; the right side is at least $b^{13(n-1)}$, and $12n + 4 + n/2 \le 13(n-1)$ because $n \ge 10^5 + 1$. Thus the retained bracket is at most $(b^4 g)^{-t}$. It follows that

$$\prod_v \int K_v \le \tau (b^4 g)^{-tkf} d_j^{-(k+1)/2}.$$

Together with the packing bound and the I^{k+1} colors this gives the required value of H_0 in (30.13), proving (30.14). □

The absolute-amplitude estimate now has a short exact deduction. Cauchy–Schwarz in the integral of the Bessel kernel gives

$$(\mathrm{amplitudeAbs}(c, c', d))^2 \le S_d S_{d^{-1}},$$

where the right second factor is for the opposite classes. In fact the frame-volume weights in the two factors cancel after the substitution $y \mapsto d_\infty^{-1} y$, and the boxed indicators do not change the integrand on the product kernel's support by the arithmetic cutoff construction. This is `amplitudeAbs_sq_le`. Applying the sequence Cauchy–Schwarz inequality and Theorem 30.9 to both inverse classes, using invariance of d_j on inversion and writing I_1, I_2 for the two row indices, gives

$$\sum_{d \in \mathcal{D}} \mathrm{amplitudeAbs}(c, c', d) \le \nu m_B \tau^2 (b^4 g)^{-tkf} (I_1 I_2 / d_j)^{(k+1)/2}. \tag{30.15}$$

Suppose now that termDiscNorm$(\gamma) \le N^f d_j$. Then $G = N^f d_j/(I_1 I_2)$. Comparing the square of the right side of (30.15) with $\nu^2 a_j(G)^2$ of the energy subsection, the ratio is exactly

$$\frac{M_0 m_B^2 N^{f(k+1)} b^{-8tkf}}{D_j} = \frac{B m_B N^f \tau^{-32t}}{D_j}.$$

By admissibility $m_B \le b^{2kf} = \tau^8$ and $N^f \le \tau$. For $j \le h$ the last ratio is at most $\tau^{9-32j}/\tau^{13-32j} \le 1$; for $j = n$ use $D_j = D \ge \tau^{44} B$ to bound it by $\tau^{9-32h-44} \le 1$. Hence the absolute-amplitude sum over this thin double coset is at most $\nu a_j(G)$, exactly `tsum_amplitudeAbs_class_le`. For a stable union of thin double cosets, only cosets containing a near term can contribute, since the boxed supports of the two cutoffs must meet whenever the absolute amplitude is nonzero. The chosen near representative of each such coset has the same gauge: the level, Z, and row indices are invariant under the two frame-fixing actions. Therefore summing this result proves (30.7) on the thin part (the disagreement of a reduced residue makes the stable amplitude zero by the covariance argument of Chapter 22).

30.10 The large-discriminant block, with its actual cell mass

For a nondegenerate near term let $x = \text{termDiscNorm}(\gamma) > 0$, hence $x \ge 1$ by integrality of the discriminant. Put $m_B = \#\mathcal{I}$, $d_B = \text{besselDegree}$, $C_G = \text{openCellConst}$, $L = 10^5 n^4$, and $P = \prod_{v,a<n} |s_{v,a}|^2$, $F_s = \text{frameScaleProd}$. The normalized real discriminants satisfy $\prod_v \mathfrak{g}(Z_v) = P^2 x$ by the change-of-basis formula. The exact cell mass of Theorem 25.19, at radius b^{2n^2}, therefore has product

$$\mathcal{M}(\gamma) = \frac{m_B^2}{d_B C_G^f}\, 2^{(k-1)f} b^{18n^3 fL} P^{-2(L+k-1)} x^{-(L+k-1)}. \tag{30.16}$$

We prove the numerical inequalities needed in `IsNear.le_classSize_and_cellMass_le`:

$$b^{-k^3 f} \le a_n(x), \qquad \mathcal{M}(\gamma) \le (\nu a_n(x)/2)^2. \tag{30.17}$$

From the near entry bound b^{5n^2}, the squared absolute values of all leading minors are at most b^{11n^3}, by expansion of the determinant and $n \le b$. Consequently $x \le \prod_v |\Phi_v(D_n)|^2 \le b^{11n^3 f}$ (the subtracted term in the nonnegative discriminant cannot increase it). Since $D \ge \tau^{44} B$ and $B \ge M_0$, we have $a_n(x)^2 \ge g^{-2hkf} x^{-(k+1)}$. This is at least $b^{-46hkf - 11n^3 f(k+1)} \ge b^{-2k^3 f}$ by the size bound on k.

For the second inequality use exactly the standing condition `packetConst_mul_mass_le`, namely

$$|\bar{S}||\mathcal{C}| g^{fh(2k+2h+1)} c_* \le (1 + b^{-9}) D d_B C_G^f \frac{(\nu F_s/m_B)^2}{\text{vol}(E)^n},$$

where $c_* = (\pi^k/((k-1)!N^k))^f |d_E|^{1/4} \text{vol}(E)^k$ and $\text{vol}(E) = (2/\text{rd}_E)^f$. Substitute (30.16) and $M_0 = |\bar{S}||\mathcal{C}| N^{-kf}$; after canceling the common factors a sufficient inequality is

$$4(1 + b^{-9}) 2^{(k-1)f} b^{18n^3 fL} P^{-2L} (F_s^2 P^{-2(k-1)}) x^{2-L} \le \tau^4 g^{fh(2h+1)} c_* N^{kf} \text{vol}(E)^n.$$

Here $x^{2-L} \le 1$, $P \ge N^{-nf}$, and by the definitions of F_s and P,

$$F_s^2 P^{-2(k-1)} = \prod_{v,a<n} |s_{v,a}|^{2(2+4(h-a))} \le b^{276n^3 f}$$

using $|\kappa_v| \le b^{23}$ and $N \ge 1$. The left side is thus bounded by $b^{f(1+k/2+138n^3 L+276n^3)}$, since $N \le b^{60n}$, $2 \le \sqrt{b}$ and $4(1 + b^{-9}) \le b$. The standing root-discriminant size condition gives $k, \text{rd}_E \le b^{1/1000}$, so $(k-1)! \le k^k$ and

$$c_* N^{kf} \text{vol}(E)^n \ge b^{-f(2k+n)/1000}.$$

As $g \geq 1$ and $\tau^4 = b^{kf}$, the desired comparison follows from

$$1 + k/2 + 138n^3(10^5 n^4) + 276n^3 + (2k+n)/1000 \leq k,$$

which holds for $k \geq 1000(2n)^9$ and $n \geq 10^5 + 1$. This proves both claims of (30.17).

Now let U be a stable set of nondegenerate terms over (c, s, c'). We choose weights adapted to open-cell integration. Partition into coarse double cosets $\Gamma_c \backslash U / \Gamma_{c'}^0$, choosing a representative d in each, chosen to be a near representative for the finer double cosets $\Gamma_c^0 \backslash U / \Gamma_{c'}^0$ if such a representative exists in the coarse coset. This is the choice in `enorm_amplitudeSum_le_classSize_top`. There is no multiplicity factor: the action of $\Gamma_c \times \Gamma_{c'}^0$ on a nondegenerate coarse double coset is free. Indeed, if $u\gamma = \gamma u'$ and $u' \in \Gamma_{c'}^0$, then u fixes γF. The dual hyperbolic rows of γF have rank h since its full frame Gram is invertible. The frame-fixing components of u are then forced to vanish by the triangular block formula, while its frame unitriangular part is forced to be the identity by invariance of the pair Gram with γF. Equivalently, write the latter identity as $A^* Z = Z$ where $uF = FA$: invertibility of Z implies $A = 1$, and the full-rank dual rows eliminate the remaining two upper off-diagonal blocks. Hence $u = 1$ and also $u' = 1$.

Replace the original product of cutoffs by the nonnegative weight that is $\omega_{c'}(d_\infty^{-1} y)$ at a chosen representative d and zero at every other term, where $\omega_{c'}$ is the quotient cutoff (22.9). To verify its orbit-sum identity, act by pairs in $\Gamma_c \times \Gamma_{c'}$. For each coset of $\Gamma_{c'}^0$ in $\Gamma_{c'}$, there is precisely one pair whose translated term is the chosen representative of its coarse coset, by the freeness just proved. The frame-fixing right factor does not affect $\omega_{c'}$, which is invariant under U_∞^0. The sum of the new weights thus equals the sum of $\omega_{c'}$ over $\Gamma_{c'}/\Gamma_{c'}^0$, which is one. The weight-change argument of Theorem 27.3 gives the same amplitude and bounds the total integrated difference between the full and cut kernels with these weights by $\nu b^{-k^3 f}/2$.

If a chosen representative contributes a nonzero cut integral then, at some point of the integrand's support, $d_\infty^{-1} y$ is boxed by the support of $\omega_{c'}$, and y satisfies the Bessel-size bound because the left cut kernel is nonzero. Some translate uy with $u \in \Gamma_c$ is boxed by the arithmetic frame-reduction assertion in Section 22.9. Then ud is near. Nearness is invariant under left and right frame-fixing unipotent translations, which leave both the frame box and Bessel-size bound unchanged. Thus the coarse coset contains a near finer representative, and the selection procedure ensured that the chosen d is such a representative.

For each such d, apply the open-cell theorem 25.19 at its normalized local components. Use the test function $\omega_{c'}$ in the product of the probability spaces of that theorem. It has absolute value at most one. The product L^2 contraction is the product of the individual contractions: verify it first on simple tensors, then on finite linear combinations by the bound for the tensor product of bounded Hilbert-space operators, and finally on the full L^2 space by density. Equivalently apply Fubini and the contraction successively to each variable, using the identity operator on the other Hilbert factors. The expectation of the norm squared of the resulting tensor integral is at most $\mathcal{M}(d)$. There exists a product translation t at which the norm squared is at most that number (if necessary take the infimum over the sample values; this yields the same bound by a limiting inequality). In the new integral put $y = d_\infty t z$, so the weight becomes $\omega_{c'}(z)$ by its $U_\infty^0 H_\infty$ invariance. This change of variables preserves Haar measure. The compact product region supplied by the local theorem contains the support of the cut integrand: the right frame box is unchanged by t, its scale bound follows from the cutoff, and the left Bessel-size bound follows from the nonzero left cut function. Restrict the full integral to this compact region. The integrated difference from the cut integral is at most the corresponding full-minus-cut absolute error with the new weights, bounded above by $\nu b^{-k^3 f}/2$. Therefore the norm of the cut term is bounded by

$$\sqrt{\mathcal{M}(d)} + \nu b^{-k^3 f}/2 \leq \nu a_n(x)$$

using (30.17). Summing the bounds over chosen near representatives, and allowing all near finer representatives on the right by nonnegativity, proves `enorm_amplitudeSum_le_classSize_top`.

For the whole stable set of level j, split into the thin part termDiscNorm $\le N^f d_j$ and its complement, both stable because the relevant leading minors and discriminant are invariant under the unipotent actions. The complement is necessarily nondegenerate, hence at level n; at this level the selected rows are all the rows of P, so both row indices are one and its class gauge is exactly $x =$ termDiscNorm. Apply the preceding open-cell estimate there and the thin-class estimate on the other part. If s does not agree, covariance under rational lifts of the reduced unipotent groups makes the whole stable amplitude zero (Chapter 22). This establishes (30.7), precisely `enorm_amplitudeSum_le_classSize`, for any filling satisfying the standing admissibility assumptions, the stipulated local Bessel-system estimates, and the already proved local open-cell estimate. The numerical energy deductions of Section 30.7 consequently require no independent global thin- or thick-class amplitude hypothesis.

30.11 Obligation ledger for this continuation

The distinction between a proof for a stipulated admissible filling datum and a proof that the required datum exists remains essential. The following statements are now supported by the written arguments above, with constants matching the source:

- **PROVED (algebra and counting).** `tsum_inv_index_pow_le_of_congr`, `HasLevel.middle_shared_vector_le`, `col_eq_zero_of_norm_lt_one`, `HasLevel.exists_frameImage_toRows2_eq_add`, and `IsLabelledTerm.eq_of_frameImage_eq` in `VolumeKernel.lean`, in the ambient fields and marked lattice of the problem. Lemma 30.1, Lemma 30.2, and Sections 30.2–30.3. The class-label injectivity is a statement about the actual quotient residue and free-column-level stabilizers.

- **PROVED (the exact weighted sums).** `HasLevel.norm_shared_vector_lt`, `tsum_invPowWeight_le_of_codim_le`, `sum_latticeDisc_inv_le`, and `sum_pairWeight_le` in `BlockBound.lean`, together with the index divisibilities used in the last lemma. Theorems 30.3, 30.4, 30.5. These close the weighted counting part of B11, conditional only on the stipulated filling arithmetic data, not on a separate weighted-count assertion.

- **PROVED (real determinant kernel).** `frameVolume_pos`, `frameVolume_mul_left`, `exists_kernel_frameVolume_le`, and `frameX0Part_unipotentOf_inv_mul`, with the exact Hua integral exponent and constant, Theorem 30.8.

- **PROVED as an implication for the stipulated real system.** `terms_of_class_spec`, `tsum_schurIntegral_le`, `tsum_amplitudeAbs_class_le` in `ThinClasses.lean`, `amplitudeAbs_sq_le` and `lintegral_boxCutoff_le_of_average` in `SchurTest.lean`, Theorem 30.9 and (30.15). This does not presume the global thin-class bound.

- **PROVED as an implication for the stipulated admissible filling.** `IsNear.le_classSize_and_cellMass_le`, `enorm_amplitudeSum_le_classSize_top` and `enorm_amplitudeSum_le_classSize`, Section 30.10, using the exact open-cell theorem of the previous continuation, the uniform summability/weight-change theorem from that continuation and its stipulated local majorant. These complete the global class-amplitude implication without a new independent packet estimate.

- **PROVED as an implication for the stipulated admissible filling.** `energy_sub_cut_le` and the entire crude energy deduction `energy_le_of_level_pairs`, `energy_le_`

`of_level_lt`, `energy_le_of_linkEnergy_le`, Section 30.7, with their exact constants. The character cancellation and flag-propagation lemmas from `CrudeEnergy.lean` were already proved in Chapter 22. Thus the crude-energy part of B11, B12 and B16 is no longer an independent hypothesis.

- **B04 remains OPEN as a datum-existence problem in this chapter.** The actual simultaneous arithmetic and local spectral choices, and the admissibility inequalities, must be verified by the construction of the filling; in particular the real Bessel system needs its actual quantitative majorant (the premise `IsCutPair.exists_majorant`) and the admissibility conditions given in `Setup.lean`.
- **B11, B12, B15 and B16 remain CONDITIONAL at their full statements.** The weighted counts, thin/thick class bounds and crude energies established here remove their corresponding independent global estimates. What remains in these obligations, absent additional chapters closing it, is the actual coherent packet comparison (including `sum_blocks_le_topKernel` and specialized zero-form/finite packet estimates), the required global-to-local quadric-limit input, the actual admissible system and the final sharp energy/span applied to those jointly realized inputs. Neither this chapter nor the arithmetic pair estimate alone proves the coherent packet upper bound (22.25).

Part IV

Lower bounds: periods, congruences and crystalline defects

Chapter 31

The determinant lower bound

Chapters 32 and 33 develop the complementary frames, finite square formulas, integral gluing, Rallis unfolding, comparison operators and the bounded Hecke annihilator arguments needed below. Chapters 11 and 34 give the normalized convergent mean and graph evaluation, exact arithmetic factors and local calculations. The genuine global period identities and their specialized nonvanishing remain explicit inputs when not proved.

We turn to the lower bound which is used in conjunction with the shortness and spanning theorems. In this part there is a second pair of mechanisms, both involving a comparison with theta lifts for an auxiliary finite-order character. The first comparison converts a nearly integral family of rows into a height estimate for the ratio of two full-dimensional determinants. The second compares the unknown determinant of a chain of lifts with determinants of single lifts, and then replaces each of these unknown determinants by a product of a special-value ratio and an integral comparison determinant. It is the last step which repays the cost of the ordinary (or idle) differentiation directions in the upper bound. The existence of adequate twists, a separate arithmetic theorem, will be considered subsequently.

We write $n = \dim_{\mathbb{C}} W > 0$, and retain $B = fa(\log b)^3$, $L = B \log b$, $H = mk\mathcal{U}$. Here the cost of one direction is the particular real number

$$\mathcal{U} = \sum_{\tau \in \Phi} (J(w_0(\tau)) - A(w_{\mathrm{nom}}(\tau), \tau)). \tag{31.1}$$

In particular it is not a positive-definite quadratic form or a norm. None of the inequalities below assumes its nonnegativity. Put $B_0 = B/(mk)$. The object of this section is the following precise estimate.

Theorem 31.1 (C)**.** *Let $y_1, \dots, y_q \in W$ be independent, where $q \geq (1 - 4b^{-2})n$. Suppose that* $\mathrm{IndexLe}(y, d_{\mathrm{ind}})$ *holds and that, in the coefficient-conjugate metrics of the setup,*

$$\mathrm{covol}_\sigma^2(\Theta_1 y_1, \dots, \Theta_1 y_q) \leq e^V \quad (\sigma \in \mathrm{Aut}\,\mathbb{C}).$$

Then

$$n(H - 3L) \leq V + 2d_{\mathrm{ind}}. \tag{31.2}$$

The statement to be reconstructed is `Data.integral_forms_long` in `New/Endgame/Long Forms.lean`. The precise estimates below designated *Formal-source input* are taken from named declarations of the compact development. We give ordinary mathematical proofs of the intervening algebraic and determinant implications, and isolate the statements of the more demanding automorphic and local computations. In particular the occurrence of a theorem from the formal source as an input is not an assertion that its independent verification or its translation from a published theorem has been completed.

31.1 Conventions for norms of conjugate coefficient vectors

Fix a level on a definite unitary group. In the coefficient model of algebraic modular forms, a form is a family of vectors in weight representations indexed by a finite adelic class set, with the appropriate covariance. For the Weyl coordinate representation used in the compact code, the pairing has a rational positive-definite matrix G (the matrix denoted `weylGramOver`). Write the class-sum pairing in schematic form as

$$\langle u, v\rangle = \sum_{h\in X} a_h\, \overline{u(h)}^{\,T} G v(h), \qquad a_h = |\operatorname{Aut}(h)|^{-1}.$$

Here X is a choice of class representatives at the specified level; different levels give different class sets and pairings. At the levels where the class stabilizers are trivial, $a_h = 1$. At other levels one clears the stabilizer orders by a uniform integer. The normalization concerning class sums is essential: it is not the usual probability measure on the finite class set.

For $\sigma \in \operatorname{Aut}(\mathbb{C})$ the norm used in the construction is

$$\|u\|_\sigma^2 = \sum_h a_h\, \overline{\sigma(u(h))}^{\,T} G \sigma(u(h)),$$

with σ applied entrywise. If u is a tuple of forms, the associated positive Gram matrix is formed from these conjugated vectors. It is in general incorrect to identify it with σ applied to the original Gram matrix. With κ denoting complex conjugation and $\sigma^* = \kappa\sigma\kappa$ as field automorphisms, one has instead

$$\sigma\langle u, v\rangle = \sum_h a_h\, \overline{\sigma^*(u(h))}^{\,T} G \sigma(v(h)). \tag{31.3}$$

This follows from the rationality of the coefficient pairing and from $\sigma(\bar z) = \overline{\sigma^*(z)}$. Thus

$$|\sigma\langle u, v\rangle| \le \|u\|_{\sigma^*}\|v\|_\sigma.$$

Much of the emphasis below on simultaneous estimates for *all* field conjugates is due to this distinction, not to the number of complex embeddings of a fixed coefficient field. These conventions match `LevelForm.normSqAt`, `LevelForm.pairingMatrix` and `LevelForm.covolumeSq` in `Lib/Adelic/CuspForms/LevelForms.lean`.

Lemma 31.2 (Mixed Gram determinant and completion)**.** *Let $v = (v_i)_{i=1}^q$ be a tuple in a finite coefficient representation as above. If its squared covolume is bounded by e^V at every conjugate, then*

$$|\sigma \det(\langle v_i, v_j\rangle)| \le e^V \quad (\sigma \in \operatorname{Aut}\mathbb{C}).$$

If $z_1, \dots, z_h$ satisfy $\|z_j\|_\sigma^2 \le e^A$ for every σ, the corresponding bound for the tuple (v, z) is e^{V+hA}.

Proof. Regard the scaled coefficient vectors as vectors in one ordinary finite-dimensional Hermitian space, so that the form displayed above is its Gram form. Equation (31.3) identifies the conjugate of the determinant with the scalar product of the exterior products of the σ^*-tuple and the σ-tuple. Cauchy–Schwarz in the exterior power bounds its modulus by the square root of the product of their squared covolumes, at most e^V. Equivalently, one uses

$$|\det(X^*Y)|^2 \le \det(X^*X)\det(Y^*Y)$$

for two rectangular matrices of equal column size. For the completion bound, if the conjugated v_i are dependent, the extended Gram determinant vanishes. Otherwise perform Gram–Schmidt

beginning with these q vectors. Each subsequent squared length is bounded by the original squared length of the added vector. Hence

$$\det \mathrm{Gram}_\sigma(v, z) \le \det \mathrm{Gram}_\sigma(v) \prod_{j=1}^{h} \|z_j\|_\sigma^2.$$

The source proves these assertions as `norm_map_det_gram_le` and `covolumeSq_elim_le` in the same file. □

We will also use the lattice property of the coefficient pairing. Let $\phi : \mathbb{C} \to \mathbb{C}_\ell$ be a field embedding and Λ_ϕ the lattice of forms of Weyl gauge at most one at ϕ. The coordinate lattices at the individual points can be calculated in integral adelic frames, rather than in an arbitrary fixed complex coordinate basis. Changing a rational frame is compatible with the Gram pairing. Consequently, for $u \in \Lambda_{\phi\circ\kappa}$, $v \in \Lambda_\phi$ the local pairing of the two coordinate vectors is integral. At the levels under consideration the class stabilizers are trivial and this implies

$$|\phi\langle u, v\rangle|_\ell \le 1. \tag{31.4}$$

Indeed after passing to integral frames the pairing becomes a sum of terms $\phi(\bar{u}_i)G_{ij}\phi(v_j)$ with integral entries; the integral class-sum property then follows from the ultrametric inequality. The source identifies these frames by `exists_mul_mem_integralAbove` and its Weyl gauge change law. We use their existence and compatibility as *Formal-source inputs*: `gramForm_mem_embValRing`, `norm_inner_le_one` (`Lib/Adelic/CuspForms/LevelForms.lean`), and `Data.classAut_eq_bot_of_le_chosenLevel` (`New/Data/ChosenPlaceLevels.lean`, used in `New/Endgame/LongForms.lean`). Notice that integrality at just one embedding for both arguments would not justify (31.4).

31.2 Heights with varying finite-place entries

In the direct application, different primes can require different twists. The height device allows the local entries of a family to vary, while preventing the arbitrary insertion of prime-dependent normalizing factors in the archimedean term. Let

$$I = \{\iota = (\ell, \iota_\ell) : \ell \text{ a rational prime, } \iota_\ell : \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell \text{ a field isomorphism}\}.$$

The ℓ-adic absolute value is normalized by $|\ell|_\ell = \ell^{-1}$. These isomorphisms are not topological. Their use as field isomorphisms follows from the axiom of choice and cardinalities of transcendence bases: both algebraically closed fields have cardinality and transcendence degree $2^{\aleph_0}$ over a countable algebraic closure of $\mathbb{Q}$. In particular any embedding of a fixed number field in either algebraically closed field extends to the requisite field isomorphism.

Definition 31.3. For a family $u = (u_\iota)_{\iota\in I}$ of complex numbers and $c \in \mathbb{R}$ write $\mathrm{Ht}_{\le c}(u)$ if there exist $Q \ne 0$ in $\mathbb{C}$, a real number α, and a real family h_ℓ indexed by primes such that

$$\begin{aligned} |\sigma Q| &\le e^{\alpha} && (\sigma \in \mathrm{Aut}\,\mathbb{C}),\\ |\iota_\ell(Qu_\iota)|_\ell &\le e^{h_\ell} && (\iota \in I),\\ \alpha + \sum_{\ell<M} h_\ell &\le c && (M \text{ sufficiently large}). \end{aligned}$$

The last sum runs over primes. We allow the h_ℓ to be signed; this is an eventual bound for prime cutoffs, not an assertion of absolute summability. This is `HeightLe` in `New/Heights/Heights.lean`.

Thus Q is a single nonzero number, uniform in ι. Even if a height bound is negative, it need not contradict a product formula when the entries u_ι differ. The construction requires precisely such a negative bound for the inverse volume ratios, to which the common determinant calculation will attach their reciprocal.

Lemma 31.4 (Height calculus)**.** *The following assertions hold.*

1. *If* $|\iota(u'_\iota)|_\ell \le |\iota(u_\iota)|_\ell$ *for all* ι*, then* $\mathrm{Ht}_{\le c}(u)$ *implies* $\mathrm{Ht}_{\le c}(u')$*; the constant* c *can be increased.*
2. *From* $\mathrm{Ht}_{\le c}(u)$ *and* $\mathrm{Ht}_{\le c'}(u')$ *follows* $\mathrm{Ht}_{\le c+c'}(uu')$*. In particular estimates add under finite products, and* $\mathrm{Ht}_{\le jc}(u^j)$ *for* $j \in \mathbb{N}$*.*
3. *If* $|\iota(u_\iota)|_\ell \le e^{h_\ell}$ *for every* ι *over* ℓ*, and* $\sum_{\ell\in F} h_\ell \le c$ *for all finite sets of primes* F*, then* $\mathrm{Ht}_{\le c}(u)$*.*
4. *If* $\mathrm{Ht}_{\le c}(1)$ *holds for the constant family, then* $c \ge 0$*.*

Proof. The first assertion uses the same witness. The second uses the product witness QQ' with archimedean bound $\alpha + \alpha'$ and finite-place bounds $h_\ell + h'_\ell$. The eventual prime cutoff works by taking the maximum of the two cutoffs. The third uses witness 1. For the last, the boundedness of all field conjugates of Q already implies that Q is algebraic: if it were transcendental, isomorphisms defined by extension of transcendence bases could send Q to a transcendental element of arbitrarily large absolute value. Put $F_0 = \mathbb{Q}(Q)$, $s = [F_0 : \mathbb{Q}]$, and $z = N_{F_0/\mathbb{Q}} Q \in \mathbb{Q}^\times$. Taking products over all embeddings of F_0 into $\mathbb{C}$ and into $\overline{\mathbb{Q}}_\ell$ respectively gives

$$\frac{1}{s}\log|z| \le \alpha, \qquad \frac{1}{s}\log|z|_\ell \le h_\ell.$$

Choose the eventual cutoff M large enough that $|z|_\ell = 1$ for every $\ell \ge M$. The rational product formula gives

$$0 = \frac{1}{s}\left(\log|z| + \sum_{\ell<M} \log|z|_\ell\right) \le \alpha + \sum_{\ell<M} h_\ell \le c.$$

The named source lemmas are `HeightLe.mono`, `HeightLe.mul`, `HeightLe.prod`, `HeightLe.pow`, `HeightLe.of_le` and `HeightLe.nonneg`. □

A useful consequence, repeatedly used below, is that if Q_ι is a positive integer prime to ℓ then inserting an arbitrary integral power of Q_ι into u_ι does not change its height bound: its image has local absolute value one. No new archimedean witness is required. In the application Q_ι is the norm of the twist conductor, not a fixed normalization scalar.

31.3 The determinant lemma with index

Let $T_\theta : W \to V_\theta$ be injective linear maps of finite-dimensional complex spaces with the prescribed metrics, for example Θ_θ. Define the squared volume

$$Z_\theta = \det(T_\theta^* T_\theta) > 0.$$

For a basis $v = (v_i)_{i=1}^n$ of W write $G_\theta(v) = \mathrm{Gram}(T_\theta v)$. A change of basis from an orthonormal basis to v shows

$$\det G_\theta(v) = Z_\theta \det \mathrm{Gram}(v), \qquad \frac{Z_\theta}{Z_1} = \frac{\det G_\theta(v)}{\det G_1(v)}. \tag{31.5}$$

Here it is important to use the same basis of the row space for every twist. The proof of Theorem C uses the following form of the determinant argument; its applicability does not require an integral basis of all row forms at a fixed finite place independent of the twist.

Lemma 31.5 (One determinant)**.** *Suppose v is a full basis and $|\sigma \det G_1(v)| \le e^{\alpha}$ for every σ. Fix a family θ_ι of twists and a family of nonzero scalars γ_ι such that $\mathrm{Ht}_{\le g_0}((\gamma_\iota\bar\gamma_\iota)^{-1})$. Let h be a nonnegative integer. Suppose the following holds for every ι over ℓ and every embedding $\phi : \mathbb{C} \to \mathbb{C}_\ell$: a matrix M_ϕ combines $T_{\theta_\iota} v$ into coefficient-integral rows at ϕ, with*

$$|\phi(\det M_\phi)|_\ell \ge e^{-e_\ell} |\phi(\gamma_\iota)|_\ell^h, \qquad \sum_{\ell \in F} e_\ell \le d_{\mathrm{ind}}$$

for every finite prime set F. Assume that the integral lattices pair as in (31.4). Then

$$\mathrm{Ht}_{\le \alpha + 2d_{\mathrm{ind}} + hg_0}\left(\frac{Z_{\theta_\iota}}{Z_1}\right). \tag{31.6}$$

Proof. Take $\phi = \iota_\ell$ and its composition with κ; the two matrices are not required to be the same. After applying ι_ℓ, the matrix of mixed pairings of the new rows is integral by hypothesis, and in the complex model it is

$$\overline{M_{\phi\circ\kappa}} G_{\theta_\iota}(v) M_\phi^T.$$

All its entries have image of absolute value at most one, hence so does its determinant by the Leibniz formula. Evaluating the determinant product and using the two determinant lower bounds gives

$$|\iota_\ell(\det G_{\theta_\iota}(v)(\gamma_\iota\bar\gamma_\iota)^h)|_\ell \le e^{2e_\ell}.$$

In view of (31.5), using $\det G_1(v) \ne 0$ as witness shows that $(Z_{\theta_\iota}/Z_1)(\gamma_\iota\bar\gamma_\iota)^h$ has height at most $\alpha + 2d_{\mathrm{ind}}$. Multiply by the hth power of the assumed inverse-square family, and apply Lemma 31.4. This is `one_determinant` in `New/Heights/OneDeterminant.lean`. □

The coefficient 2 in front of the index is therefore forced by the Hermitian determinant and is not a generous error allowance. The source treats the general situation in which the chosen lattices vary with ι, θ_ι, and ϕ. The quantifier over *all* embeddings ϕ here is intentional. In particular, an index bound at ι_ℓ only, in the absence of one at $\iota_\ell \circ \kappa$, is insufficient to reach (31.6).

31.4 Special-value data and fit twists

It will be helpful to give the exact analytic quantities to which the determinants are compared. Let M denote the set of chosen split places above $g_0 g_1$ (not the analytic cutoff on primes), and for a twist ϑ let $Q(\vartheta)$ denote its support, disjoint from M and its conjugate in the sense of the tame level datum; write $N(Q) = \prod_{w \in Q} Nw$. A *pair* $p = (\nu, z)$ of the proof consists of a tame unitary character ν of the norm-one torus of weight $-c_\tau$, unramified away from M, and an integer z with $1 \le z \le t$ and $z \le c_\tau \le 4b$. Tame part is denoted ψ. Define

$$\rho_p(\vartheta, s) = \frac{|L^{M\cup Q}(z, (\nu\vartheta^s)^{-1})|}{|L^M(z, \nu^{-1})|}, \qquad s \in \{-1, 1\}. \tag{31.7}$$

The removed local factors are to be understood in the source convention, including the conjugate places for the Hecke L-functions on the norm-one torus. The lower point p_v is the same pair with z replaced by v, for $1 \le v \le z$. For an integer $e \ge 0$ put

$$\Pi_S(\eta, e) = \left(\prod_{w \in S} Nw\right)^{e^2} \prod_{v=1}^{e} |L^S(v, \eta)|^2.$$

This is precisely `lValueProd` of `Lib/SiegelWeil/Tower/LargeSideConstantTerm.lean`; the ratio of two products at e satisfies

$$\frac{\Pi_{M\cup Q}((\nu\vartheta^s)^{-1},z)}{\Pi_M(\nu^{-1},z)} = N(Q)^{z^2}\prod_{v=1}^{z}\rho_{p_v}(\vartheta,s)^2. \tag{31.8}$$

Indeed each factor agrees and the first prefactor is a ratio of products of residue cardinalities over disjoint sets. The definitions are `Data.Pair.ratio`, `Pair.lowerPoint` and the identity `Pair.lValueProd_eq` in `New/Exchange/LValueRatio.lean` and `New/Exchange/TriangleLevelExponent.lean`.

The needed nonvanishing of these special values uses the weights of the pairs and their particular Hecke characters. We do not use GRH for them. As a *Formal-source input*, the declaration `ratio_ne_zero_and_real` of `New/Exchange/TriangleLevelExponent.lean` gives that every ratio here is nonzero and real; this follows in the code from `AutChar.HasWeight.L_ne_zero`. This property allows reciprocal families of ratios to enter the height calculus without exceptional zero entries.

The local projection condition. For an auxiliary pair $O=(W_j,V_j)$ with $\dim_E W_j=r$, let V_O denote its algebraic form space on the definite group. At each split place w outside $M\cup Q$ and its conjugate, there is a normalized Hecke polynomial operator $H_w(X)$ on V_O. The test point is

$$X_w=(a_O(w)\,Nw^{\mathrm{rankDiff}(O)-1})^{-1},$$

where $a_O(w)$ denotes the Frobenius character value in the normalization of `AuxPair.aFrobValue` for rank r. We use the precise operator in `AuxPair.heckePolyOnForms`. The definition of X_w is `AuxPair.testValue` in `New/Rows/StackedDeterminants.lean`. If e_0 denotes the chosen central projector and $Z\subset V_O$ the space of lifts, set

$$F_w=\operatorname{im}e_0\cap\ker H_w(X_w)\cap Z^{\perp}.$$

The points X_w are designed so that genuine lifts belong to the first two factors. A vector is *tested* if it is orthogonal to F_w at every such place; the false-lift subspaces depend on w. This is `AuxPair.IsTested` in `New/Selmer/FitTwists.lean`. The orthogonal complement $Z^{\perp}$ is computed with the ordinary complex pairing.

For a fixed isomorphism ι_ℓ, Z has denominator bound β when the following holds for $d_Z=\dim Z$: for every $u_1,\dots,u_{d_Z}$ integral at every ℓ-adic embedding, and every $v_1,\dots,v_{d_Z}$ integral at ι_ℓ and tested,

$$\bigl|\iota_\ell\det(\langle u_i,p_Zv_j\rangle)\bigr|_\ell\le e^{d_Zr\beta}. \tag{31.9}$$

Here p_Z is the complex orthogonal projection. This is the actual declaration `DenominatorLe` in the same file. The projection is only applied to the second family; equality with the pairing of both projections may be used at the ordinary complex metric. In this definition it would be substantially stronger to ask for an entrywise denominator bound for every integral vector, or to drop the word tested. Neither strengthening is assumed.

An m-fold twist θ has first m powers still nontrivial at every prime of its support, each power with the same level support. The family θ_ι is fit if:

1. $\ell\nmid N(Q(\theta_\iota))$ for ι over ℓ;
2. for every $j\in[1,m]$ there exist numbers β_ℓ (allowed to depend on j) such that $\sum_{\ell\in F}\beta_\ell\le B_0$ for all finite prime sets F, and the good single-lift subspace at twist θ_ι^j has denominator bound (31.9); the same β_ℓ bounds the rank-one lift lines of all pairs of rank-one type at this index. For the latter the rank in (31.9) is one;

3. for all j, both signs s, and all pairs p ramified exactly at M,

$$\mathrm{Ht}_{\le x^{999}}\left(\rho_p(\theta_\iota^j, s)^{-1}\right).$$

Here rank-one type means $m < z \le r$, tame part ω^j, and $w_j + r - t \le c \le w_j$ at all infinite places (with the rank index condition in `Data.Pair.IsRankOne`); note that the term rank-one refers to a lift *from* $U(1)$ and not the dimension of its algebraic coefficient representation. At a fixed j the same finite-place costs must handle all these auxiliary pairs. These conditions are precisely `Data.IsFitFamily` in `New/Selmer/FitTwists.lean`. The existence result `exists_isFitFamily` in `New/Selmer/FitTwistsExist.lean` is a *Formal-source input* here; its substantial arguments are developed separately later in the paper.

Definition 31.6 (Cost of an exchange factor)**.** For a complex scalar $c(\theta)$ depending on the m-fold twist, say it has cost s squares if for every fit family one has

$$\mathrm{Ht}_{\le 2stx^{999}}\left(|c(\theta_\iota)/c(1)|^{-2}\right).$$

This description is used below only for nonzero factors. It is `Data.CostsSquares` in `New/Exchange/CostOfSquares.lean`.

Lemma 31.7. *Costs add under multiplication (and may be increased). A factor whose squared modulus equals a fixed positive number times an integral power of $|\mathcal{O}_E/\mathfrak{m}(\theta)|$, for $\mathfrak{m} = M^a Q(\theta)$ with fixed a, costs zero squares. If a factor c satisfies*

$$|c(\theta)|^2 = C\Pi_{M\cup Q(\theta)}((\nu\theta^j)^{-1}, z),$$

with $C > 0$ independent of θ, and $p = (\nu, z)$ ramified exactly at M, then c costs one square.

Proof. The first assertion is Lemma 31.4. The cardinality of the residue ring is a fixed integer times $N(Q(\theta))$, so in the second assertion the squared modulus ratio is a local unit at the relevant ℓ; the constant height witness 1 suffices. For the last assertion apply (31.8) and the fit-family condition to each of the z inverse ratios, twice. The factor $N(Q)^{-z^2}$ is a local unit and does not alter the bound. Since $z \le t$, the resulting constant is at most $2tx^{999}$. These are the rules expressed in `CostsSquares.mul`, `CostsSquares.of_norm_sq_eq_mul_card_zpow` and `exists_isTameFor_costsSquares_one`. □

31.5 The correspondence of complementary rank and the finite square

There are two uses of an exchange identity: finding a spanning family of crude rows, and factoring the volume on the good row space. We give their common input in some detail. For Hermitian spaces V, V', A let $I_A^V(\xi, \Phi)$ be the theta integral over $[U(V)]$, with real datum ξ and finite datum given by Φ on the matrix residue ring. The source makes this precise as a mixed real–finite datum `modDatum`, with splitting characters on the two members having product the prescribed rank twist. The input character ν has weights μ_τ and tame finite character χ modulo a level ideal $\mathfrak{m}$. The normalized character on the large space of rank N is denoted $\mathrm{rankTwist}(\nu^{-1}, N)$; it includes a power $\lfloor N/2 \rfloor$ of the fixed splitting character. We do not suppress this normalization in comparing the special values.

Proposition 31.8 (Complementary-rank exchange; formal-source input)**.** *Let $N = \dim_E A = \dim_E V + \dim_E V'$, $\dim_E V = \dim_E V' + 2e$, $e \ge 0$. At the infinite places the negative signs of*

V, V' add up to those of A; write this number as q_A. Assume $2q_A + 2 \le N$ and $N \le 2\mu_\tau + 2q_A$ at every infinite place. Fix the index identification of the complementary blocks and a rapid real datum v. Then there are a rapid real datum x and a constant C independent of the varying modulus and character such that, for every admissible tame character $(\mathfrak{m}, \nu, \chi)$, there is $c \neq 0$ with

$$|c|^2 = C\Pi_{\mathfrak{m}}(\nu, e), \qquad I_A^V(x, \Phi) = cI_A^{V'}(v, \Psi)$$

whenever the finite data Φ, Ψ are partners. Here $\Pi_{\mathfrak{m}}$ means the product defined above with the places of $\mathfrak{m}$.

This is `exchange_theorem` in `New/Exchange/ExchangeTheorem.lean`. Its proof in the formal development reduces the nonzero constant at equal rank to the calculation `exists_equalRankThetaRatio_value` in `New/Exchange/EqualRankValue.lean`, and then applies `exchange_of_partners` in the same file. Along a hyperbolic plane the surviving constant term of the lifted character has a scalar given by local Gauss sums and a Tate integral; the size of the total scalar under repeated reduction is precisely the product Π. Its factors have the conductor exponent e^2 since adjoining successive isotropic lines adds exponents $1, 3, \dots, 2e-1$. The actual matching of these integrals and nonconstant terms is not established by this counting identity. The tower and constant-term results in `Lib/SiegelWeil/Tower/LargeSideConstantTerm.lean`, `New/Exchange/ExchangeDifference.lean` and `New/Exchange/EqualRankValue.lean` are essential background to the displayed theorem. We take the complementary-rank theorem with its stated hypotheses as an input, rather than simply appeal to a general Siegel–Weil identity without verifying normalization or ramified data.

The passage from such an equality of integrals to an exchange of operators is important. With two exterior Hermitian spaces U_1, U_2, combine them to play the role of A. The two intermediate spaces have ranks differing by $2e$. Suppose the characters on all four edges are matched so that their products are the rank twist on $U_1 \oplus U_2$. Opposite choices of signs are used in the descent edges. Pairing the composition of a lift and a descent with a functional on the exterior target gives an integral on the product of the exterior groups, with a theta integral over the intermediate group. This is the situation of Proposition 31.8. Inserting the complementary-rank equality changes the order to descent then lift through the other intermediate group.

The source takes special care about integrability in this argument, because some of the involved groups are indefinite. Its lemma `exists_integrable_kernel_bound` bounds, for rapid vectors on the exterior groups, the integral of their products against a theta kernel by a constant times a weighted norm of the real datum. The permitted real datum on the other side is obtained as a limit in that norm and approximated by finite sums of elementary tensors. The closedness of a one-dimensional span in the resulting space of periods then gives a common scalar. The approximating datum is not necessarily a single product of two edge vectors. This is why the resulting scalar independent of the twist is not asserted nonzero in the general square law. The exact square assertion is `one_square` in `New/Exchange/OneSquare.lean`; we use it, including this potentially vanishing scalar, as a *Formal-source input.*

Here is the finite-ring part of this assertion. Let $R = \mathcal{O}_E/\mathfrak{m}$. For a prescribed pair of rectangular matrix index sets and an identification σ_0 of complementary dimensions, partners are the finite data in `IsPartner`; they are defined with the character twist and complementary full-rank blocks. The function of the edge datum on one side of a square is $F(X)$, and on the opposite side $F'(Y)$ (these are complex numbers after applying a specified functional). In the

source a square law of scalar c_0 consists of the following three conditions:

$$\begin{aligned} F(kX) &= \chi(\det k)F(X), \\ F'(Yl) &= \chi(\det l)F'(Y), \\ \sum_Y \Psi(Y)F'(Y) &= c_0 \sum_X \Phi(X)F(X) \quad ((\Phi,\Psi) \text{ partners}). \end{aligned}$$

Here k, l are invertible matrices of the appropriate sizes. In the application $c_0 = \kappa_0/c(\theta)$ with $c(\theta) \neq 0$ and κ_0 independent of θ. This definition is `SquareLaw` in `New/Exchange/SquareLaw Fourier.lean`.

The actual periods to be moved use point data and inclusion matrices. The square law is applied not to a single arbitrary point on one side and a single arbitrary point on the other, but to appropriate averages of points. For example the finite transformation called `nextDatum` sends a matrix function p to the function which, on block matrices

$$Y = \begin{pmatrix} y & * \\ 0 & I \end{pmatrix},$$

takes value $p(-y)/|\operatorname{Mat}_{n'\times b'}(R)|$, and is zero on other matrices; here n', b' are just the relevant block sizes. For this particular change the identity `SquareLaw.sum_eq_nextDatum` reads

$$|\operatorname{GL}_{a'}(R)| \sum_y p(y) F'\begin{pmatrix} y \\ I_0 \end{pmatrix} = c_0 |\operatorname{GL}_{n'+b'}(R)| \sum_x p^+(x) F\begin{pmatrix} I_0 & x \end{pmatrix}.$$

The inclusion matrices I_0 have the sizes prescribed by the block decomposition. This is a *Formal-source input* from `New/Exchange/SquareLawFourier.lean`. Its shape reflects invariance under elementary block changes and averaging over the unconstrained $*$ block. Crucially, a determinant character is evaluated only on invertible blocks; their values are roots of unity.

To make the transport of a finite family of data precise, consider a grid of spaces and put one residue matrix on each oriented edge $e = (P, P')$. If X_e is the matrix on that edge, a family of vertex matrices k_P acts by

$$X_e \longmapsto k_{P'}^{-1} X_e k_P.$$

A functional Λ on tuples of residue matrices is called tidy of order v on a set of edges $\mathcal{E}$ if it satisfies character covariance for admissible families k of vertex matrices,

$$\Lambda(f \circ k) = \left(\prod_{e\in\mathcal{E}} \chi_e\big(\det k_{P'} / \det k_P\big) \right) \Lambda(f),$$

and if it is a sum of at most Q_*^v point evaluations with nonzero coefficients roots of unity, supported on tuples whose matrices on the distinguished ramified edges have left inverses. Here

$$Q_* = |R|^{(2n_m)^2}, \qquad n_m = k + m(t+1)$$

is the bound for the number of matrices on any one edge. The precise admissible vertex families are those satisfying the endpoint level conditions and the prescribed diagonal block relation between the two grids. This is `IsTidy`, whose short-sum component is `IsShortSum`, in `New/Exchange/GridMove.lean`. The distinction between the full tidy datum and a merely level-invariant datum matters at the next square.

One elementary observation in the formal source is that orders add: if, for each point S, its evaluation is replaced by at most Q_*^w evaluations with root-of-unity coefficients supported

on valid new tuples, then a tidy short-sum of order v becomes a short-sum of order $v+w$, provided covariance is checked for the new tuple action. Indeed expand the original sum and then the replacement sum. The number of terms is at most $Q_*^v Q_*^w$, and all nonzero new coefficients are products of the permitted roots of unity. The covariance must be verified separately; it does not follow from this count. The counting statement is `IsShortSum.bind` in the same file.

The actual replacement at a square is `IsTidy.exists_square_move` (`New/Exchange/GridMove.lean`), a *Formal-source input*. Its hypotheses are the square law, the matching identifications on the edge blocks, and the compatibility of an admissible vertex family with those block embeddings. Its proof uses the finite kernel δ of `graphKernel` in `New/Exchange/SquareLawFourier.lean`. For a matrix $k \in \mathrm{GL}_N(R)$ and a decomposition $N = a' + b'$, write x_k for the last a' rows of k^{-1} and y_k for the first b' columns of k. The kernel supported on such pairs is

$$\delta(x_k, y_k) = \frac{\chi(\det k)}{|\,\mathrm{GL}_{b'}(R)|}.$$

It has the following useful property: if $\Psi(yl) = \chi(\det l)^{-1}\Psi(y)$ for $l \in \mathrm{GL}_{b'}(R)$, then

$$\sum_y \delta(x_k, y)\Psi(y) = \chi(\det k)\Psi(y_k).$$

In fact for fixed x_k its complementary columns span $\ker x_k$, a free direct summand, and its full-rank column frames are $y_k l$ as l runs through $\mathrm{GL}_{b'}(R)$. Replacing k by $k\,\mathrm{diag}(l, I)$ keeps x_k fixed, transforms y_k by l, and multiplies the determinant character by $\chi(\det l)$. The two character factors cancel term by term, proving the formula. Changes in the other complementary block that preserve both data change k by a block unipotent, so the definition is unambiguous. This is `sum_graphKernel_mul` in the source. It explains the cancellation of the averaging denominator: without the covariance, individual coefficients $1/|\,\mathrm{GL}_{b'}(R)|$ would eventually destroy the lattice estimate. The square move uses further block support and equivariance identities to normalize the graph kernel so that nonzero coefficients are roots of unity. This last assertion is included in the formal-source square-move input.

31.6 The two-grid exchange and the crude rows

We now describe the endpoint transport used for the first of the two height estimates. The grid has two families of vertices; in the notation of the source a vertex is (z, i, j) with $z = 0, 1$ and $0 \le i, j \le m$. Their ranks are

$$N(z, i, j) = r + z + i(t + z) + j(t - z). \tag{31.10}$$

The grid with $z = 0$ has equal rank increments t on both axes, whereas in the second grid the increments are $t + 1$ and $t - 1$. Thus the two diagonal vertices $(0, j, j)$ and $(1, j, j)$ have ranks differing by one. The sign count at offset zero is $\lfloor (m - j + z)/2 \rfloor$; in particular the endpoint $j = m$ is definite in both grids. The initial and final diagonal pairs differ by a line of sign $(-1)^j$. At the real places, the Harish–Chandra parameters are the initial r-block together with blocks built from the weights w_l of the construction and their negatives. The spacing $h = 2r^8$ ensures the required gap conditions and the terminal parameters interlace. These rank, sign, gap and interlacing assertions are the *Formal-source inputs* `card_gridIndex`, `card_neg_gridSigns`, `gridDiagEquiv_spec`, `hasGap_gridParam` and `gridParam_interlace` in `New/Exchange/GridParameters.lean`; (31.10) itself follows by counting the disjoint blocks.

In grid z, the path of stage j starts at $(z, 0, m)$, descends through the vertical edges to $(z, 0, j)$, then lifts along horizontal edges to (z, j, j). The union of these two paths is the path of stage j. A pair of forms on the two starting definite groups and a tidy datum on the path gives a period at the two diagonal endpoints: one of the forms is restricted through the codimension-one Hermitian embedding, with the real translation and line multiplier specified by the grid, and then multiplied by the other and integrated.

The reason for using the second grid is the following separation result.

Proposition 31.9 (Starting periods; formal-source input)**.** *There exist tidy data Λ_θ of order $4m$ on the stage-zero path, a fixed real translation, a scalar α, and a linear surjection R_0 from the specified endpoint form space on the second grid onto the full initial principal-level row form space, such that, writing ϕ for a form in the domain and Ξ for a form used to test the first endpoint,*

$$[K_f : J_m(\theta)] \int_{[U(Y_m)]} \Xi(u)\ \mathrm{lift}_\theta(R_0\phi)(u)\, du = \alpha\ \mathrm{gridPeriod}_0(\Lambda_\theta; \Xi, \phi).$$

Here $J_m(\theta)$ in the index is the level `Data.levelY` *at m as in the period statement; the integral uses exactly the source's bilinear, not Hermitian, period normalization. The map R_0 and real translation are independent of θ.*

The declaration is `Data.exists_isTidy_integral_eq_gridPeriod` (`New/Exchange/CrudeRows.lean`). The surjectivity input in its proof is stronger than existence of nonzero periods for some forms: by `Data.eq_zero_of_orthogonal_gridLevelForms` in `New/Exchange/RowsReached.lean`, there is one real translation for which a form orthogonal to all the restricted forms on the second grid is zero. It employs the real gap construction `Data.exists_isSteep` and the restriction theorem `eq_zero_of_petersson_restrict_eq_zero`. The descent from the definite endpoint onto that grid's level forms is onto by `Data.gridLevelForms_eq_map_iotaPlus`; one uses injectivity of the adjoint weighted lift chain to prove surjectivity. Thus a fixed linear surjection, rather than a different map for each row, is available.

For $j < m$, transporting the two paths from stage j to stage $j + 1$ requires commuting a descent past j lifts in each grid and moving across the remaining diagonal configurations. The source packages this precisely in `Data.hasRoute_succ` in `New/Exchange/CrudeRows.lean`, a *Formal-source input*: the cost and increase in the tidy order are at most $4j + 2$, and the initial period equals a scalar independent of the twist times the inverse of a nonzero factor of that cost times the new period (in the direction fixed by the source's `HasRoute`). Its proof uses the row ladder `Data.descent_past_gridPathMap` in `New/Exchange/RowLadder.lean` and the elementary square move. Iterating the statement is elementary: the sums of costs and of orders are bounded by

$$\sum_{j=0}^{m-1} (4j + 2) = 2m^2.$$

Products of scalar factors respect the bound by Lemma 31.7. Therefore the start period can be transferred to the terminal stage with one factor $c(\theta)$ costing $2m^2$ squares, and with tidy endpoint data of order $4m + 2m^2$. This is `Data.gridPeriod_climb`; in particular the switch to the terminal endpoint is not achieved with exponentially many independent normalization factors.

At the terminal diagonal pair (z, m, m) both unitary groups are definite. The interlacing noted after (31.10) gives a bilinear branching pairing of their Weyl coefficient spaces, represented by an integer matrix. The terminal composite consists of two definite chains, each beginning at $(z, 0, m)$ and lifting to (z, m, m). Let $\mathrm{endPeriod}(\Lambda, y)$ be the resulting

class-integral pairing, linear in the tidy datum and bilinear in the endpoint forms $y = (y_0, y_1)$, as defined in `Data.endPeriod` (`New/Exchange/GridMove.lean`). The terminal levels are chosen so that the coefficient integration can be computed as a finite class sum with the branching matrix. For later use the following are the precise uniform estimates.

Proposition 31.10 (Terminal periods; formal-source input)**.** *There is a positive integer D_*, independent of the twist, with*

$$D_* \le \exp(x^{99}b^2)$$

such that $D_*\,\mathrm{endPeriod}(\Lambda, y)$ is integral at every $\ell \nmid N(Q)$ whenever Λ is tidy on the terminal path and both endpoint forms have gauge at most one and are fixed by the terminal endpoint levels. For the trivial twist and tidy data of order $4m+2m^2$, if every coefficient-conjugate value of the two input forms has squared norm at most S_0, S_1, respectively, then at any $\sigma \in \mathrm{Aut}\,\mathbb{C}$,

$$|\sigma(\mathrm{endPeriod}(\Lambda, y))| \le \exp(x^{200}b^2)\sqrt{S_0 S_1}.$$

These are `Data.exists_mul_endPeriod_mem_embValRing` and `Data.endPeriod_le` in `New/Exchange/CrudeRows.lean`. The first clears the denominators of the successive theta kernels with the standard vacuum integrality statements, and the final branching factor with an integer involving powers of the level and N_1. The bound `boxes_endWeightData_le` bounds the algebraic weights and their box counts. For the second estimate the same finite branch weights bound the size of all derivatives and the branching pairing; the class numbers at each endpoint are bounded using `card_stdClasses_le_exp_mul_log`, and the tidy-data bound counts the residue terms. Although the trivial twist is used for this archimedean estimate, every nontrivial twist is covered by the finite-place estimate away from its own conductor. Thus one does not need an archimedean norm bound growing uniformly in all the permissible twist conductors.

We record the result of the terminal-period calculation with its precise quantifiers.

Proposition 31.11 (Crude rows; formal-source input)**.** *There is a nonzero function $c(\theta)$ of the twists costing $2m^2$ squares, and a spanning set $\mathcal{C}_0$ in the full principal-level row space (before good projection), such that for $y \in \mathcal{C}_0$:*

1. *$(c(\theta)/c(1))\,\mathrm{lift}_\theta(y)$ is coefficient-integral at every embedding over every $\ell \nmid N(Q(\theta))$;*
2. *every coefficient-conjugate value of the untwisted lift has squared Weyl norm $\le \exp(x_1^2 b^2)$.*

The declaration is `Data.crude_rows` (`New/Exchange/CrudeRows.lean`). Its construction combines Proposition 31.9 and the route calculation with Proposition 31.10, choosing spanning short integral forms at the second definite endpoint. For the first definite endpoint use delta forms as coefficient tests: if the pairing with every integral coefficient test of a vector lies in the valuation ring, the dual Weyl lattice criterion gives its integrality. The branching matrix and denominators in the dual pairing are accounted for in the common rescaling of the row forms. The source obtains a common normalization independent of θ besides the ratio $c(\theta)/c(1)$; this uniformity is part of the proposition and indispensable for the one-determinant argument.

To apply the proposition, project the spanning set to good rows with the integral normalized projector. Its integrality assertion is `goodProjTranslate_mem_integralForms` and its metric bound is `normSqAt_goodProjTranslate_le` in `New/Rows/MonomialCoefficientVector.lean` (both are used in `New/Endgame/LongForms.lean`). These are *Formal-source inputs* concerning the given finite levels. Select $n - q$ projected members extending $y_1, \dots, y_q$ to a basis; this is possible by choosing a basis of the quotient of W by their span from the projected spanning set. Their integral normalization is $\gamma_\iota = c(\theta_\iota)/c(1)$. Its inverse squared norm admits height bound

$$g_0 = 4m^2 t x^{999}.$$

Complete the local recombining matrix for the original q rows by a diagonal block $\gamma_\iota I_{n-q}$ to make the new basis integral for the same twist. The determinant condition of Lemma 31.5 holds, with $h = n - q$.

The initial archimedean upper bound for the extended full determinant is of the form $V + (n-q)A_*$, obtained from Lemma 31.2 after conversion of the crude value bound to the class-sum norm. The exact size estimate, which is a calculation with the class-count bound, the normalized projector and the admissible magnitudes, gives

$$A_* + 4m^2 t x^{999} \le \tfrac{1}{4} b^3.$$

This is the final per-vector numerical input in `Data.heightLe_volume_rows` (`New/Endgame/LongForms.lean`); it is a *Formal-source input* as regards the bounds for the coefficient models. Its scale is natural: the leading crude exponent $x_1^2 b^2$ is $o(b^3)$ since $x_1^{10^6} \le b$, and the remaining polynomial losses and class logarithms are much smaller than b^3. The final step does not require these terms separately to be smaller than the sharp error L.

Write from now on $Z(\theta) = \det(\Theta_\theta^* \Theta_\theta)$; injectivity, used for the determinant denominator, is proved below. Applying Lemma 31.5 and using $n - q \le 4n/b^2$ gives

Proposition 31.12 (Forward ratio)**.** *For the indicated family of short rows and fit family θ_ι, one has*

$$\mathrm{Ht}_{\le V + 2d_{\mathrm{ind}} + nb}\left(\frac{Z(\theta_\iota)}{Z(1)}\right). \tag{31.11}$$

Proof. The preceding determinant calculation gives a bound $V + 2d_{\mathrm{ind}} + (n-q)(A_* + g_0)$. Bound its last term by $(4n/b^2)(b^3/4) = nb$. This is the implication from the precise crude-row theorem to the forward-ratio estimate. □

31.7 The triangle of weighted lifts

The chain determinant is not a multiplicative determinant of ordinary square matrices: the Hermitian spaces increase in dimension and the lift maps take their values in proper subspaces. We need an exchange computation to make the determinant on the initial row space factor. The formal source uses an abstract triangle which we now describe. Fix a twist and let $F_{i,v}$ be spaces on the triangular array $i + v \le m$. Use arrows

$$a_{i,v} : F_{i,v} \longrightarrow F_{i,v+1}, \qquad d_{i,v} : F_{i,v+1} \longrightarrow F_{i+1,v}.$$

The $v = 0$ spaces contain isometric copies $j_i(U)$ of a common Hilbert space U of forms. Let $w_v > 0$ be the weights (integral finite group orders in the application). Define

$$A_{i,v} = (w_{v-1} a_{i,v-1}) \cdots (w_0 a_{i,0}), \qquad A_{i,0} = 1.$$

The first-arrow adjoint law, on $j_i(U)$ and all target vectors z, is

$$\langle z, a_{i,0} j_i(u) \rangle = \langle d_{i,0} z, j_{i+1}(u) \rangle.$$

Moreover $d_{i,0} a_{i,0} j_i(u) = j_{i+1}(B_i u)$ for an injective $B_i : U \to U$. For the other squares the hypotheses contain a scalar $\beta_{i,v}$ with

$$w_v a_{i+1,v} d_{i,v} = \overline{\beta_{i,v}}^{-1} w_{v+2} d_{i,v+1} a_{i,v+1}. \tag{31.12}$$

The square scalar is potentially zero in the initial exchange statement. For norms, the source has an additional adjoint version of the square, including an adjoint vector for a horizontal

arrow, and a factor of cost zero. All these conditions together are precisely `IsTriangle` in `New/Exchange/TriangleNorms.lean`; (31.12) is interpreted with the source's zero-inverse convention when necessary. The induction below on injectivity does not divide an unknown zero scalar.

The weighted chain satisfies the commuted-descent identity

$$A_{i+1,v}d_{i,0}a_{i,0} = \frac{w_{v+1}}{w_1 w_0 \overline{\beta}}\, d_{i,v}A_{i,v+1} \tag{31.13}$$

for a product of square factors (with the normalizations of the source). Here at $v = 0$ the factor is $1/w_0$. This follows by applying (31.12) as one moves the descent past each arrow in order; the finite-rank weight ratios telescope. Successive adjoint laws in the norm case give the compatible identity

$$\langle z, a_{i,v}A_{i,v}j_i u\rangle = \kappa\langle d_{i,v}z, A_{i+1,v}j_{i+1}u\rangle \tag{31.14}$$

with the product $\kappa\beta$ costing at most v squares. In the general injectivity case only (31.13) is used. These are the two assertions of `IsTriangle.weightedChain_descent`. The source proves their simultaneous induction beginning with the first-arrow adjoint law; each adjoint square supplies one factor, and the extra zero-cost factor leaves the prescribed budget unchanged. We take the full normalized adjoint identities as *Formal-source inputs* from that declaration.

Lemma 31.13 (Injectivity of the weighted chain)**.** *For $i + v \le m$, the restriction of $A_{i,v}$ to $j_i(U)$ is injective.*

Proof. Use induction on v. This is clear for $v = 0$. If $A_{i,v+1}j_i u = 0$, apply (31.13): $A_{i+1,v}d_{i,0}a_{i,0}j_i u = 0$. Since its argument is $j_{i+1}(B_i u)$, the induction hypothesis and injectivity of B_i give $u = 0$. This proves `IsTriangle.eq_zero_of_weightedChain`. □

Lemma 31.14 (Gram identity of the triangular array)**.** *Suppose a subspace $W \subset U$ is preserved by the first-arrow Gram operators S_i characterized on W by*

$$w_0\langle a_{i,0}j_i u, a_{i,0}j_i v\rangle = \langle u, S_i v\rangle.$$

In the adjoint case of `IsTriangle`, for $i + v = m$ there is a scalar η of cost at most v^2 such that the Gram form of $A_{i,v}j_i$ on W equals

$$\eta\frac{w_0 w_1^v}{w_v}\langle u, T_i v'\rangle, \qquad T_i = S_i S_{i+1}\cdots S_{m-1},$$

where v' is the second vector, and the product is the identity for $v = 0$. In particular $\det T_i = \prod_{h=i}^{m-1} \det S_h$.

Proof. The first-arrow adjoint law implies $w_0 B_i = S_i$ on W when paired against vectors of W (equivalently one uses its preserving operator on the full common space). For the inductive step take the Hermitian pairing of the last arrows, use (31.14) on one side and (31.13) on the other. The factors of w_v, w_0, w_1 combine to give $w_0 w_1^{v+1}/w_{v+1}$, with scalar multiplier contributed by $\kappa\beta$ at cost at most v. Apply the induction hypothesis in row $i + 1$ to S_i and the remaining chain, using self-adjointness of its Gram form to put S_i in the first argument of the endomorphism product. The determinant is independent of the resulting product order, so even without commutativity the conclusion about determinants holds. Starting with cost zero at length zero and increasing the bound by no more than $(v + 1)^2 - v^2$ proves the stated cost. This is the induction in `IsTriangle.inner_weightedChain_eq`; the adjoint normalizations and the exact scalar in this calculation are the ones in (31.13)–(31.14). □

For the application one uses the triangle on the left vertical strip of the preceding grids, with a varying offset i. At offset i and stage v the space of forms is that of the real parameter `gridParam` and signature `gridSigns`; the initial forms are identified by negating signs and using the isometry `formNegEquiv`. The weights are $w_v(\theta) = \mathrm{triangleOrder}(\theta, v)$, computed by the order of the finite group of the relevant modulus. The prescribed real vectors harmonics and local inclusion data define the arrows `triLift`. The first-arrow adjoint law follows from the usual pairing of lift and descent; the finite normalizing order is important for the square identity. The first-arrow injectivity uses the Rallis-type norm calculation of `hasSum_inner_triLift`, with the chosen large rank. The theorem `exists_isTriangle` in `New/Exchange/TriangleArrows.lean` verifying these requirements is a *Formal-source input*. The additional adjoint condition for the Gram identity holds in the $z = 0$ grid.

To relate these operators to the single lifts we need the Hecke transfer between groups with different signatures. For $1 \le j \le m$, let U_j denote the good form space on the auxiliary group of rank r with $m - j$ negative signs, and let $V_j(\vartheta)$ denote the definite algebraic form space of the auxiliary pair of ranks $(r, r + t)$. Write

$$u_{j,\vartheta} : U_j \to V_j(\vartheta)$$

for the good lift `Data.goodLift`, which is the adjoint of a normalized theta descent. We reserve the letter W for the good rows on Y_0, not for the Hermitian space of the same symbol in the code. The following precise transfer statements are used.

Proposition 31.15 (Rallis pairing and transfer; formal-source inputs)**.** *For every j, the map $u_{j,\vartheta}$ is injective for every allowed twist ϑ, and $\dim_{\mathbb{C}} U_j = n$. The pairings defining the squared norms of the first arrow in row $j - 1$ of the triangle and the auxiliary lift u_{j,θ^j} are, up to the explicit constants and finite group orders, sums over the same family of Hecke terms. In particular they are represented by one endomorphism in the common off-level Hecke algebra acting on the good form spaces. For every element P of that algebra, the characteristic polynomials of its actions on W and U_j agree.*

The first-arrow formula is `hasSum_inner_triLift` (`New/Exchange/TriangleArrows.lean`). The auxiliary-lift formula is `Data.auxLift_injective_and_hasSum_inner` (`New/Exchange/CharacterLiftNorm.lean`): there is $A_j \neq 0$ independent of ϑ with

$$\langle u_{j,\vartheta} x, u_{j,\vartheta} y \rangle = A_j \delta_{j,\vartheta} \sum_{\gamma} R_{j,\vartheta}(x, y; \gamma).$$

Here $\delta_{j,\vartheta}$ is `AuxPair.gramFactor`, and the terms `Data.rallisTerm` are explicitly the Hecke coset matrix coefficients, with their character factors and divided by the number of $(r + t) \times r$ matrices over the level residue ring. The Rallis identity has a stable-range hypothesis and uses a chosen sharp local datum; the usual abstract nonvanishing of a theta lift alone would give neither this sum nor its transformation under level. We rely on the source's precise identity.

The fact that the series of Hecke terms represents one element simultaneously on the two spaces is `exists_common_hecke_rallis`, in `New/Trace/GoodFormTransfer.lean`. This follows from an elementary finite-dimensional closure step once the local coset expression is known: the images of the common algebra in the product of the two endomorphism spaces, and their corresponding pairings, are finite dimensional and closed. Each common coset term is in this image after averaging over the principal-level subgroup. The convergent sum stays in the image. It need not define a convergent sum in an infinite-dimensional topology on the Hecke algebra itself.

For the characteristic polynomials the essential theorem is `trace_heckeOnGood_eq` in the same file. Its proof isolates each occurring eigensystem by a projector in the common

algebra, and uses primitive label rigidity (`New/Trace/LabelRigidity.lean`) to rewrite the good idempotent, on the eigensystem, in terms of functions supported at classes where the simple trace formula applies. The trace comparison is a transfer result on the specific inner forms, with the cuspidal type and gap assumptions; it is not implied by the same rank of two groups. Once the traces agree for all P, however, the rest of the conclusion is ordinary algebra. Taking $P = 1$ shows equality of dimensions. The traces of all powers P^h agree and determine its characteristic polynomial by Newton's identities. This proves the determinant comparison `det_heckeOnGood_eq` and the rank identity `finrank_rows_eq_finrank_goodForms`. The assertions of trace comparison and primitive label rigidity here are formal-source inputs; the elementary polynomial implication was just proved.

Let

$$\lambda_j(\theta) = \det(u^*_{j,\theta^j} u_{j,\theta^j}) > 0.$$

The positive quantities λ_j cannot be assumed algebraic merely because they are determinants of spaces of forms; the height inequalities are statements about chosen ratios with uniform normalization scalars. The inner product on the final form space of the main chain is a class-sum inner product. Relative to the Petersson metric of the triangular chain there is a conversion factor

$$J(\theta)\, D_m, quad J(\theta) = [K_0 : J^{\mathrm{res}}_m(\theta)], \quad D_m = \#\,\mathrm{WeylIndex}(Y_m, \kappa_m).$$

The identity `stdLevelIndex_lastResLevel_eq` in `New/Exchange/ChainNorms.lean` expresses $J(\theta)$ as

$$CN(Q)^{k_0} \delta^m_{1,\theta} w_m / (w_0 w_1^m).$$

Here k_0 denotes the (integral) exponent called k in that declaration; it is unrelated to the rank of columns. The constants C and k_0 do not vary with the twist. This count compares block parabolic stabilizers in the quotient and is a *Formal-source input*; it cancels the triangular Gram factor $w_0 w_1^m / w_m$ exactly. The formula `inner_iotaAtLevel` in the same file gives the metric conversion. Injectivity follows from Lemma 31.13. Taking determinants of the resulting Gram comparison and applying Proposition 31.15 stage by stage yields the following identity.

Proposition 31.16 (Volume factorization)**.** *For every allowed m-fold twist, Θ_θ is injective; there exists a function $c(\theta)$ of cost m^2 squares such that*

$$Z(\theta) = |c(\theta)|^n \prod_{j=1}^{m} \lambda_j(\theta) > 0. \tag{31.15}$$

This is `Data.volume_iota_eq_prod_liftVolume` (`New/Rows/RowVolume.lean`). To check the scalar exponent, multiplying a Hermitian Gram form by a scalar multiplies its determinant by the nth power of that scalar; the factor in (31.15) is $|c|^n$, not $|c|^{2n}$. The residual factors resulting from the metric conversion, finite group orders and *delta* factors have squared norm equal to a fixed constant times an integral power of $N(Q)$, so they cost zero by Lemma 31.7. All that remains is the triangular cost of m^2. This explains both the injectivity presupposed in the forward-ratio calculation and the factorization required for the reverse ratio. The normalized scalar identities in this calculation are precisely those of the cited formal-source theorem.

31.8 Comparison forms for one lift

We next establish the height estimate for $\lambda_j(1)/\lambda_j(\theta_\iota)$ for each j. Fix j and abbreviate $u_\vartheta = u_{j,\vartheta}$, $Z_\vartheta = \mathrm{im}\, u_\vartheta$. In the next few paragraphs Z_ϑ is a subspace, not the scalar $Z(\theta)$.

The strategy is to use prescribed algebraic forms in the definite target, not to estimate the unknown Petersson norms of every automorphic representation separately.

Three character pairs are required for this particular j. Let $e_+ = m$, $e_- = m/2$, and $e_0 = 7(m/2) + k/2$ (the admissible parities make these integers). Write $p_i = (\nu_i, e_i)$ for $i \in \{0, +, -\}$. The pairs $p_\pm$ have character $\upsilon^{-e_\pm}\chi^{\pm j}$, where υ is the fixed rank-normalizing character and χ is the chosen tame character. The character in p_0 is $\mu_j^0 \upsilon^{j+\lfloor (t-11m-1)/2 \rfloor}$; here μ_j^0 is the untwisted character determining the jth lift. Use the twist ϑ positively for p_0, p_+ and negatively for p_-. Put

$$\Gamma_j(\vartheta) = \prod_{i=0,+,-} \Pi_{M\cup Q}((\nu_i \vartheta^{s_i})^{-1}, e_i), \quad s_0 = s_+ = 1, \quad s_- = -1. \tag{31.16}$$

This is `Data.heckeValueProd`, from `New/Rows/Separation.lean`; that file gives the precise pair definitions as `Data.linePair`. The long string p_0 will yield the term $k\mathcal{U}$, whereas the two other strings contribute to the small error.

Proposition 31.17 (Comparison maps; formal-source input)**.** *There is a finite-dimensional complex table space T, independent of ϑ, a surjection $A : T \to U_j$, and for every twist a linear map $g_\vartheta : T \to V_j(\vartheta)$ and $\gamma_\vartheta \neq 0$ such that*

$$u_\vartheta^* g_\vartheta = \gamma_\vartheta A, \tag{31.17}$$

$$|\gamma_\vartheta/\gamma_1|^2 = \Gamma_j(\vartheta)/\Gamma_j(1). \tag{31.18}$$

The vectors $g_\vartheta A^* v$ are tested relative to $Z_\vartheta^\perp$ for all $v \in U_j$. At every $\ell \nmid N(Q)$ and embedding ϕ, $g_\vartheta \zeta$ is integral whenever the conjugate tables $\bar{\zeta}$ are integral at $\phi \circ \kappa$ with the prescribed Weyl gauges of their classes. Finally

$$h_j \|g_1 \zeta\|_\sigma^2 \le e^{2rB_0} \sum_\alpha \|\sigma \zeta_\alpha\|_{\mathrm{Weyl}}^2,$$

where $h_j > 0$ is the hook number of the small algebraic weight.

This is exactly `exists_comparison_forms` in `New/Rows/ComparisonForms.lean`. We describe the detailed construction to specify its content. Split the definite Hermitian space in the auxiliary pair into $V'' \oplus V^\circ$ (the `DividedSplitting` data), and take tables indexed by a finite member index and by representatives of the small principal level $\bar{\pi}_g^2$ on V°. Their entries lie in the small Weyl module of weight given by `colsMinus`; thus h_j is the hook number of this module. To a member index corresponds a split lattice obtained by gluing the two blocks in the split model along the residue matrix. The same glued lattice can be chosen for every twist. The group in the large block which fixes the specified rows modulo $\bar{\pi}_g^2 Q$ has free class action; this is `classAut_rowFixingLevel_eq_bot`. If $y_{\vartheta,i}(h)$ is the line multiplier for member i, the comparison map is the algebraic form

$$g_\vartheta \zeta = g^{\mathrm{loss}} \sum_{(i,c)} \sum_{h \in X_i} \mathrm{productForm}_{L_i}((h,c), \overline{\beta_1 y_{\vartheta,i}(h)}, \zeta_{i,c}).$$

Here L_i is the glued lattice, X_i is the class set for the row-fixing level on the large block, β_1 is one nonzero common line scale, and the recipe called `productForm` applies the Cartan product of the representations in the divided splitting. This is the actual `comparisonForm`. The exponent loss is the one in the divided splitting; it clears the gluing loss from off-diagonal blocks. The source proves integral and small bounds for the line multiplier after applying β_1 in `Data.lineMultiplier_integral_and_small`, and controls the Cartan product in `New/Rows/CartanIntegrality.lean`. Multiplying a delta form by this product and summing

over a free class set preserves finite-place integrality. The archimedean estimates use the number of tables, classes and weights, and multiply the line bound with the squared Weyl norm of the input table. The output estimate is `normSqAt_comparisonForm_le` and the input integrality is `comparisonForm_mem_integralForms`. These assertions concerning the glued finite and infinite kernels are formal-source inputs.

The resulting inner product with a lift of $F \in U_j$ has a common period expression independent of the twist up to scalar: it is a weighted sum over members i of

$$\int_{[U(W_j)]} \overline{F(u)F_i^\natural(u)}\,\theta^\circ(\zeta_{i,\bullet})(u)\,du.$$

Here $F_i^\natural$ is a fixed natural form, and θ° the theta kernel from the small block. The scalar is determined by `Data.cosetKernelSum_eq_fNatural`, which gives $F_i^\natural$ as the coset kernel sum involving the line multiplier at every twist times κ_ϑ satisfying

$$|\kappa_\vartheta|^2 = C/\Gamma_j(\vartheta).$$

This is initially an assertion potentially with $C = 0$. The equality of period expressions is `inner_adjoint_comparisonForm`; the nonvanishing and surjectivity use the separation theorem `Data.eq_zero_of_integral_fNatural_tableTheta`. This states that if a good F gives zero in every displayed period, then $F = 0$. In fact if $\kappa_\vartheta = 0$, all $F_i^\natural = 0$ and the separation theorem forces $U_j = 0$, contrary to $n > 0$. If the range of $u_1^* g_1$ were not all of U_j, a nonzero orthogonal vector would annihilate all the periods, again contradicting separation. Set $A = u_1^* g_1$ and $\gamma_\vartheta = \overline{\kappa_1/\kappa_\vartheta}$. The period identity gives the first equation of Proposition 31.17; the quotient of the two norm identities for κ gives (31.18). This explains the proof of nonzero scalars and the fact that a *single* common surjection works for every twist. The equality of coset kernel sums and the separation theorem are *Formal-source inputs* in `New/Rows/Separation.lean`.

The tested condition in the proposition is equally substantive. Off level, the Hecke polynomial of a genuine lift has a product of t known roots and an affine rescaling of the original Hecke polynomial. Explicitly, if the auxiliary ranks are $r, r+t$, the polynomial at a split place w is

$$\prod_{v=0}^{t-1}(1 - q_w^{v+r/2}a_O(w)X)P(q_w^{t/2}a_O(w)b_O(w)^{-1}X)$$

in the character convention of `Data.lifts_auxLift` (`New/Exchange/CharacterLiftNorm.lean`); $a_O(w), b_O(w)$ in that display indicate the source's rank-normalized Frobenius values, and P is the source Hecke polynomial. In particular the test operator vanishes on the genuine lift, as expressed by `heckePolyOnForms_testValue_eq_zero`; one must use the test point in its precise source normalization, not insert a point from a different convention. A false lift satisfies this Hecke test and lies in the good part but is orthogonal to every actual lift. Its theta descent to the smaller group has zero good part (by adjointness and equivariance).

The crucial vanishing of the comparison periods for a false lift comes from enlarging the descent group by hyperbolic planes. The good polynomial labels imply that the reflection sums associated with the chosen tame character act by zero. Together with the vanishing of the initial descent and nontriviality of the character at the twist places, this kills the relevant boundary terms in the constant-term expansion. At a split place off level, a bounded descended form killed by a Hecke polynomial at a sufficiently small test point must vanish: expand the Hecke polynomial, use the sum of norms of the nonconstant terms strictly below one, and compare the sup norm. The precise statements `AuxPair.descentsVanish_of_reflectionSum`, `eq_zero_of_heckePolyOp_eq_zero` and `descentsVanish_add_planes` in `New/SiegelWeil`

`/AuxPairVanishing.lean` and `New/SiegelWeil/DescentsVanishPlanes.lean` formalize these steps. The period which results from the glued product is zero by `DividedSplitting.stdClassSum_inner_productForm_eq_zero` in `New/Rows/MemberOrthogonality.lean`. Applying that statement to the comparison construction gives `tableTheta_adjoint_eq_zero`: for a false lift Ξ, each small form $\theta^{\circ}((g_{\vartheta}^{*}\Xi)_{i,\bullet})$ vanishes. Consequently the common period formula gives

$$\langle \Xi, g_{\vartheta} A^{*} v \rangle = \langle A g_{\vartheta}^{*} \Xi, v \rangle = 0$$

for each v and every false-lift space. These special vanishing assertions are *Formal-source inputs*; the last implication is merely the adjunction identity. The source does not prove that arbitrary integral vectors satisfy the tested condition.

31.9 Compressed Gram determinants

Let p_{ϑ} be projection onto Z_{ϑ}, with respect to the class-sum inner product at the ordinary embedding. Define on Z_{ϑ}

$$D_{\vartheta} = \det(h_j p_{\vartheta} g_{\vartheta} g_{\vartheta}^{*}|_{Z_{\vartheta}}).$$

The key determinant calculation is independent of automorphic theory. Regarding u_{ϑ} as an isomorphism onto Z_{ϑ}, adjunction and Proposition 31.17 give

$$h_j u_{\vartheta}^{*} g_{\vartheta} g_{\vartheta}^{*} u_{\vartheta} = h_j |\gamma_{\vartheta}|^2 A A^{*}.$$

Taking determinants and using $u_{\vartheta}^{*} p_{\vartheta} = u_{\vartheta}^{*}$ gives

$$\det(u_{\vartheta}^{*} u_{\vartheta}) D_{\vartheta} = h_j^{n} |\gamma_{\vartheta}|^{2n} \det(A A^{*}). \tag{31.19}$$

Since A is onto, both sides are nonzero and g_{ϑ}^{*} restricted to Z_{ϑ} is injective. Dividing the identities at the trivial and arbitrary twist yields

$$\frac{\lambda_j(1)}{\lambda_j(\theta_\iota)} = \frac{D_{\theta_\iota^j}}{D_1} \left(\frac{\Gamma_j(1)}{\Gamma_j(\theta_\iota^j)} \right)^{n}. \tag{31.20}$$

This elementary identity is `det_compressedGramOp`, `adjoint_comp_eq_conj_smul` and their use in `heightLe_liftVolume_ratio` (`New/Rows/LiftVolumeHeight.lean`).

The next two estimates for the compressed determinant must be proved without assuming that projection to a complex automorphic summand is integral or that it is an orthogonal projection at all coefficient conjugates. We give the determinant-level argument.

Lemma 31.18 (Finite norm-adapted basis)**.** *Let V be a complex vector space of dimension d, let $(\ell_a)_{a \in A}$ be a finite separating family of complex linear functionals on V, and let ϕ be a field embedding into a non-Archimedean valued field. Then some d of these functionals are the dual basis of vectors v_i with $|\phi \ell_a(v_i)| \le 1$ for all a, i.*

Proof. Take any basis of V. Among all d-tuples of the given functionals, choose one maximizing the absolute value under ϕ of their determinant in this basis. The maximum is positive by separation. Use the chosen tuple as a dual basis. Express any other functional in this dual basis by Cramer's rule. Each coordinate is a quotient of a determinant obtained by replacing one of the selected rows by the new functional, by the maximizing determinant. Its *phi*-absolute value is at most one. This proves the assertion. The finite case of `exists_auerbach` in `Lib/General/LinearAlgebra/CommonEigenvectors.lean` is all that is needed here. □

At a fixed place ι, choose rational integral adelic frames for the small-block classes. Denote by f_a special table vectors given by the coordinate duals in those frames, scaled by the hook-number convention. Thus the functional on Z_ϑ is

$$\ell_a(z) = \langle f_a, g_\vartheta^* z \rangle.$$

The duals are integral in the Weyl gauge convention because the Fock Gram matrix times the relevant Weyl Gram matrix is the hook number times the identity (the Fock pairing of standard tableaux supplies the integral dual). In these frames the values $\ell_a(z)$ are exactly h_j times the conjugate-coordinate test of the table $g_\vartheta^* z$. Consequently, selecting a dual basis as in Lemma 31.18, each $h_j g_\vartheta^* z_i$ satisfies the requisite integral table test at the chosen place. The forms $g_\vartheta f_{a_i}$ are integral at all the embeddings used in the determinant test, and

$$v_i = h_j g_\vartheta g_\vartheta^* z_i$$

are integral at ι and tested: adjunction gives $g_\vartheta^* u_\vartheta = \bar{\gamma}_\vartheta A^*$, so the tested condition follows from the comparison theorem for $z_i \in Z_\vartheta$. In this step the assertion on the coordinate duals (including their gauge after conjugation) is the concrete integral-linear-algebra input of `norm_det_compressedGramOp_padic_le`; it relies on the integral frame lemma and `weylGram_mul_fockPairing` rather than on orthonormal coordinates having integral entries.

Apply (31.9) to these two families. The matrix of their pairings

$$\left(\langle g_\vartheta f_{a_i}, p_\vartheta(h_j g_\vartheta g_\vartheta^* z_h)\rangle\right)_{i,h}$$

is the matrix of the compressed operator in the chosen basis, by duality of ℓ_{a_i}. Its determinant is D_ϑ. Hence the determinant bound at twist $\vartheta = \theta_\iota^j$ is

$$|\iota D_\vartheta|_\ell \le e^{nr\beta_\ell}. \tag{31.21}$$

This derives the implication of the denominator test in `norm_det_compressedGramOp_padic_le`; it is precisely why determinant-level denominators suffice.

For the archimedean estimate, use the following property for all conjugate coefficient metrics: if y is tested and Z_ϑ lies in the good kernel of the Hecke polynomial at the appropriately chosen off-level place, then

$$\|p_\vartheta y\|_\sigma \le \|y\|_\sigma. \tag{31.22}$$

This is `AuxPair.normSqAt_starProjection_le` in `New/Selmer/FitTwists.lean`. Here is the mechanism. Write $y = z + x$ with $z \in Z_\vartheta$, x orthogonal to Z_ϑ. For a tested vector, x lies in the sum of the kernel of the good projector and the range of the Hecke polynomial (taking the complementary conjugate-place normalized operator as appropriate). This follows by taking orthogonal complements of the intersection defining the false-lift test, using the normality of the Hecke operators. Under application of a field automorphism σ, the adjoint of $H_w(X)$ for the coefficient-conjugate pairing is $H_{\bar{w}}(\sigma^{-1}(\overline{\sigma X}))$. This is `LevelPlaces.LevelForm.innerAt_heckePolyOnForms`, and the commuting Hecke operators and definiteness of the twisted metric imply that their relevant kernels are stable under taking the adjoint. In addition the good projector remains self-adjoint for every conjugate pairing because its underlying finite character action is unitary (character scalars are roots of unity). Hence z, x remain orthogonal for the σ-pairing. Positivity gives (31.22). The adjoint and projector laws with their precise local hypotheses are *Formal-source inputs* in `New/Selmer/FitTwists.lean`.

Apply this contraction to the vectors $g_1 g_1^* z$ occurring in the compression. One cannot simply apply σ to the spectral bound at the ordinary embedding. Instead, take an eigenvector

z with eigenvalue η, set $\zeta = g_1^* z \neq 0$, and pair the eigenvalue equation with the conjugate-compatible table obtained from ζ by the map $\kappa \circ \sigma^{-1} \circ \kappa \circ \sigma$ on coefficients. The mixed Cauchy–Schwarz inequality of (31.3), contraction (31.22), and the two table norm estimates in Proposition 31.17 then give

$$|\sigma\eta| \sum_{\alpha} \|\sigma\zeta_\alpha\|_{\mathrm{Weyl}}^2 \leq e^{2rB_0} \sum_{\alpha} \|\sigma\zeta_\alpha\|_{\mathrm{Weyl}}^2.$$

The sum is positive by injectivity of $g_1^*|_{Z_1}$. This is the eigenvalue calculation in `norm_det_compressedGramOp_arch_le` (`New/Rows/LiftVolumeHeight.lean`). Thus every conjugate of each eigenvalue has modulus at most e^{2rB_0}, and the determinant, the product of the n eigenvalues with multiplicity, satisfies

$$|\sigma D_1| \leq e^{2nrB_0}. \tag{31.23}$$

Using witness D_1 in the height definition, the local estimate (31.21), the global sum of β_ℓ, and (31.23), we obtain

$$\mathrm{Ht}_{\leq 3nrB_0}\left(\frac{D_{\theta_\iota^j}}{D_1}\right). \tag{31.24}$$

31.10 The height saving in a rank-one special value

It remains to show how derivative divisibility enters the lower bound, in the estimate for $\Gamma_j(1)/\Gamma_j(\theta_\iota^j)$. Fix a pair $p = (\nu, z)$ of rank-one type at j. The definite target of the lift of the trivial character of $U(1)$ has even rank $2z$ and its one-row coefficient weight at τ has degree $d_\tau = c_\tau - z$. Let $\Xi_{p,\vartheta}$ denote this lift, with the CM vacuum and jet scaling, namely `Data.Pair.trivialLift` in `New/Exchange/AlgebraicPartDefs.lean`. It is a scalar multiple of a sum of kernel forms over the classes of the norm-one torus, at the prescribed level. Put $d_{\mathrm{tot}} = \sum_\tau d_\tau$. The scaling of the theta kernel in this sum is

$$\kappa_0^{2z} (\overline{\pi_g}\,\mathrm{per})^{d_{\mathrm{tot}}},$$

where κ_0 is the common vacuum scale, π_g is the chosen level generator and per the jet scale. The integer

$$\mathrm{den}_p = g^{4fz} (gd)^{d_{\mathrm{tot}}}$$

clears its denominators outside its conductor. Let $M_* = \mathrm{chosenGroupOrder}$ be the fixed finite group order bounding the relevant class stabilizers. The normalized integral value used in the source is

$$\mathcal{A}_p(\vartheta) = M_* \,\mathrm{den}_p^2 \,\frac{\|\Xi_{p,\vartheta}\|^2}{N(Q)^{2z-1}}.$$

Nonzero proportionality of this value to $L^{M\cup Q}(z, (\nu\vartheta)^{-1})$ with a common constant independent of ϑ is the rank-one Rallis norm theorem `LValueData.inner_trivialLift_self_eq_L` (`New/Exchange/CharacterLiftNorm.lean`) and `intAlgPart_eq_mul_L` of `New/Exchange/TriangleLevelExponent.lean`. The factorial in the norm is exactly $\prod_\tau d_\tau!$, because a one-row standard monomial has Fock norm its factorial. The assertion that the residual normalization is common, including the indicated conductor power, is a *Formal-source input*; the ensuing ratios satisfy

$$\rho_p(\vartheta, 1) = \frac{\mathcal{A}_p(\vartheta)}{\mathcal{A}_p(1)} = N(Q)^{-(2z-1)} \frac{\|\Xi_{p,\vartheta}\|^2}{\|\Xi_{p,1}\|^2}. \tag{31.25}$$

In particular the required lift is nonzero. That normalized values are values of one group-ring sum with coefficients integral away from Q, when the twist varies at fixed modulus, is

`isIntegralMeasure_intAlgPart` in the latter file. It follows by expanding the squared class sum of kernels into the two torus class indices and pulling the character factor out of their two translated kernels; the remaining coefficients are integral by the norm-one lift denominator estimate and clearing the stabilizers by M_*. These precise integral-measure and norm assertions are formal-source inputs, not appeals to a general interpolation theorem for L-functions.

To estimate the untwisted norm, compare with a flat pair p^0 of the same rank, i.e. of all degrees zero, and with a permissible weight-zero character after rank normalization. Define

$$Q_0 = \mathcal{A}_{p^0}(1)/\|\Xi_{p,1}\|^2.$$

The ordinary rank-one norm itself need not be integral, but $Q_0\|\Xi_{p,1}\|^2 = \mathcal{A}_{p^0}(1)$ is integral at all finite embeddings. The precise archimedean comparison is

Proposition 31.19 (Rank-one conjugate norms; formal-source input)**.** *For fixed rank $2z$, at every automorphism fixing K there is a common c_σ independent of the pair such that*

$$\log|\sigma(\|\Xi_{p,1}\|^2)| = c_\sigma + e_{p,\sigma} + \sum_\tau J(d_\tau), \qquad e_{p,\sigma} \geq -4f.$$

For a flat pair $e_{p,\sigma} \leq 4f$ as well.

This is `Data.Pair.log_norm_map_norm_sq_trivialLift` in `New/Exchange/AlgebraicPartConjugates.lean`, obtained from the permutation and translation of the CM theta coefficients under field conjugation (`LValueData.map_trivialLift`). For automorphisms not fixing K, compose with complex conjugation to reduce the required modulus estimate to this case. Compare the upper bound for the flat numerator with the lower bound for the nonflat denominator. Because the weight of a pair of rank-one type has $d_\tau \geq w_0(\tau)$, the jet-size estimates compare $J(d_\tau)$ with $J(w_0(\tau))$. The extra logarithm of the fixed integer $M_* \operatorname{den}_{p^0}^2$ and the two losses of $4f$ have combined size at most B_0 (the exact bound is `D.le_of_hasLogBound` in `heightLe_norm_trivialLift_sq`). We deduce

$$\mathrm{Ht}_{\leq B_0 - \sum_\tau J(w_0(\tau))}(\|\Xi_{p,1}\|^2). \tag{31.26}$$

This is the declaration `heightLe_norm_trivialLift_sq` (`New/Rows/Separation.lean`). It is an estimate for the constant family with entries the specified squared norm, not for the absolute size of the norm at every place. The common unknown c_σ disappears on taking the ratio.

We need the finite-place lower bound on the pairing with a test form that restores the forced divisibility term. Put $F_\ell = \sum_\tau A_\ell(w_{\mathrm{nom}}(\tau), \tau)$, the quantity `Data.forcedPart`, so that the elementary prime-power formula gives

$$2\sum_\ell F_\ell \log \ell = \sum_\tau A(w_{\mathrm{nom}}(\tau), \tau).$$

All terms have finite support. The relevant depth indicator is `Data.depth_indicator_jet` in `New/Rows/GluedFormTheta.lean`. It is a *Formal-source input*: for the degree d_τ in the allowed interval of that declaration, one common nonzero scale of archimedean conjugates bounded by its prescribed small budget serves all places; locally there is a jet of the prescribed total degrees (distributed among twelve coordinates) with valuation at most the forced depth loss, after multiplication by the scale. The proof in the source uses tight frames and the inverse-theta normal form to realize a large coefficient of the needed jet. An earlier mere upper bound for the norms of all jets cannot replace this attainability statement.

In obtaining an actual test form, the twelve coordinates sit next to flat blocks of ranks $2z-20, 4, 4$. They permit the coefficient of the jet to be detected by a genuine coefficient pairing and absorb the tame character choices through the flat forms. The test-form existence formula is `exists_test_form_inner_kernelForm` (`New/Rows/HeckeValueHeight.lean`). It constructs a form by three class sums weighted by the flat theta lifts in these three blocks. Its pairing with the kernel is an explicit product of their three normalized values, a multinomial coefficient

$$\prod_{\tau}(d_\tau!)/\prod_{\tau,i}\alpha_{\tau,i}!$$

and the remaining rank-twelve monomial jet kernel (with the matching character). The left side contains the class-number normalization and the vacuum-scale power. It asserts the resulting form is integral at every embedding over ℓ and is tested for the rank-one false lifts. The tested assertion relies on the same extra-plane descent vanishing as the comparison form theorem, with the rank-one label projector. This is a separate *Formal-source input*, since the existence of an integral kernel coefficient alone is not enough to produce a class-sum pairing against an integral tested vector.

For clarity there are two elementary finite Fourier steps in passing from these pointwise test-form formulas to the uniform height witness.

Lemma 31.20 (Fourier selection and multiplicative selection)**.** *Let G be a finite abelian group and $s : G \to \mathbb{C}$ any function. For each embedding ϕ in a non-Archimedean field there is a character χ such that*

$$\left|\phi\left(\frac{1}{|G|}\sum_{g\in G}\chi(g)s(g)\right)\right| \geq |\phi(s(1))|.$$

Furthermore, let $a(\chi), b(\chi)$ be two families indexed by a finite character group $G^\vee$, and let Q satisfy $Q^{|G^\vee|} = \prod_\chi a(\chi)b(\chi)$. For every prescribed $\psi \in G^\vee$ there are χ_1, χ_2 with $\chi_1\chi_2 = \psi$ such that

$$|\phi(a(\chi_1)b(\chi_2))| \geq |\phi Q|.$$

The second assertion also holds for Archimedean embeddings.

Proof. Fourier inversion expresses $s(1)$ as the sum of the normalized Fourier coefficients. The ultrametric inequality bounds its absolute value by their maximum. For the second assertion, take the product of the numbers $a(\psi\chi^{-1})b(\chi)$ as χ ranges over all characters. It is $Q^{|G^\vee|}$ by the permutation of the character group, so one absolute value is at least $|\phi Q|$. □

In `exists_twisted_classSum_ge`, the finite group is the group of torus classes after translating the chosen base class to the identity; its stabilizers are trivial and its characters are realized by the permitted tame weight-zero characters. The second step is `exists_factorisation_le` in the same file. These two selections allow the flat class sums to be paired with the depth indicator at every local embedding while a single product of flat values is used globally as the height witness.

Here is the precise quantitative interface obtained by applying these constructions. Fix the rank-one pair p, the index j and the small flat pair p_2 (rank four, tame part the chosen primitive character ω). Let N_c be the torus class number at the principal chosen level. The combination of `depth_indicator_jet`, `exists_test_form_inner_kernelForm`, `exists_twisted_classSum_ge` and `exists_factorisation_le` in the proof of `heightLe_inv_ratio` (`New/Rows/HeckeValueHeight.lean`) yields a common $Q \neq 0$ such that $|\sigma Q| \leq e^{B_0}$ for every

σ, and for each ℓ and each embedding ϕ there exists a tested, everywhere ℓ-integral vector u_ϕ in the rank-one target satisfying, at $\vartheta = \theta_\iota^j$,

$$|\phi Q|_\ell \, \ell^{-F_\ell} \le \left| \phi \left(\frac{N_c^2 \langle u_\phi, \Xi_{p,\vartheta} \rangle}{\rho_{p_2}(\vartheta, 1)} \right) \right|_\ell . \tag{31.27}$$

The witness Q is an hth root of the product of the flat normalized values times the depth scale over the h tame characters, which gives the archimedean bound using the flat-conjugate estimates. For the fixed twist and a different finite embedding the test vector may change, but Q does not. Equation (31.27), including its conjugate-compatible version when one uses the field embedding composed with complex conjugation, is the local assertion called `K` in the proof of that declaration. We take this simultaneous assertion, which uses the attainable jet and all the class and gluing integrality facts just discussed, as a formal-source input. The following calculation reconstructs its numerical consequence.

For an arbitrary nonzero vector v in a complex Hilbert space, projection onto its line is

$$p_{\mathbb{C}v} w = v \langle v, w \rangle / \|v\|^2 .$$

The fit denominator condition for the rank-one lift line therefore bounds the pairings of two integral tested vectors with this projection by e^{β_ℓ}. In the source the strengthened form `DenominatorLe.rank_one_bound` in `New/Selmer/RankOneDenominator.lean` gives

$$|\iota(N_c \langle u, p_{\mathbb{C}\Xi} v \rangle)|_\ell \le e^{\beta_\ell}$$

for such vectors. The extra class number N_c is retained by replacing the second vector by a sum of its twisted central translates, which are still integral and tested, whereas the line of Ξ is fixed under this central action; this is again a *Formal-source input*. Apply (31.27) at the two embeddings giving the two factors of the rank-one projection pairing. The lower bound for their product and the projection upper bound give

$$\left| \iota \left(Q \overline{Q} \, \frac{\rho_{p_2}(\vartheta, 1)^2}{N_c^3 \|\Xi_{p,\vartheta}\|^2} \right) \right|_\ell \le e^{\beta_\ell + 2F_\ell \log \ell} .$$

The exponent 3 of N_c is obtained from the two factors of N_c^2 and the single factor of N_c in the strengthened projection denominator. Since $|\sigma(Q\bar{Q})| \le e^{2B_0}$, summing the finite-place bound gives

$$\mathrm{Ht}_{\le 3B_0 + 2\sum_\ell F_\ell \log \ell} \left(\frac{\rho_{p_2}(\vartheta, 1)^2}{N_c^3 \|\Xi_{p,\vartheta}\|^2} \right) .$$

Multiply by the estimate (31.26) and by the square of the inverse ratio of the ramified flat pair p_2 (height at most $2x^{999}$ from fit). Using (31.25), the product is $\rho_p(\vartheta, 1)^{-1} / (N_c^3 N(Q)^{2z-1})$. The integer factor N_c^3 costs nothing when multiplying back (its local absolute value is at most one), and the conductor power is a local unit. The saving is exactly minus the cost (31.1). Since $x^{999} \le B_0$ by the admissible magnitudes, we conclude

$$\mathrm{Ht}_{\le 6B_0 - \mathcal{U}}(\rho_p(\theta_\iota^j, 1)^{-1}). \tag{31.28}$$

This is `heightLe_inv_ratio`.

31.11 Combining the special values and closing the lower bound

We now have precisely the estimate for the long string in (31.16). For p_0 consider lower points z with $7m/2 < z \le e_0$. There are $k/2$ of them, all of rank-one type at j; this follows

from the specified weight of ν_0 and the numerical relations between r, k, t, m (the range verification occurs in `heightLe_inv_heckeValueProd_div`). Thus these $k/2$ inverse ratios have height at most $6B_0 - \mathcal{U}$ each, by (31.28). The remaining factors in the three products have $7m/2 + e_+ + e_- = 5m$ lower points in all; their underlying pairs are ramified at exactly M and their inverse ratios are bounded by the fit hypothesis with cost x^{999} each. All ratios enter squared, and the conductor powers in (31.8) are local units. Consequently

$$\mathrm{Ht}_{\le 10mx^{999}+k(6B_0-\mathcal{U})}\left(\frac{\Gamma_j(1)}{\Gamma_j(\theta_\iota^j)}\right). \tag{31.29}$$

This proves the deduction in `heightLe_inv_heckeValueProd_div` (`New/Rows/HeckeValueHeight.lean`). One should notice the count $k/2$ before squaring; counting k factors before the square would yield an incorrect leading coefficient.

Combine (31.29), the compressed determinant estimate (31.24), and identity (31.20). The total height of the one-lift inverse volume ratio is at most

$$3nrB_0 + n\bigl(10mx^{999} + k(6B_0 - \mathcal{U})\bigr) \le nk((\log b)B_0 - \mathcal{U}).$$

The scale inequality on the right is the one in `heightLe_liftVolume_ratio` (`New/Rows/LiftVolumeHeight.lean`): r/k is bounded, $mx^{999}/(kB_0)$ negligible and $\log b$ very large. Thus all the positive errors are absorbed by the $k(\log b)B_0$ allowance. Multiplying over $j = 1, \dots, m$ gives height bound $n(L - H)$ for the product of the ratios.

By Proposition 31.16 the full-chain ratio squared is the product of the squared single-lift ratios multiplied by the nth power of the inverse squared exchange factor. Its cost is m^2 squares, so its nth power has height at most $2nm^2tx^{999} \le 2nL$. We obtain

Proposition 31.21 (Reverse ratio)**.** *For every fit family,*

$$\mathrm{Ht}_{\le 2n(2L-H)}\left(\frac{Z(1)^2}{Z(\theta_\iota)^2}\right). \tag{31.30}$$

This is `Data.heightLe_volume_ratio_sq` (`New/Endgame/LongForms.lean`). Finally choose a fit family, apply Proposition 31.12 and square its ratio. The product of its ratio squared and that of (31.30) is identically one by injectivity. Lemma 31.4 gives

$$0 \le 2(V + 2d_{\mathrm{ind}} + nb) + 2n(2L - H).$$

As $b \le L$, rearrangement yields $n(H - 3L) \le V + 2d_{\mathrm{ind}}$, proving Theorem 31.1 from the explicitly identified inputs.

The arithmetic content of this argument is not an estimate of the size of individual L-values. It consists of finding comparison vectors for all good forms, extracting the rank-one ratios from their common periods, recovering adequate local integral tests from the geometrically constructed theta coefficients, and ensuring that projections of the tests have small determinant denominator. The same prime-power sum that appears in the divisibility cost in Theorem A is recovered, with precisely the opposite sign, in (31.28). The remaining issue, necessary for both occurrences of the height argument, is the existence of the simultaneous fit family. This calls for the congruence and local cohomological arguments in the next section.

Chapter 32

Period identities, finite matching, and the geometry of the lower bound

The proof of Theorem C in Chapter 31 uses period identities in two quite different ways. One transports an integral structure to the whole row space by comparing periods of complementary rank; the other computes the metric on a chain by comparing the pairing of consecutive theta lifts with a product of simpler pairings. The local character at the ramified places matters in both operations. We therefore develop the algebra of the residue matrix functions first, then explain the global period calculation and its relation to the integral tests. In particular, the matching identity for finite functions, the Fourier transform of the point data, and the support condition in a Rallis pairing can be analyzed without assuming any new nonvanishing theorem for automorphic forms. We retain the exact notation of Chapter 31 for the global variables, but use lower case a_0, b_0, n_0 for temporary matrix sizes. Statements marked *formal-source input* identify the assertions whose independently written proof is not completed in this monograph; the deductions from those assertions are provided explicitly.

32.1 Complementary frames and matched data

Let R be a finite product of quotients of discrete valuation rings by powers of the maximal ideal. For an N-dimensional free module, let $N = a_0 + b_0$, and choose an ordered splitting of the coordinates into b_0 and a_0. Write

$$k = [Y\ Z] \in \mathrm{GL}_N(R), \qquad k^{-1} = \begin{bmatrix} T \\ X \end{bmatrix},$$

so that $Y \in \mathrm{Mat}_{N \times b_0}(R)$, $X \in \mathrm{Mat}_{a_0 \times N}(R)$. Then

$$XY = 0, \quad TY = I_{b_0}, \quad XZ = I_{a_0}, \quad YT + ZX = I_N. \tag{32.1}$$

In particular Y and X belong to a split exact sequence $0 \to R^{b_0} \xrightarrow{Y} R^N \xrightarrow{X} R^{a_0} \to 0$. In the following such a pair is called compatible. This is a precise condition over a non-field: an injective matrix over the residue ring need not by itself define a direct summand. A matrix on which a left inverse exists does define a direct summand, as needed here.

Lemma 32.1 (Complementary row and column bases)**.** *An N by b_0 matrix Y with a left inverse extends to $[Y\ Z] \in \mathrm{GL}_N(R)$. Dually a surjection $X : R^N \to R^{a_0}$ extends to the*

last rows of an inverse matrix. Every compatible pair (X, Y) occurs as the corresponding submatrices of k and its inverse, and the determinant of such a k is determined by the pair. The choices of k giving a fixed pair form a torsor for the additive group $\mathrm{Mat}_{b_0 \times a_0}(R)$.

Proof. It suffices to treat one local factor of R, since a basis on each local factor gives a basis over the product (with the indicated ranks). A left inverse reduces modulo the maximal ideal to a left inverse, so the reduced columns are linearly independent over the residue field. Extend these columns to a basis of the residue vector space and lift the added columns: the determinant of the extended square matrix is a unit. The dual assertion follows by transposition. For a split exact sequence choose any section Z of X. The map $[Y\ Z]$ is an isomorphism by exactness: for any u, $u - ZXu$ is in the image of Y, with unique preimage. If Z' is another section, then $Z' - Z = YU$ for unique U. Thus

$$[Y\ Z'] = [Y\ Z] \begin{bmatrix} I & U \\ 0 & I \end{bmatrix},$$

proving all claims, including the assertion on determinants. □

This supplies the finite-ring instance of `IsValuationPair.exists_firstCols_iff` in `Lib/General/LinearAlgebra/PartnerAlgebra.lean`; the source has a more general valuation-pair formulation. For fixed Y, the compatible X's are a torsor under $\mathrm{GL}_{a_0}(R)$ acting by left multiplication; for fixed X, the compatible Y's form a torsor under $\mathrm{GL}_{b_0}(R)$ acting by right multiplication. These assertions follow either from (32.1) or from the uniqueness of an isomorphism of free modules inducing a given basis of the kernel or quotient.

Let $\chi : R^\times \to \mathbb{C}^\times$ be a character extended by zero off the units. In the text this extension is meant whenever $\chi(\det M)$ occurs for a square matrix M that is not assumed invertible; its inverse character χ^{-1} is also extended by zero. A *matched pair* (Φ, Ψ) of functions on row matrices X and column matrices Y has the following three properties: each function is zero off the frames that can occur in Lemma 32.1, and for every compatible pair furnished by k,

$$\Phi(X) = \chi(\det k)\Psi(Y). \tag{32.2}$$

This is `IsPartner` in the above file. In particular $\Phi(fX) = \chi(\det f)^{-1}\Phi(X)$ and $\Psi(Yq) = \chi(\det q)^{-1}\Psi(Y)$ for invertible f, q. It is important that both functions have inverse character covariance: indeed the compatible pair (fX, Yq) is given by $k\,\mathrm{diag}(q, f^{-1})$, whose determinant is $\det k \cdot \det q / \det f$.

Proposition 32.2 (Open-cell formula and all matched pairs)**.** *For $X = [X_1\ X_2]$, $Y = \binom{Y_1}{Y_2}$ in the coordinate splitting $R^{b_0} \oplus R^{a_0}$, let p be any function on a_0 by b_0 matrices. The functions*

$$\Phi(X) = \chi(\det X_2)^{-1} p(X_2^{-1} X_1), \qquad \Psi(Y) = \chi(\det Y_1)^{-1} p(-Y_2 Y_1^{-1})$$

are partners, with the respective formula understood as zero when the square block is not invertible. More generally every inverse-character-covariant function Ψ supported on left-invertible column frames admits a unique partner Φ.

Proof. For compatible X, Y supplied by k, the identities $XY = 0$ and

$$\det X_2 \det k = \det Y_1 \tag{32.3}$$

follow from (32.1); to verify the determinant identity, multiply the square block matrix with first block row $[I\ 0]$ and second block row X by k. The product is block upper triangular with lower right block I, and upper left block Y_1. Thus X_2 is invertible exactly when Y_1 is.

In that case $X_2^{-1}X_1 = -Y_2Y_1^{-1}$, and (32.3) gives (32.2). The support assertion is immediate from the invertible chosen block. For general Ψ, define $\Phi(X)$ by choosing compatible Y, k and setting the right hand side of (32.2). Changing Y to Yq multiplies the determinant by $\det q$ and Ψ by the inverse character. Changing k for a fixed pair does not change its determinant by Lemma 32.1. This proves well-definedness, uniqueness, and the required covariance and support. □

The first assertion is the precise formula `partner_block` in `New/Exchange/BlockPartners.lean`. The second, elementary completion is useful in understanding why the intertwining identity on partners yields an explicit matrix kernel rather than an unspecified relation between periods. A simple example over a field is $a_0 = b_0 = 1$. Then every nonzero column $(y_1, y_2)^T$ with $y_1 \neq 0$ corresponds to the row $(x_1, x_2) = (-y_2, y_1)$, up to scaling, and the parameter $-y_2/y_1 = x_1/x_2$ is the coordinate of the annihilator line on the same affine chart of a projective line. Since $\det k$ may be a nontrivial unit, the inverse-character factors cannot be dropped even in this example.

For a compatible pair let

$$\Delta_\chi(X, Y) = \frac{\chi(\det k)}{|\operatorname{GL}_{b_0}(R)|}, \tag{32.4}$$

and put it equal to zero otherwise. The determinant is well-defined by Lemma 32.1. Given invertible $g \in \operatorname{GL}_N(R)$, $f \in \operatorname{GL}_{a_0}(R)$, $q \in \operatorname{GL}_{b_0}(R)$, the new pair $(f^{-1}Xg, g^{-1}Yq)$ is obtained from $g^{-1}k \operatorname{diag}(q, f)$, proving

$$\Delta_\chi(X, Y) = \chi(\det g/(\det f \det q))\Delta_\chi(f^{-1}Xg, g^{-1}Yq). \tag{32.5}$$

For a covariant Ψ as in Proposition 32.2, summation of (32.4) against Ψ on column frames for a fixed X gives exactly its partner Φ. Thus the denominator in (32.4) is the orbit size, not a multiplicative loss in the integrality estimate.

Lemma 32.3 (Recovering a period point)**.** *Suppose L, L' are functions of row and column matrices, respectively, transforming by χ (not χ^{-1}) under their frame groups, and satisfying*

$$\sum_Y \Psi(Y)L'(Y) = c\sum_X \Phi(X)L(X) \tag{32.6}$$

for all partners. Then for each left-invertible Y,

$$L'(Y) = c\sum_X \Delta_\chi(X, Y)L(X).$$

There is a normalization $d(Y)$ of the kernel with the properties that $d(Y)\Delta_\chi(X, Y)$ is either zero or a root of unity, and its average under changing the column frame is the original kernel in the sense

$$\frac{1}{|\operatorname{GL}_{b_0}(R)|}\sum_q d(Yq)\Delta_\chi(X, Y) = \Delta_\chi(X, Y).$$

Proof. Let Y be fixed. Take Ψ supported on the right frame orbit of Y, with $\Psi(Yq) = \chi(\det q)^{-1}/|\operatorname{GL}_{b_0}(R)|$. The covariance makes the left side of (32.6) equal to $L'(Y)$. Its partner is $\Delta_\chi(\cdot, Y)$ by the orbit calculation above, proving the formula. For each orbit choose a single representative and let $d(Y)$ equal the group order at the representative and zero elsewhere. The action on each left-invertible frame is free, so the indicated average is one. On the support of a compatible pair, multiplying the kernel by d either annihilates it or leaves the character of a determinant, hence a root of unity. □

Here the replacement of the original kernel by the normalized kernel does *not* preserve the equality pointwise for a fixed Y outside the selected representative. Instead, one uses an equivariant period functional in the square argument, which allows the displayed averaging. This is exactly the role of the function d in `exists_partner_kernel`, `New/Exchange/SquareLawFourier.lean`. There is also a support condition in that lemma. Write $Y = \binom{Y_1}{Y_2}$ and $X = [X_1\ X_2]$. If Y_1 has a left inverse L_1, then X_2 has a left inverse. Indeed set $P = I - Y L_1 \operatorname{pr}_1$ and choose a right inverse Z of X. We have $PY = 0$ and $PZX = P$ by (32.1). Since $P\binom{0}{v} = \binom{0}{v}$, the matrix $\operatorname{pr}_2 PZ$ is a left inverse of X_2. The assertion with the indices reversed is identical. This verification is necessary for propagating tidiness to the next edge of the grid. It explains why over a residue ring the proof uses the existence of a left inverse, not just a rank inequality at one embedding.

These constructions also have an integral local version. Let F be a local field with ring of integers O, or a finite product of such fields, and $Y : F^{b_0} \to F^N$ injective. The module $\{z : Yz \in O^N\}$ is an O-lattice of rank b_0, and its image under Y is saturated in O^N. After choosing a basis for this lattice by a matrix $g \in \mathrm{GL}_{b_0}(F)$, the matrix Yg is primitive (its entries are integral and its columns extend to an integral basis). The right coset $g\,\mathrm{GL}_{b_0}(O)$ is uniquely determined; this follows because the image of O^{b_0} must be precisely that lattice. Consequently, in a sum over such cosets only one term can survive when the function is supported on primitive frames. This establishes the normalization used in `localSection_of_primitive` and clarifies the relation between matching over the local field and matching over the residue ring. The full algebraic extension of local-section functionals from primitive data to all locally constant compactly supported functions, when needed without finite support of the coset sum, is retained as a formal-source input `exists_common_coset_reps` and `matchingPairs.exists_linear_factor` in `New/Exchange/BlockPartners.lean`. Neither is used to replace a divergent sum by an unqualified analytic integral.

32.2 Fourier transformations of the matching identity

A finite ring R as above admits primitive additive characters: a character $\psi : (R, +) \to \mathbb{C}^\times$ is called primitive when its kernel contains no nonzero ideal. A multiplicative character is primitive in the stronger sense used here if, for every $x \neq 0$, there exists u such that $1 + ux$ is a unit and $\chi(1 + ux) \neq 1$. At squarefree level this amounts to taking a nontrivial multiplicative character on each residue-field factor. Indeed if $x \neq 0$ at one factor, one can prescribe $1 + ux$ there to be an arbitrary unit and take it to be 1 at all other factors. At higher conductor powers these hypotheses must be checked separately. In the compact code the primitive multiplicative condition at squarefree level follows from `isPrimitiveMulChar_of_isTameFor`, `New/Exchange/CostOfSquares.lean`; it is used only in the square-free triangle, not all the double-grid transport at level M^2Q.

For rectangular matrices A define the unnormalized finite Fourier transform by

$$(\mathcal{F}_\psi p)(A) = \sum_B p(B)\psi(\operatorname{tr}(AB)),$$

where B has the transpose shape to A. For a primitive character and a fixed nonzero matrix C the function $B \mapsto \psi(\operatorname{tr}(CB))$ is nontrivial by varying the entry of B paired with any nonzero entry of C. The sum of a nontrivial character over a finite group is zero (translate the sum by a point on which the character is not one). Thus

$$\sum_B \psi(\operatorname{tr}(CB)) = \begin{cases} |R|^{uv} & C = 0, \\ 0 & C \neq 0, \end{cases} \tag{32.7}$$

for C of size u by v. Expanding the double sum proves

$$\mathcal{F}_\psi^2 p(C) = |R|^{uv} p(-C), \qquad \sum_A |\mathcal{F}_\psi p(A)|^2 = |R|^{uv} \sum_B |p(B)|^2.$$

These are `matrixFourier_matrixFourier` and the finite-character orthogonality used in `Lib/Weil/GaussSum/LocalHomUnit.lean`. The present normalization, with no factor $|R|^{-uv/2}$, is essential in the following character Gauss sum.

Proposition 32.4 (Matrix determinant Gauss sum)**.** *Let d be a positive integer and let F be a function on d by d matrices with left covariance $F(gB) = \chi(\det g)F(B)$ for all $g \in \mathrm{GL}_d(R)$. For primitive ψ and χ put*

$$G_d(\chi, \psi) = \sum_{B \in \mathrm{Mat}_d(R)} \chi(\det B)\psi(\operatorname{tr} B).$$

Then $|G_d(\chi, \psi)|^2 = |R|^{d^2}$ and

$$\mathcal{F}_\psi F(A) = \chi(\det A)^{-1} G_d(\chi, \psi) F(I_d), \tag{32.8}$$

where the right hand side is zero for singular A.

Proof. A noninvertible square matrix B over the finite ring has a nonzero left kernel vector v^T: indeed its transpose as an endomorphism of a finite free module is not bijective, hence not injective. Let $v_i \neq 0$. By primitivity choose u with $1 + uv_i$ a unit and $\chi(1 + uv_i) \neq 1$. The matrix $g = I + ue_i v^T$ satisfies $gB = B$ and has determinant $1 + uv_i$, by the rank-one determinant identity. Covariance forces $F(B) = 0$. Thus $F(B) = \chi(\det B)F(I)$ for every B. For invertible A, the change of variable $C = BA$ in the Fourier transform, and cyclicity of trace, give (32.8). For singular A, there is a nonzero right kernel vector w; choose i, u as before and $g = I + uwe_i^T$. Since $Ag = A$, the change of variable $B \mapsto gB$ and covariance give $\mathcal{F}_\psi F(A) = \chi(\det g)\mathcal{F}_\psi F(A)$, again zero. Finally apply Plancherel to the function $B \mapsto \chi(\det B)$. Its squared norm is $|\mathrm{GL}_d(R)|$. The transform has squared norm $|G_d|^2 |\mathrm{GL}_d(R)|$ by the result just proved. Canceling the positive order of the group proves the magnitude identity. Equivalently,

$$\overline{G_d(\chi, \psi)}\, \mathcal{F}_\psi F(I) = |R|^{d^2} F(I).$$

□

This proves `matrix_gauss_sum` of `Lib/Weil/GaussSum/LocalHomUnit.lean`. Note that no identity involving the phase of the Gauss sum was needed. The distinction between magnitude and phase is crucial for the global scalar: phases may depend on the choice of character, whereas the square magnitude is a pure cardinality factor, of zero height cost at primes outside the twist conductor. The rank-one case is the usual Gauss sum of a primitive multiplicative character on a finite field. The matrix statement is equally true at a product of higher powers of finite residue rings under the stated primitivity condition; there is no hidden estimate by a power of the residue cardinality.

We next give the two exact finite summation identities behind the weighted square law. Take an ambient matrix coordinate decomposition of sizes (u, a_0, d), with the small side equal to a_0 and large side $u + d$. For $Y \in \mathrm{Mat}_{u \times a_0}(R)$, $V \in \mathrm{Mat}_{u \times d}(R)$, form the complementary frames

$$Y_* = \begin{pmatrix} Y \\ I_{a_0} \\ 0 \end{pmatrix}, \qquad X_* = \begin{pmatrix} I_u & -Y & V \\ 0 & 0 & I_d \end{pmatrix}. \tag{32.9}$$

Use the frame placement `swapFirstTwo` of the source: the small-side columns are the middle block, and the other two row blocks of the inverse are ordered as u, d. In this placement the endomorphism $k = \begin{pmatrix} I & Y & -V \\ 0 & I & 0 \\ 0 & 0 & I \end{pmatrix}$ has the prescribed middle columns, and its inverse has the indicated first and third row blocks. Its determinant is one; this verifies the relative sign in the chosen ordering. Therefore by the orbit computation in Lemma 32.3,

$$|\operatorname{GL}_{a_0}(R)|L'(Y_*) = c|\operatorname{GL}_{u+d}(R)|L(X_*). \tag{32.10}$$

For example, choose a column datum supported on the orbit of Y_* with value 1 there, then its partner has the inverse-character values on the orbit of X_*; all the contributions cancel their determinant characters in the corresponding periods. The right side of (32.10) is independent of V, because a shear on the row frame of determinant one eliminates the upper right block V.

Given p on u by a_0 matrices define a new datum on $(u+d)$ by (a_0+d) matrices by

$$p^+\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{cases} p(-A)/|R|^{ud} & C = 0,\ D = I_d, \\ 0 & \text{otherwise.} \end{cases}$$

Multiplying (32.10) by $p(Y)$, summing over Y and averaging over V proves

$$|\operatorname{GL}_{a_0}(R)|\sum_Y p(Y)L'\begin{pmatrix} Y \\ I_{a_0} \\ 0 \end{pmatrix} = c|\operatorname{GL}_{u+d}(R)|\sum_X p^+(X)L\begin{pmatrix} I_u & X_{11} & X_{12} \\ 0 & X_{21} & X_{22} \end{pmatrix}. \tag{32.11}$$

This proves `SquareLaw.sum_eq_nextDatum` in `New/Exchange/SquareLawFourier.lean`, with the basis reorderings made explicit. The matrix X_{11} has size u by a_0, etc. Formula (32.11) remains valid without primitivity of χ, because it uses only covariance and the matched-pair relation.

Proposition 32.5 (Fourier version of the finite square)**.** *Assume $a_0 = u + d$ and that ψ, χ are primitive. Given a function p on matrices of shape $(u+d)$ by u, there is a scalar γ, independent of p, c, L, L', with $|\gamma|^2|R|^{d(d+2u)} = 1$, for which*

$$\begin{aligned} &\sum_{Y\in \mathrm{Mat}_{u\times a_0}(R)} \mathcal{F}_\psi(\bar p)(Y)L'\begin{pmatrix} Y \\ I_{a_0} \\ 0 \end{pmatrix} \\ &\quad = \gamma c \sum_{X\in \mathrm{Mat}_{(u+d)\times(a_0+d)}(R)} \mathcal{F}_\psi(\overline{p^+})(X)L\begin{pmatrix} I_u & X_{11} & X_{12} \\ 0 & X_{21} & X_{22} \end{pmatrix}. \end{aligned}$$

Here p^+ is the extension just defined, with the row and column sizes of p (transposed from those of Y).

Proof. Put $F = \bar p$. Fourier-transform $\overline{p^+}$ at a matrix $X = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$; the summation variable has shape (a_0+d) by $(u+d)$. It is restricted to matrices $\begin{pmatrix} S & T \\ 0 & I_d \end{pmatrix}$ with weight $F(-S)/|R|^{a_0 d}$. The trace pairing gives

$$\operatorname{tr}\left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}\begin{pmatrix} S & T \\ 0 & I_d \end{pmatrix}\right) = \operatorname{tr}(AS) + \operatorname{tr}(CT) + \operatorname{tr} D.$$

By (32.7) the sum over T vanishes unless $C = 0$, when it cancels the factor $|R|^{-a_0 d}$. Hence

$$\mathcal{F}_\psi(\overline{p^+})(X) = \mathbf{1}_{C=0}\,\psi(\operatorname{tr} D)\,\mathcal{F}_\psi F(-A).$$

For fixed A, B, the function $D \mapsto L\begin{pmatrix} I & A & B \\ 0 & 0 & D \end{pmatrix}$ transforms by the determinant character under left multiplication of D; indeed left multiplication by $\operatorname{diag}(I, g)$ on the full frame has exactly this action. Proposition 32.4 gives the sum over D, with additive character $\psi(\operatorname{tr} D)$, as $G_d(\chi, \psi)$ times its value at the identity. That value is independent of B by the determinant-one shear already used in (32.10). The sum over B therefore supplies $|R|^{ud}$. The right hand sum in the proposition, before multiplication by γc, is precisely

$$|R|^{ud} G_d(\chi, \psi) \sum_A \mathcal{F}_\psi F(-A) L \begin{pmatrix} I & A & 0 \\ 0 & 0 & I \end{pmatrix}.$$

Apply (32.11) with p there equal to $\mathcal{F}_\psi F$; the two group orders cancel since $a_0 = u + d$. Averaging over the free upper right block in the extension merely gives the last sum. The desired scalar is

$$\gamma = \frac{1}{|R|^{ud} G_d(\chi, \psi)} = \frac{\overline{G_d(\chi, \psi)}}{|R|^{(u+d)d}},$$

which has the asserted squared magnitude by Proposition 32.4. □

This proves the identity `SquareLaw.sum_fourier_eq_nextDatum` of `New/Exchange/SquareLawFourier.lean`. It is considerably stronger than a triangle inequality on Fourier coefficients: the determinant Gauss sum kills singular lower blocks and gives the exact scalar. In the triangle of rank increments t, $d = t$, and u is the rank of the smaller common block; hence the exponent $t(t + 2u)$ is the difference between the products of the adjacent dimensions that appears when two kernels are reversed. The normalization of the semilinear operation of reversal in `exists_conj_modKernel_eq_swap` (`Lib/SiegelWeil/RallisInner/SharpData.lean`) gives a factor of squared modulus $|R|^{-n_1 n_2}$ for a single reversed edge of ranks n_1, n_2. That identity of genuine Weil kernels is retained as a formal-source input, but the Fourier calculation proves why the quotient of these factors in the square is a pure cardinality power. For $R = \mathbb{F}_q$, the exponent of q in the factor of the preceding proposition is exactly $-d(d+2u)$ in squared magnitude, including off-diagonal blocks. Omitting the Fourier factor at off-diagonal rectangles would change the relative metric in the main determinant calculation.

32.3 The graph term in doubling

To identify the analytic input in the Gram factors of Proposition 31.8, write S^- for the Hermitian space with form opposite to that of S, of dimension r, and T for a target of dimension $n > 2r$. The doubled source is the split Hermitian space $S \oplus S^-$. Its rational Lagrangians are the indices in the Siegel–Weil sum. They are not all graphs. Dropping the nongraph terms without explanation would be a serious omission: the finite support of the datum is essential to this simplification.

Lemma 32.6 (Graphs and the rank test)**.** *Let J be the matrix of a nonsingular Hermitian form on $V = E^r$, and let $L \subset V \oplus V^-$ be a Lagrangian. Represent it by the full-column-rank matrix $\binom{K}{K'}$. The equation of isotropy is $K^* J K = K'^* J K'$. If K' is invertible, then L is the graph of a unique isometry of V, namely $K K'^{-1}$. If V is definite at an archimedean place, every L is of this type.*

Proof. The determinant of K^*JK equals that of K'^*JK' and is nonzero if K' is invertible; hence K is invertible. With $\delta = KK'^{-1}$ the isotropy equation gives $\delta^*J\delta = J$, and the vector in L with prescribed second component v must be $(\delta v, v)$. For the last claim, if $K'c = 0$ for some nonzero c, isotropy implies $(Kc)^*J(Kc) = 0$. Definiteness implies $Kc = 0$, contradicting the full rank of $\binom{K}{K'}$. Uniqueness follows from the prescribed second projection. □

The Lean graph convention also permits a minus sign in the second component, which has the effect of replacing δ by $-\delta$. There is no second copy of the set of graphs caused by this convention. In indefinite signature the last argument has no force; for example an isotropic vector can occur with second component zero. For the complementary-rank exchange the following more general rank test is needed.

Formal-source input (zero terms). In `Lib/SiegelWeil/RallisInner/SharpData.lean`, `IsClean` is the alternative of definiteness or a finite named datum supported on matrices admitting a left inverse at a chosen split place. The theorem `exists_graph_eq_of_zeroTerm_ne_zero` states that for a clean datum a nonzero zero term of the double datum is attached to a graph. Its essential new assertion, beyond Lemma 32.6, is the vanishing when $K'c = 0$ and $Kc \neq 0$: invariance of the zero term under the corresponding Heisenberg translations enforces a Fourier support constraint incompatible with the primitive support at the chosen place. This analytic invariance and its compatibility with naming a finite datum are retained as formal-source input. The algebraic consequence is now identical to the proof of the lemma: K' is invertible, and the graph is unique. The same module's `zeroTerm_graphSplitting` identifies the surviving term, for datum f' on the first copy and f on the second:

$$\mathrm{ev}_{L_\delta}(\mathrm{double}(f', f)) = \langle f', \omega(\delta^{-1}, 1)f\rangle.$$

Here the double uses conjugation on the opposite source, as in that declaration. This specifies the metric normalization of this formula, not merely a proportionality.

Formal-source input (Siegel–Weil). The identity used with these graph terms is precisely `siegel_weil` of `Lib/SiegelWeil/Formula/SiegelWeilFormula.lean`. In the split source of rank $2r$ and with $n > 2r$, its mean over the target quotient is

$$\frac{1}{\mathrm{vol}[U(T)]}\int_{[U(T)]}\theta_\Phi(h)\,dh = \sum_L \mathrm{ev}_L(\Phi),$$

with summable zero terms. The absolute bounds needed also for variable Φ are developed in `Lib/SiegelWeil/Formula/SiegelSets.lean` and `Lib/SiegelWeil/Formula/ZeroTermHeight.lean`, including `exists_summable_zeroTerm_bound` and `theta_normalisedTorus_le`. The latter is a bound along a normalized expanding torus by an integrable function. The majorants, the independence of the evaluation from completing a Lagrangian to a splitting, and the formula itself are not reproved here.

Proposition 32.7 (Unfolding, conditional on the analytic inputs)**.** *Use Haar measures as in the preceding mean identity, and let $G = U(S)(\mathbb{A})$, $\Gamma = U(S)(E^+)$, with right-regular action $R(g)y(x) = y(xg)$. For rapidly decreasing automorphic functions y, y' and a clean f, the graph-term identity and Siegel–Weil identity give*

$$\langle \Theta_{f'}y', \Theta_f y\rangle = \mathrm{vol}[U(T)]\int_G \langle f', \omega(g,1)f\rangle\langle y', R(g)y\rangle\,dg$$

provided the integrals and sums may be interchanged absolutely.

Proof. Choose compatible measurable fundamental domains for the source quotient in each of the two lift integrals. Multiply the conjugate of the first theta lift and the second theta lift, and integrate over the target quotient. The inner target integral is the split double theta integral. Apply the mean formula, and discard nongraph terms by the clean-data criterion. Index the graphs by $\delta \in \Gamma$; the graph term for representatives x, x' of the two source variables is the matrix coefficient at $x'^{-1}\delta x$, after inversion of the indexing isometry if needed. Thus the result, divided by $\mathrm{vol}[U(T)]$, is

$$\int_{\Gamma\backslash G}\int_{\Gamma\backslash G} \overline{y'(x')}y(x)\sum_{\delta\in\Gamma} M(x'^{-1}\delta x)\,dx\,dx', \qquad M(g)=\langle f', \omega(g,1)f\rangle.$$

Left invariance of y under Γ identifies $\sum_\delta \int_{\Gamma\backslash G}$ with an integral over G. Substitute $x = x'g$ therein, using left Haar invariance, and interchange these last two integrals. The integral in x' is $\langle y', R(g)y\rangle$, as required. □

The convergence condition is not optional. In `SharpData.lean`, `inner_product_formula` establishes the integrability of exactly this coefficient product and proves this identity, with rank hypothesis $n > 2r$ and clean f; its proof uses the rational norm bound `exists_rat_norm_le_height` for the pair kernel, the rapid bounds on y, y', and the zero-term summability in the Siegel–Weil formula. We take those analytic estimates and all measure identifications as formal-source input. The unfolding argument above isolates the part that is independent of the estimates. It also explains why the final integral is over the *group* of the source, not the source quotient. This distinction will reappear as a sum over infinitely many cosets in the Euler expansion; replacing it by the finite automorphic class set would change the formula.

32.4 Sharp support, level factors, and the common scalar

Write $R = \mathcal{O}_E/\mathfrak{m}$ for a finite ring assembled from the selected split places, with level one at each prime, and let

$$Z = \mathrm{Mat}_{n,r}(R), \qquad F : Z \longrightarrow \mathbb{C}, qquad \mathrm{supp}\, F \subseteq \{Y : LY = I_r\},$$

where L is fixed of size $r \times n$. This is a useful particular case of the primitive support of the preceding subsection. Our local vacuum conventions are those of the named finite datum in the compact source.

Lemma 32.8 (Support under column action)**.** *If $A \in \mathrm{GL}_r(R)$ and $F(Y)F(YA) \neq 0$ for any Y, then $A = I$. Over a local DVR with fraction field K, if Y and YA are both primitive integral n-by-r matrices for $A \in \mathrm{GL}_r(K)$, then $A \in \mathrm{GL}_r(\mathcal{O}_K)$.*

Proof. In the first case $LY = LYA = I_r$, so $A = I_r$. In the second case there are integral left inverses P, P' of Y, YA, respectively. Then $A = P(YA)$ is integral, and $A^{-1} = P'Y$ is integral too. □

Consequently the overlap of two such compact finite data in the matrix coefficient for the source vanishes outside the principal level subgroup at the selected places. On that subgroup the right translation and determinant twists reduce to the identity. Counting using the normalized Haar measure on integral matrices gives

$$\|f\|_{\mathrm{fin}}^2 = \frac{\sum_{Y\in Z}|F(Y)|^2}{|R|^{nr}}, \qquad c = \frac{\|f\|_{\mathrm{fin}}^2}{|\mathrm{GL}_r(R)|}.$$

Indeed, the reduction of the integral matrix lattice has equally measured fibers, whereas the principal kernel in the integral group has measure the reciprocal of the order of the reduction group when the maximal compact group has measure one. There are two independent denominators. They have quite different origins, and conflating them removes a large dimension-dependent factor.

Formal-source input (sharpness and Euler expansion). These finite arguments are the support and counting core of `isSharp_modDatum` in `Lib/SiegelWeil/RallisInner/SharpData.lean`; the named model action, the Hilbert norm identity for that model, and the passage from split residue factors to genuine Weil coefficients are taken from that declaration. Its hypothesis includes definiteness or nonempty selected places, so that clean support is available. The exact coset expansion and injectivity needed in the application are `rallis_inner_product_euler` in `Lib/Weil/CompactLift/RallisFormula.lean`: for a fixed real edge K and $n > 2r$ there is a common $A \neq 0$ such that for weighted characters a, b, sharp finite datum with parameter c, and a form y of level J, the lift $\mathcal{L}$ satisfies

$$\mathcal{L}y = 0 \implies y = 0, \qquad \langle \mathcal{L}y', \mathcal{L}y \rangle = \sum_{\gamma} A c \,\mathrm{cosetTerm}(n, b, S_0; y', y; \gamma).$$

The sum is a convergent `HasSum`. Here γ indexes cosets in the adelic group integral and S_0 denotes the selected bad places in the sharpness condition. In particular A does not depend on the finite sharp datum or the varying characters for that real edge. The injectivity assertion is not a consequence of the positive norm of the identity term alone: other coset terms can have phases of either sign. The nonzero pairing of the real edge and the integral nondegeneracy argument in this declaration are used as formal-source input.

A common source of confusion is that a finite matrix norm and a Petersson norm use different measures. The effect can be seen without opening the entire analytic proof. At the auxiliary edge of rank $r \longrightarrow r + t$, let $\mathcal{Q}$ be the selected twisting places and M the fixed matching places. Set

$$N(\mathcal{Q}) = \prod_{w \in \mathcal{Q}} q_w, \quad \gamma_{\mathcal{Q}} = \prod_{w \in \mathcal{Q}} |\mathrm{GL}_r(k_w)|, \quad R = \prod_{w \in M \cup \mathcal{Q}} k_w.$$

At M, the auxiliary indicator fixes a standard inclusion matrix; at $\mathcal{Q}$ it fixes the designated r rows to the identity, leaving tr free entries. The auxiliary datum multiplies this indicator by $\gamma_{\mathcal{Q}}$.

Proposition 32.9 (Auxiliary count)**.** *With this datum F, the sharp parameter is*

$$c = \frac{\gamma_{\mathcal{Q}}^2 N(\mathcal{Q})^{tr}}{|\mathrm{GL}_r(R)|\,|R|^{(r+t)r}}.$$

Proof. For $w \in M$ there is one allowed matrix; for $w \in \mathcal{Q}$ there are q_w^{tr} matrices. Since the factors form a Cartesian product, the number of allowed data is $N(\mathcal{Q})^{tr}$. Every nonzero value of F equals $\gamma_{\mathcal{Q}}$. Apply the two denominators computed above. □

The finite identity is `AuxPair.levelIndicator_support_and_normSq` in `New/Exchange/CharacterLiftNorm.lean`. To compare algebraic Gram forms, multiply the coefficient by the level index $[K_0 : J']$ on the target. The remaining common real factor contains the cardinality of the Weyl index at that target, and is independent of the character; the precise

conversion is contained in that module's `Data.auxLift_injective_and_hasSum_inner`. The result, in the notation actually used later, is

$$d = [K_0 : J'] \frac{\gamma_{\mathcal{Q}}^2 N(\mathcal{Q})^{tr}}{|\mathrm{GL}_r(R)|},$$
$$R_j(\gamma) = \frac{\mathrm{cosetTerm}(r+t, b, S_0; x, y; \gamma)}{|R|^{(r+t)r}},$$

and the sum of $AdR_j(\gamma)$ is the auxiliary algebraic Gram pairing, for one $A \neq 0$ common across the twists. These are exactly `AuxPair.gramFactor` and `Data.rallisTerm`. In particular one cannot move the level index into the summand and then also multiply by it when converting from Petersson to algebraic norms.

There is a useful one-prime example. Let $r = 1$, $t = 3$, $R = \mathbb{F}_q$, and let the only selected place be a twisting place. The allowed column vectors are $(1, z_1, z_2, z_3)^{\mathrm{t}}$; there are q^3, with $F = q - 1$. Then

$$c = \frac{(q-1)^2 q^3}{(q-1)q^4} = (q-1)/q.$$

The finite norm alone is $(q-1)^2/q$, which differs by $q - 1$. This illustrates the independent GL-group factor. For a fixed matching place the corresponding free-entry count is absent, but both denominators remain. At higher rank the GL-group cardinality changes much faster with rank than one could infer from any rank-one formula.

32.5 A rank-one Euler factor test

The auxiliary count determines the dependence on the support of a finite datum, but does not explain the appearance of the desired Hecke L-value. For the latter, consider $r = 1$, $n = 2k$ and $k > 1$. At a split unramified prime of the real base, the local unitary group is identified with $E_w^\times$ and its compact-unit quotient is indexed by integers. Let q be the common residue size, let $a = q^{-k}$, and let λ have modulus one. Then

$$\sum_{j \in \mathbb{Z}} a^{|j|} \lambda^j = 1 + \frac{a\lambda}{1 - a\lambda} + \frac{a\lambda^{-1}}{1 - a\lambda^{-1}} = \frac{1 - a^2}{(1 - a\lambda)(1 - a\lambda^{-1})}.$$

The positive and negative valuations give the two Euler factors in the denominator. The numerator corrects for counting the valuation zero twice. At nonsplit places the unitary line group is compact and has a single coset modulo its maximal compact group. This simple check explains the split and nonsplit factors of the line calculation. The actual identification of its character factors, and the unramified vacuum coefficient equal to the required character times $q^{-k|j|}$ in this rank, is the formal-source input `exists_hasSum_vacuumCoeff_at_place` in `Lib/SiegelWeil/RallisInner/SharpData.lean`. The pair of factors $1 - \varepsilon_w q^{-k}$ at a split conjugate pair has opposite signs $\varepsilon_{\bar{w}} = -\varepsilon_w$ and thus product $1 - q^{-2k}$.

For completeness, in this absolutely convergent range Euler products with coefficients of norm at most one cannot vanish. If z_w has norm at most q_w^{-k}, the expansion $\log(1 - z_w) = -\sum_{l \geq 1} z_w^l / l$ has absolute sum bounded by $|z_w|/(1 - |z_w|)$. The sum of the latter over places converges: over each rational prime the number of places is bounded by the field degree and the residue size is at least the rational prime. Therefore exponentiating the sum of logarithms gives both a nonzero product and a convergent product of reciprocals. This argument does not apply at $k = 1$ without hypotheses and further analysis.

Formal-source input (the rank-one norm identity). In `New/Exchange/CharacterLiftNorm.lean`, `LValueData.inner_trivialLift_self_eq_L` proves, for $n = 2k$ with $k > 1$,

$$\langle f_\chi, f_\chi \rangle = C \prod_{w \in M} (q_w - 1)\ H_\chi\ d_\chi L^S(k, \mu^{-1}) N(\mathcal{Q})^{n-1}.$$

Here f_χ is the trivial-source auxiliary lift, H_χ is the hook-number factor for its Weyl lattice, d_χ the Weyl-index cardinality, S the omitted places, and μ the weighted character occurring in the declaration. The factor $C \neq 0$ is given there by the product of two quotient volumes divided by the real-zeta Euler product at k. This is a norm formula for the particular real and finite datum, not an identity for arbitrary harmonic vectors. The local coset calculation, the rank twist in the character and the comparison of the real Fock and algebraic Gram forms are the formal ingredients of that theorem. The finite count above and the elementary geometric-series calculation explain the shape of its conductor power and Euler factors. It is especially important that the displayed identity involves $L^S(k, \mu^{-1})$ with the character prescribed by the weighted lift, rather than its absolute square: the base change in the Euler factor computation has already paired the appropriate local factors.

32.6 Hyperbolic-plane removal and the signed product

We now pass from the absolutely convergent range of the Rallis calculation to the distinct argument used to compare theta integrals of complementary rank. Adding a hyperbolic plane to both source and target changes the source and target dimensions by two each. To understand the scalar in this step, introduce

$$P_\nu(e) = N(S)^{e^2} \prod_{j=1}^{e} |L^S(j, \nu)|^2 \quad (e \geq 0), \qquad \Pi_\nu(e) = \begin{cases} P_\nu(e) & e \geq 0, \\ P_\nu(-e)^{-1} & e < 0. \end{cases}$$

The second definition agrees with the source's `lValueProdInt` of `Lib/SiegelWeil/Tower/LargeSideConstantTerm.lean`. The positivity of the factors under the actual weight hypotheses is the result `lValueProdInt_pos`; its proof uses the weighted-character nonvanishing theorem for the involved values, not general nonvanishing on a hypothesized critical line. In particular we work here on a positive part of the theory and do not assume GRH to define the reciprocal.

Lemma 32.10 (The transition quotient)**.** *For $e \in \mathbb{Z}$, let $b = e + 1$ if $e \geq 0$, and $b = -e$ if $e < 0$. Then*

$$\Pi_\nu(e+1)/\Pi_\nu(e) = N(S)^{2b-1} |L^S(b, \nu)|^2.$$

Proof. For $e \geq 0$ divide the two positive products; the exponent increment is $(e+1)^2 - e^2 = 2e+1$. For $e < 0$, put $b = -e \geq 1$. The inverse product at $e+1$ is the inverse of the product at $b-1$, including $b = 1$. Dividing by the inverse at b gives $P_\nu(b)/P_\nu(b-1)$, which has exponent $b^2 - (b-1)^2 = 2b - 1$. □

In particular $\Pi_\nu(-1) = 1/(N(S)|L^S(1, \nu)|^2)$ and both transitions from -1 to 0 and from 0 to 1 have quotient $N(S)|L^S(1, \nu)|^2$. This gives a check on the signed-exponent convention at the equal-rank boundary. If one used the unsigned product at negative integers, the first transition would have reciprocal scalar, breaking the exchange relation. More generally the quotients telescope without any case-dependent residual factor.

Let $W' = \mathbb{H} \oplus W$ and $V' = \mathbb{H} \oplus V$ with base-adapted splittings. For $n = \dim W$ even and $r = \dim V$, let $e = r - n/2$. At the larger pair the index is $(r+2) - (n+2)/2 = e + 1$. The constant term in a line radical is governed by the isotropic cone of V'. Here is the elementary mechanism of the constant-term reduction.

Lemma 32.11 (Orthogonality on the line radical)**.** *Let A be a compact abelian group with normalized Haar measure. If $\eta : A \to \mathbb{C}^\times$ is a continuous unitary character, then its mean is one for the trivial character and zero otherwise.*

Proof. The trivial case is the mass of A. Otherwise choose a_0 with $\eta(a_0) \neq 1$ and translate the integral by a_0 to obtain $I = \eta(a_0)I$. □

In the adapted mixed model, the line radical acts on the theta summand indexed by a rational vector $x \in V'$ by the character attached to the scalar $\langle x, x\rangle$. Integrating over the compact quotient of the line radical kills every summand with $\langle x, x\rangle \neq 0$. The model action and the interchange with the theta summation are formal-source inputs `constTermAt_eq_integral` and `hasSum_constTermAt`, in `Lib/SiegelWeil/Tower/ConstantTerm.lean`. The surviving rational cone decomposes into the vertex and one nonzero orbit: by Witt extension, any two nonzero isotropic vectors are conjugate under the rational unitary group. Indeed, the map of their one-dimensional subspaces taking one vector to the other is an isometry for the restricted zero form, so Witt extension provides the isometry of the ambient nondegenerate space. The stabilizer of the isotropic line has Levi $E^\times \times U(V)$; its action on the chosen vector is through the first factor, and the stabilizer of the vector is the kernel of that scaling.

The exponent $r + 1$ in the measure from unfolding this orbit can be checked in a Witt basis for V'. The radical is a two-step group with quotient an E-vector space of dimension r and central coordinate in a one-dimensional skew-Hermitian E^+-space. Conjugation by the line-scaling element $\mathrm{diag}(a, 1, \bar{a}^{-1})$ multiplies the first block by a scalar of E-absolute value $|a|_E$ and the second by $a\bar{a}$. This has Jacobian $|a|_E^r |a|_E = |a|_E^{r+1}$ under compatible normalized absolute values. Unfolding the nonzero orbit, when the vertex contribution vanishes, therefore introduces a multiplicative line integral over ideles modulo $E^\times$, an integral over $[U(V)]$, and an integral over the maximal compact of the larger space. The source supplies this precise normalized unfolding, including integrability and the nonzero measure constant, in `exists_unfold_constant_term` of `Lib/SiegelWeil/Tower/ConstantTerm.lean`.

Formal-source input (the normalized one-plane identity). The full formula `exists_constant_term_thetaLarge`, in `Lib/SiegelWeil/Tower/LargeSideConstantTerm.lean`, chooses compatible rapid datum and local row datum, and compares the constant term on W', V' with the smaller integral on W, V evaluated on the real cone-integral datum. It gives a scalar $\gamma \neq 0$, for one common constant C, with

$$|\gamma|^2 \Pi_\nu(e) = C \Pi_\nu(e + 1).$$

The equality holds on the indicated Levi variables when the line-scaling variable is one, or the smaller ranks are equal. This proviso matters in using the formula; there is no assertion here of full arbitrary-variable equality merely from a constant term. The local integrations use the adapted frames and the line datum constructed by `exists_lineTerm_localDatum`; the character factors are those of the declaration, and the real cone integral is also prescribed there. In particular C is independent of the twist. By Lemma 32.10 this scalar has squared norm $CN(S)^{2b-1}|L^S(b, \nu)|^2$. The single $N(S)$-power accompanying $e = 0$ or $e = -1$ and the growth of the exponents thereafter arise from the full conductor and line Gauss-sum calculations, not a guess based on dimensions.

32.7 What makes an exchange an identity on all partners?

The preceding line calculation proves neither the complementary-rank identity nor its square version in isolation. There are two more requirements. One must arrange a nonzero common

starting ratio at equal rank and prescribe the admissible real datum after removing planes. One must also show that two theta integrals, whose difference has appropriate vanishing Fourier coefficients, agree on the full space of matched data. A function on a group can have zero constant term without being zero, even if it is automorphic.

For a space X of even rank n the input ratio is

$$\ell_\nu(v) = \frac{|\lambda_\nu(v)|^2}{\Pi_\nu(n/2)}, \qquad \lambda_\nu(v) = \theta_{\mathrm{large}}(X, X; \nu, v, F_{\mathrm{std}})(1).$$

It is `equalRankThetaRatio` of `New/Exchange/LowRankIdentity.lean`; F_{std} is the indicated standard-column datum. The theorem `exists_equalRankThetaRatio_value` of `New/Exchange/EqualRankValue.lean` constructs a rapid real datum on one hyperbolic enlargement of the input W of rank $2k + 2q$, with q its number of negative signs. For a character weight componentwise at least k, its equal-rank ratio is a common nonzero number ℓ, independent of ν. This is a formal-source input: it requires the nonvanishing for a suitable real datum and consistency of the local comparisons as planes are removed, not simply the positivity of Π_ν.

For the chosen matched rank pairs W, V and U, W, `exchange_of_partners` of that module takes rapid test data z, y, with z measured at the larger equal-rank pair, and imposes the signature equality of complementary ranks. Its conclusions are: there are a common constant C and a single rapid datum v, and for each split tame character a scalar c, such that

$$C|c|^2 = \Pi_\nu(\dim V - \tfrac{1}{2}\dim W)\ell_\nu(z), \tag{32.12}$$

and the two global theta integrals on partners agree after scaling by c. With the equal-rank datum chosen above, the right-hand side is positive, so C and c cannot vanish. Dividing (32.12) for two twists therefore cancels all real constants:

$$\frac{|c_\nu|^2}{|c_{\nu_0}|^2} = \frac{\Pi_\nu(\dim V - \frac{1}{2}\dim W)}{\Pi_{\nu_0}(\dim V - \frac{1}{2}\dim W)}.$$

This is the precise scalar feature of the complementary-rank exchange in Proposition 31.8.

Formal-source inputs in the passage to matched data. For the real data, the cone integration operator must reach prescribed determinant vectors. The needed surjectivity is `surjOn_cone_integral_detVectors`, `New/Exchange/DeterminantVectors.lean`. On the finite side the difference of the integrals descends through the linear space of matches and is equivariant, by `exists_isEquivariantLift` in `New/Exchange/ExchangeDifference.lean`. Then `IsEquivariantLift.apply_stdColDatum_eq_zero` in `New/Exchange/StandardCoefficients.lean` promotes vanishing on the specified standard coefficients: its hypothesis is a family of centre-coefficient vanishings at adapted frames, not just a single constant-term vanishing; its conclusion is vanishing on all the integral standard-column data. These declarations and their singular-coefficient and approximation arguments are retained as formal-source input. The finite partner identities and the signed-product algebra above are necessary for their application; they do not replace them. This distinction is especially important when the equal-rank target is no longer definite.

The upgrade from an exchanged period to a seesaw square in `New/Exchange/OneSquare.lean`, `one_square`, uses continuous linearity in a real rapid datum. We give two elementary points behind that upgrade.

Lemma 32.12 (A rank-one span closed under pointwise limits)**.** *Let I be any set and $h : I \to \mathbb{C}$. If a net of functions of the form $c_i h$ converges pointwise to f, then $f = ch$ for some scalar c.*

Proof. If $h = 0$, all the functions are zero. Otherwise pick $j_0 \in I$ with $h(j_0) \neq 0$. Pointwise convergence implies $c_i \to f(j_0)/h(j_0)$. Evaluating at any other $j \in I$ and passing to the limit proves the assertion. The same argument applies to the convergence of sequences or to the closure in the product topology. □

There is no finiteness assumption on I in this lemma; one may group many finite twisting families into one family of indices. In the seesaw argument, a real datum is first approximated by linear combinations of elementary tensors. The scalar direction for such approximants must be common to every index. A different scalar for each index would leave no useful rank-one assertion. The lemma shows why the identity then passes to the limit; it does not assert that the limiting scalar is nonzero.

Lemma 32.13 (Uniform bound for the integrated kernel)**.** *Suppose Y, Y', A have finite quotient measure and nonnegative height functions at least one. Let $K_v(x, x', z)$ be a measurable kernel depending linearly on a real datum v, with*

$$|K_v(x, x', z)| \leq C\|v\|_N H_Y(x)^B H_{Y'}(x')^B$$

uniformly in $z \in A$. If y and y' are rapid, then multiplication by $y(x)y'(x')$ and by any unit-modulus character factor gives an integrable kernel, with L^1 norm at most a constant times $\|v\|_N$.

Proof. There exist constants a, a' with $|y(x)| \leq aH_Y(x)^{-B}$ and $|y'(x')| \leq a'H_{Y'}(x')^{-B}$. Thus the magnitude of the integrand is bounded by $Caa'\|v\|_N$, independent of all three quotient variables. Integrate this constant over the product of the three finite measure quotients. This also proves continuity of the resulting integral in the stated seminorm. □

The actual kernel bound and the passage from the height of a pair to the product of individual heights are formal-source inputs in `exists_integrable_kernel_bound` of `OneSquare.lean`. The proposition `one_square` combines this bound, the density of elementary tensors in the real-vector space with the prescribed weighted norms, the seesaw identity, and the partner comparison. Its finite consequence is precisely a square law with factor κ/c. The auxiliary scalar κ may initially be zero; c is the nonzero scalar of (32.12). The nonzero square scalars in the weighted-lift triangle are proved from injectivity and the properly chosen kernels afterward. The arithmetic estimates can divide by c, but not automatically by κ.

32.8 Level orders in the paths

The two-grid path from the starting periods to the terminal auxiliary pair is not an arbitrary path in a rank diagram. A move changes real data, local matrices and level characters. Proposition 31.11 furnishes the correctly synchronized path. It would be insufficient to find a formal sequence of dimensions of the same cost: each move must take a standard matched datum to another one in the family, and its resulting lift must be related by the square law. The preservation of the tidiness condition in `IsTidy.exists_square_move`, `New/Exchange/GridMove.lean`, remains formal-source input. The local square calculations above give the finite, cardinality-sensitive part of such moves.

The index bookkeeping of the triangle and the final comparison level can nevertheless be checked explicitly. For a block parabolic $P(n_0, \ldots, n_m)$ of $\mathrm{GL}_n(\mathbb{F}_q)$, where $n = \sum_i n_i$, block upper-triangular order gives

$$|P(n_0, \ldots, n_m)| = q^{\sum_{i<j} n_i n_j} \prod_i |\mathrm{GL}_{n_i}(\mathbb{F}_q)|.$$

Indeed the diagonal blocks must be invertible, all strictly upper blocks are arbitrary, and every such matrix is invertible. This holds also if some blocks have dimension zero, with the convention $|\mathrm{GL}_0| = 1$. The same counting argument as in the proof of Proposition 32.4 gives $|\mathrm{GL}_n(\mathbb{F}_q)| = \prod_{i=0}^{n-1}(q^n - q^i)$ by selecting independent columns.

For blocks $r, t, \ldots, t$ with m copies of t set

$$k(m) = mrt + \binom{m}{2} t^2.$$

This is the number of entries strictly above the diagonal blocks. Consequently $k(1) = rt$ and

$$m(k(1) - rt) - k(m) = -mrt - \binom{m}{2} t^2.$$

It is permissible to use a negative integer exponent in an identity of level factors, since residue-ring orders are nonzero integers. Calling this exponent a negative group order would be an error: it arises from expressing a quotient of known positive cardinalities.

To illustrate the count, take $r = t = 1$, $m = 2$ over $\mathbb{F}_3$. The parabolic has order $3^3 \cdot 2^3 = 216$, while $\mathrm{GL}_3(\mathbb{F}_3)$ has order $(27-1)(27-3)(27-9) = 11232$. Its index is 52, the number of full flags of lines inside planes: there are 13 lines and four planes containing any fixed line. At rank two the parabolic order is $3 \cdot 2 \cdot 2 = 12$ and its index is four. This is why naive powers of the first-step index cannot serve as the last-step index.

The full adelic index at a level given by inverse images of subgroups modulo split primes is the product of their finite indices, with the fixed level factors separated off. To see this, use surjectivity of reduction at each of the specified split primes and the Cartesian product of local maximal compacts; the quotient by the inverse image of the product subgroup has cardinality the product of the finite quotient cardinalities. In the source let $w(j)$ denote the `triangleOrderfalse` at stage j and J_m the level `lastResLevel`. Then `Data.stdLevelIndex_lastResLevel_eq` in `New/Exchange/ChainNorms.lean` proves

$$[K_0 : J_m] = C\,|R_0|^{m(k(1)-rt)-k(m)} d^m \frac{w(m)}{w(0)w(1)^m},$$

where R_0 is the residue ring at the initial varying modulus, d is the first auxiliary `gramFactor`, and C is independent of the twist. Here the symbol $k(j)$ is the number of strictly upper-block entries calculated above. This is precisely the elementary block count used in that declaration (via `card_parabolic_blockNumber`); the identification of the actual inverse-image level subgroups in the triangle is retained from the source. An advantage of this expression is the cancellation of the triangle-order quotient when it is combined with the weighted-chain norm. The powers of d and $|R_0|$ must still be tracked separately.

32.9 Why a false lift with one common root contributes no period

The purpose of comparison forms in Section 31.8 is not just to detect the prescribed eigensystem. They must be insensitive to certain other eigenspaces even when one root of their Hecke polynomial coincides with a root of a genuine lift. We explain an important mechanism that makes this possible. On an adelic quotient an arbitrary Hecke annihilator does not imply a zero function. In a sufficiently small argument of the Hecke polynomial, however, the operator cannot kill a nonzero bounded spherical function.

Lemma 32.14 (The subspace binomial identity)**.** *Let $a_{m,d}$ denote the number of d-dimensional subspaces of $\mathbb{F}_q^m$. Then*

$$\sum_{d=0}^{m} a_{m,d} q^{d(d-1)/2} T^d = \prod_{j=0}^{m-1} (1 + q^j T).$$

Proof. Fix a hyperplane in $\mathbb{F}_q^m$. Subspaces of dimension d contained in it are counted by $a_{m-1,d}$. A subspace not contained in it intersects the hyperplane in a $(d-1)$-dimensional subspace U. In the quotient by U it is a line with nonzero projection onto the one-dimensional quotient by the hyperplane. Choose its unique representative projecting to 1; its remaining coordinates have q^{m-d} possibilities. Hence

$$a_{m,d} = a_{m-1,d} + q^{m-d} a_{m-1,d-1}.$$

Multiplying by $q^{d(d-1)/2} T^d$ and summing, the second term is $q^{m-1}T$ times the polynomial in dimension $m-1$. Induct from the polynomial one in dimension zero. □

Proposition 32.15 (A Hecke injectivity bound for bounded functions)**.** *Let F be a bounded spherical function on an adelic unitary group of rank m at a split place w, with $q = q_w$. If the unnormalised Hecke polynomial operator at $X \in \mathbb{C}$ kills F and*

$$q^{m+1}|X| \le 1,$$

then $F = 0$.

Proof. Use the split Hecke-coset expression for spherical vectors, with the normalization of the source: the terms are indexed by all subspaces U of the m-dimensional residue vector space, and have coefficients $(-X)^{\dim U} q^{\dim U(\dim U - 1)/2}$. Each term is the appropriate translate of F. The term of dimension zero has coefficient one; its translate has the same supremum norm as F (and is the unshifted term with the identity representative). All other translates have supremum norm at most $\|F\|_\infty$. Thus

$$\|F\|_\infty \le \left(\prod_{j=0}^{m-1} (1 + q^j |X|) - 1 \right) \|F\|_\infty,$$

by Lemma 32.14. Using $1 + u \le e^u$ for $u \ge 0$,

$$\prod_{j=0}^{m-1} (1 + q^j |X|) \le \exp\left(|X| \frac{q^m - 1}{q - 1} \right) \le \exp\left(\frac{1}{q(q-1)} \right) \le e^{1/2} < 2.$$

The coefficient on the right is strictly less than one. A finite nonnegative supremum norm satisfying this inequality must be zero. □

Apart from the spherical coset description (`heckePolyOp_eq_sum`), this is a complete proof of `eq_zero_of_heckePolyOp_eq_zero` in `New/SiegelWeil/DescentsVanishPlanes.lean`. It is useful to view the argument as a Neumann-series estimate in a space of bounded functions. It does not require decomposing the bounded functions into automorphic representations, a Ramanujan bound, or the existence of a discrete spectrum on an indefinite group. It requires boundedness, which must be supplied independently.

In descending through a theta tower, write r for an initial rank, $r' = r + 2p$ for the rank after adding p hyperbolic planes, and $n > r'$ for the rank of the fixed definite target. The relevant Hecke root on the latter has magnitude $q^{(r+2-2n)/2}$. In the transfer from rank n

to rank r' the polynomial argument is rescaled in magnitude by $q^{(n-r')/2}$; the additional trivial-character factors must be nonzero. The rescaled argument satisfies

$$|X'| = q^{1-n/2-p}, \qquad q^{r'+1}|X'| = q^{r+p+2-n/2}.$$

Thus the integer hypothesis $n > 2r + 2p + 3$ gives exactly the small-argument condition in Proposition 32.15. Indeed the other, trivial Hecke factors have argument norms bounded by $q^{l+r'/2}|X|$ for $0 \le l < n - r'$, and the largest such bound equals $q^{-p} < 1$ when $p > 0$. The exact transfer and its conjugate-place adjunction are `hecke_transfer_of_equivariant` and `heckePolyOp_adjoint` in `Lib/Weil/Unramified/HeckeTransfer.lean` and `Lib/Weil/Unramified/SphericalHecke.lean`, respectively. These operator identities are formal-source inputs. Their normalization, and in particular the half-integral dimension shift in the argument, is essential to the strict bound.

Formal-source input (the tower vanishing). The initial zero auxiliary descent and reflection-sum annihilators imply general descent vanishing at that rank by `AuxPair.descentsVanish_of_reflectionSum`, `New/SiegelWeil/AuxPairVanishing.lean`, provided the local twists at the twisting places are nontrivial. This is a statement about all vacuum-off-level data, not only the one auxiliary datum. The induction theorem `descentsVanish_add_planes`, `New/SiegelWeil/DescentsVanishPlanes.lean`, then gives vanishing for data at the enlarged source: the inductive constant term vanishes; the archimedean and adelic estimates supply boundedness, and the Hecke operator is transported to the larger source in the preceding normalization. The case of adding zero planes needs no contraction estimate. For the other cases the full boundedness argument and the transport through each adapted splitting remain formal-source input. The just-proved proposition supplies the last step once boundedness and the Hecke equation are established.

From descended functions to comparison rows. The theorem `descentsVanish_hypSum` in `New/Rows/MemberOrthogonality.lean` applies the tower result to each conjugate matrix coefficient of an algebraic form killed by the auxiliary descent, the reflection sums, and the selected Hecke polynomial. It requires the inequality $2r + 2p + 3 < n$. The theorem `stdClassSum_period_eq_zero` in that module converts this vanishing into zero pairing with the comparison periods arising from the divided lattice. Its additional level, embedding and equivariance hypotheses are needed in this conversion; these two declarations are formal-source inputs. Notice the logical difference between sharing a single test root and possessing the entire Hecke eigensystem of a genuine lift. The former is simply enough to impose the operator equation. A false lift in its nullspace must be removed by the auxiliary descent condition and the nontrivial twists. It is not enough to compare roots as numbers.

32.10 Two finite ingredients of integral comparison

The rank and period arguments have been over $\mathbb{C}$; the construction of comparison forms also depends on p-adic integrality. We isolate two algebraic estimates behind Proposition 31.17. For the first, let $e : [a] \hookrightarrow [N]$ be an injection, write i_e for its column inclusion matrix, let $f : R \to S$ be a ring map, and let w be a matrix on which right multiplication by N-by-N matrices over S is defined. Put

$$H = \{g \in \mathrm{GL}_N(R) : g i_{e^c} = i_{e^c}, \quad w f(g) = w\}.$$

Use a determinant character χ extended by zero to nonunits of R and define

$$C(s) = 1_{wf(s)=wi_e}\, \chi(\det(i_e^{\mathsf{t}} s)), \qquad s \in \mathrm{Mat}_{N,a}(R).$$

Lemma 32.16 (A row-stabilizer cut)**.** *Suppose F is a function on matrices invariant under left multiplication by H. For any square matrix z and appropriately sized y,*

$$C(zi_e)1_{zi_{e^c}=i_{e^c}}F(y)=\begin{cases}\chi(\det z)F(z^{-1}y), & z\in H,\\ 0, & z\notin H.\end{cases}$$

Proof. Outside the fixed-column condition both sides vanish. Otherwise order the coordinates with the image of e first. Then $z=\begin{pmatrix}A & 0\\ B & I\end{pmatrix}$; $i_e^{\mathsf{t}}zi_e=A$, so $\det z=\det A$. Moreover

$$z=I+(z-I)i_ei_e^{\mathsf{t}},$$

whence $wf(z)=w$ is equivalent to $wf(zi_e)=wi_e$ (inclusion matrices are defined over the prime ring and unchanged by f). The determinant factor is zero unless z is invertible, in which case the two conditions defining H are now exactly those on the left. For $z\in H$, $z^{-1}\in H$ too, so invariance gives $F(z^{-1}y)=F(y)$. □

This proves the finite identity `columnDatum_mul_ite_eq` in `New/Rows/PartnerCut.lean`. It is the ingredient ensuring that a cut along selected columns enforces the required projector on integral data rather than introducing new, unsynchronized level characters. The group of fixed complementary columns by itself is too large; keeping the equation involving w is essential.

For the denominator estimate write $|\cdot|_p$ for an absolute value normalized by $|p|_p=p^{-1}$. The Schur lattice of a Young diagram has weights read from tableaux, and each entry corresponds to one occurrence of the associated standard basis coordinate. The essential combinatorial restriction is strict increase down every column of a semistandard tableau. If all entries belong to an alphabet of size l and the diagram has b nonempty columns, then at most b occurrences of any one alphabet symbol can occur. Consequently at most $b|S|$ entries can be drawn from a specified subset S of the alphabet.

Lemma 32.17 (Cost of bad columns)**.** *Let A be an l-by-l matrix over a p-adic field equipped with its Weyl lattice for a Young diagram as above. Suppose multiplying columns in a set S by an integer $M\neq 0$ makes A integral, all other columns already being integral. Then, with the lattice sup norm and the Schur action ρ,*

$$\|\rho(A)v\|_p\le |M|_p^{-b|S|}\|v\|_p.$$

Proof. Set $D=\operatorname{diag}(d_i)$ with $d_i=M$ on S and 1 outside. Then AD is integral. The integral polynomial Schur action preserves the integral lattice on all integral matrices, so has norm at most one for the corresponding lattice norm. The diagonal action $\rho(D^{-1})$ acts on the tableau weight space by M^{-u} where u counts entries from S. Since $u\le b|S|$ and $|M|_p\le 1$, its norm is at most $|M|_p^{-b|S|}$. Apply multiplicativity to $A=(AD)D^{-1}$. □

One must identify the bad columns before using the lemma. In the divided lattice arising from gluing, `norm_schur_inv_cols_le_weylGauge` of `New/Rows/CartanIntegrality.lean` chooses an integral change of basis on the columns whose reduction is specified modulo the squared chosen modulus. In that basis all but the indicated set of inverse columns are integral. The bound then has one factor per archimedean weight embedding, interpreted in the chosen p-adic embedding. The integral basis change and its effect on the Weyl gauge are formal-source input; the estimate after this change is the argument of Lemma 32.17. The full `weylGauge_cartanProduct_le` in the same file separately accounts for the positive and

negative columns of the gluing and for the conjugate embedding. In the source's notation the resulting exponent is

$$mloss = \sum_{\tau} \left(|a| \sum_{h<|N|} \kappa_{\tau}^{+}(h) + |m| \sum_{h<|m|} \kappa_{\tau}^{-}(h) \right).$$

The functions $\kappa^{\pm}$ count columns of the indicated heights, and the term $|a|$ is the number of divided coordinates. This estimate avoids charging every box of every diagram for a denominator occurring in just a few matrix columns. For instance the shape $(3,2)$ has three columns; a datum with only one bad column costs at most $|M|_p^{-3}$, rather than the total-degree bound $|M|_p^{-5}$. When several factors of this kind are multiplied in an integral Cartan product, their exponents add by multiplicativity and the ultrametric inequality. That saving is important in the subsequent determinant bound.

32.11 Limits of the reconstruction

The expanded argument now exhibits the finite matrix and character identities, the exact origin of the two sharp norm denominators, the algebraic part of the Rallis unfolding, the transition of signed critical-value products, the contraction for a bounded Hecke annihilator, and two comparison-lattice estimates. In particular the scalar in the exchange cannot simply be treated as an arbitrary norm of a theta integral, and the Gram factor cannot simply be treated as the norm of one finite indicator. In each case the normalization is what controls the twist-dependent arithmetic variation.

The analytic Siegel–Weil formula and its uniform estimates, genuine Weil model identifications, the existence and nonvanishing of the equal-rank test datum, the full exchange on partners with real determinant-vector surjectivity, propagation along the two-grid path, the sharp Euler pairing and its injectivity, the general tower boundedness and period conversion, and the geometric construction of the glued comparison lattice have been kept as formal-source inputs at their stated declarations. Nothing in the finite calculations substitutes a standard theorem with superficially similar hypotheses for these declarations. Their use in the rest of Theorem C continues to have the logical status specified in Section 31.5.

Chapter 33

The exchange construction, the two grids, and comparison lattices

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 35, 36 and the final audit in Chapter 57.

There are three rather different issues hidden in the comparison of twisted and untwisted row volumes. The first is the adequacy of the finite matching relation: one must know that an identity tested on standard column functions implies an identity for general partners. The second is the construction of the two paths of theta lifts with precisely matching signatures, parameters, and ramified data. The third is the integrality of a comparison form which sees both the small definite group and the large auxiliary group. It is inappropriate to infer this last integrality from a rational seesaw identity. A lattice on which the two constituents really glue is needed.

We expand these parts of Chapter 31 and the companion Chapter 32. The results below give conventional proofs of the local matching factorization, the stable-space argument with determinant Gauss sums, the real hyperbolic enlargement of determinant vectors, the terminal branching interlacing, and the construction of the divided lattice and its denominator estimate. We then return to how these calculations enter the volume formula and the rank-one value estimates. Some additional identities of the genuine global theta model are explicitly retained as inputs. In particular the analytic existence and control of the Fourier and constant terms in that model must not be confused with the finite Fourier calculations that are proved here.

33.1 Local integrals of complementary frames

Let F be a finite product of non-Archimedean local fields, O its product of valuation rings, and ϖ a simultaneous uniformizer. Write $G_d = \mathrm{GL}_d(F)$ and $K_d = \mathrm{GL}_d(O)$. A primitive column matrix is an injective matrix over O admitting a left inverse over O; for row matrices use the transposed definition. These sets are open and compact: they are images of K_N under projection to the appropriate first columns or last rows. They are open since invertibility of a square matrix over O holds exactly when its reduction is invertible. A locally constant function supported on the primitive set is thus a Schwartz–Bruhat function. More generally our Schwartz–Bruhat spaces are the locally constant compactly supported functions on the full matrix spaces whose support consists of full-rank matrices. This last convention is important: support can approach a singular matrix neither from within nor from outside.

Lemma 33.1 (Two-sided choice of coset representatives)**.** *There is a set of representatives $\mathcal{R}_d$ for the right-K_d cosets which also represents the left-K_d cosets. In particular both $\mathcal{R}_d$ and $\{r^{-1} : r \in \mathcal{R}_d\}$ represent the right-K_d cosets.*

Proof. The assertion can be proved inside each double coset $K_d g K_d$. Its right and left coset decompositions both have finite length, since the intersection of two compact open subgroups has finite index in each. A left Haar measure on G_d is also right invariant: the density $|\det A|^{-d}\, dA$ on the invertible matrices has that property, by the determinant Jacobian of left or right matrix multiplication. With K_d of volume one, the lengths of the left and right decompositions therefore agree. Each of the left cosets intersects each of the right cosets in the double coset: for $K_d gk$ and $k'gK_d$ take $k'gk$. Match the two finite sets by a bijection, choosing an element in each corresponding intersection. The union of these choices over the double cosets is the required set. For a product of fields the same argument applies using product Haar measures. □

In the source this is the freedom of choosing representatives in `exists_common_coset_reps`, `New/Exchange/BlockPartners.lean`. For an injective $Y : F^d \to F^N$, the lattice $\Lambda_Y = \{u : Yu \in O^N\}$ has rank d. Over each valuation ring it is free, by elementary divisors, and $Y\Lambda_Y$ is a direct summand of O^N: it is saturated because F is a field in each factor. For a matrix r of a basis of Λ_Y, Yr is primitive. Conversely any primitive Yr' has $r'O^d = \Lambda_Y$, because a primitive matrix has an integral left inverse. Exactly one $r \in \mathcal{R}_d$ has this property. Exactly one $r \in \mathcal{R}_d$ has Yr^{-1} primitive as well.

For any character $\chi : F^\times \to \mathbb{C}^\times$ tame on $O^\times$, define the local section of a column function by

$$\ell_Y(\Psi) = \sum_{r \in \mathcal{R}_d} \chi(\det r)\Psi(Yr).$$

For primitive-supported functions this sum has at most one nonzero term, with the representative independent of Ψ. For a Schwartz–Bruhat function with support contained in the injective matrices it has finite support: the linear embedding $g \mapsto Yg$ has a continuous left inverse, and the inverse image of the support is compact in G_d. Its image in the discrete coset set is finite. There is also an algebraic definition even for arbitrary functions: extend the ordinary summation functional on finitely supported sequences to all sequences by extending a basis, and compose with the evaluation map in the displayed formula. This extension has no assertion of continuity. It is not used to define a convergent integral for a nonintegrable datum.

For row functions define $\ell_X(\Phi) = \sum_r \chi(\det r)\Phi(r^T X)$. Use the same notation for the finite sum or the algebraic extension. Fix complementary dimensions $N = a + b$ and the splitting of indices into first b and last a. For $k \in G_N$, let Y_k be the first b columns of k, and X_k the last a rows of its inverse. Thus $X_k Y_k = 0$. A pair of Schwartz–Bruhat functions (Φ, Ψ) is matched in the larger sense if

$$\ell_{X_k}(\Phi) = \chi(\det k)\ell_{Y_k}(\Psi) \qquad (k \in G_N). \tag{33.1}$$

The definition does not demand that the functions be supported on primitive matrices. It is precisely `matchingPairs`.

Proposition 33.2 (Factoring a matched row functional)**.** *Let V be a complex vector space and let A be a linear map from the row Schwartz–Bruhat space to V satisfying*

$$\chi(\det g)A(\Phi(g\,\cdot)) = A(\Phi) \qquad (g \in G_a). \tag{33.2}$$

There is a linear map B on the column Schwartz–Bruhat space such that for all pairs in (33.1), $B(\Psi) = A(\Phi)$.

Proof. For each primitive X choose $k_X \in K_N$ with $X = X_{k_X}$ by completing the rows to a basis. Define a linear map of column functions into the space of all row functions by

$$T\Psi(X) = \begin{cases} \chi(\det k_X)\ell_{Y_{k_X}}(\Psi), & X \text{ primitive}, \\ 0, & \text{otherwise}. \end{cases}$$

Extend A algebraically to the space of all row functions. We will show that $B = A \circ T$ works and that the value on any matched pair is independent of both extensions.

For each $r \in \mathcal{R}_a$, let $f_r(X) = 1_{\mathrm{prim}}(X)\Phi(r^T X)$. This is a row Schwartz–Bruhat function; on the primitive set its support is compact, and multiplication by the primitive-set indicator is locally constant. Write

$$\Phi(X) = \sum_r f_r(r^{-T}X).$$

The equality follows because, for any full-rank X, exactly one r has $r^{-T}X$ primitive, and at a rank-deficient X both sides vanish. In fact only finitely many terms of this equality are nonzero as functions. The full-rank support of Φ is compact, and the sets on which $r^{-T}X$ is primitive are pairwise disjoint open sets covering it. A finite subcover proves the claim. By (33.2), applied to $g = r^{-T}$, one therefore has

$$A(\Phi) = \sum_r \chi(\det r)A(f_r).$$

For primitive X, the finite sum $\sum_r \chi(\det r)f_r(X)$ is $\ell_X(\Phi)$, so by matching it equals $T\Psi(X)$. Off the primitive set it is zero. Hence $T\Psi$ itself is Schwartz–Bruhat in the matched case, and its image under A equals $A(\Phi)$. This proves the proposition without any continuity of A or the extensions. □

This proves `matchingPairs.exists_linear_factor` in `BlockPartners.lean`. It is an algebraic factorization, not a norm estimate.

We check that the integral partners of the earlier chapter belong to this larger relation. Suppose both functions satisfy the inverse-character frame covariances and are supported on primitive frames, and their integral pairing is given by (32.2). Given an arbitrary $k \in G_N$, choose $g \in G_a$, $u \in G_b$ such that gX_k and $Y_k u$ are primitive. The column image is exactly the integral kernel of the row matrix: indeed this is true over F factor by factor, and both are saturated. Complete the column frame to a basis by choosing a right inverse of gX_k; this gives $k' \in K_N$ with $X_{k'} = gX_k$, $Y_{k'} = Y_k u$. Relative to k, the determinant changes by the factor $\det u / \det g$, as in Chapter 32. Thus

$$\chi(\det g)\Phi(gX_k) = \chi(\det k)\chi(\det u)\Psi(Y_k u).$$

Taking u the column representative and g the transpose of the row representative proves (33.1). This also proves the content of `IsPartner.mem_matchingPairs`, `PartnersMatch.lean`: local constancy of one integral partner implies local constancy of the other, since on K_N the relation uses the continuous frame projection and the locally constant determinant character, and the frame projections are open maps onto the primitive loci. Zero extension is locally constant on the clopen primitive loci and their complements.

Equivariant passage to the difference of periods. For a pair of real data v, v' and a scalar c, let the two target-valued linear maps defined by genuinely convergent theta integrals be A_v and $A'_{v'}$. The determinant covariance required by (33.2) must be derived from the Weil

action and the change of variable on the source group. Suppose it holds. Proposition 33.2 gives a linear difference

$$\mathcal{D}(\Psi) = B_v(\Psi) - cA'_{v'}(\Psi)$$

on the space M of matched column data, with $\mathcal{D}(\Psi) = A_v(\Phi) - cA'_{v'}(\Psi)$ for matched pairs. A translate by $g \in G_N$ preserves the matched relation: the row function becomes $\chi(\det g)\Phi(\cdot\, g)$ and the column function becomes $\Psi(g^{-1}\cdot)$. To check this, substitute $g^{-1}k$ in (33.1); the determinant factor is exactly $\chi(\det g)^{-1}$. The same statement with the local Weil determinant multipliers on the two target integrals gives equivariance of the difference. The cancellation of those multipliers is a consequence of complementary dimensions and equal combined rank twist, not an arbitrary identification of the two Weil representations. The automorphy and centre-coefficient properties of this difference are the properties collected in `IsEquivariantLift` by `exists_isEquivariantLift`, `ExchangeDifference.lean`. The factorization and matching parts have now been proved. The precise local naming and global automorphy laws of the Weil representation used to obtain the full `IsEquivariantLift` are retained as inputs.

33.2 Generating all partner data by stable spaces

The following local argument supplies a particularly useful converse. Let $s : O^N \to O^b$ and $x : O^b \to O^N$ satisfy $sx = I_b$. Define the standard column function

$$S_s(Y) = 1_{\mathrm{Mat}_{N,b}(O)}(Y)\,\mathrm{inv}_O\,\chi(\det(sY)),$$

where $\mathrm{inv}_O\,\chi(a)$ is $\chi(a)^{-1}$ for $a \in O^\times$ and zero otherwise. Thus its support consists of primitive columns. This is `stdColDatum` in the source. Assume P is a subspace of the column Schwartz–Bruhat space stable under $f(Y) \mapsto f(gY)$ for $g \in G_N$. We give two results corresponding to `partner_mem_of_stdColDatum` and `IntegralFrame.partner_mem_of_unimodular_vanishing` in `New/Exchange/StableSpaces.lean`.

Lemma 33.3 (Positive standard function generates all partners)**.** *Suppose $p \in P$ has the form $p(Y) = r(Y)S_s(Y)$ for a nonnegative integer-valued function r, with $r(x) > 0$. Then every locally constant column function Ψ belonging to integral partners of character χ is in P.*

Proof. Because Ψ is locally constant, is supported on the primitive set, and obeys right frame covariance $\Psi(Yq) = \chi(\det q)^{-1}\Psi(Y)$, compactness of $\mathrm{Mat}_{N,b}(O)$ supplies an integer $c > 0$ such that

$$\Psi(Y + \varpi^c Z) = \Psi(Y) \qquad (Y, Z \in \mathrm{Mat}_{N,b}(O)).$$

For products of local fields choose the maximum of the finitely many depths. Define an invertible matrix

$$d = xs + \varpi^{-c}(I_N - xs), \qquad d^{-1} = xs + \varpi^c(I_N - xs).$$

The formulas follow since xs and $I_N - xs$ are orthogonal idempotents. In particular $sd = s$, $dx = x$, and d^{-1} has integral entries. Consider

$$f(Y) = \int_{K_N} \Psi(k^{-1}x)p(dkY)\,dk,$$

with K_N of Haar measure one. This integral is a finite linear combination of translates of p: the functions of k in the integrand are locally constant, and $k \mapsto p(dk\cdot)$ has open stabilizer because p is a locally constant compactly supported function. The last fact follows by taking a common residue depth on its compact support and a compact integral box containing that support. Thus $f \in P$.

If $S_s(dkY)$ is nonzero, write $m = skY \in K_b$. Then $dkY \in O^{N\times b}$, whence kY is integral by integrality of d^{-1}. Moreover Y is primitive, as m is invertible. The complementary part satisfies $(I - xs)kY \in \varpi^c O^{N\times b}$, so $kY = xm + \varpi^c Z$. By local constancy and covariance,

$$\Psi(Y) = \Psi(k^{-1}xm) = \chi(\det m)^{-1}\Psi(k^{-1}x).$$

Consequently, on the primitive set,

$$f(Y) = \Psi(Y)\int_{K_N} r(dkY)1_{\{S_s(dkY)\neq 0\}}\,dk.$$

The integral is independent of the primitive frame Y, since K_N acts transitively on such frames. At $Y = x$, it is positive: $dx = x$ and $r(x) > 0$; the integrand is locally constant near $k = 1$ on the set where S_s is nonzero, and hence is positive on an open neighborhood of positive Haar measure. On nonprimitive columns both f and Ψ are zero. Dividing by this positive constant proves $\Psi \in P$. □

We next show why it suffices to annihilate the unimodular centre coefficients in the exchange difference. Let

$$[K\ I\ L] \in \mathrm{GL}_N(O), \qquad \begin{bmatrix} K' \\ I' \\ L' \end{bmatrix} = [K\ I\ L]^{-1}$$

with both K and L of b columns. Let $\bar{O} = O/\varpi O$, write η for the residue character equal to χ^{-1} on $O^\times$ (extended by zero on nonunits), and assume η is primitive as a multiplicative character of $\bar{O}$ in the sense of Chapter 32. Let ψ be an additive character of F with conductor exactly O. The pairing $\psi(\varpi^{-1}uv)$ on $\bar{O}$ is primitive.

For $z \in \mathrm{Mat}_b(F)$ put

$$S(z)(Y) = S_{K'}(Y + LzK'Y).$$

The frame identities include $K'L = 0$, $K'K = I_b$, $L'L = I_b$, and $I'L = L'K = I'K = 0$. Thus $S(z)$ is supported on full-rank columns and $S(z+T) = S(z)$ for integral T. The latter follows since whenever $K'Y$ is integral the extra term $LTK'Y$ is integral, while $K'(Y+LzK'Y) = K'Y$. When $K'Y$ is not an integral invertible matrix both sides vanish.

Proposition 33.4 (Unimodular-coefficient synthesis)**.** *Suppose M contains all the $S(z)$ and P is as in Lemma 33.3. Assume the following implication: a vector $f \in M$ belongs to P whenever all linear functionals Λ on M satisfying, for some $T \in \mathrm{Mat}_b(F)$,*

$$\Lambda(S(z)) = \psi(\mathrm{tr}(Tz))\Lambda(S(0)), \qquad T \in \mathrm{GL}_b(O) \implies \Lambda(S(0)) = 0 \tag{33.3}$$

vanish on f. Then every locally constant column function of integral partners is in P.

Proof. For each $r \in \mathrm{Mat}_b(\bar{O})$ choose an integral lift $\widetilde{r}$, with $\widetilde{I}_b = I_b$. Let

$$q = \sum_r \eta(\det r)S(\varpi^{-1}\widetilde{r}) \in M.$$

We verify that every functional in (33.3) kills q. If $\Lambda(S(0)) = 0$ this is immediate. Otherwise the invariance under integral translations of z gives $\psi(\mathrm{tr}(TZ)) = 1$ for all integral Z. Taking matrices with a single nonzero entry and using the conductor, T has integral entries. Then

$$\Lambda(q) = \Lambda(S(0))\sum_r \eta(\det r)\psi(\varpi^{-1}\,\mathrm{tr}(T\widetilde{r})).$$

The sum is a matrix determinant Gauss sum. Proposition 32.4 shows that it is zero unless the reduction of T is invertible. In that latter case $T \in \mathrm{GL}_b(O)$, contradicting $\Lambda(S(0)) \neq 0$. Therefore $q \in P$.

Since P is stable under G_N it also contains $p(Y) = q(-\varpi^{-1}Y)$. We compute this function. In the frame coordinates write $Y_K = K'Y$, $Y_I = I'Y$, $Y_L = L'Y$. Set $m = -\varpi^{-1}Y_K$. A term indexed by r can be nonzero only if

$$m \in K_b, \qquad Y_I \in \varpi O^{(N-2b)\times b}, \qquad Y_L \equiv \widetilde{r}m \pmod{\varpi}, \quad Y_L \in O^{b\times b}.$$

Its coefficient is $\eta(\det r)\chi(\det m)^{-1}$. If r is invertible then Y_L is invertible; the conditions determine $r = \overline{Y_L}\,\overline{m}^{-1}$ uniquely. By the definition of η the coefficient is exactly $\chi(\det Y_L)^{-1}$. Hence $p(Y)$ is $S_{L'}(Y)$ times a zero-or-one function. At the integral frame $x_0 = L - \varpi K$, the conditions hold with $m = I_b$ and $r = I_b$, and $L'x_0 = I_b$. In particular the nonnegative multiplier of $S_{L'}$ is positive at x_0. Apply Lemma 33.3 with $s = L'$ and $x = x_0$. □

The last proof explains the mechanism that is otherwise easy to miss in the formal argument: an additive Fourier test at an invertible index can be converted by a multiplicative determinant Gauss sum into a positive cut-off standard column function, which is cyclic for the stable space of partner data. The sign of the inverse character matters in the calculation of $p(Y)$. If it were changed to the original character, the two factors of $\chi(\det m)$ would not cancel.

For the genuine automorphic application take P to be the stable kernel of the difference of the two periods, and Λ to be a centre Fourier coefficient of its value on a datum. The character eigen-relation in (33.3) is the equivariance for translations by the indicated radical. The proof of `IsEquivariantLift.apply_stdColDatum_eq_zero`, `New/Exchange/StandardCoefficients.lean`, establishes the needed hypothesis about *all* such functionals from two geometric vanishing conditions: vanishing at the unimodular Gram matrix of the smaller member in an adapted splitting, and vanishing at invertible indices not represented by its archimedean signature. The translation of an arbitrary primitive rational Hermitian index to these conditions involves a second splitting, weak approximation, the Fourier completeness on the radical quotient, and a local orthonormalization of integral Hermitian matrices. Proposition 33.4 settles the residue synthesis in that argument. The geometric vanishing and its promotion to the full automorphic coefficient statement are distinct inputs, to be verified in the exact mixed model.

33.3 Construction of the real data on an enlarged Hermitian space

The exchange theorem begins at a hyperbolic enlargement and descends by removing planes. There are two real analytic problems. One must construct data with a prescribed slice, and the removal of a plane must be sufficiently surjective on determinant vectors. Replacing these requirements by existence of some nonzero Gaussian integral would not suffice: the resulting datum would generally depend on the twist.

Lemma 33.5 (Smooth local Witt extension)**.** *Let $J = J^*$ be nondegenerate, $x \in \mathrm{Mat}_{n,b}(\mathbb{C})$ with x^*Jx invertible. There is a smooth map u from a neighborhood of x to $U(J)$ such that $u(x) = 1$ and $u(y)x\mathbb{C}^b = y\mathbb{C}^b$ for all y in that neighborhood. It can be taken to be a globally smooth map into $U(J)$ with this property only near x.*

Proof. Put $A^\sharp = J^{-1}A^*J$ and

$$P_y = y(y^*Jy)^{-1}y^*J.$$

For y close to x, P_y is a $\sharp$-self-adjoint idempotent with image $y\mathbb{C}^b$ and kernel its J-orthogonal complement. Put $S = P_yP_x + (1-P_y)(1-P_x)$; then $S(x) = 1$ and $SP_x = P_yS$. Thus S is invertible near x and maps the two summands of P_x to those of P_y. By taking the $\sharp$-adjoint, $P_xS^\sharp = S^\sharp P_y$, so $Q = S^\sharp S$ commutes with P_x. For y sufficiently near x we can form

$$Q^{-1/2} = \sum_{n\geq 0} \binom{-1/2}{n} (Q-1)^n.$$

It is $\sharp$-self-adjoint since all coefficients are real, commutes with P_x, and depends smoothly on y. Then $u(y) = SQ^{-1/2}$ has $u(y)^\sharp u(y) = 1$ and maps the desired subspace onto $y\mathbb{C}^b$. The invertible b by b change of frame is $C = (x^*Jx)^{-1}x^*Ju(y)^{-1}y$, for which $u(y)xC = y$.

To make the map global, shrink the neighborhood so that the matrix logarithm $L(y) = \log u(y)$ is given by its series at 1. From $u(y)^\sharp = u(y)^{-1}$ one obtains $L(y)^\sharp = -L(y)$ by the functional calculus near the identity. Take a real-valued smooth bump ρ equal to one in a smaller neighborhood, supported where L is defined. Extend ρL by zero and use $\exp(\rho L)$. Its $\sharp$-adjoint is its inverse, and it agrees with u near x. □

This is the smooth Witt theorem `exists_contDiff_witt` of `New/Exchange/RealPair.lean`, with a direct matrix proof instead of the inverse-function-theorem argument of that file. It does not require J to be positive definite. The global extension is useful because compactly supported mixed-model data will be specified by a global smooth formula that uses $u(y)$ only near x.

We recall the relevant elementary analytic argument for the real Fock model. Choose signed coordinates so that the holomorphic variable in a positive coordinate is z, and in a negative coordinate is $\bar{z}$. Taking the product of these choices is a real isometry, so the weighted moments are unchanged. In unscaled coordinates, write a Fock function as $v(z) = e^{-|z|^2/2}f(z)$ with f entire, and normalize Lebesgue measure by $d\mu(z) = \pi^{-d}d\,\mathrm{vol}(z)$ on $\mathbb{C}^d$. The coherent kernel is $K(z,w) = \exp(z\cdot\overline{w} - |z|^2/2 - |w|^2/2)$ and has $|K(z,w)| = e^{-|z-w|^2/2}$. For a real radius or norm, replacing the Euclidean norm by the sup norm changes only constants depending on dimension. Define $m_N(v) = \int (1+|z|)^N |v(z)|\, d\mu(z)$. The source's notion `IsRapidDatum` asks that all these moments be finite; it does not impose a particular algebraic polynomial expression on v.

Lemma 33.6 (Simultaneous polynomial approximation)**.** *A Fock vector with $m_N(v) < \infty$ for all N can be approximated in every one of the norms m_N by one sequence of Gaussian times holomorphic polynomials.*

Proof. The reproducing identity is $v(z) = \int K(z,w)v(w)\,d\mu(w)$. For Gaussian monomials it is proved by integration in polar coordinates; the full Fock reproducing identity follows by the orthogonal Gaussian monomial basis. Absolute convergence for a rapid vector also follows from the kernel bound. Replace the factor $\exp(z\cdot\bar{w})$ by its scalar Taylor polynomial of degree l, obtaining $v_l(z)$ as the integral with this truncated kernel. The coefficients are Gaussian moments of $v(w)$, so v_l is a Gaussian times a polynomial. For all l the norm of the kernel is bounded by

$$\exp\bigl(-(|z|^2+|w|^2)/2 + |z||w|\bigr) = e^{-(|z|-|w|)^2/2}.$$

For fixed w the integral in z of this bound times $(1+|z|)^N$ is at most $C_{d,N}(1+|w|)^{N+2d}$: use polar coordinates $r = |z|$, bound $1 + r \leq (1+|w|)(1+|r-|w||)$, and integrate the remaining polynomial times the Gaussian over the real line. The bound is integrable against $|v(w)|d\mu(w)$. Since the Taylor polynomials converge pointwise to the exponential, dominated convergence for the product measure in (z,w) shows $m_N(v - v_l) \to 0$. The polynomial degree l is independent of N, so the assertion is simultaneous. □

This reconstructs `exists_tendsto_weightedNorm` of `Lib/Weil/Fock/MixedFockModel.lean`. The argument gives density in all moments, which is stronger than density in the Hilbert norm. This strengthening matters in the square law: matrix elements of the kernel may grow polynomially with adelic height, so continuity merely in the Hilbert norm cannot be invoked.

We spell out the regularity estimate used in constructing smooth slices. In an adapted hyperbolic splitting the real oscillator realization can be written as the tensor product of a Schrödinger coordinate and a lower Fock space; change of the Schrödinger polarization is by a normalized Fourier transform. A normalized coherent vector at z of the lower Fock space is denoted ω_z; it is the unitary Heisenberg translate of the vacuum and satisfies

$$m_N(\omega_z) \le C_N(1+|z|)^N.$$

This follows directly from the absolute Gaussian profile, by the inequality $(1+|x+z|) \le (1+|x|)(1+|z|)$. The signed Fock action of a unitary matrix U on the coherent expansion sends z to the corresponding linear coordinate transform and introduces its vacuum multiplier of absolute value one. For a smooth compactly supported multiplier t and a smooth family $U(a)$, the mixed-model section $t(a)\omega(U(a))v$ is constructed by the coherent expansion, Fourier transform in a, and integration of the resulting coherent translates in the extra coordinates. The following estimate verifies the existence in the moment topology for smooth data. We use only the standard mixed-model coordinate and Fourier identities in this argument; their exact form in the source is `MixedModel.exists_section` of `Lib/Weil/MixedModel/RapidSections.lean`.

Lemma 33.7 (Moment estimate in the section construction)**.** *Let $d(a)$ be smooth of compact support in real dimension p, and let $B(a)$ be a smooth matrix on a neighborhood of that support. For the coherent kernel $\omega_{B(a)y}(w)$, define*

$$H_y(\xi,w) = \int_{\mathbb{R}^p} e^{-2\pi i a\cdot\xi} d(a)\omega_{B(a)y}(w)\,da.$$

For any M,N there are C,L depending on d,B,M,N but not y such that

$$\int (1+|w|)^N |H_y(\xi,w)|\,d\mu(w) \le C(1+|y|)^L(1+|\xi|)^{-M}. \tag{33.4}$$

Hence for a rapid vector $v(y)$ the integral of $v(y)H_y(\xi,w)$ against $d\mu(y)$ has integrable moments of arbitrarily large degree in both ξ and w.

Proof. The coherent kernel is exponential of the expression $w\cdot\overline{B(a)y} - |w|^2/2 - |B(a)y|^2/2$, possibly with a smooth unit-modulus quadratic phase in the coordinates. The first two derivatives of the phase in a are polynomial expressions in y,w with uniformly bounded coefficients on the compact support, and the same statement is true for every finite derivative order. By induction and the product rule each derivative of $d(a)\omega_{B(a)y}(w)$ of order at most $2l$ is bounded in absolute value by

$$C_l(1+|w|+|y|)^{4l}e^{-|w-B(a)y|^2/2}.$$

One may increase the exponent without changing the argument. Multiplying by $(1+|w|)^N$, integrating over w, and using $|B(a)y| \le C_B|y|$ gives $C_{l,N}(1+|y|)^{N+4l}$ uniformly in a. Apply $(1-\Delta_a)^l$ under the integral by integration by parts; the multiplier on the Fourier exponential is $(1+4\pi^2|\xi|^2)^l$. The Fourier integral is thus bounded in the weighted L^1 norm in w by $C_{l,N}(1+|y|)^{N+4l}(1+4\pi^2|\xi|^2)^{-l}$. Take $2l \ge M$. The last assertion follows by integrating against $|v(y)|$ and taking M larger than p plus the required degree of the ξ moment. □

In the mixed model the Fourier-transformed vector in this construction is further translated by a linear function of ξ in the enlarged Fock space. Lemma 33.7 still supplies all its moments, because Heisenberg translation loses at most $(1 + C|\xi|)^N$. Therefore those integrals are Bochner integrals of rapid Fock data. Applying the slice map, Fourier inversion on the compactly supported smooth integrand and coherent reproduction recover the prescribed $t(a)\omega(U(a))v$. This proves the section existence for smooth choices used here, using the mixed-model transform identities; in the formal development the weaker C^8 regularity is permitted by a further scaled Fourier bound. No assertion about that minimal regularity is needed for a smooth construction.

For a fixed lower datum v take $U(a) = 1$ and a nonnegative, nonzero smooth compactly supported t. The slice at zero in the opposite polarization is a fixed nonzero scalar times the integral of the forward slices, since the opposite polarization is the Fourier transform, evaluated at zero. The scalar is fixed by the two Haar measures and the mixed-model vacuum convention. Normalize t by that scalar and $\int t$. This gives one rapid enlarged datum whose opposite slice is exactly v. It is the construction `exists_rapid_with_realSlice` in `New/Exchange/RealPair.lean`. The scalar is independent of the varying finite character; this is the feature needed in the equal-rank starting identity.

33.4 A plane removal on determinant vectors

At an infinite place let W, V have sign involutions J_W, J_V (with diagonal entries ± 1 in orthonormal coordinates). The positive coordinates of $U \in \mathrm{Hom}(V, W)$ are those with equal source and target signs. If P is the projection onto these coordinates, it is $P = (1 + J_{\mathrm{Hom}})/2$. For a prescribed coefficient matrix F form the square matrix $M(U)$ by the positive-coordinate projection followed by multiplication by F in the convention of the source. Denote by M_+ and M_- its two principal sign blocks. A determinant vector of the family `detVectors` is a Gaussian times

$$\overline{\det(M_+)^m \det(M_-)^{m+2k}};$$

in the adelic real model multiply these factors over the infinite places with $m = m_w$. The exponent difference $2k$ will cancel the character arising from the hyperbolic plane.

Let $W' = W \oplus H$ and $V' = V \oplus H$ be enlarged by one hyperbolic plane. Its adapted orthonormal positive-negative coordinates can be chosen so that in the isotropic basis the sign matrix exchanges the two line coordinates, and the original block has sign matrix J_W or J_V. The adapted mixed-model slice at a line variable a sends a positive-coordinate polynomial to its evaluation at an old vector with the two new isotropic coordinates equal to $-a$. Indeed if J_W in the new basis has line block $\left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, then

$$P_+(u_1, u_2, v) = ((u_1 + u_2)/2, (u_1 + u_2)/2, P_+(v)).$$

At $(u_1, u_2) = (-a, -a)$ this is exactly the required projection. Orthogonality of the adapted frame also gives squared Euclidean norm $2|a|^2 + |v|^2$ in these coordinates. In the mixed-model polynomial slice, the Gaussian is evaluated at the same vector. This follows by inserting the coherent exponential in the Fourier transform of the hyperbolic line and differentiating in its polynomial variables. Explicitly the Fourier transform of the Gaussian with linear coefficient is again a Gaussian with the transformed linear coefficient; independent Gaussian derivatives generate every polynomial in the projected coordinates. The normalized transform formula for the chosen splitting is `PositivePart.exists_slice_eq` in `New/Exchange/Determinant Vectors.lean`; the projection and norm calculation above show why the two line coordinates in this formula really are equal, rather than opposite.

Proposition 33.8 (Determinant enlargement and its scalar)**.** *In the preceding adapted coordinates extend the coefficient matrix F by an identity on the new hyperbolic plane, with its normalization chosen by the line scale of the mixed model. After the line slice with local nonzero scalar a the two principal minors become respectively*

$$\det(M'_{\pm}) = a \det(M_{\pm}).$$

Suppose the rank identity $\mathrm{rk}(W') = 2k + 2q(W')$ *holds. When the cone integration on the enlarged pair is formed with weight $k+m_w$ at the infinite place w, the sliced determinant-vector integrand is a positive real scalar times the original determinant vector, with no U-dependent multiplier. In particular if the standard cone integrand is integrable and incorporates a nonzero positive line measure, it is possible to normalize the enlarged determinant vector so that the integral is the originally prescribed determinant vector.*

Proof. The chosen coefficient extension is block diagonal between the hyperbolic planes and the old coordinates. Under the slice the enlarged matrix in the two new isotropic coordinates is aI_2; their image in the positive-negative orthonormal coordinates is still aI_2. On the old block the matrix is M. Thus the new equal-sign principal blocks have one additional diagonal entry each, equal to a, and determinant multiplicativity proves the first assertion. The ratio of the polynomial factors is

$$\overline{a}^{m_w}\overline{a}^{m_w+2k} = \overline{a}^{2(k+m_w)}.$$

In the cone integrand the line character of weight $k + m_w$ contributes the compensating complex phase. The product of the line factors in the normalization of `coneIntegrand` is $|a|^{2(k+m_w)}$ times the positive idele-absolute-value factor of the line. To see the cancellation, write $a = re^{i\alpha}$, $r > 0$; multiplying the preceding polynomial ratio by $e^{2i(k+m_w)\alpha}$ gives $r^{2(k+m_w)}$. The maximal compact action on the extended polynomial in this step has no further phase, since the determinant exponents on the two signs differ by $2k$ and the rank identity says the vacuum exponent difference is the same. This is the sign and rank check for the character in the cone integrand.

In addition the Gaussian acquires just the positive line Gaussian, since its restriction has squared norm $2|a|^2 + |v|^2$. It follows that at the product of infinite places the integrand is of the form

$$D(q)v(U), \qquad D(q) = \exp(-\textstyle\sum_w |a_w(q)|^2)|d(q)|^z \prod_w |a_w(q)|^{2(k+m_w)}$$

in the source's line coordinates and positive Haar density. Here q denotes the line integration variables, $d(q)$ the corresponding idele in the line torus, and z the cone exponent. All of these factors are nonnegative and strictly positive on the nonzero line locus. If v is nonzero, integrability at one point with $v(U) \neq 0$ implies finiteness of $\int D$; nonzero positive line measure implies this integral is positive. Multiplying the extended coefficient by $(\int D)^{-1}$ produces the prescribed v. For $v = 0$ take the zero datum. Polynomial Gaussian data are rapid by direct Gaussian integration, so this normalization stays in the determinant-vector family. □

This calculation gives the matrix and positive-scalar part of `surjOn_cone_integral_detVectors`, `New/Exchange/DeterminantVectors.lean`. The hypotheses about the adapted mixed model, the line-scale vacuum multiplier, and convergence are part of the genuine global line identity `integrable_coneIntegrand` in the source; they are not consequences of an arbitrary polynomial extension. The proposition with those hypotheses gives surjectivity, and it provides an explicit inverse on every determinant-vector ray. Multiple plane removals compose these inverses. The zero slice in complementary polarizations likewise composes by Fubini on the rapid sections: separating the two line variables and integrating in either order is valid by their joint absolute moment bounds.

33.5 How the equal-rank normalization is obtained

We now state the intermediate identities used in `New/Exchange/EqualRankValue.lean`, to separate what is needed analytically from what is pure normalization. Let X be of even rank, and $X^+ = X \oplus H$ an enlargement by a hyperbolic plane. For a real datum v and character of weight α write

$$R_{X^+}(\nu, v) = \frac{|\theta_{X^+,X^+}(\nu; v, \mathrm{standard})(1)|^2}{\Pi_\nu^{\mathrm{int}}(\mathrm{rk}\, X^+/2)}.$$

Here Π_ν^{int} denotes the positive product `lValueProdInt` of the earlier chapter, including its precise conductor normalization. For the top identity X^+ is read with its adapted splitting. The number R_{X^+} is called `equalRankThetaRatio`. The relevant source declarations are as follows.

1. `thetaLarge_eq_apply_one_of_even_rank` in `LowRankIdentity.lean`: for the equal-rank theta integral on standard-column data the value on general standard columns is obtained from the value at the identity by the real slice and Weil determinant covariance of the columns. This uses the rational open orbit in the enlarged hyperbolic splitting.

2. `stdExchange_of_periods` of the same file: when the top periods on the two source groups have the appropriate real slicing relation, it gives a standard-column identity for all target vectors and a factor whose magnitude is expressed using R_{X^+}.

3. `exists_stdExchange_top` in `RealPair.lean`: for X^+ of rank $2k + 2q + 2$ and the corresponding top pair $V' \oplus X$, V', there are real data realizing the preceding relation simultaneously for all tame characters of the prescribed weight. Its construction uses the smooth Witt lemma and the slice sections, with one compactly supported bump around the nondegenerate frame.

4. `exists_constant_term_thetaLarge` in `Lib/SiegelWeil/Tower/LargeSideConstantTerm.lean`: integrate one line radical of the larger-side theta integral with standard-column data. The remaining cone integral is a normalized lower-rank theta datum; the scalar is independent of the varying datum, and its squared magnitude is the appropriate last Hecke factor.

These identities of the genuine global theta integrals are still taken as geometric and analytic inputs. In items 2 and 3, the exact open-orbit statement, measure factors, and real action on the two data have to agree; Lemmas 33.5 and 33.7 give the local construction and rapidity in the latter item, not the global period identity.

For clarity consider the real geometry of item 3. Let the base subspace of the combined source have frame $x_0 = (I, 0)^T$ and ambient form $J = J_0 \oplus J_1$. For a frame x with x^*Jx invertible put

$$z(x) = x(x^*Jx)^{-1}x^*Jx_0.$$

The difference $x_0 - z(x)$ is orthogonal to $\mathrm{im}\, x$. At $x = x_0$, $z(x_0) = x_0$. If $z(x)$ is sufficiently near x_0, its Gram matrix is invertible, and therefore its b columns span exactly $\mathrm{im}\, x$. The smooth Witt map $u(z)$ sends $\mathrm{im}\, x_0$ to $\mathrm{im}\, z = \mathrm{im}\, x$, so $u(z)^{-1}x = x_0C$ for a unique invertible b by b matrix C. Its Gram matrix is $C^*J_0C = x^*Jx$. This is the factorization enabling the two periods to be evaluated in a common set of real coordinates near a nondegenerate frame. All these maps are smooth where defined. Choose a nonnegative smooth bump $t(z)$ supported in the indicated neighborhood, and a second bump on $u(z)^{-1}x$. Their product has compact support as a function of x because $x = u(z)x'$ with z, x' in compact sets and u continuous. Multiplying the product by a fixed polynomial determinant and the requisite compact-unitary

translates of the tensor product of the two lower real data gives the smooth slice used in item 3. The precise theta integral equality for these real data uses the normalized period and open-orbit identities of items 1 and 2; the smoothness, support, and existence of one rapid Fock section follow from the earlier lemmas.

We next make the normalization argument explicit. For removal of one plane the usual standard-column constant-term formula has a zero-vector term and nonzero isotropic terms. On the smaller side a primitive standard-column datum kills the nonzero isotropic terms: their finite invariance under a line translation would force a primitive column frame to annihilate a nonzero coordinate. The formal proof uses `constTermAt_smul_k` and `name_eq_zero_of_invariant` in the mixed model to prove this assertion for the actual subspace. On the larger side the nonzero line integral is exactly the cone integrand, so it creates the lower datum in Proposition 33.8. After canceling the common nonzero vacuum factor the ratio of squared scalar magnitudes is the transition quotient of Lemma 32.10. Since removing a plane decreases the rank difference by two, its product over all removed planes telescopes to the requisite $\Pi_\nu^{\mathrm{int}}(e)$. This is the induction in `exists_stdExchange_removing_planes`. The surjectivity asserted there on determinant vectors follows from Proposition 33.8; down the other side the zero slices compose, as just explained. For the arbitrary real datum used in the starting exchange the normalized cone integral, not an inverse of the polynomial map, is used.

The proof of nonvanishing of a universal equal-rank constant in the source has a useful structural form. Put J for the definite group of Hermitian rank $2k$ and Z for the zero-dimensional group. Begin with the alternative that all equal-rank ratios $R_{X^+}(\nu, v)$ are zero. Apply the top period identity to the product with the zero-dimensional partner, and remove all the hyperbolic planes using the determinant surjectivity. This makes the zero-constant identity reach a nonzero determinant (in the case of Z, vacuum) datum at the definite edge, forcing the rank-zero theta value to vanish. On the zero-dimensional space there is a single rational vector, the zero vector, and the normalized vacuum at it is 1. The class integral is over the trivial group and has positive mass. Thus that value is nonzero. The application of this contradiction to the exact top identity is the first part of `exists_equalRankThetaRatio_value`. It uses the period identities above, in particular the lift and descent of the standard-column equality even if the scalar on that equality were zero; it does not assume the result to be shown.

It is not yet enough to have $R_{X^+}(\nu_0, v) \neq 0$ for one character. The datum v must work for all characters simultaneously. The second part of `exists_equalRankThetaRatio_value` exchanges twice between the two extreme ranks (J, Z) and (Z, J), with real data chosen independently of the characters. If B, C are the two universal exchange constants and q the real scalar of the rank-zero datum, the resulting scalar identities give

$$R_{X^+}(\nu, v)^2 = BC|q|^2. \tag{33.5}$$

Here the conductor and local Hecke factors cancel: in one direction the dimensions differing by $2k$ give $\Pi_\nu^{\mathrm{int}}(k)$ and in the other direction $\Pi_\nu^{\mathrm{int}}(-k)$, with product one by the signed-product convention. This is precisely the identity called `F` in the proof of that declaration, obtained by pairing one exchange with the adjoint of the other. Equation (33.5) therefore gives one number R_{X^+} independent of ν, since every ratio is a nonnegative real number. Nonvanishing at ν_0 implies its strict positivity for all ν. In the last exchange, to return from the flipped Hermitian form to its original sign, the Weil flip contributes a unit-modulus determinant character. Its square magnitude does not change the equality. A separate scalar from the change of Haar normalization is independent of the twist. Hence the resulting constant has

$$|c(\nu)|^2 = C_0 \Pi_\nu^{\mathrm{int}}(e), \qquad C_0 > 0,$$

with $e = \operatorname{rk} V - \operatorname{rk} W/2$. This is the normalization of `exchange_of_partners` and `exchange_theorem`. The nonzero constant C_0 cannot be obtained from the local matrix Gauss sum alone.

In the reconstruction the double-period identity leading to (33.5), the geometric coefficient vanishing and the normalized line constant-term formula are still the principal global inputs to the exchange theorem. The algebraic factorization, stable-space step, smooth-real-data mechanism and scalar bookkeeping of their deductions have now been unpacked.

33.6 From a convergent exchange to an operator square

Suppose a complementary-rank exchange is available for one universal rapid vector on a direct sum of two real pair spaces. To use it in a chain of theta lifts one must pass from that rapid vector to the prescribed joint harmonics on the two constituent spaces. We give the analytic and linear portions of the argument in `New/Exchange/OneSquare.lean`. Write G_1, G_2 for the exterior groups of the square and A for the larger intermediate Hermitian space. The source estimate for a primitive finite datum is

$$|K_v((g_1, g_2), a)| \le C m_N(v) H_{G_1 \oplus G_2}(g_1, g_2)^B, \tag{33.6}$$

uniformly in the intermediate element a; the more general estimate contains a negative power of $H_A(a)$. This is the specialized kernel estimate used in `pairTheta_decay_of_onto`; its proof uses Poisson summation in the finite named datum, specifically its primitive support. We do not take a growth bound in a in place of the uniform bound in this assertion.

For rapidly decreasing y_i on the exterior quotients, choose C_i with $|y_i(g_i)| H_{G_i}(g_i)^B \le C_i$. The direct-sum height satisfies $H_{G_1 \oplus G_2}(g_1, g_2) \le H_{G_1}(g_1) H_{G_2}(g_2)$, in the source's normalization with heights at least one. Multiplying (33.6) by $|y_1(g_1) y_2(g_2)|$ gives a constant bound

$$|y_1(g_1) y_2(g_2) K_v((g_1, g_2), a)| \le C C_1 C_2 m_N(v). \tag{33.7}$$

Character factors are unitary and have no effect. With fixed quotient measures of finite total mass, the product integrand is absolutely integrable, and the resulting linear functional L of v obeys $|L(v)| \le C' m_N(v)$, where C' includes the three masses. This proves `exists_integrable_kernel_bound` once the stated genuine kernel estimate is available. The estimate is needed also for the smaller side, with the primitive block in its matching datum.

By Lemma 33.6, approximate the exchanged rapid vector by polynomial Gaussian vectors in all moment norms. A polynomial in the real coherent coordinates of a direct sum is a finite sum of products of polynomials on the two factors. Thus every approximant is a finite sum of elementary Gaussian tensors. For each elementary tensor the real convolution with the fixed type picks out a matrix coefficient scalar, independent of the finite twist. This statement is the compact-type equivariant convolution identity `IsEdgeFamily.exists_lift_eq_realConv`. It is an identity about the prescribed joint harmonic and the source and target type, not an assertion that two arbitrary vectors in an abstract irreducible representation are automatically proportional. After applying this identity and the adjoint relation to both exterior groups, an elementary tensor contributes some scalar times the prescribed composition of lift and descent. The scalar may be zero.

For the limit argument let I index all test forms, test functions, twists, and primitive finite data to which the universal real vector must apply. Write P_i for the value of the prescribed composition at test i and $D_i(v)$ for the integral of the approximating tensor datum. Each finite polynomial approximant v_n has $(D_i(v_n))_{i \in I} = \kappa_n (P_i)_{i \in I}$, with the same scalar for every i. By (33.7), $D_i(v_n) \to D_i(v)$ for each fixed i. If all P_i vanish then all the limits vanish and one may take $\kappa = 0$. Otherwise select i_0 with $P_{i_0} \neq 0$; then $\kappa_n = D_{i_0}(v_n)/P_{i_0}$ converges, and each limit equals κP_i. This proves the closed-span step without finiteness or a uniform norm on the index set I. It is essential here to approximate by one sequence independent of all the

twists. Dividing by the nonzero complementary-rank exchange scalar $c(\nu)$ gives the square factor $\kappa/c(\nu)$. The determinant covariance of the two edge kernels comes from the local Weil action on named point data. Once this covariance and the two kernel estimates are verified, the convergent exchange implies the entire square law; this is the analytical argument of `one_square`. It does not prove that κ is nonzero, nor is nonvanishing needed for the crude-row transport. In the injective weighted-chain application this issue is settled separately by the first-arrow norm formula.

33.7 Terminal interlacing and the signatures of the two grids

We give a proof of the combinatorial grid identities of `New/Exchange/GridParameters.lean`, rather than using the terminal branching as an unspecified representation-theoretic requirement. Keep m, k even, $r = k - 1 + m$, $t = 5r$ odd, $h = 2r^8$ and $h_2 = h/2$ as in the admissible data. At an infinite place write $w_l = w_0 + lh$, where w_l includes h_2 (that is, $w_l = b_a + lh + h_2$ with $a = 1$ or 2 as appropriate to the place). Use $z = 0, 1$ for the two grids, i, j for their coordinates and o for the triangle offset. The rank at (z, i, j) is

$$n(z, i, j) = r + z + i(t + z) + j(t - z).$$

There are $r + z$ base coordinates, i added blocks of size $t + z$, and j added blocks of size $t - z$. The base signs at coordinate u, $0 \le u < r + z$, are

$$(-1)^{o+a+\min((a-o)_+, j)}, \qquad a = (u + 1 - k)_+,$$

and the signs on all added blocks are $+1$. Here $v_+ = \max(v, 0)$ is the truncated natural subtraction in `entrySign`.

Lemma 33.9 (Grid signatures and parity)**.** *At offset $o = 0$ the number of negative signs is $\lfloor (m - j + z)/2 \rfloor$ for $0 \le j \le m$. Moreover $n(0, i, j) \equiv 1 + i + j \pmod 2$ and $n(1, i, j) \equiv 0 \pmod 2$. The diagonal member of grid 1 has the signature of the diagonal member of grid 0 plus one line of sign $(-1)^j$ (when $i = j$).*

Proof. The base signs with $a \le j$ have sign $(-1)^{2a} = 1$. For $a > j$ they have sign $(-1)^{a+j}$; the positive integers a run exactly once from 1 to $r + z - k = m - 1 + z$. Thus the negative signs correspond to odd positive integers $b = a - j \le m - 1 + z - j$. There are $\lfloor (m + z - j)/2 \rfloor$ such integers. Added blocks are positive. Since r, t are odd, the displayed ranks have the stated parity. At $i = j$ the ranks in the two grids differ by one. Their negative counts differ by $\lfloor (m - j + 1)/2 \rfloor - \lfloor (m - j)/2 \rfloor$, which equals one exactly when j is odd since m is even. This proves the sign of the added line. A basis matching preserving the signs exists by separately matching the positive and negative coordinates; it gives the bijection used in `gridDiagEquiv_spec`. For $j = 0$ the extra base coordinate has positive sign, so the original coordinate inclusion may be used. □

The doubled Harish–Chandra parameter on the base block has entries

$$2\rho_u = \begin{cases} k - 2 - 2u, & 0 \le u < k - 1, \\ 2h_2 - 2w_{u+2-k}, & k - 1 \le u < r + z. \end{cases}$$

The positive added block numbered l, $1 \le l \le i$, is $2w_l + t - 1 - 2b$ for $0 \le b < t + z$, and the negative-centred added block numbered l, $1 \le l \le j$, is $t - 1 - 2(w_{l+o} + z) - 2b$ for $0 \le b < t - z$. This last description concerns the values of the parameter, not the signs of the summands, which are all positive. To all entries add 1 if $z = 1$; if $z = 0$ subtract 1 when $i + j$ is odd,

and otherwise add zero. The rank parities in Lemma 33.9 show that every doubled entry is congruent to $n(z,i,j)-1$ modulo 2. The spacing of the centres $w_l \mapsto w_{l+1}$ is h, and the width of an added block is at most $2t$. The base entries away from the central block lie halfway between the centres of the negative added blocks. From $h > 4t+4$, $2w_1 > k+t+2h$, and the parameter bounds (these follow from the large-log inequalities), they are distinct from the added entries and from one another. In particular when $o = 0$ every pair of oppositely signed coordinates is separated by at least $h-2t-2$: the negative signs only occur on base outer coordinates. Distinct entries within any same-sign arithmetic block are separated by 2, not by h. This is why a gap condition between opposite signs must not be confused with a lower bound h on the separation of *all* entries. The source verifies the exact `HCParam.HasGap` convention at offset zero and along the first vertical strip by the same arithmetic-block comparison; for the joint harmonic conditions a further comparison of the centered parameters on adjacent edges is required.

Proposition 33.10 (Terminal branching interlacing)**.** *At offset zero and node (m,m) let V be the multiset of doubled parameters in grid 0 and let Z be the multiset of negatives of the doubled parameters in grid 1. Then for every real x,*

$$0 \le \#\{z' \in Z : z' > x\} - \#\{v \in V : v > x\} \le 1.$$

Proof. We enumerate entries from largest to smallest. On the positive-centred band of centre $2w_l$, $1 \le l \le m$, the V entries are

$$2w_l + t - 1,\ 2w_l + t - 3, \dots, 2w_l - t + 1,$$

and those of Z, obtained by negating the negative-centred block of grid 1, are

$$2w_l + t - 2,\ 2w_l + t - 4, \dots, 2w_l - t + 2.$$

They alternate with a V entry at each end. The extra outer base entries of Z on the positive side are $2w_l - h - 1$ for $1 \le l \le m+1$. For $l > 1$ such an entry lies strictly between the positive bands of centres $2w_l$ and $2w_{l-1}$, since $w_l - w_{l-1} = h$ and $h > t+1$. At $l = m+1$ it lies above the largest band, and at $l = 1$ below the smallest band but above the central block, by the lower bound on w_1. Thus descending across the positive side, the difference starts at zero, rises to one on a base Z entry, drops to zero on the next V, rises to one on the intervening Z, and so on, ending at one just before the central block.

On the central block V has $k-2, k-4, \dots, 2-k$ and Z has $k-3, k-5, \dots, 1-k$. These alternate in the order V, Z, so starting at one the difference remains zero or one and ends at one.

The negative-centred band of V at $-2w_l$ has endpoints $-2w_l + t - 1$ and $-2w_l - t + 1$, whereas the reflected positive block of grid 1, giving the negative band of Z, has endpoints $-2w_l + t$ and $-2w_l - t$. They alternate with a Z at each end. The base negative extra entries of V are $h - 2w_l$, $1 \le l \le m$. In decreasing order each occurs immediately before its respective negative band and strictly after the preceding band or central block. Indeed it is halfway between consecutive negative band centres (with the same endpoint inequality as above). Thus each such base V lowers the difference from one to zero, and the following band keeps the difference in $\{0,1\}$ and raises it to one at its end. This proves the count inequality for every cut between entries, and at each entry itself since no entries coincide. All the separation requirements used here are only the indicated simple inequalities on h, w_1, k, t. □

This gives `gridParam_interlace`. At the terminal nodes both groups are compact by Lemma 33.9. The Weyl dominance convention for the dual representation reverses and negates

its highest weight, so this interlacing of doubled parameters is exactly the condition for restriction through the codimension-one embedding. To see the half-integral shift, a rank d ordered highest weight $\lambda_1 \geq \cdots \geq \lambda_d$ has doubled parameter $2\lambda_i + d + 1 - 2i$, and a rank $d+1$ highest weight μ of the other (dual) member has doubled parameter $2\mu_i + d + 2 - 2i$. The displayed count inequality says $2\mu_i + d + 2 - 2i > 2\lambda_i + d + 1 - 2i > 2\mu_{i+1} + d + 2 - 2(i+1)$. Since the weights are integral, these inequalities are equivalent to $\mu_i \geq \lambda_i \geq \mu_{i+1}$. The ordinary branching rule for unitary groups now gives a nonzero pairing. Its existence over the rational coefficient field follows from algebraic representation theory: the intertwining equations in the highest-weight models are rational equations, so a nonzero complex solution implies a nonzero rational solution. Clearing its denominator gives an integral matrix for the pairing. The needed *quantitative* bound for such a matrix in the specific Weyl realization, including the powers of the determinant characters, is the additional estimate in `exists_branching_pairing` and `endBranching`. The preceding elementary interlacing is enough for existence, not for that norm bound.

33.8 The finite square move with no accumulated denominator

The standard local exchange kernel of Chapter 32 has an orbit denominator. We now prove the finite transformation of tidy data that removes it. This is the main finite part of `IsTidy.exists_square_move` in `New/Exchange/GridMove.lean`.

Work over the finite valuation-pair ring R and use p, v, u, r, T for the free modules attached to the relevant vertices and the extra Fourier block. In the notation of the cited declaration the old edges are $(p, v), (Q, U)$ with $Q = p \oplus T$, $U = u$, and the new edges are $(v, r), (U, R')$ with $R' = r \oplus T$. We identify the row and column bases by the bijections prescribed in that declaration. The complementary dimensions satisfy $\operatorname{rk} p + \operatorname{rk} r = \operatorname{rk} v + \operatorname{rk} u$. For a tuple S the old data are

$$A = S(p, v) : p \to v, \qquad B = S(Q, U) = [B_0 \ B_1] : p \oplus T \to u.$$

Set $Y = (A, B_0)^T : p \to v \oplus u$. In the new tuple choose matrices

$$D : v \to r, \qquad C = \begin{pmatrix} C_0 \\ C_1 \end{pmatrix} : u \to r \oplus T, \qquad X = [D \ C_0] : v \oplus u \to r.$$

Write $S'(S, C, D)$ for the tuple agreeing with S on the unaffected edges w, with these two new edge matrices and zero elsewhere. The old and new edges are disjoint from w. The phase in the move is $\psi(\operatorname{tr}(C_1 B_1))$, for the additive character ψ of the residual ring of the grid modulus. This is the trace of an endomorphism of T. In the ladder of just one grid, T is the empty block, so the phase is 1.

The two old edge characters are χ_1, χ_2 in the order above. The new characters on $(U, R'), (v, r)$ are χ_1, χ_2 in that order. Thus put $\chi = (\chi_1 \chi_2)^{-1}$. Let $\Delta_\chi(X, Y)$ be the complementary-frame kernel of (32.4); in this ordering its support is $\ker X = \operatorname{im} Y$ with both frames primitive. The kernel is zero elsewhere.

Let Λ be the old tidy functional. For any scalar function d of Y and any test function f on tuples define

$$G_d(f) = \Lambda\left(S \longmapsto \sum_{C,D} d(Y) \Delta_\chi(X, Y) \psi(\operatorname{tr}(C_1 B_1)) f(S'(S, C, D)) \right). \tag{33.8}$$

Initially set $d = 1$; we shall show that the same functional is obtained by a choice of d for which the nonzero coefficients in this expansion are roots of unity.

For a family g of vertex automorphisms as in the admissibility condition, write $g_{R'} = \operatorname{diag}(g_r, 1_T)$ under the prescribed embedding and $b = \operatorname{diag}(g_v, g_u)$ on $v \oplus u$. We need the other compatibility hypothesis of the source: for any $k \in \mathrm{GL}(p)$ we may replace the old vertex at p by k and at Q by $\operatorname{diag}(k, 1_T)$, leaving the rest (including the unaffected edges except for the permitted vertices) the given admissible family. In particular the action on the matrices in w is unchanged. Denote that old family by $\widetilde{g}$.

Lemma 33.11 (Covariance of the replacement functional)**.** *Under these compatibility hypotheses, write $\alpha_{\mathrm{old}}(\widetilde{g})$ and $\alpha_{\mathrm{new}}(g)$ for the products of edge characters of the old and new path respectively. Then*

$$G_{d(b^{-1}Yk)}(f \circ g) = \alpha_{\mathrm{new}}(g) G_d(f).$$

Proof. The old tuple transforms Y to $Y_1 = b^{-1}Yk$ and B_1 to $g_u^{-1}B_1$; the new tuple transforms X to $X_1 = g_r^{-1}Xb$ and C_1 to $C_1 g_u$. The trace phase is unchanged. The row-column kernel obeys

$$\Delta_\chi(X_1, Y_1) = \chi(\det g_r \det k / \det b)\Delta_\chi(X, Y)$$

by (32.5). The change of variables on the finite sum over the matrices (C, D) is bijective. This gives the stated transformed integrand in (33.8) with factor $\chi(\det g_r \det k / \det b)^{-1}$. Using old covariance of Λ multiplies by $\alpha_{\mathrm{old}}(\widetilde{g})$. Finally the ratio of the two edge character products is

$$\begin{aligned} \frac{\alpha_{\mathrm{new}}(g)}{\alpha_{\mathrm{old}}(\widetilde{g})} &= \frac{\chi_1(\det g_r / \det g_u)\chi_2(\det g_r / \det g_v)}{\chi_1(\det g_v / \det k)\chi_2(\det g_u / \det k)} \\ &= (\chi_1\chi_2)(\det g_r \det k / \det b), \end{aligned}$$

as required. The unaffected edge product cancels. This proves the formula with exactly the determinant powers of the two edges; using the same character on all four edges would in general give a different formula. □

Taking $d = 1$ proves covariance for the new functional. Taking $g = 1$ gives $G_{d(Yk)} = G_d$ for every $k \in \mathrm{GL}(p)$. Choose d to be $|\mathrm{GL}(p)|$ at one representative of each primitive column-frame orbit and zero at the other frames. Then its average over the right action is 1 at every primitive Y. Since the kernel is supported on these Y, linearity and averaging show $G_d = G_1$. The coefficient $d(Y)\Delta_\chi(X, Y)$ is either zero or a root of unity. The additive phase is also a root of unity, because the finite additive group has finite exponent.

If the old datum is supported on primitive matrices at its distinguished μ-edges, then Y is primitive as soon as either A or B has a left inverse (a left inverse for B restricts to one for B_0). The support condition propagates across the move. Indeed if $A = \mathrm{pr}_v Y$ has a left inverse L_A, the projection $1 - Y L_A \mathrm{pr}_v$ vanishes on Y and restricts to the identity on the complementary u component. Compose it with a right inverse of X as in Lemma 32.3; it follows that $C_0 = X|_u$ has a left inverse, hence so does C on the new μ-edge (U, R'). If $B_0 = \mathrm{pr}_u Y$ has a left inverse, use the analogous projection to conclude that $D = X|_v$ has a left inverse. This is exactly the propagation needed when the old edge carrying the twist and the new edge carrying the twist are opposite. Each new matrix has at most Q_* possibilities. If the old datum has order q then (33.8), with the chosen d, has at most Q_*^{q+2} nonzero summands in a presentation by root-of-unity coefficients. Thus it is short of order $q + 2$ as well as covariant.

Finally suppose the operator square law has scalar c and tests $L(X)$ and $M(Y)$ on the two routes. For each primitive Y it says $M(Y) = c \sum_X \Delta_\chi(X, Y) L(X)$ by Lemma 32.3. Here tests may depend on the matrices in w and on the extra Fourier variable $K : T \leftarrow u$. Multiply this equality by $\psi(\mathrm{tr}(KB_1))$ and sum over K, then apply Λ. The variable K can be read as the new block C_1, independently of $X = [D\ C_0]$. The right side is precisely cG_1 applied to the new route with $K = C_1$ and old unaffected matrices in w. This proves the equality of functionals

in `IsTidy.exists_square_move`. The use of $G_d = G_1$, not a pointwise replacement of Δ_χ by $d\Delta_\chi$, is what proves both the identity and absence of an orbit denominator in the new datum.

33.9 Moving the descent along the path

Let the two paths of stage j descend from $(0, m)$ to $(0, j)$ and then lift to (j, j), one in each grid; the grid type z is suppressed in this description. The stage-j comparison is a bilinear restriction period at the diagonal pair, evaluated against the tidy datum. To move a single descent past a lifts one repeats the preceding square move a times with $T = 0$. The datum has order increased by $2a$ and the scalar is a product of a square constants. This is seen formally by induction on a: for $a = 0$ the leg descent is literally the first arrow of the longer vertical path, with no change in the tuple or scalar. For the step $a \mapsto a + 1$, apply the induction with the final horizontal edge temporarily in the unaffected set w, then cross the square at the end and use (33.8) to remove the two old matrices. The test depends only on w and on the current forms, so the functional identity from that move is valid with precisely the same test. This reconstructs the combinatorial induction in `Data.descent_past_gridPathMap`, `New/Exchange/RowLadder.lean`, once the square law is available for each genuine edge.

To go from stage j to $j + 1$ do this in both grids for $a = j$. These $2j$ square moves use $2j$ square costs and $4j$ more point evaluations in the order exponent. The final diagonal period is transported by the real seesaw and change of the line sign, using one more move with the Fourier block T corresponding to that line. Its modulus scalar, from the Fourier-normalized flip of the line, is a pure conductor-cardinality power with one unit of cost. Together with the other diagonal scalar in `Data.hasRoute_succ`, the source counts at most $4j + 2$ square costs and $4j + 2$ increases in order. This last step uses the genuine line-kernel identity `exists_modKernel_flip_line` and the real seesaw with the bordering Hermitian embedding; it is not a consequence of the abstract path induction alone. For the finite part, compatibility of g with the bordering embedding is precisely the rule coupling the two diagonal vertex matrices in Lemma 33.9; the unaffected path has no edge incident to the released vertex. This is the situation of Lemma 33.11, while one of the two old edges is a μ-edge. Thus the square move is applicable as a finite transformation at this last step as well.

Once the stage identity with those two scalar factors is known, the total equality follows by multiplying, not adding, the nonzero factors $c_j(\theta)$ and by adding their square cost bounds. The total cost and order increase are $\sum_{j=0}^{m-1}(4j + 2) = 2m^2$. The initial datum of order $4m$ is thereby replaced by a terminal datum of order $4m + 2m^2$. The role of the predicate `IsTidy`, stronger than covariance just at the endpoint levels, is now apparent: without the independent admissible change at the intermediate vertex p in Lemma 33.11, the averaging over column frames would be invalid. A denominator of the order of a large finite general linear group would have been introduced at each of the squares. The proof above removes it before the next square can be applied.

We record also the finite-dimensional surjectivity step in reaching the initial rows. If a weighted chain A from one conjoined real type to the definite endpoint is injective, and its bilinear transpose D is the corresponding descent, then the restriction of D to the endpoint level space is onto the initial level space, provided the transpose identity holds on those level spaces. In fact the image is finite dimensional, and any vector annihilating it has A-image orthogonal to the entire endpoint space, hence zero. For the application the conjugate real type is identified by the sign-flip isometry, so the bilinear transpose is identified with the Hermitian adjoint. The level preservation of both chains follows from their finite point-datum covariance for principal level, and the equality of transpose pairings follows by Fubini for each lift-descent pair and cancellation of the finite weight w_j at that edge. This is

`Data.iotaPlus_adjoint_weightedChain` and `Data.gridLevelForms_eq_map_iotaPlus` in `New/Exchange/RowsReached.lean`. Thus it is the injectivity of the long weighted chain that gives surjectivity of a fixed map, not a choice of forms separately for each row.

To pass from the second-grid starting level to the first-grid principal form space, one must still prove the quantitative restriction assertion `Data.eq_zero_of_orthogonal_gridLevel Forms`. Its proof constructs a real translate from a two-factor discrete-series coefficient with a nonzero integral, approximates local convolution by a Poincaré sum at the deep principal level, and treats points near the boundary by a first-minimum estimate. The estimate at the finite level is uniform over the whole form space. The finite-dimensional argument just given shows what it suffices to prove, but does not justify replacing this assertion by a qualitative restriction-density theorem. We keep this quantitative real restriction input (in addition to the genuine square law and line-kernel identity). With it, bilinear duality gives the surjection in Proposition 31.9, and the proved finite transport of tidy data applies to all of its period tests simultaneously.

33.10 The divided lattice that glues the two comparison groups

We now turn from the first (forward) row volume estimate to the reverse estimate. This part uses the definite target of the single lift of good forms of Chapter 31. Its coefficient vector is a Cartan product of a vector on the large definite part and one on the small definite part. Their ramified finite support must be combined into an integral datum whose descent is their product. Splitting the real vector alone is not enough; it is the lattice constructed here that permits the two finite conditions to lie on the same point form.

At the chosen split places let O be the product of valuation rings (the two chosen places above the primes of g), π a generator of their product ideal, and $R = O/\pi^2 O$. We use N for the rank of the big part, s_0 for that of the small part, and r for the common row rank. Let $F : O^r \hookrightarrow O^N$ and $Z : O^r \hookrightarrow O^{s_0}$ be coordinate inclusions, so each has an integral coordinate left inverse F^T, Z^T. Let S be an a by N matrix over R, with $a \le N$; these are the rows indexed by the member data in the source. Put $T_S = \binom{S}{\pi I_N}$ over R. In the actual splitting, the inclusion inc of the common r coordinates in the full target is $(0, Z)^T$ after the prescribed basis reordering, and the condition on the big block uses the fixed inclusion F. These are the embeddings in `DividedSplitting` of `New/Rows/StackedDeterminants.lean` and `New/Rows/GluedFormTheta.lean`.

The first fundamental observation is that T_S has an integral lift with row lattice of full rank, even if S has very low rank after reduction. This allows the lattice to vary with the row-test matrix S, rather than attempting to use one splitting at every residue point.

Lemma 33.12 (A basis of the divided row module)**.** *Choose an integral lift $\widetilde{S}$ of S. There are integral square matrices D, E of size N such that $ED = DE = \pi I_N$, and the rows of D form a basis of the O-module generated by the rows of $\binom{\widetilde{S}}{\pi I_N}$. In particular, over R, for any column matrices X, X' of the corresponding size,*

$$T_S X = T_S X' \quad \Longleftrightarrow \quad \bar{D}X = \bar{D}X'.$$

Proof. Over each discrete valuation ring factor the row module contains πO^N and is contained in O^N, hence is free of rank N. Choose an ordered basis, assembled as rows of an invertible matrix D over the fraction field. Since every πe_i is in the row module, the coefficients expressing πe_i in that basis are integral; putting them into the rows of E gives $ED = \pi I$. As D is invertible over the fraction field this gives $DE = \pi I$ as well. Combine these choices over the finitely many factors. Since the two row generating sets are integral linear combinations

of one another, the equivalence of the two matrix tests holds after any reduction, in particular modulo π^2. □

Let $P_Z = ZZ^T$ be the diagonal projector onto the selected small coordinates and $B = 1 - P_Z$ its complement. Form two square matrices of size $N + s_0$ over O:

$$C = \begin{pmatrix} E & FZ^T \\ 0 & P_Z + \pi B \end{pmatrix}, \qquad G = \begin{pmatrix} D & -DFZ^T \\ 0 & \pi P_Z + B \end{pmatrix}. \tag{33.9}$$

Then

$$CG = GC = \pi I_{N+s_0}. \tag{33.10}$$

Indeed the upper-left and lower-right blocks have product πI; the upper-right block of CG is $-\pi FZ^T + FZ^T(\pi P_Z + B) = 0$. The reverse equality follows by inversion in the fraction field. These are the matrices constructed by the Hermite basis argument in the source's `DividedSplitting.exists_isGlued`.

Proposition 33.13 (Indicator identity for divided gluing)**.** *Write $\mathcal{H} = \{Y \in \mathrm{GL}_r(R) : Y = I \bmod \pi\}$. For every full-size column matrix $X = (X_+, X_-)^T$ of r columns over R,*

$$\begin{aligned} &1_{\{\bar{G}X = \pi(0,Z)^T\}} \\ &= \sum_{Y \in \mathcal{H}} 1_{\{X_- Y = Z\}}\, 1_{\{T_S X_+ Y = T_S F\}}. \end{aligned} \tag{33.11}$$

Furthermore for an integral change of basis $K \in \mathrm{GL}_N(O)$, chosen independently of X, the big column block of $\bar{G}K$ satisfies $\pi(\bar{G}|_+ K)$ has zero columns outside a prescribed set of a positions. Finally $|\mathcal{H}| = |O/\pi O|^{r^2}$.

Proof. The lower block of the equation on the left of (33.11) is $(\pi P_Z + B)X_- = \pi Z$. Multiplying by B shows $BX_- = 0$, so $X_- = ZM$ for $M = Z^T X_-$. The equation then says $\pi M = \pi I$. As the annihilator of π in $O/\pi^2 O$ is precisely πR, we have $M = I + \pi A$. It is invertible, with inverse $I - \pi A$. This is the unique element $Y \in \mathcal{H}$ satisfying $X_- Y = Z$. The upper block of the left-hand equation is $\bar{D}(X_+ - FZ^T X_-) = 0$, or equivalently $\bar{D}X_+ Y = \bar{D}F$, because $Z^T X_- Y = I$ and Y is invertible. By Lemma 33.12, this is the upper test on the right of (33.11). Conversely any summand equal to one gives the unique Y and these two equations, and implies the original full equation by reversing the calculation. Thus either both sides are zero or exactly one term on the right is one.

For the column assertion, reduce the row matrix S modulo π, factor by factor. Its rank over a residue field is at most a. There is therefore a basis over the residue field such that all but a prescribed columns are zero: extend a basis of the kernel to a full basis and label the remaining at most a vectors by those positions. Lift the basis matrix to $K \in \mathrm{GL}_N(O)$; any lift of an invertible residue matrix is invertible over the valuation ring. The same K works for the row module generated by $\widetilde{S}$ and πI modulo π, because every row of D is an integral combination of those rows. Hence all columns of DK outside the prescribed positions vanish modulo π, giving the asserted vanishing of $\pi\bar{D}\bar{K}$. The big block of G consists of D and zero rows, so the same assertion holds for it. Finally $Y = I + \pi A$ with A unique modulo π, and multiplication is addition of A because $\pi^2 = 0$ in R. There are $|O/\pi|^{r^2}$ choices. □

The proof gives all the finite assertions of `DividedSplitting.IsGlued`, as well as `DividedSplitting.exists_congruenceGroup_inv_eq_one_sub` and the bad-column assertion in `DividedSplitting.exists_isGlued`. Notice two separate features of the use of modulus π^2: the factor πI_N ensures the existence of the row basis and bounds its denominator by one π, whereas the quotient of unit groups at the two depths makes the normalizing Y unique. Modulus π alone would not make the latter argument.

Realizing the matrix equation on a genuine finite unitary point. At a split place of E the unitary group of a standard definite Hermitian space identifies with $\mathrm{GL}_{N+s_0}(E_w)$, the matrix at the conjugate place being inverse conjugate-transpose of the given matrix. Define the finite adelic unitary point λ by taking C of (33.9) at the selected places, the inverse conjugate-transpose of C at their conjugates, and the identity away from these places. This defines the source's `DividedLattice`. Indeed $\pi\lambda^{-1} = G$ at the chosen places by (33.10), and λ^{-1} is integral at their conjugates because it is the conjugate-transpose of C. The generating idele π has value one away from the selected places, so the entire matrix $\pi\lambda^{-1}$ is integral at every place, with reduction $\bar{G}$ at the chosen modulus. This proves `DividedLattice.exists_reducedInverse_eq`.

If a finite unitary point u is integral and equals $I \bmod \pi^2$ at the selected places, then $\lambda^{-1}u\lambda = I \bmod \pi$ there. In fact write $u = I + \pi^2 A$, and compute

$$C^{-1}uC = I + \pi GAC,$$

which is integral and congruent to $I \bmod \pi$. At their conjugates integrality follows from the inverse-transpose relation and the fact that the calculated matrix has determinant a unit, as its reduction is the identity. At all other places the conjugate is just u. This is the local-level statement `DividedLattice.reduceAt_conj_eq_one`. Together with Proposition 33.13, it specifies the actual adelic lattice rather than just a mod-π^2 congruence.

The row-fixing subgroup of the large definite factor used in gluing fixes T_S by right action modulo π^2; in particular it fixes its lower block πI_N. Thus each of its matrices is congruent to $I \bmod \pi$. This proves the containment `rowFixingLevel_le_levelOf` in the principal chosen level. All rational stabilizers of this subgroup are trivial by the free-level result of the earlier adelic chapter, since they are contained in rational stabilizers at the chosen level of the appropriate rank. The source uses this freeness when taking a class sum, so no inverse stabilizer appears in the comparison operator.

33.11 The loss of the Cartan product

Consider a rational prime ℓ not dividing the twist modulus, and the valuation ring $\mathcal{V}$ of a chosen coefficient embedding into $\mathbb{C}_\ell$. Let M be a nonzero rational integer belonging to the chosen product ideal (in the application take $M = g$). Multiplication by M clears denominators at each chosen split place. Because π times the whole inverse matrix of λ is integral, all columns of $M\lambda^{-1}$ are integral at every finite place. The particular bad-column assertion of Proposition 33.13 shows more: after the basis change K in the big column block, all but a of the columns of $\lambda^{-1}|_+K$ were already integral, without the factor M. Indeed $\pi G|_+K = 0 \bmod \pi^2$ outside the specified positions is equivalent to $G|_+K = 0 \bmod \pi$, and $\lambda^{-1} = \pi^{-1}G$ at the chosen places. At conjugate places the original inverse was integral, and the basis change can be taken trivial away from the chosen places. This proves the missing divisor-clearing assertion used in Chapter 32.

For comparison, let ρ_τ be the polynomial Schur action of shape with columns of multiplicities $\kappa_\tau(h)$, at an infinite weight label τ, now construed over $\mathcal{V}$ through its corresponding coefficient embedding. If A is a rectangular matrix with only columns in a set J failing integrality and MA_j is integral for $j \in J$, then the bound of Lemma 32.17 is valid for this rectangular polynomial Schur action as well. Indeed $A = (AD)D^{-1}$ with the diagonal matrix D on the column space; the integral polynomial action of AD is lattice-preserving and the action of D^{-1} is diagonal in tableau weights. Each symbol can occur at most once in a column, so its exponent is at most $\sum_h \kappa_\tau(h)$. Thus the loss is

$$|M|_\ell^{-|J|\sum_h \kappa_\tau(h)}.$$

To include the finite point h_+ of the big factor, factor its inverse column matrix as $\lambda^{-1} i_+ h_+^{-1} = (\lambda^{-1} i_+ K)(K^{-1} h_+^{-1})$, with i_+ the column inclusion. The compact matrix K^{-1} and its inverse are both integral, so its Schur action is an isometry for the lattice sup norm. The norm of $\rho_\tau(K^{-1} h_+^{-1}) y$ is consequently the gauge of the coefficient vector y at h_+, not the possibly much larger raw norm of y. This is the effect of the integral change of basis on the Weyl gauge in `norm_schur_inv_cols_le_weylGauge`, `New/Rows/CartanIntegrality.lean`. The proof of that declaration includes the multiplicative determinant twist; it cancels in this block because the prescribed Cartan product uses determinant exponent zero on the two factors and the product's exponent is the one dictated by its weight. We retain here the integral polynomial Schur model of the earlier adelic coefficient chapter.

Proposition 33.14 (Cartan gluing estimate)**.** *Let V_+, V_- be the coefficient modules of the big and small factors with multiplicities $\kappa_\tau^+(h), \kappa_\tau^-(h)$ as in the divided splitting. For their Cartan product $C(y, z)$ (the small factor with conjugate weights), at the point* $\mathrm{diag}(h_+, h_-)\lambda$ *the Weyl gauge is bounded by*

$$|M|_\ell^{-L} \, \|y\|_{h_+,\ell} \, \|\bar{z}\|_{h_-,\ell\circ\mathrm{conj}}, \qquad L = \sum_\tau \left(a \sum_{h<N} \kappa_\tau^+(h) + s_0 \sum_{h<s_0} \kappa_\tau^-(h) \right). \tag{33.12}$$

Proof. The big part has at most a bad columns, so the preceding estimate over all weight labels gives the first exponent. For the small part it is enough to use all its s_0 columns with denominator at most M. Since its action is by the conjugate matrix, the coefficient lattice to be used is that for the embedding composed with complex conjugation, applied to $\bar{z}$. One cannot silently commute the chosen embedding with conjugation. Apply the same diagonal-weight estimate to those s_0 columns, after the conjugate embedding and the inverse-transpose relation for the unitary factor. The source's Cartan product in the integral polynomial model is an integral bilinear map and compatible with block Schur actions: on the weight monomials this is the product of the two families of minors. Hence the gauge of the product is at most the product of these two bounds by the ultrametric inequality. The resulting exponent is L. □

This proves the finite divisor part of `weylGauge_cartanProduct_le`, with the prescribed divided splitting and its sign convention. The estimate is much less than bounding the full polynomial degree of the product by charging each box for the pole. For the large factor a bad column can occur strictly at most once in every Young column, which gives the coefficient a in place of the full rank N. This economy is necessary because the eventual comparison form must have small Archimedean norm after its denominator has been cleared.

33.12 Descent of a glued point form

We now trace how the indicator identity produces the two kernels in `DividedSplitting.theta_productForm`, `New/Rows/GluedFormTheta.lean`. Let O_θ be the auxiliary pair; the chosen split level is $M^2 Q$ where Q is the twist conductor and the modulus M corresponds to π. Denote by b, b' the characters of the two definite factors satisfying $bb' = O_\theta.b$, and by $K_+(g, h; y)$ the kernel on the big block with datum $T_S X = T_S F$, including its determinant twist at Q. Denote by $K_-(g, h'; z)$ the kernel on the small block with datum $X_- = Z$. Both use the prescribed harmonic polynomial in the conjugate coefficient vector. For a point form with large and small coefficient vectors y, z at $\mathrm{diag}(h, h')\lambda$ the normalized theta descent has finite datum

$$1_{\{\bar{G}X = \pi(0,Z)^T\}} \, \chi_Q(\det X_e)$$

at the ramified places; e selects the fixed rows for the twist. At M the finite matrix of the point has precisely the displayed support, since the integral argument of the point kernel is $\lambda^{-1}X = \pi^{-1}GX$. At Q the finite point λ is the identity, so the character factor is unchanged. The multiplier of the Weil action at M is

$$\beta_\lambda = \chi_{O_\theta.a}(\det_M \lambda)|\det_M \lambda|^{-r/2},$$

where $\det_M \lambda$ is the idele restricted to the selected places, and $\chi_{O_\theta.a}$ denotes the space character of the pair with the row rank r. This is `DividedLattice.weilScalar`. The exponent $-r/2$ is the Jacobian exponent for the Weil change of variables on that block. It is important that λ, $\det_M \lambda$ and the value of this multiplier are independent of the varying tame twist on Q.

By Proposition 33.13, the indicator is the sum over $Y \in \mathcal{H}$ of the product of the two block indicators, evaluated after the right column action by Y. Choose a lift $\widetilde{Y}$ at the selected places of the row group; it is congruent to 1 mod M and is trivial at Q. Consequently the determinant character of the twist has value one on it. The genuine finite naming identity for theta kernels changes the row-group point g to $g\widetilde{Y}$. The joint-harmonic polynomial of the Cartan product factors into the two polynomial kernels on the two orthogonal definite blocks. In the tensor-product realization of theta, the rational summation variables of the two blocks are independent, so the theta sum for the product datum is the product of the theta sums. Absolute convergence, for the rapid data at issue here, justifies multiplying them. These give the formula

$$\theta(\mathrm{productForm}_\lambda(h,h';y,z))(g) = \overline{\beta_\lambda \sum_{Y\in\mathcal{H}} K_+(g\widetilde{Y},h;y)K_-(g\widetilde{Y},h';z)}. \tag{33.13}$$

This explains the two kernels and the single Weil scalar in `DividedSplitting.theta_pointForm` and `DividedSplitting.theta_productForm`. The indicator identity and character compatibility have been proved above. In (33.13) we use the exact point-form evaluation and local Weil change-of-polarization law of the underlying normalized theta construction, and the compatibility of the chosen polynomial basis with block restriction. These are genuine model identities; a different normalization of the finite Weil action could change β_λ and would not be sufficient. The formal proof verifies them by the identities for `pairWeil_name_on`, `modDatum_factor`, `modKernelI_block` and `mixedCartan_mulVec_cartanProduct`; the needed statements are precisely these ingredients, not a new existence theorem for the glued lattice.

If $y(h)$ is the big-part coefficient vector depending on its class, let $C_i(g)$ be its coset kernel sum, summing $K_+(g,h;y(h))$ over the row-fixing level; the index i records S and the determinant-row choice. If c is a scalar, taking the class sum of the point forms with coefficient $\overline{cy(h)}$ and applying (33.13) gives

$$\overline{\beta_\lambda c}\sum_{Y\in\mathcal{H}} \overline{C_i(g\widetilde{Y})K_-(g\widetilde{Y},h';z)}.$$

Every sum here is finite, by definiteness of that group and finiteness of its class set. This is `DividedSplitting.theta_stdClassSum_productForm`. If it is integrated against a row form of the principal chosen level, then the right action by $\widetilde{Y}$ can be changed in the integral without changing that form, because $\widetilde{Y}$ belongs to this level. Thus the finite congruence sum contributes exactly $|\mathcal{H}|$ in the final inner product. This factor is a multiplicity, not a denominator, and the independence from the twist will enter the following comparison.

33.13 Rank-one reflections and the small-root descent argument

The comparison forms must not merely be integral. A good form on the large definite group can share a root of the Hecke polynomial of a lift without being in that lift. The denominator theorem applies to its complement only after the test vector is orthogonal to these false lifts. Two different facts are used to arrange this orthogonality: at a ramified place a reflection-sum annihilator promotes one vanishing descent to vanishing for all local data, while off level a small-root comparison propagates vanishing up the hyperbolic tower. We expand the algebraic parts of both.

Lemma 33.15 (Annihilator or standard frame)**.** *Let O be a discrete valuation ring with fraction field F, residue field k, and standard coordinate inclusion $J : O^r \hookrightarrow O^n$. Suppose $x : F^r \to F^n$ is injective. There is an integral matrix A annihilating x, with the following property. If all the entries of AJ reduce to zero in k, then there is $b \in \mathrm{GL}_r(F)$ such that xb is integral and congruent to J modulo the maximal ideal.*

Proof. By the lattice argument at the beginning of this chapter there is $c \in \mathrm{GL}_r(F)$ such that $P = xc$ is primitive. Choose a left inverse L of P over O, and set $A = PL - I_n$. Then $Ax = 0$. If $AJ \equiv 0 \bmod \varpi$, we have $\bar{P}\bar{L}\bar{J} = \bar{J}$. As $\bar{J}$ has rank r, so does the square $\bar{L}\bar{J}$, hence LJ is invertible over O. Take $b = cLJ$; the asserted congruence holds. If the assumption on AJ fails, a row of A gives an integral row l annihilating x for which the reduction of lJ is nonzero. □

This proves `AuxPair.exists_annihilator_or_congr_inclMatrix`, `New/SiegelWeil/AuxPairVanishing.lean`. It selects an annihilator of the entire local frame, not merely of a residue matrix. The distinction matters for the action on an arbitrarily small coset around x.

For a nonzero row $l : k^r \to k$, define the subgroup

$$H_l = \{\, I_r + ul : u \in k^r, \quad 1 + lu \neq 0 \,\} \subset \mathrm{GL}_r(k).$$

It is indeed a subgroup, since $(I + ul)(I + vl) = I + (u + v + u(lv))l$ and the inverse is $I - (1 + lu)^{-1}ul$. Its determinant is $1 + lu$, by the determinant lemma for a rank-one perturbation. Let χ be the relevant determinant character. In any representation ρ of this finite group,

$$\frac{1}{|H_l|} \sum_{h \in H_l} \chi(\det h)^{-1} \rho(h)$$

is the projection onto the simultaneous $\chi(\det h)$ eigenspace of H_l: translation in the sum gives the required eigenrelation, and on that eigenspace every summand acts as the identity. The numerator is the reflection sum `reflectionSum` of `Lib/LocalGL/Green/FrobeniusOrbits.lean`.

Suppose now l is the integral row provided by Lemma 33.15, with $lJ \not\equiv 0$. For every integral column u of length r and $h = I_n + Jul$ with $1 + lJu \in O^\times$, we have $hx = x$. The matrix h and its inverse are integral. Consequently h fixes every integral-box coset of the form $x + \varpi^k O^{n \times r}$, even for k large. Its reduction has J-columns within the image of J and induces $I_r + \bar{u}\overline{lJ}$ on them; every member of the latter group is obtained in this way. The other entries of the matrix are unconstrained in the level precisely because the indicated columns determine the block in the source's parabolic. If $D_x(y)$ is a period linear in a form y, and equivariance of the local theta action on this fixed coset gives $D_x(\rho(h)y) = \chi(\det h) D_x(y)$, summing the normalized reflection operator in this identity yields $D_x(y) = D_x(P_{H_l} y)$. Thus $D_x(y) = 0$ whenever the specified reflection sum annihilates y. At a twisting place where the action on the frame is by a scalar determinant character alone, the same argument can use a

single h for which $\chi(1 + lJu) \neq 1$: the map $u \mapsto lJu$ is onto the residue field, so nontriviality of that character ensures such an h. The period is then equal to a nontrivial scalar times itself and must vanish. For the row convention instead of the column convention use the transpose construction. This is the elementary reflection and nontrivial-character argument at M and Q in `AuxPair.descentsVanish_of_reflectionSum`.

If no such annihilator occurs, the frames are congruent to the standard frame after a change of source basis, by Lemma 33.15. The exact proof that the associated finite coset period reduces to the specified auxiliary descent is a further step in the same declaration; it uses the named-datum translation law, the source-equivariance law and decomposition of arbitrary Schwartz–Bruhat data into such local coset functions. The genuine normalizations in that transport, as well as the approximation in the real Fock datum, are needed to deduce `DescentsVanish` from these local calculations. The rank-one reflection calculation does not assume the conclusion of that theorem.

The small-root propagation through one or more planes. Let V be the definite Hermitian space of rank n, let W have rank r, and let $W_s = W \oplus H^s$ with rank $r + 2s$, for $s \leq p$. A descent of a form on V against a rapid Weil datum is a function on W_s. Suppose all the lower descents on W_{s-1} vanish. Integrating the descendent function on the radical along an adapted last line expresses its constant term as the lower descent, by the mixed-model identity for the zero term. Thus in forming the descendent function we may subtract that zero term. This is the identity `DescentsVanish.integral_sourceTerm_eq_zero`, which uses Iwasawa coordinates with respect to the adapted plane and the assumption that the finite datum is the vacuum off the allowed places. The induction in `descentsVanish_add_planes` requires a boundedness estimate for the difference between the original theta summand and its zero term. To see how the estimate is used, denote by $F(g, h)$ the theta kernel on the larger target and by $F_0(g, h)$ its radical average. If

$$\sup_{g,h} |F(g, h) - F_0(g, h)| \leq C$$

after the chosen short-vector and finite-coordinate reductions, then the function $D(g) = \int F(g, h)y(h)\, dh$ is bounded by $C\|y\|_{L^1}$, since the lower term integrates to zero. The same estimate with an integrable majorant in h in place of the constant C suffices. The boundedness derived in the source uses the short rational isotropic vector of W_s, the Witt theorem to choose its splitting, then translation of the named finite data and cancellation of the zero term. This is the application of `exists_norm_name_le` with exponent zero after subtracting the average. It relies on a genuine adelic estimate and remains a quantitative input in this reconstruction.

Once boundedness is known, the Hecke root calculation is completely explicit. Let w be a split spherical place of norm q, and suppose the original Hecke polynomial annihilates the form at an argument X of absolute value $q^{r/2+1-n}$, the value `AuxPair.testValue` used by the source. The Hecke transfer identity at the conjugate place (the adjoint in the integral pairing exchanges the two places) gives a product of trivial factors and a smaller Hecke polynomial on the descendent function on W_s at argument X' of modulus

$$|X'| = q^{(n-(r+2s))/2}|X| = q^{1-n/2-s}.$$

Here character values have modulus one after the appropriate unitary normalization. All the trivial factors have the form $1 - q^{j+(r+2s)/2}\zeta X$ with $|\zeta| = 1$ and $0 \leq j < n - r - 2s$. Their largest possible absolute norm of the term subtracted from one is

$$q^{n-r-2s-1+(r+2s)/2}|X| = q^{-s} < 1$$

for $s \geq 1$. Hence none of these factors vanishes. For the remaining polynomial,

$$q^{r+2s+1}|X'| = q^{r+s+2-n/2} \leq 1$$

whenever $n > 2r + 2p + 3$, since then $n \geq 2r + 2p + 4$ and $s \leq p$. The finite Hecke-operator contraction proof of Proposition 32.15 applies to this polynomial on bounded spherical functions on W_s. Thus the descent is zero. Iteration in s proves the propagation from level zero to level p, under the just-stated boundedness and genuine Hecke transfer inputs. It is important to use $s \geq 1$ at the trivial-factor cancellation; at $s = 0$ the initial vanishing is instead the special auxiliary-descent condition, not a strict inequality $q^0 < 1$.

The exact transfer identity is `hecke_transfer_of_equivariant`, `Lib/Weil/Unramified/HeckeTransfer.lean`, and its adjoint is `heckePolyOp_adjoint`. Combined with the reflection mechanism and the analytic boundedness estimate, the above proves the tower conclusion of `descentsVanish_hypSum` in the actual application to conjugate matrix coefficients of algebraic forms. The further period conversion in `stdClassSum_period_eq_zero`, `New/Rows/MemberOrthogonality.lean`, uses that a class sum on the smaller block paired with a translate of the algebraic matrix coefficient is an auxiliary theta period on the larger descent W_p. To derive it, expand a kernel vector in matrix coordinates, form the orthogonal block product of the corresponding two Weil data, and integrate against the conjugate coefficient of the given form. The twist level and the gluing conjugation property proved after Proposition 33.13 make that integrand fixed by the required compact subgroup. The precise global period identity identifying this integral with the class sum (the statement `period_eq_zero` of `Lib/Shimura/ComplexTheta/PeriodVanishing.lean`) is a further model input. It must preserve both levels and conjugate weights. No eigenform uniqueness principle at one Hecke root can substitute for it.

33.14 The comparison map and the precise scalar it sees

We return to the notation $u_\vartheta : U_j \to V_j(\vartheta)$ for a single good lift. The small definite group is denoted V^- and the large one is V^+. Its coefficient space has a fixed positive Hermitian Gram matrix, the one of the integral Weyl model. Let I be the finite set of member indices, and $\mathcal{C}^-$ the class set of V^- at the squared chosen level M^2. Let T be the Hilbert direct sum of the small coefficient spaces indexed by $I \times \mathcal{C}^-$. This is `Data.Tables` of `New/Rows/ComparisonForms.lean`; T is independent of the twist conductor. Write $\Xi(i, c)$ for a vector of this table at (i, c).

For each i take the glued point λ_i of Proposition 33.13 and its multiplier β_i. Let $y_{\vartheta,i}(h)$ be the large-factor line multiplier vector, and c_* the line scale of the source (one fixed nonzero scalar). The comparison map $g_\vartheta : T \to V_j(\vartheta)$ is the following sum of actual point forms:

$$g_\vartheta(\Xi) = g^L \sum_{i,c} \sum_{h \in \mathcal{C}_i^+} \mathrm{productForm}_{\lambda_i}(h, c; \overline{c_* y_{\vartheta,i}(h)}, \Xi(i, c)). \tag{33.14}$$

The set $\mathcal{C}_i^+$ is the class set of the row-fixing level of the big group associated with S_i. All its stabilizers are trivial, as proved above. The scalar g^L, with L as in (33.12), is the fixed denominator clearing of the Cartan product. This is exactly `Data.comparisonForm`. For all appropriate coefficient places, the norm bound on the large factor of the form is the integral-line-scale condition `lineMultiplier_integral_and_small` of the source. Specifically it asserts that $\overline{c_* y_{\vartheta,i}(h)}$ has gauge at most one at every coefficient embedding at a prime prime to $Q(\vartheta)$; in the untwisted case the squares of the Archimedean conjugate coefficient norms are bounded by $\exp(rB_0)$, with B_0 the small budget of Chapter 31.

By Proposition 33.14, every summand of (33.14), including the factor g^L, is integral if the conjugate table coefficient $\overline{\Xi(i,c)}$ has gauge at most one at the conjugate embedding for its class c. The point-form operation itself preserves such integral vectors: it puts their translated coefficients on one adelic class and vanishes on the others, with only an integral compact action and a unit-modulus finite level character on its support. All the point forms are well-defined because of the prescribed level conditions in the auxiliary pair. Therefore the finite sum in (33.14) is integral. This proves `comparisonForm_mem_integralForms` from the line-scale assertion and the genuinely constructed divided lattice, not from an abstract integral representation of the comparison map.

Write $C_{\vartheta,i}(g)$ for the coset kernel sum corresponding to the large line vectors in this comparison. The essential geometric identity in `Data.cosetKernelSum_eq_fNatural`, `New/Rows/Separation.lean`, is

$$F_i(g) = \eta_\vartheta C_{\vartheta,i}(g), \qquad |\eta_\vartheta|^2 = C/\Gamma_j(\vartheta), \tag{33.15}$$

for a fixed family of functions F_i (the `fNatural`) independent of ϑ. The number C is a fixed real constant, not initially known nonzero in the proof. The identity is an application of complementary-rank exchange to the large carrier and to the flat line kernels. Its actual construction also uses the determinant polynomials of `New/Rows/Carrier.lean`; the three line-pair products in Γ_j must appear exactly, since they are the special values that repay the idle directions. We retain this exact period equality as an input.

For a table subfamily indexed by $c \in \mathcal{C}^-$, let $T_i(\Xi)(g)$ denote the small-block sum of conjugate kernels (`Data.tableTheta`), with datum equal to the indicated small coordinate inclusion at M^2 and the prescribed polynomial of the conjugate coefficient. It too is independent of ϑ. Integrate the descent identity (33.13) first against a good row $F \in U_j$, then use (33.15). The class sum over h gives $C_{\vartheta,i}$, the sum over c gives $T_i(\Xi)$, and the congruence translates each give the same integral by principal-level invariance of F. Since the inner product is conjugate-linear in the first variable, this yields

$$\bar\eta_\vartheta \langle F, u_\vartheta^* g_\vartheta \Xi\rangle = |\mathcal{H}| g^L \bar c_* \sum_i \overline{\beta_i} \int \overline{F(g')F_i(g')}, T_i(\Xi)(g'), dg'. \tag{33.16}$$

This is the identity `inner_adjoint_comparisonForm`. Every factor on the right is independent of ϑ. The genuine theta point-form identities isolated in the preceding subsection assure the indicated normalization; the finite grouping and phase cancellation have now been shown.

One additional separation theorem for the fixed functions is needed:

$$\Big[\ int\overline{FF_i}\, T_i(\Xi) = 0 \text{ for all } i, \Xi\ \Big] \quad\Longrightarrow\quad F = 0. \tag{33.17}$$

This is `Data.eq_zero_of_integral_fNatural_tableTheta`. A crucial part of its proof is the statement `exists_fNatural_approx` in `New/Rows/NaturalFormNear.lean`: for $n_* =$ `nStar`, some function K_0 with nonzero $K_0(1)$ can be uniformly approximated by linear combinations of translates $F_i(gy)$ at points where the first adelic minimum of the moved lattice exceeds its specified cut-off. The error is at most $e\max(1,\|y\|)^{6n_*}$, with $4e(5b)^{7n_*} + e < |K_0(1)|$. The function is formed from the nonzero carrier theta lift and holomorphic minors; reproduction and a separate isolation estimate convert orthogonality to a contradiction. The archimedean localization and quantitative approximation, including the first-minimum threshold, are necessary to use this statement and remain inputs. This separates the problem of detection by actual theta products from the linear algebra of the table map.

Proposition 33.16 (Scalar and surjectivity of the comparison)**.** *Assume the genuine identities* (33.15)–(33.16)*, separation* (33.17)*, and* $\dim U_j > 0$*. Put* $A = u_1^* g_1$*. Then* $A : T \to U_j$ *is*

surjective and

$$u_\vartheta^* g_\vartheta = \gamma_\vartheta A, qquad \gamma_\vartheta = \overline{\eta_1/\eta_\vartheta}, \qquad |\gamma_\vartheta/\gamma_1|^2 = \Gamma_j(\vartheta)/\Gamma_j(1).$$

Moreover, suppose for every false lift z at an allowed split place the small-block periods of the table $(g_\vartheta)^* z$ all vanish, i.e. $T_i((g_\vartheta)^* z) = 0$ for every i. Then $g_\vartheta A^* y$ is orthogonal to these false lifts for every $y \in U_j$.

Proof. If one $\eta_\vartheta = 0$, (33.15) makes every $F_i = 0$. Then (33.17) makes every $F \in U_j$ zero, a contradiction. Thus every $\eta_\vartheta \neq 0$ and the constant in (33.15) is positive. Comparing (33.16) for ϑ and 1 gives $\overline{\eta_\vartheta} u_\vartheta^* g_\vartheta = \overline{\eta_1} A$ by nondegeneracy of the inner product on U_j. The formula for the absolute value is immediate from (33.15).

If F is orthogonal to $\operatorname{ran} A$, the right side of (33.16) is zero on every table. Choose a table supported at one member index i and allow its small-class coefficients to vary freely. All other $T_{i'}$ then vanish. Since $|\mathcal{H}| g^L \bar{c}_* \bar{\beta}_i \neq 0$, the integral in (33.17) is zero for every such table. By separation $F = 0$; finite dimensionality of U_j gives surjectivity.

For a false lift z, let $v = g_\vartheta^* z$. By hypothesis $T_i(v) = 0$ for all member indices. Equation (33.16) at the trivial twist shows $Av = 0$, hence $\langle z, g_\vartheta A^* y \rangle = \langle A g_\vartheta^* z, y \rangle = 0$ for all y, as desired. □

The last hypothesis in Proposition 33.16 is exactly the consequence of the tower-vanishing and period-conversion assertions in the preceding subsection. Indeed a false lift belongs to the good idempotent, the kernel of the polynomial at the small test root, and the orthogonal complement of $u_\vartheta U_j$. The first property gives the reflection-sum annihilators, the third the vanishing auxiliary descent. The class-sum period conversion then kills tableTheta$((g_\vartheta)^* z)$. This is `tableTheta_adjoint_eq_zero` in the source. The proposition shows why testedness is asserted on the special range $g_\vartheta A^* U_j$, not on the entire form $g_\vartheta(\Xi)$ for arbitrary tables. The proof of `exists_comparison_forms` has now been reduced to explicit finite construction, norm, and the genuine fixed-carrier and orthogonality statements.

33.15 The Archimedean budget of the comparison operator

The remaining size requirement for this construction, at the trivial twist, is $h_- \|\sigma(g_1(\Xi))\|^2 \leq e^{2rB_0} \sum_{i,c} \|\sigma \Xi(i,c)\|^2$, where h_- is the hook factor of the small coefficient space. We detail why the large denominator exponent L is still affordable here. The untwisted line-scale hypothesis says that each big coefficient vector has squared conjugate norm at most e^{rB_0}. Cartan multiplication has norm at most the product of coefficient norms in the integral Fock-Weyl realization, with the same weights as the gauge estimate; this follows either from tensoring the Fischer forms and taking the Cartan component or from the polynomial Gram calculation in the adelic coefficient chapter. The point-form norm at any class is at most M_* times that coefficient norm, with $M_* = \text{chosenGroupOrder}$, because the point-form construction sums over a stabilizer of order dividing M_*. The finite character values have modulus one also after every field automorphism, and the Schur matrices are mapped with the conjugate coefficient coordinates rather than with a new choice of bases.

Let H_V be the number of target classes at the untwisted level, H_I the number of member indices, H_- the number of small definite classes and H_+ an upper bound on the number of classes of each big row-fixing level. Summing first over h, then over (i,c) in (33.14), the triangle inequality and Cauchy–Schwarz give the pointwise bound

$$g^L M_* e^{rB_0/2} H_+ \sqrt{H_I H_-} \left(\sum_{i,c} \|\sigma \Xi(i,c)\|^2 \right)^{1/2}.$$

The squared class-sum norm of the resulting form is bounded by its squared maximum times H_V (class weights are at most one). Thus

$$h_-\|g_1(\Xi)\|_\sigma^2 \le h_- H_V H_I H_- H_+^2 M_*^2 g^{2L} e^{rB_0} \sum_{i,c} \|\sigma\Xi(i,c)\|^2. \tag{33.18}$$

These bounds hold for the fixed conjugate coefficient-space Hermitian form under the field automorphism, not for the image under the automorphism of a positive real norm inequality. The former is what is required in the compressed determinant.

We check the remaining logarithm along the admissible family of Chapter 54. The class bounds used in `normSqAt_comparisonForm_le` in `ComparisonForms.lean` are $\log H_V, \log H_-, \log H_+ \le b$ eventually; the analogous bound holds for the rank $6m+1$ flat class set occurring in H_I. Indeed each is bounded by the class-count estimate at the principal level M^2 in the relevant rank (all ranks are bounded by a power of x), and the row-fixing and comparison levels contain that principal level. The logarithm of the finite residue ring at M^2 is $2f \log g$, with f the number of infinite places. By `Data.MemberIndex` and `Data.LineChoice` the other choices in H_I consist of a matrix with $3m$ rows and $N = \operatorname{rk} V^+$ columns over this ring, and for every infinite place a list of $2m$ row choices among $6m+1$ repeated $h_2 - \lfloor (t+1)/2 \rfloor$ times. Their logarithmic cardinalities are bounded by

$$6mNf \log g + 2mfh_2 \log(6m+1) = x^{O(1)}$$

uniformly along that family. Since the small coefficient multiplicities are each h, as in `ComparisonForms.lean`, the total size of its Young diagrams and the logarithm of their hook number h_- are bounded by fixed powers of x. The group-order denominator M_* also has logarithm bounded by a fixed power of x, by the free-level bound of the adelic chapter.

For the nonpolynomial part of L, the source's `card_detIndex` expresses $\sum_{h<N} \kappa_\tau^+(h)$ as the determinant-index count minus the reduced half gap; the identity `card_detIndex_add_bigCharExp` bounds this count by $w_j(\tau) \le 2b$ eventually. The negative-column sum is at most $s_0 h$ per weight label. Since $a = 3m$ in the actual splitting, this gives

$$L \le 6fmb + fs_0^2 h.$$

Here $a = 3m$ refers to the number of divided row coordinates, not the prime-sum cutoff (which is still denoted by a elsewhere). The second term is polynomial in x. Also $\log g \le 22 \log b$. Write $Y = \log b = x^\varepsilon$, and for this estimate denote the large prime-sum cutoff by a_{pr}. Then

$$rB_0 = \frac{r}{k} \frac{f a_{\mathrm{pr}} Y^3}{m} \quad \ge \frac{fmbY^3}{4},$$

using $r \ge k$, $a_{\mathrm{pr}} \ge xb$ and $m^2 \le 4x$ for large x. Therefore $2L \log g = O(fmbY) = o(rB_0)$. All the other logarithms in (33.18) amount to $O(b) + x^{O(1)} = o(rB_0)$ by Lemma 54.1. Consequently their sum is below rB_0 for all sufficiently large x. We recover precisely

$$h_-\|g_1(\Xi)\|_\sigma^2 \le e^{2rB_0} \sum_{i,c} \|\sigma\Xi(i,c)\|^2. \tag{33.19}$$

This proves the budget calculation in the norm bound of the comparison forms. It is an asymptotic verification on the cofinal admissible family, which is enough for the limiting argument. It uses the fixed untwisted line-scale norm estimate and the indicated class-count, tableau and integral point-form bounds. In particular it does not infer such a norm estimate uniformly at all twist conductors from the bound for the trivial conductor.

33.16 Two further finite calculations for the rank-one values

In the height estimate for $\Gamma_j(1)/\Gamma_j(\vartheta)$ there are two types of rank-one input. One is the arithmetic size of a normalized integral kernel coefficient; the other is the identification of the pairing of a test form with an actual Hecke value. The first is a delicate attainability result (the depth indicator in Chapter 31). The second involves the exact normalization of class sums, which we detail here.

Fix a pair $p = (\nu, z)$ with target rank $2z$ and the remaining data of `LValueData`. Let Q denote a fixed squarefree twist modulus and consider its tame weight-zero characters ϑ. Write I_Q for the finite indexing set consisting of the chosen torus class set and the local groups of units at Q. The class sum defining $\Xi_\vartheta = \mathrm{trivialLift}(H, \vartheta)$ can be expanded by decomposing each principal-level torus class into these finitely many translates. If g_i, $i \in I_Q$, denote these translates, the finite character occurs only in the factor $\vartheta(\det g_i)$ of each kernel coefficient. Thus, after fixed normalization, write

$$\Xi_\vartheta = \sum_{i \in I_Q} \vartheta(g_i) K_i \tag{33.20}$$

with K_i the prescribed point kernel at modulus Q independent of ϑ. A choice of inverse of the character just interchanges i, j in the formulas below. This is the class expansion `trivialLift_apply`; in the source's proof the nontrivial character factors are those of the Weil determinant multiplier on the split torus. In particular the point-kernel data must be kept at the fixed modulus Q, including for the trivial character viewed at that modulus. One may not replace K_i by the point kernels of a varying truncated level.

Put $d_p = \mathrm{denomBound}$ as in Chapter 31 and $M_* = \mathrm{chosenGroupOrder}$. The normalized value is $\mathcal{A}_p(\vartheta) = M_* d_p^2 \|\Xi_\vartheta\|^2 / N(Q)^{2z-1}$. On inserting (33.20),

$$\mathcal{A}_p(\vartheta) = \sum_{i,j \in I_Q} c_{ij}\, \vartheta(g_j/g_i), \quad c_{ij} = \frac{M_*}{N(Q)^{2z-1}} \langle d_p K_i, d_p K_j \rangle. \tag{33.21}$$

This uses $\overline{\vartheta(g_i)} = \vartheta(g_i)^{-1}$, valid since the characters have finite order. It follows without assuming that an arbitrary automorphism of $\mathbb{C}$ commutes with complex conjugation.

The reason the coefficients in (33.21) are integral at primes prime to Q is as follows. For every such embedding the two point kernels multiplied by d_p have gauge at most one, by `weylGauge_pointKernel_le_one` of `New/Exchange/TriangleLevelExponent.lean`. This applies both to the embedding and its composition with conjugation. The Gram pairing of two such coefficient vectors is integral in the integral Gram model (one factor is evaluated at the conjugate embedding). The class-sum pairing divides each contribution by the order of its finite rational stabilizer. These orders divide M_*, and therefore M_* clears every denominator in the pairing. Finally the conductor power $N(Q)^{2z-1}$ is a unit at the given embedding. Hence every c_{ij} belongs to the corresponding valuation ring. This proves the finite-measure calculation in `isIntegralMeasure_intAlgPart`. It uses the point-kernel integrality and the fixed-modulus expansion in the genuine Weil model, and not an interpolation theorem for special values. Once the exact Rallis norm formula for the normalized characters is known, it is an integral measure for the removed Hecke L-values as well.

For completeness we explicate the three-class-sum test formula. In the following calculation the flat blocks have all coefficient degrees zero. Suppose a factor of even rank m_0 has flat theta lift Ξ_0, normalization d_0, and conductor Q_0 prime to the considered coefficient prime. Let $\mathcal{C}$ be the prescribed finite torus class set and let $\mu_c = 1/|\operatorname{Aut}(c)|$ be its weights. Write $\mathcal{N}_c = \sum_c \mu_c$. A vector-valued test family P_h, indexed by the classes h on the flat target, is

combined with its flat lift by the operator

$$\mathcal{X}(P) = \frac{M_* d_0^2}{N(Q_0)^{m_0-1}} \sum_h \mu_h \, \Xi_0(h) P_h.$$

The placement of a complex conjugate in the class sum is determined by which entry of a Gram inner product is linear; the displayed placement agrees with using the flat Gram functional $\langle 1, \Xi_0(h)\rangle$ in a linear functional of P. This is the operation `Data.Pair.xiSum`. It preserves integrality of P_h at the coefficient prime: $d_0\Xi_0(h)$ is integral by summing the integral normalized point kernels (and accounting for any source stabilizer by its fixed normalization); one remaining factor of d_0 is integral, $M_*\mu_h$ is integral, and the conductor factor is a unit. In the flat case the Gram coefficient factor is just a scalar, so this calculation is componentwise in the valuation ring.

Let L be a complex linear functional of the test vectors and suppose $L(P_h) = C_1 K_{g_0}(h)$ with K_{g_0} the corresponding flat kernel coefficient complementary to Ξ_0. The reproducing class-sum identity in the source is

$$\mathcal{N}_c \kappa_0^{m_0} \sum_h \mu_h \, \Xi_0(h) K_{g_0}(h) = \|\Xi_0\|^2, \tag{33.22}$$

after the included twisted central normalization of K_{g_0}. Here κ_0 is the vacuum scale and the torus level is the *chosen* level, not the level with additionally varying Q_0. To see the mechanism of (33.22), expand Ξ_0 as the sum of translates of a single point kernel on torus classes. The normalized flat lift is fixed by the twisted central action. Pairing each translated kernel with Ξ_0 therefore gives the same number, and their weighted sum is the squared norm of Ξ_0. Canceling the total class mass and the vacuum power gives the displayed identity with the correct supplementary norm factor. The exact twisted center action and normalized point-form adjunction are `inner_trivialLift` and `twistedCentralOp_trivialLift` used in `Pair.smul_apply_xiSum` of `New/Rows/HeckeValueHeight.lean`. They are genuine theta identities; the class-sum argument proves the scalar extraction once they hold. It follows directly that

$$\mathcal{N}_c \kappa_0^{m_0} L(\mathcal{X}(P)) = C_1 \mathcal{A}_{p_0}(\vartheta_0).$$

For a rank-one pair of rank $2z$ in the application choose three auxiliary flat blocks of ranks $2z-20, 4, 4$ and a remaining block of size twelve. The sum of these ranks and twelve is $2z$. The finite residue datum fixes the appropriate columns at the chosen places and has determinant character the product of the four block characters, as required by the direct-sum Weil law. Choose the degree distribution $\alpha_{\tau,i}$ of the twelve-jet, so that $\sum_i \alpha_{\tau,i} = d_\tau$. When a power of a linear expression of degree d_τ is expanded in these coordinates, the coefficient of the monomial of multi-index α_τ is

$$\binom{d_\tau}{\alpha_\tau} = \frac{d_\tau!}{\prod_i \alpha_{\tau,i}!}.$$

Equivalently this is the ratio between the Fock Gram pairing of the degree-d_τ vector and that of the single monomial under the chosen polynomial normalization. Real tensor multiplication over the places multiplies these factors. On the flat blocks the monomial is 1 because their weights are zero. Therefore the initial point form of this monomial has, by the block product identity, kernel pairing equal to this multinomial product times the product of the three flat kernels and the twelve-coordinate kernel. Apply the class-sum operation $\mathcal{X}$ once for each flat block, with the other factors held fixed. Each application is linear and changes one flat kernel

into its normalized value at the cost of the factor on the left of (33.22). Since all three use the chosen torus level and the sum of their ranks is $2z-12$, the final equality has exactly

$$\mathcal{N}_c^3\kappa_0^{2z-12}\langle u,K_g\rangle=\left(\prod_\tau\frac{d_\tau!}{\prod_i\alpha_{\tau,i}!}\right)\mathcal{A}_{p_1}\mathcal{A}_{p_2}\mathcal{A}_{p_3}\,K_\alpha^{(12)}(g). \tag{33.23}$$

The three normalized values are evaluated at the chosen auxiliary twists matched with the twelve-coordinate jet character, two of them trivial or flat auxiliary twists as in the source. This is the factorization at the end of `exists_test_form_inner_kernelForm`. In particular the exponent $2z-20+8=2z-12$ of the vacuum factor represents only the three flat blocks; charging the twelve-coordinate kernel again would give an incorrect power. The construction gives an integral test vector if the normalized monomial vector of the twelve-block has gauge at most one and the indicated block-column point data are admissible. The source constructs such a vector by the integral harmonic-polynomial identity `exists_int_vector_harmonicPoly_eq`. The orthogonality of the vector to the false rank-one lifts is the small-rank application of the preceding reflection-sum and larger-rank period vanishing. These statements about the actual representation and period support are still needed to obtain testedness; the class-sum and multinomial computation alone does not imply it.

We recall why the depth acquisition is a separate problem. The chosen scale of `Data.depth_indicator_jet`, say t_0, must be nonzero, small under every Archimedean conjugate, and good at all finite embeddings in the sense that at each one some degree distribution and some integral point produce an attainable jet coefficient. If one has that input and (33.23), the finite character selection and multiplicative selection in Lemma 31.20 produce the required K inequality of the source. The first selection uses Fourier inversion on the finite torus class group, giving one twist at a chosen point whose Fourier coefficient is at least its original value. The second uses a product of the flat values over the full character group. Specifically for functions a,b on that character group, take a root V with $V^{|G|}=\prod_{\eta\in G}a(\eta)b(\eta)$. For a prescribed product character χ, multiply $a(\chi\eta^{-1})b(\eta)$ over $\eta\in G$; this is exactly $V^{|G|}$. Hence for every coefficient embedding one factor has norm at least $|V|$, while for every Archimedean conjugate a uniform upper bound on all the a,b directly bounds $|V|$. A priori the selected factors may vary with the coefficient embedding, but the height witness V is fixed. This is the reason that the witness in the height proof does not depend on the chosen test form u.

The remaining novel ingredient is precisely the attainability assertion of `depth_indicator_jet`, which uses the CM inverse-series and Frobenius packing argument. An estimate that bounds every coefficient above cannot provide a nonzero $K_\alpha^{(12)}(g)$ of the necessary local size. The uniform exact normalized Rallis pairing for the rank-one values and their conjugate norm formula (`LValueData.map_trivialLift`, `Data.Pair.log_norm_map_norm_sq_trivialLift`) must also be supplied by the actual CM theta model. Equations (33.21) and (33.23) show what has been reduced to elementary finite arithmetic and where these two sources of information enter the height argument. With those genuine inputs, plus the testedness and the rank-one projection bound, the calculation in Chapter 31 gives the height saving of one direction.

33.17 Summary of the strengthened reconstruction in C

The following list distinguishes the completed constructions of this chapter from their remaining geometric inputs.

1. The local algebraic factorization through column partners, including the two-sided lattice coset representatives and compactness proof, is now established (Proposition 33.2); the synthesis of a stable space from unimodular coefficient vanishing is proved in Proposition 33.4.

The required automorphic equivariance and actual geometric coefficient vanishing must hold on the normalized theta difference, not just on the finite data.

2. The real smooth local Witt map, simultaneous moment approximation and smooth-section estimate are proved, and the determinant-block calculation gives the positive scalar of the cone integration. The exact top-period identity, normalized mixed-model line constant term and its convergence and global invariant measures still have to be supplied for the compatible data. The scalar induction and double-exchange argument explain the nonzero common normalization but rely on these genuine global identities.

3. The finite tidy square transformation, including the removal of the frame-orbit denominator by covariance, is now proved; the terminal interlacing, signatures, and square-cost recursion are explicit. The genuine real square law with its kernel height and type-convolution inputs, and the global quantitative restriction map reaching all initial rows, are still used from the specified declarations. The latter is needed before the finite transport can establish the crude-row theorem.

4. The divided lattice gluing, including its realization at the split places, level conjugation, indicator identity, and sharp bad-column denominator bound are proved (Propositions 33.13 and 33.14). The finite assembly and scalar/surjectivity argument for the actual comparison operator are expanded, as is the Archimedean budget calculation. They use the genuine block theta evaluation, line-scale estimate and fixed-carrier separation, and the testedness requires the exact tower-vanishing period conversion.

5. The reflection sum and the small-Hecke-root induction have been made explicit; the latter still needs boundedness after subtraction of the correct constant term, the genuine Hecke transfer, and its period conversion. The finite character measure and flat-block test pairing arithmetic are reconstructed subject to the stated normalized theta and point-form identities. The depth-indicator attainability and normalized rank-one Hecke values and their Galois comparisons remain the specialized arithmetic inputs.

These are concrete advances beyond identifying the modules in which the original proofs occur. In particular the finite gluing and square transformations do not require a rational theta theorem or an assumption that the desired integral bounds already hold. The unresolved parts, on the other hand, are not removable merely by citing a classical theta correspondence in a stable range. They are precisely the analytic and normalized integral comparison statements for the genuine data of the proof.

Chapter 34

Closing the period, transfer, and depth calculations

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 35, 36 and the final audit in Chapter 57.

This chapter concerns the specific normalizations and nonvanishing estimates needed in the lower bound. We separate a calculation established below (PROVED) from an appeal to a published theorem with matched hypotheses (REFERENCE), and from a dependency for which neither has been supplied (OPEN). A calculation proved relative to an already identified model identity retains the dependency on that identity. No finite-dimensional proof of a determinant comparison can replace a theorem identifying the automorphic operators being compared.

34.1 Vanishing at a nonnegative nondegenerate centre

Here is the local-to-global argument for `centreCoeff_eq_zero_of_not_negative` in `New/Exchange/NonnegativeVanishing.lean`. Use the splitting of X over Z along b lines. Write p_Z, q_Z for the numbers of positive and negative signs, respectively, and let Y have $q_Z + b$ negative signs and at least one positive sign. Let $\mu_\tau < p_Z$ at every infinite place and let $\delta = \nu\upsilon^b$. Assume integrability of the terms of the coefficient, in the sense of `HasIntegrableTerms`. If T is nonsingular Hermitian and is not totally negative definite, the coefficient of the theta integral at $\jmath T$ is zero.

Proof. The central unipotent acts on a mixed-model summand at a frame x by the additive character paired with $x^* J_Y x$. Integrating this character against the opposite character at T over the compact adelic additive quotient leaves only frames with $x^* J_Y x = T$. The assumed integrability permits this integration term by term. If there is no such rational frame, the coefficient vanishes. Otherwise Witt's extension theorem acts transitively on the rational frames with this Gram matrix. Writing H for the stabilizer of one such frame, unfolding the orbit sum gives the integral on $H(F)\backslash U(Y)(\mathbb{A}_F)$ of the corresponding mixed-model term. These are the reductions `centreCoeff_integral_eq_tsum` and `centreCoeff_eq_integral_stabilizer`. One needs their compatible quotient measures, but no particular overall constant for vanishing.

Choose τ such that T_τ has a positive eigenvalue. The image frame w has a positive vector. An element of the real unitary group takes that vector, after adjusting its length, to a positive coordinate vector. Apply this element to the frame and write the resulting frame again as w. Its projected image in the negative coordinate subspace Y^- has rank at most $b - 1$: a

nonzero combination of its columns is in Y^+. Given an arbitrary coherent-state coordinate $y \in \mathrm{Hom}(Z_\tau, Y_\tau)$, there are at most q_Z further orthogonality constraints on a vector of Y^-, namely orthogonality to the columns of y indexed by negative coordinates of Z. Since

$$\dim Y^- - (b-1) - q_Z = 1,$$

there exists a unit vector r satisfying both sets of orthogonality constraints. With $P = rr^*$ the orthogonal projection in the ordinary positive coordinate metric, put $c_\eta = 1 + (\bar\eta - 1)P$ for $|\eta| = 1$. This element of the maximal compact fixes w, acts trivially on the positive block, and has determinant $\bar\eta$ on the negative block. It fixes the other infinite places.

For the Fock vacuum v_0, the compact weight formula in `realPairWeil_polyVector` yields

$$\omega_\mu(1, c_\eta)v_0 = (\bar\eta)^{\mu_\tau - p_Z} v_0 = \eta^{p_Z - \mu_\tau} v_0.$$

Thus covariance of the Heisenberg representation gives, for $v_y = \pi(y)v_0$,

$$\omega_\mu(1, c_\eta)v_y = \eta^{p_Z - \mu_\tau} v_{\,y + (\bar\eta - 1)Py}.$$

In the sign convention of the source, the holomorphic coordinate $\mathcal{H}$ conjugates an entry when the two signs agree and leaves it unchanged otherwise (`holCoord_eq`). The only possibly nonzero entries of Py are in negative target rows and positive source columns. Consequently $\mathcal{H}(y + (\bar\eta - 1)Py) = \mathcal{H}(y) + (\bar\eta - 1)Py$. The Euclidean length of the whole coordinate y is unchanged by c_η. The coherent-state evaluation formula is

$$v_y(z) = e(z)e(y)\exp(\overline{\mathcal{H}(y)} \cdot \mathcal{H}(z)),$$

with the vacuum Gaussian e and the chosen Heisenberg central character. Hence, for fixed z, evaluation of $\omega_\mu(1, c_\eta)v_y$ is the boundary value of an entire function of η divisible by $\eta^{p_Z - \mu_\tau}$. That exponent is positive, so the circle mean is zero. The vector-valued circle mean is zero, since the evaluations separate the coherent-state model. Coherent states are total: a vector orthogonal to all of them has identically zero reproducing holomorphic function.

Return now to the unfolded integral. The subgroup C obtained by conjugating the real compact stabilizer of w back to the original frame is a compact subgroup of $H(\mathbb{A}_F)$. It has Haar probability measure. Disintegrate the quotient integral by H, and average the mixed-model term over this subgroup; the modulus of the right action of a compact subgroup on Haar measure is one. The source-term equivariance for a target element fixing x (the source Levi equal to the identity) expresses this average, at fixed remaining data, as a continuous linear functional on the real Fock representation invariant under the compact stabilizer. For every coherent-state vector v_y its value equals the value of the circle average constructed above, hence is zero. Such a functional is zero by totality. Therefore every compact mean in the quotient integral is zero. The absolute integrability assumption justifies the disintegration and averaging. □

A separate circle is chosen for each coherent vector: there is no claim that one circle annihilates all the real K-types. The proof establishes the actual nonsingular coefficient vanishing, conditional on the mixed-model coefficient and equivariance reductions and the stated integrability.

34.2 The two line slices and the conductor factor

Consider the large-side step from W, V to W', V' obtained by adjoining a split line-pair to both, with $\dim W = 2d$, $\dim V = r$, and $e = r - d$. We use the notation of `Lib/SiegelWe`

`il/Tower/ConstantTerm.lean`, `LineGaussSum.lean`, and `LargeSideConstantTerm.lean`. The two slices are indexed by $\beta = 0, 1$; they evaluate the mixed variable on the two opposite isotropic coordinates, Fourier-dual to each other. The residual Weil model is the one for W, V.

Let S be a set of split places with no conjugate pairs, $Q = \prod_{w \in S} q_w$, and let χ_w denote a nontrivial multiplicative character of the residue field at w. The standard row datum is at the split modulus S. It is extended by zero when its determinant is a nonunit. All multiplicative Haar measures on local unit groups are normalized to give them volume one; for additive Gauss sums the residue ring has additive volume one. Fixed changes of the latter convention, and fixed frame or vacuum scalars, are retained in the constants below.

Lemma 34.1 (The local Gauss factor)**.** *Let $\mathcal{O}$ have residue cardinality q, uniformizer π, and ψ have conductor $\mathcal{O}$. If χ is a nontrivial residue-field character regarded as a function on $\mathcal{O}$ vanishing on nonunits, then*

$$I(a) = \int_{\mathcal{O}} \chi(z)^{-1} \psi(az)\, dz = \begin{cases} q^{-1}\chi(\pi a) G, & v(a) = -1, \\ 0, & v(a) \neq -1, \end{cases}$$
$$G = \sum_{u \in k^\times} \chi(u)^{-1} \bar{\psi}(u), \qquad |G|^2 = q.$$

Here $\bar{\psi}(u) = \psi(\pi^{-1}\tilde{u})$ and the integral is understood with value zero at z nonunit. In particular $|I(a)|^2 = q^{-1}$ on its support.

Proof. For $v(a) \geq 0$ the integral is the average of a nontrivial character, zero. For $v(a) < -1$, on each unit residue class the integral over $\pi\mathcal{O}$ of the additive character is zero. For $v(a) = -1$ decompose into the $q - 1$ unit residue classes and substitute $u \mapsto (\pi a)^{-1} u$. Finally,

$$|G|^2 = \sum_{t \in k^\times} \chi(t)^{-1} \sum_{v \in k^\times} \bar{\psi}((t-1)v) = (q-1) - \sum_{t \neq 1} \chi(t)^{-1} = q.$$

□

In the line slice of the standard determinant datum the adapted integral frames turn the determinant, up to an integral unit, into the product of three diagonal determinants: the line coordinate at w, the remaining standard row determinant, and the second line coordinate being Fourier integrated. The off-diagonal blocks are unconstrained integral variables. Consequently the surviving determinant character on the first line is the inverse unit character of a_w; the integration on the second line is Lemma 34.1 at argument $\bar{a}_w$. Equivalently, in the notation $\alpha = \mathrm{scaleIdele}$ and $a = \xi/\alpha$, the finite part of the slice is the lower row datum times a scalar

$$\lambda_\beta G_S \, 1_{\text{integral off } S \cup \bar{S}}(a_f) \, \frac{\chi^{-1}(a_S)}{\chi^{-1}(\varpi \bar{a}_S)}, \qquad |G_S|^2 = Q^{-1}. \tag{34.1}$$

Each unit character in this expression is zero off its unit support, and the displayed quotient means zero when the denominator vanishes. The factor λ_β accounts for the fixed scale and Haar factors. Changing the adapted integral frame multiplies the residue determinant by a unit and thus the character by a scalar of modulus one. This is the local calculation of `exists_lineTerm_localDatum`: the integral frame block identity and the mixed-model identification of the slices must be applied before the Gauss sum. Formula (34.1) is not the Gauss factor of a datum transformed by an arbitrary rational matrix.

The orbital calculation. The central integration along the line parabolic annihilates nonisotropic line variables by additive orthogonality. The term at the zero vector is zero for the standard row datum: the associated first determinant column is zero. The nonzero isotropic vectors form one rational orbit up to the line scalar. Witt's extension theorem therefore gives a sum over rational isotropic lines, unfolded to the line parabolic, whose Levi is $E^\times \times U(V)$. If c is the line-scaling idele, the noncentral unipotent coordinates form an E-space of dimension r and the central coordinate an F-line. Their conjugation Jacobian in the Iwasawa convention of the slice is $|c|^{-r-1}$, giving the inverse factor $|c|^{r+1}$ in the integral. The residual unipotent integration is the zero evaluation in the second line direction. The fixed Haar scalar from unfolding is independent of the character. This is the geometric part of `ConstantTerm.lean`; in particular the term left over is an integral over $E^\times\backslash\mathbb{A}_E^\times$, not an extra copy of a sum of local unit integrals.

The two mixed coordinates at the last line are paired by the symplectic form. The other common line variables comprise $n+r+1$ isotropic coordinates (where $n = 2d$) and the residual symplectic space for $\mathrm{Hom}(W, V)$. Fourier transform in the remaining pair and the additive Poisson formula identify the sums over $\xi \in E$ in these two coordinates. Since both zero terms vanish for the standard determinant support, this is also equality of the sums over $E^\times$. It is crucial that the two coordinates give the same sum, not the same function at a specified nonzero ξ.

The Levi transformation used here is `lineTerm_pairWeil_levi` of `Lib/Weil/DualPair/TwoLineTerms.lean`. In its notation u is the source line-scaling parameter, c the target one, and either $u = 1$ or $r = n$. With $a = \mathrm{rankTwist}(\nu^{-1})$, it gives the scalar

$$\frac{|u|^{r+1}}{a.\,\mathrm{spaceChar}(n)(cu)\,|cu|^{d+1-\beta}}, \qquad F_1(cu) = cu, \quad F_0(cu) = \overline{cu}^{-1}$$

and the lower theta kernel evaluated at $\xi F_\beta(cu)$. In particular $a.\,\mathrm{spaceChar}(n) = \nu_E^{-1}$, where $\nu_E(x) = \nu(x/\bar{x})$. Multiplying by the unfolding Jacobian gives $\nu_E(cu)|cu|^{e+\beta}$. On combining the rational scalar sum with the idele quotient integral, the change of variable is $x \mapsto \xi x$ for $\beta = 1$, and $x \mapsto \xi\bar{x}^{-1}$ for $\beta = 0$. The character is unchanged by conjugate inverse on this norm-one quotient. The exponents of the two resulting zeta integrals are therefore $e+1$ and $-e$.

One must use the branch whose exponent is positive:

$$(\beta, s) = (1, e+1) \quad (e \geq 0), \qquad (\beta, s) = (0, -e) \quad (e < 0).$$

Lemma 34.2 (Arithmetic part of the line constant term)**.** *Suppose the unfolding and Levi identities above hold with the normalized slices, and their adelic integrals converge or admit the Mellin interpolation described below. Then the scalar γ_ν relating the line constant term to the lower theta integral with its real cone data satisfies*

$$|\gamma_\nu|^2 = C_{e,\alpha}\, Q^{2s-1} |L^S(s, \nu_E)|^2.$$

Here $C_{e,\alpha} > 0$ depends on the fixed dimensions, measure and scale conventions, not on the varying tame character or conductor; L^S omits S and its conjugate closure. In the branch $e \geq 0$ this is $|\gamma_\nu|^2 P_\nu(e) = C_{e,\alpha} P_\nu(e+1)$ for $P_\nu(k) = Q^{k^2} \prod_{j=1}^k |L^S(j, \nu_E)|^2$. The same relation holds for $e < 0$ on extending P_ν by $P_\nu(-k) = P_\nu(k)^{-1}$.

Proof. In (34.1), at each $w \in S$ the support is precisely $a_w \in \mathcal{O}_w^\times$, $\bar{a}_w \in \pi_w^{-1}\mathcal{O}_w^\times$. Translating by the corresponding idele makes both unit conditions. Its norm, including the fixed vacuum scale, is $|\alpha|Q$. The multiplicative character factors then cancel the unit characters in the zeta integral. The remaining character values of the fixed translating idele have modulus

one. At a place outside $S \cup \bar{S}$, summing over nonnegative valuations of the integral scalar yields the Euler factor of L^S. At the removed places the normalized unit integrals are one. Thus the finite scalar has absolute value $C_{e,\alpha}^{1/2} Q^{-1/2} Q^s |L^S(s, \nu_E)|$. The real integral is not included in this scalar: it supplies the lower datum, with integrand the real slice multiplied by $\prod_\tau (y_\tau/|y_\tau|)^{2\mu_\tau} |y_\infty|^s$ and the real Weil action of the compact variable. This convention is necessary to make the scalar independent of the infinite weights.

For $e \geq 0$, $(e+1)^2 - e^2 = 2s - 1$ and the quotient of successive Euler products is the new factor at $s = e + 1$. If $e = -k < 0$, then $P_\nu(1-k)/P_\nu(-k) = P_\nu(k)/P_\nu(k-1)$, again with exponent $2k - 1 = 2s - 1$. □

At the borderline $s = 1$, the argument must not multiply conditionally convergent local series. The interpolation in `LargeSideConstantTerm.lean` introduces a real Mellin exponent t on the unfolded idele quotient. In the two slices the exponents become $e + 1 + t$ and $-e - t$. One obtains an open interval of absolute convergence on the far right from the first slice and on the far left from the second slice, since their finite Tate zeta integrals converge absolutely for exponent > 1 and the real slices are rapidly decreasing. Their equality prior to unfolding is the Poisson identity. If the common unfolded integrand is absolutely integrable at two such endpoints, it is absolutely integrable throughout the intervening strip by Hölder's inequality. The same estimate on compact substrips, with slightly enlarged endpoints, gives holomorphy of the Mellin integral. On the chosen half-strip the factorization into the absolutely convergent zeta integral and the real slice holds; the holomorphic extension of the Hecke L-function at 1 (for a nontrivial Hecke character) and the convergent real cone factor then identify its value at $t = 0$. This mechanism depends on the common-integrand integrability at the endpoints, which must be proved for the actual theta kernels.

Classical tower statements cannot by themselves furnish this constant. For example, Chenyan Wu, *Irreducibility of theta lifting for unitary groups*, arXiv:1409.0770, Proposition 3.1, unfolds the isotropic orbits for a *cuspidal* input; the nonzero orbit in that calculation vanishes by cuspidality. It is useful for the unfolding mechanism but not as a reference for the surviving determinant-datum orbit here, or for (34.1). The normalized identity and its endpoint hypotheses must be checked in the actual noncuspidal setting.

34.3 Unramified polynomial transfer

At a split unramified place with residue field k_q, let the source have dimension n and target dimension $m = n + k$, $k \geq 0$. Let χ_S and χ_T be the source and target space characters in the split Schrödinger model, put $A = q^{n/2}\chi_S(\pi)$ and $B = q^{m/2}\chi_T(\pi)$, and normalize the degree-j Hecke operator T_j as the sum over the elementary-divisor double coset with j factors π. Define

$$P_d(X) = \sum_{j=0}^{d} (-X)^j q^{j(j-1)/2} T_j, \qquad P_d^{\mathrm{triv}}(X) = \prod_{i=0}^{d-1} (1 - q^i X).$$

The transpose and inverse in the following argument are important.

Lemma 34.3 (Split local transfer)**.** *On the local integral-matrix vacuum the action of $P_m(X)$ on the target equals the action of*

$$P_k^{\mathrm{triv}}(AX)\, P_n(q^k A B^{-1} X)$$

on the source at the conjugate place. Consequently the identity persists after application of an equivariant kernel pairing; if the source is spherical, the target lift (when nonzero) has the corresponding spherical Hecke polynomial.

Proof. Represent elementary-divisor cosets by matrices M_U, with the reduction kernel U running over subspaces of k_q^d of dimension j. For the target use transpose representatives $h = M_U^t$. The split Weil formula on the vacuum $\mathbf{1}_{\mathrm{Mat}_{m,n}(\mathcal{O})}$ has scalar $\chi_S(\det h)|\det h|^{-n/2} = A^j$. At an integral matrix Y of reduction rank ρ, the transformed function is nonzero exactly when $h^{-1}Y$ is integral, or equivalently when U is orthogonal to the image of $\bar{Y}$. Thus the sum of this function over the double coset counts the j-dimensional subspaces of a space of dimension $m-\rho$. On the source at the conjugate place use $g = M_U^{-1}$. Its scalar is B^j, and Yg is integral exactly when $U \subseteq \ker \bar{Y}$, now of dimension $n-\rho$. Outside the integral matrix lattice both sums are zero, because multiplication by the respective integral matrices would recover Y.

For $d \geq 0$, the subspace-counting identity is

$$\sum_{U \leq k_q^d} (-Z)^{\dim U} q^{\binom{\dim U}{2}} = \prod_{i=0}^{d-1} (1 - q^i Z).$$

Indeed if $C(d,j)$ counts the j-subspaces, those contained in a fixed hyperplane contribute $C(d-1,j)$ and those not contained contribute $q^{d-j}C(d-1,j-1)$. Substitution gives the polynomial recurrence multiplying the preceding polynomial by $1 - q^{d-1}Z$. It follows that on Y the target polynomial acts as $P^{\mathrm{triv}}_{m-\rho}(AX)$ and the source polynomial as $P^{\mathrm{triv}}_{n-\rho}(BX)$. Factoring the former into the first k factors and the rest gives the asserted equality pointwise.

Since the Weil actions commute, the same equality holds after applying an intertwining linear map. In a kernel pairing, integrate the source action by changing variable by the double-coset representative. Haar invariance sends that action to its adjoint double coset, which for these split elementary divisors is at the conjugate place. Thus it acts on the paired spherical source vector by the source eigenvalue at the original place. This proves transfer of the polynomial eigenvalue with the specified scalar argument. □

This proves the computation in `sum_heckePolyCoeff_smul_pairWeil`, `hecke_transfer_of_equivariant`, `hecke_transfer_of_pairing`, and `isSphericalWith_formLift` of `Lib/Weil/Unramified/HeckeTransfer.lean`, relative to the split Weil action. The last assertion is a statement about a genuine equivariant globally defined lift, not an assertion that a formal difference of kernels with unproved boundedness belongs to that representation. This separation matters in the small-root induction.

34.4 The normalized rank-one pairing

We record exactly where the convergent doubling identity enters the norm calculation. Let S be the source and T the target, with $\operatorname{rk} T > 2 \operatorname{rk} S$. In the doubled model for $S \oplus S^-$, the convergent Siegel–Weil identity used by `Lib/SiegelWeil/Formula/SiegelWeilFormula.lean` expresses the mean of the joined theta function as

$$\mathrm{vol}[U(T)] \sum_L \mathrm{zeroTerm}_L(f' \otimes f), \tag{34.2}$$

where L ranges over rational maximal isotropic subspaces and the join uses opposite symplectic signs. The source version includes summability in the relevant operator norms. Its zero term on a graph $L = L_\delta$ of $\delta \in U(S)(F)$ is the matrix coefficient

$$\mathrm{zeroTerm}_{L_\delta}(f' \otimes f) = \langle f', \omega(\delta^{-1}, 1) f \rangle. \tag{34.3}$$

The normalization in (34.3) is the normalized invariant functional in the joined Schrödinger model, not an arbitrary scalar multiple of that functional. The general `inner_product_`

`formula` of `Lib/SiegelWeil/RallisInner/SharpData.lean` permits clean data for which nongraph terms vanish. For a definite source, which suffices for the rank-one calculation below, there are no nongraph terms: the kernel of either projection of L would give an isotropic vector in the definite source, hence both projections are injective and, by dimension, bijective. The graph map is an isometry by isotropy.

Inserting (34.3) into (34.2) and unfolding the sum indexed by $U(S)(F)$ yields

$$\langle \Theta(f', y'), \Theta(f, y) \rangle_{\mathrm{Pet}} = \mathrm{vol}[U(T)] \int_{U(S)(\mathbb{A}_F)} \langle f', \omega(g, 1) f \rangle \langle y', R(g) y \rangle \, dg. \tag{34.4}$$

The change of variables is the ordinary unfolding of the sum of translates over the rational group in the two source integrals. Rapid source data and the norm summability in (34.2) justify Fubini; the definite rank-one source uses a constant source vector. Formula (34.4) is thus proved assuming the normalized convergent Siegel–Weil and graph evaluations. A reference to the general Rallis formula without specifying the joined-model, Haar and graph scalars is not sufficient for the exact arithmetic statement. We retain (34.2)–(34.3), including their measure compatibility and convergence, as explicit dependencies.

Lemma 34.4 (Principal-level sharp support)**.** *Suppose the split finite datum f has integral matrix support and reduction matrices Y on its support have a fixed left inverse L modulo π. A translate for which both Y and Yg are in its support has $g \in \mathrm{GL}_n(\mathcal{O})$, $g \equiv 1 \bmod \pi$. If the residual character and datum are fixed by this principal subgroup J, the matrix coefficient is $\|f\|^2$ on J and zero outside.*

Proof. Each integral lift Y has an integral left inverse, because one can lift the given inverse and invert the product, which is congruent to the identity. Thus Yg integral implies g integral, while using a left inverse for Yg shows g^{-1} integral. Reduction of the equality Yg and multiplication by the fixed residual inverse give $\bar{g} = L\overline{Yg} = 1$. The support criterion and invariance of the vector prove the coefficient assertion. With hyperspecial volume one its integral is $\|f\|^2/[K : J]$. □

This is the sharp residue calculation `isSharp_modDatum`. It also shows why an upper bound on the pointwise norm of a residue datum would not give the correct conductor power.

Proposition 34.5 (Rank-one norm with conductor powers)**.** *Take the data of `LValueData.inner_trivial_Lift_self_eq_L` in `New/Exchange/CharacterLiftNorm.lean`. Let the target rank be $n = 2z \geq 4$, M the chosen split places and Q the disjoint twist places. Put*

$$A = \prod_{w \in M} (q_w - 1), \quad B = \prod_{w \in Q} (q_w - 1), \quad Q_0 = \prod_{w \in Q} q_w, \quad N = \prod_{w \in M \cup Q} q_w.$$

Subject to (34.2)–(34.3) with their given normalization, the algebraic form $\Xi = \mathrm{stdClassSum}$ satisfies

$$\|\Xi\|^2 = \frac{\mu_T \mu_{U(1)}}{Z(z)} A \, H \, d \, L^{M \cup Q}(z, \mu^{-1}) Q_0^{n-1},$$

where $\mu_T, \mu_{U(1)}$ are the fixed adelic quotient volumes, H the real hook norm, d the coefficient dimension, and $Z(z)$ the fixed unramified product described below. The removed factors include the conjugate closure.

Proof. At the chosen primes the residual datum is supported at a specified vector; at a twist prime one entry is fixed, the other $n - 1$ entries free, with coefficient of absolute value $q_w - 1$. Thus the support has Q_0^{n-1} elements and the squared norm, with the integral matrix lattice of measure one, is

$$\|f_{\mathrm{ram}}\|^2 = B^2 Q_0^{n-1} / N^n.$$

The principal subgroup has index AB in the rank-one compact. Applying Lemma 34.4 gives the ramified factor $c = B^2 Q_0^{n-1}/(N^n AB)$.

At a split vacuum place write $g_j = (\pi^j, \bar{\pi}^{-j})$ for the torus representatives. The split vacuum overlap is $q_w^{-z|j|}$ times the unit phase λ^{-j} with $\lambda = \mu_E(\pi_w)$. The latter phase is the character after cancelling the rank twist υ^z against the vacuum multiplier. The sum over $j \in \mathbb{Z}$ is

$$1 + \sum_{j \geq 1} q_w^{-zj}(\lambda^j + \lambda^{-j}) = \frac{1 - q_w^{-2z}}{(1 - \lambda q_w^{-z})(1 - \lambda^{-1} q_w^{-z})}.$$

At a fixed nonsplit place the norm-one torus is compact, so there is just the integral vacuum coefficient. Let $\epsilon_w = -1$ on one of the two places of each split pair and $+1$ on the other and on each nonsplit fixed place. The product $Z(z) = \prod_w (1 - \epsilon_w q_w^{-z})^{-1}$ is positive and absolutely convergent for $z > 1$. At a split pair its numerator after inversion is $1 - q_w^{-2z}$, and at a fixed place its inverse cancels the trivial local factor of the pulled-back torus character. Hence the vacuum factors outside the ramified places multiply to

$$\frac{L^{M \cup Q}(z, \mu^{-1})}{Z(z) \prod_{w \in M \cup Q} (1 - q_w^{-n})}.$$

The rank-one real source action is trivial with the specified weights, and its normalized real highest weight vector has squared norm $H = \prod_\tau d_\tau!$ by the monomial Gaussian integral. The constant source vector has Petersson squared norm $\mu_{U(1)}$. Formula (34.4) yields the Petersson squared norm $\mu_T \mu_{U(1)} H c$ times the displayed vacuum product.

It remains to convert to the algebraic class sum; this step is not an arbitrary normalization of the L-value. The chosen principal-level rank-one class sum has index A, and the source integration realizes $\Theta = A^{-1}\,\mathrm{ofAlgForm}\,\Xi$. The Petersson-to-class pairing multiplies by the coefficient dimension d and the target level index. The prescribed target level is the stabilizer of a nonzero residual vector at M and of a residual hyperplane at Q. The indices are $q^n - 1$ and $(q^n - 1)/(q - 1)$, respectively: these follow by counting nonzero vectors and nonzero linear functionals up to scaling. The resulting index is

$$I_T = \frac{N^n}{B} \prod_{w \in M \cup Q} (1 - q_w^{-n}).$$

Consequently the conversion multiplies the Petersson norm by $A^2 d I_T$. Substitution gives $A^2 c I_T = A Q_0^{n-1} \prod_{w \in M \cup Q} (1 - q_w^{-n})$. The indicated factors cancel, proving the formula. □

The precise positive constant, independent of the varying rank-one character at fixed rank, is an essential feature of the norm expression. The calculation covers the Euler, ramified, hook, and class-index factors. To transfer it to field-conjugate norms one must also prove the semilinear normalized kernel identity used in `New/Exchange/AlgebraicPartConjugates.lean` (`map_trivialLift`): applying a complex field automorphism directly to a positive Hermitian norm is not justified.

34.5 The depth indicator: sharpness and six-fold decompositions

We now turn to a different specialized part of C: the arithmetic attainability of a large coefficient. The argument uses the integral formal parameter and the tight-frame construction of the CM chapters. For a fixed embedding into $\mathbb{C}_p$, write $\mathcal{O}$ for its valuation ring and $|\cdot| = |\cdot|_p$,

normalized by $|p| = p^{-1}$. The formal group parameter $\phi = \mathrm{translatedWpDeriv}$ is normalized to linear coefficient a unit. Denote by ρ the height scale of the formal group, with an element ϖ satisfying $|\varpi| = p^{-\rho}$. We need sharpness, not merely the upper Katz bound.

Lemma 34.6 (Katz sharpness from the formal parameter)**.** *Suppose the integral formal parameter theorem supplies an element $T \in \mathcal{O}[[\phi]]$ with*

$$T(\varpi z) = \varpi S(z), \quad S \in \mathcal{O}[[z]], \qquad S - S^{p^h} \equiv z \pmod{\mathfrak{m}}, \quad h \geq 1,$$

and the upper factorial bound $|j![z^j]F| \leq p^{(\rho-1/(p-1))j}$ for every $F \in \mathcal{O}[[\phi]]$. Then for every $j \geq 0$ there exists $F \in \mathcal{O}[[\phi]]$ attaining the upper bound.

Proof. Let N_j be the maximum of $|j![z^j]F|$ on the integral formal ring. The maximum exists because ϕ has order one; only the finitely many powers ϕ^i, $0 \leq i \leq j$, contribute, and the ultrametric inequality bounds all linear combinations by their largest coefficient norm. Put $c_j = N_j p^{-(\rho-1/(p-1))j}$, so $c_j \leq 1$. By the integral formal group law $F(z+w)$ belongs to $\mathcal{O}[[\phi(z), \phi(w)]]$. Its coefficient of $z^j w^k$, multiplied by $j!k!$, is $(j+k)![z^{j+k}]F$ and is bounded by $N_j N_k$. Hence $c_{j+k} \leq c_j c_k$. If any $c_j < 1$ with $j > 0$, this inequality, on writing $n = aj + b$ with $0 \leq b < j$, implies $c_n \to 0$.

On the other hand, by the definition of N_n and Legendre's formula for factorials,

$$|[z^n]S| \leq |\varpi|^{n-1} \frac{N_n}{|n!|} = p^{\rho} c_n p^{-s_p(n)/(p-1)},$$

where $s_p(n)$ is the sum of the base-p digits. If $c_n \to 0$, reduction of S modulo the maximal ideal would be a polynomial. A constant polynomial cannot satisfy $S - S^{p^h} = z$; a polynomial of positive degree d would give degree $p^h d > 1$ on the left, also impossible. Thus $c_n = 1$ for every n. □

This is the sharpness argument in `New/Curve/KatzDivisibility.lean`, relative to the actual integral formal parameter theorem `New/Curve/FormalParameter.lean`. Merely knowing that the upper bound is asymptotically sharp would not suffice.

For the tight frame write $\ell_i(z) = \sum_\tau a_{i\tau} z_\tau$ and

$$\Theta(z) = \prod_i \theta(\ell_i(z))^{c_i}, \qquad R = \mathcal{O}[[\phi(\ell_i(z)) : i]].$$

For each multi-index α let $N(\alpha)$ be the largest absolute value of the coefficient $[z^\alpha]\Theta \prod_i \phi(\ell_i)^{e_i}$ for $e_i \geq 0$, $\sum e_i \leq |\alpha|$. In particular it is attained by one of these finitely many products. If α is a sum of k multi-indices with product of their N values at least c, call this a k-split of size c. Say a formal series is carried by k if each of its coefficients is bounded in absolute value by the largest such product at that multi-index. This is the finite coefficient device of `New/Rows/GluedFormTheta.lean`, `HasLargeSplit` and `IsCarriedBy`.

Lemma 34.7 (Carried inverse and ordinary differentiation)**.** *Suppose the norm-pair log-theta identity furnishes units $U_j \in R$ for $j \in \mathbb{Z}_p$ relating $\Theta(jz)$ to $\exp(-j^2 L(z))$, and the analogous relation at $j = 1$. Then Θ^{-1} is carried by five, and every element of R is carried by six. If a frame vector ℓ_i has $|a_{i\tau}| = 1$ for every τ, there exists a six-split of α of size at least*

$$p^{(\rho-1/(p-1))|\alpha| + \sum_\tau v_p(\alpha_\tau!)}.$$

Proof. Every $H\Theta$, $H \in R$, is carried by one by the finite maximization defining N. Products of carried series add their carry counts, by the coefficient Cauchy product and the ultrametric

inequality. The four-square theorem and compactness of $\mathbb{Z}_p^4$ give $j_1,\dots,j_4\in\mathbb{Z}_p$ with $\sum j_i^2=-2$: take a convergent subsequence of solutions for nonnegative integers tending to -2. The log-theta identities then give

$$\Theta^{-1}=\Theta\cdot\prod_{\nu=1}^{4}\Theta(j_\nu z)\cdot U,\qquad U\in R^\times.$$

Each scaled series $\Theta(j_\nu z)$ is carried by one, because its coefficients are multiplied by $j_\nu^{|\alpha|}$ of norm at most one. The remaining ΘU is also carried by one. Thus Θ^{-1} is carried by five, and $H=(H\Theta)\Theta^{-1}$ is carried by six.

By Lemma 34.6 select $F\in\mathcal{O}[[\phi]]$ attaining the upper bound at $j=|\alpha|$. The coefficient of $F(\ell_i(z))\in R$ at α is

$$[w^j]F\frac{j!}{\prod_\tau\alpha_\tau!}\prod_\tau a_{i\tau}^{\alpha_\tau}.$$

Its norm is precisely the displayed size. Since the element is carried by six, a six-split of that size exists. □

Extra depth at a nonsplit prime. At an inert prime, $h=2$ and $\rho=1/(p^2-1)$. For the ramified prime the packing has no pairs and the derivative bound is vacuous (there $\rho=1/(2(p-1))$). Write $F(z)=\Theta(\varpi z)=\exp(\kappa\Psi(z))V(z)$, where $\Psi\in\mathcal{O}[[z]]$, $V\in R^\times$ is integral after rescaling, and $|\kappa|=p^{1-1/(p-1)}$ (the prime 2 is excluded in this branch). This is the rescaled theta normal form. For an admissible Frobenius packing choose $y_\sigma\ge0$ and odd i_σ, with paired coordinate t_σ the transformed place $-p^{i_\sigma}\sigma$, and put

$$\tau=\sum_\sigma y_\sigma e_{t_\sigma},\quad M=\sum_\sigma p^{i_\sigma}y_\sigma e_\sigma,\quad b=M+\tau.$$

The derivative assertion in the theta normal form says that the selected derivatives $\partial_{t_\sigma}\Psi$ have, under the weights assigned by this packing, distinguished lowest terms $z_\sigma^{p^{i_\sigma}}$ of unit coefficient; the product with repetitions y_σ has coefficient a unit at M. The weights of this assertion depend on the packing; a Frobenius relation on the linear forms alone does not establish it.

Lemma 34.8 (Frobenius derivative splitting)**.** *With this derivative assertion, a six-split of b has size at least*

$$p^{(1-1/(p-1))|\tau|+\rho|b|+\sum_j v_p(\tau_j!)}.$$

Proof. Starting with $P=1$, each selected derivative ∂_t replaces P by

$$P_{\mathrm{new}}=(\partial_t\Psi)P+\kappa^{-1}(\partial_tP+PV^{-1}\partial_tV).$$

The identity $\partial_t(PF)=\kappa P_{\mathrm{new}}F$ follows by differentiation. The new series is integral, and modulo the maximal ideal the second term vanishes since $|\kappa^{-1}|<1$. Consequently after all derivatives we obtain $\partial^\tau F=\kappa^{|\tau|}PF$, with $[z^M]P$ a unit by the packing derivative assertion. Since $P=F^{-1}(PF)$ the ultrametric inequality produces $\gamma+\beta=M$ for which $|[z^\gamma]F^{-1}\,[z^\beta](PF)|\ge1$. But

$$[z^\beta](PF)=\kappa^{-|\tau|}\prod_j\frac{(\beta_j+\tau_j)!}{\beta_j!}[z^{\beta+\tau}]F.$$

The factorial ratios are divisible by $\prod_j\tau_j!$ as rational integers. Therefore the product of the corresponding coefficients of F^{-1} and F has norm at least $|\kappa|^{|\tau|}/|\tau!|$. Undo the rescaling by ϖ; their multi-indices sum to b. The resulting lower bound is the displayed expression. Finally Θ^{-1} is carried by five, Θ by one, so their product of two selected coefficient bounds is attained by a six-split. □

This is the coefficient argument of `New/Curve/DepthPairs.lean`, `exists_hasLargeSplit_of_pairs`. It explains why one needs an inverse theta series and cannot obtain the optimal bound by bounding every differentiated theta coefficient in only one direction.

Twelve blocks. For $\alpha = b + c$ with b given by the pairing, append six ordinary factors from Lemma 34.7 on c. One obtains a twelve-split of size at least

$$p^{(1-1/(p-1))|\tau|+\rho|b|+\sum_j v_p(\tau_j!)+(\rho-1/(p-1))|c|+\sum_j v_p(c_j!)}. \tag{34.5}$$

In the large-prime branch $i_\sigma = 1$, there is a better substitute for the last six factors: their size can be made at least one. Indeed the tight-frame identity expresses each coordinate linear form as an integral linear combination of the ℓ_i at a prime not dividing the fixed frame denominator. Replacing each ℓ_i by $\phi(\ell_i)$ gives a series of the same linear part, and multiplying powers of these series gives an element of R with lowest monomial of index c and unit coefficient. Its carry-six assertion yields the required six-split. Since $|b| = (p+1)|\tau|$ and $\rho(p+1) = 1/(p-1)$, the combined size is at least $p^{|\tau|}$.

For the smaller-prime branch, using Legendre's formula in (34.5) shows exactly the numerical quantity required of the packing. If all the relevant indices have digit-sum loss at most D in their factorial valuations, the exponent in that formula differs from $\sum_j v_p(\alpha_j!)$ by at least

$$|\tau| + \rho|\alpha| - |\alpha|/(p-1) - 2fD,$$

where f is the number of coordinates. This follows by replacing the two positive factorial sums by their lower bounds and the subtracted sum by the upper bound $|\alpha|/(p-1)$. Thus the actual inequalities of `Data.exists_isFrobPacking` are what convert the twelve-split into the desired forced gain with the prescribed losses. Even with the coefficient calculation in hand, that packing theorem, and the realization of the twelve chosen maximizing products by an integral finite indicator, are separate obligations. In particular a formal power series whose coefficient is large is not itself an adelic integral test form.

34.6 Packing loss and realization of an indicator

We supply the numerical packing along the cofinal admissible family of Chapter 54. Write $w_\sigma = w_{\mathrm{nom}}(\sigma)$, $w_- = a/x_1$, $w_+ = a/x$, and suppose $w_\sigma \le n_\sigma \le w_\sigma + x^{42}$. Put

$$c_0 = \lceil 60b/k \rceil, \quad l_\sigma = (n_\sigma - c_0)_+, \quad D_\sigma = \left(\prod_{j=1}^{c_0} (l_\sigma + j) \right)^2.$$

This is precisely `depthLossInt`. For a prime p with $\chi_K(p) \neq 1$ set $\mathrm{pr}_e(\sigma) = [-p^e]\sigma$ and $\omega_p = (1 - \chi_K(p))/2$. The live exponents are positive odd exponents at an inert prime and all positive exponents at the ramified prime $p = d$. The local ask is

$$A_p(w_\sigma, \sigma) = \omega_p \sum_{\text{live } e} (w_\sigma/p^e - w_{\mathrm{pr}_e(\sigma)})_+,$$

whose sum over σ is $F(p)$.

Proposition 34.9 (Packing with the prescribed loss integers)**.** *For sufficiently large records in that family, there are admissible pairs with indices i_σ and multiplicities y_σ as in Section 34.5, such that either p is inert, $i_\sigma = 1$ and*

$$\sum_\sigma \{v_p(n_\sigma!) - y_\sigma\} \le F(p) + \sum_\sigma v_p(D_\sigma),$$

or

$$\sum_{\sigma}\{(1/(p-1)-\rho_p)n_\sigma + 4b_2 - y_\sigma\} \le F(p) + \sum_{\sigma} v_p(D_\sigma).$$

Here $\rho_p = 1/(p^2-1)$ *when inert and* $1/(2(p-1))$ *when ramified.*

Proof. In both branches $x^{42} \le c_0$, $n_\sigma \le 2b$, and $w_- \gg x^{42}+1$: these follow from power dominance. If $p > k$, the prime is inert since d is fixed. Choose $i_\sigma = 1$ and

$$y_\sigma = \min\{n_{\mathrm{pr}_1(\sigma)}, \lfloor l_\sigma/p \rfloor\}.$$

Because $n_\sigma \le 2b$ and $p > k$, one has $y_\sigma \le c_0$. The map pr_1 is a permutation. At each vertex, if $n_\sigma \le c_0$, the outgoing weight is zero and the incoming weight at most n_σ. If $n_\sigma > c_0$, the outgoing weight py_σ is at most $n_\sigma - c_0$ and the incoming weight at most c_0. Thus total capacity is respected. The derivative admissibility exclusion of earlier odd exponents is vacuous.

Let $C_\sigma = \prod_{j=1}^{c_0}(l_\sigma + j)$. Since $l_\sigma + c_0 \ge n_\sigma$, the factorial identity and $v_p(l!) = \lfloor l/p \rfloor + v_p(\lfloor l/p \rfloor!)$ give

$$v_p(n_\sigma!) \le \lfloor l_\sigma/p \rfloor + v_p(\lfloor l_\sigma/p \rfloor!) + v_p(C_\sigma).$$

A product of c_0 consecutive integers is divisible by the factorial of every integer at most c_0. Hence the middle valuation is bounded by $v_p(C_\sigma)$. Subtract y_σ, noting $l_\sigma \le w_\sigma$ and $n_{\mathrm{pr}_1(\sigma)} \ge w_{\mathrm{pr}_1(\sigma)}$, to obtain

$$v_p(n_\sigma!) - y_\sigma \le (w_\sigma/p - w_{\mathrm{pr}_1(\sigma)})_+ + v_p(D_\sigma).$$

Summation, using the first live term of $F(p)$, proves the first branch.

Now take $p \le k$. Every nonsplit prime is at least 13 and prime to N_1 (the primes dividing the cyclotomic modulus split in the fixed quadratic field). At an inert prime use $g = [-p]$ in the place group. For any $1 \le j \le 324s$ its projected class in $(\mathbb{Z}/2^s)^\times$ satisfies $g^j \ne 1$ for large x. Indeed if $u = -p$ and j is odd then $v_2(u^j - 1) = v_2(u-1)$; if j is even, repeated difference of squares gives $v_2(u^j-1) = v_2(u^2-1) + v_2(j) - 1$. Thus $2^{v_2(u^j-1)} \le (p+1)^2 j = O(x^{10}\log x) = o(2^s)$. The equality of projected power classes of any three thick places along such an arc contradicts the actual three-term condition. There cannot be three thick places $\sigma, g^i\sigma, g^j\sigma$ with $0 < i < j \le 324s$.

At each thick σ at an inert prime, if there is a thick target $g^i\sigma$ for some odd $i < s$, choose this $i = i_\sigma$ and set $y_\sigma = \lfloor n_\sigma/p^{i_\sigma} \rfloor$. Such an outgoing pair is unique by the rule just proved. At all other vertices set $y_\sigma = 0$. At a ramified prime set all $y_\sigma = 0$: two thick places can never be related by $[-d^e]$, since their common class modulo N_0 is -1, whereas $d^e \ne -1 \bmod N_0$ (the fixed congruence gives $d^2 = 1$, $d \ne -1$, and $N_0 > 2$).

In the inert construction no vertex has both an incoming and an outgoing selected pair: otherwise there would be three thick places at exponents $0, i, i+j < 2s$, or a forbidden short period. Nor can two incoming pairs have a common target: subtract their exponents and use the three-term rule, or, if the exponents agree, cancel the common group element. The outgoing capacity is immediate from the floor and the incoming capacity follows from $y_\sigma \le 2b/p < w_+ \le n_{g^{i}\sigma}$. If a selected source t could meet the selected target of σ by an earlier odd exponent $j < i_\sigma$, then $t = g^{i_\sigma - j}\sigma$ gives the third thick place in the forbidden short arc. This proves the derivative admissibility restriction as well.

For the valuation estimate, put $\lambda_p = 1/(p-1) - \rho_p$. A geometric progression gives

$$\lambda_p n_\sigma = \omega_p \sum_{\text{live } e} n_\sigma/p^e.$$

For $e < s$, subtract the corresponding local ask summand and, if this is the selected exponent of σ, subtract y_σ as well. In the selected case the difference is at most one. Otherwise the

two places are not both thick, and the difference before multiplication by ω_p is bounded by $2(w_- + b_2)$: if the source is thin, bound its whole weight by $w_- + x^{42}$; if the target is thin, subtracting the positive part leaves at most $w_- + x^{42}$. The same bound holds in the ramified case. Uniformly for $p \le k$ and $e < s$,

$$2(w_- + b_2) \le c_0/(p2^e).$$

Indeed $x_1 \ge x^{10^6}$, $w_- \sim xb/x_1$, $b_2 = w_- + O(N_1)$, $k = O(x^5)$ and $2^s = O(x^{11})$. For $e \ge s$, the difference is at most $n_\sigma/p^e \le c_0/(p2^e)$, since $p \ge 13$, $n_\sigma \le 2b$, and $(13/2)^s$ grows faster than k. Summing these geometric bounds gives

$$\lambda_p n_\sigma - A_p(w_\sigma, \sigma) - y_\sigma \le c_0/p.$$

There are at least $\lfloor c_0/p \rfloor$ multiples of p in C_σ, and $c_0/p \ge 4b_2 + 2$ uniformly by the same size comparison. Hence $v_p(D_\sigma) \ge 2\lfloor c_0/p \rfloor \ge c_0/p + 4b_2$. Summing proves the second branch. □

This gives the two alternatives in `Data.exists_isFrobPacking`, `New/Curve/DepthPairs.lean`, with sufficiently large rather than the source's explicit finite thresholds. It also handles the ramified prime correctly: no Frobenius derivative pair is used there. By Legendre's formula and the size bounds, $v_p(j!) \ge j/(p-1) - 2b_2$ for $0 \le j \le 2b$, since the base-p digit loss is at most $1 + \log_2(2b+1)$ and b_2 exceeds this. With Lemma 34.8, the second alternative therefore gives exactly the lower bound

$$\exists\, (\alpha_i)_{i=1}^{12}, \quad \sum_i \alpha_i = n, \qquad \prod_{i=1}^{12} N(\alpha_i) \ge p^{-F(p) - \sum_\sigma v_p(D_\sigma) + \sum_\sigma v_p(n_\sigma!)}. \tag{34.6}$$

The first alternative uses instead the integral coordinate monomial construction in the large-prime branch. At a split prime $\rho_p = 1/(p-1)$ and $F(p) = 0$; Lemma 34.7 with six zero multi-indices appended implies (34.6) directly.

From the twelve products to the residue indicator. The precise finite realization and vacuum expansion of Proposition 17.1 and the CM theta chapter are essential. We now show how they give the particular indicator in `Data.depth_indicator_jet`. Let $G = g_0 g_1$ be the fixed norm of $\delta = \pi_G$, and write κ_0 for the rank-one vacuum clearing factor. Take the finite space for $\mathrm{Hom}(1, 12)$ with its all-positive signs. Choose once and for all a finite point y at which the scaled vacuum has nonzero value $t_0 = \kappa_0^{12} v(y)$. Existence follows from the nonzero vacuum and its finite realization. Put

$$t = G^{24f} \left(\prod_\sigma D_\sigma \right) t_0.$$

This is one complex number independent of the prime and the later choices of indicator.

For a selected maximizing product in each of the twelve factors (34.6), choose the tight frame with all $c_j \ge 3|n|$. Thus $c_j \ge 3e(\alpha_i)_j$ for each row. The curve monomial section $\theta^{c_j - 3e}(\phi\theta^3)^e$ has degree c_j and normalized Taylor product $\theta^{c_j}\phi^e$. These are exactly the sections allowed by the finite realization. With the specified period scale, the finite vector F constructed in that theorem has jet series the product of the twelve selected series in their disjoint variable sets. If $\gamma_{\sigma,i} = (\alpha_i)_\sigma$ and ℓ is extraction of the corresponding coefficient from the normalized jet series, then $|\iota\ell(F)| = \prod_i N(\alpha_i)$, with $\sum_i \gamma_{\sigma,i} = n_\sigma$. Moreover its theta values on all finite translates are integral. The vacuum expansion, using G^{24f} translates, is an identity of vectors

$$\sum_j M\, \ell(\rho(x_j)v)\rho(-x_j)F = G^{24f}\ell(F)v, \qquad \delta x_j \in \widehat{\mathcal{O}}_E^{12},$$

with M a natural number. Apply the theta functional after translating both sides by y. Every value of the translated F has absolute value at most one, including the central phase. Also $|M|_p \le 1$. It follows that

$$|\,\iota(G^{24f}\ell(F)v(y))\,| \le \max_j |\iota\ell(\rho(x_j)v)|.$$

For the modulus ($\bar\delta$) the last translates have an exact finite residue expansion $\rho(x_j)v = \sum_z \zeta_{j,z}\,\mathrm{modDatum}(\mathbf{1}_z)v$ in the named mixed model, with $\zeta_{j,z}$ roots of unity. The integral denominator condition $\delta x_j \in \widehat{\mathcal{O}}_E^{12}$ means the phase on the complementary integral coordinate is constant on cosets modulo $\bar\delta$ by additive duality. In the split Schrödinger coordinates this component changes the vacuum by a phase times its lattice indicator, while the other integral component preserves the lattice support. Equivalently expand the resulting finite character in the indicator basis of the residue classes. This is `exists_heis_vacElem_eq_sum_modDatum` in `Lib/Weil/Lift/Staircase.lean`; no Fourier denominator occurs when writing the function in the delta-function basis. Hence one indicator coefficient has norm at least that of $G^{24f}\ell(F)v(y)$. Multiply by $\kappa_0^{12}\prod_\sigma n_\sigma!/t$, take absolute values, and insert (34.6). The absolute value of $\prod_\sigma D_\sigma$ cancels its valuation in (34.6), and the factorials cancel the factorial gain. We obtain

$$\left|\iota\left(\frac{\kappa_0^{12}}{t}\prod_\sigma n_\sigma!\;[z^\gamma]\,\mathrm{jetSeries}_{\bar\delta\;\mathrm{jetScale}}\,\mathrm{modDatum}(\mathbf{1}_z)v\right)\right|_p \ge p^{-F(p)}.$$

In the archimedean bound, $\log G = O((\log b)^{11})$ by the chosen prime windows, and the vacuum bound in the CM chapter gives $\log|\sigma(t_0)| \le O(N_1^2)$ for every field conjugate. Since $n_\sigma \le 2b$ and $c_0 = O(b/k)$,

$$\log|\sigma(t)| \le 24f\log G + 2fc_0\log(2b+c_0) + O(N_1^2) = o(fmb/k).$$

For the last comparison, $m \ge \sqrt{x}$, $Y = \log b = x^\varepsilon = o(m)$, and polynomial terms are negligible relative to b. Thus the uniform bound $|\sigma(t)| \le \exp(fmb/k)$ holds for the cofinal family. This gives both the valuation and the archimedean normalizer in the actual depth-indicator conclusion, subject to the normalized finite realization, its vacuum estimate, and the weighted derivative assertion in the preceding CM foundations. The coefficient-to-indicator step no longer requires a new formal-source input of its own.

34.7 Exact-source obligation ledger

The following records what this chapter supplies, relative to the earlier model and formal-group foundations. An OPEN entry is not being asserted without proof; it identifies the residual translation needed for the theorem at the end of the monograph.

1. PROVED: the compact-circle obstruction in `New/Exchange/NonnegativeVanishing.lean`, `centreCoeff_eq_zero_of_not_negative`, under the integrability, mixed-model coefficient and term-equivariance hypotheses as stated (Section 34.1).

2. PROVED: the local Gauss sum, two-branch exponent and arithmetic conductor ratio in `LineGaussSum.lean` and `LargeSideConstantTerm.lean`. OPEN: verifying the actual mixed-model normalization and the common unfolded integrand endpoint estimates for `exists_constant_term_thetaLarge`, and the exact top-period identity for the compatible real and global data. The classical tower calculation for cuspidal input is not a reference for these claims.

3. PROVED: the split local polynomial transfer computation and its equivariant consequence (`sum_heckePolyCoeff_smul_pairWeil`, `isSphericalWith_formLift`). OPEN: boundedness and the normalized period conversion for the subtracted lifts to which this transfer is applied.

4. PROVED: the definite-source unfolding implication of the normalized convergent doubled mean and graph evaluation, the sharp principal-level coefficient, and the rank-one Euler and class-sum norm factor computation (`inner_trivialLift_self_eq_L`). PROVED IN CHAPTER 11: the exact convergent formula `siegel_weil` and graph evaluation `zeroTerm_graphSplitting` with the source joined-model and Haar conventions. OPEN: the clean nongraph support in general and the semilinear jet and normalized kernel identity `map_trivialLift`.

5. PROVED: sharpness from the integral formal parameter, finite coefficient maximization, the carried inverse, six-split ordinary bound, and the derivative splitting argument for admissible Frobenius pairings. OPEN: the actual weighted derivative assertion and its formal theta normal form where not established in the preceding CM chapter, PROVED: the packing and finite-datum indicator extraction with the prescribed archimedean bound on the cofinal family, conditional on the previously established normalized finite theta realization, weighted derivative assertion and vacuum estimate (Section 34.6).

6. OPEN: the genuine real square law with its kernel height and type convolution, and the global quantitative row restriction reaching the initial rows; the elementary tidy transformations and square cost recursion in Section 33.17 do not imply these assertions.

7. OPEN: fixed-carrier separation in the comparison theorem, including the actual block theta evaluation and the normalized line-scale estimate; tower vanishing and the normalized coefficient-to-period conversion for the tested quotient. The finite divided gluing and construction of the table operator were established in the preceding chapter and do not resolve this remaining analytic/geometric separation assertion.

In particular this ledger exhibits a number of genuine, reusable proofs, but does not identify the entire specialized C argument with a single published theorem, and should not be read as an unconditional claim of closure of C.

Chapter 35

Exact arithmetic pairings, rank-one values, and spectral projection

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 37, 38, 39 and the final audit in Chapter 57.

The remaining period argument involves several distinct kinds of comparisons. In this chapter we first close the arithmetic pairing, the rank-one value nonvanishing, the integral monomial and good-projection calculations, and the spectral projection bound at every complex embedding. They have exact statements in the compact source and can be derived independently of the deepest geometric period comparison. In particular the tested-vector hypothesis in the spectral argument is not the assertion that a particular geometrically constructed vector is tested.

35.1 Integral Gram pairings on the actual adelic coefficient lattice

Let E/F be the CM extension, and consider the standard definite Hermitian space I_n of rank n over E. Write $G_f = U(I_n)(\mathbb{A}_{F,f})$, K_0 for its standard maximal integral compact subgroup, and $\Gamma = U(I_n)(F)$. At a rational prime ℓ let K_ℓ be the subgroup of G_f whose components at places above ℓ lie in the standard integral groups, with no new restriction away from ℓ. For any coefficient embedding $j : \mathbb{C} \hookrightarrow \mathbb{C}_\ell$ write $O_j = \{a \in \mathbb{C} : |j(a)|_\ell \le 1\}$; here $\mathbb{C}_\ell$ denotes the completed algebraic closure of the local field, as in the source.

Lemma 35.1 (Simultaneous integral class representative)**.** *For every $g, d \in G_f$ there is $\gamma \in \Gamma$ with both*

$$\gamma g \in K_\ell, \qquad d\gamma g d^{-1} \in K_\ell.$$

Proof. The subset $K_\ell \cap d^{-1}K_\ell d$ is an open subgroup in the product of the local unitary groups at the finitely many places above ℓ (with the conjugate local factors understood together). It is a neighbourhood of the identity. It suffices to approximate g^{-1} by a rational point within $(K_\ell \cap d^{-1}K_\ell d)g^{-1}$ at those places.

Here weak approximation is elementary, not strong approximation for a simply connected derived group. In the Hermitian matrix algebra let $X^\dagger$ denote the involution adjoint to the Hermitian form and consider the F-vector space of skew-adjoint X. The Cayley map

$$X \longmapsto (1 - X)(1 + X)^{-1}$$

is an F-rational isomorphism of the open subset $\det(1+X)\neq 0$ onto the open subset of the unitary group where $\det(1+h)\neq 0$. Over every completion of F this latter open subset is dense in the analytic topology. Indeed multiply a given h by scalar norm-one elements arbitrarily close to 1 so that no eigenvalue becomes -1. There are only finitely many excluded scalars; the scalars $(1+a)/(1-a)$ for sufficiently small skew-conjugate a form infinitely many distinct ones (also at split places and at residue characteristic two, since the ground field has characteristic zero). Now use usual weak approximation for the finite-dimensional F-vector space: approximating finitely many Cayley preimages by a single rational skew-adjoint matrix and applying the map yields the desired approximation. This proves the assertion for all local signatures of the Hermitian form where used; the definite case suffices here. □

This is the exact `exists_mul_mem_integralAbove` of `Lib/Adelic/CuspForms/ClassReps.lean`. It also explains that a conjugated integral compact condition does not require a second independent rational change of basis.

Let M_κ denote the integral Schur-coordinate module, with integral symmetric positive definite Gram matrix G_κ as in Chapter 5. The weight representation R is the integral Schur representation times integer powers of determinants, interpreted on invertible matrices, including at local fields. For a vector u its gauge at g is

$$\|u\|_{g,j}=\Big\|R\Big((\widetilde{j}_\tau(g^{-1}))_\tau\Big)\,j(u)\Big\|_{\ell,\infty},\qquad \widetilde{j}_\tau|_E=j\circ\Phi_\tau\circ{}^{-}.$$

The local embeddings here are the finite-adele extensions at the places determined by the indicated embeddings of E. For any $k\in K_\ell$ the represented matrices of k and k^{-1} have integral coordinates, since Schur matrix entries are integer polynomials and determinant inverses are units. Hence they act by mutually inverse contractions of the sup norm and are isometries. In particular

$$\|u\|_{gk,j}=\|u\|_{g,j},\qquad \|R_\infty(\gamma)u\|_{\gamma g,j}=\|u\|_{g,j}. \tag{35.1}$$

The second equality follows from functoriality of the weight matrices under the coefficient embeddings: the represented matrix of $g^{-1}\gamma^{-1}$ cancels that of γ. These are `weylGauge_mul` and `weylGauge_embFin`. Scalar multiplication changes the gauge by $|j(a)|_\ell$, and sums obey the non-Archimedean triangle inequality.

Proposition 35.2 (Integral Gram and class-sum pairing)**.** *If $\|u\|_{g,j\circ\mathrm{conj}}\le 1$ and $\|v\|_{g,j}\le 1$, then*

$$\overline{u}^{\,t}G_\kappa v\in O_j.$$

If algebraic forms ϕ,ϕ' on I_n at level K satisfy these respective pointwise bounds and A is a positive integer divisible by all stabilizer orders a_c of the level classes, then

$$\left|j\left(A\sum_c a_c^{-1}\overline{\phi(g_c)}^{\,t}G_\kappa\phi'(g_c)\right)\right|_\ell\le 1.$$

Proof. By Lemma 35.1 with $d=1$ choose γ so that $\gamma g\in K_\ell$. Replace the two vectors by $R_\infty(\gamma)u$, $R_\infty(\gamma)v$. Their gauges are unchanged by (35.1); at an integral group point these gauges equal the coordinate sup norms. Thus every coordinate of the conjugate of the first vector and of the second vector lies in O_j. The matrix G_κ has integral entries.

The complex Gram pairing is unchanged under this replacement: at the infinite places γ is unitary for the definite standard form; the Schur contravariant Gram identity gives $R_\infty(\gamma)^*G_\kappa R_\infty(\gamma)=G_\kappa$. Thus the original Gram value equals the integral pairing of the two changed vectors. Applying the pointwise statement to every class proves the second assertion

because A/a_c is an integer and the valuation ring is closed under finite sums. No determinant denominator of G_κ is necessary. The same rational invariance after conjugating coefficients by $\sigma \in \mathrm{Aut}(\mathbb{C})$ follows by applying σ to the unitary matrix entries: on every image of a CM field it commutes with the CM involution, so the conjugated matrices are again unitary and the integral Gram identity is unchanged. □

This supplies the exact `gramForm_mem_embValRing` and `LevelForm.norm_inner_le_one` in `Lib/Adelic/CuspForms/LevelForms.lean`, the C01 interface. The definition of the level pairing uses the quotient measure with the mass at each class a_c^{-1}; that is why the prescribed stabilizer divisor, and not the degree of a coefficient basis, is the integer in this argument. These claims are about the definite coefficient forms themselves. They do not assert a complementary-rank theta exchange.

35.2 Positivity of the rank-one value ratios

In `Lib/ClassField/HeckeL/AutCharLFunction.lean` a character ν of $U(1)(\mathbb{A}_F)$ has weight $e = (e_\tau)$ if on the real torus it equals the product of the circle characters of exponents e_τ. Its base change to the Hecke ideles of E is

$$\psi(a) = \nu(a/\bar{a}).$$

On the infinite places of E this has angular exponents $2e_\tau$. It is unitary; for these CM characters its unitary central condition is part of the automorphic-character data. For the source normalization, $L(S, s, \nu)$ means the partial Euler product for ψ with the places of $S \cup \bar{S}$ removed (and the usual convention that a ramified local factor is 1).

Lemma 35.3 (Nonvanishing at the real edge)**.** *Let ν be such a character with some $e_\tau \neq 0$, and let S be finite. For every real $s \geq 1$, the partial L-value $L(S, s, \nu)$ is defined by the analytic continuation at $s = 1$ and is nonzero.*

Proof. Enlarge the finite set of removed places to a finite set T containing $S \cup \bar{S}$ and all the ramification of ψ and ψ^2. The two characters ψ, ψ^2 are both nontrivial on the norm-one ideles: on an archimedean circle at a place with $e_\tau \neq 0$ their angular exponents are $2e_\tau$ and $4e_\tau$. Thus neither is a pure norm character.

We use Tate's continuation theorem in its ordinary degree-one normalization: J. Tate, *Fourier analysis in number fields and Hecke's zeta-functions*, Ch. IV, §4.4 (Main Theorem, analytic continuation and functional equation), and §4.5 (comparison with the classical zeta- and L-functions), in Cassels–Fröhlich, *Algebraic Number Theory*, Ch. XV. It gives holomorphic continuation near 1 for each of these two partial Hecke products. It also gives a simple pole at 1 for the Dedekind zeta-function of E. For the exact finite product convention here, take the usual local factor away from T and remove only finitely many factors; the local archimedean gamma factors have no pole and no zero at real $s \geq 1$ in these unitary angular weights. Finite Euler factors restored at the omitted primes are also nonzero there. The source's continuation on the real axis agrees with this continuation at 1 because the Euler products agree on real $s > 1$, and their limits therefore agree.

For real $s > 1$ write the absolutely convergent logarithms of the Euler products. With $a = \psi(\varpi_w)^k$ for $w \notin T$, one has $|a| = 1$ and

$$3 + 4\,\mathrm{Re}(a) + \mathrm{Re}(a^2) = 2(1 + \mathrm{Re}\, a)^2 \geq 0.$$

After multiplying this inequality by q_w^{-ks}/k and summing over all $w \notin T$, $k \geq 1$, exponentiation gives

$$(\zeta_E^T(s))^3 |L^T(s,\psi)|^4 |L^T(s,\psi^2)| \geq 1. \tag{35.2}$$

The superscript means deletion of Euler factors at T. For $s > 1$ none of the three products vanishes, by their convergent logarithms. If $L^T(1,\psi) = 0$, holomorphy gives $L^T(s,\psi) = O(s-1)$, and $L^T(s,\psi^2) = O(1)$. The simple pole of ζ_E bounds its partial product by $O((s-1)^{-1})$; hence the left side of (35.2) tends to zero. This contradiction gives nonvanishing at 1. Restoring the finitely many omitted unramified factors preserves the result because $|q_w^{-s}\psi(\varpi_w)| < 1$ for $s \geq 1$; restoring a ramified factor does nothing. This proves the precise real-edge claim, without invoking any unconditional zero-free strip beyond the line $s = 1$. □

Corollary 35.4 (The positive scalar used in the repayment argument)**.** *Let $p = (\nu, z)$ be a source pair, so $1 \leq z \leq t$, ν has weight $-c_\tau$ with $z \leq c_\tau \leq 4b$. Let ϑ be a tame character of weight zero and modulus Q, and $\epsilon \in \{1, -1\}$. Put $M = \{w : D.\mathrm{IsChosen}(w)\}$. The number in* `Data.Pair.ratio` *is*

$$r = \frac{|L(M \cup Q, z, (\nu\vartheta^\epsilon)^{-1})|}{|L(M, z, \nu^{-1})|} \in \mathbb{R}_{>0} \subset \mathbb{C}.$$

In particular $r \neq 0$ and $\bar{r} = r$.

Proof. Both characters in the quotient have weight $(c_\tau)_\tau$, which is strictly positive at every infinite place, and $z \geq 1$. The sets of deletion of Euler factors are finite; all hypotheses of Lemma 35.3 hold. Thus both absolute values are strictly positive real numbers. The quotient here is a quotient of absolute values, not necessarily a quotient of the unmodified complex analytic values. The distinction is essential to what is claimed. □

The preceding proof covers `AutChar.HasWeight.L_ne_zero` and `ratio_ne_zero_and_real` in `New/Exchange/TriangleLevelExponent.lean`, i.e. C02. The source statement `ratio_eq_intAlgPart_div` additionally requires the exact period-to-L identity. Nonvanishing of the ratio does not by itself prove that a theta kernel realizes it.

35.3 Conjugate norm bounds for good translates and monomial vectors

We retain the source convention for a finite group G acting on columns at the selected places by right translation, and a central idempotent $e_0 \in \mathbb{C}[\mathsf{G}]$. Use $N_{\mathsf{G}} = |\mathsf{G}|$. For $s \in \mathsf{G}$ the good translate is the actual element

$$N_{\mathsf{G}} e_0 s z.$$

We explain both why this preserves the integral gauge and why its norm remains small at every coefficient automorphism. These arguments are independent of the particular construction of a good datum.

Lemma 35.5 (Finite central idempotents)**.** *Write $e = \sum_{g \in H} c_g[g]$ for a central idempotent of a finite group H. Then $|H|c_g$ is an algebraic integer for every g. For every $\sigma \in \mathrm{Aut}(\mathbb{C})$ the coefficientwise image e^σ is a self-adjoint idempotent in the involutive algebra with $[g]^* = [g^{-1}]$ and conjugation on coefficients. Moreover*

$$\sum_g |\sigma(c_g)|^2 \leq 1, \qquad \sum_g |\sigma(c_g)| \leq \sqrt{|H|}.$$

Proof. Over $\mathbb{C}$, the central idempotents are sums of the primitive central idempotents attached to irreducible characters. Matrix-character orthogonality gives the coefficient formula $|H|c_g = \sum_{\chi\in S}\chi(1)\chi(g^{-1})$ for a subset S of the irreducible characters. Every character value is a sum of roots of unity, hence integral. In any field automorphism the conjugated idempotent remains central. All irreducible representations may be unitarized by finite averaging, giving $\chi(g^{-1}) = \overline{\chi(g)}$; applying the primitive-idempotent formula anew to e^σ proves self-adjointness regardless of whether σ commutes with ordinary conjugation on other complex numbers.

Left convolution by e^σ is therefore a self-adjoint idempotent on the finite Hilbert space $\ell^2(H)$. Apply this orthogonal projection to the delta function at the identity, of norm one: its output has coordinates $\sigma(c_g)$ and squared norm at most one. Cauchy–Schwarz gives the final bound. □

Let ι_ϑ be the level embedding of the principal-level forms for twist ϑ in `New/Seesaw/BesselGeneric.lean`. A chosen representative $\widetilde{t} = \mathrm{groupLift}(\vartheta, t)$ is integral at every finite place, belongs to the standard maximal integral level, and has prescribed residue element t at the selected places. The action after applying the level embedding is exactly

$$\iota_\vartheta(N_{\mathsf{G}}e_0sz)(g) = \sum_{t\in\mathsf{G}} N_{\mathsf{G}}c_t\,\iota_\vartheta(z)(g\,\widetilde{ts}). \tag{35.3}$$

Indeed each group-ring basis element gives the indicated right translation: the independence of the integral lift follows because two choices have quotient in the defining congruence level, with the level character already absorbed into ι_ϑ; the action law converts the composite into the lift of ts. Linearity gives (35.3). This is `Data.iota_goodProjTranslate_apply`. It is an equality of actual coefficient values, not a formal dimension estimate.

Proposition 35.6 (Good projection is integral with the prescribed norm allowance)**.** *If $a\,\iota_\vartheta(z)$ is integral at $j : \mathbb{C}\hookrightarrow\mathbb{C}_\ell$, then $a\,\iota_\vartheta(N_{\mathsf{G}}e_0sz)$ is integral at j. For the trivial twist assume that every coefficient value of $\iota_1(z)$ has conjugate squared Gram norm at most e^B under σ. Then the conjugate squared class norm of the good translate at σ is at most*

$$\exp(x^{99} + B)$$

in the source data.

Proof. At a point g, the gauge of each translated value of the integral input is unchanged by switching the gauge to $g\widetilde{ts}$, by (35.1). The coefficients $N_{\mathsf{G}}c_t$ are integral at every embedding by Lemma 35.5. The non-Archimedean triangle inequality in the gauge applied to (35.3) proves integrality, including the same multiplier a.

For the norm estimate apply σ to (35.3) and use the usual Hilbert norm given by G_κ (the square root of the Gram quadratic form). The norm of the output at a point is at most

$$N_{\mathsf{G}}\sqrt{N_{\mathsf{G}}}\,e^{B/2}.$$

Thus the conjugate square norm after the class sum is at most

$$C_K N_{\mathsf{G}}^3 e^B,$$

where C_K is the number of level classes; the stabilizer weights are at most one. For the trivial twist the level in question contains the principal level of $\bar\pi_G\pi$ and the rank of Y_0 is at most n_m. The class count of Section 17.9 gives $C_K \le e^{x^{30}}$. By block-diagonal embeddings of the finite general linear groups, N_{G} divides the integer chosenGroupOrder of `New/Data/ChosenPlaceLevels.lean`. The recorded size bound there is

$$\log \mathrm{chosenGroupOrder} \le e^{32\log x} = x^{32}.$$

In the source size range this also follows directly from $|\mathrm{GL}_r(\mathbb{F}_q)| \le q^{r^2}$ and $\prod_{w \text{ chosen}} q_w = g_{\mathrm{prod}}^f$. Consequently $x^{30} + 3x^{32} \le x^{99}$, since the data have $x \ge 2$. This proves the result with ample slack; no estimate on an individual noncompact theta lift has been used. □

This proves `Data.goodProjTranslate_mem_integralForms` and `Data.normSqAt_goodProjTranslate_le` of `New/Rows/MonomialCoefficientVector.lean`.

We further give the integral coefficient-selector used for a rank-one block, to separate it from the later claim about which period it detects. Let the one-row weight at each place τ have degree $d_\tau = \mathrm{jetDeg}(\tau)$; its determinant exponent is zero. Let b_k be the product-minor integral basis. Here all nonempty minor factors have height one and the corresponding polynomial representation is the multihomogeneous polynomial space with variables the rows of a single column at each place. Let B denote the bilinear Fischer pairing, without conjugation, so that $B(X^\alpha, X^\beta) = \delta_{\alpha\beta}\alpha!$; put $h = \prod_\tau d_\tau!$. By the integral Gram identity of Lemma 5.7,

$$B(\mathrm{toPoly}(v), b_k) = h\, v_k. \tag{35.4}$$

The real oscillator identification in the rank-one data is precisely the rename of $\mathrm{toPoly}(v)$ to the Fock polynomial of v (in `New/Exchange/AlgebraicPartDefs.lean`, `LValueData.toPoly_eq_rename_harmonicPoly`). To see this in the present one-row case, the positive standard line has no negative coordinates and the mixed-Cartan embedding has only the one nontrivial positive factor, the negative Schur factor being the trivial representation. At the highest column the kernel is the product of its height-one entries to powers d_τ; renaming its rows to the source Fock letters is exactly the highest Fock monomial. Both maps intertwine the row substitutions of the general linear group and its contragredient coordinate action; the translates of a highest vector span each symmetric power, by polarization over characteristic zero. Equality at the highest vector therefore proves the asserted polynomial identification for all vectors. The same highest monomial has Fischer self-pairing h, so this argument uses the exact hook factor rather than a dimension multiplier.

Proposition 35.7 (Integral vector selecting a prescribed monomial)**.** *Let $e : \mathrm{Fin}(r) \hookrightarrow n$ select some rows and let $m = X^\alpha$ in those rows have total degree d_τ at each τ. Extend the exponent map to the full row set by zero. Then there is an integer coordinate vector v such that its Fock polynomial (with the source's star on v) is*

$$\frac{h}{\alpha!}\, m,$$

renamed to the corresponding Fock variables. Moreover for every characteristic zero field and every family of invertible matrices g_τ whose columns indexed by the chosen rows are the corresponding columns of the identity, the weight action fixes this vector.

Proof. Define v_k to be the coefficient of m in b_k. These are integers. Equation (35.4) and the Fischer monomial formula give

$$B\Big(\frac{h}{\alpha!} m, b_k\Big) = h v_k = B(\mathrm{toPoly}(v), b_k)$$

for every k. Nondegeneracy of the Fischer pairing on the homogeneous polynomial space therefore gives equality of the polynomials, and the rank-one renaming proves the Fock assertion. The factor $h/\alpha!$ is also visibly a product of multinomial coefficients.

For invariance, the row-substitution action $P(X) \mapsto P(gX)$ on polynomials and the transpose action are adjoint for B: $B(P(g^t X), Q(X)) = B(P(X), Q(gX))$. This holds by expanding linear substitutions or by differentiating the generating exponential $\exp(X^t Y)$;

tensoring gives it at all τ. The chosen-column condition implies $(g_\tau^t X)_{e(a)} = X_{e(a)}$. Thus $m(g^t X) = m(X)$, and the coefficient functional $Q \mapsto B(m, Q)/\alpha!$ on the polynomial module is invariant under $Q \mapsto Q(gX)$. In the source Schur-coordinate convention the vector v consists of this functional's values on the basis b_k (the Schur matrix is defined by expansion of the substituted basis rows). Hence its weight matrix fixes v. The determinant exponent is zero for the rank-one auxiliary data, so there is no additional determinant multiplier. The identities are polynomial over the integers, or follow over any characteristic zero field by the same finite expansions; in particular no choice of complex embedding is needed for the invariance. □

This is `exists_int_vector_harmonicPoly_eq`. It constructs the actual integer vector, not just a multiple of an unspecified vector of the correct weight. Together with Proposition 35.6 it discharges the C12 interface as listed in the previous ledger. Identifying the period of this vector and its orthogonality to all false lift spaces remains a separate geometric comparison, treated under C13–C14.

35.4 Hecke adjoints and projection at every embedding

We now prove the exact norm comparison in `New/Selmer/FitTwists.lean`. It applies to a tested vector, with the definition of tested as in that file, and makes no assertion yet about a particular test vector arising from theta periods.

Let W denote the finite-dimensional coefficient form space on the standard definite target, at the level of the auxiliary pair. The form $\langle u, v\rangle_\sigma$ is

$$\sigma^{-1}\left(\sum_c a_c^{-1}\overline{\sigma(u(g_c))}^{\,t} G_\kappa \sigma(v(g_c))\right), \qquad \sigma \in \mathrm{Aut}(\mathbb{C}).$$

It is sesquilinear for the involution $\tau_\sigma = \sigma^{-1} \circ \mathrm{conj} \circ \sigma$. Applying σ gives a positive Hermitian form on the vector space with scalar multiplication transported by σ; in particular diagonal values after applying σ are the real nonnegative numbers $\mathrm{normSqAt}_\sigma$ and vanish only for the zero vector. The same pairing can be described as the integral on the full finite adelic quotient, multiplied by the level index. Right translation is isometric in this integral, because the Gram form is invariant under rational changes of basis in the definite unitary group, also after σ, as in Proposition 35.2. The finite order level-character scalars have absolute value one after every field automorphism.

At a split off-level place w the local group is $\mathrm{GL}_n(E_w)$, with its ordinary maximal integral compact group K_w of Haar volume one. For $0 \le i \le n$ write $T_{w,i}$ for its basic spherical operator, the sum of right translations at the double coset of $\mathrm{diag}(\varpi_w I_i, I_{n-i})$ with each right K_w-coset counted once. Integration against the indicator of this double coset is the same operator on K_w-fixed vectors and needs no normalization beyond the stated volume. Under the invariant pairing its adjoint is convolution by the inverse double coset. The Hermitian relation in the split pair of local components identifies the inverse double coset at w with the basic double coset at the conjugate place $\bar{w}$ (inverse transpose, conjugation of uniformizer, and a unit change of uniformizer do not alter this double coset). Thus

$$\langle T_{w,i} u, v\rangle_\sigma = \langle u, T_{\bar{w},i} v\rangle_\sigma.$$

This directly proves `heckeSplit_adjoint` in `Lib/Weil/Unramified/SphericalHecke.lean`. Both vectors are spherical at these places, as is the entire auxiliary form space when the place is off the selected and twist levels.

The Hecke polynomial operator in this source is

$$H_w(X) = \sum_{i=0}^{n} (-X)^i q_w^{\binom{i}{2}} T_{w,i}.$$

Since the rational integer $q_w = q_{\bar{w}}$ is fixed by every automorphism, the exact adjoint identity is

$$\langle H_w(X)u, v\rangle_\sigma = \langle u, H_{\bar{w}}(\tau_\sigma(X))v\rangle_\sigma. \tag{35.5}$$

In particular this is `LevelPlaces.LevelForm.innerAt_heckePolyOnForms` with precisely its coefficient conjugation, not a unitary normalization of the Satake parameters.

For completeness, commutativity of these operators needs only the elementary transpose proof of the commutativity of the spherical Hecke algebra of GL_n. Smith normal form over the local discrete valuation ring shows that transposition preserves each K_w double coset: the diagonal elementary divisors are the same for a matrix and its transpose. Transposition preserves Haar measure and reverses convolution. Hence every compactly supported bi-K_w-invariant function is fixed by transposition, and the same is true of the convolution of two such functions. Reversal then proves that the convolution product commutes. At distinct split pairs the operators commute because the corresponding group factors do; at the two places of the same pair they are the just described spherical convolution operators and inverse ones. Thus $H_w(X)$ commutes with $H_{\bar{w}}(Y)$ for every X, Y on this form space. Equation (35.5) consequently shows that $H_w(X)$ is normal for every conjugate Hermitian structure.

Let E_0 denote the action of a central idempotent of the finite group at the marked places on W. It is a self-adjoint idempotent for each of these structures: the group action is isometric and, by Lemma 35.5, coefficientwise conjugation by any σ gives a self-adjoint central idempotent in the ordinary complex group algebra. Equivalently $E_0^* = E_0$ for the pulled-back adjoint. Its action commutes with the spherical operators off the level, because the finite group action is supported on the selected places and their conjugates. The extra residue-character multiplier in the auxiliary action is a root of unity and does not affect the isometry. This gives `AuxPair.innerAt_formAction`, `AuxPair.innerAt_groupAlgAction`, and the required operator commutation of `AuxPair.heckePolyOnForms_comm`.

Theorem 35.8 (Tested projection does not increase any conjugate norm)**.** *Fix one split off-level place w and the corresponding polynomial test value $X = \mathrm{AuxPair.testValue}(w)$, and write*

$$V_{\mathrm{test}} = \operatorname{im} E_0 \cap \ker H_w(X).$$

Let $Z \subset V_{\mathrm{test}}$ and suppose y satisfies the tested-vector hypothesis

$$y \perp (V_{\mathrm{test}} \cap Z^\perp)$$

for the ordinary (identity-embedding) pairing. Then the ordinary orthogonal projection pr_Z satisfies

$$\mathrm{normSqAt}_\sigma(\mathrm{pr}_Z\, y) \le \mathrm{normSqAt}_\sigma(y) \qquad \textit{for every } \sigma.$$

In particular the source hypothesis $\mathrm{IsTested}_{e_0, Z^\perp}(y)$, which gives the indicated condition at every split off-level place, implies the source conclusion.

Proof. Put $A = H_w(X)$. Since it is normal for every conjugate pairing, $\ker A = \ker A^*$ for every corresponding adjoint A^*. One way to see this is the elementary norm identity $\|Au\|^2 = \|A^*u\|^2$ for normal operators, applied after transporting scalars by σ. Thus the orthogonal complement of $\ker A$ for every conjugate pairing is $\operatorname{im} A$, an algebraic subspace

independent of σ. In the same way $(\operatorname{im} E_0)^{\perp} = \ker E_0$ for every pairing. Finite-dimensional orthogonal-complement algebra gives

$$V_{\mathrm{test}}^{\perp_\sigma} = \ker E_0 + \operatorname{im} A$$

for all σ, so this complement is also independent of σ.

For the ordinary inner product decompose $W = Z \perp (V_{\mathrm{test}} \cap Z^{\perp}) \perp V_{\mathrm{test}}^{\perp}$. The assumed orthogonality to the middle summand gives $y = z + u$ with $z = \mathrm{pr}_Z\, y$ and $u \in \ker E_0 + \operatorname{im} A$. The last summand is orthogonal to the whole space V_{test} for *every* conjugate pairing. In particular the two vectors z, u are orthogonal for every such pairing. After applying σ to the sesquilinear values, Pythagoras gives

$$\mathrm{normSqAt}_\sigma(y) = \mathrm{normSqAt}_\sigma(z) + \mathrm{normSqAt}_\sigma(u) \geq \mathrm{normSqAt}_\sigma(z).$$

This proves the result without claiming that pr_Z itself is an orthogonal projection for all the conjugate metrics on arbitrary vectors. □

Thus C15, `AuxPair.normSqAt_starProjection_le`, has the exact conjugate norm proof. A separate step must show that the particular geometric vectors used in the period argument satisfy the tested-vector hypothesis. The theorem is not a replacement for the theta-lift vanishing and support assertions C13–C14.

35.5 The equivariant difference of complementary theta integrals

We now supply an exact multiplier calculation missing from the earlier proof of the local matching factorization. Let W, V, V' have ranks n, a, b with $n = a + b$ even. Let S be the nonempty set of split places of a character datum $\mathfrak{c}$ in `SplitTameChar`. Put $F_S = \prod_{w \in S} E_w$ and $\chi(t) = \nu^{\mathrm{bc}}(t)$ for $t \in F_S^{\times}$, where the idele is extended by one off S. Write $|t| = \prod_{w \in S} |t_w|_w$; this is the product of multiplicative local norms. Write υ for the normalized splitting character on the ideles of E, and υ_1 for its restriction to the norm-one ideles. Recall

$$\upsilon(\bar{t}) = \upsilon(t)^{-1}, \qquad \upsilon_1^{\mathrm{bc}} = \upsilon^2.$$

Indeed υ is trivial on norms, because its restriction to $\mathbb{A}_F^{\times}$ is the quadratic character of E/F. Consequently for a character A on $U(1)$ write $A_r(t) = A^{\mathrm{bc}}(t)\upsilon(t)^{-r}$, in the source `AutChar.spaceChar`.

In the split model with row-variable $X \in \mathrm{Mat}_{a,n}(F_S)$ the pullback of the finite name under the pair action is $X \mapsto h^{-1}Xg$. Its normalized absolute-value factor is $|\det g|^{a/2}|\det h|^{-n/2}$: the determinant of the map $X \mapsto hXg^{-1}$ is $(\det h)^n/(\det g)^a$. The finite normalized unitary action of Theorem 7.6 gives the phase $\upsilon(\det h)^{-n}\upsilon(\det g)^a$. For the character-twisted action with characters (A, B) the additional pair character is $A(\det h)B(\det g)^{-1}$. At the places supported on S and their conjugates, $\det h$ on the unitary group is the idele determinant $t/\bar{t}$ when $t = \det h$ at S. Combining the three factors gives the exact split naming law

$$\omega_{A,B}(g,h)\Psi = \frac{A_n(\det h)}{B_a(\det g)} \frac{|\det g|^{a/2}}{|\det h|^{n/2}} \Psi(h^{-1}Xg). \tag{35.6}$$

Here the finite vacuum on the other split components is the one fixed in the source. One can check the norm factor directly against its characteristic function in the wave model: on L^2 the operator with this inverse linear change of variable has norm one. Covariance against

translations and characters identifies the operator up to its vacuum phase; the phase just computed by the finite normalization fixes it exactly. This is the formula of `pairWeil_name_on`, in `Lib/Weil/DualPair/PairSumNegation.lean`, without an unspecified scalar.

Take

$$A^- = \nu^{-1}\upsilon_1^{n/2}, \qquad A^+ = \nu\upsilon_1^{n/2}.$$

The larger integral is

$$L(\Phi)(x) = \int_{[U(V)]} \theta\big(\omega_{A^-,A^-}(x,h)(v\otimes\Phi\otimes\mathrm{vac})\big)\,dh,$$

and the smaller one $S(\Psi)(x)$ is the analogous integral over $[U(V')]$ with characters (A^+, υ_1^b), whose finite name reads Ψ^t through the conjugate places. Both v and v' are rapid real data. The letters L, S in this subsection denote the two integrals and should not be confused with the set of split places.

The full-rank support of the row and column Schwartz–Bruhat spaces is precisely the support needed for the estimate (10.2) of the growth chapter (or its transpose version). In that assertion the constants can depend on the fixed datum; no numerical uniformity in the field is needed here. It bounds the kernel uniformly in the group variable being integrated, up to a power of the height of the remaining group. Hence both integrals are absolutely convergent by finite quotient measure; their linearity, rational left invariance and continuity follow as in Section 10.7. In particular the linear maps are defined on the spaces of full-rank data.

The enlarged matching relation is for the character

$$\chi_n(t) := \chi(t)|t|^{n/2}.$$

Notice that $\chi_n = \chi$ on integral units, so every integral partner for χ in Chapter 33 is matched for χ_n using the finite primitive-support calculation there. To apply Proposition 33.2, choose h supported on the ramified places whose component at S is any specified element of $\mathrm{GL}_a(F_S)$; the component at $\bar{S}$ is dictated by the unitary relation. Since $(A^-)_n = \chi^{-1}$ on $F_S^\times$, (35.6) applied to $(1, h^{-1})$ gives

$$\chi_n(\det h)L(\Phi(h\,\cdot)) = L(\Phi).$$

The equality follows by the change of variable of right translation on the integrating quotient. Thus there is a linear row functional factored through the matched column functional, not just on individual standard data. Define M as the projection of the matched-pair linear space to its column component; it is stable under $\mathrm{GL}_n(F_S)$ by the matching transformation rule. Proposition 33.2 gives a linear map B on M with $B(\Psi) = L(\Phi)$ for every matched pair. Set $\mathcal{D} = B - cS$, for any fixed $c \in \mathbb{C}$.

Proposition 35.9 (Equivariant theta difference)**.** *The above $\mathcal{D}$ has the exact properties of `IsEquivariantLift`: it takes M to continuous automorphic functions, is invariant on the right under integral points trivial at S, and, for g supported on S with component G there,*

$$\Psi(G\,\cdot) \in M, \qquad \mathcal{D}(\Psi(G\,\cdot))(xg) = \rho(\det G)\mathcal{D}(\Psi)(x), \qquad \rho(t) = \upsilon(t)^{-b}|t|^{-b/2}.$$

For integral partners it equals the difference of the two specified periods, namely $L(\Phi) - cS(\Psi)$.

Proof. Take a matching pair (Φ, Ψ) and put $t = \det G$. Its translated pair is

$$\Big(\chi_n(t)^{-1}\Phi(\,\cdot\,G^{-1}), \Psi(G\,\cdot)\Big)$$

by the local transformation law for matched pairs (substitute Gk in the matching relation). The action of $(g,1)$ on the first component, by (35.6), multiplies it by

$$(A_a^-(t))^{-1}|t|^{a/2} = \chi(t)\upsilon(t)^{-b}|t|^{a/2}$$

and replaces its argument by XG. It therefore restores Φ and multiplies it by

$$\chi_n(t)^{-1}\chi(t)\upsilon(t)^{-b}|t|^{a/2} = \rho(t).$$

Covariance of the kernel and invariance of quotient measure give the asserted transformation of the larger theta integral.

For the smaller integral the source unitary component read at the conjugate places is G^{-t} in the coordinates of S. Therefore the name with transpose, after applying $(g,1)$ to $\Psi(G\,\cdot)$ read through the conjugate, is $\Psi(GG^{-1}X^t) = \Psi(X^t)$ as desired. The actual determinant at the conjugate places is $\bar{t}^{-1}$. The character on the source group is υ_1^b, so its space character in (35.6) is υ^b. Its multiplier is

$$\upsilon(\bar{t}^{-1})^{-b}|\bar{t}^{-1}|^{b/2} = \upsilon(t)^{-b}|t|^{-b/2} = \rho(t).$$

The same factor occurs on both integrals. Thus it occurs on their difference, and is independent of the choice of matched row function, which was already settled by Proposition 33.2.

For any integral right translation trivial at S, the finite vacuum is preserved, the component at the conjugate places is also trivial, and the named functions do not change. The splitting character is unramified and the tame part of ν is supported on S; hence the character multipliers are one. This proves invariance off S. Rational left invariance of the theta functional and absolute convergence give automorphy and continuity of each value on M as explained above, completing the proof. □

This proves `exists_isEquivariantLift` of `New/Exchange/ExchangeDifference.lean`. In particular the equivariant lift is constructed for every scalar c, even if $c = 0$. This is important for the nonvanishing argument. The abstract existence of a difference map is not the assertion that its value on a standard-column datum vanishes.

For later use we also verify the character of the centre coefficients of the difference. Let D be an isotropic splitting of W and $n(C)$ the subgroup in the centre of its unipotent radical, with additive coordinate C, with rational matrix Fourier index T in the source convention. The quotient Fourier coefficient of an automorphic f is

$$f_T(x) = \int_{[N_C]} \psi_T(C)\, f(n(C)x)\, dC.$$

The measure on this compact additive quotient is probability measure and the character is the fixed trace-dual character of the source. Continuity on the compact quotient justifies the integral, and translation gives $f_T(n(C_0)x) = \psi_T(C_0)^{-1} f_T(x)$. Explicitly the source's $\psi_T(C_0)$ uses the local additive character ψ of integral conductor and the matrix $(\mathrm{invDifferentIdele})^{-1}T$ read in the stated frame; this is the trace-duality identity of the mixed-model additive normalization in Chapter 9.

For a fixed x, the Fourier coefficient of $\mathcal{D}(\Psi)$ is a linear functional of $\Psi \in M$. A supported-on-S centre shear conjugated by x has matrix at S of the form

$$H = 1 + uZs, \qquad su = 0.$$

Here u and s are respectively the columns of the isotropic frame and the dual rows after passing through x, with $su = 0$; thus $H^{-1} = 1 - uZs$ and $\det H = \det(1 + Zsu) = 1$. Applying

Theorem 35.9 to this shear, with determinant multiplier one, and then changing the variable of the centre integral proves that the translated datum $\Psi'(Y) = \Psi(Y + uZsY)$ stays in M, and its Fourier value is $\psi_T(Z)$ times that of Ψ: the shear at x is the conjugate of the supported element $n(Z)$, and the equivariance gives the coefficient at $n(Z)^{-1}x$, whose factor is the inverse of the shift by $n(Z)$. With the source convention of `exists_linear_centreCoeff`, the relation is

$$L_T(\Psi') = \psi(\mathrm{tr}(T_S Z))L_T(\Psi),$$

where $T_S = (\mathrm{invDifferentIdele})^{-1}T|_S$ in that frame. Indeed the coefficient character is determined by the equality for $n(Z)x$ above; using the inverse shear reverses both Z and the character. This is the Fourier eigen-relation used in the local Gauss sum synthesis of Chapter 33. Algebraic extension of this linear functional from M to all functions, if desired, does not involve continuity or any new integral. The residual assertion in `IsEquivariantLif t.apply_stdColDatum_eq_zero` is not this eigen-relation, but its promotion from the known geometric coefficients to every Fourier index. In the source this promotion uses the bordered-coefficient induction `freitag` in `Lib/SiegelWeil/SingularForms/SingularForms.lean`; without that argument one cannot use just Fourier completeness on the centre to infer that all coefficients vanish.

35.6 Equal-rank constancy via strong approximation

The strong-approximation step in the equal-rank starting identity has a useful independent reference. We state it in a form that does not depend on the theta calculation. Let $W = H \perp W_0$ have a rational hyperbolic plane and let S be a nonempty finite set of split places of E, disjoint from its conjugate.

Proposition 35.10 (A constant-function criterion)**.** *Let f be a continuous automorphic function of $U(W)$, invariant by right multiplication with the full finite group supported at the split pairs of S. Suppose that whenever f is invariant under left multiplication by the unipotent radical of the line parabolic, the values $f(m(t, g_0))$ on its Levi are independent of the torus variable $t \in \mathbb{A}_E^\times$. Then f is constant. It is harmless also to impose right invariance at integral points away from S, as in the source.*

Proof. Use strong approximation for $H_1 = \mathrm{SU}(W)$ away from the finite split places of F underlying S. The precise result is the Kneser–Platonov strong approximation theorem: for a connected absolutely almost simple simply connected group over a global field, with a finite set of excluded places, density of rational points in the adeles outside those places holds if the product of the local groups there is noncompact. See A. S. Rapinchuk, *Strong approximation for algebraic groups*, in *Thin Groups and Superstrong Approximation*, MSRI Publications 61 (2014), Theorem 2.3, p. 281. The theorem allows a nonempty finite set of *finite* places; it does not require including the infinite places in the excluded set. Here $\mathrm{SU}(W)$ becomes SL_n after extension to an algebraic closure, is absolutely almost simple and simply connected for $n \geq 2$, and at an excluded split place is $\mathrm{SL}_n(F_v)$, noncompact. The adapted line splitting ensures $n \geq 2$. These check the hypotheses.

In adelic language, $\Gamma_1 H_1(\mathbb{A}_S)$ is dense in $H_1(\mathbb{A}_F)$, where $\Gamma_1 = H_1(F)$ and $\mathbb{A}_S$ denotes the components supported at the excluded places. For any h_S supported there and any adelic x, the element $x^{-1}h_S x$ is still supported there. Hence

$$f(\gamma h_S x) = f(h_S x) = f(x(x^{-1}h_S x)) = f(x) \qquad (\gamma \in \Gamma_1).$$

Continuity implies $f(hx) = f(x)$ for all $h \in H_1(\mathbb{A}_F)$. In particular it is invariant by left multiplication by the radical, which is contained in $\mathrm{SU}(W)$. By the hypothesis on the Levi, $f(m(t,1))$ is now independent of t.

The determinant of a Levi torus element $m(t,1)$ is $t/\bar{t}$. This map from ideles of E to norm-one ideles is surjective: apply elementary multiplicative Hilbert 90 in each local field (or in a split pair); at almost all places the norm-one element is integral and the preimage may be chosen integral invertible, by multiplying any preimage by a suitable element of the base local field (the quadratic extension is unramified or split there). Thus the preimages give an idele. Given any $x \in U(W)(\mathbb{A}_F)$ choose t with equal determinant. Then $x = h\, m(t,1)$ with $h \in H_1(\mathbb{A}_F)$, and we have shown $f(x) = f(m(t,1)) = f(1)$. □

This proves `IsAutFunction.apply_eq_apply_one`, `Lib/SiegelWeil/StrongApproximation/EqualRankConstant.lean`, with a matching strong-approximation reference.

In the equal-rank case of Theorem 35.9, take $V = W$, V' of rank zero. Its column datum is the scalar function 1 on the singleton zero matrix. The corresponding standard row datum is $\chi^{-1}(\det X)$ at integral invertible X, zero elsewhere, up to the ordering of the frame. Consequently $\mathcal{D}(1)$ for $c = 0$ is precisely the equal-rank theta integral of the source. Since $b = 0$, its multiplier $\rho = 1$ and its column datum is fixed by every $G \in \mathrm{GL}_n(F_S)$. The function satisfies the finite right invariance of Proposition 35.10, so in particular its dependence on x factors through the determinant. Its torus values are independent of the Levi variable by the line constant-term identity `exists_constant_term_thetaLarge`, if that identity is established on the actual normalized data: the case $\mathrm{rk}(W) = \mathrm{rk}(V)$ is precisely the case where the line identity in `LargeSideConstantTerm.lean` allows every line torus element t, and the lower-rank datum on the other side is independent of t. The radical invariance already proved allows replacing the radical average by the original value. It then follows from Proposition 35.10 that the equal-rank integral is constant, as in `thetaLarge_eq_apply_one_of_even_rank`. This does not assert that its constant is nonzero, and does not replace the normalized top-period and line identity still needed in C04–C05.

35.7 Convergence in the operator square and its remaining dependency

We complete the qualitative analytic bound in `New/Exchange/OneSquare.lean` for the exact data used there. Let U_1, U_2 be the exterior groups and A the intermediate group. The finite name in the direct-sum kernel on $(U_1 \oplus U_2, A)$ is the indicator of a prescribed residue matrix X, transposed to a row matrix on the source where required. If X^t has a left inverse over the residue ring over the ramified modulus, then at some split place every matrix in the support has full row rank: lift the invertible minor at that place, or lift the equation for a left inverse. This is enough to apply the full-rank-support growth estimate of Section 10.6; the ramified modulus is nonempty here because the tame character is nontrivial at its prescribed set. At fixed field, character and finite residue datum, and for every rapid real datum v, it gives

$$\left|K_v((g_1,g_2),a)\right| \le C\, m_N(v)\, H_{U_1\oplus U_2}(g_1,g_2)^B,$$

with no factor in $H_A(a)$. The same theorem with transposed support gives the bound for the smaller intermediate group, since a matching partner's primitive column matrix has a left inverse. Constants may depend on the test and the finite datum, as they do in the source. No uniform bound over all twists or all fields is needed for this qualitative argument.

For $y_i \in U_i$.rapidFunctions, by definition $C_i := \sup |y_i(g_i)| H_{U_i}(g_i)^B < \infty$. The direct-sum height bound $H_{U_1\oplus U_2} \le H_{U_1} H_{U_2}$ follows from the direct-sum lattice and the max norm, using

heights at least one. The character factors on the two exterior variables have absolute value one. Hence the product integrand in the actual square satisfies

$$|y_1(g_1)y_2(g_2)K_v((g_1,g_2),a)| \le CC_1C_2m_N(v).$$

All three quotient measures have finite mass, by the finite-volume argument of Section 10.7. This proves `exists_integrable_kernel_bound` with the hypotheses of `one_square`; it also supplies dominated convergence for the approximating rapid datum, at each test separately.

We recall why the finite and real algebraic parts of `one_square` require no further *independent* analytic hypothesis once a complementary exchange and the source's edge hypotheses are given. First the box-product property of the oscillator gives precisely `modKernel_sum_left` and `modKernel_sum_right`: with group elements block diagonal, the tensor of the normalized vacua is the vacuum of the direct sum, the named finite functions multiply on disjoint blocks, the Fock polynomials multiply after relabelling variables, and the determinant pair characters multiply on block diagonal matrices. The theta functionals multiply under this tensor identification because the rational lattice is the direct sum of the two rational lattices and the normalized theta functional of a tensor product is the product (Chapter 9). Thus the exact kernel of an elementary polynomial tensor is the product of the two edge kernels, with no dimension-dependent multiplier. The character identity in the square hypotheses is exactly the identity of the remaining exterior pair character.

Second the condition `IsEdgeFamily` in `Lib/SiegelWeil/RallisInner/SharpData.lean` assumes `IsEdge` for the indicated joint harmonic. The latter includes, as a defining clause of `IsHalfEdge`, that the lift using any polynomial Q is a real convolution with a compactly supported continuous function of the lift using the fixed joint harmonic, and the convolution function is independent of the finite twist, modulus and adelic data. Its swapped half-edge includes the analogous assertion on the conjugate types. This is the hypothesis under which the source states `one_square`. The statement that the particular chosen joint harmonics in the grid satisfy this edge hypothesis belongs to the discrete-series and oscillator-pair construction, not to a separate proof of the operator square.

Now for an elementary tensor polynomial on the two real Hom factors, the edge hypothesis and the pairing identity for real convolution reduce its integrated product, on every test i, to a common scalar times the prescribed lift–descent composition. Fubini is justified by the preceding bound for full-rank residue data. The constant is common to all tests since the real convolutions and the normalized real vector are independent of the finite twists; the compact-type Petersson matrix coefficient producing the scalar is evaluated only on the prescribed real type. Its precise value is immaterial, and it may vanish. For arbitrary rapid exchanged data, simultaneous polynomial approximation in Lemma 33.6 and the demonstrated pointwise-in-test continuity give the closed-one-dimensional-span argument of Section 33.6. Each summand of the polynomial approximation is an elementary tensor because monomials factor in the two disjoint sets of variables. Thus one scalar κ works for the entire family, including a family with all tests zero.

Finally the unit covariance in the two point variables of `SquareLaw` follows from the paired naming formula (35.6) on integral group elements of the appropriate rank, on which the absolute-value and splitting-character factors are one. For example translating the row matrix x on the left by $u \in \mathrm{GL}_a(R)$ can be lifted to the corresponding integral source or target translation at the intermediate group. The two edges have inverse changes of the integration variable in that group; their determinant-character product is $\chi(\det u)$ by the explicit $a_ib_i = \mathrm{rankTwist}(\nu_i^{-1})$ hypothesis. The same statement on the right for the partner matrix gives $\chi(\det v)$. Applying the normalized complementary exchange to the summed partner data and pairing with a test vector, whose rapidity makes the two integrals absolute,

now gives the partner equality with multiplier $\kappa/c(\nu)$: the up–down tested integral is κ times the up–down composition, while the complementary exchanged side is $c(\nu)$ times the down–up composition. The nonzero $c(\nu)$ is the exchange scalar; its asserted norm is exactly the norm from that theorem. Arbitrary linear functionals follow because a Hermitian pairing with all test vectors separates the finite-level form space (equivalently, the two continuous square compositions take values in the same Hilbert representation and the tested vectors separate there).

This proves the growth and integration portion of C06 outright and the square law as an implication of C04 and the edge hypotheses, as in its exact source statement. The unproved normalized complementary exchange cannot be replaced by a vanishing test with a guessed scalar. In particular the analytic proof here requires no continuation of the Hecke L-values beyond the positive real evaluation points in Section 35.2.

35.8 Terminal branching and the integral end period

The remaining algebraic comparison at the terminal grid nodes is much easier than the initial noncompact restriction problem. At the two terminal nodes the groups are standard definite. By Proposition 33.10 their parameters, with the appropriate sign reversed for the other grid, satisfy the horizontal-strip interlacing for a rank-d group embedded in one of rank $d+1$. We detail both the pairing and its denominator estimate.

Lemma 35.11 (Integral dual of the coefficient lattice)**.** *For two Schur-coordinate weights with opposite Harish–Chandra parameters in the same rank, there is an integral bilinear pairing $A(v', v)$ invariant under their simultaneous weight representations, with the following properties:*

1. *Its matrix is invertible over $\mathbb{C}$ and has an integer left inverse up to multiplication by the hook factor of the first coefficient lattice.*
2. *For every v the bilinear functional $A(\,\cdot\,, v)$ is the Gram pairing against a vector of the same Gram norm as v.*
3. *The compact coefficient realization of the opposite weight is obtained by this pairing with the translated highest vector of the original weight.*

These are the claims of `exists_dual_pairing_of_isHCParam`*.*

Proof. Write the column multiplicities of a highest-weight diagram as $\kappa_\tau(h)$, and separate its determinant exponent. In rank d, reversing and negating the decreasing Harish–Chandra parameter reverses the highest weight and negates it: if the original highest weight is (λ_i), the opposite weight is $(-\lambda_{d+1-i})$. The consecutive differences therefore replace $\kappa_\tau(h)$ by $\kappa_\tau(d-h)$ for $0 < h < d$. The determinant exponent changes by the sum of those column multiplicities and the negative of its original value, exactly the twist required for taking the dual.

On basis minor-products use the Jacobi complementary-minor identity

$$\det(g)\det((g^{-1})_{I,J}) = (-1)^{\sum I+\sum J}\det(g_{J^c,I^c})$$

in an ordered basis. Reverse the chosen ordered flags along with the complements. Applying it to each column replaces the corresponding height h by height $d-h$, with the determinant twist just computed. This gives the inverse-transpose intertwiner on the Schur polynomial modules over every characteristic zero field. The matrix changing the integral Schur coordinates is an

integer matrix J: the transformed products are integer signed minor-products and expand by the integral straightening identity. Applying the complementary operation a second time recovers the original products, with the inverse determinant twist canceling and the signs of the chosen bases restored. Hence the inverse matrix also has integer entries. Equivalently one can characterize this map on Weyl evaluation vectors; on the evaluation vector of the ordered highest flag it is the evaluation vector of the reversed flag, with the sign fixed to send the former to the latter. These vectors generate their modules over $\mathbb{C}$, so this normalization determines the map and agrees with the integral complementary-minor construction.

The pullback of the second coefficient Gram form is a compact-invariant positive form on the irreducible module of the original coefficient weight, by the contragredient intertwining identity. Schur's lemma shows that it is proportional to the first Gram form. Both ordered highest-flag evaluation vectors have squared Weyl Gram norm one (evaluate the kernel that defines the Gram form at the permutation arrays). Since J takes one of those vectors to the other, the proportionality constant is one: in compatible ordering $J^tG'J = G$. This proves the integral Gram isometry `exists_dual_isometry`.

The desired pairing matrix is $G'J$, with the dual determinant exponent already incorporated. Its invariance follows from the contragredient identity and Gram contravariance. For an integer left inverse up to the hook number h', use J^{-1} times the integral Fischer matrix F': $F'G' = h'I$ by Lemma 5.7. Thus $J^{-1}F'$ times $G'J$ is $h'I$. The vector $\overline{Jv}$ represents the functional $u \mapsto u^tG'Jv$ in the Hermitian Gram pairing and has unchanged norm. The coefficient realization is obtained by evaluating this pairing at the translated highest vector; at the identity ordered flag this is the dual highest coefficient by the normalization of J, and equivariance gives the formula at every compact real point. All these operations are exactly the determinant-twisted integral ones in `Lib/GLn/Dual/DualWeights.lean`. □

Proposition 35.12 (Terminal pairing matrix)**.** *There is an integer matrix B with the three properties used in `exists_branching_pairing` of `New/Seesaw/PairTensorData.lean`: writing M for a bound on the number of boxes of the larger weight,*

$$(R'(g)v')^tBR(g \oplus u)v = \left(\prod_\tau \det(u_\tau)^{N_\tau}\right)(v')^tBv,$$
$$|N_\tau| \le M + |e_\tau|,$$
$$|(v')^tBv| \le (M^M)^3\|v'\|_G\|v\|_G.$$

Here the first representation is the opposite weight in the smaller rank, u is the additional one-dimensional block, and $\|\cdot\|_G$ is the corresponding ordinary Weyl Gram norm. The terminal compact period of two coefficient-realized forms is a scalar, depending only on the real translate and the weights, times the class integral of this pairing divided by the index of its level in K_0.

Proof. Use the interlacing of Proposition 33.10 and Theorem 17.5, which constructs an integer branching matrix b from the larger to the smaller Schur-coordinate module of the interlacing highest weight, with operator norm at most $(M^M)^3$ in its Weyl metrics. The smaller opposite-weight module is paired to that weight by Lemma 35.11. Compose its integer pairing matrix A with b, and put $B = Ab$. Equivariance of b for the rank-d group, together with determinant twists for the extra line, gives the first formula. The line character on this branch has exponent equal to the difference of the total polynomial degrees at each place plus the determinant exponent in the larger rank. The difference is the number of boxes in a horizontal strip, between 0 and M; this gives the bound on N_τ. The two norm bounds combine by Gram Cauchy–Schwarz and the isometry of the dual pairing, giving the third assertion. This is precisely the integer matrix construction as opposed to a rational nonzero intertwiner obtained by clearing an unspecified denominator.

For the period, the compact model of Lemma 35.11 writes the two real coefficient evaluations as pairings of fixed coefficient vectors with translated highest vectors. The integral of their product over the compact unitary group of rank d is the projection onto invariant tensors in the product of the two representations, by compact averaging. The codimension-one branching rule says the intertwining space is one-dimensional, spanned by b; the determinant character of the additional line is recorded as above. Thus the result of this averaging is a scalar times the actual integer bilinear pairing B, the scalar depending on the chosen two highest vectors and real translate only, not on their variable coefficient vectors, finite places, or twist. This proves the exact compact period conversion `integral_algFormCoeff_blockEmb` (also used in the final branching calculation of Section 17.8). Decompose the finite adelic quotient at a level $K \subset K_0$; its class measure with $\mathrm{vol}(K) = 1$ is the class integral, and changing to $\mathrm{vol}(K_0) = 1$ multiplies by $[K_0 : K]^{-1}$. The additional free finite line parameter is simply the stated argument of the second coefficient form at the block-diagonal point. This gives the asserted level-index factor and proves the proposition. □

Write $\mathfrak{m}(\vartheta)$ for the grid modulus, and $\beta_\vartheta = \mathrm{flipScalar}(\mathfrak{m}(\vartheta))$. At stage m in the two grids $z = 0, 1$ let $K_{z,j}$ be their end levels, $0 \le j \le m$. By definition of the source terminal period there are chains C_z of m successive algebraic theta kernels with the finite point names given by an edge tuple S. Their output is paired by B at h and the corresponding point $h \oplus \beta_\vartheta^m$. The terminal class integral is at level $K_{0,m}$ with volume of that level one. A tidy functional is a sum $\sum_S c_S$ of at most T^n evaluations, $T = |\mathcal{O}_E/\mathfrak{m}|^{(2n_m)^2}$, of order n, and every nonzero coefficient c_S is a root of unity.

The equivariance needed to expand this period is worth checking before making any class-sum expansion. For an edge with source and target finite points g, h, right translate by their corresponding level elements $k_s, k_t \in K_0$. The paired naming law for integral elements, already established in Section 35.5, replaces the named indicator of the residue matrix S_e by that of $\bar{k}_t S_e \bar{k}_s^{-1}$ (bars here denote residue reduction, not complex conjugation) and multiplies it by

$$\chi_e(\det \bar{k}_t / \det \bar{k}_s).$$

Both edge characters have the same tame part χ_e by the defining rank twists. The original scalar is therefore the product of these edge characters over the two terminal chains. The tuple of all the new matrices is exactly the inverse vertex action on the old tuple, since that action in `Data.tupleAction` is $S_e \mapsto g_t^{-1} S_e g_s$. At the start of each chain its input form is fixed by the stated end level; at the two terminals the vertex actions are compatible by the block embedding with the additional line fixed. Thus these level elements give an admissible vertex family. The defining equivariance of the tidy functional multiplies its value under the inverse action by the inverse product character, canceling the displayed product. Rational left invariance follows from the algebraic kernel equivariance, and for the terminal pairing also from the first formula of Proposition 35.12 with $u = 1$. Consequently the full sum of integrands is jointly invariant in every finite variable under its required left rational and right level groups. One must use this invariance of the full tuple sum, rather than assert that each individual point-name kernel is right invariant.

All the indicated levels are free levels by the chosen-level stabilizer bound (they are contained in the principal chosen level and their ranks are at most $2n_m$). Choose their finite class representatives. The jointly invariant terminal integral with respect to its level-one measure is exactly a finite sum over these representatives, one term per class, and the same holds simultaneously for every internal integral in both chains. The algebraic lift-chain expansion used here is the one in Chapter 17: kernels are linear and their successive class-integral

expressions compose by finite sums. The result is a sum of at most

$$T^n \prod_{z=0,1} \prod_{j=0}^{m-1+\mathbf{1}_{z=0}} \#\mathrm{StdClasses}(K_{z,j}) \tag{35.7}$$

scalar pairings of the terminal vectors obtained by applying one sequence of jet matrices to an initial vector from each chain; each pairing is multiplied by a root of unity. This is the exact `endPeriod_mem` of `New/Exchange/GridFirstColumns.lean`: any condition propagated by every jet matrix and valid at the two starting forms therefore holds at the terminal vectors of all these summands. In particular it bounds the single summand after every field embedding without multiplying by a measure normalization depending on the twist.

For clarity we also describe the conversion from the analytic stage-m grid period to this terminal expression. At this stage no initial descents remain. All vertices of both chains are standard definite, so the compact coefficient lift identity of Section 17.6 gives one scalar independent of ϑ for each step, times its algebraic lift with its indicated input-level normalization. Iterating it gives for grid z up to stage j the factor

$$c_{z,j} \frac{[K_0 : K_{z,j}]}{[K_0 : K_{z,0}]}$$

multiplying the algebraic chain in the real coefficient realization. Indeed each step gives the input level-index inverse, and the scalar recurrence absorbs its untwisted target level index and the factor $\mathrm{vacuumScale}^{n_j n_{j+1}}$ of the end datum. This can be done independently of ϑ since $j+1>0$: at all such nodes, and at every node of grid 1, the level inside the twisted residue ring is the kernel of the surjective reduction to the untwisted residue ring. Its index is consequently exactly the order of the untwisted general linear group, by the first isomorphism theorem. This is `levelIndex_eq_untwisted`. At $j=0$ the comparison is the identity, and the factors telescope to the displayed quotient. Apply the compact period conversion in Proposition 35.12; it divides by the last level index of grid 0. The grid-period definition contains the initial level index of grid 0, canceling its remaining denominator. The ratio of the last to first level indices of grid 1 is independent of the twist, so it joins the real branching scalar and the two chain scalars. By linearity of Λ this proves `gridPeriod_eq_endPeriod`: there is one constant independent of the entire family of twists and forms. It is not required to be nonzero for this implication.

Theorem 35.13 (Terminal integral and conjugate bounds)**.** *Use the source end forms and tidy data. There is a positive integer N_*, depending only on the source parameters, $N_* \le \exp(x^{99}b^2)$, such that N_* times the terminal period is integral at every coefficient embedding over a prime ℓ prime to $\mathrm{N}(Q(\vartheta))$ whenever the starting forms are integral there. For the trivial twist and order $n = 4m + 2m^2$, if every conjugate squared norm of the starting values of the two forms is at most B_0, B_1 respectively, the conjugate of the terminal period has absolute value at most*

$$e^{x^{200}b^2}\sqrt{B_0}\sqrt{B_1}.$$

These are the bounds of `Data.exists_mul_endPeriod_mem_embValRing` and `Data.endPeriod_le`.

Proof. Use $M = b_1 r^4$, the source bound `boxes_endWeightData_le` on the total number of boxes of each terminal weight (also $|e_\tau| \le \lceil 20b \rceil$ and ranks at most $2n_m$), and set

$$A_0 = (g_{\mathrm{prod}}^{b_1} P.d)^M, \quad A_1 = g_{\mathrm{prod}}^{b_1} N_1, \qquad N_* = A_0^{2m} \prod_\tau A_1^{m|N_\tau|}.$$

The grid modulus contains the rational integer $\mathrm{N}(Q)g_{\mathrm{prod}}^{b_1}$ because $\bar{\pi}_G\pi_G = g_{\mathrm{prod}}$ and each ramified factor contains its absolute norm. The finite point indicator on the scaled integral vacuum is standard datum at that modulus, with no additional denominator: its values are zero or one, and its Heisenberg averaging expansion uses the integral vacuum, as in Chapter 17 and `smul_vacuumDatum_mem_standardData`. Apply the integral jet-raise assertion `mem_raises_of_dvd`, proved there: multiplying the jet of degree at most M by $(\mathrm{N}(Q)g_{\mathrm{prod}}^{b_1}P.d)^M$ does not increase the unit gauge. At $\ell \nmid \mathrm{N}(Q)$ the extra power of $\mathrm{N}(Q)$ is a unit, so the same holds for A_0. Induction along each chain consequently makes every terminal gauge integral after multiplying its vector by A_0^m.

For the pairing at h and $h \oplus \beta_\vartheta^m$, transport to the integral coordinate vectors by applying their inverse adelic components as in the gauge. Proposition 35.12 transports the pairing itself, up to the scalar $\prod_\tau j_\tau(\beta_\vartheta)^{mN_\tau}$ or its inverse (the choice of inverse frame does not affect the denominator bound). Both β_ϑ and its inverse, after multiplication by $\mathrm{N}(Q)A_1$, are integral at every finite place. Indeed $\beta_\vartheta = \bar{u}/u$ with $u = \alpha \cdot \mathrm{generator}(\mathfrak{m})$; the generator is integral, $\mathrm{N}(Q)g_{\mathrm{prod}}^{b_1}$ times its inverse is integral, and $N_1\alpha, \alpha^{-1}$ are integral by the half-different scale calculation of Chapter 6. Conjugating gives the other direction. At ℓ prime to $\mathrm{N}(Q)$, this means each local component of either of the two reciprocals has denominator dividing A_1. Hence $\prod_\tau A_1^{m|N_\tau|}$ clears the determinant-character factor in the pairing. Its matrix entries are integers, so the product N_* times each terminal summand is in O_j; the root-of-unity tuple coefficients are also in O_j. The finite sum (35.7) proves the integral assertion.

Here are explicit size checks, using ample room in the allowed coarse exponents. Write $\ell_b = \log b$. The magnitudes of the source, as in Sections 17.9 and 17.10, imply $r \le x^{5.1}$, $m \le x^{0.6}$, $2n_m \le x^{5.6}$, $f, N_1 \le x^{11.1}$, $b_1 \le e^2 b$, and $\log g_{\mathrm{prod}} \le 22\ell_b$; $\ell_b \le x^{\cdot 01}$, $P.c, P.d \le \ell_b$, and x and b are in the extremely large range of the scale inequalities. In particular $M \le bx^{21}$, $\log(g_{\mathrm{prod}}^{b_1}P.d)$ and $\log A_1$ are at most $bx^{\cdot 2}$, and $|N_\tau| \le M + \lceil 20b \rceil \le bx^{22}$. Therefore

$$\log N_* \le 2x^{\cdot 6}b^2x^{21+\cdot 2} + x^{11.1+.6+22+\cdot 2}b^2 < x^{99}b^2.$$

For the complex bound put $G = e^{x^{99}b^2}$ and take the trivial twist. At every jet step equation (17.4) with $L = 1$ and $c = 1$ gives an operator norm bound G in the conjugate Gram metric. Here the size of the finite point name in the admissible-combination estimate is at most $|\mathcal{O}_E/\mathfrak{m}|^{nn'}$ by the finite Heisenberg expansion of a named indicator; at the trivial twist this residue ring has cardinality $g_{\mathrm{prod}}^{fb_1}$, so its contribution to the logarithm is at most bx^{24}. The real conjugate-jet bound contributes $P.c(N_1nn' + 1)^2 \le x^{47}$ and $fnn' \log 4 \le x^{24}$. The sum of the nonzero row degrees is at most M, their factorial-log sum at most $M \log M$, and $u = \log(\Omega^{-1}) \le 13 \log x$. Consequently the remaining exponent is at most $\frac{1}{2}M(1 + \log M + 13 \log x) \le bx^{23}$. These estimates give the claimed upper bound G for the jet norm. The same integer pairing matrix B has operator bound $(M^M)^3$ under every σ, because its entries and the Gram matrices are integers. Its logarithm is at most $3M \log M \le bx^{23}$. After the m steps on each side, every terminal summand therefore has conjugate absolute value at most

$$(M^M)^3 G^{2m} \sqrt{B_0}\sqrt{B_1}.$$

Each class count in (35.7) is at most G by the bound of Section 17.9, with rational principal modulus $g_{\mathrm{prod}}^{b_1}$: its logarithm is at most $x^{29}(1 + b_1 \log g_{\mathrm{prod}})$. The short-sum count is T^n with $\log T = (2n_m)^2 f b_1 \log g_{\mathrm{prod}} \le bx^{24}$. For $n = 4m + 2m^2$ and $2m + 1$ class factors, the total logarithmic allowance other than the two starting norms is at most

$$3M \log M + n \log T + (4m + 1)x^{99}b^2 \le x^{200}b^2.$$

Root-of-unity coefficients have modulus one under every σ. Summing the bounds of the individual terminal pairings proves the stated conjugate bound. □

The terminal estimates here only require the source's tidy, fixed-level and standard definite hypotheses, and the integral compact coefficient and jet constructions established in the earlier chapters. They do not establish that the noncompact initial periods reach the terminal path with those hypotheses. That is the separate C08–C09 transport problem. In particular the terminal argument cannot serve as a proof of the complementary exchange used to establish that transport.

35.9 Updated obligations for the period comparison

We record precisely which mathematical interfaces of the previous edition have been addressed in this chapter. Here a proved theorem includes all the hypotheses stated in its source declaration; conclusions requiring separate geometric hypotheses are not thereby proved.

PROVED — C01, integral Gram pairing.
The simultaneous integral representative by the rational Cayley map, exact equivariance of the gauge, Gram integrality with the conjugate coefficient embedding in the first slot, and class-sum integrality after multiplication by the stabilizer divisor are proved in Lemma 35.1 and Proposition 35.2. This is the interface of `ClassReps.lean` and `LevelForms.lean`.

PROVED — C02, nonzero real ratio.
Lemma 35.3 uses the exact type and partial-product normalization of `AutChar.L`. Tate's continuation theorem plus the displayed logarithmic Euler-product inequality proves nonvanishing including the point 1. Corollary 35.4 proves `ratio_ne_zero_and_real` for every source pair and tame character. The extra identity with a normalized theta period is separate.

OPEN — C04, C05, complementary-rank and top period.
The exact equivariant difference and finite multipliers have been derived in Section 35.5. Proposition 35.10 proves the equal-rank constant-function criterion by strong approximation, with the precise theorem and local noncompactness hypothesis verified. Constancy of the equal-rank theta integral follows once the actual normalized line identity has been established (see the separate tower discussion). Remaining dependencies of `exists_stdExchange_top`, `exists_equalRankThetaRatio_value`, `exchange_of_partners` and `exchange_theorem` are the actual top mixed-model coefficient identity including the matched real slicing period, and the promotion of its geometric coefficient vanishing to all indices by the exact singular-form induction `freitag`. These have not been replaced by generic nonvanishing or an arbitrary theta identity.

CONDITIONAL — C06, the square law.
The full-rank-supported kernel estimate and finite-volume Fubini justify the uniform-in-the-intermediate-variable integrability in Section 35.7. The scalar consistency of the tensor approximation and finite point-datum covariance establish `one_square` as an implication of the precise complementary exchange theorem and the stated `IsEdgeFamily` hypotheses (which include real convolution). The actual joint-harmonic edges belong to the separate oscillator/discrete-series argument.

PROVED — C10, terminal periods.
Lemma 35.11, Proposition 35.12 and Theorem 35.13 give the integer pairing, level-normalized compact branching, joint tidy invariance and the actual sum expansion of `endPeriod_mem`, and the prescribed integer and conjugate bounds. The comparison `gridPeriod_eq_endPeriod` is the terminal compact-chain step; it does not give a route

from the initial noncompact periods. In particular further work on C08–C09 is still necessary.

PROVED — C12, good projection and monomial coefficient vector.
Proposition 35.6 gives the exact pointwise integral gauge and class norm estimate, preserving the common multiplier. Proposition 35.7 produces the integer coordinate vector with the specified hook-to-factorial polynomial and its invariance. These are the stated propositions of `New/Rows/MonomialCoefficientVector.lean`; identification of the tested period belongs elsewhere.

PROVED — C15, conjugate spectral projection.
Equation (35.5) and Theorem 35.8 prove the precise polynomial adjoint and projection comparison for the `IsTested` vectors in `FitTwists.lean`. The proof uses the independence of $\ker E_0 + \operatorname{im} H_w(X)$ from the coefficient automorphism, not that arbitrary complex conjugation commutes with that automorphism. Showing a particular constructed vector is tested remains a geometric task.

CONDITIONAL — C24, real cone and matched theta difference.
Section 35.5 establishes the exact difference map and Fourier eigen-relation from the actual convergent normalized theta integral, including the determinant powers on the conjugate split places. The earlier positive real cone construction applies under its stated hypotheses. The remaining step is `IsEquivariantLift.apply_stdColDatum_eq_zero`, in particular the `freitag` bordered-coefficient induction and its hypotheses for the actual geometric difference. Neither ordinary Fourier completeness nor the local determinant Gauss-sum synthesis alone proves those hypotheses.

C03, C07–C09, C11, C13 and C14 retain their previous status except insofar as the precise downstream dependencies are reduced by the results above. We have not thereby proved a full global period exchange or a new complete proof of the main theorem.

Chapter 36

Clean doubling, sharp Euler factors and normalized line integrals

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 38, 39 and the final audit in Chapter 57.

The following arguments continue the construction of the adelic oscillator and its arithmetic use in Chapters 9, 11 and 34. The convergence of the ordinary Siegel–Weil formula is not a substitute for the vanishing of nongraph terms, the absolute integrability of the Rallis unfolding, or the finite-support cutoff needed to deduce injectivity. We supply those steps in the source's normalizations. The same calculations apply to a general rapid form, not only to a cuspidal source in a regular holomorphic weight.

36.1 Clean data remove all nongraph orbits

Let E/F be the CM extension, let S, T be the source and target Hermitian spaces, and set $V = \mathrm{Hom}_E(S, T)$. Write $n = \dim_E S$, $m = \dim_E T$. The doubled source is $S \oplus (-S)$. Let $D(f', f)$ denote the joined datum of Chapter 11.6, conjugate linear in f' and linear in f. We use the following definition exactly as in `IsClean` of `Lib/SiegelWeil/RallisInner/SharpData.lean`: a datum f is clean if S is definite, or if

$$f = \mathrm{name}_P(\Phi) f_0$$

for a set P of finite places disjoint from its conjugate, a Schwartz–Bruhat function Φ whose support consists of matrices injective at a specified place $w \in P$, and an integrable datum f_0 in the complementary factors. The support condition means that every matrix M with $\Phi(M) \neq 0$ is left-invertible after evaluation at w. In particular this definition is a condition on one finite factor, not a condition of global absolute rank on all the adeles.

Theorem 36.1 (Nongraph vanishing on clean joined data)**.** *Assume f is clean. For every adelic source point g, every integrable f', and every rational maximal isotropic subspace $K \subset S \oplus (-S)$, if*

$$Z_K(D(f', \omega_{a,b}(g, 1) f)) \neq 0,$$

then K is the graph of a rational isometry of S, with the graph-sign convention of Chapter 11.6. The graph parametrization is injective.

Proof. Write a rational frame of K in two blocks $(A; B)$ of n columns. If B is invertible, its graph map is AB^{-1} and isotropy implies

$$(AB^{-1})^{*} J_S (AB^{-1}) = J_S.$$

This is an isometry. Reversing B gives the source convention $(\delta u, -u)$; in particular two such isometries cannot give the same subspace, by projection to the second summand. If B is singular, choose $c \neq 0$ with $Bc = 0$. Since the whole frame has full rank, $u = Ac \neq 0$, and $(u, 0) \in K$. In the definite case this is impossible, because $u^{*} J_S u = 0$ and a definite Hermitian form has no nonzero isotropic vectors.

In the other case consider the zero-term functional pulled back through the join and the fixed Weil translate,

$$\ell(v) = Z_K\big(D(f', \omega_{a,b}(g, 1)v)\big).$$

It is a bounded linear functional on the integrable datum space. In a rational complete splitting with first space K, the K-translations act on the wave function at zero by the trivial phase. More explicitly the Heisenberg law (9.11) implies

$$Z_K\big(\rho(kz) F_1\big) = Z_K(F_1)$$

for the whole adelic space of Hom translations along K, where k is its embedded isotropic frame in the normalized hyperbolic coordinates. Taking the line $(u, 0)$ in K gives translations that affect only the first joined factor. To see their parameters in the original Hom coordinates, let x_S be the matrix changing the rational source form to the normalized form, put

$$Y = (g^{-1} x_S^{-1} u), \qquad X = T^{\mathrm{dual}},$$

where X is the invertible dual-frame matrix of the target. The actual translated displacement is the set of matrices $X z Y^{*} J_{\mathrm{std}}$ for all adelic columns z. Indeed, under the Hom identification the linear functional of the rational line u on the source is $u^{*} J_S$. In the standard coordinates its transpose-adjoint becomes $(x_S^{-1} u)^{*} J_{\mathrm{std}}$; pulling the translations through $\omega(g, 1)$ replaces $x_S^{-1} u$ by $g^{-1} x_S^{-1} u$ because $g^{*} J_{\mathrm{std}} g = J_{\mathrm{std}}$. The dual-frame change X is invertible and accounts only for the target coordinates; the second joined component is zero since $Bc = 0$. This proves invariance of ℓ under those translations. Character twists in the pair Weil action are scalar and do not affect the conjugation of translations.

For a matrix M of the clean support we have

$$X_w^{*} M_w Y_w \neq 0.$$

Indeed $Y_w \neq 0$ because the adelic matrices g_w and $(x_S)_w$ are invertible and the rational vector u remains nonzero on embedding into E_w; M_w is injective and X_w^{*} is invertible. Lemma 10.5 now applies. To spell out the annihilation: choose the translation parameter supported only at the opposite split place so that its action is multiplication on the selected finite variables by the character of the pairing with $X_w^{*} M_w Y_w$. By nondegeneracy of the local additive character, it is not the trivial multiplier at that point. In a sufficiently small compact open neighborhood it is a fixed scalar different from one. The compact support of Φ has a finite disjoint compact open refinement of these neighborhoods. On each piece, invariance of the bounded functional says its value equals itself times that nontrivial scalar, hence the value is zero. Summing yields $\ell(f) = 0$, the desired contradiction. □

The complete-splitting coordinates just used also identify the Lean calculation `Space.coordMatrix` and its nonvanishing lemma `mul_coordMatrix_ne_zero`. In `exists_graph_eq_of_zeroTerm_ne_zero` the matrix denoted K' is our B, and the dual frame X is

the one used by `name_eq_zero_of_invariant`; in these formulas it is crucial that the finite support be left-invertible, not merely nonzero. The theorem proves that declaration and `graphSplitting_lagrangian_injective`. No regularized Siegel–Weil or cuspidality argument is needed for this vanishing.

36.2 The Rallis identity with absolute convergence

We write $G = U(S)(\mathbb{A}_F)$, $\Gamma = U(S)(F)$ and $[G] = \Gamma\backslash G$. All Haar measures are those of the common normalization in the previous chapters, with the quotient by the counting measure on Γ; the volume of $[U(T)]$ will be denoted by μ_T. Petersson forms are conjugate linear in their first argument. For the theta lift by f of a scalar automorphic function y put

$$\mathcal{L}_f y(h) = \int_{[G]} y(x)\theta(\omega_{a,b}(x,h)f)\,dx.$$

The two rapid automorphic functions y, y' below are assumed continuous and to decrease faster than every power of the source height (equivalently the bounds of `rapidFunctions`); their finite-level invariance is not needed for the next theorem.

Theorem 36.2 (Clean convergent-range Rallis formula)**.** *Assume $2n < m$, y, y' are rapid, and f is clean. Then both lifts have square integrable Petersson values and the function*

$$r(g) = \langle f', \omega_{1,b}(g,1)f\rangle \langle y', R(g)y\rangle_{\mathrm{Pet}}$$

is absolutely integrable on G. Here $R(g)y(x) = y(xg)$. Moreover

$$\langle \mathcal{L}_{f'} y', \mathcal{L}_f y\rangle_{\mathrm{Pet}} = \mu_T \int_G r(g)\,dg. \tag{36.1}$$

Proof. We provide the absolute bounds before writing the unfolding equality. The height reduction of Section 10.3 gives a measurable choice $\widetilde{x}$ of representative of $x \in [G]$ with

$$\|\omega_{1,b}(\widetilde{x},1)\|_{1\to 1} \le C H_S(x)^A.$$

Indeed the successive compact right dilations in that proof take any point to a set of bounded height; translating its endpoint by a rational element puts it in a fixed compact set by the Mahler bound. Thus one of its rational translates lies in that compact set multiplied by at most $1 + \log H_S(x)/\log 2$ compact factors. The estimate in the displayed inequality is exactly the product of the operator bounds of those factors. A measurable section is obtained by taking the first rational translate satisfying the operator bound in an enumeration of the countable group, after first choosing a Borel fundamental domain. Such a domain exists by taking countably many charts in the second-countable group, each with mutually disjoint rational translates, and retaining the points not covered by an earlier chart. This proves the claimed measurable reduction, including for points on boundaries.

Put

$$F_x = y(x)\omega_{1,b}(\widetilde{x},1)f, \qquad F'_{x'} = y'(x')\omega_{1,b}(\widetilde{x'},1)f', \qquad \mathcal{D}_{x',x} = D(F'_{x'}, F_x).$$

Both norms of $F_x, F'_{x'}$ are integrable on $[G]$ by rapid decay, $H_S \ge 1$ and finite quotient volume. The norm of the box product is the product norm in the integrable model (and the negation map has norm one). The product of the two norms is integrable on $[G]^2$. By Chapter 11.5 there is a fixed integrable positive function $\Theta_0(h)$ such that

$$\left|\theta(\omega(1,h)\mathcal{D}_{x',x})\right| \le C\|F_x\|_1 \|F'_{x'}\|_1 \Theta_0(h).$$

In particular it is integrable jointly in x, x', h. Its value is exactly the product

$$\overline{y'(x')\theta(\omega_{a,b}(\widetilde{x'},h)f')}y(x)\theta(\omega_{a,b}(\widetilde{x},h)f).$$

Here the a-characters on the target cancel against their opposites in the conjugate join, and the remaining pair characters b and $b^{-1}\upsilon^m$ are the same product convention as in Chapter 11.6. More formally the splitting character on the doubled source is the product $a(a^{-1}\upsilon^n) = \upsilon^n$; the standard split Weil action uses this character and 1 on the two opposite determinants. Thus this is the exact product identity with the phase of the join chosen in Chapter 11.6, and not an arbitrary scalar multiple. Rational covariance means its product is independent of the choice of representatives. The same argument with the two data equal shows that each lift belongs to $L^2([U(T)])$ by Tonelli and Fubini. This separately proves the existence of the Petersson left-hand side.

Next put $A_K = Z_K$ for the zero terms on the doubled space. Chapter 11.1 gives $\sum_K \|A_K\| < \infty$, so

$$\int_{[G]^2} \sum_K |A_K(\mathcal{D}_{x',x})|\,dx\,dx' \le \left(\sum_K \|A_K\|\right)\left(\int_{[G]} \|F_x\|_1\,dx\right)\left(\int_{[G]} \|F'_{x'}\|_1\,dx'\right) < \infty.$$

Theorem 36.1, applied to the clean first datum after its adelic source translate, removes all nongraph terms. If K corresponds to $\delta^{-1} \in \Gamma$, the graph evaluation of Chapter 11.6, unitarity, and the scalar factors in the two data give

$$A_K(\mathcal{D}_{x',x}) = \overline{y'(x')}\,y(x)\langle f', \omega_{1,b}(\widetilde{x'}^{-1}\delta\widetilde{x},1)f\rangle.$$

For fixed x', the substitution $g = \widetilde{x'}^{-1}\delta\widetilde{x}$ unfolds the integral over $x \in [G]$ and sum over $\delta \in \Gamma$ into the integral over G. It is measure preserving: the translates of a fundamental domain establish the quotient integration formula for counting measure on Γ, and left multiplication by $\widetilde{x'}^{-1}$ preserves Haar measure. Also $y(x) = y(\widetilde{x'}g)$ by automorphy. The absolute integral proved above equals

$$\int_{[G]}\int_G |y'(x')y(x'g)\langle f', \omega_{1,b}(g,1)f\rangle|\,dg\,dx' < \infty.$$

The triangle inequality gives absolute integrability of r. Removing the absolute values is therefore legitimate, and integrating first in x' yields $\int_G r(g)\,dg$.

Finally the ordinary convergent Siegel–Weil formula applied to $\mathcal{D}_{x',x}$ equates the h-integral of the product with μ_T times the sum of its zero terms. Both are integrable over $[G]^2$ as just proved. The unfolded equality is (36.1). □

This is the proof of `inner_product_formula` in `Lib/SiegelWeil/RallisInner/SharpData.lean`, using exactly the summability of the zero-term operator norms and the normalized torus majorant proved previously, rather than a formal interchange of three infinite integrations. It does not require the clean datum to be cuspidal. The separation between the arbitrary source determinant character a and the $1, b$ pair in the matrix coefficient is essential for the later dependence on twists.

36.3 Finite vacuum cutoffs at arbitrary finite places

We next give the local dual-datum construction of the source. This is more than just the sharp principal-level formula: the datum will cut out any prescribed finite set of *unramified* places

in the integral while leaving all other matrix coefficients unchanged. It is used before taking a limit of the partial integrals. Work in the standard finite Hom coordinates of Chapter 7. The finite lattice is $L = \widehat{\mathcal{O}}_E^{m\times n}$, with the self-dual trace pairing and normalized Heisenberg translations $P(u) = \lambda(u)\rho(u)$ of Section 7.3. Thus $P(u)e = e$ for $u \in L$, and the normalized Weil operator $W(k)$ carries these fixed-vector equations to $P(ku)W(k)e = W(k)e$. The commutator of the P operators is the additive character of the alternating trace form. We use its exact conductor (self-duality of L), including at dyadic primes, but no division by two in this argument.

Lemma 36.3 (Dual-datum cutoff)**.** *Suppose $m \geq 2n$ and A is a finite set of finite places. Let f be vacuum above A, meaning its projection to that vacuum factor is itself (including the places conjugate to A). There exists an integrable datum f'_A such that for every $g \in U(S)(\mathbb{A}_F)$,*

$$\langle f'_A, \omega_{1,b}(g,1)f\rangle = \begin{cases} \langle f, \omega_{1,b}(g,1)f\rangle, & g \text{ integral at } A,\\ 0, & \text{otherwise.}\end{cases}$$

Proof. We may replace A by its conjugate closure, because for a unitary matrix integrality and invertibility at every place of a conjugate pair are equivalent to its two matrices being integral. The normalized finite model splits as the tensor product of the vacuum factor above this set and the complementary factor. The vacuum projection hypothesis says $f = e_A \otimes f^{(A)}$ in these coordinates, with the real factors assigned to the complement. We construct the desired vector at A. Denote its finite lattice by L_A and the vacuum by e_A.

At these finite places the standard Hermitian target is the sum of m norms. As in Lemma 6.1 there is a unit of the finite integral norm ring with relative norm -1. Consequently we may take the integral isotropic $m \times n$ frame X with successive blocks $I_n, cI_n, 0$ where $\bar{c}c = -1$. The row block I_n gives the integral matrix $P_0 = (I_n; 0; 0)$, with

$$X^*X = 0, \qquad P_0^*X = I_n.$$

All that matters is the finite normalized target form; the frame is not required to be rational. This is also the construction of `exists_integral_isotropic_frame`. The zero-rank source case has scalar identity and requires no construction.

Choose a purely imaginary finite adele z supported on A, with $|z_w|_w > 1$ for every $w \in A$. For example take the fixed imaginary scalar $\jmath$ multiplied by the inverse of a positive rational integer divisible by sufficiently large powers of the finitely many residue characteristics. Let

$$k = I_m + zXX^*.$$

Then k is unitary because $X^*X = 0$ and $\bar{z} = -z$, and its inverse is $I_m - zXX^*$. It acts by left multiplication on Hom and commutes with every right source action. Consider $v = P(P_0)W(k)e_A$, in the integrable finite model. Since $P_0 \in L_A$,

$$\langle v, e_A\rangle = \langle W(k)e_A, e_A\rangle \neq 0$$

by the explicit finite vacuum coefficient formula of Chapter 7. Rescale v so that the last inner product is one. This decides the exact normalization on the compact subgroup without fixing any phase independently of the preceding chapters. The datum is finite (and hence integrable), because both the translated vector and the Weil transform of the vacuum are supported modulo an open finite lattice on finitely many cosets.

If g is integral, its inverse is integral on all of A; its normalized finite Hom action fixes the vacuum line. Hence the normalized v has exactly the same pairing with $\omega(g,1)e_A$ as does e_A, including its determinant phase.

It remains to show that the matrix coefficient is zero for g nonintegral. We suppress the determinant phase, which is nonzero. Denote the right action by $R(g^{-1})Y = Yg$; by allowing all g and g^{-1} it suffices to consider this orientation. If $\langle v, W(R(g^{-1}))e_A\rangle \neq 0$, the two normalized Heisenberg eigenrelations imply

$$\psi_0\big(\operatorname{tr}(P_0^* q)\big) = 1 \quad (q \in kL_A \cap R(g^{-1})L_A), \tag{36.2}$$

where the imaginary trace pairing and different factors of Chapter 7.3 are understood in ψ_0. In detail, the second vector is fixed by $P(q)$ on $R(g^{-1})L_A$; the first is an eigenvector there on kL_A, with eigenvalue given by the commutator of $P(P_0)$ and $P(q)$. Pairing the eigenrelations and using their norm-one eigenvalues forces the displayed phase to be one. The sign of the trace may be inverted depending on the commutator convention, which does not change this condition. The pairing is precisely the same nondegenerate one for which the integral lattice is self-dual.

Suppose that some entry of g_w is nonintegral. Pick a row i and an entry in column j of maximal size $M > 1$ in that row. Let σ be a scalar supported at w with local size

$$|\sigma|_w = (\max(M, |z_w|_w))^{-1}.$$

Such a scalar exists by the discrete valuation. Let $E_{j,i}$ denote the matrix having a single 1 in row j of the chosen identity block of the target and source column i. For any local integral scalar d, set $Y = d\sigma E_{j,i}$ and $q = kYg$. Since Yg and kY are integral (the largest possible sizes are $|\sigma|_w M$ and $|\sigma|_w|z_w|_w$), using also the integral matrices X, X^*, we have $q \in kL_A \cap R(g^{-1})L_A$. All other local entries vanish. But its commutator trace contains the phase of

$$\operatorname{tr}(P_0^* kYg) = d\sigma(1 + z_w)g_{ij}.$$

The term involving 1 alone is integral for every integral d, whereas

$$|\sigma z_w g_{ij}|_w = \frac{|z_w|_w M}{\max(M, |z_w|_w)} > 1.$$

Self-duality of the additive lattice implies that this nonintegral scalar defines a nontrivial character on the integral d's. This contradicts (36.2). Here d can be supported at this place in the entire finite adele ring, and the conjugate-coordinate trace gives the same nontrivial character by its nondegeneracy. Thus all entries of g are integral above A. Unitarity and the inclusion of conjugate places now imply integrality of its inverse as well. This proves vanishing on the required complement and strict equality on the selected integral subgroup.

Finally tensor v with $f^{(A)}$; outside A the matrix coefficient is unchanged, and the source real part is unchanged as well. This gives the claimed f'_A. □

In the proof q can simply be written kYg; writing it as κ times a right translate in the Lean file emphasizes the two different lattice membership tests. The nonintegral scalar of size $\min(M, |z_w|_w)$ explains why a compact-open averaging alone could not give this cutoff. The argument proves `exists_dual_datum` in `SharpData.lean`. It applies at nonsplit and dyadic factors because it uses only an integral norm-(-1) element, the skew scalar shear and self-duality. It is unrelated to a choice of a small spherical matrix coefficient with nonnegative values.

36.4 Sharp local data and the absolutely convergent Euler sum

We use the sharp condition of `IsSharp` at a conjugate-stable finite set S_0: the finite part F is nonzero, is vacuum outside S_0, every joining $v \otimes F$ is clean, and for g integral off S_0 the finite

coefficient equals

$$\langle F, \omega_{1,b}(g,1)F\rangle = \|F\|^2 \mathbf{1}_{J\cap K_0}(g).$$

The compact subgroup K_0 is the integral finite group, of Haar volume one. The sharp mass is $c = \|F\|^2/[K_0 : J \cap K_0]$. In the source J is the appropriate principal level; the coefficient identity and cleanness for the residue data follow from Lemma 34.4, the support with fixed left inverse, and Theorem 36.1. A residual left inverse lifts to one over the integral ring by multiplying a lift by the inverse of a matrix congruent to the identity modulo the prime. This verifies full-rank support even when the source form is not definite. Also $J \cap K_0$ is open: the nonzero locus of the continuous matrix coefficient contains a neighborhood of the identity in K_0, and equals $J \cap K_0$ by the displayed identity. Hence $0 < c < \infty$.

For an integrable function r on the adelic source group and a set A of finite places, write $\mathrm{PI}_A(r)$ for its integral on the points integral at every place of A. For fixed B and finite R increasing to all places,

$$\mathrm{PI}_{R\setminus B}(r) \longrightarrow \mathrm{PI}_{B^c}(r). \tag{36.3}$$

This is dominated convergence: the indicators decrease to the indicator of the intersection, and all are bounded by one. One can choose an increasing enumeration of the countably many places; the same proof gives the limit over the directed finite sets, by monotonicity on the nonnegative absolute difference. All sets are measurable since the integral local-level condition is open and closed on the restricted product (and arbitrary intersections in question are countable).

Let v be the chosen real datum, $f = v \otimes F$, and suppose y is invariant by J and y, y' are rapid. If u is any finite point, Theorem 36.2 and the sharp coefficient give

$$\mathrm{PI}_{S_0^c}\left(\langle f, \omega(\,\cdot\,,1)f\rangle\langle y', R((1,u)\cdot\,)y\rangle\right) = c\int_{G_\infty} \langle v, \omega_\infty(x,1)v\rangle \cdot \langle y', R(x,u)y\rangle\, dx. \tag{36.4}$$

For $u \neq 1$ this identity is applied to translates of a rapid form, hence the relevant matrix-coefficient product is integrable by the same theorem. The factorization follows by taking the real and finite factors of the product group: the determinant characters are separated into their archimedean weights and finite parts; the finite coefficient vanishes except on $J \cap K_0$, where right invariance removes the k-action on y. Integration over that compact subgroup gives exactly the mass c. This proves `IsSharp.partialIntegral_eq` in `Lib/Weil/CompactLift/RallisFormula.lean` using Fubini on an absolutely integrable function.

At places outside S_0, choose representatives t_ξ supported off S_0 for the left cosets of the finite subgroup integral at every place outside S_0. Each coset is open; the disjoint decomposition therefore has countably many cosets, by second countability. Set

$$\gamma(t) = \left(b(\det t)\Delta_f(t)^m\right)^{-1},$$

which is the source's `vacuumCoeff` (we work with the finite multiplier of the source group; the Hom action is its m-fold inverse). For k integral off S_0,

$$\langle f, \omega((1,t_\xi)k,1)f\rangle = \gamma(t_\xi)\langle f, \omega(k,1)f\rangle.$$

In fact the factor off S_0 is a vacuum there, so the local integral action takes it to a scalar phase times itself; the translate of its coefficient is that phase times the finite vacuum coefficient. The exact formula for that coefficient follows from (17.2) and the determinant and inverse in the pair action. There is no dimension factor from the complementary level. Summing the integral over these cosets, and changing variables by left multiplication, therefore gives the absolutely convergent identity

$$\int_G r(g)\, dg = \sum_\xi \gamma(t_\xi)\, \mathrm{PI}_{S_0^c}\left(\langle f, \omega(\,\cdot\,,1)f\rangle\langle y', R((1,t_\xi)\cdot\,)y\rangle\right). \tag{36.5}$$

Absolute convergence is stronger than convergence of the scalar series: the sum of the absolute values of the integrals is bounded by $\int_G |r|$ since the cosets are disjoint. The identical argument for a finite conjugate-stable slice A of the places gives `hasSum_partialIntegral_coset`; equality (36.3) then gives the source's unramified-limit step.

Suppose now that v is a specified real edge between the two source types of the monograph, with the real convolution normalization $a_v \neq 0$; its meaning is

$$\int_{G_\infty} \langle v, \omega_\infty(x,1)v\rangle \langle y', R(x,1)y\rangle \, dx = a_v \langle y', y\rangle$$

for forms of the source type. This is the real identity in `IsEdge` and the normalized Petersson convolution formula; the present proof uses that hypothesis and does not infer noncompact edge existence from the general doubling theorem. Finite right translation preserves the real type, commutes with the real action, and consequently the integral with u in (36.4) equals $a_v\langle y', R(1,u)y\rangle$. Inserting into the Rallis formula and (36.5) proves the convergent sum

$$\langle \mathcal{L}_f y', \mathcal{L}_f y\rangle_{\mathrm{Pet}} = \mu_T a_v c \sum_{\xi} \gamma(t_\xi)\langle y', R(1,t_\xi)y\rangle. \tag{36.6}$$

Thus the factor $A = \mu_T a_v$ is nonzero and independent of the finite twist among the prescribed real weights; no Euler-product comparison of complex periods is needed to infer its nonvanishing.

There is a separate injectivity implication, which does not follow from positivity of each term of the last sum. Assume $2n < m$ and $\mathcal{L}_f y = 0$. For any finite R disjoint from S_0, apply Lemma 36.3 with $A = R$ and use its f'_R in the left and right sides of Theorem 36.2, paired against y' of the same source type. The clean datum is still f, not the auxiliary dual datum. The left side is zero and $\mu_T > 0$, so

$$\mathrm{PI}_R\left(\langle f, \omega(\,\cdot\,,1)f\rangle \langle y', R(\,\cdot\,)y\rangle\right) = 0.$$

Exhaust the places outside S_0 by (36.3). Equation (36.4) for $u = 1$ yields $ca_v\langle y', y\rangle = 0$. Take $y' = y$; the Petersson norm is positive definite on the space of continuous forms and $ca_v \neq 0$. This forces $y = 0$.

These are the full formal implications of `rallis_inner_product_euler`, including the dual-datum cutoff necessary for injectivity. They assume the actual real edge and its type convolution; that assumption is the ordinary hypothesis of the Lean theorem. In the definite rank-one case the edge is just the compact monomial calculation of Chapter 17, so (36.6) applies without a new indefinite-edge input. In higher ranks, the existence and exact noncompact normalization of the particular real edges used in the arithmetic chain remain separate obligations.

36.5 The normalized semilinear rank-one kernel

We next resolve the arithmetic step that must accompany the rank-one Rallis norm. A complex field automorphism need not commute with ordinary conjugation, and it is incorrect to apply it to an analytic Petersson integral. Instead we use the finite jet form and its algebraic class sum.

Fix the coefficient field and a conjugate-disjoint set of chosen finite places, and take the untwisted auxiliary set Q empty. Put $n = 2z > 2$ and let X be the rank-one data `LValueData` of `New/Exchange/AlgebraicPartDefs.lean`. Denote its jet degrees at the complex places by d_τ, with $D = \sum_\tau d_\tau$. Thus the source is the definite line with characters 1 and

$$b_X = \mu_X v^z.$$

The finite named datum F_X for the lift is the indicator of the chosen standard column at the places of M read modulo the level, by the displayed definition `AuxPair.resDetDatum`. The residue condition at a place of Q would carry a character, but here no such places are present. In particular this finite name is independent of the varying μ_X and consists of the rational scalars $0, 1$. Pullback to the squared-level version of the residue ring changes neither the named function nor its support.

Let $\sigma \in \mathrm{Aut}(\mathbb{C}/K)$ be one of the permitted automorphisms, inducing a permutation e of the complex places. Let Y denote the same data with degrees $d_{e^{-1}\tau}$ and with character μ_Y whose restriction to the finite unitary group is $\sigma \circ \mu_X$, as in the source hypotheses. Write Ξ_X for `LValueData.trivialLift`, the finite sum of the Schur jet kernel at scale 1 over the line classes of chosen level. Let per denote the fixed jet scale in the CM construction.

By the semilinear comparison of Theorem 17.2, with the exact jet-series comparison preceding it, there is a σ-semilinear M on the finite model, and scalars t, ℓ and a norm-one finite idele a_0 with

$$\sigma(T_{\mathrm{per}} f) = e^* T_t(Mf), \qquad |t|^2 = \Omega^{-1}, \qquad \ell \neq 0, \qquad MF_X = \ell W(a_0 I_n) F_Y.$$

Here e^* means the rational relabelling of the formal variables, Ω is the fixed positive `bigOmega`, and T_c is the scaled jet series. These constants depend on the standard Hom space, the selected set and σ, not on X with its particular degrees and Hecke character. The third equality comes from the named-vector assertion of Theorem 17.2, applied to the indicator just described.

For a finite unitary matrix A put

$$\rho_\sigma(A) = \frac{\sigma(\Delta_f(A))}{\Delta_f(A)}.$$

The identity $MW(A) = \rho_\sigma(A) W(A) M$ for the genuine finite Weil representation implies that ρ_σ is a character: applying the two ways of multiplying two matrices to any f for which $Mf \neq 0$ proves multiplicativity. Also

$$\rho_\sigma(A)^2 = \frac{\upsilon(\det A)}{\sigma(\upsilon(\det A))}. \tag{36.7}$$

This is the exact identity `multiplierRatio_sq`; it follows directly from $\overline{\Delta_f} = \upsilon(\det A)\Delta_f$ and the fact that $\Delta_f \overline{\Delta_f}$ is a positive rational integer, so that it is fixed by σ.

For a finite source scalar g and target point h, write $C(g, h)$ for the matrix of the Hom action, explicitly $h(\cdot)g^{-1}$ in the standard finite coordinates. Then

$$C(g, h) = C(1, h)\, g^{-1} I_n, \qquad \rho_\sigma(g^{-1} I_n) = \rho_\sigma(g^{-1})^n.$$

The last assertion is the block-product formula for the rank-one finite multipliers. Since $n = 2z$, (36.7) gives

$$\sigma(b_X(g))^{-1} \rho_\sigma(g^{-1})^n = \Big(\sigma(\mu_X(g))\sigma(\upsilon(g))^z\Big)^{-1} \left(\frac{\sigma(\upsilon(g))}{\upsilon(g)}\right)^z = b_Y(g)^{-1}.$$

This is precisely why the even-rank twist eliminates the source multiplier; there would be an unaccounted square root without it.

Let $\omega_X(g, h)$ and $\omega_Y(g, h)$ be the respective finite pair representations. It follows from the formulas above, and $W(a_0 I_n)$ commuting with $C(g, h)$, that

$$M\omega_X(g, h) F_X = \ell\, b_Y(a_0^{-1})\, \rho_\sigma(C(1, h))\, \omega_Y(g a_0^{-1}, h) F_Y. \tag{36.8}$$

Indeed the determinant factor on the left after applying M is $b_Y(g)^{-1}$, and

$$W(C(g,h))W(a_0 I_n) = W(C(ga_0^{-1},h));$$

changing that pair representation's determinant factor multiplies it by exactly $b_Y(a_0^{-1})$. This verifies the constant including its dependence on $u = a_0^{-1}$.

The rank-one Schur jet is homogeneous of degree D: at each place a height-one column corresponds to a variable, and there are d_τ such columns. In the integral minor basis all coefficients of the Schur matrix are rational, by Chapter 8. Consequently renaming place variables and applying σ commutes with extraction of this Schur coefficient, while replacing T_1 by T_c multiplies it by c^D. More explicitly for a monomial X^α, the coefficient extractor is $\alpha![X^\alpha]$, as in Chapter 17; a rational linear combination of such homogeneous extractors commutes with σ and with the relabelling, with the factorial unchanged. The full jet after this operation is exactly `schurJet`, including its coordinate basis, not an arbitrary polynomial representation isomorphic to it. Applying it to (36.8) therefore proves the conjugate kernel identity.

Finally take the class sum over the source line. Right multiplication by $u = a_0^{-1}$ permutes its classes, because the finite adelic unitary group of a line is abelian, preserves the chosen level, and preserves each stabilizer. Thus changing the source coordinate g to gu leaves this class sum unchanged. It is a finite sum with rational (reciprocal stabilizer) coefficients, so σ can be applied termwise. We obtain the exact conclusion

Theorem 36.4 (Galois action on the normalized rank-one lift)**.** *With the data and the normalizations above there exist t, ℓ, u, independent of the degrees and the varying character, with $|t|^2 = \Omega^{-1}$, $\ell \neq 0$, and u a norm-one finite idele, such that*

$$\sigma\big(\operatorname{per}^D \Xi_X(h)_i\big) = t^D \ell\, b_Y(u)\, \rho_\sigma(C(1,h))\, \Xi_Y(h)_{e(i)}. \tag{36.9}$$

The index $e(i)$ means the same relabelling of the integral Schur basis by the transformed place permutation (with the opposite notation e^{-1} if source rather than target coordinates are numbered).

This supplies `LValueData.map_trivialLift` in `New/Exchange/AlgebraicPartConjugates.lean` using the actual Weil multiplier and class sum. It does not assert that complex absolute value commutes with σ.

36.6 Conjugate norms and the exact flat error

For the norm-square calculation let c denote ordinary complex conjugation and put $\sigma^{\#} = c\sigma c$. The two automorphisms $\sigma, \sigma^{\#}$ induce the same permutation of the complex places: if $\tau : E \hookrightarrow \mathbb{C}$ is an embedding then $c\tau = \tau^{-}$, since E is CM, and the same identity holds after composing τ with σ. Also μ_Y is a unitary automorphic character so its finite values have complex absolute value one. Consequently $\sigma^{\#}(\mu_X(g)) = \mu_Y(g)$ at the finite points also: it equals $\overline{\sigma(\mu_X(g)^{-1})} = \overline{\mu_Y(g)^{-1}} = \mu_Y(g)$. Thus the right-hand sides of Theorem 36.4 for σ and $\sigma^{\#}$ use the *same* lift Y.

For the two multiplier factors we have an even stronger equality than their moduli:

$$\rho_\sigma(A)\overline{\rho_{\sigma^{\#}}(A)} = \frac{\sigma(\Delta_f(A)\overline{\Delta_f(A)})}{\Delta_f(A)\overline{\Delta_f(A)}} = 1. \tag{36.10}$$

The integer norm-square identity is exactly what is needed at this point.

If vectors $v(h)$ and $w(h)$ satisfy the two semilinear equations

$$\sigma(v(h)_i) = a\, b(h) w(h)_{e(i)}, \qquad \sigma^{\#}(v(h)_i) = a'\, b'(h) w(h)_{e(i)}, \qquad b(h)\overline{b'(h)} = 1,$$

then

$$\sigma\langle v, v\rangle = a\overline{a'}\langle w, w\rangle. \tag{36.11}$$

To verify this expand the finite class sum in its rational stabilizer factors and expand each inner product with its rational Weyl Gram matrix. The Gram matrix is unchanged under the place relabelling: at each place it is the same integral Gram matrix for the relabelled diagram, so the total product Gram matrix has just its rows and columns permuted. The conjugate coefficient in each term is transported by $\sigma(\overline{v_i}) = \overline{\sigma^{\#}(v_i)}$. The displayed cancellation is pointwise, so all h-dependent factors disappear before summing. This is `LevelPlaces.map_inner_self_eq` in `Lib/Adelic/CuspForms/LevelForms.lean`. It is a purely algebraic identity of finite sums.

Apply (36.11) to $\mathrm{per}^D \Xi_X$, using (36.10) for the cancellation. Since each $b_Y(u)$ has modulus one, taking absolute values gives

$$\left|\sigma\|\,\mathrm{per}^D \Xi_X\|^2\right| = |\ell_\sigma \ell_{\sigma^{\#}}|\,\Omega^{-D}\|\Xi_Y\|^2. \tag{36.12}$$

Now the actually scaled `Data.Pair.trivialLift` is

$$\Xi = \kappa_0^n (\bar{\pi}_G\,\mathrm{per})^D\, \Xi_X,$$

where κ_0 is the integral vacuum scale from the CM construction. The principal chosen-prime element has $\pi_G \bar{\pi}_G = g_0 g_2 =: G_0$ (the integer `gProd`). In any complex embedding of the imaginary quadratic field, the two conjugate values are ordinary complex conjugates, so

$$|\sigma(\bar{\pi}_G)|^2 = G_0.$$

Thus (36.12) shows that, at fixed rank n, the conjugate norm of the scaled lift is a constant independent of X, times $(G_0/\Omega)^D \|\Xi_Y\|^2$. The fixed nonzero scalar includes the conjugates of κ_0^n and the two scalars ℓ. No dependence on the particular degrees is hidden in it.

For $n = 2z$ with $z \geq 2$, combine this with the rank-one norm of Proposition 34.5, now justified by Theorem 36.2. At the empty twist level, the factors at the chosen places and the two quotient volumes depend only on the rank and field data. The hook factor is $\prod_\tau d_\tau!$: the highest vector of a one-row Weyl diagram is the monomial of degree d_τ in one standard column, whose square norm in the Fock pairing is $d_\tau!$. Denote the coefficient dimension by $d_{\mathrm{W}} \geq 1$. There is therefore a fixed real constant C with

$$\log\left|\sigma\|\Xi\|^2\right| = C + \sum_\tau \left(\log(d_\tau!) - d_\tau \log(\Omega/G_0)\right) + \log d_{\mathrm{W}} + \log\left|L^M(z, \mu_Y^{-1})\right|. \tag{36.13}$$

Here the L-value is unambiguously the Euler product with both selected and conjugate places removed.

For completeness, the bound on the last summand is elementary in this range. For $q \geq 2$, $z \geq 2$ and $|\alpha| \leq 1$,

$$\left|\log\left|(1 - \alpha q^{-z})^{-1}\right|\right| \leq \sum_{k \geq 1} \frac{q^{-zk}}{k} \leq 2q^{-z} \leq 2^{2-z}\frac{2}{q^2}.$$

At most $[E : \mathbb{Q}]$ places lie over each rational prime and $q_w \geq p$ at every such place, so $\sum_w q_w^{-2} \leq [E : \mathbb{Q}] \sum_p p^{-2} < [E : \mathbb{Q}]$. This shows absolute convergence of the logarithms with bound $2[E : \mathbb{Q}]2^{2-z} \leq 4f$, where $f = [E : \mathbb{Q}]/2$. Exponentiating the log Euler series also

proves nonvanishing. In particular neither a zero-free result at $s = 1$ nor GRH is needed. This derives `abs_log_norm_eulerProd_le` in the precise range needed here.

Writing

$$R = \log d_{\mathrm{W}} + \log |L^M(z, \mu_Y^{-1})|$$

we conclude $R \geq -4f$. If every $d_\tau = 0$, the Weyl component consists only of the constant polynomial, so $d_{\mathrm{W}} = 1$ and $R \leq 4f$. These are exactly the two bounds in `Data.Pair.log_norm_map_norm_sq_trivialLift`; the remaining sum in (36.13) is $\sum_\tau \mathrm{jetSize}(d_\tau)$. The rank-one semilinear comparison and the flat error bound are thereby supplied; this is not the quite different carrier-separation statement about the norm of a noncompact comparison form.

36.7 Integral normalized value measures

We give the integral bound missing from the value-measure argument of Chapter 43.7, retaining the exact exponent on the denominator.

Lemma 36.5 (Finite named indicator denominator)**.** *Let $\mathfrak{m}$ be a split modulus of the CM field and suppose the positive rational integer qN belongs to $\mathfrak{m}$. Let Φ be a function on r-by-s residue matrices modulo $\mathfrak{m}$ whose values are integral at every coefficient embedding above primes not dividing q. Fix such a coefficient prime ℓ. Then*

$$N^{2frs} \kappa_0^{rs} \, \mathrm{name}_{\mathfrak{m}}(\Phi) \, \mathrm{vac}$$

belongs to the standard-data module of denominator N at ℓ (the module of Chapter 17 generated by the integral vacuum and shifts u for which Nu is integral above ℓ).

Proof. On the integral finite name variables the function descends to the finite additive quotient $(\mathcal{O}_E/\mathfrak{m})^{rs}$. Expand it in its additive Fourier basis,

$$\Phi(x) = \frac{1}{|A|} \sum_{\eta \in A^\vee} \widehat{\Phi}(\eta)\eta(x), \qquad A = (\mathcal{O}_E/\mathfrak{m})^{rs}.$$

The $\widehat{\Phi}$ are sums of products of Φ-values by finite roots of unity and hence are integral at the specified coefficient prime. The order $|A|$ divides $|\mathcal{O}_E/(qN)|^{rs} = (qN)^{2frs}$ because the map from the larger residue quotient onto A is a surjective map of finite additive groups. Since ℓ is prime to q, the fraction $N^{2frs}/|A|$ is integral at ℓ.

Each additive character η is represented by a finite Heisenberg shift in the opposite split coordinates. Indeed the annihilator of the integral additive lattice under the normalized trace character is the same lattice in the dual coordinates, by the different normalization of Chapter 7. Since η kills the ideal $\mathfrak{m}$ and hence the sublattice $qN\,\mathcal{O}_E$, the representing shift u satisfies qNu integral. At $\ell \nmid q$ this means Nu integral above ℓ. Applying a normalized Heisenberg modulation to the integral vacuum gives the named vector η times its vacuum support, up to a finite unit phase; the phase is a root of unity by the explicit finite Schrödinger action of Chapter 17. Thus after multiplication by N^{2frs} the displayed Fourier expansion is an integral linear combination of the prescribed translates of κ_0^{rs} vac. This is exactly the standard-data module, proving the claim. □

The proof gives the actual linear combination rather than an upper bound on the Fourier denominator at unspecified primes. It identifies the assertion `Parameters.smul_vacuumDatum_mem_standardData` from `New/Exchange/CrudeModularData.lean`.

For the point data in `TriangleLevelExponent.lean` put $n = 2z$, let $G_0 = \mathrm{gProd}$, let Q be the set of twist places, and put $q = \prod_{w \in Q} q_w$. At chosen level exponent c the ideal $\mathfrak{m}$ is the product of the c-th power of the chosen ideal and the square-free auxiliary ideal. It contains qG_0^c: each rational residue cardinality belongs to its maximal ideal, and $\pi_G \bar{\pi}_G = G_0$ with the chosen ideal generated by one of the two factors, by the chosen-space construction. Multiplying these ideal containments proves `Data.norm_mul_gProd_pow_mem_modulus`. For the *point* datum levelIndicator the exponent is $c = 1$: its functions are the indicators of the chosen column at M and only its one selected coordinate at Q, read modulo MQ. The original lift datum is written at exponent two, but the sum of translates of the point datum in `AuxPair.upsilon_eq_sum` pulls back from exponent one, so its use does not double the denominator. This distinction is needed for the stated denomBound.

Lemma 36.5, applied with $r = n, s = 1, N = G_0$, proves

$$G_0^{2fn}\,\mathrm{pointDatum} \in \mathrm{standardData}_{G_0}.$$

Its values are 0 or 1 before the integral vacuum scaling. The jet scale is $\bar{\pi}_G$ per; $\bar{\pi}_G$ is an algebraic integer and therefore has coefficient ℓ-adic norm at most one at every embedding. The clearing factor for the jets of standard data of denominator N proved in Chapter 17 is $(Nd)^D$, where D is the number of boxes below the target rank, and d the discriminant parameter in that comparison. For a one-row diagram $D = \sum_\tau d_\tau$. Substituting yields exactly

$$B = G_0^{2fn}(G_0 d)^D. \tag{36.14}$$

The standard-data gauge bound therefore states that

$$\mathrm{weylGauge}_{\ell,h}\left(B\ \mathrm{pointKernel}(g,h)\right) \le 1$$

for every g, h and every coefficient embedding above $\ell \nmid q$. This is the source `weylGauge_pointKernel_le_one`. It uses the arithmetic CM jets and Weil stability already established earlier; it is not an integrality assertion for an arbitrarily normalized analytic theta function.

We include the stabilizer and pairing factors to justify interpolation rather than only a bound on one coefficient. Put $M_G = |\prod_{w \in M} \mathrm{GL}_{rt}(k_w)|$, the `chosenGroupOrder`. At the source chosen principal level the stabilizers of the finite classes are trivial by the definite free-level argument, Chapter 8. At a target level contained in K_0 the rational stabilizer Γ_h of a finite class acts on the corresponding definite lattice. Reduction modulo the selected places injects it into $\prod_{w \in M} \mathrm{GL}_n(k_w)$: its kernel lies in the free chosen principal level and is thus trivial. Since the ranks here satisfy $n \le rt$, block embedding into $\prod_{w \in M} \mathrm{GL}_{rt}(k_w)$ proves $|\Gamma_h| \mid M_G$ by Lagrange. This is the exact divisor assertion `card_classAut_dvd_chosenGroupOrder`; replacing it by a numerical bound on the orders would not suffice for integrality.

For any two Weyl coefficient vectors integral in the corresponding gauge at h, one for the coefficient embedding composed with ordinary conjugation and the other for the original embedding, their Gram pairing is integral. Indeed choose a rational translate making h integral at the coefficient prime (the weak approximation in Chapter 17); equivariance and invariance of the Gram pairing reduce the assertion to the integral Weyl lattice itself. There the Gram matrix is integral in the minor basis and both vectors have integral coordinates. The pairing is a finite sum of products of integral numbers. This proves precisely the use of `gramForm_mem_embValRing` in the class sum.

At fixed Q, the class sum of the varying lift is a finite sum over source classes of chosen level and residue-unit lifts at Q. To see the latter factor directly, summing translates of levelIndicator by the lifted residual units gives resDetDatum: at each place of M the support is always the chosen column; at a place of Q only the unique residual unit equal to the selected

nonzero entry contributes, and its determinant character contributes the required residue character. If the entry is zero all terms vanish. Thus the point-kernel finite sum is the source `Data.Pair.trivialLift_apply`. By the character-twist identity for the finite Weil action, varying the tame twist from 1 to η in each summand contributes only the scalar $\eta(g_a)^{-1}$ at the source point of that summand. In the norm-square finite class sum the coefficient of $\eta(g_a^{-1}g_b)$ is a Gram pairing of two point kernels (with the fixed reference character); alternatively one may use the inversely ordered points by conjugating the pairing. This is the exact finite evaluation expansion of Chapter 43.7, now with the point-kernel integrality (36.14).

In particular each coefficient of

$$\frac{M_G B^2}{q^{n-1}}\|\Xi_\eta\|^2$$

is integral at every coefficient embedding above $\ell \nmid q$: B on each vector clears the gauge, M_G clears the stabilizer divisors and q^{n-1} is a unit. The value is therefore a single integral measure for all tame twists of the fixed auxiliary modulus in the precise sense of `Data.TameChar.IsIntegralMeasure`. This proves `isIntegralMeasure_intAlgPart` without presupposing integrality of the normalized L-value itself. Proposition 34.5 and Theorem 36.2 show in turn that this expression is a fixed nonzero constant times

$$L^{M\cup Q}(z,(\mu_X\eta)^{-1}).$$

The factor q^{n-1} from the norm formula cancels its displayed denominator exactly. This is the normalization `intAlgPart_eq_mul_L`. Hence the proof of the rank-at-least-four value-family normalization in Chapter 43.7 no longer needs an independent rank-one Rallis or point-kernel input. Degree zero is included; the difficult rank-two non-absolute Euler range requires the distinct rank-one value-family argument there.

Lastly, for the nonzero reality of the ratios at $z \geq 2$ in `ratio_ne_zero_and_real`, the pulled-back character on the CM field satisfies $\mu^{\mathrm{bc}}(\bar{\mathfrak{a}}) = \mu^{\mathrm{bc}}(\mathfrak{a})^{-1} = \overline{\mu^{\mathrm{bc}}(\mathfrak{a})}$. The omitted set is conjugate-stable and the Euler product converges absolutely at $z \geq 2$. Its factors at a split pair multiply to $|1-\alpha q^{-z}|^{-2} > 0$, while a fixed place of the pulled-back unitary character has trivial value on the local norm-one scalar from the real subfield and, in any case, a real unitary value ± 1 giving a positive Euler factor. The whole product is positive and nonzero by the preceding logarithmic bound. This applies just as well to the twist by η or η^{-1}; dividing numerator and denominator proves reality and nonvanishing in this range. The separate critical rank-two case $z = 1$ would require the specified nontrivial Hecke character and its analytic nonvanishing there, not the logarithmic estimate above.

36.8 Weighted bounds for the real mixed slices

We return to the two line models of Chapter 34.2. One source plane and one target plane have been separated with the adapted rational splittings A, X; write their ranks before adjoining the planes as $n = 2d$ and r, and $e = r - d$. We first verify the analytic hypothesis of the Mellin argument using the actual real Fock model.

Let $M_N(v) = \int (1+\|y\|)^N |v(y)|\,dy$ be the weighted integrable norm of Chapter 10.4 (we may use the maximum norm instead of the Euclidean norm; all the statements here are for fixed finite ranks). In the real adapted mixed model the decomposition of the original real coordinate is

$$y = La(y) + Iw(y) + Kb(y),$$

with fixed real-linear maps a, w, b, since the columns of the adapted frame form a basis. If $\mathcal{S}_x$ denotes the slice at x, then (9.11) and the real Gaussian identity give

$$\mathcal{S}_x(\rho(y)e_0) = \zeta(x,y)\rho_1(w(y))(e^{-\|x+a(y)\|_2^2}e_1), \qquad |\zeta(x,y)| = 1.$$

Here the two vacua are the normalized real Gaussians in the original and residual spaces. A fixed nonzero scalar would make no difference to the following estimates; choosing the real Gaussian of the source gives exactly the displayed formula. This is the defining slice law of `MixedModel` in `Lib/Weil/Fock/MixedFockModel.lean`.

Lemma 36.6 (Slice bound in all weighted norms)**.** *For each pair of integers $N, P \geq 0$ there is a constant $C_{N,P}$ such that*

$$(1 + \|x\|)^P M_N(\mathcal{S}_x v) \leq C_{N,P}\, M_{N+P}(v).$$

Proof. The normalized Heisenberg action is a translation times a phase of modulus one, so

$$M_N(\rho_1(w)e_1) \leq (1 + \|w\|)^N M_N(e_1).$$

For the coherent vector $\rho(y)e_0$, use the slice law and the inequality

$$(1 + \|x\|)^P \leq (1 + \|x + a(y)\|)^P (1 + C\|y\|)^P.$$

The factor $(1 + \|x + a(y)\|)^P$ times the displayed Gaussian is bounded independently of x, y, and $\|w(y)\| \leq C'\|y\|$. It follows that the weighted slice of the coherent vector is at most $C_{N,P}(1 + \|y\|)^{N+P}$. The coherent reproducing integral expresses v as an integral of those vectors against $v(y)$; the slice operator is bounded for every fixed x, and the triangle inequality in weighted L^1 gives the result. Equivalently the claim follows first for finite coherent sums and then by cutting off the reproducing integral and dominated convergence in the weighted norms. This proves the same bound for the full integrable Fock space, whenever the displayed right side is finite. □

In particular every rapid datum has rapid slices; this establishes `MixedModel.existsWeightedNorm_sliceLe` with the actual exponent $N + P$.

The compact real unitary group in an adapted pair preserves the real Gaussian up to a phase and acts by bounded isometries of the real coordinate norm. Its Weil operators therefore preserve the weighted norms up to constants uniform on the compact group, by the same coherent translation formula. The finite components of the maximal compact subgroup do not act on this real datum. For either line branch β the real cone integrand is

$$F_\beta(s; k, a) = \left(\prod_\tau (a_\tau/|a_\tau|)^{2\mu_\tau}\right)\left(\prod_\tau |a_\tau|^{2s}\right)\mathcal{S}_{(0,\dots,0,a)}(\omega_\infty(1,k)v'), \qquad a \in (\mathbb{C}^\times)^f.$$

The coordinate a occupies the last mixed slot; k ranges through a fixed maximal compact with finite Haar measure. This is `coneIntegrand` of `Lib/Weil/MixedModel/EquivariantCoefficients.lean`; the rank-dependent determinant characters in the compact real action have modulus one. The factor $\prod |a_\tau|^{2s}$ is the normalized complex idele modulus to the power s.

Let s vary on any compact subinterval $[\epsilon, B]$ of $(0, \infty)$. Lemma 36.6, with arbitrarily large P, gives

$$M_N(F_\beta(s; k, a)) \leq C\, M_{N+P}(v') \prod_\tau \frac{\max(|a_\tau|^{2\epsilon}, |a_\tau|^{2B})}{(1 + |a_\tau|)^{P/f}}.$$

Indeed $(1 + \|a\|)^P \geq \prod_\tau (1 + |a_\tau|)^{P/f}$. The multiplicative Haar measure on each $\mathbb{C}^\times$ is a constant multiple of $dr/r\, d\theta$. Each one-variable factor is integrable near zero because $\epsilon > 0$ and near infinity for $P/f > 2B$. Thus the displayed product is integrable, uniformly for the compact group and the specified interval. The same holds after inserting any fixed power of $\sum_\tau \log|a_\tau|$, by increasing P and, near zero, replacing ϵ by $\epsilon/2$. This proves joint integrability

of every moment and all the bounds on the Bochner integral; the resulting cone integral is a rapid *datum*, not just a complex number. It also proves holomorphic dependence on s in $\Re s > 0$ after application of any of the fixed bounded linear functionals controlled by a weighted norm. We have thus verified `integrable_coneIntegrand` for the actual source integrand, including the endpoint $s = 1$ (in fact for all positive s).

36.9 The normalized line orbit and the two finite slices

We now make explicit the geometric comparison used in the line constant term. This proof uses the source mixed sums of (9.12) and the zero-term covariance of Chapter 9.9.

For the source splitting A of W', the term indexed by a rational vector $x \in V'(E)$ in its mixed sum is the theta value of the residual datum $\mathrm{sourceTerm}_A(x)f$. In the coordinates of V' the parameter x is its rational vector read through the fixed adelic frame. The precise transformation of this term under the radical element with noncentral parameter c and central parameter z is

$$\mathrm{sourceTerm}_A(x)(\omega(n(c,z),1)f) = \psi\Big(\tfrac{1}{2}\operatorname{tr}(zx^*J_{V'}x)\Big)\,\rho(xc^*J_W)\,\mathrm{sourceTerm}_A(x)f. \tag{36.15}$$

Here z in the source notation is Hermitian; the fixed imaginary scalar in its radical coordinates and additive character are the same as in the mixed-Heisenberg law. This equation is `sourceTerm_radicalElt` of `Lib/Weil/DualPair/PairSplitting.lean`: inserting the block shear into the Heisenberg law gives the displacement by xc^*J_W, and the central phase is the quadratic form of the last mixed variable. No determinant-character term appears because the radical has determinant one and its Weil scalar is one, as proved in Theorem 9.5.

In particular integration over the central parameter in the radical quotient gives zero unless $x^*J_{V'}x = 0$. With the source parameterization of `lineRadical`, the multiplier of z in (36.15) is the nondegenerate additive character on the rational quotient of the central line. Its mean is therefore exactly 0 or 1. The remaining noncentral mean is the one in the definition `constTermAt`. The compact additive quotient measure is probability measure; consequently this supplies `constTermAt_eq_integral`, not a scalar multiple of it.

The sum of these term integrals agrees with the radical integral of the entire theta kernel. For clarity, the mixed expansion of Chapter 9.7 has $\sum_x |\theta(\mathrm{sourceTerm}_A(x)F)| \le C_A\|F\|_1$ by the coherent Gaussian bound. On compact subsets of the radical coordinates the Weil operator norms are uniformly bounded. The parameter quotient admits representatives in a compact subset, because it is a product of additive adelic groups modulo their rational lattices (as in Chapter 9.3); the central coordinate is the corresponding Hermitian linear quotient of the second additive group. Thus the sum of the absolute integrals over a compact set of representatives is finite. Since every summand in the mean is periodic in the rational radical parameters, this proves the interchange on the quotient. It is the exact declaration `hasSum_constTermAt`.

In the nonzero isotropic orbit, fix a rational isotropic vector k_X of the target splitting X. For each $\xi \in E^\times$, the displacement in (36.15) for $x = \xi k_X$ lies along the target-isotropic frame. Averaging the corresponding Heisenberg translations is the zero term of the target mixed model: the map from their noncentral parameter quotient to the zero-term parameter quotient is the invertible linear matrix change induced by ξ. This change is rational, hence preserves the probability quotient measure (it is the isomorphism of compact quotients in Chapter 9.7). The result is the theta value of the successive slice $\mathrm{lineTerm}_1(\xi)$, removing first the source and then the target plane. This proves the identity `constTermAt_smul_k`.

We verify the zero orbit for the standard row datum, rather than leaving it an extra vanishing hypothesis. Locally the row datum at an adapted frame of the source is the inverse

tame character of the determinant of the square matrix obtained by stacking the fixed rows on the free rows, with support on integral invertible matrices. Write the integral adapted source matrix as (K, I, L), with inverse blocks $(K'; I'; L')$. The fixed larger rows are sI', so that their product with K is 0. Evaluation of the first mixed slice at $x = 0$ imposes zero in the free-row product with K as well. The entire square stacked matrix then kills the nonzero primitive column K and cannot be invertible. This is exactly the vanishing of the finite source-term formula: the free row in the L' direction is Fourier integrated at zero, but it cannot change the vanishing column because $L'K = 0$. Hence the residual local datum at that term is zero. More explicitly the mixed evaluation is the integral over the free complementary line coordinate with the first coordinate fixed at zero, and the determinant character is zero at every point of that integral; the other factors do not affect this conclusion. Therefore $\mathrm{constTermAt}(0) = 0$ on the actual standard row data. If the target is translated at the time of evaluation, the condition that the distinguished mixed column be zero is preserved under invertible target multiplication, so this vanishing persists under all such translates. Source line Levi and noncentral radical also preserve the zero column. This is the support calculation used in `LargeSideConstantTerm.lean`; it does not follow just from character orthogonality for the nonzero orbit.

The nonzero rational isotropic vectors of V' form one orbit under rational unitary isometries by Witt extension (and the scalar of a vector is also part of the action). Grouping first by isotropic lines and then by their nonzero rational scalars therefore unfolds the sum over isotropic lines to the quotient by the line parabolic. The Levi used by the source is

$$(c, h) \longmapsto X.\mathrm{levi}(c^{-1}, h), \qquad c \in \mathbb{A}_E^{\times},\ h \in U(V)(\mathbb{A}_F);$$

it acts on k_X by c^{-1} and on the residual space by h. The radical has one vector parameter in an r-dimensional E-space and a central parameter in a one-dimensional F-space. Its conjugation by the Levi scales the vector parameter by $\bar{c}^{-1}$ and the central coordinate by $(c\bar{c})^{-1}$; their measure multipliers are respectively $|c|^{-r}$ and $|c|^{-1}$. Thus its modular factor is $|c|^{-r-1}$, giving $|c|^{r+1}$ in the unfolded Levi integral.

We record why the Iwasawa measure change is available for the specific compact subgroup of the source, rather than invoke an unspecified finite level. At each finite factor, normalize an arbitrary isotropic line vector to be primitive by multiplying by a scalar. The integral Hermitian Witt lemma of Chapter 16.6 (valid in the present split/unramified local fields, including dyadic primes) takes it to a primitive adapted line of the reference frame by an integral unitary matrix. This is a local Iwasawa decomposition into the line stabilizer and the integral compact. Archimedeanly write an isotropic vector as a sum of its positive and negative components, whose lengths are equal and nonzero; after rescaling, the compact block unitary group takes it to the reference sum of positive and negative unit vectors. Almost all finite choices may be the identity, giving the adelic decomposition. A matrix preserving the line has the usual triangular block form in the adapted basis and decomposes uniquely into its radical and Levi matrices. The ordinary Iwasawa integration formula is the quotient integration formula for $P\backslash G$ with the inverse modular weight; the compatible Haar measure is a fixed positive scalar times the chosen Haar measure on the group. With counting measure on the rational points this gives the corresponding quotient formula; the rational radical quotient has the probability measure already chosen. In particular the scalar is independent of the datum and the twisting character.

The radical integration in the target line removes the second line direction; its residual mixed datum is exactly the successive slice just obtained. Consequently the constant term, provided the unfolded integrand is absolutely integrable, equals a fixed nonzero positive scalar

times

$$\int_{E^\times\backslash\mathbb{A}_E^\times}\int_{[U(V)]}\int_{K_{V'}}|c|^{r+1}\sum_{\xi\in E^\times}\theta\Big(\mathrm{lineTerm}_1(\xi)\omega(g',X.\mathrm{levi}(c^{-1},h)k)f\Big)\,dk\,dh\,d^\times c. \tag{36.16}$$

This is the orbital unfolding of `exists_unfold_constant_term`; the quotient integrals use the actual stabilizer of the rational line, not that of an ordered basis of its complementary hyperbolic plane. To justify the initial exchange with target integration before unfolding, the standard upper row support has full right rank at each selected split place: the invertible stacked matrix provides an integral right inverse of the free rows, by taking the appropriate columns of its inverse. The qualitative kernel bound (10.2) for $N = 0$ makes the kernel bounded over all target points, uniformly for the source point in a compact set. The source radical quotient has compact representatives, hence the kernel is absolutely integrable on its product with the finite-volume target quotient. We will prove the required absolute integrability of (36.16) below before using it. The quotient unfolding of a countable orbit for an absolutely integrable function is justified by the usual disjoint-translates partition of a fundamental domain, first for nonnegative functions by Tonelli and then by absolute convergence. The same partition can be applied to the sum of absolute orbit terms once the sum of their absolute unfolded integrals is finite.

Here the two mixed conditions along the removed source and target planes are transverse except in one hyperbolic factor. The common isotropic subspace in $\mathrm{Hom}(W', V')$ has rank $n + r + 1$: in the decompositions $W' = k_A \oplus W \oplus l_A$, $V' = k_X \oplus V \oplus l_X$, the common pairs removed are the two off-central Hom blocks adjacent to those lines and a common line-to-line entry. Their remaining factor is the hyperbolic pair of the other two line-to-line coordinates. The two `lineTerm` maps are evaluation along the first and the opposite isotropic directions in this last pair, after applying the common mixed evaluation. This is verified by ordering the tensor-product Hom frame in the blocks of the source and target line directions, the residual directions and their adjoint frames. In the common evaluation Hom from the removed plane on one side to the residual space on the other has an isotropic block of dimensions n or r, respectively; the four line-to-line entries form two hyperbolic pairs, with one shared removed line in the first pair and complementary lines in the second pair. This is because the tensor-product Hermitian form pairs each source line only with its dual source line and each target line only with its dual target line. Thus the two remaining last lines are perfectly paired and their reversal exchanges the two mixed bases without changing the residual model. All common zero evaluations are over probability measures of additive rational quotients, so their order is irrelevant by Fubini. Now apply the normalized mixed theta expansion to the last hyperbolic plane in either polarization. Both sums over $\xi \in E$ equal the theta of the common residual-plus-hyperbolic datum. This proves the equality of sums `tsum_lineTerm`, including any scalar: it is the expansion of the same normalized theta, rather than equality of its two individual coefficients.

We include the finite calculation identifying the slice datum itself. Multiplying the entire stacked defining row matrix on the right by the integral adapted source frame (K, I, L) changes the inverse unit determinant character by a unit-modulus factor and replaces its fixed rows by $(0, s, 0)$. In the target let the adapted frame and its dual be the analogous integral matrices. The mixed frame columns are tensor products of the first chosen line block, the dual residual frame and the complementary line block, as in the preceding decomposition. Under the integral frame changes on the left and the right and the permutations to the block order of the removed directions, the matrix evaluated inside the determinant is

$$\begin{pmatrix} u & Z_{12} & Z_{13} \\ 0 & Y & Z_{23} \\ 0 & 0 & v \end{pmatrix}.$$

Here Y is the remaining square matrix of the lower standard rows and the fixed rows s in their prescribed order. Its block Z_{23} is padded by zero on the fixed rows; all other displayed off-diagonal free entries are the ordinary integral variables of the mixed evaluation. The diagonal scalars are $(u, v) = (a_w, t)$ in the first polarization and (t, a_w) in the second, where t is the Fourier-integrated variable at the selected place and $a_w = \xi_w/\alpha_w$ with the fixed scale $\alpha = \mathrm{scaleIdele}$. This block identity is the source's from-blocks matrix in `Lib/SiegelWeil/Tower/LineGaussSum.lean`: its first and last new columns are the two line coordinates and its middle columns the residual row variables; the prescribed rows have vanishing first and last coordinates because $I'K = I'L = 0$. The dual target change makes the off-diagonal lower entries zero in the removed mixed coordinates. The two last line entries are exchanged in the opposite polarization. The remaining integral changes are invertible and so preserve the integrality conditions in both directions.

The determinant of the displayed matrix is $u \det(Y)\, v$. Thus the row datum factors into the lower row datum at Y, the inverse unit character of a_w, the inverse unit character of t, and indicators of all the free off-diagonal entries being integral. The untwisted free entries integrate with their additive characters at frequency zero, hence with volume one. The last entry integrates against the additive character of $\bar{a}_w t$, with the normalized integral conductor (the trace-dual different is already in α). This is precisely Lemma 34.1. Concretely its support is $a_w \in \mathcal{O}_w^\times$ and $\bar{a}_w$ in a fixed inverse-uniformizer unit coset. At places outside the selected conjugate-closure the datum is vacuum and the mixed slice is the integral-lattice indicator in the new coordinate times the vacuum of the remaining space. The common scalar is the l_β of the normalized mixed vacuum; it depends on the fixed geometric splittings rather than the character. To see this explicitly, apply the same fixed mixed evaluation to the vacuum and evaluate its residual vector at zero. The coherent construction gives its real Gaussian and finite lattice indicator multiplied by a nonzero scalar, fixed once by (9.12). The named-vector evaluation is the additive Fourier integral in the adapted finite coordinates relative to this vacuum, normalized by lattice volume one. Its scalar cannot vary with the subsequently named function. The remaining determinants of integral coordinate changes give character values of units, each of modulus one, and do not change this scalar's modulus. These observations yield (34.1) for each branch with $|G_S|^2 = Q^{-1}$, including the claimed independence from the character and selected local norms of its vacuum factor. They also show that the last slice at $a = 0$ vanishes in either branch when the split tame conductor is nonempty, as is assumed in `SplitTameChar`.

Finally the standard finite row datum is fixed under all integral target points k. In its named Schrödinger coordinate the target acts by $x \mapsto k^{-1}x$. The determinant of the stacked matrix is thus multiplied by $\det k^{-1}$; its inverse unit character gains a factor $\chi(\det k)$. For the pair character on the target $\mathrm{rankTwist}(\nu^{-1})$ the tame factor is $\chi(\det k)^{-1}$ and the extra v factor is trivial on integral determinants. They cancel, and the vacuum away from the conductor is unchanged. This proves `pairWeil_stdRowDatum_eq_self`. The finite compact component of k can be dropped, leaving only the compact real action in F_β.

The exact Levi factor. The source point in (36.16) is $g' = A.\mathrm{levi}(u, g)$, with $u = 1$ or $r = n$, as in the source. Put $c_1 = cu$ and $a = \mathrm{rankTwist}(\nu^{-1})$ at the lower source rank, and write $\chi_j = a.\,\mathrm{spaceChar}(j)$ for the corresponding adelic character on scalars. The defining rank twist gives $(av).\,\mathrm{spaceChar}(j+2) = a.\,\mathrm{spaceChar}(j)$. To compute the scalar in the first branch, remove the source plane then the target plane. The zero-term law (9.13) contributes $\chi_r(u)^{-1}|u|^{(r+2)/2}$ and the target law (9.14) contributes $\chi_n(c)^{-1}|c|^{-n/2}$ because its Levi acts by c^{-1} on the line. In the second branch remove the target first, giving $\chi_n(c)^{-1}|c|^{-(n+2)/2}$, then the source, giving $\chi_r(u)^{-1}|u|^{r/2}$. When $r = n$, $\chi_r = \chi_n$; when $u = 1$ the factors of u are

trivial. Thus the two products are respectively

$$\frac{|u|^{r+1}}{\chi_n(c_1)|c_1|^{d}}, \qquad \frac{|u|^{r+1}}{\chi_n(c_1)|c_1|^{d+1}}.$$

The transformation of the line translation parameter is $\xi \mapsto \xi c_1$ in the first branch and $\xi \mapsto \xi \bar{c}_1^{-1}$ in the second, by the line action and its dual in the remaining mixed pair. This proves precisely `lineTerm_pairWeil_levi`, the scalar and the two substitutions in Chapter 34.2. No additional $|\det h|$ occurs because the lower unitary determinant has idele norm one. Also $\chi_n(c_1) = \nu_E(c_1)^{-1}$ as a consequence of even n and the prescribed rank twist, where $\nu_E(a) = \nu(a/\bar{a})$.

36.10 Absolute Mellin endpoints of the line constant term

In (36.16) insert $|c|^t$ with a complex parameter t, leaving every other factor unchanged. Call the resulting Mellin integral $I(t)$ whenever absolutely convergent. Initially consider real t, and let

$$s_+(t) = e + 1 + t, \qquad s_-(t) = -e - t.$$

We claim that it converges on both sufficiently far half-strips, using the first and second line model respectively. Here $s_\pm(t)$ are the zeta exponents of Chapter 34.2, not an analytic continuation assumption on the original integral.

For the first branch, after the Levi calculation the factor $|c|^{r+1}$ becomes $\nu_E(cu)|cu|^{e+1}$; the insertion of $|c|^t$ gives the additional fixed factor $|u|^{-t}$ and exponent $s_+(t)$ in cu. Estimate by summing absolute values over the nonzero rational parameters ξ. Upon integrating over the idele-class quotient, the substitution $x = \xi cu$ converts that sum and the quotient integral into one integral over $\mathbb{A}_E^\times$, by the counting-measure quotient formula. The product formula $|\xi| = 1$ ensures the positive modulus factor is unchanged; the character is trivial on principal ideles, so there is likewise no additional phase. The lower residual theta kernel has only the lower row datum at the chosen finite places. By its full right rank as above, (10.2) at $N = 0$ bounds its absolute value, for fixed lower source g, by $C_g M_N$ of its real datum, uniformly for every residual target point. Its integral over the residual target quotient is bounded by the same expression times the finite quotient volume. Apply Lemma 36.6 and the real cone majorant of Section 36.8. This proves that the real and compact factors of the positive majorant have finite integral whenever $s_+(t) > 0$. The finite factor, after the fixed scalar and inverse-uniformizer translation of Chapter 34.2, has absolute value supported on integral valuations at all other places and units at the selected ones. Its absolute multiplicative integral is a fixed finite product times

$$\prod_{w \notin S \cup \bar{S}} (1 - q_w^{-s_+(t)})^{-1}.$$

It converges for $s_+(t) > 1$; compare it with a finite power of the ordinary rational-prime Euler product since at most $[E : \mathbb{Q}]$ factors lie over each rational prime and $q_w \geq p$. Thus the sum of absolute orbit terms in this first branch is integrable in the desired right half-strip.

In the opposite branch the same argument uses the equality of the two rational sums proved in Section 36.9; its scalar after insertion is obtained with the substitution $x = \xi\, \overline{cu}^{-1}$. The change of variables of ideles by conjugate-inverse preserves multiplicative Haar measure (a Haar-preserving involution), sends their modulus to the inverse, and sends $\nu_E(x)$ to itself, by the norm-one quotient formula. The exponent is exactly $s_-(t)$ because the Levi denominator in this branch has the extra power $|cu|$. The same positive majorant of the lower residual theta and the real cone estimates apply with $s_-(t)$, and the finite absolute Euler product

converges for $s_-(t) > 1$. This proves absolute integrability in the desired left half-strip for the opposite sum, hence for the common *summed* integrand by the triangle inequality.

For example take $T = |e|+3$. The two endpoints $t = T$ and $t = -T$ have the respective zeta exponents > 1. Let $H(c, h, k)$ denote the common summed integrand of (36.16), including the modulus factor but not the additional $|c|^t$. The preceding estimates prove

$$\int |H(c,h,k)|\,|c|^{T} < \infty, \qquad \int |H(c,h,k)|\,|c|^{-T} < \infty.$$

For every $-T \le x \le T$, one has $|c|^x \le |c|^T + |c|^{-T}$, so the common integrand is absolutely integrable at every real exponent in between, especially 0. Holomorphy for $|\Re t| < T$ follows from dominated convergence on compact substrips: if $\Re t$ is separated from the endpoints, every derivative of $|c|^t$ introduces a power of $\log|c|$, bounded by a constant times a sufficiently small extra positive or negative exponent. The enlarged endpoint majorants above still dominate. We have thus obtained $I(t)$ as an ordinary absolutely convergent Mellin integral holomorphic on this strip. This is the common-endpoint argument, not an appeal to separate analytic continuations of unrelated formal sums.

For $s_\pm(t) > 1$ the prior unfolding and changes of variables are now justified with both absolute values and phases retained. They factor $I(t)$ as the fixed measure scalar times the finite Tate zeta factor, its translating idele character and modulus power, and the lower theta integral of the cone datum $F_\beta(s_\pm(t); k, a)$. The finite calculation in Chapter 34.2 gives its fixed conductor factor times $L^S(s_\pm(t), \nu_E)$. The translating idele contributes only the fixed norm power and a unit-modulus character; the real cone factor is holomorphic for $\Re s_\pm(t) > 0$ by Section 36.8, even after applying the bounded theta integral controlled by the weighted moments. The unitary Hecke character of the source is nontrivial, since it has the stipulated positive infinity weights in `SplitTameChar`. Tate's zeta integral theorem gives holomorphy of its Hecke L-function on $\Re s > 0$ (there is no pole at 1 for a nontrivial unitary non-norm infinity type). The standard theorem and the nonvanishing at the real value 1 used here are also established for these characters in Chapter 35.4. For the continuation argument only holomorphy is required. Removing the finite Euler factors multiplies this holomorphic function by finite polynomials in q^{-s}.

Choose the branch with $s(0) = e+1$ if $e \ge 0$, and $s(0) = -e$ if $e < 0$. Its exponent is at least 1. The two holomorphic expressions agree in their nonempty common open region of absolute convergence and, by the identity theorem, continue to agree at $t = 0$ (inside $\Re s > 0$). No conditionally convergent Euler factors at $s = 1$ were multiplied. The substitutions involving u contribute $|u|^{-t}$ and disappear at this original exponent. The conductor factor calculation in Lemma 34.2 therefore yields a nonzero scalar γ_ν satisfying

$$|\gamma_\nu|^2\, P_\nu(e) = C_{e,\alpha}\, P_\nu(e+1),$$

with the positive constant independent of the varying conductor and character, and with the lower real datum given by the actual cone integral at $s(0)$. This proves the common Mellin endpoint and line orbit needed in `exists_constant_term_thetaLarge`. The initial exchange and orbit partition at $t = 0$ use the sum of absolute values of the line contributions grouped by line (each contribution is itself the summed integral over ξ), not absolute summation in an arbitrary polarization at $s = 1$. The required grouped-line majorant is exactly the common summed-integrand estimate just proved. Thus the grouped orbit unfolding is legitimate at the endpoint. No estimate on the entire sequence of descents in the noncompact tower follows simply from this line calculation.

36.11 Exact-source status of this chapter

1. **PROVED:** `SharpData.lean`, `exists_graph_eq_of_zeroTerm_ne_zero` and `graphSplitting_lagrangian_injective` (Theorem 36.1); together with Section 11.6, this supplies the clean graph evaluation of C20.

2. **PROVED:** `inner_product_formula` in the same file, including absolute convergence on the source group and square integrability of the theta lifts (Theorem 36.2); `exists_dual_datum`, including the unitary shear and exact coefficient cutoff (Lemma 36.3); `IsSharp.partialIntegral_eq`, `hasSum_partialIntegral_coset`, `tendsto_partialIntegral`, and `rallis_inner_product_euler` with the real-edge hypothesis as stated in that theorem (Section 36.4). These close the general Rallis and sharp Euler steps of C22 and the resulting specialized algebraic rank-one norm in C16, by Proposition 34.5; the existence of the actual higher-rank noncompact edges remains a different representation-theoretic assertion.

3. **PROVED:** `LValueData.map_trivialLift` and `Data.Pair.log_norm_map_norm_sq_trivialLift` (Theorem 36.4, Section 36.6), using the previous semilinear CM comparison. The rank-one conjugate-norm component of C17 is derived, including both numerical error bounds; the carrier separation part of C17 is not a consequence of this calculation.

4. **PROVED:** `Parameters.smul_vacuumDatum_mem_standardData`, `Data.norm_mul_gProd_pow_mem_modulus`, `weylGauge_pointKernel_le_one`, and the rank-at-least-four normalized evaluation sum `isIntegralMeasure_intAlgPart` and `intAlgPart_eq_mul_L` (Section 36.7). Thus the earlier C16 and S05 rank-one normalized value-measure premises in that range are supplied. The rank-two family also requires the independently treated determinant-period and carrier comparison; S05 as a whole should not yet be recorded closed.

5. **PROVED:** `MixedModel.exists_weightedNorm_slice_le`, `integrable_coneIntegrand` in the actual weighted Fock norm, and the common Mellin-endpoint convergence and factorization given the adapted normalization of the two line slices (Sections 36.8–36.10). The preceding reconstruction gives the adapted triangular matrix and finite Fourier calculation, line central mean, Poisson exchange, orbits and Levi factors explicitly. The fixed mixed-model vacuum scalar rests on the exact normalized global oscillator construction of Chapters 9, 11; the remaining comparison is an audit of the exact adapted frame identifications in `exists_lineTerm_localDatum`, not an unresolved endpoint estimate. We retain that qualification for the full C23 interface. These results do not themselves prove C25, namely the global tower vanishing and boundedness after subtraction of all relevant constant terms, or the normalized period conversion for the actual noncompact tested quotient.

Chapter 37

The singular-coefficient induction and the standard-column reduction

This chapter closes the abstract reduction C04/C24, separating it from the mixed-signature geometric vanishing proved in the exchange chapter. The sources are `Lib/SiegelWeil/SingularForms/SingularForms.lean` (particularly `freitag`), `New/Exchange/StandardCoefficients.lean` (`IsEquivariantLift.apply_stdColDatum_eq_zero`), together with the supporting radical, opposite-frame, Iwasawa and centre-coefficient lemmas of that directory. We use the integral adelic frames and the scalar conventions of the paragraph following Proposition 35.9. In particular, the induction concerns a prescribed matrix coefficient and not arbitrary completeness of all degenerate Fourier expansions.

37.1 Centres, phases and bordered coefficients

Write D for an isotropic embedding of rank b, K for its adelic frame and $K^\flat$ for the dual row; $K^\flat K = 0$. Adjoin an isotropic line A, with frame u and row $u^\flat$, contained in the orthogonal complement of the hyperbolic pair defining D. The composite embedding has frame $F = [u\ K]$, $F^\flat = \binom{u^\flat}{K^\flat}$. We denote the opposite line by A^-, obtained by exchanging the two vectors of this hyperbolic pair; the complementary embedding B then gives the same centre construction in either choice. Thus $F^\flat F = 0$. A centre element in these coordinates is

$$n_F(M) = 1 + FMF^\flat$$

with M satisfying the centre's adjoint condition. The adjoint $*$ is conjugate transpose. Frame conjugation of this element by g with $gF = FC$ sends M to CMC^*: indeed $F^\flat g^{-1} = C^* F^\flat$. The matrix index transforms in the opposite direction, by $R \mapsto C^* R C$.

Use δ for the local scale idele of the source, $\jmath$ for its rational skew scalar, and ψ_0 for the source's standard additive character on $\mathbb{A}_E/E$. The defining phase is precisely

$$\psi_0(\delta^{-1} \operatorname{tr}(RM)).$$

This includes the prescribed scale, not an extra factor of 2. At a finite place, the self-duality statement `stdLattice.forall_psi0_mul_eq_one_iff` says that triviality against all integral test matrices is equivalent to integrality of the entries of $\delta^{-1}R$; at a conjugate pair the test is made over one member with the adjoint condition determining the other. This is the local inverse-different calculation: the character of the trace pairing has the inverse different as its annihilator, and the chosen scale identifies it with the prescribed standard lattice. Rational matrices are dense in the product of any finite family of completions. We also use the additive

strong approximation statement that rational matrices plus matrices supported in a nonempty set U of finite places are dense in the full additive adelic matrix space. This follows by the Chinese remainder theorem after clearing denominators, or equivalently from the perfect adelic trace pairing: the annihilator of the diagonal rational space is rational, and a rational annihilator of the entire local space at one place is zero. These statements apply to rectangular matrices as well as to the adjoint subspace specifying a centre.

Let f be a continuous left rational invariant function, right invariant by the finite integral level subgroup, and write

$$f_{D,T}(x) = \int_{[N_D]} f(nx)\overline{\psi_T(n)}\, dn$$

with probability Haar measure on the compact additive quotient. Similarly define $f_{F,R}$. Characters of these compact abelian groups give Fourier uniqueness for continuous functions. Fubini on the compact quotients shows that the Fourier coefficient of $n \mapsto f_{D,T}(nx)$ on the larger centre, at an index R, is $f_{F,R}(x)$ if the bottom-right block of R is T, and is zero otherwise. At a bordered index we may use

$$R = \begin{pmatrix} \sigma_0 & \sigma_1 \\ 0 & T \end{pmatrix}.$$

Here indices may be represented modulo their annihilator of the centre's adjoint subspace; the displayed triangular representative is the source's `stdRowDatum`/bordered convention. A rational shear

$$s_A(d)F = F \begin{pmatrix} 1 & 0 \\ -d^* & I \end{pmatrix} \tag{37.1}$$

fixes the D-frame and sends an index with invertible bottom-right block T to bordered form by eliminating the lower-left block. Explicitly, for a general R, take $d^* = -T^{-1}R_{21}$ when using the inverse shear. Thus, if all bordered coefficients at all rational shears $s_A(d)x$ vanish, Fourier uniqueness implies $f_{D,T}(x) = 0$. This is the argument of `centreCoeff_eq_zero_of_shearElt`; it uses only Fourier uniqueness on the larger centre, not an assertion that degenerate Fourier terms form a complete system for f itself.

37.2 The radical hypothesis

Let S be the set of finite places used in the induction, and let U be a nonempty set disjoint from the conjugation-stable set $S^{\pm} = S \cup \bar{S}$. We use e_V for the idempotent selecting a set V of places and $[g]_V$ for the element equal to g on V, identity elsewhere. The source's radical hypothesis is invariance at x under the unitary elements

$$\nu = 1 + \overline{e_U} uc - e_U a, \qquad cu = 0,$$

where x fixes the D-frame outside $S^{\pm}$ and a is chosen so that ν is unitary. It is imposed for each admissible line extension and its opposite. Taking ν in the composite centre with first-column parameter supported at U, the entries of that column are arbitrary, subject only to the adjoint relation at the conjugate place. Invariance of the bordered coefficient under these translations implies that σ_0, σ_1 vanish, by the local nondegeneracy of the character and the fact that a nonzero rational entry has nonzero image in every completion. Thus, under the radical hypothesis, only $R_0 = \mathrm{diag}(0, T)$ remains.

For later use the resulting coefficient $b_0 = f_{F,R_0}$ is invariant under every line-radical element whose coefficients are supported away from $S^{\pm}$, and actually under the whole line radical at x. Here is the calculation that explains the latter assertion. The line-parabolic radical

has parameters (c, z) with c in the orthogonal complement of the hyperbolic line pair. Its standard matrix is obtained by adding $-Icu^\flat + uc^*H_0I'$ and the central correction

$$-\tfrac{1}{2}u(\jmath^{-1}(z+z^*)+c^*H_0c)u^\flat.$$

For $c = e_U y, z = 0$, its expression is of the precise allowed form $1 + \bar{e}_U u\widetilde{c} - e_U\widetilde{a}$ (the term quadratic in y goes into $\widetilde{c}$). The central parameter is invisible at index R_0. The rational line-radical elements preserve this index, since on F they act by the upper triangular unipotent block with bottom-right identity. Left rational invariance therefore supplies every rational translation of c. Additive strong approximation for the rational coordinates of c and their U-supported values, followed by continuity, proves invariance under every c. These are the two mechanisms of the source lemmas which move a nonzero bordered coefficient through Iwasawa and replace its upper border by zero.

37.3 The exact determinant-index induction

For matrices at the finite places $S^\pm$, let

$$Q = \delta^{-1}\gamma^*T\gamma, \qquad \mathcal{D}(Q) = \prod_{v\in S^\pm} q_v^{\mathrm{ord}_v(\det Q)}$$

when Q is integral. Integrality here and throughout is entrywise. If Q is not integral, set the induction index to zero. For an integral nonsingular Q, this is the positive finite index of its image lattice, by the elementary divisor theorem. Write $x \sim_S g$ for the relation that x agrees with g at $S^\pm$ and fixes the D-frame elsewhere (including infinity). We will apply this relation with the matrices g in the line- or frame-parabolic preserving D. It is preserved under multiplication of such matching pairs.

The primitive hypothesis is the following: for every $x \sim_S g$, where $gK = K\gamma$ at $S^\pm$, if Q is integral unimodular at all these places then $f_{D,T}(x) = 0$. We assume the prescribed radical hypothesis above at matching elements, for every admissible line extension and opposite, in addition to left rational invariance and right integral level invariance. No growth estimate enters.

Theorem 37.1 (Singular-coefficient descent; Freitag's argument)**.** *Under these hypotheses, if T is invertible, then $f_{D,T}(x) = 0$ for every $x \sim_S g$ with g preserving the D-frame. In particular a right-hand argument supported on $S^\pm$ gives zero. This is the source reduction* `freitag`*.*

Proof. First, a nonzero bordered coefficient supplies a special matching element. Apply the flag-compatible Iwasawa decomposition at $S^\pm$ to the line and composite flags: multiplication on the right by a finite integral level element κ makes the composite frame block upper triangular, of the form

$$\begin{pmatrix} t & * \\ 0 & \gamma' \end{pmatrix}.$$

Such decompositions follow by taking a basis of the saturated intersection of each isotropic subspace with the self-dual lattice, compatibly in the nested flag, extending this to a hyperbolic integral basis and taking its Gram-dual partners. At a split place this is elementary row reduction of a flag of free modules over a DVR; in the unramified Hermitian case the partners are corrected using a trace-one element to make their pairings vanish. These are precisely the finite places under consideration in this decomposition. The upper-block line-parabolic matrix factors as $n_A(c, z)m$, where m acts on the composite frame by $\mathrm{diag}(t, \gamma')$ and on the

opposite composite frame by $\operatorname{diag}(t^{-*},\gamma')$. Right level invariance and the radical result permit replacement of the bordered coefficient by $b_0(x_0)\neq 0$ with $x_0\sim_S m$, upon left multiplication by $[n_A(c,z)]_{S^\pm}^{-1}$. In this replacement the D-index stays T.

There must be a matching $Y\sim_S m$ with $f_{D,T}(Y)\neq 0$. Indeed, if every such value were zero, the same would hold after translating x_0 by the U-supported additive centre and by rational centre elements, since the latter change a D-coefficient only by its character. Additive strong approximation and continuity would give zero throughout the larger centre, hence $b_0(x_0)=0$ by the nested integral. At Y we may translate by every integral D-centre element on the right; conjugating by Y and using the transformation law gives triviality of the phase with index $Q'=\delta^{-1}\gamma'^{*}T\gamma'$. Local self-duality forces Q' integral. In particular any such normalized nonzero coefficient with nonintegral Q' is impossible.

Now prove the stronger assertion that every bordered coefficient at a matching element whose composite frame block is lower triangular,

$$C=\begin{pmatrix} a & 0\\ z & \gamma\end{pmatrix},$$

vanishes unless $\gamma^{-1}z$ is integral at $S^\pm$, using strong induction on the index. Compare C to the Iwasawa upper block just constructed. The change-of-frame matrix between them is integral, since it is induced on the saturated composite lattice by the integral flag adapter. Taking its bottom blocks i,r, the two identities are

$$z=\gamma' i,\qquad \gamma=\gamma' r.$$

Thus $\gamma^{-1}z=r^{-1}i$ and $Q=r^{*}Q'r$. If $r^{-1}i$ is not integral, then r cannot be unimodular somewhere. If Q' is nonintegral, the normalized nonzero coefficient has already been ruled out. Otherwise $Q=r^{*}Q'r$ implies Q integral, and

$$\mathcal{D}(Q)=\mathcal{D}(Q')\prod_{v\in S^\pm} q_v^{\operatorname{ord}_v(\det r^{*})+\operatorname{ord}_v(\det r)}>\mathcal{D}(Q').$$

Consequently induction excludes this nonzero coefficient at the smaller normalized index. We explain explicitly how to exclude a normalized nonzero coefficient at the current index; this also justifies the induction assertion used at every earlier index.

Fix $b_0(x_0)\neq 0$ at the normalized block $m=\operatorname{diag}(t,\gamma)$, and take any d at $S^\pm$ whose row dQ is integral. Consider the difference

$$f'(x)=f(x[s_A(d)]_{S^\pm})-f(x).$$

For any $X\sim_S m$, use opposite rational shears $s_{A^-}(w)$ on X; the opposite composite frame block of the resulting matching element is

$$C_-=\begin{pmatrix} t^{-*} & 0\\ z & \gamma\end{pmatrix},\qquad z=-w^{*}t^{-*}.$$

The right translation by $s_A(d)$ is a centre parameter at this opposite embedding with top-right block d. At a bordered coefficient of the original function at this element, if it is nonzero, the lower-triangular support assertion just proved implies integrality of $\gamma^{-1}z$. Direct multiplication of $C_-MC_-^{*}$ with the bordered index gives the relative phase

$$\psi_0\bigl(e_{S^\pm}\delta^{-1}\operatorname{tr}(RC_-MC_-^{*})\bigr)=\psi_0\bigl(e_{S^\pm}(dQ)\gamma^{-1}z\bigr)=1.$$

All terms involving the upper border disappear, because the chosen opposite centre parameter has zero first column in the bordered representative. If the original bordered coefficient is zero,

its translate is likewise zero by the same character law. Thus every such bordered coefficient of the difference is zero. The rational shear criterion implies $f'_{D,T}(X) = 0$ for every $X \sim_S m$. The nested integral and additive density argument of the first paragraph give $b'_0(x_0) = 0$.

Hence $b_0(x_0[s_A(d)]_{S^\pm}) = b_0(x_0) \neq 0$. This time the ordinary composite block is lower triangular with $z = -\gamma d^*$. The same lower-triangular support assertion shows that d^*, hence d, is integral at all $S^\pm$. We may choose d successively to be each row of Q^{-1}, for which dQ is an integral standard row. It follows that Q^{-1} is integral. Together with integrality of Q this proves that Q is unimodular. But a matching $Y \sim_S m$ with nonzero D-coefficient was obtained in the first paragraph, contradicting the primitive hypothesis. This completes both steps of the simultaneous induction, with nonintegral index zero handled by the local character test and the same comparison. Finally all normalized bordered coefficients vanish, and the rational shear criterion applies at the original argument. □

The exact role of the index is worth stressing. Merely establishing integrality of Q does not reach the primitive case. The comparison $Q = r^*Q'r$ supplies the strict descent when the frame shear has a denominator; the opposite-centre calculation then eliminates every denominator of Q^{-1}. These are independent steps and both are required. Haar measures are probability measures throughout; changes of centre coordinates thus introduce no unspecified constants.

37.4 Application to an equivariant lift

We now prove the passage from the primitive coefficient to a standard-column function. This is not a second geometric vanishing statement. The source abstracts an equivariant lift $\mathscr{D}$ from the finite level Schwartz space into continuous automorphic functions; its law on an admissible integral matrix is precisely the relation used below. We work with the finite set P of split tame places indexing the standard columns, disjoint from its conjugate. Set $S = (P \cup \bar{P})^c$ in the finite places and $U = \bar{P}$. The integral frame D has standard row $s = K^\flat|_P$, and the Schwartz function Ψ_0 attached to this row is the prescribed standard-column datum of rank b.

For this passage, write h_0 for the two primitive geometric assumptions in `StandardCoefficients.lean`. In the geometric normalization, index $T = \jmath J_{V'}$ is associated with a rational Hermitian space V' of rank b; $J_{V'}$ is the rational Gram matrix in its chosen coordinates, and its finite adelic frame is denoted $x_{V'}$. The first assumption says the coefficient of $\mathscr{D}\Psi_0$ at this index is zero when the finite isometric coordinate computed from the argument is integral (right multiplication by a finite integral basis change is allowed), with all the required real signs. The second says it is zero at a rational Hermitian index whose real signs are not those in the allowed family V'. We use these assumptions only in the primitive case, and not for an arbitrary index or arbitrary test function. They are precisely the geometric inputs to be supplied by the mixed term, not asserted as independent theorems here.

37.4.1 Shear averaging

A standard-column function at P has support on those Y for which sY is invertible with integral inverse (in the integral normalizations, $sY \in \mathrm{GL}_b(\mathcal{O}_P)$); it is invariant under each integral shear of the prescribed form. The following elementary observation is the source's shear-averaging mechanism. Suppose a linear functional Λ on the Schwartz space is unchanged by

$$Y \longmapsto Y + \widetilde{u}vsY$$

for all rows v, where $s\widetilde{u} = 0$, and suppose $r\widetilde{u} = 0$. Then

$$\Lambda\big(\Psi_0(Y + \widetilde{u}rY)\big) = \Lambda\big(\Psi_0(Y)\big). \tag{37.2}$$

One proves this by averaging over a compact open additive lattice of the row variable v, chosen large enough that $(rY)(sY)^{-1}$ lies in that lattice whenever either integrand is nonzero. The support condition guarantees that $(sY)^{-1}$ exists, and compactness of the support guarantees the uniform choice of lattice. More precisely, the expression $(rY)(sY)^{-1}$ depends only on the quotient by the one-dimensional translation direction $\widetilde{u}$, because $s\widetilde{u} = r\widetilde{u} = 0$. Thus averaging the first function produces the same function pointwise as averaging the second: the difference between the two row variables is exactly $(rY)(sY)^{-1}$, and Haar translation invariance applies. The average is a finite linear combination in the locally constant compactly supported Schwartz space, so applying Λ is legitimate without a continuity assumption on Λ. This proves the assertion.

We verify the `RadicalInvariant` hypothesis for $f = \mathscr{D}\Psi_0$. Consider any of the line extensions of D needed in the descent (including the opposite line), let R be its bordered index and x fix $D.K$ away from $S^{\pm}$. The radical element has the form $\nu = 1 + \bar{e}_U uc - e_U a$ above. It is supported on $P \cup \bar{P}$, and on P its matrix is $1 + uc$ with $cu = 0$. Thus it has determinant one there, so the equivariance law applies without a determinant character. Put $x' = \nu x$, $\widetilde{u} = x'^{-1}u|_P$, and $r = cx'|_P$. Since x fixes K at $\bar{P}$, its adjoint fixes $K^\flat$ at P; hence $K^\flat x = K^\flat$ there. Likewise $K^\flat \nu = K^\flat$, by $K^\flat u = 0$, so

$$s\widetilde{u} = 0, \qquad r\widetilde{u} = 0.$$

The law of the equivariant lift identifies the original coefficient at x with the coefficient functional at x' on $\Psi_0(Y + \widetilde{u}rY)$. On the other hand this bordered functional is invariant under all shears $Y \mapsto Y + \widetilde{u}vsY$: these are its composite centre parameters with top-right row v on P, whose character is trivial since $R_{21} = 0$. Apply (37.2) and the same law to identify it with the coefficient at x' of $\mathscr{D}\Psi_0$. This supplies exactly the required radical invariance. The concatenated frame here is ordered with the new line first; starting with the source's $D + D_1$ rather than $D_1 + D$ is just the corresponding permutation of both indices and rows.

37.4.2 From the geometric primitive case to the Freitag base

Let T be rational with $T_P = \delta^{-1}T|_P$ unimodular. In particular T is invertible. Use the prescribed adjoint symmetry of the rational indices to write $B = \jmath^{-1}T$, a Hermitian matrix. At a matching element $x \sim_S g$ for the induction, with $gK = K\gamma$ at $S^{\pm}$, the action on K at the finite places is given by $q = e_{S^{\pm}}\gamma + (1 - e_{S^{\pm}})I$. Thus the base condition says

$$Q_f = \delta^{-1}q^{*}Tq \quad \text{is integral and unimodular at all finite places.}$$

Here the places in $P \cup \bar{P}$ satisfy the condition by T_P and the adjoint symmetry. The source's unimodular orthonormalization statement for the given finite scale lattice gives $k \in \mathrm{GL}_b(\widehat{\mathcal{O}}_E)$ with $k^{*}Q_f k = I$. This is the local classification of unimodular Hermitian lattices in the specified unramified and split cases: over a split pair it reduces to choosing a basis on one side and its dual on the other; over an unramified quadratic local extension the norm map on units is surjective and a unimodular Hermitian matrix admits a unit vector (use nondegeneracy of the residual trace pairing if the diagonal entries alone are not units), which splits off orthogonally, giving induction on the rank. These local normalizations agree with the chosen finite scale δ.

If the signatures of B are outside the permitted family, the second primitive hypothesis gives the vanishing. Otherwise its rational Hermitian isometry class is that of the corresponding V'.

Indeed the finite unimodular normalizations give the same local Hermitian isometry classes as the reference form, after accounting for the scale factor $\delta/\jmath$ in both; the real signatures are the chosen ones. The Hasse principle for Hermitian forms over a quadratic extension (Landherr's theorem) supplies a rational a_0 such that

$$a_0^* J_{V'} a_0 = B.$$

Apply the rational Levi automorphism to the argument so its frame coordinate becomes $q' = a_0 q$. Conjugation of the centre index then identifies the original coefficient at T with the coefficient at $\jmath J_{V'}$ at the new argument. In the notation of the finite normalizing frame this argument has coordinate $Y = x_{V'}^{-1}\kappa^{-1}a_0 q$, where κ is the source's scale idele frame correction. The defining identity of the normalizing frame gives

$$(Yk)^*(Yk) = k^* Q_f k = I$$

in those finite Hermitian coordinates. Thus Yk belongs to the integral isometry group in the precise primitive normalization used in h_0. This is the first primitive assumption, and the required coefficient vanishes. In both signature cases we have proved the base of Theorem 37.1. That theorem, with the radical hypothesis just verified, gives

$$(\mathscr{D}\Psi_0)_{D,T}(x) = 0 \tag{37.3}$$

whenever T_P is unimodular and x is supported on the complementary places.

37.5 Promotion from the primitive coefficient to the standard datum

Theorem 37.2 (The equivariant-lift reduction of the standard column)**.** *Under the hypotheses just stated, including the two primitive geometric assumptions, $\mathscr{D}$ kills every standard-column datum of the prescribed integral type. This is* `IsEquivariantLift.apply_stdColDatum_eq_zero`.

Proof. Use the stable-kernel and unimodular synthesis argument of the earlier standard-coefficient chapter (the test-functional condition (33.3)). We recall why the precise test functionals needed there vanish here. Use the opposite integral frame to the one defining the standard column; its first inverse row is $s = K^\flat$ and the complementary column is K. At an argument x supported outside $P \cup \bar{P}$, each rational centre coefficient of $\mathscr{D}\Phi$ is a linear functional $\Lambda_{T,x}$ of the datum Φ in the source's rectangular Schwartz space. Right centre conjugation and the equivariant law show that it is an eigenfunctional under the shear family of the opposite frame, with eigencharacter indexed by T_P. If this index is unimodular, (37.3) says that it annihilates Ψ_0. Hence every such test functional satisfies the primitive vanishing premise in (33.3); for an index that is not unimodular, that premise is vacuous.

Let M be the synthesis subspace given by that earlier argument. For $\Phi \in M$, its defining test-functional condition kills every rational centre coefficient of $\mathscr{D}\Phi$ at every such x. Fourier uniqueness on the compact centre then gives $\mathscr{D}\Phi(x) = 0$. This time the full rational set of centre characters is used, since the synthesis condition applies both to unimodular and non-unimodular indices. By left rational invariance and weak approximation, these arguments determine $\mathscr{D}\Phi$ globally. The weak approximation needed here is only at finitely many completions at a time: for the rational unitary group, the Cayley transform of the skew-adjoint matrix space gives a rational open coordinate chart. After multiplication by a suitable rational norm-one scalar, any finite tuple of target elements lies in this chart (avoid finitely many eigenvalue obstructions). Approximate its coordinates simultaneously by rational matrices and

apply the chart. The same argument includes archimedean places. Since the other components of the arguments can be prescribed off P and the archimedean places, the indicated left rational translates are dense in the adelic group.

Thus M is contained in the kernel of $\mathscr{D}$. That kernel is stable under the admissible integral transformations by the equivariant law. The unimodular synthesis proves that any stable subspace with the indicated test-functional property contains every integral standard-column datum of the desired rank, including those with nonunimodular stack in the prescribed support. Applying it to the kernel gives the conclusion. □

37.6 Closure record and handbook references

- The source `freitag` is replaced by Theorem 37.1: nested Fourier uniqueness, line-radical density, finite integral flag decomposition, determinant-index comparison, and the opposite-centre phase each appear explicitly. The base is unimodularity, not mere integrality.

- The source `IsEquivariantLift.apply_stdColDatum_eq_zero` is replaced by Theorem 37.2, conditional exactly on the two primitive geometric clauses of h_0; it does not itself assert those clauses. Their geometric proof belongs to the exchange chapter.

- The background theorems invoked are standard: the elementary divisor theorem over a DVR for the index formula; Fourier uniqueness on a compact abelian group (Stone–Weierstrass applied to characters); the local classification of unimodular Hermitian forms (whose needed unramified and split cases were explained above); and Landherr's local–global theorem for Hermitian spaces over quadratic number-field extensions. The latter can be consulted in W. Scharlau, *Quadratic and Hermitian Forms*, Springer, Grundlehren 270 (1985), in the treatment of Hermitian forms over number fields; cf. R. Jacobowitz, *Hermitian forms over local fields*, American Journal of Mathematics 84 (1962), 441–465, for the local classification. The global Hasse principle is Landherr, not a local-field theorem.

- C04 and C24's abstract singular-coefficient and standard-datum interfaces can therefore be read as proved relative to the stated geometric input and the previously established synthesis. Claiming the overall geometric vanishing still requires the precise primitive hypotheses of the exchange construction, not a general assertion about singular Fourier coefficients.

Chapter 38

Complementary-rank exchange: real sections, matched coefficients and the line descent

The equality of two theta kernels in a range of rank is not by itself an exchange of their periods. Three different constants occur: the scalar of the oscillator in an adapted frame, the Haar-measure ratio in the unfolding, and the square norm of the equal-rank lift. We separate these. The notation for the rapid oscillator and the line Gauss sums is that of Chapters 9, 36; in particular the vacuum v_0 has norm one and both adelic mixed-coordinate Haar measures are normalized as there.

38.1 The mixed name and its exact finite support

Let D split the source into a hyperbolic pair of size b and its residual space W_1. Write its adapted frame as $A = (K, I, L)$, its inverse as $(K'; I'; L')$ in the corresponding order; use the integral frames of `Lib/SiegelWeil/SingularForms/AdaptedHom.lean`. For a target T denote its normalizing matrix by $T.x$, with J_T the matrix in the normalized coordinates. The linear identification on Hom, with columns ordered K, I, L, is

$$(1 \otimes K'^{t},\ J_T \otimes L'^{t},\ 1 \otimes I'^{t}), \qquad t = (T.x^{-1})^{*} \otimes 1.$$

This is the identification `frameAt_homSource` followed by the change of basis in `adaptedHom Basis`. In particular $wK = y, wI = z, wL = u$ give a decomposition of the original matrix as $yK' + zI' + uL'$. The signs in the middle hyperbolic identification are units; changing these signs on both factors of a dual pairing leaves the Haar measure and the conductor lattice unchanged.

Here is the resulting finite slicing identity. Suppose the named datum in this frame has finite function

$$\Phi(w) = \Psi(wI) \prod_i 1_O(wL_i) f_i(wL_i)$$

on the selected split places, with any specified finite indicator in the wK coordinates included in Ψ. On the mixed slice at $t^{-*}\alpha x$ (where α is the conductor-scaling idele), its finite function is obtained by

$$F_x(z) = \Psi(z)\, 1_{\{\text{specified } xK \text{ integral}\}} \prod_i \int_{O_i} \psi_i(\pi_i \overline{x_i}\, u_i) f_i(u_i)\, du_i. \tag{*}$$

Off the selected split places and their conjugates one imposes the original conductor-lattice condition on x. In this notation π_i is the coefficient of the local pairing, not necessarily a uniformizer. The residual function is named on the residual real slice of the datum and multiplied by the same nonzero scalar $\kappa_{D,T}$ for all finite functions in this formula (the scalar denoted `vacuumFactor` in the source). Indeed the block identification above turns the hyperbolic Heisenberg translations into multiplications by the characters $\psi_i(\pi_i\overline{x_i}\,u_i)$; the other coordinates are unchanged. On a finite indicator and a character, the Fourier integral is thus precisely the right-hand side of (*). Linearity and the decomposition of a finite locally constant, compactly supported function into indicators of lattice cosets prove the formula for every such function. The analogous statement with an unselected local factor is the characteristic function of the self-dual conductor lattice, by Fourier orthogonality. These are the finite factors of the mixed expansion (9.12). The scalar comes from the identification of the two normalized vacuum vectors; its nonvanishing follows from unitary intertwining of the oscillator and the nonvanishing of the vacuum slice. The measures on the finite O_i have mass one, so there is no additional volume depending on a name.

For the standard square name, write $P = (K, I)N$ for the frame used to form its square matrix and denote by χ^{-1} the extension of the inverse tame character by zero outside the units. The name is

$$\Phi(w) = 1_{\{w \text{ integral}\}}\epsilon\,\chi^{-1}\big(\det((wK, wI)N)\big), \quad |\epsilon| = 1.$$

It is essential that the character is extended by zero: $\Phi(w) \neq 0$ implies the indicated square matrix is invertible over the local integer rings on which the character is taken. In this case $f_i = 1$ on O_i in (*). Orthogonality imposes integrality at the conjugate places, so altogether the mixed coefficient vanishes unless its finite argument y is integral. Its remaining finite function is

$$F_y(z) = \epsilon\,1_{\{z \text{ integral}\}}\,\chi^{-1}(\det((y, z)N)). \tag{**}$$

This accounts for both halves of the finite integrality test, not just for the place at which the determinant character was defined.

38.2 The standard radical and Levi term

Use the notation of `thetaTerm_stdDatum_spec` in `New/Exchange/LowRankIdentity.lean`. Let $p = n(c, z)m(q, g)$ in the parabolic of D, where z here denotes the central radical coordinate (not the variable z of (**)). Set $a = \dim T$, $x' = T.x^{-1}x$, $x'q = \alpha Y$, $y = k_f^{-1}Y_f$; let k be the target point relative to the fixed coordinate change. Write $\beta' = \beta v_1^b$. With the source's Hermitian trace convention, the term indexed by the rational frame x is exactly

$$\psi\Big(\tfrac{1}{2}\operatorname{tr}(zx^*J_Tx)\Big)\ (\gamma.\mathrm{spaceChar}(a))(\det q)^{-1}|\det q|^{a/2}$$
$$\cdot\,1_{\{y \text{ integral}\}}\kappa_{D,T}\,\theta_{\mathrm{real}}\Big(W_1, T; \beta, \gamma, F_y, x'c^*J_{W_1}, (g, k), \mathrm{realSlice}(D, T, k^{-1}Y)v\Big). \tag{38.1}$$

Here θ_{real} is the full theta of the residual finite name tensored with the specified real vector and with the displayed Heisenberg translation; it is not a theta function of only real variables. The base-change in source rank replaces the source splitting character by β' exactly as in the zero-term law.

To prove (38.1), first act by $n(c, z)$. In the hyperbolic mixed model the central skew-Hermitian parameter acts on the x term by $\psi(\frac{1}{2}\operatorname{tr}(zx^*J_Tx))$ and the off-diagonal parameter acts on the residual wave by translation $x'c^*J_{W_1}$. This follows by composing the Heisenberg translations in (9.11); the central half-pairing is the one of that law. The Levi action on the

same wave is precisely the zero-term determinant scalar of Theorem 9.5, with coordinate $x'q$. Moving the target point to the wave argument gives $k^{-1}Y$. Apply the finite calculation (**) and then the defining residual theta. This proves the formula, including the scalar and the character; no invariance of an unspecified local model under a compact subgroup is used.

The orthogonal-target-sum version of this calculation is worth recording. When $T = V' \oplus W_1$, the residual finite name is the product of the two finite names, the real datum is the box product, and the translation is in the V' summand only, the right-hand side separates into the lower theta and the equal-rank theta for W_1. The assertion is the product law of Corollary 9.4: the symplectic form is the orthogonal direct sum, so all Heisenberg phases and the normalized theta sums multiply. This is the content of `thetaOnRealData_boxProd_eq_mul`.

38.3 Absolute coefficient extraction for the standard square name

The Fourier coefficient of a period needs more than integrability of the total theta kernel. In this particular standard-name situation, the sum of the absolute values of the mixed terms is integrable in the target variable. Here is the strengthened growth argument.

Fix the source translate. The support condition on the translated name is that the matrix obtained by applying w to the translated (K, I) is right-invertible. For the standard name this is exactly the invertibility just established, since a parabolic element maps the frame (K, I) to itself by a block triangular invertible matrix. Work initially with a finite collection of rational mixed terms and multiply each term by an arbitrary scalar of absolute value at most one. The estimates of the absolute mixed expansion of Chapter 9.7 bound each such finite phased sum by the same weighted oscillator norm, independently of the collection and the phases.

In a target Siegel set at large height the reduction argument of Chapter 10 supplies a short rational isotropic vector d. Choose rational source directions in the span of (K, I), scaled to be integral in the adapted frame. Translations in the target direction d, with these source directions, have zero K-projection: the directions are orthogonal to K. At rational translation parameters each individual mixed term is invariant. Indeed the central phase on its rational argument is the trivial character on F; the mixed argument is unchanged and its residual theta undergoes a rational translation. As in the proof of the growth estimate one averages each finite phased sum over the compact additive-parameter quotient. The orthogonality of averaging annihilates its named datum unless the prescribed affine functional of the square matrix vanishes on the chosen short line. That functional cannot vanish on every direction of (K, I) for a right-invertible square matrix. Thus the averaged finite component vanishes. Taking successive finite differences of the real translates, or equivalently applying the weighted estimates to the derivatives of the real translates, gives, for every N,

$$\left| \sum_{x \in B} \eta_x T_x(h) \right| \le C_{N,g} M_{N'}(v) H_T(h)^{-N}, \qquad |\eta_x| \le 1.$$

The constants in the difference estimates depend on the fixed source translate, not on B or the phases; the proofs use the absolute mixed expansion and its weighted-norm bound at each step. Choose the phases of the finitely many terms so that the terms become nonnegative. Exhausting the rational index set by finite sets gives the same bound for the full sum of absolute values. Since the target quotient has finite volume, this proves absolute integrability of the series. It is the strengthened assertion `hasIntegrableTerms_integrated_name`, under its right-invertibility hypothesis, rather than the assertion only for the whole theta.

Consequently taking the Fourier coefficient in the compact central radical quotient commutes both with the series and with target integration. Additive character orthogonality applied to (38.1) retains exactly the rational frames with the prescribed Gram matrix. For nonsingular Gram matrix, all such frames, if any, form one rational target-unitary orbit, by Witt extension; its stabilizer fixes the frame pointwise and is the unitary group of its orthogonal complement. Summing over that orbit and integrating unfolds to the stabilizer quotient with the quotient Haar measure. If the Gram matrix is not represented there are no terms. These are the hypotheses needed when invoking the ordinary stabilizer unfolding, not assertions of formal cancellation in a divergent series.

38.4 A smooth section of real mixed evaluation

We prove the precise real interpolation used in `MixedModel/RapidSections.lean`. Let x run through the real mixed-argument space, let ξ be a rapidly decreasing residual oscillator vector, let τ be smooth and compactly supported, and let $U(x)$ be a smooth family of residual symplectic transformations on a neighbourhood of its support. There exists a rapid full vector Λ with

$$\Lambda[x] = \tau(x)\omega(U(x))\xi. \tag{†}$$

Work in adapted real coordinates L, I, K where the normalized full vacuum has real slice $e^{-\|x\|^2}v_{0,I}$. Let ι embed the residual Fock space by the vacuum in the removed factors (so its slice is $e^{-\|x\|^2}$ times its residual vector). Set

$$G(x) = e^{\|x\|^2}\tau(x)\iota\omega(U(x))\xi.$$

This is compactly supported and smooth in every rapid weighted norm. For completeness, the coherent resolution of the residual vector expresses it as an absolutely converging integral of coherent translates of the vacuum with a rapidly decreasing coefficient. Differentiating the explicit Gaussian kernel after smooth symplectic transformation produces polynomial factors in the coherent parameter, and the transformed Gaussian has strictly positive real part on the compact set of parameters $U(x)$. Each derivative of order k is bounded in the norm M_N by a polynomial in the coherent parameter times its rapid coefficient, so dominated convergence proves the asserted differentiability. Integration by parts in the compact x support now shows that the vector Fourier transform $\widehat{G}(b)$ decreases more rapidly than every power of $\|b\|$ in each weighted norm. Use the dual character of the mixed Heisenberg law, namely $\beta(2x, b)$, and its dual Haar measure, in this Fourier transform.

Define the full vector

$$\Lambda = \int \rho(Kb)\widehat{G}(b)\,db.$$

Translations by Kb have polynomial growth in each weighted norm, so the integral is absolutely convergent in every such norm. Taking a slice is continuous in these norms by Lemma 36.6. Fourier inversion and the exact law

$$(\rho(La + Iw + Kb)v)[x] = \beta(2x + a, b)\rho(w)v[x + a]$$

therefore give (†). In the opposite polarization the zero slice is the integral of the right-hand side of (†). To check the normalization, both polarizations have real vacuum slice equal to the residual vacuum at zero. On the real mixed space the source's measure is the Gaussian reference measure, for which $\int e^{-\|x\|^2}\,dx/\pi^d = 1$ in complex dimension d. Applying the complementary Fourier formula to the vacuum determines the proportionality constant as exactly one. Equivalently, summing and integrating the compact-quotient form of (9.12)

derives the same identity with its product finite factor; cancel that nonzero common finite vacuum scalar. Thus for $U = 1$ and $\int \tau = 1$, one obtains the original ξ as the opposite zero slice, as required in the construction below. This proves the section and normalization, not merely density of its image.

38.5 Smooth Witt motion near a nondegenerate frame

We give a real local construction in place of a choice of isometries with an unproved smooth-dependence assertion. Let $T = V' \oplus W_1$ have real Hermitian matrix $J = \operatorname{diag}(J', J_1)$, and $x_0 = (1; 0)$ be the frame of the first summand. Near x_0 write a frame as $(Y_1; Y_2)$ with Y_1 invertible and $Z = Y_2 Y_1^{-1}$. Define

$$k(Z) = \begin{pmatrix} 1 & -J'^{-1} Z^* J_1 \\ Z & 1 \end{pmatrix} \begin{pmatrix} (1 + J'^{-1} Z^* J_1 Z)^{-1/2} & 0 \\ 0 & (1 + Z J'^{-1} Z^* J_1)^{-1/2} \end{pmatrix}.$$

For sufficiently small Z the inverse square roots are the convergent binomial power series, hence are smooth and self-adjoint for their respective Hermitian metrics. The two unnormalized columns of blocks are orthogonal; their Gram matrices are respectively $J'(1 + J'^{-1} Z^* J_1 Z)$ and $J_1(1 + Z J'^{-1} Z^* J_1)$. Thus $k^* J k = J$. Its first subspace is the graph of Z, exactly the span of the prescribed frame. Multiplying Z by a cutoff before applying this formula gives a globally defined smooth k agreeing with it on a smaller neighbourhood. This construction applies at each infinite place and to indefinite J', J_1 as well.

38.6 The top exchange real matching construction

Let $n = \dim W_1 = 2j$, and let μ_ν be the infinite weights of the splitting character in `exists_stdExchange_top`. For clarity write z_B for the idele supported at infinity with determinant coordinates $D_\nu = \det B_\nu$. The archimedean factor to be matched is

$$\nu^{bc}(z_B)|z_B|^{n/2} = \prod_\nu (D_\nu / \bar{D}_\nu)^{\mu_\nu} |D_\nu|^{2j}. \tag{‡}$$

Choose a nonnegative smooth compactly supported σ on the tuple of real frames, positive at x_0, supported so near x_0 that their Gram matrices are invertible, and that the preceding smooth Witt motion is valid. Let Γ extend matrix inversion smoothly on a neighbourhood of J' (using a cutoff inside the open set of invertible matrices). For any frame x put

$$y = Z(x) = x\Gamma(x^* J x) x^* J x_0, \qquad w = w(y, x) = k(y)^{-1} x.$$

Where both $\sigma(y)$ and $\sigma(w)$ are nonzero, Γ agrees with inversion, since $x^* J x = w^* J w$ is near J'. Thus y is the projection of x_0 onto $\operatorname{span}(x)$ for the metric J; it depends only on that subspace. Since y is a full frame on the support of σ, this subspace is also the span of y.

Set $\tau'(B) = \sigma(x_0 B) \prod_\nu |\det B_\nu|^{2(j+\mu_\nu)}$ and let $l = (\int \tau')^{-1}$. This integral is strictly positive and finite. Given a rapid lower vector v, the smooth-section construction yields the lower vector Λ' with mixed slice $l\tau'(B)v$ and opposite zero slice v. Given also the equal-rank vector v_1, construct the upper vector Λ with mixed slices

$$\Lambda[x] = l\, \sigma(Z(x))\, \sigma(w(Z(x), x)) \prod_\nu \overline{\det w(Z(x), x)_{\mathrm{top},\nu}}^{\,2\mu_\nu} \cdot \Omega(1, k(Z(x)))(v \boxtimes v_1). \tag{38.2}$$

This is a valid smooth section: on its support both y, w lie in fixed compacts and $x = k(y)w$ lies in a compact; the determinant power has no singularities there. The tensor product is rapid by the orthogonal sum of the weighted coherent estimates. The construction of the preceding subsection then applies.

Define $\rho(h) = \sigma(Z(h^{-1}x_0))$ on the real target group. It is nonnegative and positive on a neighbourhood of 1. If $\rho(h) > 0$, put $y = Z(h^{-1}x_0)$. The isometry $k(y)$ sends the first summand onto the span of $h^{-1}x_0$, so there is $u \in U(V')$ with $k(y)x_0u = h^{-1}x_0$. Pointwise stabilization of the orthogonal complement then gives $q \in U(W_1)$ such that

$$\operatorname{diag}(u, q^{-1})hk(y) = 1.$$

Write $p = (u, q^{-1})$. At $x = (\operatorname{diag}(p)h)^{-1}x_0B = k(y)x_0B$, the projection $Z(x)$ equals y wherever either of the two relevant slices is nonzero. Indeed in the support of the upper slice its Gram lies in the inversion neighbourhood, and in the lower slice this follows from $\sigma(x_0B) \neq 0$. Hence the two expressions vanish together outside this support. Multiplication by $\Omega(1, \operatorname{diag}(p)h)$ cancels the last factor in (38.2). Dividing the determinant weight of the upper slice by that of the lower slice gives the inverse of (‡), because

$$\frac{\bar{D}^{2\mu}}{|D|^{2(j+\mu)}} = (D/\bar{D})^{-\mu}|D|^{-2j}.$$

If $\rho(h) = 0$, an upper slice at any $h^{-1}x_0u^{-1}B$ cannot be nonzero: in its support true inversion would give the projection onto the span of $h^{-1}x_0$, contradicting the condition $\sigma(Z(h^{-1}x_0)) = 0$. Take $p = 1$ in this case.

The orthogonal-sum identity following (38.1) now says that the integrated upper residual theta equals the lower residual theta times $\rho(h)$, the inverse archimedean factor (‡), and the equal-rank period

$$\lambda(v_1) = \int_{[U(W_1)]} \theta(W_1, W_1; v_1; 1, k_1)\, dk_1.$$

The value of the latter period at other source points is equal to its value at 1 by the even-rank constancy argument of Chapter 35.10, using the line-cusp calculation below; it is not a pointwise constancy of the unintegrated theta. Thus the real matching required in `stdExchange_of_periods` follows for every source point and every real h, with the actual power of the determinant, the actual normalization of the opposite slice and a nonnegative ρ nonzero at the identity. Nothing in this construction asserts that $\lambda(v_1)$ is nonzero; that is a separate period problem.

38.7 The missing adapted-frame scalar in the line calculation

We complete the finite line computation of Chapter 36.9, in the frame prescribed by `exists_lineTerm_localDatum` of `Lib/SiegelWeil/Tower/LineGaussSum.lean`. In this computation $Q = (Q.K, Q.I, Q.L)$ is the adapted integral frame of the target and its inverse is integral. Write H for the block matrix which is 1_b on the fixed rows and Q on the other rows, with the row and column orderings of the line, residual block, and opposite line. Explicitly its columns are $(0; Q.K)$, the block $1_b \oplus Q.I$ with the indicated ordering, and $(0; Q.L)$, followed by the inverse row permutation. Let R be the source row-permutation matrix. This is exactly the source matrix `HKLI` with $K = Q.K, L = Q.L, I = Q.I$. Because the block-diagonal matrix and the two permutations are integral invertible, so are H, R and their inverses. Therefore

$$\epsilon = \chi^{-1}(\det RH), \qquad |\epsilon| = 1.$$

Multiplying the square-name matrix by these matrices preserves integrality in both directions and changes the determinant character by this fixed scalar. In the actual slicing coordinates the resulting matrix is block upper triangular with diagonal a, Y, t or t, Y, a, where a is the prescribed line coordinate, Y is the residual square name, and t is the Fourier variable of the opposite line. The off-diagonal coordinates are either zero-padded fixed rows or unrestricted integral coordinates with Fourier argument zero. In both orders the determinant is $at \det Y$. Thus the integrality tests are exactly the tests for the diagonal blocks and the free off-diagonal integral variables. Every free-coordinate integral is one in the normalized lattice Haar measure.

What remains at the nontrivial local place is the integral

$$1_O(a)\chi^{-1}(a)\int_O \psi(\pi\bar{a}\,t)\chi^{-1}(t)\,dt$$

times the lower square name. At the conjugate place the untwisted indicator at the corresponding conductor is imposed by orthogonality. The integral with character extended by zero is exactly the Gauss integral of Chapter 34.1, with absolute square $|O/\mathfrak{p}|^{-1}$ in the nonzero branch; outside the prescribed unit branch it vanishes by translation by the conductor subgroup. The finite constants are consequently the unit ϵ and the Gauss factors, never a variable depending on the residual datum. The product, denoted γ in the line lemma, has the absolute square specified there. In particular the vacuum ratio l_β of the two normalized slices is fixed by the vacuum evaluation, as in (*); it introduces no new character-dependent scalar. Combining this frame calculation with the triangular-block computation and the absolute Mellin endpoints of Chapters 36.9 and 36.10 gives the normalized constant term on the two branches. This is the adapted-frame step missing from C23.

38.8 Matching the finite and infinite periods with their Haar measures

We use the definitions of `thetaLarge` and `thetaSmall`: on a source of even rank $n+2b$ the two characters on the large side are both $\mathrm{rankTwist}_{n+2b}(\nu^{-1})$, whereas on the small side the first is $\mathrm{rankTwist}_{n+2b}(\nu)$ and the second is υ_1^b. The small name is transposed and read at the conjugate places. On the residual rank n put $A = \mathrm{rankTwist}_n(\nu^{-1})$. In the real-product formula the lower residual theta uses (A, υ_1^b), and the equal-rank one uses (A, A).

Suppose the matched real slices have the property of Section 38.6, denoting the character-dependent equal-rank period by $\lambda_{\mathfrak{c}}$. More generally suppose they satisfy the displayed hypothesis `hc` of `stdExchange_of_periods`, with $\lambda_{\mathfrak{c}}$ any given scalar. Let

$$a_0 = \det(\alpha V'.x), \qquad \kappa = \frac{\kappa_{D,V'\oplus W_1}}{\kappa_{D,V'}}|a_0|^{n/2}.$$

Both are nonzero and free of the varying character. The standard row name at the adapted frame has the unit phase $\epsilon_{\mathfrak{c}}$ arising from its integral frame change. We claim that the nonsingular central coefficients of the difference of the two standard lifts required by the promotion theorem, with scalar

$$c_{\mathfrak{c}} = R\,\kappa\,\epsilon_{\mathfrak{c}}\nu^{bc}(a_0)\lambda_{\mathfrak{c}} \tag{1}$$

are zero whenever the upper real slices are supported on Gram matrices represented by the signs of V'. Here $R \neq 0$ is a comparison of fixed quotient Haar measures, not a local character factor. Explicitly the two classes of coefficients in `IsEquivariantLift.apply_stdColDatum_eq_zero` are: the coefficient indexed by $J_{V'}$ at parabolic points with $q = \alpha V'.xY$ and $Y_f k \in U(1)$ for some $k \in \mathrm{GL}_b(\widehat{O}_E)$; and every invertible index not represented at the real places by the lower

signs. We prove precisely these two comparisons, not a claim of rational isometry for every Gram with the same real signs.

First unfold the coefficient with Gram matrix $J_{V'}$. Its upper rational frame is $x_0 = (1; 0)$ with stabilizer $U(W_1)(F)$; its lower frame is 1 with trivial stabilizer. These are the stabilizer quotients of Section 38.3. On the upper quotient integrate in stages over the closed block-diagonal group $U(V')(\mathbb{A}_F) \times U(W_1)(\mathbb{A}_F)$; in the first factor the integration is over the *whole* adelic group and in the second over its rational quotient. This distinction is forced by the pointwise frame stabilizers. To see that there is one measure constant R, write G for the upper target group, $L = U(V')(\mathbb{A}_F)$, $H = U(W_1)(\mathbb{A}_F)$, and let

$$\rho_0(g) = \rho(g_\infty)1_{\{\exists\, l_f \in (L\times H)_f:\ l_f g_f \in K_f(G)\}}.$$

By quotient integration for the closed embedded unimodular group $L \times H$, the integral over $H(F)\backslash G$ is the integral over $(L \times H)\backslash G$ of the inner integral over $L \times [H]$. This inner integral is unchanged by multiplication on the left of g by any element of $L \times H$. If it equals $\rho_0(g)$ times a scalar independent of g, the outer integration contributes one constant. Indeed if any such nonzero scalar occurs for an absolutely integrable coefficient, ρ_0 is integrable for the induced orbit measure and its integral R is positive, since ρ_0 is positive near 1. If all such scalars vanish the asserted identity holds with any fixed nonzero R. This is the measure argument of `exists_integral_stages_eq_mul`; it explains why no character-dependent mass is introduced in (1).

For g in the indicated finite coset, left multiplication by an element of $L \times H$ puts its finite component in $K_f(G)$. The upper standard name is fixed by this finite integral compact subgroup: the matrix change is multiplication of rows by an integral unitary element, the determinant character on the square name cancels $\mathrm{rankTwist}_{n+2b}(\nu^{-1})(\det g_f)$ in the pair action, and outside the named places the vacuum is fixed. This is also the direct proof of `pairWeil_stdRowDatum_eq_self`. We may thus compare terms using Y in the lower rank b coordinates with

$$B = y^{-1}Y, \qquad x = B_f, \qquad \det q = a_0 \det y \det B,$$

where $y \in L$ is the lower target point. At the finite places the upper integrality condition is exactly the lower one; if it holds, x and x^{-1} are integral since the translated frame is unitary relative to its normalized Gram. In particular $u = \det B_f$ is a finite unit idele. The full square name of the upper term is now block triangular with blocks B_f and the residual matrix: its finite named vector is

$$\epsilon_{\mathfrak{c}}\chi^{-1}(\mathrm{ev}_S\, u)(F \boxtimes F_1), \qquad F_1 \text{ the standard equal-rank name.}$$

The lower finite vector is $\chi^{-1}(\mathrm{ev}_S\, \bar{u})F$; this conjugation is the reason one reads its transpose at the conjugate places. These identities follow by expanding the determinant of the block upper triangular matrix, using the inverse-character-on-units product identity (including its zero value outside units), and applying the finite slicing formula (**). The product name is a tensor product since both the finite split evaluation and the projected vacuum respect direct sums. If the condition fails, both terms vanish.

Here is the entire scalar calculation. Write υ for the splitting character on the ideles and note $\upsilon_U^{bc} = \upsilon^2$. The upper Levi character $\mathrm{rankTwist}_{n+2b}(\nu^{-1}).\mathrm{spaceChar}(b+n)$ equals $\nu^{bc,-1}\upsilon^b$; the lower Levi character $(\upsilon_U^b).\mathrm{spaceChar}(b)$ equals υ^b. In the inverse-character half-density factors in (38.1), the ratio is therefore

$$\nu^{bc}(\det q)|\det q|^{n/2}.$$

The remaining scalar factors multiplying the lower term are

$$\begin{aligned} \text{mixed vacuum:} &\quad \kappa_{D,V'\oplus W_1}/\kappa_{D,V'},\\ \text{upper versus lower name:} &\quad \epsilon_{\mathfrak{c}}\chi^{-1}(\mathrm{ev}_S\, u)/\chi^{-1}(\mathrm{ev}_S\, \bar{u}),\\ \text{real matching:} &\quad \rho(g_\infty)\lambda_{\mathfrak{c}}/(\nu^{bc}(\det B_\infty)|\det B_\infty|^{n/2}),\\ \text{lower residual character conversion:} &\quad \nu(\det y)^{-2}. \end{aligned}$$

For the last line the real product was matched to a lower kernel with first character A, not the actual lower first character $\mathrm{rankTwist}_n(\nu) = A\nu^2$. The ratio for the same target point y is indeed $\nu(\det y)^{-2}$. Inserting $\det q = a_0 \det y \det B$ into the Levi ratio cancels this line, because $\det y$ has norm one and $\nu^{bc}(\det y) = \nu(\det y)^2$. The archimedean factor cancels with the preceding line. On the finite unit idele u we have $|u| = 1$ and

$$\chi^{-1}(\mathrm{ev}_S\, u)\nu_f^{bc}(u)/\chi^{-1}(\mathrm{ev}_S\, \bar{u}) = 1.$$

Indeed $u/\bar{u}$ is integral and its ν-value is the tame determinant character at S, by its defining restriction. This is `SplitTameChar.invOnUnits_mul_div_star`. The remaining product is exactly $\rho(g_\infty)\kappa\epsilon_{\mathfrak{c}}\nu^{bc}(a_0)\lambda_{\mathfrak{c}}$, times the lower kernel. Notice especially the exponent $n/2$ on the absolute value: replacing the Levi ratio by a determinant character would lose this factor of the normalizing frame.

If g is not in the finite block coset then the upper term vanishes for every lower integration point. In fact an integral mixed frame in a nonzero term, after multiplying by the inverse of its integral unitary normalization of the Gram, would give an integral orthonormal frame with Gram matrix 1. The integral Witt extension of Section 16.6 extends this frame to an integral unitary matrix. Multiplying by it puts the relative upper point in the stabilizer of the first summand, hence in the image of $L \times H$ modulo $K_f(G)$, a contradiction. This proves the required staged inner-integral relation also in the bad coset. Absolute integrability was checked before unfolding, so the quotient integration just established is legitimate.

It follows that the upper central coefficient on the specified Gram at the prescribed unimodular parabolic points equals the lower central coefficient times (1). For forms not represented by the lower signs at the real places (`RepresentedBy`), the lower coefficient vanishes by its rational frame expansion; the upper coefficient vanishes by its real-slice support and the mixed model formula. These are the nonsingular coefficient comparisons required in the standard-row/column annihilation argument, not only an equality at the identity of the source group.

Finally set $C = |R\kappa|^{-2}$, and recall

$$\ell(\mathfrak{c}) = |\lambda_{\mathfrak{c}}|^2/\Pi_{\mathfrak{c}}(n/2), \qquad \Pi_{\mathfrak{c}}(e) = \mathrm{lValueProdInt}(\mathfrak{c}, e).$$

Since $\nu^{bc}(a_0)$ and $\epsilon_{\mathfrak{c}}$ have modulus one, the exact relation is

$$C|c_{\mathfrak{c}}|^2 = |\lambda_{\mathfrak{c}}|^2 = \Pi_{\mathfrak{c}}(n/2)\ell(\mathfrak{c}). \tag{2}$$

The nonsingular coefficient comparison and the equivariant difference of Section 35.5 give the full standard-column identity upon applying the source's singular-coefficient induction `IsEquivariantLift.apply_stdColDatum_eq_zero`. One must make that last application to the linear functional on the column space, not assert that an arbitrary automorphic form with the indicated nonsingular coefficients zero is identically zero. With this induction, (1)–(2) are precisely `stdExchange_of_periods` under its real support alternative. Together with Section 38.6 they give `exists_stdExchange_top`. The other alternative of the source, with a negative definite lower space and the specific nonnegative upper signature, will be used for nonvanishing below.

38.9 The small-side tower descent on standard-column data

Suppose W' splits adaptedly over W along one hyperbolic plane, with V' similarly of two ranks more than V. Put $e = \dim V - \dim W/2$. Take an exchanged standard pair as in `StdExchange`. The large-side line constant term is now known in the precise normalized form of Sections 36.9–36.10 and 38.7:

$$\int_{[N]} I_{W',V'}(x)(ng)\,dn = \gamma_{\mathfrak{c}} I_{W,V}(\mathcal{C}x)(g), \quad |\gamma_{\mathfrak{c}}|^2 \Pi_{\mathfrak{c}}(e) = D_1 \Pi_{\mathfrak{c}}(e+1), \tag{3}$$

for the character-independent real cone operator $\mathcal{C}$ and $D_1 > 0$. Here the finite columns of the original integral are multiplied by the residual inverse frame I', exactly as in the source's line formula. The cone operator is defined for every rapid real datum, not just polynomial Gaussians, and its image is rapid by the weighted moment proof.

On the smaller target U the radical constant term has the mixed-index expansion over rational isotropic vectors a in U. We show directly that only $a = 0$ can contribute for these standard-column names. Suppose $a \neq 0$. Choose a rational splitting of U having isotropic line Ea; this is possible by completing the hyperbolic pair against a vector pairing nontrivially with a, then subtracting its self-pairing. In the corresponding mixed model every functional giving the constant term indexed by this line is invariant under the Heisenberg translations along the line: it is a zero-slice theta in the target polarization followed by the appropriate fixed group translate. Formula (9.11) at mixed coordinate zero proves this invariance, with the normalized dual-frame matrices giving the translated line $h^{-1}a$ in the matrix coordinate. Average the named datum under these translations over the compact dual-lattice subgroup, using the same character orthogonality as in (*), now with arbitrary source-row parameters. A nonzero finite function would require that its matrix annihilate the nonzero line $h^{-1}a$. But the transposed standard-column name has full rank on its support: the frame used to form it is primitive, and the character on its completed square matrix is zero unless the completion is invertible. Thus no nonzero line can be annihilated. This is the application of `name_eq_zero_of_invariant` to `constTermAt_smul_k`: it requires the right-invertibility of the transposed column name, not merely nonzero determinant of its fixed part. Therefore every nonzero isotropic index contributes zero.

At $a = 0$ use (*) in the opposite polarization. Its residual name is exactly the lower transposed standard column, and the multiplier is the nonzero vacuum ratio $\kappa_A = \kappa_{A,U}$. Taking the small-side line mean gives

$$\int_{[N]} I_{W',U}(y)(ng)\,dn = \kappa_A I_{W,U}(\mathrm{realSlice}(A,U,0)y)(g). \tag{4}$$

The source frame changes use a unit scalar $m_{\mathfrak{c}}$ from `thetaLarge_move`; they change neither the absolute square nor the norm of the datum. Thus, if the old exchange had scalar $c_{\mathfrak{c}}$, equations (3) and (4) give the new scalar

$$c'_{\mathfrak{c}} = c_{\mathfrak{c}} \kappa_A m_{\mathfrak{c}} / \gamma_{\mathfrak{c}}, \qquad C' = \frac{C D_1}{|\kappa_A|^2}.$$

The product $C'|c'_{\mathfrak{c}}|^2$ is $C|c_{\mathfrak{c}}|^2 \Pi_{\mathfrak{c}}(e)/\Pi_{\mathfrak{c}}(e+1)$, exactly $\Pi_{\mathfrak{c}}(e)\ell(\mathfrak{c})$ if the old product was $\Pi_{\mathfrak{c}}(e+1)\ell(\mathfrak{c})$. This closes the normalized one-plane period descent.

For determinant vectors the cone surjectivity of Proposition 33.8 now has its integrability and adapted vacuum hypotheses established; the phase cancellation gives a positive multiple of the smaller vector, and distinct choices of coefficients can be lifted by block-diagonal coefficient extension and real scaling. The adapted splitting of several hyperbolic planes can

be reordered into successive one-plane splittings: take the first pair of isotropic vectors, split its orthogonal complement, and repeat on the remaining orthogonal hyperbolic pairs. Zero real slices compose by Fubini of the normalized Fourier integrals, including their nonzero vacuum factors; on the vacuum all normalizations agree. Thus iteration proves the plane removal map of `exists_stdExchange_removing_planes` and its determinant-vector surjectivity (with the rank and weight hypotheses stated there). This statement concerns the standard exchanged data. A general subtracted theta lift on a noncompact quotient will require a separate vanishing and boundedness argument; one must not infer it from the standard-column case alone.

38.10 Circle vanishing for the negative-definite comparison

The real support condition above is appropriate for constructed top slices, but the nonvanishing proof needs the different signature condition in `stdExchange_of_periods`; here an arbitrary rapid determinant vector is used. The following supplies the requisite replacement for support.

Let the residual source have p positive and q negative signs, and let the target have $q + b$ negative signs and at least one positive sign. At an infinite place write the residual first-character weight as $m < p$. Consider an invertible Gram matrix of a frame with b columns which is not negative definite there. Its image contains a positive vector. By a real unitary change arrange that a positive unit vector in this image is a positive coordinate vector. The negative-row projections of its b columns consequently span a subspace of dimension at most $b - 1$ in the negative-coordinate space of dimension $q + b$.

For any coherent vector parameter y in the residual Hom space choose a unit negative row direction r orthogonal to that negative projection and also orthogonal to the q columns of y corresponding to negative source signs. Such a direction exists since at most $(b-1)+q < q+b$ independent equations were imposed. Put $P_r = rr^*$ for its orthogonal projection. The circle

$$k_\eta = 1 + (\bar\eta - 1)P_r, \quad |\eta| = 1,$$

is compact unitary and fixes the frame pointwise. Applied to the coherent vector $\rho(y)v_0$ the real oscillator operator of the target point k_η is

$$\eta^{p-m}\rho(y + (\bar\eta - 1)P_r y)v_0.$$

Indeed the negative-target vacuum exponent is $m - p$ on the eigenvalue $\bar\eta$, by the polynomial formula `realPairWeil_polyVector` (or the real Gaussian vacuum multiplier); covariance of the Heisenberg representation gives the shifted coherent parameter. By the orthogonality condition on the negative-source columns, $P_r y$ is supported only on oppositely signed Hom coordinates. The holomorphic Fock coordinates conjugate those coordinates. Hence every coherent evaluation of the displayed vector is a positive power η^{p-m} times a holomorphic entire function of η: the norm of its shifted parameter is unchanged on the circle, and the linear exponential in a coherent evaluation has no $\bar\eta$ term. Its circle mean is zero by the power series, since $p - m > 0$.

Any continuous linear functional on the real Fock space invariant under the compact subgroup fixing the frame must consequently annihilate every coherent vector. The choice of the circle may depend on the coherent parameter; this is allowed because the functional is invariant under the whole compact stabilizer. The coherent vectors span a dense subspace, so the functional vanishes. To apply this to the central coefficient, unfold it to its pointwise frame stabilizer as in Section 38.3. Average the term over the compact subgroup of the stabilizer at the selected infinite place; Haar measure of that subgroup is normalized to one. The integral

is unchanged by this averaging because the stabilizer is unimodular and the coefficient is absolutely integrable. At a fixed remaining target point the averaged functional on its real slice is of the invariant kind just shown to be zero (move the fixed frame to the convenient real coordinates first). Thus the unfolded coefficient is zero.

In the signature alternative of `stdExchange_of_periods`, the residual rank is $n = 2j$, the residual weight is $m = j - \mu$, and $q \le j$ with $\mu > 0$. Hence $m < j \le n - q = p$. The lower space is totally negative. Every invertible Gram not represented by it is nonnegative in some direction at some real place, and the argument above applies. This proves the alternative used in the source `centreCoeff_eq_zero_of_not_negative`. In conjunction with the matching for the represented Gram and Theorem 37.2, the top standard-column identity does not require real-slice support in this signature case.

38.11 Nonvanishing and independence of the equal-rank ratio

Let W have rank $2k + 2q$, where q is its number of negative signs. Let $\mu_\tau \ge k$ be the weights, and use the tame-character weight convention of the source (strictly positive). Choose the definite space J of rank $2k$, and an adapted space X of the same signs as W, splitting over J along q hyperbolic planes. Such a space is constructed by taking the orthogonal sum with the hyperbolic planes and reordering the signs. Set $Y = H \oplus X$ and let Z be the rank-zero space. Put

$$\mathcal{R}_Y(\mathfrak{c}, v) = \frac{|I^{\mathrm{large}}_{Y,Y}(\mathfrak{c}; v, \mathrm{standard})(1)|^2}{\Pi_{\mathfrak{c}}(\dim Y/2)}.$$

It is nonnegative and the denominator is positive by the Hecke endpoint argument. This notation is the source's `equalRankThetaRatio`. Moving an adapted space to the given one by rational isometry and the unitary normalization of its adelic frame changes the theta lift by a unit phase, not an absolute value. To see this, the induced change of rational Hom matrices sends rational translates bijectively to rational translates; the normalized Weil action identifies the two theta functionals by uniqueness on these translates, with identical vacuum-modulus normalization. Changing the variable of quotient integration by the same isometry preserves the normalized quotient measure. The remaining pair character is a unit-modulus determinant scalar, precisely `thetaLarge_move`. Thus it suffices to prove the assertion for Y.

If the family of tame characters is empty the uniform-ratio assertion is vacuous; otherwise suppose for contradiction that $\mathcal{R}_Y(\mathfrak{c}, v) = 0$ for every character and every rapid real vector. The now-established line identity and Proposition 35.10 give the exact constancy of the equal-rank theta integral. Therefore it is zero at every source point, not just at 1, for every rapid real vector.

Use the upper target $J^- \oplus Y$ and the lower target J^-, where J^- has the opposite definite signs. The top source is $X^{\mathrm{top}} = Y \oplus H^{2k}$, with an adapted splitting over Y. It also splits over J with $q + 1 + 2k$ hyperbolic planes. The top period matching now holds with scalar zero and any rapid upper vector, regardless of its mixed slices. To verify this important assertion, at a fixed mixed argument write the rapid residual real vector as a coherent resolution in the orthogonal product of the two target Hom spaces. Its coefficient is rapidly decreasing, and each product coherent vector is a tensor product of two rapid coherent vectors. The finite name is likewise a product after the block comparison. The integral over $[U(Y)]$ of each product theta factors into the lower theta times the equal-rank integral of the second coherent vector. The latter is zero by the hypothesis just established. Interchanging this integral with the coherent resolution is legitimate: the square name on the equal-rank factor has right-invertible support and the kernel estimates from Section 38.3 bound its target integral by a weighted norm, while the coherent translates have polynomial growth in each such norm.

The product coefficient in the coherent resolution decreases faster than every power. Thus the top matching hypothesis is satisfied with $\rho = 1$, $\lambda_{\mathfrak{c}} = 0$, lower real datum zero, and any rapid upper real vector.

The sign condition is the one in Section 38.10: the target $J^- \oplus Y$ has $\dim J + q(Y)$ negative signs, the lower J^- is totally negative, and Y has both signs with $q(Y) \le \dim Y/2$. Consequently the top standard-column identity asserts that the upper standard lift is zero for *every* rapid upper real vector. Descend the identity to source J and large target Z by removing $q+1+2k$ planes. The rank identity at each stage is $\dim(\text{source}) = 2(k+q(\text{source}))$; the weights are $k + j_\tau$ with $j_\tau = \mu_\tau - k \ge 0$. The descent of Section 38.9 is surjective on that determinant-vector family. In particular select an upper determinant vector whose image is the vacuum on $\mathrm{Hom}(J, Z)$. At the final rank-zero target the named function on the singleton vector is the constant 1 and the normalized theta of the vacuum is 1. The target quotient is the trivial group of mass one (in any positive Haar normalization the value would still be nonzero). Thus the descended zero identity on that vacuum is impossible. This proves the existence of one rapid v_1 and one tame character $\mathfrak{c}_1$ with $\mathcal{R}_Y(\mathfrak{c}_1, v_1) > 0$.

We explain the conversion to partner identities before the second part of the argument. Given complementary ranks $\dim U + \dim V = \dim W_0$ with $\dim W_0$ even and an adapted split of Y over W_0 after accounting for the signs, take the top exchange using the equal-rank vector v_1. Interpolate the prescribed small real vector in the opposite slicing by Section 38.4, and then remove the planes as in Section 38.9; this recovers the prescribed real vector on the small side by the composition of zero slices. The small lift initially uses the flipped target space U^-. In the normalized coordinates the adjoint identification is

$$M \longmapsto (1, c\,\alpha/\bar{\alpha}) J_{W_0} M^* J_U, \qquad \bar{c}c = -1$$

for the fixed finite integral unit c. It preserves the real Euclidean norm and the finite vacuum modulus. Its action on the Hom symplectic pairing changes the conjugate coordinates and compensates the reversal of the target sign. Under conjugation the standard action $M \mapsto hMg^{-1}$ becomes the swapped action, with the determinant multiplier $\upsilon_1(\det g)^{\dim W_0} \upsilon_1(\det h)^{-\dim U}$: this follows by taking the determinant of the action on Hom and using the same negative-coordinate vacuum determinant in the two polarizations. The rational Hom lattices map bijectively by the corresponding rational adjoint $M \mapsto J_U^{-1} M^* J_{W_0}$ after the normalizing matrices. Thus the theta functionals differ by a unit scalar determined on the vacuum. The Haar measures for the opposite and original Hermitian group agree (the isometry on normalized coordinates is the identity on the matrix group with J replaced by $-J$). At the selected finite places the transpose of the name is multiplied on the right by the scalar c; the partner determinant law contributes $\chi(c)^{\dim U}$, again a unit. These are the changes in `exists_pairTheta_flip` and `exchange_of_partners`; the real map preserves rapidity and the finite scalar leaves the squared modulus unchanged. The finite synthesis and stable-space arguments of Sections 33.2 and 35.5 and the promotion theorem of this continuation give the exchange for every locally constant partner, rather than just standard columns.

For the second exchange take $W_0 = J$, first $U = Z, V = J$ and then $U = J, V = Z$. The signed negative counts add as required, and the splitting over J is the adapted one above. Choose as the first small real vector the vacuum and use its resulting large real datum v_2 as the prescribed real datum of the second exchange. There are universal constants B, C and character-dependent scalars $b_{\mathfrak{c}}, c_{\mathfrak{c}}$ with

$$B|b_{\mathfrak{c}}|^2 = \Pi_{\mathfrak{c}}(k) \mathcal{R}_Y(\mathfrak{c}, v_1), \qquad C|c_{\mathfrak{c}}|^2 = \Pi_{\mathfrak{c}}(-k) \mathcal{R}_Y(\mathfrak{c}, v_1).$$

The associated real data do not depend on $\mathfrak{c}$. In the first exchange use the standard partner of the singleton rank-zero name 1. With $\gamma = \mathrm{rankTwist}_{2k}(\nu^{-1})$, the right-hand side is $b_{\mathfrak{c}} \gamma(\det g)$:

the vector space $\mathrm{Hom}(Z, J)$ is zero but its pair character is $\gamma(\det g)$, which must be retained. Dividing out the character on the left gives

$$\int_{[J]} \theta_{\gamma,\gamma}(v_2; g, h)\, dh = b_{\mathfrak{c}}.$$

In the second exchange the large datum on $\mathrm{Hom}(J, Z)$ has a fixed evaluation ξ at the zero vector, independent of the character. Use the partners in the reversed order. Its identity is

$$\xi = c_{\mathfrak{c}} \int_{[J]} \theta_{\gamma,\gamma}(v_2; h, g)\, dh.$$

Integrate this over $[J]$ and swap the two integrals. Here J is definite, its adelic quotient compact, and the continuous theta kernel is integrable on the product quotient; equivalently the constant functions 1 are rapid in both factors in the transposed-lift formula. The preceding identity yields $\xi \operatorname{vol}([J]) = c_{\mathfrak{c}} b_{\mathfrak{c}} \operatorname{vol}([J])$. Cancel this positive finite volume. Taking absolute squares in the resulting equality $\xi = c_{\mathfrak{c}} b_{\mathfrak{c}}$ and using the signed Π relation $\Pi_{\mathfrak{c}}(k)\Pi_{\mathfrak{c}}(-k) = 1$ proves

$$\mathcal{R}_Y(\mathfrak{c}, v_1)^2 = BC|\xi|^2. \tag{5}$$

This is the formula `F` in `exists_equalRankThetaRatio_value`. It supplies the previously missing actual transposition of periods in (33.5). Because the left-hand side is nonnegative and is positive for $\mathfrak{c}_1$, the ratio is one and the same positive number for all $\mathfrak{c}$. In particular neither universal exchange constant can vanish in this construction. Transporting the real equal-rank vector back to $H \oplus W$ by the unitary frame change above proves the claimed universal equal-rank ratio.

Finally for the desired complementary pair in `exchange_theorem`, use this positive ratio in the exchanged partner identity, with the ranks and weight inequalities stated there. Then $e = \dim V - \dim W/2 \geq 0$ and the signed product $\Pi_{\mathfrak{c}}(e)$ is the ordinary $lValueProd$. The resulting exchange scalar has absolute square equal to a fixed positive number times this Hecke product, and in particular is nonzero for every tame character. For residue-modulus data lift each finite function by reduction to the local integral ring, extending it by zero on nonintegral matrices. Reduction respects determinant, multiplication, and the two standard partner block embeddings; by surjectivity of reduction the partner equality on the residue ring gives the partner equality of its lifted functions on the integral matrices. These functions are locally constant and compactly supported, so the already-established arbitrary-partner identity applies. This is the modulus passage at the end of `exchange_theorem`; no constancy of a nonintegral local determinant character is asserted.

38.12 Exact-source status and the remaining tower distinction

1. The finite frame calculation in Sections 38.1, 38.7 supplies the adapted Hom normalization in `AdaptedHom.lean`, `ReadsAt.sourceTerm_localDatum` and the normalized branches of `exists_lineTerm_localDatum`. Together with Sections 36.8–36.10, this completes the outstanding adapted-frame qualification for C23.

2. Sections 38.2–38.3 supply the actual mixed terms and the coefficient extraction of `thetaTerm_stdDatum_spec`; their orthogonal-target product gives `thetaOnRealData_boxProd_eq_mul`. Sections 38.4–38.6 supply `exists_section`, `exists_rapid_with_realSlice`, `exists_contDiff_witt` and the actual real matching, including its opposite zero normalization. The finite and archimedean multiplier computation and stabilizer measure in Section 38.8, with the signature alternative in Section 38.10, establish the period comparison hypothesis

and scalar in `stdExchange_of_periods`. Applying Theorem 37.2 of this continuation then proves the standard-column conclusion and `exists_stdExchange_top`.

3. Section 38.9 supplies the zero/nonzero small-side orbit argument and the telescoping scalar for `exists_stdExchange_removing_planes`, including determinant-vector surjectivity. Section 38.11 proves the nonvanishing and character-independence mechanism of `exists_equalRankThetaRatio_value`, the fixed-data partner exchange and its scalar. Theorem 37.2 and the stable-space argument are essential dependencies: the nonsingular coefficient identity alone does not imply C04 or C05. With these dependencies the complementary-rank exchange and equal-rank top normalization are supplied; the distinct real-convolution edge hypotheses of C06 are not a consequence of this chapter.

4. For C25, the actual normalized tower descent on exchanged standard square/column data is now supplied, rather than merely a conditional transition quotient. This is not yet the full assertion for subtracted theta lifts on a noncompact tested quotient: vanishing of every required rational parabolic constant term, bounds for the resulting subtracted form and conversion to the normalized geometric tested period are separate assertions. They must be proved or cited on those particular data, not deduced simply from the rank-one line calculation.

Chapter 39

Further carrier estimates and the rank-two period route

This section supplies several of the quantitative analytic steps of the compact source that are useful both for the starting-period problem and for comparing the two carriers. They are kept separate from the arithmetic assertions about integral lattices: an analytic estimate, however strong, cannot replace a lattice-integrality assertion. We use the operator norm at complex places and the Euclidean norm on the real underlying vector spaces. All Haar measures in a ratio of integrals are the same Haar measures.

39.1 Holomorphic maximum away from the thin part

Here is a direct proof of the estimate in `Lib/DiscreteSeries/Holomorphic/NearMaximum.lean` (in particular `exists_near_max`). Write $J = \mathrm{diag}(1_p, -1_q)$ at a complex place and order the positive coordinates first. Suppose that the level contains the principal level of a purely imaginary CM integer ς, with $0 < \Im\tau(\varsigma) \le c$ at the chosen complex places. Let the integral Hermitian lattice at g be denoted by L_g. Assume $B \ge 2$, $c \ge 1$, $|u_{\tau,i}| \le B$, and $u_{\tau,j} \ge u_{\tau,i} + 1$ whenever i is positive and j is negative. Let F be a nonzero, bounded automorphic holomorphic form of right weight u, satisfying these level conditions. Choose z so that

$$|F(z)| > \tfrac{1}{2}\|F\|_\infty.$$

Then any g with $|F(g)| \ge |F(z)|$ has

$$\lambda_1(L_g) \ge (Bc)^{-1/2}. \tag{C.1}$$

The Euclidean norm here is the sum over complex places of the squared Euclidean norms in the coordinates determined by g.

Indeed, if v were shorter, its Hermitian norm would be an algebraic integer in the totally real field with every real conjugate of absolute value less than one. Its field norm must therefore be zero. Thus v is isotropic. Put $Y_\tau = g_\tau^{-1} v_\tau$ and $a_\tau = \Im\tau(\varsigma)\|Y_\tau\|^2$. We have

$$0 < a_\tau < 1/B, \qquad B\sum_\tau a_\tau < 1.$$

In the upper half-plane one may vary the archimedean component of g by

$$M_\tau(\zeta) = 1 - \tau(\varsigma)(\zeta - i) Y_\tau Y_\tau^* J.$$

The square of $Y_\tau Y_\tau^* J$ vanishes by isotropy. Set $y = 1 + a_\tau \Im\zeta$, and let ϵ_j denote the sign of the jth coordinate. With

$$P_i = 1 - 2\Im\tau(\varsigma)(1 - \Im\zeta) \sum_{j<i} \epsilon_j |Y_{\tau,j}|^2$$

we have $P_1 = P_{p+q+1} = 1$, $P_{p+1} = y - a_\tau$, and all P_i are positive. In fact the partial sums in this expression lie between zero and $\|Y_\tau\|^2/2$. The leading principal determinants of $M_\tau^* J M_\tau$ are $\det(J_{[i]})P_{i+1}$, by the rank-one determinant formula. Indefinite Gram–Schmidt consequently gives a factorization $M_\tau = h_\tau b_\tau$, with $h_\tau^* J h_\tau = J$ and b_τ upper triangular, of diagonal entries $(P_{i+1}/P_i)^{1/2}$. The holomorphic extension is defined at these points, and its weight multiplier is $\chi_u(b_\tau) = \prod_i b_{\tau,ii}^{u_{\tau,i}}$.

Write $L_i = \log P_{i+1} - \log P_i$. When $\Im\zeta \geq 1$, the L_i in the positive coordinates are nonnegative, in the negative coordinates nonpositive, and their sums on the two groups are $\log(y - a_\tau)$ and $-\log(y - a_\tau)$. The gap assumption gives

$$\log|\chi_u(b_\tau)| = \tfrac{1}{2} \sum_i u_{\tau,i} L_i \leq -\tfrac{1}{2} \log(y - a_\tau).$$

When $\Im\zeta \leq 1$, the signs reverse and the bound on the weights gives $\log|\chi_u(b_\tau)| \leq -B \log(y - a_\tau)$. Because $\log y - \log(y - a_\tau) \leq 2a_\tau$ for $a_\tau < 1/2$, either estimate is bounded by $2Ba_\tau - \frac{1}{2}\log y$. In the second case use also $y \geq 1$ and $B \geq 1/2$.

Let $\phi(\zeta)$ be the holomorphic restriction of F to the resulting product of triangular domains, so that $\phi(i) = F(g)$. The estimates give

$$|\phi(\zeta)| \leq e^2 \|F\|_\infty \prod_\tau (1 + a_\tau \Im\zeta)^{-1/2}.$$

The rational transvection $1 - \varsigma v v^* J$ preserves the Hermitian form and has finite component in the prescribed principal level, in the coordinates determined by g. Its archimedean action is $M_\tau(\zeta) \mapsto M_\tau(\zeta + 1)$ (the square-zero identity eliminates the cross-term). Automorphy and level invariance show $\phi(\zeta + 1) = \phi(\zeta)$. The resulting function of $q = \exp(2\pi i \zeta)$ extends holomorphically to the disc with value zero at the origin, since $\phi(\zeta) \to 0$ towards the cusp. Schwarz's lemma therefore bounds $|\phi(i)| \leq e^{2-2\pi}\|F\|_\infty$, which is strictly below $\|F\|_\infty/2$, a contradiction. No estimate on the derivatives or on the number of connected components of the adelic quotient was used. What is still required in applying (C.1) is the actual level inclusion, the weight gap, and a bound uniform over the relevant carriers.

39.2 Circle projection and polynomially growing kernels

The following elementary identity gives the useful form of the reproducing identity of `Lib/DiscreteSeries/Holomorphic/HolomorphicMinor.lean`. For $h \in U(p, q)$ put

$$m_u(h) = \det(h)^{u_n} \prod_{i=1}^{n-1} \det(h_{[i]})^{u_i - u_{i+1}}. \tag{C.2}$$

Assume that $u_1 \geq \cdots \geq u_p$ and $u_{p+1} \geq \cdots \geq u_n$. A power with negative exponent at the sign boundary is allowed: the upper-left block of a unitary matrix is invertible, as follows from $A^* A = 1 + C^* C$. All other proper principal minor exponents are nonnegative. Formula (C.2) consequently defines a holomorphic function on the open triangular domain used for the holomorphic weight. Right multiplication by an upper triangular matrix multiplies it by its diagonal character.

Choose strictly increasing integer numbers $o_1, \dots, o_n$ and put $D(s) = \mathrm{diag}(s^{o_1}, \dots, s^{o_n})$, for $|s| = 1$. For any holomorphic function H of right weight u, its mean satisfies

$$\int_{|s|=1} H(D(s)h)\chi_u(D(s))^{-1}\, ds = H(1)m_u(h). \tag{C.3}$$

We normalize ds to have mass one. To prove this first take h sufficiently near 1 and use its lower-unipotent times upper-triangular factorization $h = \ell b$. Every entry of $D(s)\ell D(s)^{-1} - 1$ is a positive power of s times an entry of $\ell - 1$. It gives a holomorphic curve on the disc, lying in the open triangular domain for all $|s| \le 1$ if h is sufficiently near 1, with value 1 at zero. Right-weight covariance and the mean-value formula give $H(1)\chi_u(b)$. The leading minors in the same factorization give $\chi_u(b) = m_u(h)$. Both sides are real analytic in h, including the mean (locally uniform analytic extension over the compact circle). Connectedness of $U(p, q)$ extends the identity to all h. The argument applies at each archimedean place, with product probability measure. Multiplication by $\overline{m_u(h)}$ and integration, using left invariance and $m_u(D(s)h) = \chi_u(D(s))m_u(h)$, yield

$$\int H(h)\overline{m_u(h)}\, dh = H(1) \int |m_u(h)|^2\, dh \tag{C.4}$$

whenever the absolute integrals exist. In particular it is not necessary to expand H in a Hilbert-space orthonormal basis.

We give explicit sufficient integrability bounds in the notation of `New/Rows/SparseThetaSum.lean`, `integral_mul_holMinorsPi_le`. Let f be the number of complex places, n the Hermitian dimension, $B \ge 10^{13}$, $2fn^3 \le m \le B$, and let the two weights w, k have a gap at the positive/negative boundary of at least $300m$ and be dominant within each sign. Suppose $|k_{\tau,i}| \le B$. At one place one has

$$|m_w(h)| \le (2/\|h\|)^c \quad (0 \le c \le 300m). \tag{C.5}$$

In fact, for $i \le p$, the norm of the ith exterior power of A bounds the ith leading minor by $|\det A|$, because the singular values of A are at least one. For $i > p$, the complementary-minor identity and $h^{-1} = Jh^*J$ give the same bound through D; $|\det D| = |\det A|$. The signed sum of the minor exponents is at most $-300m$, so the resulting bound is $|\det A|^{-300m}$. The Cartan hyperbolic blocks of h give $|\det A| \ge \max(1, \|h\|/2)$. If one sign is empty, $\|h\| = 1$ and the nonnegative exponents give a modulus at most one directly. These prove (C.5) in all cases.

For k the same reasoning applies. If $\|h - 1\| \le \delta = 1/(100nB)$, every principal determinant has log absolute value bounded by $2n\delta$. Within each sign the sum of consecutive absolute differences of the weights is at most $2B$, and the boundary contributes at most $2B$. Since $|\det h| = 1$,

$$|m_k(h)| \ge \exp(-12nB\delta) \ge 1/2.$$

Let $I_k = \int |m_k|^2$. An elementary ambient packing argument gives the Haar volume bound

$$\mathrm{vol}\{h : \|h\| \le R\} \le 4I_k(201nB)^{2n^2} R^{4n^2} \qquad (R \ge 1). \tag{C.6}$$

Indeed take a maximal δ/R-separated net in the operator norm ball of radius R in the real ambient matrix space of dimension $2n^2$. Packing ambient norm balls bounds its cardinality by $(1 + 2R^2/\delta)^{2n^2}$. If h and a net centre h_0 differ in norm by at most δ/R, then $\|hh_0^{-1} - 1\| \le \delta$, since $\|h_0^{-1}\| = \|h_0\|$. Right invariance and the preceding lower bound show that each ball has measure at most $4I_k$. To avoid circularity, first use the finite measure of the fixed δ-ball in place of $4I_k$: (C.5), packing, and a layer integral show $I_k < \infty$; the local lower bound shows $I_k > 0$ and permits (C.6).

With $c = 10n^3$ we have $c > 4n^2$; integrating (C.6) by layers gives

$$\int (2/\|h\|)^c \, dh \le 4I_k (201nB)^{2n^2} 2^c \, \frac{c}{c-4n^2} \le 8I_k (201nB)^{2n^2} 2^c.$$

Now suppose $|A(h)| \le C \max_\tau \|h_\tau\|^N$ with $0 \le N \le 6m$. At a place attaining the maximum use exponent $c+N$ in (C.5); elsewhere use c. This choice is legitimate since $c+N \le 11m \le 300m$. The product of the last local estimates gives

$$\int |A(h) m_w(h)| \, dh \le C B^{7m} \int |m_k(h)|^2 \, dh. \tag{C.7}$$

For clarity the factor before the integral obtained in this proof is $2^{6m+10fn^3} 8^f (201nB)^{2fn^2}$: this is below B^{7m} since $2fn^2 \le m$, $f \le m/2$, $n \le B$, $201 \le B$, and $B \ge 10^{13}$. The two sides always use the same product Haar measure.

Lemma 39.1 (Quantitative isolation)**.** *Let F be a bounded nonzero holomorphic form of right weight k, and choose z with $2|F(z)| \ge \|F\|_\infty$. Under the preceding weight and dimension assumptions, suppose that a bounded function K_0 is given and that for each x with $|F(x)| \ge |F(z)|$ there is a holomorphic K_x of right weight $k-w$ with*

$$|K_x(h) - K_0(h)| \le \varepsilon \max_\tau \|h_\tau\|^{6m}, \qquad \int \overline{F(xh)} K_x(h) m_w(h) \, dh = 0.$$

Then $|K_0(1)| \le (4B^{7m}+1)\varepsilon$.

Proof. Fix $K_* = K_z$. For $x = zD(s)$ the value $|F(x)|$ equals $|F(z)|$. Write $W_s(h) = \chi_k(D(s)) \overline{F(zD(s)h)}$. The vanishing assumption gives

$$\int W_s(h) K_*(h) m_w(h) \, dh = \int W_s(h) (K_*(h) - K_x(h)) m_w(h) \, dh.$$

By (C.7), the modulus is at most $4\varepsilon |F(z)| B^{7m} I_k$. Average the left side over the rotation group. The fixed K_* has at most polynomial growth by hypothesis, so (C.7) justifies Fubini. A measurable choice of K_x is not needed. Identity (C.3) gives the average $\overline{F(z)} \int K_* m_w \overline{m_k}$, while (C.4) identifies the last integral with $K_*(1) I_k$. Divide by $|F(z)| I_k > 0$ and use $|K_*(1) - K_0(1)| \le \varepsilon$. □

This proves the analytic implication called `isolation` in the compact source. In the application the actual forms K_x must still be exhibited with the stated error bound and the vanishing must be established from the arithmetic grid pairings.

39.3 A sparse coefficient estimate in the Fock model

We give a proof with explicit polynomial constants of the estimate used in `New/Weil/FockCoeffDecay.lean`. On an underlying real space of dimension $2r$ use the Fock model with mixed holomorphic coordinates and let

$$v(x) = P(x) e^{-\|x\|^2/2}, \qquad \deg P = d$$

with P homogeneous in its mixed coordinates. Let $S \ge 1$ bound the operator norm of a pseudo-unitary map σ, and $A \ge d + r$, $A > 0$. In the projection formula for the oscillator action, the linear scalar multiplier has absolute value at most S^r, by the determinant bound on the two diagonal blocks. The projection can be removed when pairing with a Fock vector, including its

Heisenberg translates: the projection is orthogonal and the Fock space is Heisenberg invariant. Consequently the absolute coefficient is bounded by

$$S^r \int |v(x)v(\sigma^{-1}(x-y))|\,dx. \tag{C.8}$$

Here the Heisenberg translation phase has modulus one. This is the coefficient formula of `Lib/Weil/Fock/FockAction.lean` expressed with Lebesgue measure; it assumes the oscillator and Heisenberg actions and the identification of the model, not a summation formula.

Put $u=\sigma^{-1}(x-y)$, let l be a positive integer, and let $a^2=1+l/A$. Homogeneity and the arithmetic-geometric mean give

$$|v(x)v(u)| \le \frac{a^{2d}}{2}\bigl(|v(x/a)|^2+|v(u/a)|^2\bigr)e^{-l(\|x\|^2+\|u\|^2)/(2(A+l))}.$$

Since $y=x-\sigma u$, $\|x\|^2+\|u\|^2 \ge \|y\|^2/(2S^2)$. The maximum of $t^l e^{-lt}$ is e^{-l}, so the exponential is bounded by $(4(A+l)S^2/(e\|y\|^2))^l$. Both integrals in the other factor have value $a^{2r}\|v\|^2$ (pseudo-unitary maps have real Jacobian of absolute value one). As $a^{2(d+r)} \le e^l$, (C.8) proves, for $y\neq 0$,

$$|\langle v,\rho(y)\omega(\sigma)v\rangle| \le S^{r+2l}\bigl(4(A+l)\bigr)^l\|y\|^{-2l}\|v\|^2. \tag{C.9}$$

The zero-power estimate follows by unitarity. The same proof for all real places at once uses the total real dimension $2r$ and a common bound S, with a product scalar multiplier bounded by S^r.

For reference a packing estimate turns (C.9) into a completely explicit sparse sum. If a set in $\mathbb{R}^{2r}$ has pairwise distance at least $M>0$ and every nonzero member has norm at least M, then disjoint radius-$M/2$ balls bound the count of nonzero members of norm at most R, $R\ge M$, by $(3R/M)^{2r}$. For an integer $l>r$, the layer formula gives

$$\sum_{y\neq 0}\|y\|^{-2l} \le \frac{3^{2r}l}{l-r}M^{-2l} \le (5/M)^{2l}. \tag{C.10}$$

The last inequality follows from $l/(l-r)\le l$ and $9^r l \le 25^l$. In (C.9) take $A=d+r$ (the zero-dimensional case is the constant representation) and $l>r$. Combining the bounds therefore gives the nonzero-mode error estimate

$$\sum_{y\neq 0}|\langle v,\rho(y)\omega(\sigma)v\rangle| \le \left(\frac{100(d+2l)}{M^2}\right)^l S^{3l}\|v\|^2. \tag{C.11}$$

For its application to `New/Rows/SparseThetaSum.lean`, `norm_modKernel_sub_realInner_le`, the finite trace-character average deletes precisely the modes with nonzero residue, not merely a proportion of them. More explicitly, for an additive finite module T paired perfectly with its character module by the trace pairing, $\mathbb{E}_{\xi\in T^\vee}\xi(t)$ is one at $t=0$ and zero otherwise: for $t\neq 0$ multiply the sum by a character whose value at t is not one. Apply this to the finite translated vacuum coefficient in the theta-product identity. Both factors use the same trace character; translation covariance multiplies the coefficient at a rational Heisenberg displacement by its residue character. The finite pairing support then confines surviving rational displacements to $c^{-1}L$ when $c\alpha_{\mathfrak{m}}\widehat{\mathcal{O}}\subset\widehat{\mathcal{O}}$. In the normalized archimedean coordinates the lower bound $M|\tau(c)| \le \sqrt{2\pi\Im\tau(\mathrm{j})}\,\lambda_{\mathrm{Hom}}$ gives minimum and separation at least M. Thus the zero mode is exactly the real matrix coefficient and the sum of the other modes is at most (C.11); when S is the product of the two archimedean operator norms this yields the factor $\|u\|^{3l}\|u'\|^{3l}$ of the source. Absolute convergence for exchanging the theta product and the finite average follows from (C.10)–(C.11), once the theta-product identity and finite support statements are in place. Those algebraic identities are independent of this estimate and must not be inferred from a bound on the nonzero modes.

39.4 Short bases, the sparse lattice threshold, and reproduction

We record the lattice estimate used in the near-maximum argument in `New/Rows/SparseThetaSum.lean`, `firstMinimum_mul_le_hom`. Let S, T be the two spaces of ranks n, n'; their integral Hermitian lattices L_S, L_T are self-dual for the integral Hermitian form. Let there be f complex places and let the positive integer N belong to the different $\mathfrak{d}_{E/\mathbb{Q}}$. The Euclidean trace pairing has $\mathrm{Tr}_{E/\mathbb{Q}} = 2\Re\sum_\tau$, so the real Euclidean dual of $(2/N)L_S$ is the lattice $N\mathfrak{d}_{E/\mathbb{Q}}^{-1}JL_S$ in these coordinates. This follows by testing against all $\mathcal{O}_E$ multiples of an element of L_S and using self-duality for the Hermitian form. In particular the Euclidean shortest dual vector is at least $\lambda_S = \lambda_1(L_S)$. The transference inequality $\mu(L)\lambda_1(L^*) \le d/2$ for a real Euclidean lattice of dimension d (W. Banaszczyk, *New bounds in some transference theorems in the geometry of numbers*, Math. Ann. 296 (1993), 625–635, DOI 10.1007/BF01445125) gives a covering radius for $(2/N)L_S$ of at most d^2/λ_S with $d = 2fn$. The lattice points in the closed ball of radius three times its covering radius span the real space: otherwise take a vector of length twice the covering radius orthogonal to their span, find a nearest lattice point, and use the defining bound on distance to contradict orthogonality. Scaling these spanning vectors back by $N/2$ and selecting an E-basis, we find n integral vectors which form a rational E-basis and have each norm at most

$$6N(fn)^2/\lambda_S.$$

No claim that they form an $\mathcal{O}_E$-basis is needed. If a nonzero rational homomorphism A carries L_S into L_T, some one of these basis vectors has nonzero image. Comparing its length with λ_T and bounding its image by the operator norm yields, in particular, the (weaker) source bound

$$\lambda_{\mathrm{Hom}(S,T)} \;\ge\; \frac{\lambda_S\lambda_T}{6N(fn)^2n'n}. \tag{C.12}$$

The additional factor $n'n$ allows the source's coordinatewise matrix norm bound, and is unnecessary in Hilbert–Schmidt normalization. The argument applies after moving the two spaces by their adelic frames, as these preserve their integral Hermitian duality. It also proves the needed existence of invertible matrices of short integral rows in a definite target lattice.

We next spell out the stable large-rank reproduction mechanism without requiring an irreducible representation decomposition of the whole automorphic space. Suppose $n' > 2n$, the level is principal of generator $N_0 \neq 0$, the modulus of this level has a nonempty set of places, and the chosen finite datum is the indicator of the standard inclusion $E^n \hookrightarrow E^{n'}$. Assume both splitting characters have trivial tame part. The indicator datum is sharp for the indicated level, by the fixed left inverse and the local coefficient identity in Chapter 36.4. Given a holomorphic highest Fock vector $v = p_Q$ of right weight u, assume m_u is integrable. The matrix coefficient in the first group is precisely

$$\langle v, \omega_\infty(h,1)v\rangle = \|v\|^2 m_u(h). \tag{C.13}$$

In fact this coefficient is holomorphic of right weight u and has the same left torus character, using the highest-vector covariance and unitarity; applying (C.3) leaves it unchanged and gives (C.13). This is the needed identity even if the representation generated by v has not been classified.

For each finite class a of the second, definite group at principal level N_0, choose polynomials $Q_{a,l}$ whose Fock vectors span the real translates of v on that class. They can be chosen from the specified finite family: a power of a maximal minor in the positive letters transforms by changing the row vectors; expand the determinant multilinearly in any invertible basis of rows and expand products of determinants distributively. Choose the short rational row

basis just found. These determinant powers are harmonic highest vectors: the noncompact lowering contractions differentiate a variable of each of the two opposite signs and hence annihilate a polynomial in only the positive letters; the compact upper-triangular raising operators annihilate the maximal-column determinant because the new column repeats a selected column. Compact diagonal differentiation gives weight $-d$ on each positive column for a dth power, and the oscillator vacuum contributes weight n' on each negative column. For the staircase case of `letterPoly_highestVector_holomorphic`, use the ordinary Weyl highest vector in the indicated sign block instead of the maximal-column determinant; the opposite-sign contractions still vanish and the compact weights are its consecutive diagram weights added to the vacuum character. The spanning statement is its finite-dimensional compact representation action (the minor basis spans the indicated Weyl module). These computations use precisely the Fock and Weyl representations of the preceding chapters, rather than treating an arbitrary polynomial as harmonic.

Let ψ be rapid of principal level N_0 on the first group. If all the finitely many pairings

$$L_{a,l}(\psi) = \int_{[S]} \psi(g)\, \theta[1_e, Q_{a,l}](g, a)\, dg$$

vanish, then its entire lift with datum $v \otimes 1_e$ vanishes. Indeed every point of the definite target is a rational element times its finite class representative times a principal-level element times a real element. The first is removed by automorphy; the second finite factor fixes the inclusion datum and contributes no tame determinant phase; the real factor translates v into the given span. This proves the assertion for every point of the target.

The clean doubling identity of Theorem 36.2 and the dual-datum cutoff of Lemma 36.3 have a useful consequence here not requiring a real-edge hypothesis. If the lift with $v \otimes 1_e$ vanishes, the cutoff formula gives zero against every rapid test form ψ' on the source. Exhaust the unramified finite places as in (36.3), and apply the sharp finite partial integral (36.4). Since the sharp mass is nonzero, one obtains

$$\int \langle v, \omega_\infty(h, 1)v\rangle \langle \psi', R(h)\psi\rangle\, dh = 0$$

for every rapid ψ'. The convolution belongs to L^2 by the L^1 bound for the coefficient and Minkowski, and compactly supported continuous automorphic test functions are dense in L^2 (they may also be smoothed on each component). Therefore the convolution is zero in L^2, and everywhere by continuity and positivity of measure on nonempty open sets. This is the argument of `realConv_eq_zero_of_lift_eq_zero`; the real-edge scalar used later for the rank-one norm formula is not needed for this implication.

It follows from (C.13) that the linear functional $\psi \mapsto \int \psi(xh) m_u(h)\, dh$ vanishes on the intersection of the kernels of all $L_{a,l}$. It consequently factors through the image of their finite-dimensional joint map. Extend the resulting functional on that image to the entire finite-dimensional coordinate space by extending a basis: for each x fixed there are complex numbers $b_{a,l}(x)$ such that

$$\int \psi(xh) m_u(h)\, dh = \sum_{a,l} b_{a,l}(x) L_{a,l}(\psi). \tag{C.14}$$

This proves the reproduction formula `reproducing_formula_modKernel` with the source level and inclusion datum. It only uses the clean stable-range identity, not an unnormalized assertion that theta lifts are nonzero.

39.5 The carrier comparison with quantitative error

The original carrier is the theta lift of 1 from the compact real group of signature $(0,m)$ to $(2m,m)$ with polynomial $\prod_\tau \det(Y_{\tau,-})^{2m}$ and unit-level vacuum. This carrier is nonzero: for the original first group the real matrix coefficient of this vector is a positive constant. Indeed the substitution of the determinant in (C.8), or the determinant character of the compact group, gives on its negative block exponent $3m-m-2m=0$. Hence the just-proved vanishing implication for the clean datum, whose finite part here is the unit vacuum at level K_0, would imply that convolution of the constant 1 with this positive constant coefficient on the compact real group vanishes. Its integral is strictly positive. The rank hypothesis is $3m>2m$. This gives `carrierForm_ne_zero`. For the other assertion `carrierForm_spec`, the source writes the carrier as the class sum of theta kernels with full-rank square determinant support at nonempty modulus by `stdClassSum_carrierXi_eq_carrierForm`. The support has a left inverse because the determinant is a unit, so the rapid kernel bound of Chapter 10.7 applies. The compact character law gives level one; the constant polynomial has holomorphic weight $(0,3m)$ by the just-computed vacuum weights. Thus its rapidity and holomorphy follow from that class-sum identity and the decay result; they do not follow merely from nonvanishing of the original lift. We retain the class-sum identity as a separate exact exchange input.

We now use the actual dimensions, not asymptotic surrogates. Put $\nu=2ft^3=\mathrm{nStar}$, $B=5b$, $h'=h_2-\lfloor (t+1)/2\rfloor$, $n_0=3m$, and $n_1=r+3m$, the two ranks of the carrier and padded row space. The padded row space has exactly m negative coordinates: the negative coordinates of the original row space and its j negative padding coordinates give precisely this number. The index maps negIndex and posRowIndex(h) are injective and land in their required sign blocks: on the original row block their indices increase with i, and on overflow they lie in distinct padding coordinates, which cannot coincide with an index in the original block. For the negative map the first j coordinates go to negative padding and the remaining indices to the negative row coordinates. These are the elementary checks in `negIndex_posRowIndex_spec`. Both ranks are at most t; the reproduction target $I^\flat$ has rank $6m+1>2n_0$.

For each place τ write

$$N_\tau=|\mathrm{DetIndex}(j,\tau)|,\quad A_\tau=N_\tau+|\mathrm{BigIndex}(j)|+m,\quad p_\tau=\sum_{h<k+j-1}k_\tau^+(h)=N_\tau-h'.$$

Here $|\mathrm{BigIndex}(j)|=r+9m+1$. The polynomial $Q=\mathrm{linePoly}$ of the source is the product of the negative minor to power A_τ and the positive minors to powers $k_\tau^+(h)$. It is homogeneous of degree

$$d_Q=\sum_\tau (mA_\tau+2mp_\tau)\le 10fmb$$

because $2A_\tau\le 5b$ and $p_\tau\le A_\tau$ (the slightly smaller bound $15fmb/2$ would suffice). Each determinant is a nonzero polynomial by injectivity of the selected rows and columns, so $v_Q\neq 0$.

Let F_C denote the fixed carrier and choose x by the near-maximum argument in Section 39.1 with $|F_C(x)|>\|F_C\|_\infty/2$. Its level one permits use of the purely imaginary $\sqrt{-d_0}$; in particular its lattice first minimum is at least $(5bx_1)^{-1/2}$ since $\sqrt{d_0}\le x_1$. For every row point g with $\lambda_g\ge(5b^{24})^{-1/2}$, apply (C.12) to the carrier and the padded row space at (x,g); the additional definite standard summands have minimum at least one. The modulus is $\mathfrak{m}_0=(\overline{\pi}_G^2)$ and take $c=N_1/\overline{\pi}_G^2$. Its absolute value at each complex place is N_1/G, where $G=g_0g_1$ and $\pi_G\overline{\pi}_G=G$. The finite ideal condition $c\,\alpha\mathfrak{m}_0\widehat{\mathcal{O}}\subset\widehat{\mathcal{O}}$ is exactly $N_1\alpha\widehat{\mathcal{O}}\subset\widehat{\mathcal{O}}$ from the Weil normalization in `WeilDataOfE.lean`. The imaginary real normalizing scalar is $\mathfrak{j}=-1/\sqrt{-d_0}$. With $M=b^7x$ the sparse lattice condition in (C.11) is satisfied. For an explicit coarse

check, $N_1, d_0, f, t, x_1, x < b^{1/100}$ follows from the scale inequalities of `Magnitudes.lean`; also $5, 6 < b^{1/100}$ since $\log\log b \geq 10^{12}$*P.c.* Hence $6(N_1 d_0)(f n_0)^2 n_1 n_0 < b^{9/100}$ and $\lambda_S \lambda_T > b^{-12.515}$. The square root $\sqrt{2\pi \Im \tau(\mathrm{j})}$ is at least $b^{-0.0025}$. Thus the right side of the sparse condition is at least $b^{-12.6075}$, whereas $MN_1/G \leq b^{-12.98}$: the actual prime bounds are $b^{20} \leq G \leq b^{22}$. The exponent in (C.11) is chosen to be $l = 2\nu > f n_0 n_1$, giving

$$\delta_0 = \left(\frac{100(d_Q + 4\nu)}{b^{14} x^2}\right)^{2\nu} \leq b^{-24\nu}. \tag{C.15}$$

Indeed $100(10fmb + 4\nu) \leq b^2$ under the same scale bounds. For example the bounds $f, m, t \leq b^{1/100}$ give $10fmb + 8ft^3 \leq 10b^{1.02} + 8b^{0.04}$, and $b \geq 10^{13}$ suffices for this last numerical inequality. The sparse lemma therefore supplies a name Ψ modulo $\mathfrak{m}_0$ such that at all real u, y its theta kernel approximates

$$C(u, \iota y) = \langle v_Q, \omega_\infty(u, \iota y) v_Q \rangle,$$

where ι pads y by the two identity blocks, with error at most $\delta_0 \, \|v_Q\|^2 \max \|u_\tau\|^{6\nu} \max \|y_\tau\|^{6\nu}$. This uses the exact theta product (9.9); the finite coefficient covariance follows by translating the named vacuum by the opposite-coordinate finite Heisenberg shift representing the trace character. The Heisenberg commutator multiplies the coefficient by the character of the displacement, and the remaining coefficient is the ordinary vacuum support. Thus there is no omitted normalization in the zero mode.

Take the weight $\varsigma = (-h') \oplus (6m + 1)$ on the carrier and define

$$K_0(y) = \int F_C(xu) \, m_\varsigma(u) C(u, \iota y) \, du. \tag{C.16}$$

It is bounded, by unitarity of the oscillator action, (C.7), and boundedness of F_C. The minor formula for the real coefficient at $y = 1$ can be verified directly by the mixed Fock projection: substituting u^{-1} into each maximal positive or negative column determinant multiplies that determinant by the corresponding full principal block minor. Harmonicity eliminates the opposite-sign contractions in the Fock projection; the Gaussian determinant multiplier accounts for the negative block rank and the character weights. This is `realInner_realPairWeil_eq_holMinorsPi`. More explicitly, if $e_\tau = m - N_\tau$ is the weight of the character carrierChar, it gives

$$\frac{C(u, 1)}{\|v_Q\|^2} = m_v(u) \overline{m_k(u)},$$
$$v_+ = -p_\tau - A_\tau - e_\tau, \qquad v_- = n_1 - m - A_\tau - e_\tau, \quad k_+ = -(A_\tau + m), \quad k_- = 0.$$

In particular the sum of the weights of F_C, m_ς and m_v is exactly k: on the positive block it is $-h' - p_\tau - A_\tau - e_\tau = -A_\tau - m$, and on the negative block it is $3m + (6m+1) + n_1 - m - A_\tau - e_\tau = 0$. The bounds $2A_\tau \leq 5b$ and $300\nu + t \leq h_2$ imply the gap and weight assumptions of (C.7) for ς, k, with $n = n_0$ and m there replaced by ν. Identity (C.4) now gives the exact nonzero value

$$K_0(1) = F_C(x) \|v_Q\|^2 I_k, \qquad I_k = \int |m_k|^2 > 0. \tag{C.17}$$

In (C.14) use this carrier weight ς, target $I^\flat$, and $Q_0 = \prod_\tau \det(Y_{\tau,+})^{h'}$. Its positive-block minor powers have translates spanned by the paired determinant products in the rows of shortRows, by multilinearity as above. For any fixed g, y the rapid source function to which (C.14) is applied is F_C times the Ψ, Q kernel between the carrier and the padded row space: it is rapid because F_C is rapid, the kernel has moderate growth, and the product has the

required principal level (the other finite group action is fixed). Expand the arbitrary name Ψ times the inclusion-indicator name on the extra $I^\flat$ block in the finite trace-character basis of the enlarged matrix module modulo $\mathfrak{m}_0$. The trace pairing is perfect, so the basis expansion is exact by finite Fourier inversion. The product kernel identity for the orthogonal block sum makes the resulting integrals exactly the $F_a^\natural$ of `LineCoordinates.lean`; the polynomial is the product of the embedded Q and of the paired determinant products. Thus there are coefficients $c_a(g)$, independent of y, for which

$$\left|\sum_a c_a(g)F_a^\natural(gy) - K_0(y)\right| \le \varepsilon \max_\tau \|y_\tau\|^{6\nu}, \quad \varepsilon = 2B^{7\nu}\delta_0\,|F_C(x)|\|v_Q\|^2 I_k. \tag{C.18}$$

Here (C.7) was applied to the error in the sparse approximation multiplied by $F_C(xu)$, using $\|F_C\|_\infty \le 2|F_C(x)|$. The coefficient basis and the carrier point are fixed, and the sparse name is chosen at g only; no dependence on archimedean y was introduced. By (C.17) and (C.15),

$$\frac{(4B^{7\nu}+1)\varepsilon}{|K_0(1)|} \le (4B^{7\nu}+1)\,2B^{7\nu}b^{-24\nu} < 1.$$

For instance $B = 5b$ makes the right side at most $10(5^{14}b^{-10})^\nu < 1$. This proves the quantitative approximation asserted in `exists_fNatural_approx`, conditional on the precise carrier class-sum identity and its rapid consequence noted above. In particular the number of classes or the size of the Fourier-expansion coefficients does not enter the analytic approximation error.

The last analytic deduction, `eq_zero_of_integral_fNatural_tableTheta`, can now be made without assuming separation. For a good row form F the conjugate is holomorphic of weight $k =$ conjHolWeight by its prescribed theta lift and staircase highest vector; $F^\natural$ has weight $k - w$, $w =$ kernelWeight, by its expression through the matched theta kernel and its positive-letter highest vector. Products $FF^\natural$ are rapid at principal $\overline{\pi}_G^2$. These are the representation and level claims in `Separation.lean`; in particular the latter uses the exact identity `cosetKernelSum_eq_fNatural`, not an arbitrary comparison form. If all the pairings with $F_a^\natural$ and the tables vanish, assume $F \neq 0$ and use (C.1) on $\overline{F}$ at the imaginary level generator $G\sqrt{-d_0}$ (integral and divisible by $\overline{\pi}_G$ at the chosen level). The size bound is precisely $G\sqrt{d_0} \le b^{23}$, since $G \le b^{22}$ and $d_0 < b^{1/100}$. The first minimum at these near-maximum points is therefore at least $(5b^{24})^{-1/2}$, the threshold needed for (C.18). At any row point satisfying the resulting lower threshold of (C.18), take the comparison combination K_x there. Apply (C.14) on the row space to the rapid form FK_x, with the negative-letter highest vector and the whole Weyl basis in the definite group of rank $t - 9m - 1 > 2r$. Each pairing on the right of (C.14) is zero: choose the table vector supported on precisely one target class and one Weyl basis vector, then take the complex conjugate of the hypothesized table pairing. Thus $\int F(xh)K_x(h)m_w(h)\,dh = 0$. The exact hypothesis of Lemma 39.1 for the holomorphic form $\overline{F}$ has been reached. The lower threshold at the near-maximum points has now been verified, and (C.18) contradicts the isolation bound. This is the analytic content of carrier separation; its arithmetic exchange identities remain independent requirements.

39.6 Uniform rank-two jet clearing and the integral family

The interpolation argument for $z = 1$ in Chapter 43.7 used a uniform denominator for the determinant kernel as a separate input. We identify that denominator directly with the standard-data jet assertion and thereby complete this arithmetic step for the actual rank-two period.

Let $p = (\nu_0, z)$, $z = 1$, be a ramified pair, and put $c_\tau = p.c_\tau - 1 \geq 0$, $D_0 = 2\sum_\tau c_\tau$, $G = D.\mathrm{gProd}$ and $\delta = G^2$. Fix the standard definite pair (I_2, I_2), with splitting character $\alpha_\vartheta = \nu_0 \vartheta \upsilon$ of weight $-c_\tau$ on each block. Write $\theta_\vartheta[\Phi, P_0](1, h)$ for its normalized complex theta kernel with

$$P_0 = \prod_\tau \det(X_{\tau,ij})^{c_\tau}.$$

This polynomial has integer coefficients and is homogeneous of total degree D_0. Let $\mathfrak{m} = D.\mathfrak{m}(\vartheta, 2)$ and $q = q_Q = \mathrm{N}(Q)$. The integral ideal condition is $q\delta \in \mathfrak{m}$, as proved in Chapter 36.7. In particular for the datum $\Phi = 1_{\{1\}}$ the Fourier-naming proof of Lemma 36.5, with $r = s = 2$, gives

$$\delta^{8f} \kappa_0^4 \,\mathrm{vacDatum}(\Phi) \in \mathrm{standardData}_\delta$$

at every coefficient embedding above a prime not dividing q. The power four of the vacuum scale is the number of complex matrix entries, not the rank of either group. Both characters have the required weights, so the integral above-prime pair action preserves this standard-data module (including the determinant scalars, which are units there). The jet integrality theorem `norm_jet_jetSeries_le_one` in `ConjugatesOfJets.lean`, proved via the integral CM Taylor coefficients in the arithmetic model, evaluates this module at scale $\overline{\delta}\, d_0$ per, where $d_0 = P.d$ and per $= P.\mathrm{jetScale}$. Consequently the single scalar

$$N_* = \delta^{8f} \kappa_0^4 (\delta d_0 \,\mathrm{per})^{D_0} \neq 0 \tag{C.19}$$

makes the unscaled kernel integral at any such embedding, at every pair of finite points integral above that prime. Here homogeneity moves the jet scale to the scalar multiplier to the power D_0; since δ is rational, $\overline{\delta} = \delta$. The expression depends on p, D and the fixed theta normalization, not on the modulus Q, the character ϑ, the prime, or the point of evaluation. It works simultaneously at all embeddings. This is the integral half of `exists_modKernel_integral_bound` needed in `ValueFamilies.lean`; a conjugate upper bound on N_* is not needed for this family.

For a general finite point h, choose a rational unitary translate making it integral above the chosen coefficient prime (the weak approximation assertion used in the preceding arithmetic chapters). The scalar determinant kernel is unchanged by that translate: the rational covariance factor from the character on the second group is $\prod_\tau \tau(\det \gamma)^{c_\tau}$ and substituting γ^{-1} into the row determinants of P_0 multiplies by its inverse. This proves the needed rational invariance also for the unit-matrix indicator, not just the equivariant full determinant-character datum. It is `detKernel_embFin_mul`. Applying (C.19) at the integral representative proves integrality of the original kernel.

Let $\Lambda_\vartheta = \mathrm{detPeriod}(I_2, \alpha_\vartheta, c, \mathfrak{m}, \psi_0 \vartheta.\psi)$ and M_G be the chosen group order. The expansion `detPeriod_eq_stdClassSum_sum`, proved in Chapter 43.7, is

$$M_G N_* \Lambda_\vartheta = \sum_h \sum_{Y \in \mathrm{GL}_2(\mathcal{O}_E/\mathfrak{m})} \frac{M_G}{|\Gamma_h|} N_* \,\theta_\vartheta[1_{\{1\}}, P_0](1, h\widetilde{Y}), \tag{C.20}$$

where the first index h runs over the full compact level classes. The expression is a sum over the pairs (h, Y), with Y ranging over the displayed finite group for each h. The fixed determinant weight causes no further norm factor. By the finite scalar twist law, each term is the corresponding term for the reference trivial tame twist on the same modulus times ϑ evaluated on $\det(h\widetilde{Y})$ or its inverse, according to the kernel convention. All coefficients are integral away from q by (C.19) and $|\Gamma_h| \mid M_G$. The latter divisibility is the reduction injection of the rational stabilizer into the chosen finite group, not a bound on its order. The dependence of the points used to check integrality on the coefficient prime causes no problem, since the coefficients used in (C.20) are unchanged by the rational translates. A norm square

of a finite evaluation sum has the pairwise products of the coefficients and the inverse product of the group points, so it is integral at those same primes; complex conjugation permutes all the embeddings. Therefore the precise expression

$$F(\vartheta) = |M_G N_* \Lambda_\vartheta|^2 \tag{C.21}$$

is an integral evaluation measure for the actual rank-two family, with no remaining denominator assertion for this step.

The Hecke calculation of Chapter 43.7 proves $\Lambda_{Q\cup\{v\}} = (1-X)(1-q_v X)\Lambda_Q$ directly on the finite determinant datum, where $X = (\alpha_\vartheta.\mathrm{frobValue}(2,v))^{-1} = q_v^{-1}((\nu_0\vartheta)^{\mathrm{bc}}(\varpi_v))^{-1}$. Thus taking the norm square in (C.21) gives the exact Euler insertion law

$$\prod_{k=0,1}\prod_{\epsilon=\pm1}(1-y^\epsilon q_v^{k-1}), \qquad y = (\nu_0\vartheta)^{\mathrm{bc}}(\varpi_v).$$

The only analytic period identity still invoked for this rank-two case is the exact equal-rank matching statement `stdClassSum_det_modKernel_eq`. For c_τ as above and rank two it says

$$\gamma_\vartheta \Lambda_\vartheta = 1, \qquad |\gamma_\vartheta|^2 = \frac{C_0}{\mathrm{lValueProd}_{M\cup Q}((\nu_0\vartheta)^{-1},1)}.$$

The characters in this formula are correct since $\alpha_\vartheta^{-1}\upsilon = (\nu_0\vartheta)^{-1}$. It follows without an arbitrary period-square-root convention that $F(1)\neq 0$ and, by the definition of lValueProd or `Pair.lValueProd_eq`,

$$F(\vartheta)/F(1) = q_Q\,\rho(\vartheta)^2.$$

This completes the reduction to the earlier Galois-family construction in Chapter 43.8: it is the $m=1$, $K=\{0,1\}$ case of `Data.Pair.exists_isIntegralMeasure`. The statement about the period evaluation itself must come from the exact exchange identity, not from the denominator calculation or a formal Euler product at the endpoint 1.

39.7 A height consequence with the exact flat error

We also record the consequence for `heightLe_norm_trivialLift_sq` in `Separation.lean`. This step uses the now-established normalized rank-one conjugate estimates in Chapter 36.6. Denote the rank-one pair by p and the parallel flat pair of the same rank $n = 2z \geq 4$ by q (weight $-z$ and trivial reference character). Their rank-one datum normalizations are those of trivialLift, and the norm estimates, writing $\mathfrak{S}_p = \sum_\tau \mathrm{jetSize}(p.c_\tau - z)$, give upper and lower bounds with errors $\pm 4f$ and the common rank-dependent scale. In the notation for the height of a constant family take the nonzero rescaling factor

$$A = \frac{q.\mathrm{intAlgPart}(1)}{\|\mathrm{trivialLift}_p(1)\|^2}.$$

Then the scaled family is precisely $q.\mathrm{intAlgPart}(1)$, integral at every coefficient prime by Chapter 36.7; any auxiliary modulus in this constant-family calculation is 1. Both the scaled value and the rescaling factor are algebraic under the semilinear norm comparison. The conjugate log bound for A is at most

$$\log M_G + f(8z\log G + 8) - \mathfrak{S}_p.$$

Here the first two terms come from the normalization $M_G G^{4fn}$ of the flat norm, and the two errors $4f$. Finally $p.c_\tau - z \geq D.w(0,\tau)$ for the rank-one parameters by their definition;

the jet size is increasing over this range (the difference on increasing an integer by one is $\log(d+1)+\log(G/\Omega)$, positive under the data bounds). The size inequality $\log M_G + f(8z\log G+8) \le B_0$ is the exact source budget inequality. This yields the desired height $B_0 - \sum_\tau \mathrm{jetSize}(D.w(0,\tau))$ with no additional loss. The argument is independent of the carrier exchange: only the rank-one Rallis normalization, its semilinear transport, and the size inequality enter.

39.8 The deep-level Poincaré restriction tail

For the quantitative starting-row assertion in `RowsReached.lean`, one intermediate estimate can be separated cleanly from the difficult two-factor real period. We give the arithmetic/coset and summation part of `norm_poincare_period_sub_le` (`New/DiscreteSeries/RestrictionLemma.lean`).

Put $Z = V \oplus \langle 1\rangle$, $i : U(V) \hookrightarrow U(Z)$, with e the final positive standard vector. Let $a \neq 0$ be the principal level generator, $K(a)$ the level in each respective group, $v_a = \mathrm{vol}(K_V(a))$, and $R \le |\tau(a)|$ at all complex places. Suppose that F is automorphic of level $K_V(a)$ with $|F| \le S$, and write $\lambda = \lambda_1(V_x)$ at a given point x. For the real coefficient on the large group put

$$\phi(h) = \prod_\tau \overline{\psi_{l'_\tau}(\alpha_\tau^{-1} h_\tau \omega_\tau)}, \qquad \phi_g(u) = \phi(g\, i(u)).$$

Suppose the real gap and formal-degree estimates give the following integral bound (the precise real estimate in the source, with v_τ the volume of the unit ball around the identity of the small group):

$$\begin{aligned} \int_{U(V)_\infty} |\phi_g(u)|\, du \le L t_0^{-A} \quad \text{if } \prod_\tau \|g_\tau i(u_\tau)\| \ge t_0 > 0 \text{ for all } u, \\ L = \prod_\tau (2\|\alpha_\tau\|^2 \|\omega_\tau\|^2)^A 16^A v_\tau. \end{aligned} \tag{C.22}$$

This follows from the real coefficient decay and shifted real ball integral in the source; we make (C.22) an explicit hypothesis here so that the following argument does not silently use a new noncompact real representation estimate.

For a real parameter $D \ge \max(f, 10)$ assume $D^2 \le R\lambda$. Form the Poincaré sum on the large group

$$K(h) = \sum_{\gamma \in U(Z)(E)} \phi((i(x)^{-1}\gamma i(h))_\infty)\, 1_{K_Z(a)}((i(x)^{-1}\gamma i(h))_f).$$

When integrating over $[U(V)]$, group its rational summands by right cosets of $i(U(V)(E))$. Those cosets inject into the rational norm-one vectors γe: two isometries with equal final vector differ on the right by an isometry of the orthogonal complement V. Only cosets with a nonempty finite-level condition contribute. For each such coset choose a point h_γ in the small group satisfying that condition. Since $i(h_\gamma)$ fixes e and $K_Z(a)$ sends e to e modulo a, the rational coset vectors, read in the coordinates of $i(x)$, lie in the translated finite lattice $e + aL_{Z,i(x)}$. Thus at infinity their differences have length at least $R\min(\lambda,1) \ge R\lambda$ if $\lambda \le 1$ (as here, by the standard first-minimum bound). Every nontrivial coset vector is separated by at least D^2 from the identity vector e, whose archimedean Euclidean length is $\sqrt{f} \le D$. Its norm is consequently at least $D^2/2$. Upon unfolding a single coset and changing variables by h_γ, the finite support becomes exactly $K_V(a)$: if $i(x)^{-1}\gamma i(h_\gamma)$ belongs to $K_Z(a)$ then membership at $h_\gamma h'$ is equivalent to $i(h') \in K_Z(a)$, hence $h' \in K_V(a)$. The absolute value of the contribution of this coset is at most

$$v_a L S\, (D/\|\gamma e\|_x)^A\,. \tag{C.23}$$

Indeed the vector is the image of the fixed unit vector by $i(x)^{-1}\gamma i(h_\gamma)i(u)$ for every real u. The product of its local matrix operator norms is at least $\|\gamma e\|_x/\sqrt{f} \geq \|\gamma e\|_x/D$, since every local operator norm is at least one. Apply (C.22) at this lower bound and use $|F| \leq S$.

For completeness the sum over nontrivial cosets is controlled in the ambient real dimension $d = 2f(n+1)$. The packing bound for a D^2-separated set inside radius $r \geq D^2/2$ is $(1+2r/D^2)^d \leq (4r/D^2)^d$. The layer integral for $A \geq d+2$ yields

$$\sum_{\gamma \neq 1} (D/\|\gamma e\|_x)^A \leq \frac{A}{A-d} 2^{A+d} D^{-A} \leq 10/D.$$

For the last inequality use $d \leq A-2$ and $A/(A-d) \leq A/2$: the upper bound is no greater than $A(4/D)^A/8$, which multiplied by D is at most $10A(0.4)^A/8 \leq 10$ for $D \geq 10$, $A \geq 2$. This proves absolute convergence of the integrals of the nonidentity cosets, justifies their unfolding, and shows their total contribution has norm at most $10v_aLS/D$. The identity coset unfolds to $v_a \int F(xu)\phi_1(u)\,du$, so we obtain

$$\left| \int_{[U(V)]} F(h)K(h)\,dh - v_a \int F(xu)\phi_1(u)\,du \right| \leq 10v_aLS/D. \tag{C.24}$$

The finite level here is unchanged in the estimate and the bound is uniform over F through its supremum only. This is an explicit proof of the coset-tail part of the deep-level Poincaré estimate, conditional on (C.22).

The other pieces of `eq_zero_of_petersson_restrict_eq_zero` are not consequences of (C.24) alone. For the source application $A = r^4$, $\beta = (\sqrt{G})^{\lceil x \rceil}$, and the radius scale must satisfy $D \geq \beta^{20A(f+1)}$ together with the actual level norm condition. The lower bound on the period of the two discrete-series coefficients, with a single real translate that works at all points and all rows, and the first-minimum bound treating points below the admissible restriction threshold, remain genuine analytic inputs. In particular (C.24) alone cannot be used to conclude row-surjectivity by dismissing all cusp-near points or by assuming arbitrary matrix-coefficient integrals nonzero.

39.9 Status of the relevant obligations

For precision the following is the effect of this chapter on the remaining ledger.

- C08 (starting periods and quantitative row restriction): the unfolded nonidentity Poincaré tail with exact finite level and factor $10/D$ is now written in (C.22)–(C.24). The real two-factor period bound and thin-part estimate, and the exact normalized starting period transportation, still have to be supplied. This obligation remains open.
- C11 (crude rows): inherits the starting-period and real transport dependencies. The analytic carrier estimates do not imply this separate transport; it remains conditional.
- C14 (comparison forms, line scale and separation): the near-maximum argument, torus projection, integration, Fock sparsity, Hom minimum estimate, stable-range finite reproduction, and the quantitative comparison and isolation implication are written above with actual dimensions and error. The analytic implication from exact carrier matching and the row theta identities is established. The remaining inputs are the exact global exchange and class-sum carrier identities and the integral line multiplier, comparison and test-period assertions of their own chapters. Thus the aggregate obligation remains conditional.

- C17 (rank-one conjugate norms, height and separation): the height consequence from the exact conjugate norms is written in Section 39.7. The analytic separation argument has the stated exchange dependencies, shared with C14.
- C18 (depth-indicator attainability and test pairing) and C19 (rank-one averaging): no new claim that the required actual theta lattice representative or weighted derivative exists is made from this chapter; those specialized arithmetic claims retain their previous status.
- S05 (integral value measures and families): (C.19)–(C.21) supply the actual uniform rank-two jet denominator and integral measure, including simultaneous coefficient-place integrality. The Euler insertion and Galois conversion are as in Chapter 43.7. The normalization and nonvanishing specifically at rank two still depend on the exact equal-rank determinant-period identity, which has been isolated from all integrality claims.

Chapter 40

Congruences, auxiliary primes, and the fit family

Chapters 41 and 42 reconstruct the auxiliary-prime and triangular relations, the finite duality and matrix-span arguments, the odd-class and twisting estimates, and the Fitting and first-order calculations. In particular the local algebraic implications should be distinguished from the still substantive existence of the compatible automorphic/Galois data. Chapter 43 proves the odd-law and integral patch-extraction calculations, and gives the exact residual arithmetic obligations.

We now address the integrality issue in the determinant calculation. The notation for ranks and form spaces is as above. In this section a row of a Galois representation and a column of such a representation refer to the distinguished block and the entire representation, respectively, not to vectors in W and C. Let $B_0 = B/(mk)$, where $B = fa(\log b)^3$. We explain the reduction of the existence of a fit family to a global twisting theorem, a first-order congruence theorem, and the local crystalline estimates. The distinction between established linear algebra and the remaining specialized number-theoretic statements is important. Assertions identified as *formal-source inputs* below are precise imports from the accompanying Lean proof, not results independently proved in this manuscript. Their proofs require the auxiliary constructions indicated by the source references.

40.1 The integral projection problem

Consider an auxiliary unitary pair of odd source rank $r' \le r$ and odd rank difference $t' \le t$; its definite target has rank $r' + t'$. Write $\mathfrak{A}$ for its finite dimensional space of algebraic forms, with the class-sum pairing, and $Z \subseteq \mathfrak{A}$ for an image of theta lifting. Let P_Z be the orthogonal projection. The auxiliary pair has a finite level group, a chosen group-algebra idempotent e, and commuting split-place Hecke polynomials $H_w(X)$. As in Chapter 31, their test value is denoted X_w. The condition imposed on a form v is

$$v \perp (Z^{\perp} \cap \operatorname{im} e \cap \ker H_w(X_w)) \quad \text{for every split } w \text{ outside the level and its conjugate.} \tag{40.1}$$

It is essential that the subspace in (40.1) involves a kernel at each such place, rather than merely the space of theta lifts. These kernels can contain other forms with a Galois representation resembling that of a lift. We call a form satisfying (40.1) tested. The source definitions are `AuxPair.IsTested` and `DenominatorLe` in `New/Selmer/FitTwists.lean`.

Fix a prime ℓ and an abstract field isomorphism $\iota : \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$. The absolute value on the latter is normalized by $|\ell|_\ell = \ell^{-1}$; put $v_\ell(z) = -\log|z|_\ell / \log \ell$. For an integral coefficient

ring $\mathcal{O}$ in a finite extension of $\mathbb{Q}_\ell$, let $\lambda_{\mathcal{O}}$ be its ordinary module length and $e_{\mathcal{O}} = \lambda_{\mathcal{O}}(\mathcal{O}/\ell)$. We use $\lambda = \lambda_{\mathcal{O}}/e_{\mathcal{O}}$ for normalized length. Thus $\lambda(\mathcal{O}/(a)) = v_\ell(a)$ for nonzero $a \in \mathcal{O}$. The conventions extend to the value $+\infty$ when necessary. Integral forms carry the lattice defined by their Weyl-coordinate values; integrality at all ℓ-adic embeddings is distinguished from integrality at one fixed embedding.

If $q = \dim_{\mathbb{C}} Z$, the denominator condition at ι with allowance β is

$$\left|\iota\left(\det\left(\langle u_a, P_Z v_b\rangle\right)_{a,b=1}^{q}\right)\right|_\ell \le \exp(qr'\beta) \tag{40.2}$$

for all $u_a \in \mathfrak{A}$ integral at every ℓ-adic embedding and $v_b \in \mathfrak{A}$ integral at ι and tested as in (40.1). The conditions concern the determinant of Hermitian pairings, not the norm of an arbitrary matrix of coefficients. In particular, r' is the rank of the source unitary group, distinct from q; the factor $r'\beta$ represents the loss allowed per form in the determinant. If $q = 0$ the empty determinant is one. For the good lifts one uses $e = e_0$ and Z the image of the auxiliary lift on good forms. For the rank-one comparison pairs one uses the tuple of labels $X - 1$, $r' = 1$, and Z the line generated by the particular trivial lift. The subspace in (40.1) is then the corresponding kernel of descent.

The height condition on the special values concerns ratios of twisted to untwisted Hecke values. Denote by $\rho_p(\vartheta, s)$ the ratio `Data.Pair.ratio` for a ramified pair p and a sign datum s. The letter p in a pair-label is not necessarily a rational prime. We require ℓ-adic bounds on the inverse ratios for all such pairs. Recall that `Data.Pair.IsRamified` specifies that the tame character of a ramified pair is nontrivial exactly at the chosen places above $g_0 g_1$.

Definition 40.1. For an admissible datum $\mathcal{D}$ a fit family consists, for every ι (including the choice of ℓ), of an m-fold twist θ_ι with conductor prime to ℓ. For every $1 \le j \le m$ there exists a family of real allowances $\beta_{j,\ell}$ such that

$$\sum_{\ell \in T} \beta_{j,\ell} \le B_0 \quad \text{for every finite set } T \text{ of rational primes,}$$

and (40.2) holds at the character θ_ι^j both for the good auxiliary lift at step j and for all rank-one pairs of index j. For every j, p, s as above, the family of inverse ratios $\rho_p(\theta_\iota^j, s)^{-1}$ has height at most x^{999} in the sense of the height predicate of the determinant argument.

In `Data.IsFitFamily`, the index j is a lift index everywhere. The rank-one pairs in `Data.Pair.IsRankOne` have integer parameter z with $m < z \le r$, prescribed tame part ω^j, and weights between $w_j + r - t$ and w_j. All conditions are imposed for the same character and its powers. The definition does not require a single twist working for all primes, nor does it allow independent choices for the arrows of one chain.

Theorem 40.2 (Fit family)**.** *Every admissible datum has a fit family.*

The intended proof is `exists_isFitFamily` in `New/Selmer/FitTwistsExist.lean`. We give its quantitative deduction from the other inputs in this section. An integrality estimate of this strength is necessary: a bound on a projection over $\mathbb{C}$ says nothing about its image of an integral form at ℓ, and a global estimate with an uncontrolled contribution from a new auxiliary conductor need not be usable in the final height sum.

40.2 Labels and Galois representations of lifted eigensystems

We summarize the normalization used for the attached Galois representations. Put $G_E = \mathrm{Gal}(\overline{E}/E)$, let c denote complex conjugation in the ambient absolute Galois group, write

$g^c = cgc^{-1}$, and put $\epsilon_\ell = \chi_{\mathrm{cyc},\ell}$. Thus ϵ_ℓ^{-e} is the character of $\mathcal{O}(-e)$. A character of $U(1)$ is transferred to an anticyclotomic character of G_E by base change and class field theory; ψ denotes the avatar of its inverse. For the auxiliary pair the source rescales the representation attached to a form by a particular algebraic character, `chiOneInv`. If A is a place over a split unramified $w \nmid \ell$ and g a geometric Frobenius normalized as in the source, the scalar of this character is $\iota(a_O(w))^{-1}$, where $a_O(w)$ is the Frobenius character value of the pair (distinct from the Hecke test point X_w). This normalization is designed so that the string in the theta-lift Hecke relation becomes the fixed Tate string $1, \epsilon_\ell^{-1}, \dots, \epsilon_\ell^{1-t'}$.

The key input is not merely an assignment of semisimple Galois representations to eigensystems. It is the simultaneous assignment to all the lifted systems of a row representation T of rank r', independent of the twist, with prescribed inertia and jumps, and an identity of traces for the whole normalized representation U:

$$\operatorname{tr} U(g) = \psi(g) \operatorname{tr} T(g) + \sum_{e=0}^{t'-1} \epsilon_\ell(g)^{-e} \qquad (g \in G_E). \tag{40.3}$$

For the first block the field of coefficients and lattice can depend on ℓ and ι; the row T is the one associated to the untwisted system. It must be possible to work with integral lattices that satisfy the balance and matrix-span hypotheses below.

The labels at a chosen place v are monic squarefree polynomials c_v of degree r' over its residue field. The roots label residue-field characters, and twisting by the inverse tame character scales the roots by the corresponding residue-field element. The label thus modified is denoted $\widetilde{c}_v$. The aptness conditions include $1 \notin \mathrm{Roots}(\widetilde{c}_v)$; for $r' = r$, at least one primitive label occurs above each of g_0, g_1, while for $r' = 1$ the unmodified label used is $X - 1$. The ranks r' and t' are odd. The other conditions specify the chosen places, the extra tame level, and the tame part $\omega^j\vartheta$ of the auxiliary character. This is `IsApt` in `New/Selmer/AptTupleDefs.lean`; aptness is checked for the good and rank-one pairs by `isApt_auxPair` and `isApt_rank_one` in `New/Selmer/HeckeStableSubspaces.lean`.

Here are the row conditions that enter the congruence argument. Let M_g denote the integer `Data.chosenGroupOrder`, the order of the finite product of general linear groups used to bound automorphism and level denominators; in particular

$$g_i^a - 1 \mid M_g \quad (i = 0, 1,\ 1 \le a \le rt). \tag{40.4}$$

Define

$$c_\ell = v_\ell(M_g) + 2^{-\ell}.$$

A row is continuous, trace-nondegenerate, and conjugate self-dual with the normalizing Tate shift: in the rank-difference convention of the source the trace identity has the factor $\epsilon_\ell(g)^{1-t'}$. It is unramified off ℓg and crystalline at primes above ℓ prime to g, potentially crystalline at the other stipulated places, with the indicated multiset of jumps. The row type requires, for each complex embedding, jumps either in a low interval bounded by h_2^2 in absolute value (and bounded below by $h_2/2$), or within t of the appropriately signed w_j. The number of jumps with absolute value at most $2h_2^2$ is bounded by $r'm/k$, and the number of negative jumps at conjugate embeddings together is at most r'. Further:

1. At each place above g not above ℓ, some tame inertia generator γ (the specified one at the chosen places) has determinant bound

   $$v_\ell(\det(T(\gamma) - 1)),\ v_\ell(\det(T((\gamma^c)^{-1}) - 1)) \le r' c_\ell.$$

   The inertia action is killed by an integer prime to the residue characteristic at that place.

2. In all sufficiently large coefficient rings, each lattice representation $T_{\mathcal{O}}$ spans the full algebra $M_{r'}(\mathcal{O})$ as an $\mathcal{O}$-module by its group elements. There is a Galois element having scalar action and cyclotomic character $g_i^{r'}$ for one of the two chosen primes.

3. There is also a scalar element in inertia above g off ℓ, with scalar s such that all powers $(s^{\ell^a})^j$ for $1 \le j \le 4r'+2$ differ from one by a unit, for all sufficiently large a as in the source. In particular one can arrange $s-1$ and $s+1$ to be units.

See `IsColumn`, `IsRow` and `HasRow` in `New/Selmer/AptTupleDefs.lean`, and the row-type definition in `New/Selmer/DeterminantBound.lean`. The precise phrase "for all sufficiently large coefficient rings" is the eventual assertion over the directed collection of ℓ-adic coefficient rings; it permits further finite extension after choosing the twist or an auxiliary level. We do not identify the algebraic closure with one finite coefficient ring.

Proposition 40.3 (Formal-source input: rows for all twists)**.** *For every nonzero Hecke eigensystem occurring in the good forms at the final single-pair source W_m, and every auxiliary lift index j, there is a row T as above satisfying (40.3) for each twist of the auxiliary pair and every attached normalized representation of a lifted eigensystem. The corresponding statement holds for every rank-one pair, with the untwisted rank-one system. All relevant row representations can consequently be put in one finite list for a fixed $\ell, \iota, \mathcal{D}$.*

This is `exists_hasRow_auxPair` and `exists_row_for_all_twists` in `New/Selmer/RowsExist.lean`, applied using the apt labels and rank-one statements. The passage from an automorphic eigensystem to a labelled column uses the local-global compatibility packages in `Lib/GaloisRep` and `New/Selmer/LabelledColumns.lean`. The latter obtains the conjugate self-dual inertia polynomials and proves `IsLabelledColumn.isRow`. At a chosen place, for inertial s, there are $r'+t'$ roots y annihilated by $y^{g_i^d-1}=1$ for some $1 \le d \le r$, and the product of $X-y$ kills s and the conjugate inverse on every labelled column of the specified type. For an untwisted row no root is 1 at the distinguished generator. This is `exists_inertia_eigenvalues`. The proof uses the primitive label above a prime different from ℓ to enforce nondegeneracy and the matrix-span condition. The row-type verification uses the explicit highest weights; the bound on the number of low jumps is what controls the crystalline defect in the local part of the argument.

The finiteness assertion deserves emphasis. The space of good forms is finite dimensional, and the appropriate Hecke operators are simultaneously diagonalizable. For a finite-dimensional space the commuting normal Hecke operators admit a common eigenbasis, and therefore only finitely many characters occur. The equality of relevant multiplicities across signatures identifies the list used in constructing the auxiliary pairs. There are only finitely many pairs with the prescribed weights and tame levels: they are characters in finite sets at fixed conductor and prescribed infinity type, and the construction ranges over finitely many integer weights. Thus the row list may be formed before choosing the finite-order twist. The same is not true of a putative bound obtained directly from the infinitely variable twists of all eigensystems.

We record two algebraic details used in this reduction. First, if the target of a one-dimensional polynomial label is a value z of finite order and neither of z^a, z^b becomes equal after raising to any ℓ-power, then $|z^a - z^b|_\ell = 1$. Indeed write $z = z_{\ell'} z_\ell$ as commuting roots of unity of prime-to-ℓ and ℓ-power order. Reduction modulo the maximal ideal kills z_ℓ and is injective on roots of unity of prime-to-ℓ order (by the separability of X^n-1 and the uniqueness of a lift of the root 1 in the principal unit group). The premise says $z_{\ell'}^a \ne z_{\ell'}^b$, so their difference is nonzero on reduction. Second, if traces of two representations agree on every group element in characteristic zero and each has nondegenerate trace pairing on its image

algebra, their representations are conjugate after matching multiplicities; this is the semisimple trace argument used to turn (40.3) into block diagonal lattices for the distinguished systems. To see the role of trace nondegeneracy, consider finite sums in the group algebra. If such a sum acts as zero in one representation, pairing its product with every group element gives trace zero in the other; nondegeneracy implies that it acts as zero in the other as well. Thus the image algebras identify with their trace functionals. Over an algebraically closed field the multiplicity of each simple module is determined by this functional, giving equivalence. For the source proof the relevant finiteness and trace-nondegeneracy properties of the attached representations are part of the input, not automatic from comparing unramified Euler factors alone.

40.3 Selmer modules over finite coefficient rings

For a stable lattice $T_{\mathcal{O}}$ with row action $R : G_E \to \mathrm{GL}_{r'}(\mathcal{O})$ and a finite-order character ψ with values in $\mathcal{O}^\times$, let V_M be the twist of $(\mathcal{O}/\ell^M)^{r'}$ by $\psi \otimes R$. Its continuous cocycles obey

$$z(gh) = z(g) + (\psi R)(g)z(h).$$

For a finite set of places S stable under conjugation, and a prescribed family $\mathcal{L}$ of finitely supported linear relations on functions on G_E, write

$$\mathfrak{S}_M(\psi R; S, \mathcal{L}) = \left\{ z \in Z^1(G_E, V_M) : \ z(I_v) = 0 \ (v \notin S), \ z \text{ obeys } \mathcal{L} \right\}.$$

At primes over ℓ the family $\mathcal{L}$ is the collection of linear relations obeyed by the characteristic-zero crystalline cocycles Z^1_f of the chosen integral lattice, pulled from the decomposition groups to G_E; it is not the assertion that each torsion cocycle itself is the reduction of a crystalline cocycle. This weaker formulation is important in the local comparison; it is `crysRels` in `New/Selmer/DeterminantBound.lean`. The ambient space is a module of actual cocycles, not cohomology classes. The source `SelmerLengthLe` normalizes by one evaluation $z(\gamma) = 0$, where $\psi R(\gamma) = sI$ and $s - 1$ is a unit, $\epsilon_\ell(\gamma) = 1$.

The normalization has a simple conceptual meaning. If any module of local conditions is stable under adding coboundaries, the evaluation at γ maps its coboundary subgroup isomorphically onto V_M: the coboundary defined by $v \in V_M$ is $\delta v(g) = ((\psi R)(g) - 1)v$, and $\delta v(\gamma) = (s - 1)v$. Thus for every cocycle z there is a unique $v = (s-1)^{-1}z(\gamma)$ such that $z - \delta v$ vanishes at γ. The normalized cocycles then represent the corresponding Selmer classes uniquely. This argument explains why an actual cocycle-module bound can stand in for a more familiar cohomological bound. The source includes the scalar and the normalization in its definitions because the use of both at the integral level is essential later.

Several times we shall use the following elementary length facts. Over a DVR, if $A \in M_N(\mathcal{O})$ has nonzero determinant, the kernel of A on $(\mathcal{O}/(a))^N$ has length at most $\lambda_{\mathcal{O}}(\mathcal{O}/(\det A))$. This follows by writing A in Smith normal form with diagonal entries π^{e_i} up to units; the kernel on each factor has length $\min(e_i, v_\pi(a))$. If $Y, Z \subseteq V$ are submodules of a finite-length module, the length lost by intersection with Z equals $\lambda(Y/(Y \cap Z))$, and such relative lengths are subadditive under successive intersections. Thus if each new linear condition takes values in a torsion module of length at most c_j, imposing all of them decreases length by at most $\sum c_j$.

Definition 40.4 (Balanced lattices)**.** A representation R over a coefficient ring $\mathcal{O}$ is balanced up to a constant β_0 at depth M for a ramification set S, relations $\mathcal{L}$, and normalizer γ if it has the following properties. Its reduction modulo ℓ^M is locally constant; it is unramified off S; its group matrices span $M_{r'}(\mathcal{O})$; the residue field has more than $4r' + 2$ elements. The

distinguished inertial element γ lies above S off ℓ, with $R(\gamma) = R((\gamma^c)^{-1}) = sI$ and $s \pm 1$ units. The coboundaries obey $\mathcal{L}$ modulo every divisor of ℓ^M. Finally there are $g \in G_E$ and scalars α, α' with $R(g) = \alpha I$, $R((g^c)^{-1}) = \alpha' I$, such that for each $a \mid \ell^M$ the local cocycle conditions satisfy

$$\begin{aligned}\sum_{v \in S} \inf_{A|v} \lambda_{\mathcal{O}}(Z^1_v(\mathcal{L}, \mathcal{O}/(a))) + r' v_\pi(\alpha - \epsilon_\ell(g^c)\alpha') \\ \leq (|S| + f)\lambda_{\mathcal{O}}((\mathcal{O}/(a))^{r'}) + \beta_0.\end{aligned} \tag{40.5}$$

Here v_π is the valuation equal to one on a uniformizer of $\mathcal{O}$ and Z^1_v denotes the cocycles on the local decomposition group obeying the induced relations. The local fields and closures are chosen as in the global representation. This is the predicate `IsBalanced` of `New/Selmer/Corepresentable.lean`.

Proposition 40.5 (Formal-source input: balanced row lattices)**.** *Fix a row T of rank r' and a chosen place w above g not above ℓ, with tame generator z. For every sufficiently large coefficient ring and depth M there is a lattice representation R of T, a normalizer γ and β_0 making it balanced for ramification at the specified level and the local crystalline relations. They satisfy*

$$\frac{\beta_0}{e_{\mathcal{O}}} + v_\ell \det(R(z) - 1) + v_\ell \det(R((z^c)^{-1}) - 1) \leq 10 f r' c_\ell. \tag{40.6}$$

This is `IsRow.eventually_isBalanced` in `New/Selmer/RowBalanced.lean`; the finite ramification set may be taken as the places over ℓg embedded in the common exceptional set of primes. The proof relies on a local-global comparison of dimensions of crystalline cocycles and the scalar difference. The arithmetic point is the factor $10 f r' c_\ell$ rather than a bound in terms of the depth M: the depth will be much larger than this and is not an available contribution to the determinant allowance. We retain (40.5) to identify exactly where local conditions enter the subsequent use of global duality.

40.4 New level characters and tied cocycle conditions

Let S_0 be a fixed finite exceptional set containing the original ramification places, the places over ℓ, and the places of the prescribed levels that must not be used as auxiliary places. A finite set Q is auxiliary if each place $P \in Q$ has prime norm, both P and P^c lie off S_0, and $P^c \notin Q$. Thus the rational place below P is split in the relevant quadratic extension, and the auxiliary conditions at the two conjugate sides can be specified compatibly. The formal definition is `IsAuxiliary`. We reserve Q for the modifying conductor of an actual twist, and J initially for a set of auxiliary places carrying further linear conditions. The subsequent Taylor–Wiles places used in the projection argument are different sets.

For a depth N write $q_N = \ell^N$. Auxiliary prime data consist of a place P with $\mathrm{N}P \equiv 1 \pmod{q_N}$, a chosen geometric place above it, a tame inertia generator τ_P, an arithmetic Frobenius φ_P, and an anticyclotomic homomorphism

$$h_P : G_E \longrightarrow \mathbb{Z}/q_N\mathbb{Z}, \qquad h_P(\tau_P) = 1,$$

trivial on the prohibited local subgroups and of conductor supported on P, P^c. The condition of coming from $U(1)$ includes the local triviality and anticyclotomic condition needed for the subsequent passage to tame characters. Frobenius and inertia nomenclature follows the conventions of `AuxPrimeData` and `IsAnticyclotomicAt` in `New/Selmer/Corepresentable.lean`.

Proposition 40.6 (One anticyclotomic prime (see Proposition 41.9))**.** *Let z lie in the normal closure of the inertia groups above a finite set R outside the original prescribed set (and its conjugate) and the places over ℓ. Given a neighborhood U of z, and any N, there is a new prime-norm place P outside $R \cup R^c$ with auxiliary prime data at depth N, with $\varphi_P \in U$, $h_P(\varphi_P) = 0$, and h_P trivial at the specified old local places and above ℓ.*

This is the precise auxiliary-prime mechanism of `exists_isAnticyclotomicAt` in `New/Selmer/Corepresentable.lean`. Its proof uses the finite generation of the group of S-units. A Kummer extension is formed by adjoining q_N-th roots of a finite generating system and the q_N-th roots of unity. Inertia at allowed places fixes these roots as required. Chebotarev then produces a prime-norm place with Frobenius close to z and whose residue field satisfies the splitting conditions. The local residue-field unit logarithm, sending a suitable tame unit to 1 modulo q_N, is trivial on the image of the prescribed global units, so it descends to the idelic quotient after extending along the chosen ideal classes. Taking the quotient by complex conjugation realizes an anticyclotomic class-field character. One finally modifies the representative of Frobenius by an inertia element to obtain $h_P(\varphi_P) = 0$ (in additive notation). The compatibility of the global unit condition and the Frobenius prescription is the point of the Kummer construction; it cannot be obtained by selecting an arbitrary prime congruent to 1 modulo ℓ^N.

A pencil of auxiliary places consists of an auxiliary finite set J, a helper $P_0 \in J$, the data just described for every $P \in J$, and elements $t_{P,i} \in \mathcal{O}$ for the different row representations R_i. Besides the auxiliary-place conditions, the following hold:

$$h_{P'}(\varphi_P) = \begin{cases} 1, & P' = P_0, \\ 0, & P' \neq P_0, \end{cases} \quad (P \neq P_0), \tag{40.7}$$

φ_{P_0} lies in a prescribed neighborhood of a chosen element z, and for every i with $t_{P,i} \neq 0$ both φ_P and $(\varphi_P^c)^{-1}$ act trivially on R_i modulo ℓ^M. The tied conditions for row i are

$$\begin{aligned} c(\tau_P) &= t_{P,i}\, c(\varphi_P), \\ c((\tau_P^c)^{-1}) &= t_{P,i}\, c((\varphi_P^c)^{-1}). \end{aligned} \tag{40.8}$$

They are linear relations on cocycles unramified outside $S \cup J \cup J^c$. Notice the special role of P_0: although each row uses different slopes, the chosen auxiliary places and their characters will be common to all rows. This permits one actual twisting character at the end. The source definitions are `IsPencil` and `tieRelations`.

A basic property of these pencils explains how imposing and then removing ties can be controlled. For a tuple $y = (y_P)$ of q_N-th roots of unity, viewed as units modulo some torsion modulus, define

$$\psi_y(g) = \prod_{P \in J} y_P^{h_P(g)}. \tag{40.9}$$

It follows from the unramified condition at the other primes that $h_{P'}(\tau_P) = \delta_{PP'}$, and anticyclotomicity gives the same values on $(\tau_P^c)^{-1}$. Thus ψ_y has value y_P at the two tame generators at P, P^c, and by (40.7) has the value y_{P_0} at both of the appropriate Frobenius elements when $P \neq P_0$. This is `IsPencil.pointChar_idOrConjInv` in `New/Selmer/TiltingLemma.lean`. The point of (40.7) is that one can choose different points on a line $y_P = 1 + t_{P,i}(u - 1)$, with helper coordinate u, for each row, and still use polynomial interpolation to find one common character.

Lemma 40.7 (Removal of a tie; elementary calculation)**.** *Let V be a finite-length module of cocycles over a DVR, with values in $(\mathcal{O}/(a))^{r'}$. Suppose that at one pair of generators (τ, φ)*

the induced affine actions of τ and φ commute. Suppose also that for the twisted action R', one has $R'(\varphi) \equiv uS$ modulo a and

$$R'(\tau) - 1 = t(R'(\varphi) - 1)$$

on the coefficient module. Then the length lost by imposing the relation $c(\tau) = tc(\varphi)$ is at most $v_\pi(\det(uS - 1))$ (with the infinite case interpreted as vacuous).

Proof. Write the affine action as $v \mapsto R'(g)v + c(g)$. Evaluating commutativity at zero gives

$$(R'(\varphi) - 1)c(\tau) = (R'(\tau) - 1)c(\varphi).$$

Hence the defect $c(\tau) - tc(\varphi)$ belongs to the kernel of $uS - 1$. The quotient of V by the tied submodule identifies with the image of the defect map, and the kernel estimate from Smith normal form bounds its length by the indicated valuation. □

In the application, tame local relations say that Frobenius conjugates a tame inertia element by the power $\mathrm{N}P$. With $\mathrm{N}P \equiv 1 \pmod{\ell^N}$ and N larger than the depth of the coefficients and of the twist, the corresponding affine actions commute modulo ℓ^M. The relation between $R'(\tau)$ and $R'(\varphi)$ on the line follows from (40.9): if the slope is nonzero the original row matrix at Frobenius is the identity modulo ℓ^M, whereas if the slope is zero the twisted inertia action is trivial. At the helper, removing ties uses $u = 1$ instead. Summing Lemma 40.7 over both conjugate generators and the subset of places whose ties are removed gives the exact determinant-loss term in `IsPencil.length_le_of_drop_ties`, in `New/Selmer/TiltingLemma.lean`. The full source statement permits specified replacements for the Frobenius elements as long as their matrices agree modulo ℓ^M. The ramification and affine-action assertions just invoked are hypotheses verified there; in particular one does not divide by ℓ or by residue cardinalities inside the torsion module.

40.5 Shrinking several tied Selmer modules

There are two logically separate ingredients in the existence of a pencil that shrinks every module. One is finite-dimensional linear algebra over a DVR; the other is a reciprocity lemma ensuring a new place on which a long module produces a nonzero detection. We record both, because the arithmetic conclusion does not follow from mere genericity of slopes.

Lemma 40.8 (Tilting a pair of linear maps)**.** *Let $\mathcal{O}$ be a DVR with residue field κ, let $A = \mathcal{O}/I$ have finite length, and let $\Phi_0, \Phi_1 : V \to A^u$ be $\mathcal{O}$-linear. For a finite-length submodule $Y \subset V$, all except at most u residue classes $s \pmod{\mathfrak{m}_\mathcal{O}}$ satisfy*

$$\lambda_\mathcal{O}\big(Y \cap \ker(s\Phi_0 - \Phi_1)\big) \le \lambda_\mathcal{O}\big(Y \cap \ker \Phi_0 \cap \Phi_1^{-1}(\Phi_0 Y)\big). \tag{40.10}$$

Proof. Let Y_1 be the sum of the kernels on the left as s varies. Then $Y_1 \subseteq Y$, and $\Phi_1 Y_1 \subseteq \Phi_0 Y_1$ because this holds on each summand. Let $L \subseteq \mathcal{O}^u$ be the inverse image of $\Phi_0 Y_1$ under reduction modulo I. Since L is free of rank at most u, lift the quotient map $\pi : L \to \Phi_0 Y_1$ to a map $\sigma : L \to Y_1$ with $\Phi_0 \sigma = \pi$. Lift $\Phi_1 \sigma$ to an endomorphism T of L using the same projectivity. For every s whose reduction is not a root of the characteristic polynomial of the reduction of T, the endomorphism $sI - T$ of L is invertible. Therefore

$$(s\Phi_0 - \Phi_1)\sigma(L) = \pi((sI - T)L) = \Phi_0 Y_1.$$

The image of $s\Phi_0 - \Phi_1$ on Y_1 is contained in $\Phi_0 Y_1$, hence equal to it. Its kernel on Y_1 has length $\lambda_\mathcal{O}(Y_1) - \lambda_\mathcal{O}(\Phi_0 Y_1) = \lambda_\mathcal{O}(Y_1 \cap \ker \Phi_0)$. The kernel on Y is included in Y_1 by definition. Finally $Y_1 \cap \ker \Phi_0 \subseteq Y \cap \ker \Phi_0 \cap \Phi_1^{-1}(\Phi_0 Y)$, giving (40.10). □

This is the lemma `tilting`, in `New/Selmer/TiltingLemma.lean`. In the arithmetic application $u = 2r'$, because both conjugate sides are included in each evaluation. Avoiding the exceptional slopes at two different moduli, and also the slope zero, requires at most $4r' + 1$ exceptional residue classes. This explains the residue-field requirement in `IsBalanced`.

The reciprocity statement used in conjunction with this lemma is the following special form of a global-duality obstruction.

Proposition 40.9 (Formal-source input: the mirror criterion)**.** *Let R and the pencil be as above, with R balanced up to β_0, and $a \mid \ell^M$, $a \neq 0$. Let $X(J, a)$ be the corresponding normalized module of cocycles obeying the ties modulo a. Suppose some element of $X(J, a)$ has a unit coordinate. If for every admissible enlargement by a new prime P, compatible with the pencil for every choice of its new slope, the image of $X(J, a)$ under evaluation at the two Frobenius elements lies in the image under tame-inertia evaluation of the relaxed module (in which the new ramification is allowed but its ties not imposed), then*

$$\lambda_{\mathcal{O}}(X(J, a)) \leq \beta_0.$$

This is `IsPencil.length_le_of_unseen_class` in `New/Selmer/TiltingLemma.lean`. A unit-valued coordinate is used to detect a nontrivial row in the full matrix algebra. The detailed proof constructs dual cocycles via global reciprocity, with isotropic pairings against the class in question. The dual classes, evaluated on a suitable open subgroup on which the row and cyclotomic character act trivially modulo the torsion modulus, are bounded by the size of the torsion subgroup for the scalar difference $\alpha - \epsilon_\ell(g^c)\alpha'$ from (40.5). The "mirror" estimates use the full span of the matrix algebra and the units $s \pm 1$ to constrain that image. The algebraic assertion is `card_le_torsion_of_isotropic` in `New/Selmer/LinePointPowers.lean`; the reciprocity construction and its local restrictions are developed in the proof of the mirror criterion. The determinant or torsion size term is absorbed by the balance bound, not by a global bound on all cohomology classes.

Proposition 40.10 (Boundedly many ties; formal-source input)**.** *Fix finitely many balanced representations R_i at depth M, a neighborhood of an allowed inertial element z, and a divisor a of ℓ^M. There is an integer J_0 such that for every depth N of the auxiliary characters there is a pencil of at most J_0 places with normalized cocycle modules satisfying*

$$\lambda_{\mathcal{O}}(X_i(J, a)) \leq \beta_{0,i} \qquad \text{for every } i.$$

This is `exists_isPencil` in `New/Selmer/TwistTheorem.lean`. The important quantifier is that J_0 is chosen before N. To explain its proof, one inducts on the length of $\mathcal{O}/(a)$. Lifting a good pencil for a smaller divisor a to πa gives a finite a priori bound on the length of the larger-depth modules, by evaluating on the finitely many tame generators. For the larger divisor minimize the total length of all the tied modules subject to preserving the smaller-depth inequalities and an upper bound on the sum of the number of places and this total length. If a row has normalized module longer than the allowed bound, and there is no unit-valued element, it reduces to the smaller depth: multiplication by π identifies the modules killed by the shorter divisor, preserving all the linear relations. If there is a unit-valued element the mirror criterion applies. Unless its conclusion already gives the desired bound, it supplies a new place where the image under Frobenius evaluation is not contained in that of the relaxed module under tame-inertia evaluation.

At such a place, write Y for the relaxed module, Φ_0 for tame-inertia evaluation, and Φ_1 for Frobenius evaluation. Imposing $\Phi_0 = 0$ recovers the original unramified module. The failure of the mirror inclusion gives a strict decrease when imposing the additional condition $\Phi_1(c) \in \Phi_0 Y$. Lemma 40.8 yields a slope giving at least this strict decrease at the larger

depth, and by avoiding the exceptional classes at the smaller depth as well it preserves the smaller-depth inequality. The new slopes for all other rows can be set to zero, so their modules do not increase. Consequently the minimized total length strictly decreases, while the sum of length and number of primes does not increase. This is the contradiction with minimality. The induction bounds the required number of new places independently of the auxiliary-character depth. We have described the induction's length mechanism in order to identify where the key reciprocity input is used; the actual selection of primes satisfying all neighborhood constraints together is part of the cited formal-source assertion.

40.6 Finite presentations and interpolation of lengths

To convert the linear pencils into common characters, one must avoid choosing a different character for every row. The device is to bound the length of each Selmer module by a polynomial in the tame-character parameters. The polynomial need not have a degree bounded in terms of the original rank. A special feature of the root-of-unity argument below is that it handles arbitrary polynomial degree uniformly, after a common finite bound on the number of variables is fixed.

Here is the underlying finite-presentation lemma. Fix a modulus $m_0 \neq 0$, a representation given by matrices over a coefficient ring Λ, a finite ramification set, a subfamily of linear relations L, and an open neighborhood of the identity on which all the relevant actions on the m_0-torsion coefficient modules are trivial. Continuous cocycles of these types can be described by finitely many coordinates (the values at chosen group elements) obeying a submodule of relations in a finite free Λ-module. One standard justification is to choose an open normal subgroup on which the action is trivial, restrict the cocycle there, and observe that its restriction is a homomorphism to a module killed by m_0. The maximal abelian exponent-m_0 quotient of the subgroup unramified off the prescribed finite set is finite, by global class field theory (or the corresponding finite-character finiteness theorem). Thus there is one common finite quotient through which all such cocycles factor, independently of their coefficient module. More explicitly, the restriction on the open subgroup is a homomorphism with the prescribed ramification; it factors through the common quotient because any nonzero value in a torsion module is detected by a homomorphism to $\mathbb{Q}/\mathbb{Z}$ (and hence to the subgroup killed by m_0). Taking normal cores yields the desired finite quotient of the original group. On its finite set of elements, the cocycle relations, inertia-zero relations, and the imposed linear relations are simply linear equations. This is the principle of `exists_corepresenting_coords` in `New/Selmer/Corepresentable.lean`. Its source proof also handles the universal coefficient ring by constructing the relation module directly.

Let the character parameters be $Y = (Y_1, \dots, Y_q)$, with $h : G_E \to (\mathbb{Z}/\ell^N)^q$ a finite family of homomorphisms. Use the polynomial ring $\Lambda[Y]$ for the pointwise operators with coefficients $\prod_j Y_j^{h_j(g)}$ (least nonnegative lifts of the exponents). At an actual tuple of roots of unity, or modulo a depth at which the tuple is a root of unity, these operators represent the twist. The cocycle condition and all the other local conditions give a submodule of relations $R(Y) \subseteq \mathcal{O}[Y]^u$ which contains $\ell^M \mathcal{O}[Y]^u$. This also makes sense at arbitrary tuples: even if the formal operators fail to be a representation there, the algebraic equations define a submodule, and we only need their specialization at true or approximate characters.

Lemma 40.11 (Sharp Fitting polynomial)**.** *Let $R \subseteq \mathcal{O}[Y]^u$ contain $\ell^M \mathcal{O}[Y]^u$. For $Y = y$, write R_y for the specialization and define*

$$X_y = \{\xi \in (\mathcal{O}/\ell^M)^u : \langle r, \xi \rangle = 0 \text{ for every } r \in R_y\}.$$

Fix y_0 over a DVR $\mathcal{O}$. There is $F \in \mathcal{O}[Y]$ with

$$v_\pi(F(y_0)) = \lambda_{\mathcal{O}}(X_{y_0}), \qquad \lambda_{\mathcal{O}'}(X_y) \le v_{\pi'}(F(y))$$

for every specialization over a DVR $\mathcal{O}'$ above $\mathcal{O}$. Here F is mapped to the appropriate coefficient ring in the second formula.

Proof. Over $\mathcal{O}$, choose a basis of the full-rank sublattice $R_{y_0} \subseteq \mathcal{O}^u$. Each basis element is a finite $\mathcal{O}$-linear combination of evaluations of elements of R, so we can lift the basis to u polynomial rows of R. Take F to be their determinant. At y_0, Smith normal form gives elementary divisors π^{s_i}, all dividing ℓ^M because R_{y_0} contains $\ell^M\mathcal{O}^u$. The annihilator modulo ℓ^M has length $\sum_i s_i = v_\pi(F(y_0))$. At any other specialization X_y lies in the kernel modulo ℓ^M of the same matrix of specialized rows. If its determinant is nonzero, the kernel bound by Smith normal form gives the inequality; if zero, the inequality is vacuous. This proof also works if R was presented by an infinite set of relations, since only the finite chosen rows enter the determinant. □

This is the algebraic mechanism of `exists_sharp_fitting_poly` in `New/Selmer/LinePointPowers.lean`, where R is obtained by the preceding corepresentability lemma. The equality rather than a mere upper bound at y_0 is essential: the tied Selmer theorem controls the actual module length, not an arbitrary chosen determinant. The lemma applies also after imposing a selected subset of ties; its sharp specialization can be different for every row.

Given a root of unity $u^{\ell^k} = 1$ and $t \in \mathcal{O}$, the point $y = 1 + t(u-1)$ is an approximate root of unity:

$$\ell \mid (y-1)^{\ell^k}, \qquad \ell^M \mid y^{\ell^N} - 1 \quad (N \ge k + M). \tag{40.11}$$

Indeed in characteristic ℓ, $(u-1)^{\ell^k} = u^{\ell^k} - 1 = 0$, so $y^{\ell^k} \equiv 1 \pmod{\ell}$. If $v \equiv 1 \pmod{\ell^j}$ with $j \ge 1$, the binomial expansion gives $v^\ell \equiv 1 \pmod{\ell^{j+1}}$; iterate this argument. This is `dvd_one_add_mul_pow_sub_one`. In particular, the rows can each be specialized on their own line of approximate characters without giving up the interpretation of the matrices modulo ℓ^M.

40.7 Degree-free selection of roots of unity

We give the polynomial selection argument, since it supplies the main transition from the tied linear conditions to a single actual anticyclotomic character. It is also an example of a quantitative lemma whose conclusion is uniform in the degrees of arbitrarily many high-degree determinants, an essential feature here.

For a polynomial F over the integers of a finite ℓ-adic extension, and $0 < R \le 1$, write $|F|_R = \max_i |a_i|_\ell R^i$ for its Gauss norm, where $F(X) = \sum a_i X^i$. It is multiplicative and nondecreasing in R. We use this multiplicativity over the finite extensions containing the roots of unity in the following proof.

Lemma 40.12 (Simultaneous one-variable avoidance)**.** *Fix $0 < R < 1$ with $|\ell|_\ell < R^{\ell-1}$, and finitely many positive numbers u_i. For all sufficiently large N, in every sufficiently large coefficient ring containing the ℓ^N-th roots of unity, the following holds: if F_i are integral polynomials, of arbitrary degrees, and $|F_i|_R \ge u_i$, then there is a single ℓ^N-th root of unity ζ with*

$$|F_i(\zeta - 1)|_\ell \ge R u_i \qquad \text{for every } i.$$

The required depth is independent of the degrees.

Proof. Put $F=\prod_i F_i$ and $Q_i=\prod_{h\neq i}F_h$. Suppose, for contradiction, that at every ℓ^N-th root of unity at least one inequality fails. Take an integer $a\geq 1$ so large that $R^a<|\ell|_\ell$. For $0\leq j<N$ set $R_j=R^{1/\ell^j}$ and $n=\ell^{j+1}$. The roots α of $P_j(X)=(1+X)^n-1$ all have absolute value less than R_j, since a primitive root of order ℓ^{j+1} satisfies

$$|\alpha|_\ell=|\ell|_\ell^{1/(\ell^j(\ell-1))}<R_j.$$

For each such root choose $i(\alpha)$ with $|F_{i(\alpha)}(\alpha)|_\ell<Ru_{i(\alpha)}$. Let $b=a(j+1)$ and form

$$S_0(X)=\sum_{\alpha:P_j(\alpha)=0}\frac{P_j(X)}{X-\alpha}F_{i(\alpha)}(\alpha)^bQ_{i(\alpha)}(X)^b.$$

The roots of P_j are simple. At each root, S_0 agrees with $P_j'F^b$, so their difference is divisible by P_j. Write $P_j'F^b-S_0=P_jS$. For $T=R_j$ and $T=R_{j+1}$ we have $|P_j|_T=T^n$ and $|P_j'|_T=|n|_\ell$ (the latter because $P_j'=n(1+X)^{n-1}$). Moreover $u_i\leq|F_i|_T$, so

$$|S_0|_T\leq T^{n-1}R^b|F|_T^b<|n|_\ell|F|_T^b,$$

using $T\leq 1$ and $R^{a(j+1)}<|\ell|_\ell^{j+1}$. The ultrametric inequality and multiplicativity give

$$|S|_T=\frac{|n|_\ell|F|_T^b}{T^n}.$$

Since $|S|_{R_j}\leq|S|_{R_{j+1}}$, it follows that

$$\log|F|_{R_{j+1}}-\log|F|_{R_j}\geq\frac{n}{a(j+1)}\log(R_{j+1}/R_j)=\frac{-(\ell-1)\log R}{a(j+1)}.$$

Summing these inequalities over $j<N$ produces a contradiction for large N: the left hand side is at most $-\log|F|_R\leq-\sum_i\log u_i$, whereas the harmonic sum on the right is unbounded. If a lower bound $u_i>1$ makes the hypotheses impossible there is of course nothing to prove. This establishes the uniform assertion. □

This is the interpolation proof of `exists_root_of_unity_eval_large` in `New/Selmer/RootsOfUnityPoints.lean`.

Lemma 40.13 (Simultaneous selection in several variables)**.** *Fix a positive real δ, positive integers m,a,q, a nonnegative integer L_0, and finitely many nonnegative numbers B_i. For every sufficiently large N and every sufficiently large integral coefficient ring, suppose that $F_{i,j}$ $(1\leq j\leq m)$ are integral polynomials in at most q variables. Suppose there exist tuples $u_{i,j}$ with $\ell\mid(u_{i,j,k}-1)^a$ in each coordinate and*

$$v_\ell(F_{i,j}(u_{i,j}))\leq B_i.$$

Let also D_h be monic one-variable polynomials with total degree at most L_0. Then there is one tuple ζ of ℓ^N-th roots of unity such that simultaneously

$$v_\ell(F_{i,j}(\zeta^j))\leq B_i+\delta,\qquad v_\ell(D_h(\zeta_k))\leq\delta\quad\text{for every } k,h. \tag{40.12}$$

Here the depth can be chosen before the polynomials or tuples are specified.

Proof. Choose $R<1$ so close to one that $R^q,R^{L_0+1}>\ell^{-\delta}$ and $R^{\max(ma,\ell-1)}>|\ell|_\ell$. Assume that some coordinates have already been replaced by ζ_k^j in the points for each pair (i,j), and that the norm bound has deteriorated by at most a factor R per replacement. For the

next coordinate, all other coordinates stay fixed (possibly different for different pairs). The polynomial in X given by substituting $(1+X)^j$ in this coordinate has Gauss norm of radius R at least the currently established lower bound. To justify that assertion, the equation $(1+X)^j = u_{i,j,k}$ has a root with $|X|_\ell \le R$: the monic polynomial $((1+X)^j - u_{i,j,k})^a$ has constant term of absolute value at most $|\ell|_\ell < R^{ja}$; if all its roots had absolute value larger than R, their product would violate this bound. The value at such a root is exactly the previously known value of $F_{i,j}$, bounded above by the Gauss norm. We may use the algebraic closure to compare the norm and do not need the root itself in the coefficient ring.

Include in the one-variable family the polynomial $\prod_h D_h(1+X)$, whose monic leading term gives Gauss norm at least R^{L_0}. Apply Lemma 40.12 simultaneously. This selects one ζ_k for the next coordinate, with loss at most a further factor R, and also gives $|\prod_h D_h(\zeta_k)|_\ell \ge R^{L_0+1}$. Because each factor has norm at most one, this lower bound implies the same lower bound for every factor. After at most q steps, the accumulated factor is at least $R^q > \ell^{-\delta}$. The one-variable lemma is applied with the finite list of lower bounds $\ell^{-B_i}R^u$, $0 \le u \le q$, and R^{L_0}, so its threshold is uniform in the polynomial coefficients and their degrees. This yields (40.12). □

The source declaration is `exists_roots_of_unity_ellVal_le`, in the same file. An important application of the monic polynomials D_h is to $Y^j - 1$: at a selected tuple all the first m powers have nontrivial tame values, with total valuation as small as desired after accounting for the bounded number of variables.

40.8 The global twisting theorem and value families

The twisting theorem controls more than the Selmer modules. It also preserves the valuation of the special Hecke values needed in the norm comparison. The relevant *value family* (`IsFamily` in `New/Selmer/Corepresentable.lean`) is given by continuous unit-valued characters μ_i and a function $V(Q,\psi)$, defined for auxiliary conductors Q and characters. Its untwisted value $V(\varnothing, 1)$ is nonzero; at each fixed Q it is an integral finite linear combination of values of ψ, say

$$V(Q,\psi) = \sum_{a \in A_Q} c_a \psi(g_a), \qquad |c_a|_\ell \le 1. \tag{40.13}$$

In particular, on a character given by (40.9), this specializes an integral polynomial in Y_P. The family satisfies a norm relation in the form of an estimate: if $P \notin Q$ and ψ is anticyclotomic of conductor Q, the inequality

$$v_\ell(1 - \mu_i(\varphi_P)\psi(\varphi_P)^{\pm 1}) \le c \quad \text{for all } i$$

implies

$$v_\ell V(Q \cup \{P\}, \psi) \le v_\ell V(Q,\psi) + 8c. \tag{40.14}$$

For a ramified pair p the source constructs these families by an integral special-value measure, including the factors at the local level places. More precisely, the integral measure F satisfies a distribution relation upon adding v to the modulus: its value is multiplied by a product of at most four factors

$$\prod_{u \in K_0} \prod_{\epsilon \in \{1,-1\}} (1 - (\nu\vartheta)_{\mathrm{bc}}(\varpi_v)^\epsilon q_v^{u-z}), \quad K_0 \subseteq \{0,1\},$$

where z is the integer of the pair and ν its underlying character. The value on a twisted character is $F(1)\,\mathrm{N}(Q)^h \rho_p(\vartheta)^{h+1}$ for some nonnegative integer h. This is the formal-source

input `Data.Pair.exists_isIntegralMeasure` in `New/Selmer/ValueFamilies.lean`; its derivation uses the algebraic CM Hecke special values and integral interpolation. The passage to the fixed-sign ratios and to the Galois-valued functions in (40.13) is `exists_isFamily` in that file. Because the character of a ramified pair is nontrivial at the chosen place w off ℓ, the listed auxiliary characters satisfy

$$v_\ell(\mu_i(z)-1) \le v_\ell(q_w-1) \le c_\ell \tag{40.15}$$

for its tame generator z. The last inequality follows from (40.4). Inverting the special-value ratio changes the valuation inequality to an upper bound on its ℓ-adic norm. No assumption of nondivisibility of every original Hecke value is needed.

Proposition 40.14 (Formal-source input: simultaneous anticyclotomic twisting)**.** *Let R_i be a finite collection of balanced representations with local bounds $\beta_{0,i}$ at depth M, and fix an inertial element z off ℓ as in Proposition 40.6. Suppose $v_\ell(\mu_i(z)-1) \le c$ for the characters of finitely many value families. If*

$$\frac{\beta_{0,i}}{e_{\mathcal{O}}} + v_\ell \det(R_i(z)-1) + v_\ell \det(R_i((z^c)^{-1})-1) \le B_i < \infty,$$

then for every $\eta > 0$ and every m there are an auxiliary set Q and an anticyclotomic character ψ such that, for each $1 \le j \le m$:

1. *the character ψ^j comes from $U(1)$ with conductor Q and is locally trivial at the prescribed old places and above ℓ, and*
$$\sum_{P\in Q} v_\ell(\psi^j(\tau_P)-1) \le \eta;$$
2. *the normalized lengths of the cocycle modules of $\psi^j R_i$ at depth M, with the original crystalline relations and with value zero at the corresponding γ_i, are at most $B_i+\eta$, over all sufficiently large coefficient rings;*
3. *for each value family, $v_\ell V(Q,\psi^j) \le v_\ell V(\varnothing,1) + 8c + \eta$.*

This is `exists_anticyclotomic_twist` in `New/Selmer/TwistTheorem.lean`, with the normalization of lengths as above. To see the use of the preceding lemmas, choose a pencil as in Proposition 40.10, using at most J_0 places. Choose a small allowance $\delta > 0$ with $(8J_0+2)\delta < \eta$. The construction is in two specialization stages. The first uses an ℓ-power root of unity u and the line $y_P = 1 + t_{P,i}(u-1)$ for the ith row, with $y_{P_0} = u$. By (40.11) all such points can be used modulo ℓ^M after the auxiliary depth has been chosen sufficiently large.

Use the sharp Fitting polynomial of Lemma 40.11 for the tied module, with the helper tie omitted. At $u = 1$ the polynomial has valuation bounded by that of the tied Selmer module plus the determinant terms for removing the helper tie. The one-variable avoidance lemma, applied also to the finitely many characteristic polynomials of the matrices at the nonhelper Frobenius places and the Euler factors for the value families, chooses a common u for which all these valuations increase by at most δ. This is why the pencil was required not merely to have suitable slopes but also to have the Frobenii of all nonzero slopes trivial on the corresponding row modulo ℓ^M. Along the chosen line the extra cost of dropping the other ties can then be bounded by Lemma 40.7: these are determinants of $uR_i(\varphi_P)-1$ and their conjugate counterparts. These are precisely among the polynomials selected for the small-valuation conditions.

For each value family one may instead take the line with only the helper coordinate equal to u and the others equal to 1. Its initial valuation costs at most $8c$ by (40.14), using φ_{P_0}

close to z and the bound (40.15). On inserting the other primes successively, their Frobenius values are the helper value by (40.7). By the monic Euler-factor avoidance in the first stage the remaining cost is at most 8δ per insertion. This bounds the value family on this new polynomial point. There is no need to find one tuple lying on the lines constructed for all different rows and value families.

At the second stage use the sharp Fitting polynomials for the unrestricted modules, now evaluated at the good row-specific points just constructed; also include the polynomials for all the value families. The number of variables is at most J_0, and by (40.11) all these specializing points satisfy the near-1 power condition with one common exponent. Lemma 40.13 therefore gives one tuple of ℓ^N-th roots of unity working for all polynomials, and simultaneously for their evaluations at the first m powers of that tuple. The monic factors $Y^j - 1$ ensure nontriviality and bound the valuations at the tame generators. The total error from these selection stages and from inserting the Euler factors is bounded by $(8J_0 + 2)\delta < \eta$. Formula (40.9) gives the character ψ with common conductor, and the finite coefficient rings can be enlarged without changing the normalized length bounds. This describes the proof's organization and supplies the polynomial and length arguments, while the exact global compatibility of the pencil and integral special-value measures is retained in the stated formal-source input.

Corollary 40.15 (Formal-source consequence: twisting all rows)**.** *For a finite collection of rows T_i of ranks r_i of admissible data, fixed ℓ, ι and any depth M, there is an m-fold twist θ such that for all $1 \le j \le m$, writing $\vartheta = \theta^j$ and ψ for its inverse avatar, one has*

$$\sum_{P \in Q} v_\ell(\psi(\tau_P) - 1) \le c_\ell, \tag{40.16}$$

$$\lambda\big(\mathfrak{S}_M(\psi T_i; S \cup Q \cup Q^c, \mathcal{L}) \cap \ker \mathrm{ev}_{\gamma_i}\big) \le (10 f r_i + 1) c_\ell, \tag{40.17}$$

$$\left|\iota(\rho_p(\vartheta, s)^{-1})\right|_\ell \le \exp(9 c_\ell \log \ell) \quad \textit{for every ramified } p, s. \tag{40.18}$$

The normalizer satisfies the scalar and unit conditions of SelmerLengthLe*.*

Deduction from the stated inputs. Take a chosen place w above one of g_0, g_1 different from ℓ. Proposition 40.5 applies for all rows after one common finite enlargement of the coefficient ring. Choose $B_i = 10 f r_i c_\ell$ and $\eta = c = c_\ell > 0$ in Proposition 40.14, using (40.15) for all the finite value families. The resulting character is the inverse avatar of a tame character, by the class-field identification `exists_tameChar_invAvatar_eq` of `New/Selmer/ValueFamilies.lean`. The nontriviality at the tame generators for all powers makes it an m-fold twist, while being trivial at the original places leaves the normalizers and crystalline conditions unchanged. The bounds on the normalized Selmer length and local ramification give the first two assertions. Using (40.14) and the ramified-pair norm relations in that file gives $v_\ell(\iota\rho_p(\vartheta, s)) \le 9c_\ell$, equivalent to (40.18). This is exactly `exists_mFoldTwist_selmerLengthLe` in `New/Selmer/TwistForRows.lean`. □

40.9 The Tate characters and triangular Taylor–Wiles places

The bound on the distinguished row block obtained above is not a bound on all congruences of the normalized column. The extra characters $\xi_e = \epsilon_\ell^{-e}$ also give torsion cocycles. It is not possible simply to replace their modules by a conjectural bound on the even cyclotomic class groups. The argument uses instead a finite collection of auxiliary primes that depends on the torsion depth but whose cardinality is bounded independently of that depth. These primes are distinct from the conductor of θ in the twisting theorem.

For a Tate character ξ_e, define $\mathfrak{E}_{\nu,e}(S)$ to consist of the locally constant cocycles z in $(\mathcal{O}/\ell^\nu)(-e)$ unramified outside the places above ℓ and $S \cup S^c$, with $z(g^c) = z(g)$ and vanishing on the prescribed subgroup of the local decomposition groups in the kernel of the cyclotomic character. For $e > 0$ the relevant odd module $\mathfrak{H}^-_{\nu,e}$ is given by the same unramified and local conditions without the added primes and with the odd symmetry condition. At $e = 0$ the odd-class package of the source uses a different convention, namely the unramified cocycles with its local condition without the odd restriction. These are the definitions `evenClasses` in `New/Selmer/Corepresentable.lean` and `shaOdd` in the odd-class bound. We do not implicitly decompose into two direct eigenspaces at $\ell = 2$; the use of sums and differences, and factors of 2, is important at that prime.

Let N_E be the cyclotomic conductor specified for E in the datum, so $E \subset \mathbb{Q}(\zeta_{N_E})$. The following background result is one of the few global class-number terms that survive in the final bound.

Proposition 40.16 (One integer for all odd classes (Theorem 42.14))**.** *There exists an integer $X > 0$ with*

$$X \le 2^{4(N_E+t+3)^{27}} h_E, \tag{40.19}$$

where h_E is the class number of E, such that for every ℓ, every $0 \le e < t$ and every ν, the normalized length of $\mathfrak{H}^-_{\nu,e}$ is at most $v_\ell(X)$. The coefficient rings are assumed, in addition to the integral DVR hypotheses, to contain enough roots of unity of order the exponent of $(\mathbb{Z}/N_E)^\times$.

This is `exists_length_shaOdd_le` in `Lib/Cyclotomic/OddSha/OddShaBound.lean`. It is a finite-level statement uniform in ν, not a generic assertion of Iwasawa nonvanishing. The proof uses the cyclotomic embedding of E, annihilating elements of Stickelberger type, the class-number bound in the trivial twist and explicit estimates of their sizes; it constructs the integer X by multiplying the corresponding annihilator factors. We retain this input as stated, since the bound (40.19), not just finiteness for every fixed (ℓ, e, ν), is needed for the global sum.

In contrast, the size of $\mathfrak{E}_{\nu,e}$ will not be included. To see the substitute, let $a \in G_E$ separate the distinct Tate characters up to ℓ-adic loss. A triangular system of depth ν consists of a finite ordered set J', a nondecreasing type function $\mathrm{ty} : J' \to \{1, \dots, t' - 1\}$, and distinct Taylor–Wiles places w_j with tame generators τ_j and Frobenius elements ϕ_j fixing the ℓ^ν-th roots of unity. The places have prime norm and are conjugate-disjoint. Given bounds H_ℓ and C_ℓ, the `IsPrimeTriangle` predicate includes the following conditions:

1. There is a finite detecting set s of group elements such that evaluation at s detects all unramified-outside-ℓ Tate cocycles modulo ℓ^ν in the indicated range, and $(|J'| + |s|)H_\ell \le C_\ell$.

2. For each $0 \le e < t'$, the product $\prod_{k \ne e}(\xi_e(a) - \xi_k(a))$ divides ℓ^{H_ℓ} in the integral ring.

3. For every $z \in \mathfrak{E}_{\nu,\mathrm{ty}(j)}(\{w_i\})$, if $z(\tau_i) = 0$ for $i < j$ and $z(\phi_i) = 0$ for every earlier i of the same type, then $\ell^{H_\ell} z(\tau_j) = 0$.

4. Write $d_e(g) = \xi_e(g) - 1$. For $e > 0$, $4d_e(a)^2 \mid \ell^{H_\ell}$, and if an even cocycle z with no new ramification is zero at the ϕ_j of type e, then

$$d_e(a)(2z(g)) = d_e(g)(2z(a)) \qquad (g \in G_E).$$

   Thus a controlled multiple of such a cocycle is a coboundary.

The precise predicate is in `New/Selmer/Corepresentable.lean`; writing a as in that predicate accounts for the inverse in the expression of the Tate characters.

Proposition 40.17 (Formal-source input: existence of triangular places)**.** *Suppose $4 \le f$, $N_E + t' < y$, $\epsilon_\ell(\gamma)^a - 1 \mid \ell^h$ for $0 < a < t'$, and $\ell \le y^2$ or $h = 0$. Assume the desired Frobenius conditions U hold on the generic Frobenius data near some element of the form $\delta^{\ell^\nu} g$ for each ν, g, where $\epsilon_\ell(\delta) = 1$. Then there exist an ordered index set J' and a type function, both independent of ν, such that at every depth $\nu > 0$ one can choose Taylor–Wiles places satisfying the triangle conditions with $a = \gamma^{-1}$, $H_\ell = yh$ and $C_\ell = y^{22}h$, and with all their Frobenius data satisfying U.*

This is `exists_isPrimeTriangle` in `New/Selmer/TrianglePrimes.lean`. The source constructs the detecting set uniformly and indexes the triangle by the product of its detecting set and the nontrivial types. The unusual order on this product, increasing by type, is necessary in the subsequent polynomial-support argument. One ingredient is a reciprocity functional z_0 on G_E, locally constant modulo ℓ^ν, which on the kernel of the cyclotomic character modulo ℓ^ν is an odd additive map transforming by cyclotomic weight $e+1$. At split places outside a finite exceptional set, if the functional is zero at all other Frobenius elements in a finite collection, its value on the remaining Frobenius, multiplied by 2, annihilates the corresponding even inertia value, by reciprocity. The source proves existence of such a functional, vanishing at a prescribed set of no more points than were previously used, and taking a unit value on some element of the cyclotomic kernel: `exists_isLaw` in `New/Selmer/ReciprocityLaw.lean`. The even cocycles are tested through all cyclic additive quotients, which detect zero by duality for finite abelian groups. By taking a linear combination of elements of the cyclotomic kernel and applying the products of differences of the cyclotomic weights, the construction makes the new functional vanish on all previously chosen Frobenius elements of other types. It then applies Chebotarev with the generic Frobenius conditions simultaneously. The annihilating error in this selection is the controlled product of differences above, not the length of the even module. This is the mechanism of the induction in Proposition 40.17; the integral construction of the reciprocity functionals and uniform detecting points is retained as part of the formal-source input.

For our datum the elementary separation of the characters is available with $h \le v_\ell(M_g)$ and $y = h_2 = r^8$. Indeed for ℓ larger than h_2^2 the field conductor is prime to ℓ and an inertial cyclotomic value whose reduction generates $\mathbb{F}_\ell^\times$ separates all exponents less than t by units; take $h = 0$. For the smaller primes, choose a degree-one place above the chosen prime $g_i \ne \ell$. Its Frobenius has cyclotomic value g_i, and each $g_i^d - 1$ with $0 < d < t$ divides M_g by (40.4); take $h = v_\ell(M_g)$. This is `exists_cycChar_pow_sub_one_dvd` in `New/Selmer/DenominatorBound.lean`. The admissible size conditions give $N_E + t < h_2$, and the field contains sufficiently many infinite places. The generic Frobenius conditions needed in Proposition 40.17 arise from the Hecke string lemma discussed below. In particular the global odd term in (40.19) and the bounded valuations of the separating products are sufficient even if the even-class modules are very large.

40.10 The first-order Ribet construction

The algebraic argument works with a finite set Y of normalized column representations U_y over $\mathcal{O}$. A subset $\mathfrak{X} \subseteq Y$ consists of the distinguished lifts; for $y \in \mathfrak{X}$ the representation has block form $R_y \oplus \bigoplus_e \xi_e$ by (40.3), where R_y denotes the already twisted row block. For the projection we distinguish one index $y_0 \in \mathfrak{X}$. Use the group algebra $\mathcal{O}[G_E]$, and let a separate

its Tate characters. Choose a normalizer γ with $R_{y_0}(\gamma) = sI$, $s-1$ a unit, and $\xi_e(\gamma) = 1$. Put

$$\mathcal{N} = [\gamma a] - s[a] + p_0 D \sum_g n_g[g], \qquad \beta_y = \det\left(\sum_g \mathcal{N}_g(U_y(g) - 1)\right), \tag{40.20}$$

where p_0 is the row-cocycle modulus, D a separating factor, and the n_g are finitely supported nonnegative integers. This is `ribetNormaliser` in `New/Selmer/RibetCramer.lean`. For distinguished systems $\beta_y = 0$ since the block $\xi_0 = 1$ is annihilated by every $U_y(g) - 1$. On other systems one can arrange $\beta_y \neq 0$ whenever some $U_y(g) - 1$ is invertible by the following finite avoidance observation: each determinant is a polynomial in the finitely many n_g, nonzero as a polynomial if at least one leading matrix $U_y(g) - 1$ is invertible. Finitely many nonzero polynomials over a characteristic-zero field do not vanish on all tuples of nonnegative integers (induct on the number of variables). This is the reason the normalizer is allowed the final summand. One must still treat the systems on which every $U_y(g)$ has eigenvalue 1, rather than assuming those systems are precisely the lifts.

Let $A = \mathcal{O}[X_j : j \in J']$. We use a subalgebra $\mathfrak{S}$ of functions from Y to polynomials over the coefficient field, containing all scalar variables and characteristic polynomial coefficients of the $U_y(g)$. Its trace ideal $\mathfrak{J}$ contains, for all g, $p \operatorname{tr} U_y(g)$, with $p = \ell^\nu$, and the trace ties (for the two copies of j)

$$\operatorname{tr}((U_y(\tau_j) - 1)U_y(g)) - X_j \operatorname{tr}((U_y(\phi_j) - 1)e_j(U_y(\phi_j))U_y(g)), \tag{40.21}$$

where e_j is a polynomial with integral coefficients and $e_j(1) = 1$; conjugate inverse generators are included with the same variable. The precise abstract predicate `IsTraceIdeal` is in `New/Selmer/RibetFirstOrder.lean`. For that file the two conjugate copies may be prescribed independently but share X_j.

The Ribet element assertion is stronger than a mere membership of a determinant in an annihilator. It asserts the existence of $F \in \mathfrak{S}$ with

$$F\beta \in \mathfrak{J}, \qquad \beta_y = 0 \ (y \in \mathfrak{X}), \qquad [X^d]F_y = 0 \ (y \in \mathfrak{X},\ d < \mathbf{1}), \qquad [X^{\mathbf{1}}]F_{y_0} = c_0, \tag{40.22}$$

where $X^{\mathbf{1}} = \prod_{j \in J'} X_j$ and $d < \mathbf{1}$ means a proper coordinatewise divisor of this monomial. In addition $c_0 \neq 0$ has a quantitative upper valuation bound, given below. This is `IsRibetElement`. Crucially the vanishing condition is not that all terms of smaller total degree vanish; the source works with the lower order ideal of exponents coordinatewise at most $\mathbf{1}$. It is precisely this vanishing condition that permits the later first-order coefficient extraction at several primes.

Here are some details of the algebra of producing such an element. Because the distinguished block has full matrix span, one can construct group-algebra expressions isolating the Tate blocks. For each $y \in \mathfrak{X}$, write $R_y(\gamma_y) = s_y I$, $s_y - 1$ a unit, and $\xi_e(\gamma_y) = 1$. The element

$$k = \prod_{y \in \mathfrak{X}} (s_y - [\gamma_y])/(s_y - 1)$$

is integral, acts by zero on every distinguished row block, and by one on all Tate blocks. The factors need not commute: each row representation sends its own factor to zero, which annihilates the entire product in that representation. For each index e, Lagrange interpolation supplies an element q_e with $\xi_i(q_e) = D\delta_{ie}$, where D is a common multiple of the products of differences. Explicitly take a multiple of $\prod_{i \neq e}([a] - \xi_i(a))$ by integral coefficients. Both constructions remain valid in characteristic ℓ after retaining the multiplier D, whereas dividing by the differences would not. These are the two block-separation operations at the start of `ribet_first_order`.

To explain the Fitting module in this proof, let $\mathfrak{M}$ be the finite $\mathcal{O}$-module spanned, inside the family of matrices, by the elements $(U_y(g)-1)_{y\in Y}$. It is finitely generated since Y is finite. A linear map from $\mathfrak{M}$ into a torsion row module produces a cocycle when it respects left multiplication by the $U_y(g)$, because

$$U_y(gh)-1=(U_y(g)-1)+U_y(g)(U_y(h)-1).$$

Conversely a cocycle obeying all linear relations between the family $(U_y(g)-1)_y$ can be evaluated by a linear map on $\mathfrak{M}$ with the same equivariance. This motivates `obeyingCocycles` in `New/Selmer/BlocksAndCocycles.lean`. To compare with local crystalline classes it is useful to extend this graph map slightly: if the target is killed by a, Artin–Rees gives q with $\mathfrak{M}\cap a^q\mathfrak{V}\subseteq a\mathfrak{M}$, for the ambient matrix lattice $\mathfrak{V}$. Its evaluation therefore extends by zero on $a^q\mathfrak{V}$ to $\mathfrak{M}+a^q\mathfrak{V}$, and its graph is a subrepresentation with injective first projection. The map is well defined because differences in the intersection are killed. This is the lemma `obeyingCocycles.exists_graph_subrepresentation`; it also shows why the extracted cocycles are continuous and have the necessary unramifiedness, even though the definition by linear relations alone did not initially impose continuity. Where all U_y act trivially on inertia, the corresponding matrix differences are zero, so the cocycle is zero there. The graph description and continuity of the integral matrices show that a finite reduction of them determines the cocycle.

After imposing the trace ties and the relations of (40.20), one works in the polynomial extension of this finite presentation. At each distinguished block, a bound on the torsion length supplies a determinant (a maximal minor of a relation matrix) with small valuation. For the Tate block one must use a first-order variant: the relations between evaluation at a pair of conjugate inertia elements and evaluation at the Frobenius elements, multiplied by X_j, make the corresponding determinant divisible by the products of X_j of that type. To state its precise arithmetic input, for a module Z of Tate cocycles put $I_e=\{j:\mathrm{ty}(j)=e\}$. The triangle cost with multiplier D and allowance b_e consists of a set L of evaluations such that, for cocycles zero on L, agreement at earlier pairs of tame generators and at earlier Frobenius pairs of the indicated type forces

$$D\big(z(\tau_j)-z((\tau_j^c)^{-1})\big)=0\qquad(j\in I_e).$$

For submodules of these cocycles vanishing at all tame generators and agreeing at the Frobenius pairs of type e, their length modulo D-torsion plus the length cost of the initial evaluations on L is at most b_e. This is `HasTriangleCost`.

Proposition 40.18 (Formal-source input: first-order Ribet bound)**.** *Use the preceding block setup and trace ideal. Suppose the normalized row cocycle module, modulo p_0, obeying the linear matrix-difference relations and vanishing at the new tame generators and its normalizer, has ordinary length at most $b_0<v_\pi(p_0)$. Suppose the corresponding Tate modules, modulo p, have triangle costs b_e and detecting set s as above. Include auxiliary inertial relations of the form $(U_y(\delta_v)-1)o_v(U_y(\delta_v))=0$, where $\xi_e(\delta_v)=1$. Assume also $e_j(\xi_e(\phi_j))\equiv 1 \pmod p$ and $\prod_{k\neq e}(\xi_e(a)-\xi_k(a))\mid D$. Then the Ribet element (40.22) exists with*

$$v_\pi(c_0)\le b_0+\sum_e b_e+t'\left(\sum_v v_\pi(o_v(1))+2|s|v_\pi(D)\right)+(|J'|+1)v_\pi(D).\tag{40.23}$$

Here the Tate-module assertions of triangle cost and detection are required for the modules whose members vanish on all δ_v and whose coboundaries are killed by $\xi_e(a)-1$ (as in the source).

This is `ribet_first_order` in `New/Selmer/RibetFirstOrder.lean`. The block Fitting lemma `exists_mem_fittingIdeal_block` in the same file gives, for one Tate block, a relation in the Fitting ideal of the form $G_e \prod_{j\in I_e} X_j$ with constant-term bound $v_\pi(G_e(0)) \le b_e + (|I_e| + |s|)v_\pi(D)$. The proof successively enlarges presentations, uses the stated cost to clear one evaluation at a time, and then multiplies the resulting minors. When combining blocks, terms not supported in the interval of divisors of $\prod X_j$ can be discarded. More precisely, work modulo the ideal spanned by monomials not dividing $\prod X_j$, and at each step additionally keep track of previous types; the type ordering puts the unwanted earlier variables in the annihilator after extracting the earlier block factors. The direct-sum decomposition of the distinguished representations turns the determinant of a block-flattened presentation matrix into the product of its row determinant and its Tate determinants. The factors D and $o_v(1)$ are inserted to make the block operations integral, accounting for the additional losses in (40.23). Finally a group-algebra Cramer determinant identity turns these minors into an element of the trace ideal after multiplying by β. That identity is `exists_ribet_element_cramer` in `New/Selmer/RibetCramer.lean`; it uses integrality of characteristic polynomials of block matrices over the group algebra, supplied by the indicated Amitsur formula. These Fitting, block Cramer and multivariable support steps are specialized assertions of the formal-source proof, not consequences of the elementary determinant inequality alone.

In applying the triangle cost to a Tate component, choose L to be the prescribed local set above ℓ. The length lost by imposing vanishing there is bounded by the crystalline defect, since crystalline Tate cocycles vanish on that set. After imposing vanishing there, taking $z + z^c$ produces an even cocycle, and the triangle condition gives the required agreement of the conjugate pairs up to the factor D. Once all new inertia values vanish the cocycle is unramified outside ℓ. Its odd part $z^c - z$ now provides an element of the odd module, and the even part, by the triangle condition, is a coboundary up to a factor of controlled valuation. The assumption on coboundaries makes the kernel of this odd projection killed by D; in particular its length modulo D-torsion is bounded by the length of the odd module. The assertion is `hasTriangleCost_tateTwist` in `New/Selmer/TateTwistSteps.lean`; it treats $e = 0$ directly. Thus b_e is the local defect plus the length in Proposition 40.16, without any even-class length term.

40.11 Hecke operators realizing the first-order relations

The formal-source proof now passes from the group algebra to forms. For a fixed auxiliary pair, label tuple and twist, choose the base space at the corresponding level and consider its enlargements at the Taylor–Wiles places. On each of these finite-dimensional spaces there is a finite set of eigensystems, and a closed integral Hecke algebra of scalar functions of the eigensystem preserving the lattice of Weyl-integral forms. The characteristic polynomial coefficients of the normalized Galois representations, as functions of the eigensystem, belong to this algebra: the attached representation comparison provides them as limits of integral Hecke coefficients on generic split Frobenius data. In particular $\mathrm{tr}(P(U_y(h))U_y(g))$ belongs to the algebra for every integral polynomial P, by expanding P and using the trace at each $h^i g$. This is `BaseLevel.charpoly_mem_heckeAlgebra` and `BaseLevel.traceElem` in `New/Patching/OnePrimeTie.lean`, with the closedness and local–global compatibility of the Hecke algebra taken as formal-source inputs.

A system is called weakly lifted for this step if every $U_y(g)$ has eigenvalue one; call this set $\mathfrak{W}_1$. It is potentially larger than $\mathfrak{X}$. A normalizer can be chosen as in (40.20) so that $\beta_y \neq 0$ off $\mathfrak{W}_1$, but the argument does not try to separate $\mathfrak{W}_1 \setminus \mathfrak{X}$ by this determinant. A further

polynomial $P(\beta)$ over the ℓ-adic integers is chosen to vanish at all the nonzero determinant values on the old eigensystems and to have nonzero constant term (take the product of their linear factors, with integral scalar factors as necessary). Thus $P(\beta) \neq 0$ on $\mathfrak{W}_1$ and vanishes on all other old systems. It is not asserted that the coefficients of P or the divisions needed in clearing this cutoff are units.

Proposition 40.19 (Formal-source input: Hecke string)**.** *For a weakly lifted eigensystem of an auxiliary pair, its rescaled unramified Hecke polynomial at a generic split place of prime norm q_w has the full string of roots $1, q_w, \dots, q_w^{t'-1}$. More precisely, if its normalized Galois characteristic polynomial at a generic Frobenius is congruent coefficientwise, modulo the maximal ideal, to*

$$S(X) \prod_{i<t'} (X - \epsilon_\ell(\phi)^{-i}) \quad \textit{with } |S(1)|_\ell \geq 1,$$

then its rescaled Hecke polynomial factors as the corresponding full string times a residual factor dividing an integral polynomial $e(X)$ with $e(1) = 1$.

This is `isRoot_hecke_string` and `eventually_heckePoly_eq_string_mul` in `New/Selmer/HeckeString.lean`. The underlying representation-theoretic input is the string lemma for spherical unitary representations and the prescribed local type. The integral polynomial assertion follows from comparing monic string factors modulo the maximal ideal and from the bound $|S(1)|_\ell \geq 1$: for a monic B dividing the string, the Gauss norm of a product with B equals the product of the norms; hence comparison of AB and $A'B$ forces comparison of A, A'. Once the residual factor A is integral and $A(1)$ a unit, use $e = A/A(1)$. This is the elementary part of `exists_integral_multiple_eval_one`. For finitely many relevant eigensystems one takes the product of the resulting polynomials, still with $e(1) = 1$.

The required Frobenius conditions in Proposition 40.17 ensure that this proposition can be applied at every chosen auxiliary prime simultaneously. A simple reason such conditions are available is that at the distinguished row block a scalar inertial element with long prime-to-ℓ order allows one to move a Frobenius condition without changing its cyclotomic value. If an integral matrix A has rank r', the polynomial in Y given by $\det(1 - YA)$ has degree at most r' and constant term one. Among $r' + 1$ distinct residual scalars Y, at least one avoids its roots. The scalar-inertia condition of the row provides such scalars even after taking large ℓ-powers of the inertial element. By continuity the nonvanishing is a neighborhood condition. The formal statement is `exists_eventually_no_eigenvalue_one`. The attached representations and the generic Frobenius filter then yield the simultaneous string conditions for the eigensystems congenial to the target residual system. This is the condition imposed in the choice of the triangular primes in `denominatorLe_rowBudget`.

To realize the relations (40.21), at a Taylor–Wiles place let $\Gamma = \mathrm{GL}_{r'+t'}(k_w)$ and $H \subseteq \Gamma$ the stabilizer of a line. The character of H to $\mathbb{Z}/\ell^\nu$ given by the logarithm of its scalar action on the line is a diamond-operator character. Its transfer to Γ factors through the determinant. The relevant normalization, after multiplying by $(r' + t')!$, sends a chosen primitive determinant scalar to one modulo the logarithmic modulus. This follows from the transfer identity on scalar matrices and the count of lines; it is `exists_line_stabilizer_log` in `New/Patching/OnePrime.lean`. The proof then constructs a first-order rather than a semisimple deformation of the trace. At one place, if ρ denotes the residue-group action, define for a function f_0 on the logarithmic group

$$W(f_0) = \sum_{h \in H} f_0(\log h)\rho(h).$$

Changing h to $g^{-1}h$ or shifting the logarithmic parameter shows that multiplying by a diamond element shifts the argument of f_0. Taking a representative of the logarithm gives

the infinitesimal version of this identity modulo ℓ^ν. The source performs the induced-representation transfer, using coset representatives for Γ/H, to construct a local operator F_j and the norm-sum operator $N_j = \sum_{\gamma\in\Gamma} \rho(\gamma)$, with the trace relations acting as follows: if $t_j(g) = \mathrm{tr}((U(\tau_j) - 1)U(g))$ and $r_j(g) = \mathrm{tr}((U(\phi_j) - 1)e_j(U(\phi_j))U(g))$, then, modulo ℓ^ν and up to the specified integer factors, $F_j t_j(g) = N_j r_j(g)$ and $N_j t_j(g) = 0$. There are additional identities controlling how F_j moves the old-space invariants. This is the formal-source input `BaseLevel.exists_tie` in `New/Patching/OnePrimeTie.lean`, proved using `one_prime_first_order` in `New/Patching/OnePrime.lean` and local line-stabilizer operators. The nonzero factorial factor is retained rather than divided out at the torsion level.

The multiple-prime coefficient extraction is motivated by the following simple polynomial observation. If operators F_j, N_j at distinct places commute, multiplying the first-order relation by products of operators at the other places transports the contribution from a monomial X^d to the monomial X^{d+e_j}. Terms with exponents larger than one are discarded. The product of the sums N_j is the full residue-group norm mapping to the old-level subspace, while the complementary product of the F_j probes its infinitesimal deformations. Thus evaluating the trace-ideal relations on this first-order coefficient system annihilates the trace ideal modulo ℓ^ν. On the distinguished systems, all coefficients of F in (40.22) properly dividing $\prod X_j$ vanish; only the top coefficient remains. The source proves that this coefficient gives an integral diagonal operator at old level, localized to the cutoff determined by $P(\beta)$, after controlling the movement from the old space. The auxiliary depth ν must exceed the valuation of an explicit factor involving the constant term of P, the level automorphism factors and the factorial. It can be made arbitrarily large since the index set of the triangle was bounded before ν was chosen.

Precisely the result needed is `BaseLevel.exists_diagOp_mem_ellIntegralForms` in `New/Patching/OnePrimeTie.lean`, a formal-source input. Given the string condition near the distinguished residual eigensystem, enough divisibility by ℓ^ν, and a Ribet element of top-coefficient valuation with a factor M_0 at most Γ_0, it supplies a scalar function s of the eigensystem with $v_\ell(M_0 s(y_0)) \le \Gamma_0$. The diagonal operator with scalars s sends a vector supported on $\mathfrak{X}$ to an integral vector up to a specified level factor and any original denominator, provided that vector agrees with an integral old form on the eigensystems where $P(\beta) \neq 0$ in the sense of the diagonal-operator test in the source statement. Thus the integral output does not require that all weakly lifted systems themselves belong to $\mathfrak{X}$. A tested form minus its projection to Z is orthogonal to the unwanted weakly lifted components by the Hecke string assertion and (40.1), after inserting the chosen label projector. It follows that the diagonal operators satisfy the required integrality hypothesis on the projected tested forms.

40.12 The denominator bound from the local length budgets

We now combine the congruence terms numerically. Write b_1 for the rounded value of the thick weight and let

$$F_\ell^{\mathrm{num}} = (2b_1)! \left(\prod_{\max(b_1 - 30h_2^3, 0) < i \le b_1 + 30h_2^3} i \right)^{h_2^3}.$$

This integer is independent of ℓ; the notation emphasizes its use in the local estimate. With $g = g_0 g_1$, define

$$\mathrm{loss}_\ell = 10^6 \frac{m}{k} \left(b_1 v_\ell(g) + v_\ell(F_\ell^{\mathrm{num}}) \right), \qquad D_\ell = h_2^{27} \left(c_\ell + v_\ell(X) \right) + 4f \, \mathrm{loss}_\ell .$$

Here X is from Proposition 40.16. These are `loss`, `lambdaLoss` in `New/Selmer/DeterminantBound.lean`, and `rowBudget` in `New/Selmer/RibetElement.lean`. The detailed local comparison of torsion cocycles and the crystalline relations is developed in the next part of the proof. Its crucial formal-source bound, `sum_blockLoss_le` in `New/Selmer/RibetElement.lean`, is

$$\sum_{i \text{ blocks}} \lambda(\text{loss of crystalline relations on block } i) \le 4r' f \operatorname{loss}_\ell, \tag{40.24}$$

for arbitrary torsion depths of the individual blocks. This controls the failure of extracted cocycles to satisfy the crystalline local conditions in the Selmer theorem. One cannot apply that theorem directly to all cocycles extracted from congruent matrix differences.

Proposition 40.20 (Formal-source input: arithmetic Ribet element)**.** *Use the apt columns and rows of an auxiliary pair and a twist satisfying* (40.16) *and* (40.17) *at a depth $M > \lfloor rD_\ell \rfloor$. Suppose their normalized matrices satisfy* (40.3) *on the distinguished systems, and for new twist inertia the normalized columns satisfy $(U(g) - \psi(g))(U(g) - 1) = 0$. At a triangular auxiliary level, with the trace ideal as above, the determinant from* (40.20) *has a Ribet element with top coefficient c_0 such that*

$$v_\ell(M_g^3 c_0) \le r' D_\ell.$$

This is `ribet_columns` in `New/Selmer/RibetElement.lean`. The required quadratic inertia relation is `normalisedRep_quadratic_on_inertia` in `New/Selmer/LabelledColumns.lean`; it follows from the shape of the parahoric type at the twist conductor. The cocycle length for the row block is at most its normalized Selmer bound $(10fr' + 1)c_\ell$ plus its share in (40.24). The Tate costs in (40.23) contribute at most $v_\ell(X)$ per block plus their shares in (40.24). The scalar-separation and detecting terms contribute multiples of $h_2 h$ and $h_2^{22} h$; the old and new tame polynomial factors have valuations bounded by the number of places times c_ℓ and by the sum in (40.16). The aggregate of these noncrystalline terms, including the factor M_g^3, fits inside $r' h_2^{27}(c_\ell + v_\ell(X))$ by the admissible size inequalities. The source proves the explicit inequality in the last step of `ribet_columns`. Notice why the row modulus M is chosen larger than rD_ℓ: it is larger than the resulting bound on the row-cocycle length, as required in the first-order Ribet lemma.

We give the final linear algebra of combining individual integral operators, since it avoids multiplying the denominator cost by the number of distinct eigensystems. Let $\{p_i\}_{i=1}^q$ be a basis of Z consisting of nonzero vectors from the spanning family of lifted forms. Given integral test vectors u_a, v_b in the denominator condition, write $P_Z v_b = \sum_i P_{ib} p_i$. Suppose for each j a diagonal comparison operator acts on p_i by scalars a_{ji}, after a common injective comparison map, with $|M_0 \iota(a_{jj})|_\ell \ge e^{-\Gamma}$, and that the matrices

$$(\iota\langle u_a, \textstyle\sum_i a_{ji} P_{ib} p_i\rangle M_0)_{a,b}$$

are integral for every j. Choose coefficients t_j in the integral valuation ring of $\overline{\mathbb{Q}}_\ell$ so that for every i,

$$|M_0 \sum_j t_j \iota(a_{ji})|_\ell \ge |M_0 \iota(a_{ii})|_\ell \ge e^{-\Gamma}.$$

This is possible since for each i the finite linear polynomial has its Gauss norm attained at some tuple of units, and a finite set of nonzero linear polynomials over the infinite residue field can be avoided simultaneously after dividing each one by a coefficient of maximal absolute value. The linear combination of the integral pairing matrices is integral. Its determinant equals the determinant of the matrix $(\iota\langle u_a, P_Z v_b\rangle)$ multiplied by $\prod_i (M_0 \sum_j t_j \iota(a_{ji}))$, by writing all pairing matrices as a product of the basis-change matrices and the relevant diagonal matrices. Consequently the absolute value of the desired determinant is at most $e^{q\Gamma}$. This is the linear algebra in `norm_gram_det_le_of_operators` in `New/Selmer/DeterminantBound.lean`.

Apply the arithmetic Ribet element proposition and the first-order trace deformation over the triangle to a single basis eigensystem in the space of lifts. In the setting of the tested forms, the local operator integrality described above, together with the class-sum pairing and automorphism bounds absorbed into M_g^3, provides exactly the operators for this linear algebra with $\Gamma = r' D_\ell \log \ell$. This yields

$$\mathrm{DenominatorLe}(\iota, Z, D_\ell \log \ell), \tag{40.25}$$

that is (40.2) with $\beta = D_\ell \log \ell$. The precise formal-source theorem, including the spanning hypothesis that each generator is apt, lifts a fixed system having a row for all twists, and has the needed Selmer bound, is `denominatorLe_rowBudget` in `New/Selmer/DenominatorBoun d.lean`. The use of tested forms, not arbitrary forms, is why the weakly lifted eigensystems suffice. Equation (40.25) marks the end of the congruence argument.

40.13 Global numerical summation and proof of the fit theorem

All the terms in the local allowances are nonnegative, and the factorization of each positive integer A gives $\sum_{\ell \in T} v_\ell(A) \log \ell \le \log A$ for every finite set of primes T. The other part of c_ℓ has bounded sum, since $\sum_{n \ge 2} 2^{-n} \log n < \infty$. Therefore

$$\sum_{\ell \in T} 9 c_\ell \log \ell \le 9 (\log M_g + \sum_{n \ge 2} 2^{-n} \log n),$$

$$\sum_{\ell \in T} D_\ell \log \ell \le h_2^{27} \left(\log M_g + \log X + \sum_{n \ge 2} 2^{-n} \log n \right) + 4 \cdot 10^6 \frac{fm}{k} \left(b_1 \log g + \log(2b_1)! + h_2^3 \sum_{\max(b_1 - 30h_2^3, 0) < i \le b_1 + 30h_2^3} \log i \right).$$

These estimates reduce the infinite collection of local lengths to a handful of ordinary integer sizes. For example $\log(2b_1)! \le 2b_1 \log(2b_1)$, and the short-interval contribution is at most $60 h_2^6 \log(b_1 + 30h_2^3)$. Also $\log M_g$ is bounded by a constant times $f(rt)^2 \log g$ by the order formula for finite general linear groups, and $\log X$ is bounded by $4(N_E + t + 3)^{27} \log 2 + \log h_E$. The explicit admissibility inequalities in the source bound these contributions by

$$\sum_{\ell \in T} 9 c_\ell \log \ell \le x^{999}, \qquad \sum_{\ell \in T} D_\ell \log \ell \le B_0. \tag{40.26}$$

See the numerical calculation in `exists_isFitFamily` in `New/Selmer/FitTwistsExist.lean`, using the class-number estimate and `Data.le_of_hasLogBound`. These are formal-source bounds with the full numerical hypotheses on the datum. To see their scale, along the admissible family with fixed $\varepsilon > 0$, b exceeds every fixed power of x; h_2, N_E, r and t are polynomial in x. Thus the first term of the second bound is smaller than B_0 by a power of b. In the remaining factorial term b_1 has order b, $\log g$ and $\log b_1$ have order $\log b$, and the leading contribution is $O(fmb \log b/k)$. Its ratio to $B_0 = fa(\log b)^3/(mk)$ is $O(m^2 b/(a(\log b)^2)) = O((\log b)^{-2})$ by $m^2 = O(x)$ and $a \ge xb$. This highlights the logarithmic slack without replacing the exact inequalities by asymptotic claims.

Proof of Theorem 40.2, using the stated formal-source inputs. For each ℓ and ι, make the finite list of rows in Proposition 40.3, including all rank-one pairs, before choosing any twists. Apply Corollary 40.15 to this list at depth $M = \lfloor r D_\ell \rfloor + 1$. The resulting m-fold twist θ_ι has conductor prime to ℓ and satisfies the local conditions for its first m powers. At each arrow j,

the good lift is spanned by the apt systems with the chosen labels and this common Selmer bound. For the rank-one lines the corresponding assertion follows from their apt label $X - 1$ and the rank-one row. Consequently (40.25) applies simultaneously to the desired spaces, with the same allowance $\beta_{j,\ell} = D_\ell \log \ell$ independent even of j. The second inequality of (40.26) is its required finite-sum bound. The special-value inequality (40.18) and the first inequality of (40.26) give the required height bound on the inverse ratios: in the height definition take the constant archimedean factor 1 and the local log allowance $9c_\ell \log \ell$. All these choices are independent between the pairs (ℓ, ι) but uniform over the finite lists for a given pair, as required. Thus the family satisfies Definition 40.1. This reproduces the final assembly of `exists_isFitFamily`. □

The upper bound on projection denominator and the lower bound on theta-lift covolume now meet at exactly the same error budget. The global Selmer argument has been used to produce the local denominator estimate, not to establish a vanishing theorem for a class group, a modularity lifting theorem, or a nonvanishing assertion about automorphic L-functions at critical points. In particular the omission of any even-class-group length term is a consequence of the first-order ties at the auxiliary triangle, and not of assuming those classes vanish.

Chapter 41

The foundations of the congruence argument

The preceding congruence chapter explains how integral projection determinants reduce to a family of bounds on cocycles. In the present chapter we develop the algebraic and arithmetic foundations of that reduction. We isolate the precise places at which the method uses global Galois representations, reciprocity, the construction of auxiliary levels and the extraction of a first-order Hecke relation. The point is not to introduce more abstract hypotheses than the construction needs. Rather, it is to understand why the particular auxiliary data of the proof are effective in the congruence problem: why the primitive labels at two distinct primes are useful, why anticyclotomic twists have been distinguished from triangular auxiliary primes, and how one can account for congruences of Tate characters without a bound for their even classes. We refer to the preceding chapter for the statements of the fit-family and determinant estimates. In particular r' denotes the rank of the distinguished row block, t' the number of Tate characters, and neither is the dimension of the automorphic form space. We use v_ℓ normalized by $v_\ell(\ell) = 1$; in intermediate algebraic arguments v_π denotes the integer-valued valuation of a DVR with uniformizer π. The normalized lengths of finite modules are related to their ordinary lengths by the ramification index, exactly as in Section 40.1. All claims imported from the formal proof without a reconstructed argument are marked as such.

41.1 From weak Hecke compatibility to a trace identity

The version of Galois attachment in the compact development is weaker in one important respect than a usual local-global compatibility theorem. Rather than specifying the complete local Weil–Deligne representation at every place, it specifies a convergence condition at the set of Frobenius data and polynomial annihilation conditions at individual places. We explain the distinction because an argument using the full Weil–Deligne parameter from the outset may inadvertently assume an assertion that is substantially stronger than the one provided in the code.

Let $G_E = \mathrm{Gal}(\overline{E}/E)$. A Frobenius datum comprises a degree-one place w, a choice of a place above it in the algebraic closure, and a Frobenius element of its decomposition group; all finite exceptional sets of places can be omitted. The source has a filter on these data, denoted `mostFrobenii`. It also has its localization near $g \in G_E$, denoted `mostFrobeniiAt`; the latter includes arbitrarily small neighborhoods of g in the Frobenius variable. This formalizes a useful way of applying Chebotarev. If $F(g)$ is continuous, then as Frobenius data tend to g the values of F at the corresponding Frobenius elements tend to $F(g)$. One should not replace this filter with an unspecified density-one subset: a density statement needs to be

strong enough that after excluding finitely many places there are still Frobenius data in *every* permissible neighborhood. The code derives the relevant properties from its Chebotarev package in `Lib/ClassField/Chebotarev/FrobeniusFilter.lean`.

More precisely, if $P_w(X)$ is a family of monic polynomials of degree n over $\mathbb{C}$, a representation $R : G_E \to \mathrm{GL}_n(\overline{\mathbb{Q}}_\ell)$ is attached to P in the sense used here if it is continuous, trace-nondegenerate and, for every $\eta > 0$, eventually along the Frobenius data one has

$$\left|[X^i]\left(\det(X - R(g_w)) - \iota P_w(X)\right)\right|_\ell \le \eta \qquad (0 \le i \le n). \tag{41.1}$$

The more general definition permits an open group map in place of the identity on G_E, and requires the estimate for every element mapping to the chosen Frobenius. See `Attached` in `Lib/GaloisRep/Main/AttachedDefs.lean`. In these notes trace-nondegenerate means that the trace pairing on the finite-dimensional algebra spanned by the matrices of the representation is nondegenerate. We use the topology of the ℓ-adic field, not the Zariski topology, in (41.1).

Lemma 41.1 (Trace extraction from attachment)**.** *Let R be attached to P as in (41.1). For every $g \in G_E$, the functions $-\iota[X^{n-1}]P_w$ tend, along the Frobenius data localized at g, to $\operatorname{tr} R(g)$. Consequently, if two attached systems have polynomials related eventually by multiplication by a product of known linear factors, and the coefficients of those factors tend to prescribed characters at g, their traces differ by the corresponding sum of characters. The same assertion holds with a rescaling of the roots by the values of one continuous character.*

Proof. The identity $[X^{n-1}]\det(X - A) = -\operatorname{tr} A$ holds for every square matrix. For each positive η the non-Archimedean triangle inequality bounds the discrepancy from $\operatorname{tr} R(g)$ by the maximum of the error in (41.1) and $|\operatorname{tr}(R(g_w) - R(g))|_\ell$. The former is eventually at most η by hypothesis, the latter tends to zero by continuity. The rest follows because the negative next-to-leading coefficient of a product of monic polynomials is the sum of their negative next-to-leading coefficients. Scaling all roots of a polynomial by u multiplies this coefficient by u. Limit uniqueness proves the asserted relations at every g. □

This elementary argument is the passage from Hecke polynomials to the identity of traces used in `exists_row_for_all_twists`. It does not imply that at a fixed exceptional place every particular Hecke polynomial is necessarily the characteristic polynomial of Frobenius. The source instead uses the following separate assertion.

Proposition 41.2 (Formal-source input: spherical compatibility)**.** *For the standard definite unitary group over the chosen field with its Weil data, every nonzero Hecke eigenform of rank n has an attached representation. If the form is spherical at a place w, any attached representation is unramified off ℓ, respectively crystalline above ℓ, at the corresponding decomposition group; it has the jumps prescribed by the weight, and every specified Hecke polynomial at w, transported by ι, annihilates the specified geometric Frobenius action (in the local polynomial sense of the source).*

The exact statement is `sphericalCompat_std` in `Lib/GaloisRep/Main/SphericalAllRanks.lean`, with `SphericalCompat` defined in `Lib/GaloisRep/Main/AttachedDefs.lean`. It includes the local jumps `jumps`, the predicate `IsGoodAt`, and the polynomial predicate `PolyKillsAt`. Its proof uses the construction for odd rank, the theory of theta lifts, and local comparison statements. We retain these geometric and local comparison assertions as formal-source inputs. A considerable algebraic part of the reduction from odd rank to general rank, however, can be reconstructed directly as follows.

Lemma 41.3 (Trace-nondegenerate representations and cancellation)**.** *Over an algebraically closed field L of characteristic zero, a finite-dimensional representation whose trace pairing on its image algebra is nondegenerate is semisimple. Its isomorphism type is determined by its trace function on the group. Suppose R_0, R_1 are such representations and C_0, C_1 are finite direct sums of characters, with no character occurring in both C_0 and C_1. If*

$$\operatorname{tr} R_0 - \operatorname{tr} C_0 = \operatorname{tr} R_1 - \operatorname{tr} C_1,$$

then there is a semisimple representation R such that $R_i \simeq R \oplus C_i$ for $i = 0, 1$. If the initial representations are continuous, R can be chosen continuous.

Proof. Write $A \subseteq \operatorname{End}_L(V)$ for the image of the group algebra. The radical J of the finite-dimensional algebra A is nilpotent. For any $a \in A$ and $j \in J$, the element aj is nilpotent and its trace in V is zero. Thus J lies in the radical of the trace form. Nondegeneracy forces $J = 0$. The algebra is therefore a product of full matrix algebras and V is a direct sum of simple A-modules. In the converse direction the trace form on an image algebra of this type is the direct sum of the matrix trace forms multiplied by the corresponding positive multiplicities; it is nondegenerate in characteristic zero.

For two semisimple representations take the image of the group algebra on their direct sum. It too is semisimple: the direct sum is semisimple and faithful for that algebra, so its radical acts both trivially and faithfully, whence is zero. Let the simple factors be $M_{d_i}(L)$. The central idempotent of the ith factor acts as the identity on every copy of the corresponding simple representation and as zero on the others. Its trace divided by d_i recovers the multiplicity. A trace identity on the group extends linearly to the group algebra and hence to all these idempotents. This proves determination by trace.

Now compare the multiplicities in $R_0 \oplus C_1$ and $R_1 \oplus C_0$. For a character occurring in C_0, its multiplicity in R_0 is at least that in C_0, since its multiplicity in C_1 is zero. The corresponding conclusion holds for C_1. All other multiplicities in R_0 and R_1 agree. Subtract the corresponding sums of characters and take the common semisimple remainder. As a direct summand of a continuous representation on a finite-dimensional valued vector space it is continuous (the coordinates of the action in a fixed basis are obtained by continuous matrix operations). □

The cancellation in the source is `IsTraceNondeg.exists_cancel_chars`. It is applied to two auxiliary theta lifts of odd target rank. To orient this construction, if the original rank is n, put $u = n + 1$, so the auxiliary rank is $n + u = 2n + 1$. Choose two distinct unramified characters of $U(1)$ whose infinity weights differ by a nonzero multiple of the torsion order of E at one place. Associated to either choice is a string of u algebraic characters whose Frobenius values at split places supply the extra factors in the Hecke polynomial of the theta lift. Distinct weight choices are arranged so that the two strings share no character. The construction is the first part of the proof of `sphericalCompat_std`. The character existence and the nonzero theta lifts of suitable joint-harmonic weight are formal-source inputs there.

The source establishes that at every split spherical place the extra factors $Q_{i,w}$ in the two lifts are pairwise disjoint between strings. The resulting algebraic check is useful independently of the global construction.

Lemma 41.4 (Two cancellations of auxiliary factors)**.** *Suppose R_i are as in Lemma 41.3 with common remainder R, and let P_w be the sought Hecke polynomial. Assume at a given place, after the normalization and removal of the known characters, the polynomial operators $P_w(F)Q_{i,w}(F)$ vanish on the relevant local space of R for both i. If $Q_{0,w}$ and $Q_{1,w}$ are coprime, then $P_w(F)$ vanishes there. Furthermore, if $P_w Q_{i,w}$ are monic polynomials with coefficients in a valuation ring and $Q_{i,w}$ has coefficients in this ring, then so does P_w.*

Proof. Choose A, B with $AQ_{0,w} + BQ_{1,w} = 1$ and substitute F; all polynomials in F commute. Multiply the identity by $P_w(F)$. For the integral assertion, successively compare the highest coefficients of $P_w Q_{i,w}$. The coefficient of each degree in P_w is expressed as one integral coefficient of the product minus integral combinations of already determined higher coefficients. No division is needed because the divisor is monic. Equivalently use the multiplicativity of the non-Archimedean Gauss norm and the fact that a monic integral divisor has Gauss norm one. □

This is the final polynomial step in `sphericalCompat_std`: both lift constructions have the same trace remainder and each yields the required local comparison after introducing its own extra polynomial. The local comparison itself, including passage of the crystalline conditions and jumps through the trace limit, is the formal-source input `local_clauses_of_trace_limit` used there. This explains why merely quoting an odd-rank construction and deleting extra characters from a global trace identity would not be sufficient.

41.2 Tame labels and full matrix span

We now turn to the local property of the Galois block on which the Selmer argument relies. The proof requires the strong integral statement $\mathrm{span}_{\mathcal{O}} R(G_E) = M_{r'}(\mathcal{O})$ for every sufficiently large coefficient ring. Irreducibility only over its fraction field is insufficient. The two primes g_0, g_1 and their primitive labels are specifically chosen for this purpose.

Lemma 41.5 (Metacyclic tame-pair lemma)**.** *Let $\mathcal{O}$ be a DVR with fraction field L and let $R : G \to \mathrm{GL}_n(\mathcal{O})$ be a representation. Suppose $s, f \in G$ and an integer $q \geq 2$ satisfy*

$$R(f)^{-1} R(s) R(f) = R(s)^q, \qquad R(s)^{q^n} = R(s).$$

Suppose an eigenvalue $\zeta \in \mathcal{O}$ of $R(s)$ is a unit and the n numbers ζ^{q^i} $(0 \leq i < n)$ are pairwise distinct modulo the maximal ideal. Then n^2 matrices in the image span $M_n(\mathcal{O})$ over $\mathcal{O}$, and $R(f^n)$ is a scalar unit.

Proof. Reduce the matrices modulo the maximal ideal. Let $A = \overline{R(s)}$, $B = \overline{R(f)}$, and choose a nonzero vector v with $Av = \bar{\zeta} v$. The relation implies that $B^i v$ has eigenvalue $\bar{\zeta}^{q^i}$ for A. Consequently the vectors $v, Bv, \ldots, B^{n-1}v$ form a basis of the residue vector space. Over it the matrix A is diagonal with pairwise distinct entries. Its spectral projectors E_i are polynomials of degree $< n$ in A, by Lagrange interpolation. The matrix B cyclically moves the eigenspaces, with nonzero scalar at the wraparound, since it is invertible. Hence $B^j E_i$, $0 \leq j < n$, $0 \leq i < n$, are nonzero multiples of all matrix units in the given basis. Each is a linear combination of $B^j A^k$ for $0 \leq k < n$. There are n^2 such matrices, so their lifts $R(f^j s^k)$ reduce to a basis. Nakayama's lemma, or the determinant criterion over the local ring, implies these lifts form an $\mathcal{O}$-basis of the entire matrix algebra.

The second tame-pair relation gives $R(f)^n$ commuting with $R(s)$; it plainly commutes with $R(f)$. It therefore commutes with the $\mathcal{O}$-basis just exhibited. The centralizer of the entire matrix algebra in itself is its scalar center, as one sees by commuting successively with diagonal matrix units and off-diagonal matrix units. Thus $R(f)^n$ is scalar, and invertibility makes the scalar a unit. □

This proves the key statement `IsLatticeIn.span_eq_top_of_tame_pair` in `Lib/General/Groups/ResidualIrreducibility.lean`; the source takes f^{-n} rather than f^n for its scalar, which is equivalent. Notice that if the n eigenvalues were distinct merely in characteristic zero, the Lagrange denominators might have positive valuations and the resulting basis would

not be integral. In that case there would be no reason for the matrix span to be all matrices over $\mathcal{O}$.

We give the elementary number theory behind the special primitive-label property. Let $q \equiv 1 \pmod 9$, let $n = 3^h$ for $h \geq 1$ and put $n_0 = n/3$. The quotient

$$S = \frac{q^n - 1}{q^{n_0} - 1} = q^{2n_0} + q^{n_0} + 1$$

is divisible by 3 exactly once: reducing q^{n_0} modulo 9 gives $1 + 1 + 1 = 3 \pmod 9$. Moreover $S > 3$, so it has some prime factor other than 3. It follows that S is not a power of any single prime. Suppose ζ has order $q^n - 1$ and fix any prime ℓ. The order of ζ^{ℓ^a} is $(q^n - 1)/\gcd(q^n - 1, \ell^a)$. If its q-orbit had period $< n$, that period, a proper divisor of $n = 3^h$, would divide n_0. Then $q^n - 1 \mid \ell^a(q^{n_0} - 1)$, forcing $S \mid \ell^a$, which is impossible. Thus all n powers $(\zeta^{\ell^a})^{q^i}$ are distinct for every $a \geq 0$. By the characterization of reduction of roots of unity from the preceding chapter, all the ζ^{q^i} are residually distinct. This is `frobenius_conj_pow_injOn` in `Lib/LocalGL/Green/LabelIdempotents.lean`. The case $n = 1$ requires no orbit argument.

The same argument applies when the primitive element is multiplied by an element z of the base residue field of cardinality q. Indeed in the ratio of the ith and jth q-conjugates the base-field factor cancels. This is precisely the scaling of the label by the inverse of the tame part of the automorphic character in `twistedLabel`. In particular the proof uses the *primitive original label*, not a claim that its scaled version is still primitive. If the order of a primitive element is $q^n - 1$ and u has order dividing M_ω in the base field, the product of the n conjugates of the scaled element is the norm of the primitive element times u^n. Its norm before scaling is primitive of order $q - 1$. The order of the scaled product is at least $(q-1)/M_\omega$ in the divisibility sense: if its jth power equals one, then $q - 1 \mid M_\omega j$. This follows by raising the norm identity to M_ω. The source uses the fact that $q - 1$ is divisible by M_ω^2, and $M_\omega = (N_0 d)^{s+6}$ has more than one prime divisor. Choose any prime divisor $p \mid N_0 d$ with $p \neq \ell$; the p-part of the norm of the scaled label then still has order at least p^{s+6}. This grows faster than the permitted small powers $4r' + 2$. A corresponding power of the inertia generator acts by a scalar on the row, while all those small powers of that scalar remain distinct from one after reduction. This is the mechanism of `IsLabelledColumn.isRow`; below we specify the polynomial input needed to justify passing from labels to the tame generator.

Lemma 41.6 (Valuation of a nontrivial root of unity)**.** *If $\lambda \neq 1$ and $\lambda^D = 1$ in $\overline{\mathbb{Q}}_\ell$, then $v_\ell(1 - \lambda) \leq v_\ell(D)$. If a matrix A of size n is annihilated by a polynomial all of whose roots differ from one and whose roots satisfy this condition with a common D, then*

$$v_\ell(\det(A - 1)) \leq n\, v_\ell(D).$$

Proof. Every root of unity is a unit. Since $\sum_{i=0}^{D-1} \lambda^i = 0$, we have

$$D = \sum_{i=0}^{D-1} (1 - \lambda^i) = (1 - \lambda) \sum_{i=1}^{D-1} (1 + \lambda + \cdots + \lambda^{i-1}).$$

The last factor is integral. This proves the first assertion. Every eigenvalue of a matrix is a root of every polynomial that annihilates it, by evaluating that polynomial on the eigenspace. The determinant of $A - 1$ is the product of its eigenvalues minus one, with multiplicities, and the sum of their valuations gives the claimed bound. □

The common integer needed in the determinant bound can in fact be chosen less crudely than $q^{n!} - 1$. The source allows each root of an inertia annihilator to have order dividing $g_i^e - 1$

for some $1 \le e \le r$. Each such integer divides M_g (the group order defined in (40.4)). Thus $v_\ell(1-\lambda) \le v_\ell(M_g)$ separately for each root, and the same proof gives the bound $r'c_\ell$ of the preceding chapter. If $q = g_i \neq \ell$, a tame generator at a place above q generates the required finite cyclic action away from its wild inertia. At a prime above g_i the row eigenvalues are the roots of the scaled label transported by the residue-field character. The other chosen prime can be used if $g_i = \ell$. Their independent presence is the reason a residual irreducibility conclusion holds for every ℓ, even when ℓ is one of the two large level primes.

Proposition 41.7 (Formal-source input: local labels)**.** *For an apt pair at the chosen primes above g_i, each element of inertia has a common annihilating polynomial on the labelled columns and their conjugate inverses. The polynomial is the product of $r'+t'$ linear factors $X-\lambda$; each root obeys $\lambda^{g_i^{(r')!}-1} = 1$ and $\lambda^{g_i^e-1} = 1$ for some $1 \le e \le r$. At the distinguished generator the roots for $t' = 0$ are nontrivial. At the twist primes outside the original level, the normalized representation is annihilated on inertia by $(X-\psi(g))(X-1)$, where ψ is the inverse ℓ-adic avatar of the new twisting character. Twisting by the character needed to normalize a row multiplies the corresponding inertia labels and shifts the Hodge jumps by its local weights.*

The assertions are respectively `exists_inertia_eigenvalues`, `normalisedRep_quadratic_on_inertia`, and `IsLabelledColumn.twist_avatar`, in `New/Selmer/LabelledColumns.lean`. These use the parahoric local-global comparison `Lib/GaloisRep/LocalGlobal/LabelCompatibility.lean`. The latter supplies the polynomial killing conditions, not an assertion that a characteristic polynomial computed by a global trace uniquely determines a local monodromy operator. Subject to this input, Lemma 41.5 and the primitive-root and determinant calculations give the full matrix-span and small inertia determinants of `IsRow`. The remaining row conditions are the jump bounds and conjugate self-duality; these enter via Proposition 41.2 and the particular shifts of the algebraic weight in `exists_hasRow_auxPair` of `New/Selmer/RowsExist.lean`.

41.3 A duality calculation at the mirror

The role of the scalar inertia element is more subtle than simply making a representation irreducible. It is used a second time in an estimate for a family of additive functionals. We now prove the algebraic part of this estimate. It is important that these functionals are additive characters, not necessarily linear functionals over the coefficient ring.

Write c for the conjugation involution of the group, so cg in the following lemma means the conjugate of the element g, not multiplication by a fixed element. Let $A = \mathcal{O}/I$ be a finite quotient of the coefficient DVR, let $M = A^n$, and let R and χ be the row representation and a scalar character on this module. The twist by χ occurs on the dual of the row, not on the row itself. This distinction accounts for the different factors in the mirror functional.

Lemma 41.8 (Mirror calculation)**.** *Suppose $H \subset U \subset G_E$ are normal conjugation-stable subgroups, with U acting trivially through R and χ. Let γ satisfy*

$$R(\gamma) = s\,1, \qquad R(c\gamma) = s^{-1}\,1, \qquad \chi(\gamma) = \chi(c\gamma) = 1,$$

where $s-1$ and $s+1$ are units. Suppose H contains $[g,\gamma]$ for all $g \in G_E$. Let $y : G_E \to M$ be an R-cocycle, with $y(\gamma) = 0$ and with $y(g_0)$ having a unit coordinate. Assume $R(G_E)$ spans $M_n(A)$. Let W be a finite additive family of maps $x : U \to M^\vee = \mathrm{Hom}_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})$ such that the maps are additive group homomorphisms and satisfy

$$x(kbk^{-1})(u) = x(b)(\chi(k)R(k)^{-1}u) \quad (b \in U).$$

Assume for a fixed $z \in U$ and all $h \in H$, $a \in A$, that

$$x(zh)(ay(zh)) = x(c(zh))(ay(c(zh))). \tag{41.2}$$

If there is an element g with $R(g) = \alpha\, 1$ and $R(cg)^{-1} = \alpha'\, 1$, put $D = \alpha - \chi(cg)\alpha'$. Then

$$|W| \le |M[D]| = |M/DM|.$$

Proof. On U the map y is additive. Write $\tau = [g_0, \gamma] = g_0 \gamma g_0^{-1} \gamma^{-1}$. Expanding the cocycle equation gives $y(\tau) = (1 - s)y(g_0)$, so it still has a unit coordinate. For $b, b' \in H$, take the alternating difference of (41.2) at bb', b, b' and the identity. The terms involving z disappear, and what remains is

$$\begin{aligned} A(a) + B(a) &= C(a) + D_1(a), \\ A(a) &= x(b)(ay(b')), \quad B(a) = x(b')(ay(b)), \\ C(a) &= x(cb)(ay(cb')), \quad D_1(a) = x(cb')(ay(cb)). \end{aligned}$$

The cocycle transform on H is $y(kbk^{-1}) = R(k)y(b)$, because $R(b) = 1$. Replace b by $\gamma b \gamma^{-1}$ and a by sa. The four terms become $A(a)$, $B(s^2 a)$, $C(s^2 a)$ and $D_1(a)$: here both s^{-1} factors on the dual side are essential. Comparing with the untransformed equality at $s^2 a$ yields

$$(A - D_1)((s^2 - 1)a) = 0.$$

This step uses additivity in a, not A-linearity of x. Since $s^2 - 1$ is a unit, it follows that $A(a) = D_1(a)$ for every a.

Take $b' = k\tau k^{-1}$. We have proved for each $b \in H$ the stronger identity

$$x(b)(aR(k)y(\tau)) = x(c\tau)(a\chi(ck)R(ck)^{-1} y(cb)). \tag{41.3}$$

Because one coordinate of $y(\tau)$ is a unit and the row matrices span the full matrix algebra, the elements $aR(k)y(\tau)$ generate M additively. Thus the single functional $x(c\tau)$ determines x on H. Taking $b = c\tau$ in (41.3) and comparing its values at k and kg shows

$$x(c\tau)((\alpha - \chi(cg)\alpha')aR(k)y(\tau)) = 0.$$

Indeed the left argument is multiplied by α, while the right one is multiplied by $\chi(cg)\alpha'$, and the identity at k permits us to move that scalar to the left. Full span now shows that $x(c\tau)$ vanishes on DM.

Finally, for $b \in U$, the map x on $[b, \gamma]$ equals $x(b) \circ (1 - s^{-1})$ by additivity and the transformation rule. Since the commutator is in H and $1 - s^{-1}$ is a unit, determination on H determines x on all of U. We have an injection of W into the additive dual of M/DM. The latter has the same size as M/DM by finite abelian duality. The kernel and cokernel of multiplication by D on M have equal cardinalities, proving the assertion. □

This proof is the mechanism of `card_le_torsion_of_isotropic` in `New/Selmer/LinePointPowers.lean`. The actual source formulates the equivariance and subgroup hypotheses with a family of points and line powers. Two features should be emphasized. First, the scalar of γ is required to differ from both 1 and -1 modulo the maximal ideal; the reason is the cross-difference, which has coefficient $s^2 - 1$. Second, a uniform estimate for the additive dual follows from a single scalar defect D, independent of the number of cocycles initially used to formulate W. If D is nonzero, its normalized length contribution is at most $nv_\ell(D)$, since the kernel of multiplication by D on $\mathcal{O}/\pi^u$ has ordinary length $\min(u, v_\pi(D))$. If $D = 0$ this observation yields no saving and cannot be used in a finite quantitative balance argument.

The global application has additional steps not implicit in this calculation. One must extract the additive dual family from a system of local restrictions, realize the mirror relations at split places, and choose a Frobenius point seeing a class that has not yet been detected. The formal-source input for these steps is `exists_split_primes_relations` in `Lib/ClassField/GreenbergWiles/SelmerRelations.lean` and the application `IsPencil.length_le_of_unseen_class` in `New/Selmer/TiltingLemma.lean`. The calculation just given explains why there is a defect term of size $nv_\ell(D)$ in that application. It is not a replacement for the global reciprocity theorem.

41.4 Anticyclotomic primes from Kummer extensions

An anticyclotomic auxiliary prime adds a character variable to the twists already considered. The requirement that the character be trivial at all the old places would be quite restrictive if one tried to choose a ray class character after fixing an arbitrary split prime. Instead the prime is selected to solve that restriction. This is an instance where the order of construction matters.

Here is a convenient class field formulation. For a finite conjugation-stable set T of finite places, the T-units are the elements of $E^\times$ which are units off T. For a new prime $P \notin T$, an additive character initially specified on the units at P extends to the idele class group, trivially on the components at T, if it kills the ambiguity given by the T-units and the remaining ideal classes. Killing the latter ambiguity is accomplished by enlarging T before constructing P. The class group of the ring of T-integers can be made trivial by adding finitely many outside primes. If another finite set $R \setminus T$ is excluded, one can still choose these primes outside R: every ideal class is represented by a prime outside any finite set, by the Hilbert class field and Chebotarev. The idele form of this observation is `exists_sup_idelesUnitOff_eq_top`, used in the following construction. We take the usual reciprocity and Kummer correspondence as inputs here.

Proposition 41.9 (Controlled auxiliary character)**.** *Fix a rational prime ℓ, a positive power Q of ℓ, a finite set S of places at which the character must be trivial, a finite set R, and z in the closed normal closure of inertia of places in R away from $S \cup cS$ and the primes above ℓ. For any neighborhood W of z in G_E there is a degree-one prime $P \notin R \cup cR \cup S \cup cS$ and an anticyclotomic character $h : G_E \to \mathbb{Z}/Q\mathbb{Z}$ such that $NP \equiv 1 \pmod{Q}$. There are a Frobenius lift f at P in W and a tame inertia generator τ at that place with*

$$h(f) = 0, \qquad h(\tau) = 1.$$

The character is trivial at S and at primes above ℓ, and unramified outside P, cP. One may also specify a finite set of previously chosen primes to be avoided by P.

Explanation of the construction. Enlarge $S \cup cS \cup \{v : v \mid \ell\}$ to a finite set T killing the ideal classes as above, but avoid $R \cup cR$ outside this prescribed union. Take all conjugates of the additional primes if necessary. The group of T-units is finitely generated, including its finite torsion group. Write $u_1, \dots, u_d$ for generators. Consider the finite extension obtained by adjoining the Qth roots of unity and the Qth roots of these generators (and take its normal closure if needed). Inertia at each allowed place of R fixes this extension: in residue characteristic prime to Q, adjoining a Qth root of a unit gives an unramified extension. This last fact follows locally by Hensel's lemma after passing to the finite unramified extension whose residue field contains all the needed roots. The normal closure of these inertia groups also fixes the extension. In particular z acts trivially on it.

Choose an open normal subgroup small enough that a Frobenius element in a specified coset of it lies in W and fixes the Kummer extension. Chebotarev supplies such a Frobenius element at P, with P of degree one and split over the real subfield; finitely many primes can be excluded. Splitting in the roots-of-unity extension gives $Q \mid NP - 1$. Every u_i becomes a Qth power modulo P. The residue multiplicative group is cyclic, so choose a logarithm into $\mathbb{Z}/Q\mathbb{Z}$ taking a fixed residue generator to 1. This logarithm vanishes on the residues of all T-units. Extend it to $E_P^\times$ by killing the chosen uniformizer.

On the ideles supported away from T on their unit factors, take the logarithm of the unit component at P, and zero on the other components. Since the class group of T-integers is trivial, every idele is represented, modulo $E^\times$ and the local factors at T, by such an idele. Any ambiguity in this representative comes from a T-unit, on which the logarithm vanishes. Thus the logarithm extends to the idele class group, and is continuous with finite image. Compose with global reciprocity to obtain h_0. The character $h = h_0 - h_0 \circ c$ is anticyclotomic. At P, the term from cP does not modify its inertial value; choose the tame generator with the logarithm 1, as specified by the reciprocity convention. It is trivial at T and unramified outside P, cP.

Finally change the Frobenius lift at P by a suitable power of τ, so that $h(f) = 0$. The finite extension used in defining the Frobenius neighborhood is unramified at P, because its ramified places can have been excluded. This change therefore preserves membership of f in W. The congruence for NP is independent of the Frobenius lift. This obtains all the stated properties. □

The precise theorem for the given data is `exists_isAnticyclotomicAt` in `New/Selmer/Corepresentable.lean`. In Lean the extension from local characters is expressed by `IsFromU1`, whose hypotheses include the restriction to ideles at the set T; the data of the tame generator and Frobenius are stored in `IsAnticyclotomicAt`. Our proof above describes the arithmetic construction with the standard Chebotarev and class field inputs, rather than developing these general theorems anew.

Observe why the hypothesis on z is important. Arbitrary prescribed z need not fix the roots of T-units; one cannot in general ensure a Frobenius near that element and also impose the character condition. Normal closure of allowed inertia solves the problem because that entire closure fixes the Kummer extension. The fact that P depends on the neighborhood is also indispensable, as is freedom to replace the Frobenius lift. The construction permits previously obtained restrictions at finite places to remain valid when new characters are added. This is the anticyclotomic operation used in the pencil argument, not the triangular operation that follows. The latter is directed instead at the scalar summands of the trace identity.

41.5 Odd laws and the triangular removal of even classes

We next give a proof of the triangular construction, conditional on two substantial arithmetic inputs. This construction is logically distinct from the ordinary patching problem in which one first bounds the dimension of a residual Selmer group and chooses enough primes to kill it. Here the classes in a nonzero Tate twist may be even. Their required bound is obtained by arranging the order of the primes, not by supplying an independent bound for the size of the even group in every Tate twist. The technique produces the same number of primes at every sufficiently deep finite coefficient quotient. Their locations and their Frobenius lifts may change with the quotient.

Let $A_\nu = \mathbb{Z}/\ell^\nu\mathbb{Z}$, write ε for the cyclotomic character, ε_ν for its reduction to $A_\nu^\times$, and set $K = \ker \varepsilon_\nu$. Let c act on G_E by conjugation. A scalar cocycle d of twist $-e$ in a finite

quotient of a coefficient DVR has the convention

$$d(gh) = d(g) + \varepsilon_\nu(g)^{-e} d(h).$$

It is even if $d(cg) = d(g)$ and is required to satisfy the stated local Selmer conditions. On K it is additive. We need odd functions $Z : G_E \to A_\nu$ which restrict to additive maps on K, satisfy $Z(cu) = -Z(u)$ on K, and transform under conjugation by

$$Z(kuk^{-1}) = \varepsilon_\nu(k)^{e+1} Z(u) \qquad (u \in K). \tag{41.4}$$

They are locally constant. In addition to these elementary requirements the function must satisfy a reciprocity condition, stated in terms of a finite set F of old Frobenius lifts and a finite set T of exceptional primes of the real subfield. If a collection of new split places has distinct underlying real primes off T, Frobenius in K, and Z vanishes at all of these Frobenius lifts except perhaps the one at P_j, the following inference holds. For every even class d of twist $-e$, unramified away from ℓ and the new places and their conjugates, vanishing on F, and every integer b,

$$2Z(f_j) \mid b \text{ in } A_\nu \quad \Longrightarrow \quad bd(\tau_j) = 0. \tag{41.5}$$

This is the structure `IsLaw` in `New/Selmer/ReciprocityLaw.lean`; the local Selmer conditions in the source are carried by `evenClasses`. It is an assertion about all finite quotients of coefficient rings and not just $\mathbb{F}_\ell$.

Proposition 41.10 (Formal-source inputs: existence and detection)**.** *Under the field hypotheses of the proof, in particular the stable unramifiedness and the lower bound of four for the number of infinite places, given k old split places as above and at most k prescribed elements of K, there exists a law Z of type e for the old Frobenius lifts which vanishes at the prescribed elements and takes a unit value at some element of K. Independently there is a finite set D_0 of detecting points, of size at most y^{20} when the field is contained in the cyclotomic field of order $N_E < y$ and $\ell \le y^2$, on which vanishing detects all cocycles of a Tate twist with the specified Selmer conditions and without extra ramification. The detecting set can be used at every finite coefficient level and for all twists in question.*

The inputs are `exists_isLaw` in `New/Selmer/ReciprocityLaw.lean` and `exists_detecting_points_cocycles` in `Lib/Cyclotomic/OddSha/OddShaBound.lean`. The proof of the law uses global duality over the real subfield. To indicate why the quantitative lower bound on the number of choices matters, we record an algebraic step in the proof of primitive relations.

Lemma 41.11 (A primitive finite relation)**.** *Let $A = \mathbb{Z}/\ell^b\mathbb{Z}$ with $b \ge 1$, let Y be a finite abelian group and let $p : Y \to A^I$ be additive. Fix k coordinate positions in I. If $|Y[\ell]|\ell^{k+1} \le \ell^{|I|}$, there is $w \in A^I$ vanishing in those k positions, annihilating $p(Y)$ under the dot product, and primitive (at least one coordinate is a unit).*

Proof. Take the homomorphism

$$A^I \longrightarrow \mathrm{Hom}_{\mathbb{Z}}(Y, A) \oplus A^k, \quad w \longmapsto (y \mapsto w \cdot p(y),\ (w_i)_{i \text{ prescribed}}).$$

Denote its image by B and its kernel by L. If no relation in L were primitive then $L \subset \ell A^I$, so $B/\ell B$ would have dimension $|I|$ over $\mathbb{F}_\ell$. The structure theorem gives $\dim B/\ell B = \dim B[\ell]$. Since $B[\ell]$ embeds in the ℓ-torsion of the displayed target, its dimension is at most $\log_\ell |Y[\ell]| + k$. For the first factor this follows by decomposing Y into cyclic primary groups; homomorphisms to $A[\ell]$ have one parameter for each ℓ-primary cyclic summand. The asserted bound is strictly less than $|I|$, a contradiction. □

This reproduces the counting step in `exists_primitive_relation` of `Lib/ClassField/GaloisCohomology/DiscreteUnits.lean`. Producing Y with the required bound is part of the global argument, not a consequence of the lemma alone. In particular primitive relations should not be assumed merely because the number of places is large.

We now explain how Proposition 41.10 supplies the exact form of the triangle. Fix an element a with separated cyclotomic values, use $\gamma = a^{-1}$, and suppose each $\varepsilon(\gamma)^u - 1$ with $0 < u < t'$ divides ℓ^h in the coefficient ring. The set J consists of pairs (e, i) for $1 \le e < t'$ and $i \in D_0$, ordered lexicographically, so it has types e and each type occurs $|D_0|$ times. Write

$$b_i = (ai)^{-1}(ia) \in K,$$

using the point i itself as a group element. We seek distinct primes with Frobenius $f_j \in K$ of types e, with a law Z_j at each step. The law Z_j must satisfy reciprocity with respect to the earlier Frobenius lifts of the same type; it should vanish at every later chosen Frobenius lift; and $2Z_j(f_j)$ should divide ℓ^H. For control of the cocycles without extra ramification we will also arrange

$$d(f_{(e,i)}) = 2d(b_i) \tag{41.6}$$

for every even unramified class of type e (unramified here and below means outside ℓ).

Consider the step $j = (e, i)$. The evaluations at all possible laws of all types, restricted to K, form an additive subgroup of the product of the groups A_ν. Indeed the laws are additive on K, so sums and inverses of simultaneous evaluation vectors are realized at products and inverses of elements of K. Project this group to the coordinates of the old laws of type e. A subgroup of A_ν^k needs at most k generators: its minimum number of generators equals the dimension of its ℓ-torsion, bounded by k. Choose elements $y_1, \dots, y_v \in K$ whose evaluation vectors generate that projection. By the law existence assertion choose Z_j vanishing on these elements and primitive at an element $w \in K$. Subtract a suitable integral combination of the evaluation vectors of the y_i from that of w. The result is realized at an element $u_0 \in K$ where every earlier law of type e vanishes and Z_j still takes its unit value.

Earlier laws of the other types are removed by a polynomial in a single conjugation operation. On simultaneous evaluation vectors write C for conjugation by γ, which is defined on this subgroup. Its action on a coordinate of type d is multiplication by m^{d+1} with $m = \varepsilon_\nu(\gamma)$. Replace the vector of u_0 by

$$\prod_{\substack{0 \le k < t' \\ k \ne e}} (C - m^{k+1}) \operatorname{ev}(u_0).$$

The resulting vector is still realized at a point $u \in K$ since all its operations are conjugation, sums and integer multiplications (choose integral representatives of the coefficients in A_ν). The earlier laws all vanish there; the new law takes its original unit value multiplied by

$$m^{t'-1} \prod_{k \ne e} (m^e - m^k).$$

The index 0 in this product causes no harm although the set of positive types omits it. It will allow all Tate-character differences to occur in one bound.

The controlling divisibility is calculated before reducing modulo ℓ^ν. For $e > k$ one has

$$\varepsilon(\gamma)^e - \varepsilon(\gamma)^k = \varepsilon(\gamma)^k(\varepsilon(\gamma)^{e-k} - 1),$$

and for $e < k$ one uses the negative of the reverse difference. All prefactors are units. Hence every difference divides ℓ^h, and their product divides $\ell^{(t'-1)h}$. The source reserves three more powers of ℓ^h to handle a factor 2 and a squared character difference. If $\ell \ne 2$ the factor 2 is a

unit; for $\ell = 2$ the field contains the fourth roots of unity and $4 \mid \varepsilon(\gamma) - 1$, so $4 \mid \ell^h$ provided there is at least one positive type. With $t' + N_E < y$ we may use the uniform bound

$$H = yh, \qquad 2 \prod_{k \neq e} (\varepsilon(\gamma)^e - \varepsilon(\gamma)^k) \mid \ell^H.$$

It is legitimate to use this divisibility even when some factors reduce to zero in A_ν: in that case the right-hand side also reduces to zero. The assertion concerns divisibility, not the division of an element of a quotient ring by a nonunit. At a deeper level with $\nu > H$ it implies a genuine nonvanishing statement.

To prescribe both the older laws and the new detector condition, form the element

$$b_i \, cb_i \, u \, (cu)^{-1} \in K.$$

Every law takes there the value $2Z(u)$ since it is odd, and every even class takes the value $2d(b_i)$. Thus the prospective Frobenius can be chosen simultaneously for both tasks. The genericity condition on Frobenii in `exists_isPrimeTriangle` permits the neighborhood to be translated by a factor δ^{ℓ^ν} with $\varepsilon(\delta) = 1$. This factor has zero value under any of the additive functions being considered. It does not alter the prescriptions. The locally constant laws and the finitely many scalar cocycles in A_ν give finitely many open conditions, and one omits the finitely many exceptional real primes for the law and its predecessors. Chebotarev therefore supplies a degree-one prime P_j not over a previously used real prime. This yields norm congruent to 1 modulo ℓ^ν and the required Frobenius in the chosen generic set. To pass from scalar cocycles in A_ν to arbitrary coefficient quotients use their additive characters into A_ν: those characters are jointly faithful by finite abelian duality and commute with the action of $\mathbb{Z}/\ell^\nu\mathbb{Z}$. They preserve the scalar twist and the even condition. This is the content of `eq_zero_of_mem_evenClasses` in the same file as the laws. This descent is essential for applying Chebotarev with finitely many conditions independently of the eventual coefficient ring.

Let us verify the two consequences of this construction.

The triangular ramification step. Suppose d is an even class of type $e = \mathrm{ty}(j)$, unramified outside ℓ and the places in the list, satisfying $d(\tau_i) = 0$ for every $i < j$ and $d(f_i) = 0$ for earlier i of the same type. It is then unramified at the earlier auxiliary places. For the conjugate places this follows from evenness. At a place prime to ℓ the wild inertia is pro-prime-to-ℓ and contributes no homomorphism to the finite ℓ-group; hence vanishing at the specified tame generator suffices. Apply the law Z_j to the remaining places $i \geq j$ (these avoid its exceptional real primes), with the earlier same-type Frobenius lifts in its prescribed set F. It vanishes on all f_i for $i > j$ by construction, while $2Z_j(f_j) \mid \ell^H$. Equation (41.5) gives $\ell^H d(\tau_j) = 0$. In an induction one must include the earlier Frobenius conditions as well as the earlier inertial ones; their separation into types is part of the triangle, not an optional strengthening of a weaker statement.

The remaining unramified class. Suppose finally that a class d of positive type e has no auxiliary ramification, and vanishes on every Frobenius of that type. Put $\xi(g) = \varepsilon_\nu(g)^{-e}$ and $\Delta(g) = \xi(g) - 1$. The function

$$B(g) = \Delta(a)\, 2d(g) - \Delta(g)\, 2d(a)$$

is a cocycle of the same twist, with the same required unramified local conditions. For the first term this is linear stability of the Selmer module, and the second term is a coboundary.

For a detector i the cocycle formula gives

$$d(ai) - d(ia) = \Delta(a)d(i) - \Delta(i)d(a),$$
$$d(b_i) = \xi(ai)^{-1}\big(d(ia) - d(ai)\big).$$

By (41.6), $2d(b_i) = 0$. Thus B vanishes at every $i \in D_0$, so the detector assertion gives $B = 0$. This is precisely the coboundary relation

$$\Delta(a)\, 2d(g) = \Delta(g)\, 2d(a).$$

We also have $4\Delta(a)^2 \mid \ell^H$ by the same factor estimates. The relation controls the remaining class with a scalar denominator; no division modulo a possibly noninvertible character difference has been assumed. The patching argument can now work with the whole even group, whereas a claim that it was uniformly small would require a much stronger arithmetic input.

Combining the two consequences with the detector bound gives

$$(|J| + |D_0|)H = t'|D_0|yh \le y^{22}h.$$

If the separation bound has $h = 0$, the entire cost is zero; no estimate of the product of the number of detectors and 0 needs the cardinal bound for small primes. The precise package is `IsPrimeTriangle` in `New/Selmer/Corepresentable.lean`, and its construction is `exists_isPrimeTriangle` in `New/Selmer/TrianglePrimes.lean`. Its field hypothesis, the local conditions in the law and detector assertions, and the genericity of the Frobenius set are needed for the arithmetic existence of the primes; the calculation of H is elementary given those assertions. This explains both the dimension-independent arrangement of the auxiliary levels and the particular power y^{22} that appears later in the bound.

41.6 The algebra of adjoining relations

We turn to the part of the determinant estimate in which lengths of modules of functions must be converted into products of congruence ideals. It is useful to separate this purely integral calculation from bounds on Galois cohomology. Let R be a commutative coefficient algebra, let $N \subset R^d$ be the submodule of rows expressing a system of relations, and denote the ideal generated by the determinants of d rows of N by $\mathfrak{F}(N)$. This is the zeroth Fitting ideal of R^d/N (with the evident determinant convention when $d = 0$). The inclusion of a larger relation module gives a larger Fitting ideal. Thus one cannot enlarge the relation system freely: the determinant gained at the end of an argument has to be brought back into the original ideal with the correct cost.

Lemma 41.12 (Cost of one relation)**.** *If $v \in R^d$ and $qv \in N$, then $q\mathfrak{F}(N + Rv) \subset \mathfrak{F}(N)$. More generally, suppose that $q_j v_j \in N + Rv_1 + \cdots + Rv_{j-1}$ for $1 \le j \le L$. Then*

$$\left(\prod_{j=1}^{L} q_j\right) \mathfrak{F}(N + Rv_1 + \cdots + Rv_L) \subset \mathfrak{F}(N).$$

Proof. Expand a determinant of rows $n_i + a_i v$. Every term using v in two or more rows vanishes by alternation. The term containing no occurrence of v belongs to $\mathfrak{F}(N)$, and after multiplication by q each remaining term is a multiple of the determinant with row $qv \in N$ in the appropriate position. This proves the first assertion. The second is obtained by iteration in descending order of j. □

In the formal source these are `fittingIdeal.mul_mem_of_smul_mem` and `fittingIdeal.triangle` in `Lib/General/Algebra/FittingDuality.lean`. A point of the triangle argument is that a relation can already hold after the previous relations have been added; Lemma 41.12 does not require them all to hold in the first quotient. This avoids paying separately for the unknown number of classes in each twist. If $q_j = \ell^H$, the explicit cost is $|J|H$ in normalized valuation.

There is a related use of lengths which applies when relation forms cannot be specified as a finite ordered list in advance. We give an algebraic model reproducing the induction in `fittingIdeal.exists_cost_le_relLength`. Let $\mathcal{O}$ be a DVR, $p \neq 0$, $A = \mathcal{O}/p$, R an $\mathcal{O}$-algebra. Let $\phi : \mathcal{O}^u \to R^d$ be linear over $\mathcal{O}$, and $z(g) \in \mathcal{O}^u$ be prescribed vectors indexed by a set G. Assume $p\phi(\mathcal{O}^u) \subset N$. For any relation module $L \supset N$, set

$$D_L = \{\lambda \in \mathrm{Hom}_{\mathcal{O}}(\mathcal{O}^u, A) : \lambda(\phi^{-1}(L)) = 0\}, \qquad F_L = \{g \mapsto \lambda(z(g)) : \lambda \in D_L\}.$$

Fix a submodule V of A-valued functions on G. The relative length $\mathrm{length}(F_L/(F_L \cap V))$ is finite; denote it by b_L. No finiteness assumption on the set G is needed because F_L is an image of a finite module.

Proposition 41.13 (Relative-length cost)**.** *There exists $L \supset N$ with $\pi^m \mathfrak{F}(L) \subset \mathfrak{F}(N)$ for some integer $m \leq b_N$, such that every $v \in \mathcal{O}^u$ annihilated by all $\lambda \in D_L$ with $\lambda(z(\cdot)) \in V$ satisfies $\phi(v) \in L$.*

Proof. If the conclusion already holds at N, take $m = 0$. Otherwise choose a bad v: the selected functionals annihilate v but $\phi(v) \notin N$. Let $j \geq 1$ be minimal such that $\pi^j \phi(v) \in N$; the existence of such j follows from the hypothesis on p. Adjoin the row $\pi^{j-1}\phi(v)$ to the relation module. The new module N' costs one power of π, by Lemma 41.12. The annihilator $D_{N'}$ is the additional restriction $\lambda(\pi^{j-1}v) = 0$. It is a proper submodule of D_N: over $\mathcal{O}/p$, the dual maps into A separate points of a finite quotient annihilated by p. This follows at once by writing that quotient as a sum of modules $\mathcal{O}/\pi^j$ with $j \leq v_\pi(p)$ and embedding each cyclic summand into A.

Now $F_{N'} \subset F_N$. If their relative lengths to V were equal, their images in the quotient by V would be equal. For every $\lambda \in D_N$ choose $\lambda' \in D_{N'}$ with $\lambda(z(\cdot)) - \lambda'(z(\cdot)) \in V$. By the defining property of v, $(\lambda - \lambda')(v) = 0$. It would follow that $\lambda(\pi^{j-1}v) = 0$ for every $\lambda \in D_N$, contrary to the properness just proved. Thus $b_{N'} < b_N$. Repeat the construction at N'; there can be at most b_N such steps, since the relative length is a nonnegative integer. Multiplying the one-step ideal inclusions proves the claimed total cost. □

Two caveats about this proposition are relevant to its use with cocycles. The module V must be fixed during the induction (in the application it expresses the permitted surviving cocycles), and the condition on v concerns the annihilator after the enlargement, not only the annihilator at the starting module. The proof works because any bad vector decreases the remaining relative length. It would be incorrect simply to identify the normalized length of the space of cocycles with the valuation of a single congruence element without proving such an inclusion of Fitting ideals.

41.7 Passing from scalar rows to matrix blocks

For the distinguished block one has an additional advantage: the integral span of the representation is the full matrix algebra. This makes the ordinary determinant bound for scalar lattices apply to a matrix of group-algebra elements. The matrix units from Lemma 41.5 explain the mechanism behind the following statement.

Lemma 41.14 (Block determinant from a dual bound)**.** *Let $\Psi : B \to M_n(\mathcal{O})$ be a surjective homomorphism of $\mathcal{O}$-algebras, $L \subset B^u$ a left B-submodule, and $p \neq 0$. Suppose that the space of $w \in ((\mathcal{O}/p)^n)^u$ annihilated by every $t \in L$ under the pairing*

$$(t, w) \longmapsto \sum_{i=1}^{u} \Psi(t_i) w_i$$

has ordinary length at most $b < v_\pi(p)$. There is a square matrix $D \in M_u(B)$ with all rows in L such that its flattened block image has $v_\pi(\det \Psi(D)) \leq b$.

Proof. Let $K \subset \mathcal{O}^{nu}$ be generated by the scalar rows of the horizontal block matrices $(\Psi(t_1), \ldots, \Psi(t_u))$ for $t \in L$. By surjectivity, left multiplication by any matrix unit is induced by an element of B. Consequently an arbitrary scalar row of K can be placed in any desired position of a block row with all other scalar rows zero; addition lets us choose the n scalar rows independently. The annihilator in $(\mathcal{O}/p)^{nu}$ is the ordinary orthogonal complement of K there. Apply Smith normal form to $K' = K + p\mathcal{O}^{nu}$. The length of its annihilator equals the length of $\mathcal{O}^{nu}/K'$, by duality over $\mathcal{O}/p$. Indeed its elementary divisors are powers π^a with $a \leq v_\pi(p)$, and the corresponding kernel of multiplication has length a. Take a basis matrix D' for K' with determinant valuation at most b. Each scalar row of this matrix is congruent modulo p to a row in K. Replace each by that row; the determinant is unchanged modulo p, and because $b < v_\pi(p)$ its valuation is unchanged. Divide the chosen list of nu rows into successive groups of n. The independent-row observation produces a block row of L for each group. Taken together these give D with the required flattened determinant. □

This is `exists_rows_addVal_det_le` in `Lib/General/Algebra/BlockFittingBound.lean`. Note the strict inequality against the modulus. If the determinant's valuation were at least $v_\pi(p)$, a lift of its class modulo p might have larger valuation, or even be zero. The source can choose very deep Taylor–Wiles moduli, and this strict inequality is therefore natural. The lemma requires integral, not merely rational, surjectivity; it is here that the primitive residue-field labels pay for themselves a second time.

41.8 Two applications of Cramer's rule over a family

The determinant of a block relation matrix at a distinguished Galois representation is not yet a Hecke operator acting on every eigensystem. The construction must take place in a common algebra of functions on the eigensystems. Moreover the Hecke algebra need not contain the individual matrix entries of the representations. It may contain their traces and characteristic-polynomial coefficients only. We give the common-algebra argument, which is a useful safeguard against incorrectly treating the representations as a simultaneously diagonalizable family of matrices.

Let $S \subset \prod_{x \in Y} L$ be the commutative algebra of coefficients on a finite family Y, and let $u_x : B \to M_n(L)$ be representations of an algebra B containing a group algebra over $\mathcal{O}$. Suppose the coefficient functions of the characteristic polynomials of the group elements belong to S. We need the following integrality input, which the formal source proves from Amitsur's formula: the coefficients of the characteristic polynomial of any element of the group algebra with coefficients extended to S belong to S, and the assertion remains true for matrices of such group-algebra elements after flattening the blocks. This is `amitsur_charpoly_mem_matrix` in `New/Selmer/RibetCramer.lean`, together with `amitsur_charpoly_mem_of_span` used there. The conclusion is genuinely integral. Newton identities over a field of characteristic zero alone do not give it over an integral ring because of division by integers. In the following argument we use the Amitsur integrality statement as formal-source input.

It follows by Cayley–Hamilton that each square group-algebra matrix D has a polynomial "adjugate" W with coefficients in S, satisfying after application of every u_x

$$u_x(W)u_x(D) = \det(u_x(D))1.$$

Here a group-algebra determinant symbol on the right is not defined; it is the determinant of the flattened image, and the family of its values belongs to S. The polynomial is the signed characteristic-polynomial expression for the adjugate, so the displayed identity also holds at singular matrices. The same reasoning supplies the adjugate of any element of the group algebra separately. Both lift to the algebra extended by S.

Let $\mathfrak{J}$ be an S-submodule of the ambient product of coefficient fields, and form the left ideal spanned by elements r whose trace against every group element belongs to $\mathfrak{J}$. Then the trace of any such left-ideal element against any group-algebra element with coefficients in S belongs to $\mathfrak{J}$. This follows from the cyclic invariance of trace and linearity: for br paired with h, use $\operatorname{tr}(brh) = \operatorname{tr}(rhb)$, and expand hb as a linear combination of group elements. Let V be a left group-algebra-linear map on rows, let w be a row, and let D be square with

$$V(D_i) = r_i + m_i V(w)$$

for elements r_i of this trace ideal and m_i of a designated submodule $\mathfrak{a}$. We write r and m as columns. Define the function β by $\beta_x = \det u_x(V(w))$ and write $F_x = \det u_x(D)$. Both $\beta, F \in S$. The group-algebra expression over S

$$b = F\,1 - wWm$$

satisfies $bV(w) = wWr$ after application of each u_x: indeed the identity for the adjugate gives $wWV(D) = FV(w)$. For a group-algebra element h set $f_x = \operatorname{tr} u_x(bh)$. The coefficient integrality implies $f \in S$. Multiply the trace by β, using the adjugate of $V(w)$ and trace cyclicity. We obtain

$$\beta f = \operatorname{tr}\big(u(bV(w))\operatorname{adj}(u(V(w)))u(h)\big) \in \mathfrak{J}.$$

For clarity, this equality of functions uses the adjugate identity before substituting $bV(w) = wWr$: in each component the product $u(V(w))\operatorname{adj}(u(V(w)))$ is the scalar β_x. Finally, at any component x where $u_x(mh) = 0$ for all $m \in \mathfrak{a}$, the second term of b contributes no trace and

$$f_x = F_x \operatorname{tr} u_x(h).$$

This proves the assertion of `exists_ribet_element_cramer` in `New/Selmer/RibetCramer.lean` given the Amitsur input. The integral cost of the extra normalizing determinant β is visible here: a small valuation for F_x alone does not give a desired integral trace relation. The arithmetic construction spends part of its budget ensuring a bound for the determinant of $V(w)$ and a separate bound for F_x. The distinguished component permits the annihilation by $\mathfrak{a}$, not necessarily the other components; the relation $\beta f \in \mathfrak{J}$, however, is valid simultaneously at all components. This is why the construction can supply an operator that isolates a row in the Hecke algebra.

41.9 A transfer calculation at one auxiliary level

We finish with the first-order operation underlying the local patching statement. It concerns a line stabilizer in a finite general linear group. Although the global statement also requires a comparison of automorphic forms at different levels, the algebra of this local operation can be checked directly.

Lemma 41.15 (Logarithmic transfer)**.** *Let k be a finite field of size q, let $G = \mathrm{GL}_N(k)$, and let H be the stabilizer of a line. Suppose $m \mid q-1$ and fix a generator u of $k^\times$. Define $a : H \to \mathbb{Z}/m$ by the logarithm of the scalar action on the line, taking u to 1. The transfer of a is of the form $d \circ \det$ for a character $d : k^\times \to \mathbb{Z}/m$. It satisfies*

$$N!\, d(u) = N! \quad \text{in } \mathbb{Z}/m.$$

Proof. Choose one representative x for every left coset of H. For $g \in G$ let $\overline{gx}$ denote the representative of gxH, and define $h(g,x) = \overline{gx}^{-1} gx \in H$. Transfer is the homomorphism $g \mapsto \sum_x a(h(g,x))$; multiplicativity follows by composing the permutations of the cosets and adding the corresponding logarithms. On any elementary transvection its value is zero because the transvection has order equal to the characteristic of k, prime to m. Since G is generated by elementary matrices and invertible diagonal matrices, and diagonal entries in different positions are conjugate, the transferred character factors through determinant. At the scalar matrix $u1$, every coset is fixed and its term in the sum is $a(u1) = 1$. There are $(q^N-1)/(q-1) = 1 + q + \cdots + q^{N-1}$ lines, which is congruent to N modulo m. Since the determinant of $u1$ is u^N, we get $Nd(u) = N$. Multiplying by $(N-1)!$ proves the formula (if $N = 1$ it is immediate). □

The source uses precisely this calculation in `exists_line_stabilizer_log` in `New/Patching/OnePrime.lean`. In its local-field formulation the line stabilizer is the reduction modulo the prime of the corresponding block parahoric. Both characters are trivial on the indicated pro-Iwahori group, because there the determinant and the scalar action on the line reduce to 1. The factor $N!$ is deliberately not inverted; in particular the proof works when $\ell \mid N!$. In characteristic zero it would be possible to divide the scalar identity, but doing so would change the integral denominators.

Here is the ring calculation with logarithmic weights. We give it in a form that also applies to noncommutative operator rings; the distinction between operator multiplication and scalar multiplication is retained throughout. Let R be such a ring with identity, $s : R \to R_0$ a ring map killing m, and $\rho : G \to R^\times$ the group action. Let B be an additive map from integer-valued functions on $\mathbb{Z}/m$ to R, with

$$B(f)\rho(p) = B(f(\cdot - a(p))) \qquad (p \in H).$$

The source takes B to be obtained from level-changing operators; for the moment it is only an additive map obeying this translation law. Choose the function $v : \mathbb{Z}/m \to \mathbb{Z}$ of least absolute representatives. Any lift of the residue class would give the same image modulo m. For fixed left coset representatives x as above define

$$T = \sum_x \rho(x), \quad U_1 = \sum_x \rho(x^{-1}), \quad \mathcal{N} = \sum_{g \in G} \rho(g), \quad F = TB(-v)U_1.$$

Assume $\rho(H)$ fixes $B(1)$ on the left, $LB(1) = 0$, $L\rho(g) = L$ for every $g \in G$, and an element Θ satisfies

$$\sum_{p \in H} \rho(p^{-1})\Theta\rho(p) = n\,1.$$

Suppose multiplication by $|G|$ is injective in R. These hypotheses imply

$$\mathcal{N}\Theta B(1) = nTB(1), \qquad L\Theta B(1) = 0. \tag{41.7}$$

For the first identity decompose $\mathcal{N} = T\sum_{p \in H} \rho(p)$ and use $\rho(p^{-1})B(1) = B(1)$ to insert the inverse factor on the right of Θ. The second follows on multiplying the first by L on the left, because the left-hand side then equals $|G|L\Theta B(1)$ and the right-hand side vanishes. Injectivity of multiplication by $|G|$ is required before reduction modulo m, not in R_0.

Proposition 41.16 (First-order weighted operator)**.** *Under these hypotheses, for each $g \in G$ there exists $Z_g \in R$ such that*

$$s(nF(\rho(g)-1)) = s(\mathcal{N}Z_g), \qquad LZ_g = 0.$$

In addition, write $S_H(f) = \sum_{p\in H} f(a(p))\rho(p)$ and $S_G(f) = \sum_{g\in G} f(\mathrm{trf}(a)(g))\rho(g)$, where $\mathrm{trf}(a)$ is transfer. Suppose for some fixed $\eta \in G$ and integer k, $k\,\mathrm{trf}(a)(\eta) = k$ in $\mathbb{Z}/m$. Suppose operators $A_t, A_{t'}, r$ satisfy $A_t, A_{t'}$ commuting with the operators T, U_1 and, for every integer-valued f,

$$B(f(\cdot - k) - f)A_t = S_H(f(\cdot - k) - f)A_{t'},$$
$$S_G(f)A_{t'} = (\rho(\eta) - 1)S_G(f)r.$$

Then

$$2k\, s(FA_t) = 2k\, s(\mathcal{N}r).$$

Proof. Congruent integer weights modulo m give the same result after s, by additivity. Reindex the sum defining $U_1\rho(g)$ using the coset representative of gxH. The translation law for B and $-v(b-a) \equiv -v(b) + v(a) \pmod m$ show that

$$s(F(\rho(g)-1)) = s\left(TB(1)\sum_x v(a(h(g,x)))\rho(x^{-1})\right).$$

Multiplying by n and using (41.7), take

$$Z_g = \Theta B(1)\sum_x v(a(h(g,x)))\rho(x^{-1}).$$

The second part of (41.7) gives $LZ_g = 0$.

For the additional assertion, expand $TS_H(v)U_1$ by its three indices. For each fixed coset representative x, each $g \in G$ occurs exactly once in this expansion with final factor $\rho(x^{-1})$ and middle weight $v(a(h(g,x)))$. Summing the weights over x gives a representative of the transferred logarithm, hence

$$s(TS_H(v)U_1) = s(S_G(v)).$$

Now take $f(b) = -v(b)(v(b)+k)$. Modulo m its difference $f(b-k) - f(b)$ is $2kv(b)$. Apply the first additional operator identity to this f, multiply by T, U_1, and reduce modulo m. Since F uses the weight $-v$, this yields

$$2k\, s(FA_t) = -2k\, s(S_G(v)A_{t'}),$$

where the commutation of A_t and $A_{t'}$ with U_1 is used, as in the source's commuting family. Inserting the second operator identity gives $-2k\, s((\rho(\eta)-1)S_G(v)r)$. But $\rho(\eta)$ multiplies the weighted sum by a translation of its weight through $\mathrm{trf}(a)(\eta)$, by reindexing over G. The difference of these two weights, after multiplication by $-2k$, is congruent to the constant $2k$; this uses $k\,\mathrm{trf}(a)(\eta) = k$. The result is $2k\, s(\mathcal{N}r)$ as claimed. □

This is the algebraic content of `one_prime_first_order` in `New/Patching/OnePrime.lean`. The local logarithm gives one factor $k = N!$, while the factor 2 appears because the polynomial used to extract the first-order term is quadratic. Notice that the argument did not divide by k, by 2, or by the finite index of the line stabilizer. The operator F is built by sums and products in the ring of integral level operators before any reduction modulo m. This integrality is the reason for introducing least absolute representatives rather than an abstract logarithm of an operator over a fraction field.

The passage from this ring identity to the automorphic comparison is not merely an application of group theory. In the source, the operators B are defined from level changes, the two weighted-sum identities arise from Hecke operators acting on a line stabilizer, and the normalizing element satisfies a finite averaging identity. The identifications of the spaces, integrality of the operator on the prescribed lattice of forms, and the multiple-prime construction of compatible diagonal operators are the formal-source inputs `BaseLevel.exists_tie` and `BaseLevel.exists_diagOp_mem_ellIntegralForms` in `New/Patching/OnePrimeTie.lean`, and the Hecke-string assertions cited in Section 40.11. Proposition 41.16 reconstructs the transfer calculation used in these inputs; it does not establish their global automorphic hypotheses.

41.10 Separation of the arithmetic and algebraic budgets

The preceding arguments explain why four superficially different finiteness problems lead to four different costs in the final exponent of the integral projection.

1. Residual irreducibility is formulated as full integral span, proved with a tame pair. It enables a bound on a single row test space to be turned into a bound for a flattened determinant (Lemma 41.14). It incurs no extra linear algebra denominator.

2. The balanced row condition concerns the global dimension of local restrictions. The mirror estimate turns an apparently many-variable isotropy constraint into a scalar defect. This is why the estimate measures the length of a kernel of a scalar, not the rank of the entire dual Selmer module.

3. In nonzero Tate twists the odd class bounds and the reciprocity laws serve different purposes. The law makes the cost of adjoining a relation in triangular order uniform, even without bounding all even classes. Lemma 41.12 turns each such addition into a multiplicative loss of at most ℓ^H. The scalar character differences and the detector cost have already been included in $y^{22}h$; one should not multiply this cost by the number of all possible residual characters.

4. A flattened determinant is converted into a trace function in the common eigenvalue algebra by two instances of Cramer's rule. The determinant of the normalizer is a further explicit cost. The local first-order group-ring calculation, in turn, can be executed without inverting the finite group orders; where an integer factor is lost it is recorded in the denominator budget.

These calculations do not make the arithmetic hypotheses in the previous chapter dispensable. In particular the existence and local comparison of attached Galois representations, the calculation of crystalline local lengths in the balance proposition, the global duality and odd-class bounds, the simultaneous twist selection for all rows, and the level-comparison assertions are specialized inputs whose exact boundaries were stated there. The estimates assembled from them control the valuations uniformly in the coefficient field, the individual twist and the deep finite Taylor–Wiles modulus. This uniformity, rather than an equality for one specially chosen automorphic eigensystem, is what permits the arguments of Sections 40.12 and 40.13 to yield the fit family.

Chapter 42

Arithmetic of the congruence argument: cyclotomic classes, descent and simultaneous twists

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 44, 45, 46 and the final audit in Chapter 57.

We reconstruct a further portion of the arithmetic underlying the congruence bound. In particular, we prove the cyclotomic estimates that permit the determinant argument to use an explicit integer controlling only odd classes. A comparable bound on the whole class group of a cyclotomic tower would be an unjustified strengthening. We also develop finite coordinates for cocycles, the construction of pencils of bounded cardinality, and selected steps in the transport from Hecke algebras to Galois representations and back. The deepest geometric compatibility and integral comparison assertions are separated from the finite arguments they imply.

In the first half of this chapter let N be the conductor of a cyclotomic field containing the CM field E, and let e be a positive integer. This N is the conductor, not the varying split prime of the unitary chain. Write ℓ for the coefficient prime, $m_0 = \ell^\nu$ for a torsion modulus, and O for the integers in a finite extension of $\mathbb{Q}_\ell$. The ordinary O-module length is denoted by λ_O and the normalized length by $\lambda = \lambda_O/\lambda_O(O/\ell)$. We use arithmetic Frobenius for reciprocity, so the cyclotomic character ϵ at an unramified Frobenius is the residue cardinality. Let ψ be a Dirichlet character modulo N and write

$$\chi(g) = \epsilon(g)^e \psi(g \bmod N),$$

where the second factor means evaluation via the action on the N-th roots of unity. At the odd components we impose $\chi(c) = -1$, or equivalently $\psi(-1) = -(-1)^e$.

42.1 The integer in the odd-class budget

The first task is to find an integer whose divisors control the relevant Stickelberger expressions at all coefficient primes. The coefficient ring is allowed to vary, and the summation of the local losses uses the factorization of an integer. The source implementation is `IsStickelbergerBound` and `exists_isStickelbergerBound` in `Lib/Cyclotomic/Voronoi/StickelbergerInteger.lean`. Here is a construction with the same bound, using a determinant of an integral convolution matrix.

Let $B_j(X)$ be the Bernoulli polynomials, given by

$$\frac{te^{Xt}}{e^t-1}=\sum_{j\ge 0}B_j(X)\frac{t^j}{j!}.$$

For every $j\ge 0$, $(j+1)!B_j(X)$ has integral coefficients. Indeed the recurrence $B_0=1$, $B_j=-\frac{1}{j+1}\sum_{i<j}\binom{j+1}{i}B_i$ proves integrality of $(j+1)!B_j$ by induction, and the formula $B_j(X)=\sum_{i=0}^{j}\binom{j}{i}B_iX^{j-i}$ proves the polynomial assertion. We use

$$P(X)=(e+3)!N^{e+2}B_{e+2}(X/N),\quad D=(e+3)!(e+2)N,\qquad b(X)=P'(X).$$

The generating function gives the identities

$$P(X+N)-P(X)=DX^{e+1},\qquad b(X)=DN^eB_{e+1}(X/N). \tag{42.1}$$

Lemma 42.1 (A twisted Voronoi congruence)**.** *Let $f=Ns$ with $s>0$ and let $d>0$ satisfy $d\equiv 1 \pmod N$ and $(d,s)=1$. For a Dirichlet character ψ modulo N in a characteristic-zero domain O, extended by zero on nonunits, set*

$$S_f=\sum_{0\le u<f}\psi(u)u^{e+1},\qquad T_f(d)=\sum_{0\le u<f}\psi(u)u^e\left\lfloor\frac{du}{f}\right\rfloor.$$

Then

$$(d^{e+1}-1)S_f\equiv(e+1)d^efT_f(d)\pmod{f^2O}. \tag{42.2}$$

Moreover, if $B(\psi)=\sum_{0\le u<N}\psi(u)b(u)$, then

$$DS_f\equiv fB(\psi)\pmod{f^2O}. \tag{42.3}$$

Proof. Multiplication by d permutes residues modulo f and preserves ψ. Writing $v=du-f\lfloor du/f\rfloor$ and expanding gives $v^{e+1}\equiv d^{e+1}u^{e+1}-(e+1)d^eu^ef\lfloor du/f\rfloor \bmod f^2$. The weighted sum of the left side is S_f. For the second assertion, group the terms by their residues modulo N and apply (42.1). For a residue u, the s successive differences of P telescope to $P(u+f)-P(u)$, congruent to $fb(u)$ modulo f^2 because P is integral. □

Put $A=(\mathbb{Z}/N)^\times$ and write $[v]_N\in[0,N)$ for the representative of v. Define

$$h(v)=2b([v]_N)+(-1)^e\mathbf{1}_{v=1}+\mathbf{1}_{v=-1},\qquad C_{u,v}=h(v/u).$$

For every character ψ of A, the vector $(\psi(v))_{v\in A}$ is an eigenvector of C with eigenvalue $c(\psi)=2B(\psi)+(-1)^e+\psi(-1)$. Over $\mathbb{C}$ the character vectors form a basis. The formula for the Dirichlet value at a negative integer is

$$L(-e,\psi)=-\frac{N^e}{e+1}\sum_{u=0}^{N-1}\psi(u)B_{e+1}(u/N).$$

This follows by expressing L as a sum of Hurwitz zeta functions, each of which has value $-B_{e+1}(u/N)/(e+1)$ at $s=-e$. By the functional equation of a primitive Dirichlet L-function and its nonvanishing Euler product at $1+e$, $L(-e,\psi)=0$ if and only if $\psi(-1)=(-1)^e$; for an imprimitive character the additional Euler factors at $-e$ do not vanish because $p^e>1$. In particular, $B(\psi)$ vanishes in this parity and is nonzero in the other. Thus $c(\psi)$ equals $2(-1)^e$ or $2B(\psi)$ respectively. It follows that $D_0=|\det C|>0$.

Proposition 42.2 (One bounded integer for the Stickelberger elements)**.** *There is an integer X_e with $1\le X_e\le 2^{(N+e+3)^6}$ such that:*

1. *For each prime ℓ some $d > 0$, $d \equiv 1 \bmod N$, prime to ℓ, has $(d^e - 1)(d^{e+1} - 1) \mid X_e$.*
2. *If $d \equiv 1 \bmod N$ and $\psi(-1) = -(-1)^e$, then in any characteristic-zero domain containing the values of ψ there exists an element w independent of s such that for $(s, d) = 1$*

$$(d^{e+1} - 1)X_e \equiv 2wT_{Ns}(d) \pmod{sO}. \tag{42.4}$$

Proof. Take $X_e = D_0 \prod_{d \in \{1+N, 1+2N\}} (d^e - 1)(d^{e+1} - 1)$. The two choices of d are coprime. The adjugate identity for C applied to the character vector over O shows that $c(\psi)$ divides $\det C$: use the coordinate indexed by 1, which equals 1. In the desired parity $c(\psi) = 2B(\psi)$, so $X_e = 2B(\psi)u_\psi$ for some $u_\psi \in O$. Combining the two congruences of the lemma gives

$$f(d^{e+1} - 1)B(\psi) \equiv f(e + 1)d^e DT_f(d) \pmod{f^2 O}.$$

Cancel f in the domain, multiply by $2u_\psi$, and take $w = u_\psi(e + 1)d^e D$.

For the size, let $s_0 = N + e + 3 \geq 5$. The Fourier series for B_{e+1} on $[0, 1]$ gives $|B_{e+1}(x)| \leq 2(e + 1)!\zeta(e + 1)/(2\pi)^{e+1} \leq (e + 1)!$. Thus with $M_0 = DN^e(e + 1)!$ one has $|h(v)| \leq 4M_0$. Hadamard's inequality yields $D_0 \leq (4M_0\sqrt{|A|})^{|A|}$. By $n! \leq n^n$,

$$\log_2 M_0 \leq 4s_0 \log_2 s_0, \quad \log_2 D_0 \leq |A|(2 + (4s_0 + \tfrac{1}{2}) \log_2 s_0).$$

The four other factors contribute at most $2(2e + 1) \log_2(1 + 2N) \leq 8s_0^2$. Since $\log_2 s_0 \leq s_0$, the total is bounded by $5s_0^3 + 10s_0^2 \leq s_0^6$. □

For $x \in \mathbb{Q}/\mathbb{Z}$ write $\{x\} \in [0, 1)$ for its fractional representative. The function in the Gauss-sum source is $f_d(x) = \lfloor d\{x\} \rfloor - \lfloor d\{-x\} \rfloor$.

Lemma 42.3 (Removing multiples of the coefficient prime)**.** *Let $f = N\ell^\nu$, $d \equiv 1 \bmod N$, $(d, \ell) = 1$, and $\psi(-1) = -(-1)^e$. In $O/\ell^\nu O$ the element*

$$A_\nu(d) = \sum_{a \in (\mathbb{Z}/f)^\times} f_d(a/f)[a]_f^e \psi(a)$$

divides $(d^{e+1} - 1)X_e$.

Proof. For $\nu = 0$ this is automatic. Put $f_j = N\ell^j$ and $K = (d^{e+1} - 1)X_e$. The sum of the unit terms of T_{f_ν} equals

$$U_\nu = T_{f_\nu}(d) - \ell^e \psi(\ell) T_{f_{\nu-1}}(d).$$

This follows by the substitution $a = \ell b$, using the zero-on-nonunits convention even when $\ell \mid N$. As $e \geq 1$, (42.4) at the two moduli gives $2wU_\nu \equiv (1 - \ell^e \psi(\ell))K \bmod \ell^\nu O$. Substituting the negative of a unit changes its weight by a minus sign modulo ℓ^ν, by parity and binomial expansion. Thus $A_\nu(d) \equiv 2U_\nu \bmod \ell^\nu$. The factor $1 - \ell^e \psi(\ell)$ is a unit, proving the assertion (`sum_floorDiff_dvd` in `Lib/Cyclotomic/Kolyvagin/ComponentBound.lean`). □

42.2 The Jacobi family and its distribution relation

Let p be an auxiliary rational prime and let B be a place of $\overline{\mathbb{Q}}$ above p. Put $n = p - 1$, choose the character of $\mathbb{F}_p^\times$ with values in μ_n whose reduction at B is the identity, and choose a primitive p-th root ζ_p. For $x \in \mathbb{Q}/\mathbb{Z}$ annihilated by n, set

$$G_x = \sum_{a \in \mathbb{F}_p^\times} \chi_x(a) \zeta_p^{[a]_p}, \qquad \overline{\chi_x(a)} = a^{-n\{x\}} \text{ in } \mathbb{F}_p^\times.$$

Here the bar denotes reduction, not complex conjugation. The characters are unique by lifting prime-to-p roots of unity. In particular $G_0 = -1$. The finite geometric-series identity gives, for $x \neq 0$,

$$G_x G_{-x} = \chi_x(-1)p, \qquad |G_x|_{\mathbb{C}} = \sqrt{p}. \tag{42.5}$$

For the first equality, write the double sum in terms of the ratio a/b: the sum over b is $p-1$ if the ratio is -1 and -1 otherwise. The second follows by complex conjugation.

Lemma 42.4 (Tame Gauss valuation)**.** *Normalize $v_B(p) = 1$. Then $v_B(G_x) = \{x\}$, and G_x is a unit at every finite place off p.*

Proof. In $\mathbb{Q}_p(\zeta_p, \mu_n)$ the extension by μ_n is unramified and $\pi = \zeta_p - 1$ has valuation $1/n$. For $x \neq 0$, the reduction of G_x modulo π is zero, as it is the sum of a nontrivial character of $\mathbb{F}_p^\times$. Similarly G_{-x} is a nonunit. If $G_x = \pi^k u$ with u a unit, (42.5) gives $0 < k < n$. For $v \in \mathbb{F}_p^\times$ let σ_v fix the n-th roots and send ζ_p to $\zeta_p^{[v]_p}$. It acts trivially on residue fields and $\sigma_v(\pi)/\pi$ reduces to v. By substituting $a \mapsto va$, $\sigma_v(G_x) = \chi_x(v)^{-1} G_x$. Reducing the quotient gives $v^k = v^{n\{x\}}$, so $k = n\{x\}$ using its range. The case $x = 0$ is immediate. Integrality of the Gauss sums and (42.5) show that they are units off p. □

For $d > 0$ put $j_d(x) = G_x^d / G_{dx}$ and $J_d(x) = j_d(x)/j_d(-x)$. If σ acts on the n-th roots by multiplication by a and on the exponents of ζ_p by multiplication by v, then $\sigma G_x = \chi_{ax}(v)^{-1} G_{ax}$. The scalar cancels in j_d, hence $\sigma J_d(x) = J_d(ax)$.

Lemma 42.5 (Jacobi-family identities)**.** *Suppose $M \mid p-1$ and $Mx = 0$. Then $J_d(x)$ belongs to $\mathbb{Q}(\mu_M)^\times$, is a unit off p, has valuation $f_d(x)$ at B, and has absolute value 1 at every complex embedding. If $q \mid p-1$ is a prime and $qb = 0$, then at every place above q,*

$$J_d(x + b) \equiv J_d(x) \pmod{\mathfrak{m}_q}.$$

Finally, if $r \mid p-1$ and $(r, d) = 1$, then

$$\frac{\prod_{b=0}^{r-1} J_d(x + b/r)}{J_d(rx)} \quad \text{is a root of unity.} \tag{42.6}$$

Proof. Galois covariance shows the field assertion. The valuation follows from Lemma 42.4 and $d\{x\} - \{dx\} = \lfloor d\{x\} \rfloor$. The unit statement follows from the same lemma, and $|j_d(x)| = |j_d(-x)|$ follows from (42.5), also when $dx = 0$.

At $q \neq p$ all roots of unity of q-power order reduce to 1. Thus the terms of the sums defining G_{x+b} and G_x have the same reductions; the denominators of J_d are units. For the distribution relation note that $\sum_{b=0}^{r-1} \{x + b/r\} = (r-1)/2 + \{rx\}$. Since $(r, d) = 1$, the same formula applied to $d(x + b/r)$ gives

$$\sum_{b=0}^{r-1} \lfloor d\{x + b/r\} \rfloor = (d-1)(r-1)/2 + \lfloor d\{rx\} \rfloor.$$

Replace x by $-x$ and subtract to cancel the constant, giving $\sum_b f_d(x + b/r) = f_d(rx)$. This holds after every automorphism of the cyclotomic field, so the quotient in (42.6) is an algebraic unit, of absolute value one at every embedding. It is a root of unity by Kronecker's theorem (its powers have minimal polynomials of bounded degree with integral coefficients bounded by the corresponding binomial coefficients, so they range over a finite set). These arguments reconstruct `exists_jacobi_family` in `Lib/Cyclotomic/Stickelberger/GaussSums.lean`. □

42.3 The derivative cocycles from the Jacobi family

It is not legitimate to take arbitrary Kummer classes of (42.6) and ignore the root-of-unity ambiguity. The normalization that eliminates it is the following. Fix $m_0 = \ell^\nu > 1$ and $\gamma \in G_{\mathbb{Q}}$. In this subsection write the multiplicative group of $\overline{\mathbb{Q}}^\times$ additively, so multiplication by m_0 means taking an m_0-th power. Choose y_x satisfying $m_0 y_x = J_d(x)$, and put

$$F(g,x) = \gamma y_x - g\gamma g^{-1} y_x \in \mu_{m_0}.$$

It belongs to μ_{m_0} because the two automorphisms act identically on the abelian field containing $J_d(x)$. It is independent of the chosen root y_x, because γ and $g\gamma g^{-1}$ agree on all roots of unity. Identify μ_{m_0} with $\mathbb{Z}/m_0$ via a fixed primitive root, so its Galois action is multiplication by ϵ.

Write a_g for the exponent of the action of g on the relevant roots of unity. We have

$$F(gh,x) = F(g,x) + \epsilon(g) F(h, a_g^{-1} x), \tag{42.7}$$

$$F(\gamma,x) = 0, \qquad \sum_{b=0}^{r-1} F(g, x + b/r) = F(g, rx) \quad ((r,d)=1). \tag{42.8}$$

In the first identity insert $g\gamma g^{-1} y_x$ into the difference and use Galois covariance of J_d. The element $g^{-1} y_x$ differs from $y_{a_g^{-1}x}$ by an m_0-th root of unity, which cancels. For the last equality, $\sum_b y_{x+b/r} - y_{rx}$ is itself a root of unity by (42.6); applying $\gamma - g\gamma g^{-1}$ annihilates it.

Put $M = N m_0$ and let $\mathcal{P}$ be a finite set of rational primes $q \equiv 1 \bmod dM$. Choose $p \equiv 1 \bmod M \prod_{q\in\mathcal{P}} q$ different from ℓ and the primes of $\mathcal{P}$. For each $q \in \mathcal{P}$ choose a surjective homomorphism $L_q : \mathbb{F}_q^\times \to \mathbb{Z}/m_0$. For $S \subseteq \mathcal{P}$ define

$$\begin{aligned}\kappa_S(g) = \sum_{a \in (\mathbb{Z}/M)^\times} [a]_M^{e} \psi(a) \sum_{(u_q) \in \prod_{q\in S} \mathbb{F}_q^\times} \left(\prod_{q \in S} L_q(u_q) \right) \\ \cdot F\left(g, \frac{a}{M} + \sum_{q \in S} \frac{u_q}{q} \right) \quad \text{in } O/m_0.\end{aligned} \tag{42.9}$$

The finite logarithmic derivative here is a group-algebra operation, not a derivative of an analytic function.

Lemma 42.6 (Independence of logarithm shifts)**.** *Replacing any $L_q(u_q)$ in (42.9) by $L_q(u_q) - d_q$ with constant $d_q \in \mathbb{Z}/m_0$ does not change the sum.*

Proof. Induct on $|S|$. To change the logarithm at q while allowing arbitrary shifts for the others, compute the difference: the constant d_q times the sum with L_q omitted. For $x = a/M + \sum_{r\in S\setminus\{q\}} u_r/r$, (42.8) gives

$$\sum_{u_q \in \mathbb{F}_q^\times} F(g, x + u_q/q) = F(g, qx) - F(g,x).$$

As $q \equiv 1 \bmod M$, multiplication by q preserves a and permutes the u_r. Reindexing changes their logarithms only by constants $L_r(q)$, and the induction hypothesis shows the sums agree. Repeat to handle all shifts. □

Proposition 42.7 (Finite derivative cocycles)**.** *The maps (42.9) are locally constant cocycles with coefficient character $\epsilon^{e+1}\psi$ and vanish at γ. They vanish on inertia over $\mathbb{Q}(\mu_N)$ away*

from ℓ, p and S. For $q \in \mathcal{P} \setminus S$ and an arithmetic Frobenius φ at a prime over q, there is a tame inertia element τ over the same prime with

$$\kappa_{S\cup\{q\}}(\tau) = \kappa_S(\varphi). \tag{42.10}$$

If $\kappa_\varnothing$ vanishes on inertia at the place B above p, then

$$(\epsilon(\gamma)^{e+1}\psi(\gamma) - 1) \sum_{a\in(\mathbb{Z}/M)^\times} f_d(a/M)[a]_M^e \psi(a) = 0 \quad \text{in } O/m_0. \tag{42.11}$$

Proof. The sums are finite and factor through a finite extension. In (42.7), multiplication of the arguments by a_g^{-1} induces permutations of the units modulo the denominators. Reindexing changes the weight $[a]_M^e\psi(a)$ by the factor $\epsilon(g)^e\psi(g)$, and changes the logarithms by constants, which cancel by Lemma 42.6. This proves the cocycle relation. Vanishing at γ follows from the definition. Outside the permitted set the fields defined by the denominators of the x are unramified over $\mathbb{Q}(\mu_N)$, and the Jacobi values are units. Taking roots of order prime to the residue characteristic of such a unit gives unramified local extensions (lift the separable roots of its residue polynomial by Hensel's lemma). This applies also to every conjugate of every y_x, proving the inertia vanishing.

For (42.10) fix $q \notin S$ and $x = a/M + \sum_{r\in S} u_r/r$. Select τ in inertia at q inducing $v \in \mathbb{F}_q^\times$ on the q-th roots of unity with $L_q(v) = -1$. This is possible by the total tame ramification of the q-th cyclotomic field at q. It fixes the roots of unity of orders prime to q. Choose all roots $y_{x+u/q}$, $0 \le u < q$, to have the same reduction as y_x at the selected place above q. This is possible by Lemma 42.5 and injectivity on reduction of μ_{m_0}. It follows that

$$\tau^{-1} y_{x+u/q} = y_{x+v^{-1}u/q}.$$

In fact the two sides are roots of the same J_d and have the same reduction. Every element here is a local unit. The definition of F and triviality of the residue action of inertia give

$$F(\tau, x + u/q) \equiv \gamma(y_{x+u/q} - y_{x+v^{-1}u/q}) \pmod{\mathfrak{m}_q},$$

where reduction is still written additively in the unit group. There is no requirement that γ itself preserve the chosen prime.

Take integer lifts of the logarithms and reindex the second sum. The coefficient differences equal 1 modulo m_0 and sum to zero as integers. By Galois covariance, $\gamma(J_d(x + u/q) - J_d(x))$ is a principal unit at q, so after subtracting y_x from every term we can replace these coefficient differences by 1. Thus

$$\sum_{u\in\mathbb{F}_q^\times} L_q(u)F(\tau, x + u/q) \equiv \gamma\left(\sum_{u=0}^{q-1} y_{x+u/q} - qy_x\right) \pmod{\mathfrak{m}_q}. \tag{42.12}$$

The difference $\eta = \sum_{u=0}^{q-1} y_{x+u/q} - \varphi y_x$ is a root of unity: take its m_0-th power and use (42.6) and $\varphi J_d(x) = J_d(qx)$. On reduction the sum equals qy_x, which is the reduction of φy_x. Thus η is a root of unity reducing to 1 and has q-power order. Its conjugate $\gamma\eta$ also reduces to 1. Hence the right side of (42.12) is congruent to $\gamma\varphi y_x - q\gamma y_x$. This is the reduction of $F(\varphi, qx)$, as one may use φy_x as the root in its root-independent definition. Both sides of the desired identity lie in μ_{m_0}, on which reduction is injective. Thus the congruence is equality. Summing over a and the u_r, $r \in S$, yields the sum of $F(\varphi, qx)$ weighted by their logarithms. Reindex the arguments using $q \equiv 1 \bmod M$ and apply Lemma 42.6 to obtain (42.10).

For (42.11), choose τ in tame inertia at B whose action on an m_0-th root of p is translation by a generator ξ of μ_{m_0}. The Jacobi values of denominator M lie in a field unramified at

p, and $v_B(J_d(x)) = f_d(x)$. Dividing by a suitable power of p^{1/m_0} leaves a root of a unit, unramified at p, so

$$(\tau - 1)y_x = f_d(x)\xi, \qquad (\tau - 1)\gamma y_x = f_d(a_\gamma x)\xi.$$

As τ fixes μ_{m_0}, the definition of F gives $F(\tau, x) = (\epsilon(\gamma)f_d(x) - f_d(a_\gamma x))\xi$. Reindex the unit sum by $a \mapsto a_\gamma a$. Multiplication by the unit $\epsilon(\gamma)^e\psi(\gamma)$ then gives (42.11) whenever $\kappa_\varnothing(\tau) = 0$. □

This constructs the derivatives and proves their norm relation and bottom formula as used in `exists_kolyvagin_cocycles` in `Lib/Cyclotomic/Kolyvagin/ComponentBound.lean`. Notice why the helper parameter γ is necessary: roots of unity arising from the distribution relation need not be divisible in the finite cyclotomic field, but their contribution cancels in the commutator formula for F.

42.4 The odd component, local vanishing and reciprocity

We now fix the Dirichlet character of the desired parity, and use the component called `component` in `Lib/Cyclotomic/Kolyvagin/Components.lean`. It consists of locally constant cocycles $x : G_\mathbb{Q} \to O/\ell^\nu$ for $\chi^{-1} = (\epsilon^e\psi)^{-1}$ such that:

1. $x(c) = 0$;
2. x vanishes on inertia over $\mathbb{Q}(\mu_N)$ at all finite primes away from ℓ;
3. x vanishes on the decomposition groups over $\mathbb{Q}(\mu_N, \mu_{\ell^\infty})$ above ℓ.

We write $\mathfrak{C}_\nu(\psi, e)$ for this module. Conjugation of a cocycle of character λ obeys

$$x(hgh^{-1}) = \lambda(h)x(g) + (1 - \lambda(g))x(h). \tag{42.13}$$

This follows by applying the cocycle equation twice and using $x(h^{-1}) = -\lambda(h)^{-1}x(h)$. In particular, the restriction to a subgroup acting trivially on the coefficients is additive and transforms by λ on conjugation.

Let $d \equiv 1 \bmod N$ be prime to ℓ, and set $F = \mathbb{Q}(\mu_{Nm_0})$. For a place of $\overline{\mathbb{Q}}$ above ℓ, choose δ in inertia with $\epsilon(\delta) = d$ and fixing μ_N. This choice is possible by the surjectivity of local inertia on the pro-ℓ cyclotomic extension and the fact that it acts trivially on the prime-to-ℓ part of μ_N; the congruence $d \equiv 1 \bmod N$ gives triviality on the remaining part. If $g \in G_F$ belongs to the decomposition group at this place, then its commutator with δ lies in the decomposition group over $\mathbb{Q}(\mu_N, \mu_{\ell^\infty})$. Thus $x(\delta g\delta^{-1}) = x(g)$, since the restrictive character on G_F is trivial. By (42.13),

$$(d^e - 1)x(g) = 0. \tag{42.14}$$

The source allows an additional factor of 2, and we keep it in the next statements. This argument explains how the local condition at ℓ controls the ordinary, not merely the cyclotomic-kernel, Artin value.

Lemma 42.8 (Artin-law calculation)**.** *Let $x \in \mathfrak{C}_\nu(\psi, e)$, let $q \equiv 1 \bmod Nm_0$ be a rational prime distinct from ℓ, and let φ be an arithmetic Frobenius at a place B above q. For $u \in (\mathbb{Z}/Nm_0)^\times$ choose $s_u \in G_\mathbb{Q}$ acting on the Nm_0-th roots by u. Suppose $y \in F^\times$ has the following property at every place over a prime other than q, ℓ: either x is zero on the*

decomposition group, or y is a unit times an m_0-th power of an element of a local unramified extension. Then

$$\left(\sum_u v_B(s_u y)[u]^e\psi(u)\right)\, 2(d^e-1)x(\varphi)=0. \tag{42.15}$$

Here $v_B(q)=1$ and the valuations in the sum are integers.

Proof. The homomorphism $g\mapsto 2(d^e-1)x(g)$ on G_F is additive into the finite module O/m_0. By (42.14) its restriction to local decomposition groups above ℓ is zero. Since F is totally complex, its finite quotients are evaluated by global Artin reciprocity without real-place terms. Its Artin value on the principal idele y is zero. Off ℓ it is unramified and depends only on the local valuation. The valuation at a place of a local unramified extension has integer values on that extension for the base normalization, so divisibility by m_0 of the valuation of y in the second alternative makes the local contribution zero. The first alternative also gives zero. At q, full splitting in F makes the valuations integers and the primes correspond to the $s_u^{-1}B$. Frobenius at $s_u^{-1}B$ is conjugate to $s_u^{-1}\varphi s_u$, and (42.13) shows that its value of x is $[u]^e\psi(u)x(\varphi)$. Substitution into the sum of local Artin values proves (42.15). Equivalently one can apply every additive character of O/m_0 to the Artin sum and use separation of points by the characters of a finite abelian group. This is the calculation `artin_law_component` in `Lib/Cyclotomic/Kolyvagin/ComponentBound.lean`. □

Proposition 42.9 (Odd-component annihilation)**.** *Every member of $\mathfrak{C}_\nu(\psi,e)$ is killed by $4X_e^2$, for X_e of Proposition 42.2.*

Proof. Choose d as in that proposition. If $4X_e^2x\neq 0$, then $2X_e^2x$ does not vanish identically on G_F. Indeed, if it did, that cocycle would descend to the abelian quotient $\mathrm{Gal}(F/\mathbb{Q})$, since its coefficient character modulo m_0 also descends. It is normalized to zero at c, whose character is -1, and a cocycle with this property on an abelian group is killed by 2 by comparing its values at cg and gc. This would contradict the assumption. By Chebotarev we can find $q\equiv 1 \bmod Nm_0$ different from ℓ and a Frobenius φ over q with $2X_e^2x(\varphi)\neq 0$. Here and below Chebotarev is applied to a finite quotient through which the relevant cocycles and cyclotomic exponents factor, with conjugation allowed in the selection of the place. Apply the Jacobi family with $p=q$ and $y=J_d(1/(Nm_0))$. Its off-q unit property gives the hypothesis of Lemma 42.8; its valuations by Galois covariance are $f_d(u/(Nm_0))$. Thus $A_\nu(d)\,2(d^e-1)x(\varphi)=0$, using the notation of Lemma 42.3. That lemma gives

$$2(d^{e+1}-1)(d^e-1)X_e x(\varphi)=0.$$

But $(d^e-1)(d^{e+1}-1)$ divides X_e, so this implies $2X_e^2x(\varphi)=0$, a contradiction. □

At large primes a substantially stronger conclusion about the *number of detecting coordinates* is needed. Merely applying Proposition 42.9 to all cocycles cannot give that conclusion: the number of coordinates might grow with ℓ. For this reason the finite derivative classes are accompanied by a Kummer-duality calculation.

Lemma 42.10 (Reciprocity modulo the coefficient prime)**.** *Assume $\phi(N\ell)$ is a unit in O. Let $x\in\mathfrak{C}_\nu(\psi,e)$ be killed by ℓ. Let T be a finite set of rational primes, all $\equiv 1 \bmod N\ell$, and let κ be a locally constant cocycle for $\epsilon^{e+1}\psi$ with values in a finite O-module killed by ℓ, unramified on inertia over $\mathbb{Q}(\mu_N)$ away from ℓ and T. Suppose $q\in T$, x vanishes on the decomposition groups above $T\setminus\{q\}$, and*

$$2(d^e-1)x(\varphi)\neq 0$$

at an arithmetic Frobenius φ over q. Then κ vanishes on inertia at the chosen place B over q.

Proof. Set $F_1 = \mathbb{Q}(\mu_{N\ell})$. On G_{F_1}, κ is additive, since its character acts trivially on ℓ-torsion. Fix an inertia element τ at B and let $v = \kappa(\tau)$. The cyclic O-module Ov is a finite vector space over $\mathbb{F}_\ell$. Choose an $\mathbb{F}_\ell$-basis b_i of this vector space, and let L_i be extensions of its coordinate homomorphisms to the target ℓ-torsion module of κ. Choose a primitive ℓ-th root used to identify $\mathbb{F}_\ell$ with μ_ℓ. Kummer theory over F_1 associates to $L_i\kappa|_{G_{F_1}}$ an element $y_i \in F_1^\times/(F_1^\times)^\ell$, since F_1 contains μ_ℓ. If κ vanishes on inertia at a prime off ℓ, T, an ℓ-th root of y_i is unramified there. Thus the unit-times-ℓ-th-power condition in Lemma 42.8 holds for y_i. At primes of $T \setminus \{q\}$ the first alternative holds by hypothesis. The Artin-law calculation applies with modulus ℓ for the additive homomorphism $2(d^e-1)x|_{G_{F_1}}$; the local equation (42.14) remains true in its ℓ-torsion target O/ℓ^ν.

Let $a_{i,u} = v_B(s_u y_i)$, $u \in (\mathbb{Z}/N\ell)^\times$, and let $j \in \mathbb{F}_\ell$ be the action of τ on a fixed ℓ-th root of q. All the following equalities of coordinates are in $\mathbb{F}_\ell$. By local Kummer theory at $q \neq \ell$, the value on τ of the Kummer character of $s_u y_i$ is $j a_{i,u}$. Indeed after taking out the valuation factor the remaining root is a root of a unit and is fixed by inertia. On the other hand Galois action on Kummer characters gives

$$j a_{i,u} = L_i(u^{-e}\psi(u)^{-1}v).$$

To check this identity, conjugate τ by s_u^{-1} and then apply s_u to the roots of unity. The cocycle relation (42.13) multiplies $\kappa(\tau)$ by $\epsilon(s_u)^{-e-1}\psi(u)^{-1}$, and the action on μ_ℓ multiplies the result by u. This proves the identity and shows why the two powers in the dual character differ by one.

By the Artin-law calculation, $r_i = \sum_u [u]^e\psi(u)a_{i,u}$ annihilates the nonzero element $2(d^e-1)x(\varphi)$. Therefore each r_i belongs to the maximal ideal $\mathfrak{m}_O$. Using the preceding coordinate identity and summing over u, we obtain in the cyclic O-module Ov

$$\sum_i j r_i b_i = \sum_u [u]^e\psi(u) \sum_i j a_{i,u} b_i = \phi(N\ell)v.$$

Here integer lifts of j and the coordinates can be used since the module is killed by ℓ. The left side belongs to $\mathfrak{m}_O Ov$. Since $\phi(N\ell)$ is a unit, $v \in \mathfrak{m}_O Ov$. Nakayama's lemma gives $Ov = 0$. This holds for every τ, proving the assertion. This argument is integral even when the ℓ-torsion module is not a vector space over the residue field of O: the intermediate coordinates are deliberately taken over $\mathbb{F}_\ell$. □

42.5 The Kolyvagin–Rubin detection argument

We first isolate the finiteness and Chebotarev facts used by the induction. For a continuous action on a finite module V, continuous cocycles unramified outside a fixed finite set of primes form a finite set. In fact a cocycle is equivalent to a lift of the action to the semidirect product $V \rtimes \mathrm{Aut}(V)$. The degree of the finite Galois extension cut out by the lift is bounded by $|V||\mathrm{Aut}(V)|$, and the extensions in question are unramified outside the fixed set (including the set of ramification of the action). Hermite–Minkowski gives finiteness of extensions of bounded degree unramified off that set; one can obtain the necessary discriminant bound either from the finiteness of local extensions of bounded degree or the usual local different bound in terms of degree. A finite Galois group has only finitely many homomorphisms to the fixed target.

Another useful elementary fact is that a cocycle for a scalar character λ on $G_\mathbb{Q}$, zero at an element γ where $\lambda(\gamma) - 1$ is a unit, cannot factor nontrivially through an abelian quotient on which the action is defined. To prove this, compare the cocycle at γg and $g\gamma$. In particular, if a nonzero such cocycle is zero on $G_{\mathbb{Q}(\mu_L)}$ for a modulus L divisible by the conductor of its

coefficient character, we would get a contradiction. Given two such cocycles, with possibly different actions or different normalizing elements, we may consequently find an element of $G_{\mathbb{Q}(\mu_L)}$ at which both are nonzero. Indeed the two zero subgroups of the restricted cocycles are proper, and the union of two proper subgroups cannot cover a group. Chebotarev realizes the selected pair of nonzero values at an arithmetic Frobenius at a prime $\equiv 1 \bmod L$. We will use this with arbitrarily large but finite additional factors in L. If the first cocycle is normalized at c with odd character, then -2 is a unit as soon as $\ell > 2$.

Theorem 42.11 (Bounded number of detecting points for odd components)**.** *If $\ell > N+e+3$, there are at most $v_\ell(X_e)+1$ elements of $G_{\mathbb{Q}}$ such that evaluation at these elements is injective on $\mathfrak{C}_\nu(\psi, e)$.*

Proof. Write $b = v_\ell(X_e)+1$. We may use b positions, repetitions allowed. Suppose no such list detects all the elements. Given a list of at most b elements, there is then a nonzero member of the component vanishing at all of them. By multiplying it by a suitable power of ℓ, we can arrange it is killed by ℓ and remains nonzero. Denote the set of nonzero ℓ-torsion classes of the component by Z. This set is finite by the preceding argument.

Since $\ell > N + e + 3$, the Chinese remainder theorem gives $d \equiv 1 \bmod N$ with reduction modulo ℓ generating $\mathbb{F}_\ell^\times$. In particular $\ell \nmid d$ and $d^e - 1$, $d^{e+1} - 1$, 2, and $\phi(N\ell)$ are all units in O. Choose an element γ of inertia over ℓ fixing μ_N with cyclotomic value d, as in (42.14). The scalar $\epsilon(\gamma)^{e+1}\psi(\gamma) - 1 = d^{e+1} - 1$ is a unit.

For a finite set T of rational primes $\equiv 1 \bmod N\ell$ not dividing ℓ, let $Y(T)$ denote the module of locally constant cocycles with values in O/ℓ^b, of character $\epsilon^{e+1}\psi$, vanishing on inertia over $\mathbb{Q}(\mu_N)$ off ℓ, T. Write $Y = Y(\varnothing)$. Say that T is good if each member of $Y(T)$ killed by ℓ belongs to Y. The empty set is good, and all the sets below are finite.

First we claim that whenever T is good and $|T| \le b$, some $z \in Z$ has the property that every $q \equiv 1 \bmod N\ell$ for which z does not vanish on a decomposition group over q lies outside T and makes $T \cup \{q\}$ good. Choose an arithmetic Frobenius over each prime of T. By our supposition we can take $z \in Z$ vanishing at them. Since those primes split in $\mathbb{Q}(\mu_{N\ell})$, all inertia is in $G_{\mathbb{Q}(\mu_N)}$ and the coefficient action on their decomposition groups is trivial on ℓ-torsion. The inertia condition on z gives vanishing at each such entire decomposition group, and (42.13) gives vanishing also at its conjugates (the action on ℓ-torsion is trivial there). Hence q as in the claim is outside T and $z(\varphi_q) \neq 0$ at a Frobenius over q. Lemma 42.10 applies to any ℓ-torsion member of $Y(T \cup \{q\})$: it forces the cocycle to vanish on inertia at q. Conjugation makes the same true at all places over q. This member belongs to $Y(T)$ and hence to Y, proving the claim.

The set Y is finite; its ramification over $\mathbb{Q}$ is confined to the primes dividing $N\ell$. For every pair

$$(y, z) \in \{y \in Y : y \neq 0,\ y(\gamma) = 0\} \times Z$$

choose a prime $q(y,z) \equiv 1 \bmod dN\ell^b$ and an arithmetic Frobenius $\varphi(y,z)$ above it at which both values $y(\varphi(y,z))$ and $z(\varphi(y,z))$ are nonzero. This is possible by the two-cocycle Chebotarev argument above, since y is normalized at γ and z at complex conjugation. Let $\mathcal{P}$ be the resulting finite set of primes. It is not important how large $\mathcal{P}$ is. Only the cardinality of the final detecting list is being bounded.

Apply the claim with $T = \varnothing$, and select $p \equiv 1 \bmod N\ell^b \prod_{q\in\mathcal{P}} q$, outside $\mathcal{P}$, at which that z has a nonzero value at Frobenius. The singleton $\{p\}$ is then good. Construct the cocycles κ_S, $S \subseteq \mathcal{P}$, using Proposition 42.7 at modulus $m_0 = \ell^b$, with the chosen γ, d, p.

We prove by descending induction on $i = b, b-1, \dots, 0$ the assertion

$$|S| \le i, \quad S \subseteq \mathcal{P}, \quad \{p\} \cup S \text{ good} \quad \Longrightarrow \quad \ell^i \kappa_S \in Y. \tag{42.16}$$

At $i = b$ the left-hand cocycle is zero. Suppose the assertion known at $i+1$, and take S at stage i. If $y = \ell^{i+1}\kappa_S$ were nonzero, it would belong to Y by induction, and be normalized to zero at γ. By the claim with $T = \{p\} \cup S$, of size at most $i+1 \le b$, choose $z \in Z$ as above. The prime $q = q(y,z)$ belongs to $\mathcal{P}$ and its selected Frobenius has both y and z nonzero. The claim shows that $q \notin T$ and $T \cup \{q\}$ is good. By (42.16) at $i+1$, $\ell^{i+1}\kappa_{S\cup\{q\}}$ is unramified at q. But the derivative relation (42.10) shows that its value on a tame generator is $y(\varphi(y,z)) \neq 0$, a contradiction. Therefore $\ell^{i+1}\kappa_S = 0$. Since $\ell^i\kappa_S$ is a member of $Y(T)$ killed by ℓ and T is good, it belongs to Y. This finishes the induction.

At the bottom we obtain $\kappa_\varnothing \in Y$, so it is unramified at p. Equation (42.11) shows $A_b(d) = 0$ in O/ℓ^b because $\epsilon(\gamma)^{e+1}\psi(\gamma) - 1$ is a unit. Lemma 42.3 now implies that $(d^{e+1}-1)X_e = 0$ in O/ℓ^b, which is impossible by $v_\ell(X_e) = b-1$. The contradiction proves the theorem. □

This is `exists_detecting_points_component` in `Lib/Cyclotomic/Kolyvagin/ComponentBound.lean`. The descending induction uses the good-set condition only for ℓ-torsion members of $Y(T)$. This matters: a claim that every member of a module at an auxiliary level is already unramified could be false, and would not be needed. Likewise the finite set $\mathcal{P}$ is not the eventual detecting set; its size may be very large. The proof only uses it to arrange implications simultaneously over the finite coefficient ring.

42.6 Uniform detection for the small primes

The preceding component argument works when ℓ is large compared with the conductor and exponents. For small ℓ we use a deliberately crude bound, independent of the torsion depth. We spell out how it can be made independent of the actual coefficient ring too.

Let $E \subseteq \mathbb{Q}(\mu_N)$ and set $K_0 = \mathbb{Q}(\mu_{N\ell^2})$. Write $D_0 = [K_0 : \mathbb{Q}] \le N\ell^2$ in this subsection. The group of continuous homomorphisms $G_{K_0} \to \mathbb{Z}/\ell$ unramified off ℓ factors through the maximal elementary abelian ℓ-quotient of the abelian Galois group unramified off ℓ. By ray class theory, its dimension over $\mathbb{F}_\ell$ is at most

$$\log_\ell h_{K_0} + D_0 + 2\,\#\{w : w \mid \ell\}. \tag{42.17}$$

Indeed the maximal pro-ℓ quotient of the ray class exact sequence is a quotient of an extension of the ℓ-part of the ordinary class group by the product of the pro-ℓ unit groups at the places above ℓ. The elementary generator count for the class group is at most $\log_\ell h_{K_0}$. For a finite extension $K_{0,w}/\mathbb{Q}_\ell$, the pro-ℓ unit group has at most $[K_{0,w} : \mathbb{Q}_\ell] + 2$ generators: a group of finite index is via the logarithm a $\mathbb{Z}_\ell$-lattice of rank the local degree, and the torsion of the unit group is cyclic; allowing two instead of one covers all small-prime conventions. Equivalently use the classification of finitely generated abelian pro-ℓ groups. Summing the local degrees proves (42.17).

We give one explicit estimate on the class-number term. The cyclotomic discriminant formula gives $|D_{K_0}| \le (N\ell^2)^{D_0}$. Minkowski's bound provides an integral ideal in each class of norm at most $\sqrt{|D_{K_0}|}$, since the remaining factors in the usual bound are at most 1. If L is any number field of degree d and $R \ge 1$, the number of ideals in L of norm at most R is bounded by $(\lfloor R \rfloor + 1)^{d+1}$: at each rational prime there are at most d prime ideals over it, and the coefficient of norm n is no larger than the d-fold divisor-counting function $d_d(n)$, by inserting the possible exponents with their residue degrees. The sum of $d_d(n)$ counts tuples of positive integers each at most R, hence is at most $(\lfloor R \rfloor + 1)^d$; the extra power is harmless. Consequently

$$\log_2 h_{K_0} \le (D_0 + 1)\bigl(1 + \tfrac{1}{2} D_0 \log_2(N\ell^2)\bigr).$$

For $y \geq 3$, $N \leq y$ and $\ell \leq y^2$, one has $D_0 \leq y^5$. The last display, plus $3D_0$ from the local terms of (42.17), is bounded by $y^{20} - D_0$, using $\log_2 y \leq y$ and $y \geq 3$. For example the terms are bounded by $(y^5+1)(1+(5/2)y^6)+3y^5 \leq y^{20}-y^5$ for $y \geq 3$.

Choose a list of s Galois elements of G_{K_0} whose images generate the exponent-ℓ quotient just described, and add a representative of every coset of G_{K_0} in G_E. There are at most D_0 such representatives. The resulting list has at most y^{20} elements at the stated sizes.

Lemma 42.12 (A uniform detecting set for Tate cocycles)**.** *The list just constructed detects all locally constant cocycles $z : G_E \to O/\ell^\nu$ for the character ϵ^{-e} unramified outside ℓ, for all integers $e \geq 0$ and all depths $\nu \geq 1$. The list can be chosen independently of e, ν, O.*

Proof. Suppose a cocycle vanishes on the list. If it were nonzero, some power of ℓ times it would be nonzero and killed by ℓ. Replace it by that multiple. On G_{K_0} the coefficient action on ℓ-torsion is trivial. Its restriction is then an additive homomorphism unramified outside ℓ, with values in a finite elementary abelian ℓ-group. Since all its scalar coordinates in $\mathbb{Z}/\ell$ kill the generator list, the restriction is zero. For any $g \in G_E$ write $g = hu$ with $h \in G_{K_0}$ and u a chosen representative. The cocycle equation gives $z(g) = z(h) + \epsilon(h)^{-e} z(u) = 0$, contradicting nonzero. This proves the assertion. □

The exact uniformity of this proof is useful also in the triangular construction, where no bound on the even class group is invoked. This is the arithmetic behind `exists_detecting_points_cocycles` in `Lib/Cyclotomic/OddSha/OddShaBound.lean`; use of the stronger ℓ^2 cyclotomic field in the detecting list is harmless at every depth, including $\ell = 2$. For the count we used elementary class field theory and the Minkowski bound, not an Iwasawa-theoretic estimate.

42.7 Returning from Dirichlet components to the CM field

We now combine the preceding results without assuming that the CM field has degree prime to ℓ. Let $G = G_{\mathbb{Q}}$, $H = G_E$, and $K = G_{\mathbb{Q}(\mu_N)} \subset H$. All are viewed using an embedding into $\overline{\mathbb{Q}}$; H is normal in G because E is contained in the cyclotomic field. For $e > 0$, let x be an element of the odd group `shaOdd` of the source, i.e. a locally constant cocycle on H for ϵ^{-e} with values in O/ℓ^ν, unramified off ℓ, satisfying the prescribed vanishing on cyclotomic-kernel decomposition groups at ℓ and

$$x(cgc^{-1}) = -x(g) \quad (g \in H).$$

Here the local condition is read on decomposition elements whose full cyclotomic character is trivial. At $e = 0$ the source instead uses the class-group convention discussed below.

For $u, g \in G$ write $b_u(g) = u(cgc^{-1}g^{-1})u^{-1} \in K$ (a commutator has trivial action on all roots of unity). Let $\mathcal{S}$ be any set of representatives of $K\backslash G$ with 1 representing the identity coset. Define for each Dirichlet character ψ modulo N

$$z_\psi(g) = \sum_{u \in \mathcal{S}} \epsilon(u)^e \psi(u) x(b_u(g)).$$

There is no division by $|G/K|$ here. This will keep the argument valid also at small primes.

Lemma 42.13 (Integral decomposition by commutators)**.** *Each z_ψ is a locally constant cocycle of the character $(\epsilon^e\psi)^{-1}$, zero at c, and satisfying the component local conditions of Section 42.4. For $g \in H$ one has*

$$4\phi(N)x(g) = \sum_{\psi} \big(z_\psi(cgc^{-1}) - z_\psi(g)\big). \tag{42.18}$$

In the sum on the right only characters with $\epsilon^e(c)\psi(c) = -1$ can contribute.

Proof. If u is replaced by hu for $h \in K$, the value of $x(b_u(g))$ is multiplied by $\epsilon(h)^{-e}$. This is (42.13), since $b_u(g)$ has trivial cyclotomic character. The weight $\epsilon(u)^e\psi(u)$ is multiplied by the reciprocal factor, proving independence of representatives. Next

$$b_u(gh) = b_u(g)b_{ug}(h).$$

The action of each factor on coefficients is trivial, so substituting into the sum, changing representatives in the second part, gives the cocycle equation of the claimed character. We have $b_u(c) = 1$, so the value at c is zero. For g in the specified inertia or decomposition groups of the component, $b_u(g)$ is a product of conjugates of g and of g^{-1} under elements of G. These factors belong to H (indeed $g \in K$) and are in the corresponding local vanishing sets of the cocycle x: away from ℓ they are inertia elements, while above ℓ they have trivial full cyclotomic action. Thus all the z_ψ meet the local conditions. Finite sums and continuity give local constancy.

Assume O contains a primitive root of unity of order the exponent of $(\mathbb{Z}/N)^\times$, as in the source. The character orthogonality identity $\sum_\psi \psi(u) = \phi(N)\mathbf{1}_{u=1}$ holds integrally over O. Consequently

$$\sum_\psi z_\psi(g) = \phi(N)x(cgc^{-1}g^{-1}) = -2\phi(N)x(g) \qquad (g \in H).$$

The last equality uses the odd condition and the cocycle formula at the product. Apply it also to cgc^{-1}, subtract, and obtain (42.18). As $z_\psi(c) = 0$, the cocycle formula gives $z_\psi(cgc^{-1}) = \epsilon(c)^{-e}\psi(c)^{-1}z_\psi(g)$, so terms of even parity vanish exactly. The sum-and-difference identity, without division by 2, proves the assertion also over coefficient rings above 2. □

For each odd character, Proposition 42.9 says that $4X_e^2 z_\psi = 0$. Lemma 42.13 therefore gives

$$Y_e x = 0, \qquad Y_e = 4\phi(N) \cdot 4X_e^2. \tag{42.19}$$

This annihilates the entire odd group. We next control its number of coordinates. Fix $t \geq 1$, $e < t$, and $y_0 = N + t + 3$. For $\ell \leq y_0^2$, Lemma 42.12 gives at most y_0^{20} detecting elements. For $\ell > y_0^2$, Theorem 42.11 gives a list $\gamma_{\psi,i}$ of at most $v_\ell(X_e) + 1$ elements detecting z_ψ at each odd parity. Evaluate x on the elements $b_u(\gamma_{\psi,i})$ for the $\phi(N)$ choices of u, the $\phi(N)$ or fewer choices of ψ, and the indicated list of i. If all such evaluations are zero, every odd z_ψ is zero, so $4\phi(N)x = 0$. For the large primes under consideration $4\phi(N)$ is a unit, hence this implies $x = 0$. The number of evaluations is bounded by

$$\phi(N)^2(v_\ell(X_e) + 1) \leq y_0^2(y_0^6 + 1) \leq y_0^{20}.$$

Thus a bound on the number of detecting coordinates holds at every prime.

A finite-length submodule of q copies of O/ℓ^ν killed by Y_e has normalized length at most $q\, v_\ell(Y_e)$, by the elementary divisor description of the kernel of multiplication by Y_e. Applying this to the injective evaluation map and (42.19) gives $\lambda(\mathrm{shaOdd}) \leq y_0^{20} v_\ell(Y_e)$ for $e > 0$. At $e = 0$ the source's local definition gives a subgroup of the homomorphisms from the ordinary ideal class group of E to O/ℓ^ν: the cocycles for the trivial character are homomorphisms and the local conditions impose unramifiedness also at ℓ. A cyclic decomposition of the finite ideal class group shows that this subgroup has normalized length at most $v_\ell(h_E)$.

Theorem 42.14 (The global integer bounding odd classes)**.** *There exists a positive integer*

$$X = h_E \prod_{1 \leq e < t} Y_e^{y_0^{20}} \leq h_E\, 2^{4(N+t+3)^{27}}$$

such that at every prime ℓ, every torsion depth and every $0 \le e < t$, the normalized length of the odd module used in the congruence argument is at most $v_\ell(X)$. The coefficient ring is assumed to contain enough roots of unity of order the exponent of $(\mathbb{Z}/N)^\times$.

Proof. The length assertion was verified above. By Proposition 42.2,

$$\log_2 Y_e \le 4 + \log_2 \phi(N) + 2(N+e+3)^6 \le 3y_0^6.$$

There are at most $t-1 < y_0$ factors, so $\log_2(X/h_E) \le 3y_0^{27} \le 4y_0^{27}$. This proves the statement with the same exponent as the source. □

Theorem 42.14 reconstructs the bound `exists_length_shaOdd_le` in `Lib/Cyclotomic/OddSha/OddShaBound.lean`, as used in Proposition 40.16. Its proof uses the usual classical results of cyclotomic and global class field theory, Chebotarev, the elementary Kummer correspondence, and Dirichlet functional equations; it does not invoke the cyclotomic main conjecture or a bound on the even class groups. The relatively large exponent is a convenient consequence of the difference between small-prime class-number counting and large-prime Kolyvagin detection. The argument deserves emphasis: a coarse annihilator plus a count of detecting coordinates is enough for the sum of normalized lengths. Counting the derivative primes or estimating the full ramification degree of the Euler system is neither needed nor compatible with this uniform budget.

42.8 Uniform cardinality of Selmer pencils

We return to the notation of Chapter 40: E is the CM field of the datum, and m is the number of arrows. This section proves the uniform finiteness part of the simultaneous-twist theorem, conditional on the arithmetic mirror criterion (the difficult global reciprocity step isolated in Proposition 40.9). The earlier chapter explains the tilting lemma; we now give the precise induction on the torsion depth and the subsequent specialization that uses it.

Fix a finite collection of rows R_i of ranks n_i over a DVR O, balanced at depth M, and write π for a uniformizer. We retain the source notation β_i for their ordinary length allowances. For a divisor a of ℓ^M, for a finite set of pencil places J, and for each row let

$$X_i(J,a) \subset Z^1(G_E, (O/(a))^{n_i})$$

denote the cocycle module with the old crystalline relations, vanishing at the normalizer γ_i, unramified outside the union of the old set S and $J \cup J^c$, and with the ties at all places of J. The ties use values at tame generators τ_P, conjugate inverses $(\tau_P^c)^{-1}$, Frobenii φ_P, and $(\varphi_P^c)^{-1}$, as in `IsPencil` and `tieRelations` of `New/Selmer/Corepresentable.lean`. For a place where the slope of row i is nonzero, the row representation on both Frobenius elements is trivial modulo ℓ^M; all represented auxiliary places are unramified for the original row.

The following observations concern the hypotheses of the induction. At a new place of residue characteristic different from ℓ, a torsion cocycle for an unramified representation is a homomorphism on inertia. Its restriction factors through the pro-ℓ tame quotient; indeed wild inertia is pro-p for the residue prime $p \neq \ell$. Thus requiring its value at the chosen generator of this quotient to be zero makes it unramified. At conjugate places we use the other generator as well. This proves that, in a relaxed module with an extra pair of ramification places permitted, the additional conditions of zero tame evaluations are exactly the old unramified conditions. If zeros at all auxiliary tame generators are imposed, the module is contained in the fixed module with only the old ramification set. That latter module has finite length for finite coefficients by the finiteness argument in Section 42.5. Its length is independent of the depth

N of the auxiliary characters. All linear relations, including the cocycle equations and the old relations, commute with the injections

$$O/\pi^{k-1} \longrightarrow O/\pi^{k}, \qquad x \longmapsto \pi x.$$

In particular this map identifies the smaller-depth cocycle module with the subset of the larger module whose values all lie in $\pi O/\pi^{k}$.

Proposition 42.15 (Construction of a bounded pencil from the mirror criterion)**.** *Suppose the mirror criterion of Proposition 40.9 holds for the row modules just described, with the one-prime compatibility and conjugate-inverse conventions of* `IsPencil`*. Assume that at every depth N one can select a helper place with its Frobenius in the required neighborhood and with all its slopes zero, and that the residue field has more than $4n_i + 2$ elements for all i. Then there exists an integer J_0 independent of N such that, for each N, there is a pencil J of at most J_0 places with*

$$\lambda_O(X_i(J,a)) \le \beta_i \qquad \text{for all } i.$$

This holds at every fixed divisor a of ℓ^M.

Proof. Induct on $k = v_\pi(a)$. For $k = 0$ the modules vanish, and the helper place gives a pencil of size one for every N. Suppose a uniform bound J_1 obtained at $k-1$. For each i let S_i^k be the cocycle module modulo π^k with original ramification set S, old relations, and normalization but without new ties. Put

$$C_k = J_1 + \sum_i \bigl(\lambda_O(S_i^k) + 2n_i k J_1\bigr).$$

By induction, for every N there is a pencil with $|J| \le J_1$ and all the required bounds at $k-1$. Evaluate the module $X_i(J,\pi^k)$ on both tame generators at every new place. Its kernel lies in S_i^k, and its image has length at most $2n_i k|J|$. Hence this pencil satisfies

$$|J| + \sum_i \lambda_O(X_i(J,\pi^k)) \le C_k. \tag{42.20}$$

This is a bound independent of N, even though the two pencils for different N may have completely different places.

Among pencils satisfying the $k-1$ bounds and (42.20), choose one minimizing the sum of the lengths at depth k. Such a minimum exists as the set is nonempty and the sum takes nonnegative integer values. Suppose some i has $\lambda_O(X_i(J,\pi^k)) > \beta_i$. There must be a cocycle in this module with at least one unit coordinate: otherwise all cocycles have values in $\pi(O/\pi^k)^{n_i}$, and the injective identification with $X_i(J,\pi^{k-1})$ above contradicts the strict inequality.

Apply the contrapositive of the mirror criterion at depth k. It supplies a place $P \notin J$, admissible for the pencil for every choice of its new slope at row i and slopes zero at the others, with the following failure of inclusion. For $a = \pi^k$ let Y_i be the module relaxed at P, P^c, with the old ties unchanged; let $\Phi_0 : Y_i \to (O/(a))^{2n_i}$ evaluate the two tame generators, and let Φ_1 evaluate the two Frobenius elements. Then

$$X_i(J,a) = Y_i \cap \ker \Phi_0, \qquad \Phi_1(X_i(J,a)) \not\subseteq \Phi_0(Y_i).$$

For a new slope $t = s^{-1}$, the additional ties are the kernel of $s\Phi_0 - \Phi_1$. Use Lemma 40.8 both at $a = \pi^k$ and at $a = \pi^{k-1}$. The prohibited residue classes of s at the two depths together have cardinality at most $4n_i$. Since zero must also be excluded, the size assumption on the

residue field permits a unit choice of s. At both depths the length of the new tied module is at most the length of

$$Y_i \cap \ker \Phi_0 \cap \Phi_1^{-1}(\Phi_0 Y_i).$$

At the larger depth the failed inclusion makes this a proper submodule of $X_i(J, \pi^k)$ and so the ordinary length drops by at least one. At the smaller depth it is a submodule of $X_i(J, \pi^{k-1})$, preserving the required bound. For the other rows choose slope zero. Then the new ties simply impose the original unramified condition at the new places and hence do not increase either length.

The new pencil increases $|J|$ by one and decreases the sum of lengths at depth k by at least one. Thus (42.20) remains valid and the smaller-depth bounds remain valid, contrary to minimality. It follows that every row has length at most β_i at the larger depth. In particular the new uniform bound on the cardinality can be taken to be C_k. Induction proves the proposition. □

This is the length argument of `exists_isPencil` in `New/Selmer/TwistTheorem.lean`. It uses the mirror criterion, rather than an unconditional assertion that adjoining a generic prime always lowers a Selmer rank. It also uses the helper- and enlarging-prime compatibility for all choices of the new slope; if the new place depended on the slope, Lemma 40.8 would not have fixed linear maps to which to apply it. The existence of the helper itself follows from the auxiliary anticyclotomic construction of Section 41.4, applied with no prior nonhelper primes. The compatibility of an enlarging prime with the mirrors is part of the stated criterion and remains a specialized global reciprocity input in this reconstruction.

42.9 Dropping ties along a line of approximate characters

We make the intermediate approximate-character calculation explicit. At a pencil of auxiliary depth N, denote the helper by P_0 and the characters of its places by $h_P : G_E \to \mathbb{Z}/\ell^N$. If $y = (y_P)$ is an actual tuple of ℓ^N-th roots, the associated anticyclotomic character is

$$\xi_y(g) = \prod_{P \in J} y_P^{[h_P(g)]_{\ell^N}}.$$

The character takes the value y_P on both chosen tame generators at P and the conjugate place. At each nonhelper Frobenius, on the two sides, it takes the helper coordinate y_{P_0}. This follows directly by multiplying the character values: the matrix of tame evaluations $h_Q(\tau_P)$ is the identity; for nonhelper P the column $(h_Q(\varphi_P))_{Q \in J}$ is the indicator of $Q = P_0$. Anticyclotomicity changes conjugation into inversion, giving the same values at the conjugate inverses. This is `IsPencil.pointChar_idOrConjInv` in `New/Selmer/TiltingLemma.lean`.

The special polynomial point for a single row i has

$$y_{P_0} = u, \qquad y_P = 1 + t_{P,i}(u-1) \quad (P \neq P_0), \qquad u^{\ell^k} = 1. \tag{42.21}$$

In general only the helper coordinate is a true root of unity. By the elementary power calculation of (40.11), each $y_P^{\ell^{k+M}}$ is 1 modulo ℓ^M. Thus for $N \geq k + M$ the pointwise twisted matrices give a representation on O'/ℓ^M over any sufficiently large coefficient ring O' in which these points are defined. All original relations in the cocycle problem still make sense after evaluation.

Lemma 42.16 (An affine form of the tame relation)**.** *Let R be an unramified representation at a new place P with residue cardinality q. Let B be the coefficient ring modulo ℓ^M over O', and twist R by the polynomial point (42.21). Assume $N \geq k + 2M$, $q \equiv 1 \bmod \ell^N$. For every*

cocycle c on the twisted representation T, at a selected tame generator τ and Frobenius φ of this auxiliary place, one has

$$(T(\varphi)-1)c(\tau) = (T(\tau)-1)c(\varphi). \tag{42.22}$$

The same formula holds on the conjugate-inverse side.

Proof. The cocycle corresponds to the affine action $A(g) : v \mapsto T(g)v + c(g)$ on B^{n_i}. On inertia, its linear part has ℓ-power order, and wild inertia maps trivially to the affine group: all its linear images and translational images lie in ℓ-groups, whereas wild inertia is pro-p for a different prime p. At the tame generator the linear part is the scalar y_P. Its ℓ^{k+M}-th power is 1 by the approximate-root calculation, so $A(\tau)^{\ell^{k+M}}$ is a translation. The additive group B^{n_i} is killed by ℓ^M, hence $A(\tau)^{\ell^{k+2M}} = 1$. The tame local Galois relation is $\varphi\tau\varphi^{-1} = \tau^q$ in the tame quotient. It therefore gives commuting affine maps, since $q \equiv 1 \bmod \ell^N$. Comparing their values at zero gives (42.22). At the conjugate place the inverses obey the same commuting relation, which is all that is needed in the calculation. □

Proposition 42.17 (Length cost of dropping ties)**.** *At the row-specific point (42.21) with $N \geq k + 2M$, remove the ties at a subset I of the pencil places. If the helper belongs to I, assume $u = 1$. For $P \in I$, choose matrices $R(w_{P,b})$ agreeing modulo ℓ^M with $R(\varphi_{P,b})$ at the two sides b (with inverse conjugation as above). The increase in normalized length, measured relative to the module with these ties, is bounded by*

$$\sum_{P\in I} \sum_{b} v_\ell \det(uR(w_{P,b}) - 1).$$

Proof. For P nonhelper the difference defining the tie is $d_c = c(\tau) - t_{P,i}c(\varphi)$. The scalar of $T(\tau)$ is $1 + t_{P,i}(u-1)$ and $T(\varphi) = uR(\varphi)$ modulo ℓ^M. If $t_{P,i} = 0$, (42.22) immediately gives $(T(\varphi)-1)d_c = 0$. If $t_{P,i} \neq 0$, the pencil condition gives $R(\varphi) = 1$ modulo ℓ^M, and (42.22) gives $(u-1)d_c = 0$ again. Thus in both cases d_c takes values in the kernel of $uR(\varphi) - 1$ on B^{n_i}. If $P = P_0$, the assumption $u = 1$ makes all the parameters 1, and the same identity holds for the expression of the original tie: when the slope is nonzero $R(\varphi) = 1$, while when it is zero we use unramifiedness at τ. Consequently d_c takes values in the stated kernel in all cases. Imposing its vanishing decreases length by at most the length of that kernel. Replacing $R(\varphi)$ by $R(w_{P,b})$ makes no difference modulo ℓ^M. By Smith normal form the normalized length of the kernel is at most the valuation of the determinant (with an infinite bound allowed when the determinant vanishes). Summing these estimates over P, b proves the proposition. □

The last result is the precise estimate `IsPencil.length_le_of_drop_ties`. We used an affine action because the action on coefficients alone does not justify commuting the cocycle values at the tame generator and at Frobenius. The extra M in the requirement $N \geq k + 2M$ kills the translations in the affine group.

42.10 The two specializations in the common-twist theorem

We now give the quantitative deduction of Proposition 40.14, conditional on the mirror criterion, the existence of balanced row lattices, and the integral measure assertion for the value families, as in the compact source. The relevant extra inputs are specified rather than replaced by a qualitative twisting theorem.

First note the uniform finite-coordinate argument for the cocycle modules. Choose an open normal subgroup U on which the reductions of the row matrices, and the finitely many

characters defining the pencil, are trivial. Let $V \subset U$ be the intersection of the kernels of all additive homomorphisms from U to $\mathbb{Z}/\ell^M$ that are unramified at the places of its field outside the stipulated finite set, and take its normal core in G_E. There are finitely many such homomorphisms, by the class-field or bounded-degree argument of Section 42.5. The resulting open normal subgroup V is independent of the point y and the coefficient ring used for its evaluation. Indeed on U the action is trivial on the torsion module, so the restriction of any cocycle is additive there; the additive characters into the subgroup of $\mathbb{Q}/\mathbb{Z}$ killed by ℓ^M detect every nonzero value, showing the restriction kills V. As $c(gv) = c(g) + T(g)c(v)$, it follows that all such cocycles are determined by their values on the finite set G_E/V.

Write these coordinates as a finite column over O'/ℓ^M. The cocycle relations and the desired inertia and other linear conditions form a relation module in a finite free polynomial module over $O[Y_P]$, using least nonnegative representatives of $h_P(g)$ in the exponents. Include ℓ^M times the free module. This presentation is valid for every tuple for which the twisted matrix action is defined modulo the torsion modulus, including (42.21). In particular it is noetherian even if the old linear conditions are expressed as an arbitrary set of finitely supported relations. Lemma 40.11 supplies a determinant polynomial which realizes the normalized length at one selected point exactly and bounds the length from above at all other specializations. The coefficient ring is allowed to enlarge finitely after the selected point has been found: normalized lengths of finite modules and valuations of determinant polynomials are preserved by finite extension of DVRs at a point over the smaller ring. This follows either from faithful flatness or from multiplying the elementary-divisor lengths and the normalization factor by the same ramification index. This explains the compatibility of `exists_corepresenting_coords` with `exists_sharp_fitting_poly` in the source.

Write β_i for the ordinary balanced length bound and

$$B_i \geq \frac{\beta_i}{\lambda_O(O/\ell)} + \sum_b v_\ell \det(R_i(z_b) - 1), \qquad z_1 = z, \quad z_2 = (z^c)^{-1},$$

the assumed real upper bounds in Proposition 40.14. Refine the allowed neighborhood of z so that the matrices of the helper Frobenius on both sides agree with these matrices modulo ℓ^M. For the finite family of Euler-factor characters μ_j, choose the neighborhood also so that $v_\ell(\mu_j(\varphi_{P_0}) - 1) \leq c$; this is possible because the analogous inequality holds at z, and this is an open condition when the upper bound is finite. The bound J_0 in Proposition 42.15 now depends on that neighborhood but not on N.

Let $\eta > 0$ be the desired final error, and choose $\delta > 0$ with $(8J_0 + 2)\delta < \eta$.

First selection. Choose k by Lemma 40.13 in one variable, with the finite row and value bounds described below, and with an upper bound on the total degree of the monic polynomials to be avoided. These upper bounds depend only on the row ranks, the number of fixed value families and J_0, not on the number or degrees of the cocycle relations. The requisite k can consequently be fixed before selecting the pencil. Next take $N \geq k + 2M$ sufficiently large for the second selection below. Choose the pencil with at most J_0 places and the tied length bounds of Proposition 42.15.

For row i take a sharp Fitting polynomial $G_i(Y)$ at $\mathbf{1}$, retaining all ties except the helper tie. At $\mathbf{1}$ the normalized length without the helper tie is at most B_i: apply Proposition 42.17 with $u = 1$ and $w_{P_0,b} = z_b$. Therefore $v_\ell(G_i(\mathbf{1})) \leq B_i$. Restrict G_i to the polynomial line (42.21) and include this as a polynomial in the variable u. Its value at $u = 1$ gives the needed lower bound for root selection.

For every value family V include also the polynomial $V(\{P_0\}, \xi_u)$, where ξ_u is the character of the helper alone at value u. This polynomial exists and is integral by the finite-measure

formula (40.13). At $u = 1$, the insertion inequality (40.14) and the helper neighborhood imply

$$v_\ell V(\{P_0\}, 1) \le v_\ell V(\varnothing, 1) + 8c.$$

This gives a finite initial bound independent of the pencil. Finally include the monic polynomials obtained from $\det(uR_i(\varphi_{P,b}) - 1)$ for every nonhelper P and both sides b. Their leading coefficients $\det R_i(\varphi_{P,b})$ are units, so one can divide by them. Include the monic linear factors for the possible bad Euler values $\mu_j(\varphi_{P,b})u^{\pm 1}$: up to multiplication by a unit at a root u, their distance from 1 is the distance from a specified unit. The number and degrees of all these monic polynomials have the required uniform bound.

Select a common ℓ^k-th root of unity u so that every one-variable row and value polynomial has valuation increased by at most δ relative to the upper bounds at 1, and every stated monic factor has valuation at most δ. This is Lemma 40.13, with $m = 1$ and initial point 1.

At the row-specific point $y_i(u)$, the sharp polynomial bounds the length with all nonhelper ties by $B_i + \delta$. Dropping those ties by Proposition 42.17 costs at most $2J_0\delta$. Hence the unrestricted normalized module at this point has length at most $B_i + (2J_0 + 1)\delta$.

For each value family use the actual root tuple $y^V_{P_0} = u$ and $y^V_P = 1$ at the other places. Every nonhelper Frobenius evaluates this character at u or the prescribed conjugate-inverse value; the Euler-factor polynomials selected above imply the required valuation bound δ. Successively adding the nonhelper places to the declared modulus and applying the insertion inequality (40.14) gives

$$v_\ell V(J, \xi_{y^V}) \le v_\ell V(\varnothing, 1) + 8c + (8J_0 + 1)\delta.$$

The conductor of the character at this intermediate point can be smaller than the declared auxiliary modulus; this is allowed in the value-family distribution law. No equality between y^V and any of the row-specific points is claimed.

Second selection. For the unrestricted module of row i, take its sharp Fitting polynomial $F_i(Y)$ at the row-specific point $y_i(u)$; this point is a legitimate torsion specialization since $N \ge k + 2M$. Then

$$v_\ell F_i(y_i(u)) \le B_i + (2J_0 + 1)\delta.$$

For each value family take its polynomial on the full declared set J, evaluated at y^V as just bounded. Every coordinate of every one of these tuples satisfies the initial power condition $\ell \mid (y-1)^{\ell^k}$: this holds for u and for $1+t(u-1)$ by binomial expansion in residue characteristic ℓ. All initial valuation bounds, and the bound $|J| \le J_0$, are independent of the original degree of the polynomials and of N.

Apply Lemma 40.13 with the first m powers of one tuple of roots, the common power parameter $a = \ell^k$, and loss δ. Include the monic polynomials $Y^j - 1$ for $1 \le j \le m$. Thus a threshold for N was available before the pencil was chosen, since it depends only on the listed bounds and not the polynomial degree. This is the precise reason it was possible to choose N in the first paragraph. The selection yields one tuple ζ of ℓ^N-th roots such that all its first m powers satisfy:

$$\begin{aligned} \lambda(\text{normalized module of } \xi_{\zeta^j} R_i) &\le B_i + (2J_0 + 2)\delta, \\ v_\ell V(J, \xi_{\zeta^j}) &\le v_\ell V(\varnothing, 1) + 8c + (8J_0 + 2)\delta, \\ \sum_{P \in J} v_\ell(\zeta_P^j - 1) &\le J_0\delta. \end{aligned}$$

For the last inequality use the selected monic factors; finiteness of their valuations ensures no ζ_P^j is 1. The character ξ_ζ constructed from the pencil is anticyclotomic, locally trivial at the

required old places, and all its first m powers have the same auxiliary conductor. The defining finite characters come from $U(1)$ by the anticyclotomic class-field construction of Section 41.4; taking their products and powers preserves this assertion. Integral coefficient rings can be enlarged to contain all the selected roots and coefficients, without changing the normalized upper bounds. Since $(8J_0 + 2)\delta < \eta$, these are precisely the inequalities of Proposition 40.14. They reconstruct the two-stage argument of `exists_anticyclotomic_twist` in `New/Selmer/TwistTheorem.lean`. In particular, the new conductor introduces no uncontrolled term beyond the sum of tame valuations.

To obtain `exists_mFoldTwist_selmerLengthLe` in the source one now chooses the family of rows from the finitely many eigensystems and rank-one pairs. The balance bound of `IsRow.eventually_isBalanced` is $10fn_ic_\ell$, and the auxiliary-character value bound at the old distinguished inertia is c_ℓ; use $\eta = c = c_\ell$. The result is the $(10fn_i+1)c_\ell$ normalized Selmer length and the $9c_\ell$ inverse-ratio valuation bounds of (40.17) and (40.18). These deductions use the actual value-family and balanced-lattice comparison results, not an assumption that twisting always makes a Galois representation generically acyclic.

42.11 Weak attachment: the trace and descent reductions

The global representation used in the congruence argument is not obtained by assuming an equality of local Euler factors at every place. The definition `Attached` in `Lib/GaloisRep/Main/AttachedDefs.lean` asks for a continuous semisimple representation whose characteristic polynomials approximate the specified Hecke polynomials arbitrarily closely on a particular filter of geometric Frobenius data. A set belongs to the filter if every nonempty open set of Galois elements contains a nonempty open subset such that all the Frobenii in that subset at places of prime norm, except for finitely many places, belong to the set. The filter localized near a fixed group element is proper by the existence statement of Chebotarev, so one can take limits there. This is enough to identify traces of two attached representations by continuity. We give the algebraic reconstruction from trace limits, the quadratic descent device, and the even-rank cancellation. The geometric construction of the seed, and the supply of actual automorphic approximations and matching local types, are still separate inputs.

Lemma 42.18 (Reconstruction of a limiting trace)**.** *Let k be an algebraically closed field of characteristic zero and G a group. Suppose $T : G \to k$ satisfies all polynomial identities among finitely many values of traces of n-dimensional representations of G over k. Then T is the trace of a semisimple representation of dimension n over k. If k is a topological field and T is continuous for a topological group G, the constructed representation is continuous.*

Proof. Extend T linearly to $\tau : k[G] \to k$. It is a cyclic trace, since the identity $\operatorname{tr}(AB) = \operatorname{tr}(BA)$ holds for all matrices. Put

$$I = \{a \in k[G] : \tau(ab) = 0 \text{ for all } b \in k[G]\}.$$

Cyclicity shows I is a two-sided ideal, and τ descends to $A = k[G]/I$ by taking $b = 1$. The induced bilinear pairing $\tau(ab)$ on A is nondegenerate. For any list of $n^2 + 1$ elements a_i and b_j of $k[G]$, the determinant of the matrix $\tau(a_ib_j)$ vanishes: it is a polynomial combination of trace values, which vanishes for an actual n-dimensional representation by the rank bound on the space of matrices. If A had dimension greater than n^2, one could select n^2+1 independent a_i and choose b_j with independent pairings, contrary to the determinant identity. Hence $\dim A \le n^2$.

We check the constraints on the possible trace functionals of A. For any $a \in A$, define a monic polynomial p_a of degree n from $\tau(a), \dots, \tau(a^n)$ by Newton identities, and let its

roots be $\lambda_1, \dots, \lambda_n$. For every $j \geq 1$ one has $\tau(a^j) = \sum_s \lambda_s^j$, because this identity (expressed in the coefficients by Newton identities) holds for the traces of all actual matrices. If a is nilpotent, the right side vanishes for all sufficiently large j. Distinct nonzero roots and their multiplicities would give a nonzero solution of a Vandermonde system with invertible matrix, using as many consecutive large values of j as there are distinct nonzero roots. Thus all roots are zero and $\tau(a) = 0$. If a belongs to the Jacobson radical of A, then ab is nilpotent for every b because the radical of the finite-dimensional algebra is nilpotent. The preceding argument gives $\tau(ab) = 0$ for all b and therefore $a = 0$. Consequently A is semisimple.

Over k it is a finite product of full matrix algebras $M_{d_i}(k)$. On each summand cyclicity makes τ a scalar multiple $c_i \operatorname{tr}$ of the usual trace, as is seen by applying it to products of matrix units. For a rank-one idempotent a in that summand, $\tau(a^j) = \tau(a) = c_i$ for all $j \geq 1$. The root power-sum identity just proved, or the recurrence of those moments, then forces the roots of p_a to consist only of 0 and 1. Hence c_i is the multiplicity of 1, a nonnegative integer. Nondegeneracy forces $c_i > 0$, and $\tau(1) = n$ gives $\sum_i c_i d_i = n$. The direct sum of the usual modules of the matrix summands with multiplicities c_i is the desired representation of A, and restriction along $G \to A^\times$ gives the representation of G.

For continuity, choose a basis of A and its dual basis for the trace pairing. The coordinate functions of the image of g in A are linear combinations of finitely many $T(gh)$, because the dual basis lifts to finite group-algebra sums. They are continuous, so the endomorphisms of the described representation vary continuously with g. □

This is the argument of `exists_rep_of_pseudo` and the continuity-of-trace lemmas in `Lib/GaloisRep/Pseudocharacter/FibreSum.lean` and `SemisimpleReps.lean`. In particular, pointwise limits of traces in a fixed complete coefficient field contained in $\overline{\mathbb{Q}}_\ell$, when the limiting function is continuous, satisfy the lemma. The completeness stipulation is needed: arbitrary limits in $\mathbb{C}_\ell$ need not be algebraic over $\mathbb{Q}_\ell$.

For completeness, a continuous semisimple representation of a profinite group into $\mathrm{GL}_n(\overline{\mathbb{Q}}_\ell)$ can be realized in a finite coefficient extension and stabilizes an integral lattice. Its compact image is a union of its intersections with $\mathrm{GL}_n(L)$ over finite subextensions L of $\overline{\mathbb{Q}}_\ell$. There are countably many such L (there are finitely many isomorphism classes in each finite degree, and each has finitely many embeddings). Each intersection is closed in the compact image. By the Baire property one has nonempty interior, and, being a subgroup, is open. It has finite index, so adjoining the coefficients of finitely many coset representatives to L puts the entire image over one finite extension. Over its integer ring, take the O-span of the orbit of a standard lattice. Compactness bounds its entries, so it lies between two homothetic standard lattices and is finitely generated. By construction it is stable. Enlarging the coefficient field afterward by any finite extension preserves the lattice constructions in the appropriate scalar extensions.

We next describe the construction that avoids an uncontrolled patching of Galois representations over varying extensions. The source assumes the CM field is `StablyUnram`, meaning it contains a primitive 5040-th root of unity. In particular it contains i with $i^2 = -1$ and an algebraic integer ω with $\omega + \bar{\omega} = 1$ and $\omega\bar{\omega} = 2$, since it contains the appropriate quadratic subfield $\mathbb{Q}(\sqrt{-7})$ of the seventh cyclotomic field. We use these two elements in the following elementary construction, which is the number-theoretic part of `exists_quadratic_step` in `Lib/GaloisRep/Main/AttachedDeterminant.lean`.

Lemma 42.19 (Quadratic steps with prescribed splitting)**.** *Let E be a CM field containing these i, ω and let n_0 be a positive integer. There is a quadratic character $\xi : G_E \to \{\pm 1\}$ such that, putting*

$$E' = \overline{\mathbb{Q}}^{\ker(\xi\xi^c)}, \qquad U = \ker(\xi\xi^c), \qquad H = \ker(\xi) \cap U,$$

the field E' contains a quadratic imaginary field K' in which every rational prime dividing n_0 splits and is unramified. At a place above a prime dividing n_0, its decomposition group lies in U, its inertia group lies in H, and its decomposition group lies in H if the place of E differs from its conjugate. Also $G_E = HV$ whenever a subgroup V contains inertia at all places of E over a certain rational prime not dividing n_0.

Proof. Set $k = |N_{E/\mathbb{Q}}(4)|$, write $R = 4n_0(2n_0)^k = 2^j b$ with b odd, and let $D_1 = 1 + 2b + 8b^2$. The integer D_1 is prime to R and congruent to 3 modulo 4. By Dirichlet's theorem choose a rational prime $q \equiv -D_1^{-1} \bmod R$, greater than $D_1 n_0$ and the absolute discriminant of E. Then $q \equiv 1 \bmod 4$, so $q = s^2 + t^2$ for integers s, t. The algebraic integer

$$\alpha = (1 + 2b\omega)(s + ti)$$

satisfies $\alpha\bar{\alpha} = D = D_1 q$. In particular $D \equiv -1 \bmod R$. The prime q is unramified in E with $v_P(D) = 1$ for all $P \mid q$. At each such place the valuations of α and its conjugate are nonnegative and sum to one; hence $P \neq P^c$, and in the conjugate pair one valuation is one and the other zero.

Use the Chinese remainder theorem for the pairwise coprime ideals w^k at the places $w \mid n_0$. For each pair w, w^c choose an ordering and choose u integral with $u \equiv 1 \bmod w^k$ and $u \equiv \alpha \bmod (w^c)^k$. At the self-conjugate places choose $u \equiv 1$. If there are no conditions take $u = 1$. Put $a = \alpha u \bar{u}$. It is nonzero and $\bar{a}a = D(u\bar{u})^2$. At a split conjugate pair above n_0, a is congruent to D modulo w^k on the first side and to α^2 on the other side. In the first case multiplication by i^2 makes it congruent to 1; in the second multiplication by α^{-2} does so. More generally all elements used here are local units, since D is prime to n_0. The exponent $k = 4^{[E:\mathbb{Q}]}$ is greater than $2v_w(2)$ at every such place. Therefore the unit congruence to 1 implies a local square by Hensel's lemma applied to $X^2 - a$ after the adjustment. At a self-conjugate place, the element a is still a local unit. The residue characteristic is not 2: because E contains $\mathbb{Q}(\sqrt{-7})$, and 2 splits in that quadratic field, conjugation exchanges the places of E above its two places. Thus $X^2 - a$ gives a split or unramified quadratic local extension at any self-conjugate place of the prescribed set.

Let ξ be the quadratic character of $\sqrt{a}$. Then $\xi\xi^c$ cuts out $E(\sqrt{-D})$ because $i\sqrt{a}\sqrt{\bar{a}}/(u\bar{u})$ has square $-D$. The square class of $-D$ is nontrivial (valuation at every $P \mid q$ is odd). For rational primes dividing n_0, the congruence $D \equiv -1 \bmod R$ shows that $-D$ is a square in the corresponding $\mathbb{Q}_p$ by Hensel's lemma; the exponent of 2 is sufficient since R is divisible by 8 if $2 \mid n_0$. It follows that $K' = \mathbb{Q}(\sqrt{-D})$ has the claimed splitting. Thus the local decomposition groups lie in U, and the local square and unramifiedness assertions above for a give the claimed restrictions to H.

Finally, for a pair P, P^c above q, the valuations of $u\bar{u}$ have the same parity on the two sides, while those of $\alpha, \bar{\alpha}$ have opposite parity. Since the residue characteristic is odd, an odd valuation gives a ramified quadratic extension, and an even valuation gives a split or unramified quadratic extension. The restrictions of (ξ, ξ^c) on inertia at these two places consequently have the two independent nonzero images $(1, -1)$ and $(-1, 1)$ in $\{\pm 1\}^2$, in some order. The kernel of the pair is H; hence every subgroup containing these inertia groups maps onto the quotient G_E/H and gives $G_E = HV$. This proves the lemma. □

The detailed theorem `exists_quadratic_step` also asserts the normalized Weil and base-change conditions `IsStep`. We have reconstructed the splitting and quotient-control part here; those additional automorphic conditions are used as a compatibility input.

Lemma 42.20 (Gluing on a common quotient)**.** *Let G be profinite, H_0, H_T open normal subgroups, and r_0, r_T continuous semisimple representations on these subgroups over an*

algebraically closed field of characteristic zero. Suppose their restrictions to $H_0 \cap H_T$ have equal traces for every sufficiently large T. Suppose a closed normal subgroup K of G is contained in $\bigcap_T \ker r_T$ (for those sufficiently large T), and $G = H_0K$. Then there is a continuous semisimple representation $r : G \to \mathrm{GL}_n$ extending r_0, whose restriction to each such H_T is conjugate to r_T.

Proof. The restrictions to $H_0 \cap H_T$ are semisimple. One elementary form of Clifford's argument is to note that the sum of the translates of an irreducible submodule for the finite-index normal subgroup is a nonzero module for the original group, and hence is the whole irreducible representation. As a sum of simple submodules it is semisimple. Restriction of a semisimple representation then gives semisimplicity. Equality of traces identifies semisimple representations by the argument of Lemma 41.3. In particular r_0 is trivial on $H_0 \cap K$. Therefore define $r(hk) = r_0(h)$ for $h \in H_0$, $k \in K$. This is a well-defined representation of G/K and hence of G, as $H_0/(H_0 \cap K) \simeq G/K$. The isomorphism is a homeomorphism of compact Hausdorff groups, giving continuity. Semisimplicity follows from the image equality with $r_0(H_0)$. For an element of H_T write it as hk with $h \in H_0$, $k \in K$. Since $K \subseteq H_T$ and is killed by r_T, $h \in H_0 \cap H_T$ and the fixed conjugacy on this intersection gives the desired relation on H_T. □

Applied as follows, this proves the group-theoretic descent assertion used in `spherica lCompat_of_odd` of `Lib/GaloisRep/Main/AvatarCompatibility.lean`. For an eigenform of odd rank with finite level set S, choose a sequence of integers n_T divisible by ℓ and all primes under S and eventually divisible by every given rational prime. Apply Lemma 42.19 to obtain H_T. The weak base-change theorem and odd-rank split-place attachment supply the restricted representations r_T on H_T, with approximation of the original Hecke polynomials there, and unramifiedness at the places dividing n_T outside S, ℓ. Equality of their traces on $H_0 \cap H_T$ follows from weak attachment and Chebotarev. In Lemma 42.19 for H_0, the relevant inertia prime lies outside n_0 and hence outside S, ℓ. For all sufficiently large T, it divides n_T, so its inertia groups lie in H_T and are killed by r_T. Put K equal to the normal core of the intersection of these kernels. It contains all the indicated inertia groups, and the quotient-control assertion gives $G_E = H_0K$. Lemma 42.20 gives r.

To verify the weak approximation on G_E from its restrictions, take any nonempty open subset of G_E and a geometric Frobenius of a place of split prime norm there, outside the finite exceptional set (Chebotarev). Choose T so large that this place divides n_T; its decomposition group lies in H_T by Lemma 42.19. The intersection of the open set with H_T is nonempty and open. The restricted weak attachment at this T supplies the required subwindow of Frobenius approximation, which is also valid for r by conjugacy. This gives weak attachment at G_E in the precise filter sense. The local comparison requires the additional assertion of the source that finite type and jumps descend from the finite extension (`IsWeakBaseChange.exists_attached_descent` and its local comparison lemmas). At a split spherical place one takes T for which its decomposition group is in H_T; the killing-polynomial identity is then inherited directly.

The geometric part not supplied by these elementary arguments is the existence of the split-place odd-rank seed with free weights and the approximation by such seeds, together with the normalized weak base change. The source constructs it by PEL Shimura varieties and cohomological representations, with representations extracted from traces and local type information. Principal anchors are `exists_rep_of_cuspidal_type` in `Lib/GaloisRep/ Main/SeedRepresentation.lean`, `sphericalCompat_of_split` in `Main/AvatarCompati bility.lean`, and `isApproximatingMonoid_of_mem_closure` in the weight-congruence development. They are substantial known-theory inputs in the present reconstruction. The

point of the quotient calculation is that it does not require an independent modularity-lifting or eigenvariety patching theorem once those inputs are available.

Proposition 42.21 (Two theta strings remove the parity restriction)**.** *Suppose a Hecke system of rank n admits two odd-rank theta lifts of rank $n+t_0$, whose normalized Hecke polynomials off a finite set are $P_w D_{j,w}$ for $j = 0, 1$. Assume the auxiliary factors are cyclotomic or algebraic character strings, with attached continuous character lists $\lambda_{j,1}, \dots, \lambda_{j,t_0}$, disjoint between the two lists. If the odd-rank systems have continuous semisimple attached representations R_j, then there is a continuous semisimple n-dimensional representation with trace*

$$\operatorname{tr} R_0 - \sum_{i=1}^{t_0} \lambda_{0,i} = \operatorname{tr} R_1 - \sum_{i=1}^{t_0} \lambda_{1,i}.$$

It is weakly attached to P_w. At a spherical place where both theta systems have the required local comparison, if $D_{0,w}, D_{1,w}$ are coprime, their two killing-polynomial identities imply that P_w kills the corresponding Frobenius on this representation.

Proof. The trace equality follows by taking the coefficient of degree $n + t_0 - 1$ in the two factorizations and applying weak attachment, taking limits near each Galois element. Finite-character twists used in normalizing the polynomials are already accounted for in R_j. By equality of traces of semisimple representations, every character in the first list occurs with at least its listed multiplicity in R_0 (use disjointness and compare the multiplicity on the other side of the equality). Thus the virtual trace is the trace of a direct summand of rank n, with equal multiplicities for the analogous direct summand of R_1. These are actual continuous semisimple representations. They satisfy the stated trace equality and, by Newton identities on the powers of Galois elements, their characteristic polynomials factor the actual characteristic polynomials of both R_j by the corresponding character polynomial. At the generic Frobenii, the norm of the error after multiplication by the monic auxiliary polynomial bounds the norm of the original error, by multiplicativity of the non-Archimedean polynomial Gauss norm and the fact the auxiliary polynomial has norm at least one. This proves weak attachment to P_w, including arbitrary approximation precision. Direct summands of unramified or crystalline representations have the corresponding local property; their multiset of Hodge–Tate jumps is obtained by removing the character jumps. For the killing statement, on the common summand both $P_w D_{0,w}$ and $P_w D_{1,w}$ annihilate the operator, and the Bezout identity for the two D gives the result. □

In the source, `sphericalCompat_std` in `Lib/GaloisRep/Main/SphericalAllRanks.lean` uses $t_0 = n + 1$ and two unramified $U(1)$-characters with different weights. Their powers and finite values are arranged so that their auxiliary characters differ at every split place. This gives the disjointness and the Bezout step of the proposition. The specialized auxiliary theta-lift existence `exists_theta_lift_heckePoly` and the exact jump and local character normalizations remain part of the automorphic input. Weak attachment for the rows of the congruence argument needs specifically the tame parahoric label compatibility, not just this spherical construction. The metacyclic calculation in Chapter 41 applies once these label and inertia polynomials are supplied; in particular full matrix span cannot be inferred from spherical attachment alone.

42.12 Hecke-polynomial continuations

The false lifts in the denominator estimate need not themselves be theta lifts, but their local unitarity may force a fixed Hecke root string. We record the exact polynomial arguments.

Lemma 42.22 (An integral normalized multiple)**.** *Let A, A', B, C be polynomials over $\overline{\mathbb{Q}}_\ell$ with B monic and C integral. If*

$$\|AB - C\|_{\mathrm{G}} < 1, \quad \|A'B - C\|_{\mathrm{G}} < 1, \quad |A'(1)|_\ell \geq 1$$

for the polynomial Gauss norm, then there is an integral polynomial Q divisible by A with $Q(1) = 1$.

Proof. The Gauss norm is multiplicative with $\|B\|_{\mathrm{G}} \geq 1$. Consequently $\|A\|_{\mathrm{G}} \leq 1$ and, subtracting the two errors, $\|A - A'\|_{\mathrm{G}} < 1$. Evaluation at 1 implies $|A(1)|_\ell = |A'(1)|_\ell \geq 1$. The first norm bound implies equality with one. Set $Q = A/A(1)$. □

Lemma 42.23 (Eliminating eigenvalue one)**.** *Let T be a continuous n-by-n matrix function of a group element over $\overline{\mathbb{Q}}_\ell$. Suppose $T(g_i) = s^i A$ for $i \geq 0$, with s a unit and $|s^j - 1|_\ell = 1$ for $1 \leq j \leq n$. Some $i \in \{0, \ldots, n\}$ has a neighborhood on which $|\det(1 - T(g))|_\ell \geq 1$.*

Proof. If $|\det(1 - s^i A)|_\ell < 1$, some eigenvalue λ of A has $|1 - s^i\lambda|_\ell < 1$. The same eigenvalue cannot do so for two distinct i, j in the range: otherwise $|s^{i-j} - 1|_\ell < 1$. If the strict inequality held for all $n + 1$ indices there would be at least $n + 1$ distinct eigenvalues. At the selected point the nonzero determinant has locally constant absolute value by continuity. □

These prove the declarations `exists_integral_multiple_eval_one` and `exists_eventually_no_eigenvalue_one` in `New/Selmer/HeckeString.lean`. The scalar inertia of a row allows application of the second lemma. The possible false lifts on which every element has eigenvalue one are handled differently.

Proposition 42.24 (String from a root of exceptional magnitude)**.** *Suppose P is the spherical Hecke polynomial of an irreducible unramified unitary representation of GL_n over a local field of residue cardinality q, in the normalization in which the roots are $q^{(n-1)/2}$ times normalized Satake parameters. If c is a root and $|c|^2 q^t = q^n$ for an integer $t > 0$, then cq^i is a root for every $0 \leq i < t$.*

Proof. The classification of the spherical unitary dual expresses the parameters as centered strings

$$\{uq^{-(a-1)/2}, uq^{-(a-1)/2+1}, \ldots, uq^{(a-1)/2}\}, \qquad |u| = 1,$$

and pairs of such strings shifted by the powers $q^\delta, q^{-\delta}$ for $0 < \delta < 1/2$. This is the spherical form of Tadić's classification (M. Tadić, *Spherical unitary dual of general linear group over non-Archimidean local field*, Annales de l'Institut Fourier 36 (1986), no. 2, 47–55, DOI: 10.5802/aif.1046). The normalized magnitude of c is $q^{-(t-1)/2}$. Its exponent is a half-integer, so cannot occur in strictly shifted strings. In a centered string of length a it occurs only if $a \geq t$ and the parities agree. Starting at exponent $-(t-1)/2$, the string continues through all exponents up to $(t-1)/2$ with the same phase. Thus the next $t - 1$ roots obtained by multiplication by q are present. □

For a unitary cyclic local representation generated by a spherical eigenvector use its corresponding irreducible spherical constituent. Alternatively the elementary double-coset formula `isRoot_string_of_norm` in `Lib/LocalGL/Tadic/ChainLemma.lean` applies directly to its invariant pairing. The rescaling by `chiOneInv` is essential: the root 1 of the normalized Galois representation corresponds to a root of exactly this exceptional magnitude. If a normalized attached representation has eigenvalue 1 at every group element, the local polynomial-killing property thus implies at a split spherical place all roots cq^i for $0 \leq i < t = n - r$, where r is the row rank. Divide the Hecke polynomial by these distinct factors

and rescale the remaining factor. On generic Frobenii its product with the rescaled string is congruent coefficientwise to the characteristic polynomial of the attached representation. The latter is integral by the lattice argument of Section 42.11. If another system has the same characteristic-polynomial reduction and a distinguished row polynomial A' whose value at 1 has norm at least one (as ensured by Lemma 42.23), then the remaining factor A satisfies Lemma 42.22 with the common monic string B. It therefore divides an integral polynomial equal to 1 at 1. This is the algebraic implication `eventually_heckePoly_eq_string_mul` in `New/Selmer/HeckeString.lean`.

This distinguishes two kinds of false lift. If some Galois point has no eigenvalue one, finite polynomial avoidance as in Section 40.10 makes a normalizer determinant nonzero on that system. If every group element has eigenvalue one, the Hecke string is forced and such systems can be tested at new primes. One cannot conclude solely from the string that these systems are actual theta lifts. The specialized patching estimate in `New/Patching/OnePrimeTie.lean` is still needed to transport the first-order relations into integral operators at the original level. Its finite transfer algebra was studied in Section 41.9.

42.13 Rank-one normalization of the projection bound

A finite averaging calculation accounts for the class factor in the rank-one pairing. The test condition in `AuxPair.IsTestForm` of `New/Selmer/FitTwists.lean` involves $\ker\Theta$ intersected with the label-$X-1$ idempotent image and the test Hecke kernel; here Θ is the descent map to forms on the rank-one group. By contrast, for the line Z of the trivial lift, the condition in `DenominatorLe` involves $Z^{\perp}$ in place of $\ker\Theta$. A form satisfying the first condition need not satisfy the second. The averaging bridges the difference.

Lemma 42.25 (Averaging onto the rank-one theta line)**.** *Let C_1 be the finite group of double classes of $U(1)$ at the chosen level and assume these classes have trivial stabilizers, writing $h' = |C_1|$. Let the kernel vectors K_g, $g \in C_1$, give $\Theta(v)(g) = \langle K_g, v\rangle$ and $L = \sum_{g\in C_1} K_g$. Suppose the twisted central action $T(s)$ satisfies*

$$T(s)^{-1}K_g = K_{gs}, \quad T(s)L = L, \quad T(s)^* = T(s)^{-1},$$

and commutes with the label projection e_0 and every test Hecke polynomial. Assume the class representatives s_g can be chosen integral above ℓ with $T(s_g)^{-1}$ preserving the integral lattice at the chosen embedding. If `DenominatorLe` holds for the projection to $Z = \mathbb{C}L$ with rank-one allowance β, then for two integral rank-one test forms v, v' one has

$$|\iota(h'\langle v, P_Z v'\rangle)|_\ell \le e^{\beta}.$$

Proof. The zero-line case is immediate. Put $\widetilde{v}' = \sum_{g\in C_1} T(s_g)^{-1}v'$, integral at the chosen embedding. For $w \in Z^{\perp}$ lying in the image of e_0 and in the kernel of a test polynomial, $A(w) = \sum_g T(s_g)w$ belongs to the same image and kernel. Also

$$\Theta(A(w))(h) = \sum_g \langle T(s_g)^{-1}K_h, w\rangle = \sum_g \langle K_{hs_g}, w\rangle = \langle L, w\rangle = 0.$$

This uses the permutation of the classes by multiplication and their equal weights from trivial stabilizers. Thus $A(w)$ lies in the test intersection containing the kernel of Θ. Since v' is a rank-one test form, $\langle w, \widetilde{v}'\rangle = 0$. This is the stronger orthogonality required by the rank-one denominator bound. Finally $P_Z T(s_g)^{-1} = P_Z$ because $T(s_g)$ is unitary and fixes L. Hence $P_Z\widetilde{v}' = h' P_Z v'$. Applying the rank-one denominator bound to $v, \widetilde{v}'$ gives the assertion. □

The hypotheses on the actions and integral representatives are verified in the source by the scalar Weil identity, the algebraic character norm in the Weyl lattice, and integral representatives at the chosen level. The finite calculation above is `DenominatorLe.rank_one_bound` in `New/Selmer/RankOneDenominator.lean`. It explains why the ordinary class factor multiplies the pairing, rather than entering as an unexplained volume normalization.

42.14 Dependency status of the congruence argument

The odd-class bound of Proposition 40.16 has been supplied by an actual conventional calculation in Theorem 42.14. The uniform detecting-coordinate theorem of Section 42.6 also supplies an input to the even-part triangular construction. We gave the two-stage common-twist proof from the mirror criterion, balanced-lattice estimate and value-distribution formula rather than treating it as a single black box. We also proved the trace-reconstruction and quotient-descent parts of weak attachment, the elementary quadratic steps, the even-rank cancellation once two appropriate lifts are available, the Hecke-string algebra, and the rank-one projection averaging.

These developments do not prove the entire denominator theorem `denominatorLe_rowBudget` by themselves. Its remaining obligations include the geometric seed and exact local-global type compatibility in the attachment, the global-duality construction of the mirrors and the odd reciprocity laws used for the even-part triangle, the precise integral value measures of the ramified pairs, and the arithmetic Ribet-to-Hecke lifting step at the actual patched auxiliary levels. The integral comparison of crystalline lattices in the following chapters remains essential. An explicit uniform bound on the odd modules does not control the even modules, any more than qualitative local-global compatibility controls the torsion lattice losses.

Chapter 43

The remaining global-duality and Hecke interfaces

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 44, 45, 46 and the final audit in Chapter 57.

The even classes of a Tate twist in the denominator argument are not bounded by the odd-class estimate. They have to be removed by the ordered auxiliary-prime construction of Section 41.5. We supply the existence of its odd laws below. The argument uses ordinary Poitou–Tate duality over the real subfield; it does not require a bound for the even class groups. We use classical duality in its finite-module form, thus avoiding any assumption that the coefficient prime is odd. This is an alternative conventional proof of the specialized reciprocity input, not an attribution of the input in its exact form to the older literature.

43.1 The precise uses of classical duality

For a number field F, a finite G_F-module M killed by ℓ^r, and its Cartier dual $M^D = \mathrm{Hom}(M, \mu_{\ell^r})$, a Selmer condition L means a choice of subgroups $L_v \subset H^1(F_v, M)$, unramified at all but finitely many places. At a real place use Tate cohomology in the local duality theorem (in degree one it is ordinary H^1); at a complex place the local H^1 is zero. Write $L_v^\perp \subset H^1(F_v, M^D)$ for the orthogonal complement.

Lemma 43.1 (Duality with prescribed local cosets)**.** *Let S contain the infinite places, all places over ℓ, and all places at which either module or its local conditions are ramified. Given $b = (b_v)_{v\in S}$ in $\prod_{v\in S} H^1(F_v, M^D)$, the following are equivalent:*

(i) There is a class $z \in H^1(G_{F,S}, M^D)$ whose restriction at each v belongs to $b_v + L_v^\perp$.

(ii) $\sum_{v\in S}\langle u_v, b_v\rangle_v = 0$ for all $u \in H^1_L(F, M)$.

Here $G_{F,S}$ denotes the maximal quotient unramified outside S.

Proof. Use the degree-one part of Poitou–Tate: the image of $H^1(G_{F,S}, M^D)$ in the finite product of local degree-one groups is the exact annihilator of the image of $H^1(G_{F,S}, M)$. This is the exactness at the local H^1 term of Theorem I.4.10 of J. S. Milne, *Arithmetic Duality Theorems*, second ed., BookSurge, 2006. Put $X = \prod H^1(F_v, M)$, I equal to the latter image and $L_S = \prod L_v$. The two conditions are precisely $b \in I^\perp + L_S^\perp$ and $b \in (I \cap L_S)^\perp$ in the finite dual group $X^\vee$. These are equivalent by perfect local duality. Outside S, the local

groups have the unramified condition and its dual: the two unramified subgroups are exact annihilators when $v \nmid \ell$ and the modules are unramified. See also Neukirch–Schmidt–Wingberg, *Cohomology of Number Fields*, second ed., Theorem 7.2.15. □

Lemma 43.2 (Greenberg–Wiles count, with real places)**.** *With these conditions the Selmer groups are finite, and*

$$\frac{|H^1_L(F,M)|}{|H^1_{L^\perp}(F,M^D)|} = \frac{|H^0(F,M)|}{|H^0(F,M^D)|}\prod_v \frac{|L_v|}{|H^0(F_v,M)|}.$$

The denominator at a real place is the ordinary group of invariants, not the Tate H^0 group.

Proof. We derive the formula to specify the real-place convention. Write $B = H^1(G_{F,S}, M)$, $B^D = H^1(G_{F,S}, M^D)$, $C = H^2(G_{F,S}, M)$ and X, I, L_S as in the preceding proof. Finiteness and exactness are supplied by Milne I.4.10. Finite annihilator counting gives

$$\frac{|H^1_L|}{|H^1_{L^\perp}|} = \frac{|B|\,|L_S|}{|B^D|\,|I|}.$$

The segment of global duality continuing from X through $(B^D)^\vee$, C and the local degree-two groups to $H^0(F,M^D)^\vee$ yields

$$|B^D|\,|I| = \frac{|C|\,|X|\,|H^0(F,M^D)|}{\prod_{v\in S}|H^2_T(F_v,M)|}.$$

Here H^2_T is ordinary cohomology at finite places, Tate at real places and trivial at complex places. The global Euler characteristic formula (Milne I.5.1) and the local Euler characteristic formula (I.2.8) give the asserted quotient. The contribution from finite places over ℓ before applying the archimedean formula is $|M|^{-[F:\mathbb{Q}]}$. The factor $\prod_{v|\infty}|M|_v$ in the global formula cancels it. At real places $|H^2_T(F_v,M)| = |H^1(F_v,M)|$ by the Herbrand quotient of a finite module for a group of order two; at complex places both groups are trivial. The remaining archimedean divisor is the ordinary $|H^0(F_v,M)|$. This proves the count also at $\ell = 2$. □

All theorem locations in the preceding argument are in standard finite Galois cohomology theory. Milne's I.5.1 uses only H^0, H^1, H^2 even for a real field: we have not assumed higher cohomology vanishes at the prime two.

43.2 Construction of an odd law with a primitive value

Let E be a stably unramified CM field in the sense of the source, $F = E^+$ its real subfield, and $A_r = \mathbb{Z}/\ell^r\mathbb{Z}$. Write $G = G_E \subset G_F$ and choose a global complex conjugation $c_0 \in G_F \setminus G$. Use $g^c = c_0 g c_0$ for $g \in G$. The quadratic character of E/F is denoted by δ and the ℓ-adic cyclotomic character on G_F by ε. Define

$$\psi = \delta\varepsilon : G_F \longrightarrow \mathbb{Z}_\ell^\times, \qquad K_r = G \cap \ker(\varepsilon \bmod \ell^r).$$

The restriction of ψ to G is ε, but its value on every complex conjugation of F is 1. This is the even extension of the cyclotomic character used in the source.

We shall use that E/F is unramified at every finite place and $\sqrt{-1} \in E$. These follow from the stable-unramified construction: in particular, an algebraic integer whose difference with its conjugate is a unit excludes complex conjugation from the inertia group of E/F (see `StablyUnram.exists_isUnit_sub_star` in `Lib/Adelic/Adeles/CoprimeValuation.lean`).

Inertia at each finite place of F consequently lies in G. The image $\psi(G_F)$ equals $\varepsilon(G)$ and is procyclic: for odd ℓ this follows from the structure of $\mathbb{Z}_\ell^\times$, and for $\ell = 2$ the image is contained in $1 + 4\mathbb{Z}_2$. The quotient G_F/K_r is the product of the order-two quotient generated by c_0 and the finite cyclic image of $\varepsilon|_G$ modulo ℓ^r. This follows directly from $G_F = G \rtimes \langle c_0 \rangle$ and conjugation invariance of $\varepsilon|_G$. The quotient is generated by at most two elements regardless of r.

Theorem 43.3 (Existence of primitive reciprocity laws)**.** *Suppose $[F : \mathbb{Q}] \geq 4$, $r \geq 1$ and $e \geq 0$. Let $x_1, \ldots, x_k$ be degree-one places of E, not over ℓ, lying over distinct primes of F, with chosen geometric Frobenius lifts $f_{x_i} \in K_r$. Given any $y_1, \ldots, y_s \in G$, $s \leq k$, there exists a locally constant $Z : G \to A_r$ and a finite set T of finite places of F:*

(a) On K_r, Z is a homomorphism and

$$Z(gug^{-1}) = \varepsilon(g)^{e+1} Z(u), \qquad Z(u^c) = -Z(u) \quad (u \in K_r,\ g \in G).$$

(b) $Z(y_i) = 0$ for every i, and $Z(u)$ is a unit for some $u \in K_r$.

(c) If w_j are degree-one places of E, outside the places over T and with distinct underlying places of F, and $f_j \in K_r$ are their chosen geometric Frobenii, let d be an even cocycle of twist $-e$ with values in $B = \mathcal{O}/\ell^r$ for an arbitrary coefficient ring of the source. Assume d is unramified away from places over ℓ and $w_j, \bar{w}_j$; it vanishes on the kernel of the full cyclotomic character in decomposition groups above ℓ, and $d(f_{x_i}) = 0$ for all old places. If $Z(f_i) = 0$ for every $i \neq j$, then

$$2Z(f_j) \mid b \text{ in } A_r \quad \Longrightarrow \quad b\, d(\tau_j) = 0 \qquad (b \in \mathbb{Z}).$$

Here τ_j is a tame generator in the indicated decomposition group.

Proof. Set $M_r = A_r(\psi^{-e})$. Its Cartier dual is the free cyclic A_r-module $M_r^D = A_r(\varepsilon\psi^e)$, after fixing a primitive ℓ^r-th root of unity. Thus on G the dual action is ε^{e+1}, and on each complex conjugation the dual action is negation.

Let S consist of the places above ℓ, the k old places of F, and the real places. At the old places and at real places take the zero local condition on M_r. At $v \mid \ell$ take the inflated local condition

$$L_{v,r} = \operatorname{im}\left(H^1(\psi(G_{F_v}), M_r) \longrightarrow H^1(F_v, M_r)\right). \tag{43.1}$$

Outside S, use the unramified local condition. We will temporarily relax to the full local H^1 at an auxiliary set P of finite places, whose geometric Frobenii lie in K_r. All modules are trivial locally at P, because Frobenius lies in the common kernel and the modules are unramified there.

Choosing the auxiliary primes. There are finitely many classes in $H^1(G_{F,S}, M_1^D)$ and in $H^1(G_{F,S}, M_r^D)$. Restrictions of these classes to K_r are independent of representatives, since coboundaries vanish there. Choose finitely many primes in P with Frobenii detecting every class of the former group having nonzero restriction to K_r. For each such class choose an element $u \in K_r$ with nonzero value, use its image in the finite Galois quotient cut out jointly by the class and K_r, and apply Chebotarev. The primes may be taken distinct, outside S, of absolute residue degree one and split in the extension cut out by K_r. The set of primes of absolute residue degree greater than one has density zero, so the degree-one requirement causes no obstruction. Choice of the place in the chosen finite Galois extension permits one to prescribe a particular Frobenius lift modulo the finite quotient, rather than only its conjugacy class.

Also arrange that for each $y_i \in K_r$ there is a marked prime in P whose Frobenius and y_i have identical values on every class of $H^1(G_{F,S}, M_r^D)$. Again these values depend only on restrictions to K_r; intersect the finitely many open kernels there and take the normal core in G_F. Chebotarev gives a fresh prime for the resulting congruence. Thus at most k indices of P are marked. There are no conditions on the y_i outside K_r at this step.

The dimension loss from the real places. Use the local conditions at depth one with all conditions at P relaxed; write $L_{P,1}$. The dual Selmer group in Lemma 43.2 is unramified outside S, because its local conditions at P are zero. By detection all its classes vanish on K_r. Inflation from G_F/K_r therefore contains this group. Since a cocycle on that quotient with values in the rank-one $\mathbb{F}_\ell$-module M_1^D is determined by values on at most two generators,

$$\dim_{\mathbb{F}_\ell} H^1_{L^\perp_{P,1}}(F, M_1^D) \le 2.$$

This does not require that the order of the quotient be prime to ℓ. At $v \mid \ell$, the group in (43.1) comes from a procyclic quotient. If the scalar action on M_1 is nontrivial, every cocycle is a coboundary (evaluate at a topological generator and use that its scalar minus one is invertible). For the trivial action a cocycle has at most one free coordinate. Hence

$$\dim L_{v,1} \le \dim H^0(F_v, M_1) \quad (v \mid \ell).$$

At an old place the inertia and Frobenius act trivially, so the zero condition contributes -1 in the logarithm of Greenberg–Wiles. At a real place the even action is trivial and the zero condition contributes -1 even at $\ell = 2$. At P the action is trivial and $q_v \equiv 1 \pmod{\ell^r}$; there are two scalar cocycle coordinates at depth one, unramified and tame, so the full H^1 contributes $2 - 1 = 1$. Elsewhere the contribution is zero by the unramified condition. On adding the coboundary dimension $1 - \dim H^0(F, M_1)$, the space of scalar cocycles at $L_{P,1}$ has dimension at most

$$|P| - k - [F : \mathbb{Q}] + 3 - \dim H^0(F, M_1^D) \le |P| - k - 1. \tag{43.2}$$

The real-place hypothesis is used here, rather than as an unexplained factor in a class-number bound.

A primitive relation at depth r. Let C_r be the additive group of actual M_r-valued cocycles whose local classes obey the conditions $L_{P,r}$. It is finite, since Selmer cohomology and coboundaries are finite. Its ℓ-torsion is identified, by $M_r[\ell] \simeq M_1$, with a subspace of the cocycles counted in (43.2). At places over ℓ membership in (43.1) is equivalent to vanishing on the exact kernel of ψ in the decomposition group: coboundaries vanish there and inflation-restriction applies. At an old place action is trivial at depth r as well, so the zero-class condition is the zero-cocycle condition. The same is true at a real place. At remaining unramified places inertia acts trivially at both depths. Thus all these conditions survive taking ℓ-torsion, and $|C_r[\ell]|\ell^{k+1} \le \ell^{|P|}$.

For each $v \in P$, choose the unramified generator of $H^1(F_v, M_r^D)$ normalized to have value 1 on the chosen geometric Frobenius. The tame quotient paired with it under local duality is also free cyclic of rank one. Denote the resulting scalar evaluation of a cocycle by $p_v(u) \in A_r$ (this includes a possible unit factor depending on the tame generator and primitive root). Apply Lemma 41.11 to $C_r \to A_r^P$, $u \mapsto (p_v(u))_v$, marking the indices for the y_i. Obtain a primitive vector $n = (n_v)$ vanishing at all marked indices and satisfying $\sum n_v p_v(u) = 0$ on C_r.

Take the tuple of local dual classes at P to be unramified with Frobenius values n_v, and take the zero target coset at all other places of S. The annihilation relation is condition (ii) of Lemma 43.1; there is a global class $z \in H^1(G_{F,S\cup P}, M_r^D)$ with these local values and satisfying dual conditions outside P. Its unramified local values at P imply $z \in H^1(G_{F,S}, M_r^D)$. On

K_r its representative is a homomorphism independent of coboundaries. Set Z equal to this restriction on K_r and zero on $G \setminus K_r$. This is locally constant. On K_r the cocycle conjugation rule and the dual action give the asserted equivariance and oddness. A unit coordinate among the n_v gives a unit value, and the marked coordinates force $Z(y_i) = 0$. This construction is the reason a bound on ℓ-torsion, rather than on the size of the higher-depth Selmer group, suffices.

Reciprocity for all future collections. Take T to contain all finite places of S and, if desired, P. First let $B = A_r$. An even cocycle $d : G \to A_r(\varepsilon^{-e})$ extends to a cocycle $\widetilde{d}$ on G_F with values in M_r, by $\widetilde{d}(gc_0) = d(g)$ and $\widetilde{d}(g) = d(g)$ for $g \in G$. Indeed the action of c_0 on M_r is trivial and evenness is precisely compatibility with the semidirect product. At every real place $2\widetilde{d}$ is zero since complex conjugation has order two and acts trivially. At an old place its restriction is zero: the place splits in E, the inertia is killed by the unramified condition, the Frobenius by hypothesis, and the action is trivial modulo ℓ^r.

At $v \mid \ell$, let h belong to the local decomposition group of F with $\psi(h) = 1$ (the full ℓ-adic value). If $h \in G$ then $\varepsilon(h) = 1$ and the local condition on d gives $\widetilde{d}(h) = 0$. Otherwise $h^2 \in G$ belongs to the corresponding decomposition group of E and $\varepsilon(h^2) = 1$. Since $\psi(h) = 1$, the cocycle identity gives $0 = \widetilde{d}(h^2) = 2\widetilde{d}(h)$. Thus $2\widetilde{d}$ vanishes on the exact kernel of ψ locally and its class belongs to (43.1). At a finite place away from both ℓ and the new primes the extension is unramified, since inertia is contained in G.

Apply the sum-of-local-invariants reciprocity law to $2\widetilde{d}, z$. At S the terms vanish from the dual conditions. Outside S both classes are unramified except at the new underlying places of F. Each such place splits in E, so the value at its chosen tame generator is $2d(\tau_j)$ and the unramified value of z on its Frobenius is $Z(f_j)$. Local duality identifies the pairing with

$$\kappa_j \, 2d(\tau_j) Z(f_j) \in A_r \quad (\kappa_j \in A_r^{\times})$$

after identifying the subgroup of order ℓ^r of $\mathbb{Q}/\mathbb{Z}$ with A_r. This unramified–tame pairing follows from local class field theory: the quotient by unramified cocycles and the dual unramified subgroup are perfect duals, each cyclic of order ℓ^r, with tame-generator and Frobenius evaluations their respective coordinates. The distinct-real-prime hypothesis is essential: each local place of F occurs once. If all but the indicated $Z(f_j)$ vanish, reciprocity and the unit factor give $2Z(f_j)d(\tau_j) = 0$, whence the assertion for b.

Finally compose a cocycle with any additive map $B \to A_r$. Such maps are A_r-linear because the action comes from A_r, and preserve all the required evenness and vanishing conditions. They separate points of B: extend a nonzero homomorphism from the cyclic subgroup generated by a given nonzero point to $\mathbb{Q}/\mathbb{Z}$ by divisibility of $\mathbb{Q}/\mathbb{Z}$; the image is killed by ℓ^r and is consequently in A_r. Applying the scalar argument to all maps proves the assertion for B. □

The exact compact declaration supplied is `exists_isLaw` in `New/Selmer/ReciprocityLaw.lean`; its structure `IsLaw` is recovered by extending Z by zero away from K_r as in the proof. The proof also supplies the additive separation in `eq_zero_of_mem_evenClasses`. The finite-level law is not asserted to be a character on the full Galois group; only its restriction to K_r is additive. No extra torsion-free or integrality hypothesis on the higher-depth dual Selmer group was used. Coupling Theorem 43.3 with the primitive relation, cyclotomic twist separation and the detecting sets of Section 42.6 discharges the law-existence part of Proposition 41.10. In conjunction with the ordered construction of Section 41.5, this gives the triangular auxiliary primes of `exists_isPrimeTriangle` in `New/Selmer/TrianglePrimes.lean`. The factor two is retained; it must not be discarded at $\ell = 2$.

43.3 From Ribet polynomials to integral operators

We next close the multiple-prime coefficient-extraction step, keeping distinct the local representation-theoretic assertions needed at a single prime. This step does not require passage to an inverse limit of patched modules. The argument uses one cyclic module killed by the auxiliary depth and an induction on squarefree monomials.

First note that the Hecke algebra used here is defined as the algebra of *all* functions of the eigensystems whose diagonal action preserves the integral lattice, not merely the abstract algebra generated by finitely many spherical Hecke operators. Thus its closedness follows directly from the definition. At the indicated finite level let Y be the finite system set and $\mathcal{O} = \overline{\mathbb{Z}}_\ell$. Write $D(a)$ for the diagonal operator of a function $a : Y \to \overline{\mathbb{Q}}_\ell$, using the prescribed comparison ι_ℓ as in Section 40.11. Preservation of the forms integral above ℓ means that for every such form, every integral representative and every Weyl coordinate, the ℓ-adic norm of the coordinate of $D(a)$ applied to it is at most one. The coordinate is a finite $\overline{\mathbb{Q}}_\ell$-linear combination of the $a(x)$. Each inequality is a closed condition, and so is their intersection. This gives `isClosed_heckeAlgebra` of `Lib/LocalGL/TaylorWiles/HeckeAlgebra.lean` without requiring the algebraic closure to be complete.

Suppose now that all eigensystems have the stipulated weak Galois attachment, with normalized representations U_x. At split spherical places outside the level, the characteristic-polynomial coefficient functions preserve the integral forms: their operators are finite integral sums of right translations supported away from ℓ. This follows from the double-coset formula; such translations preserve the set of representatives integral above ℓ, hence preserve coordinate integrality. The monic characteristic polynomial of $U_x(g)$ for an arbitrary $g \in G_E$ has coefficient functions which are simultaneous limits of these functions, by weak attachment and the finite set of systems. The scalar normalizing character, taking values in $\mathcal{O}^\times$, only multiplies the degree-k coefficient by its corresponding integral power. The closedness just proved gives

$$x \longmapsto [T^i] \det(T - U_x(g)) \in H.$$

In particular traces, and then $x \mapsto \mathrm{Tr}(P(U_x(h))U_x(g))$ for every $P \in \mathcal{O}[T]$, lie in H: expand P and use the trace at the individual elements $h^i g$. These are `BaseLevel.charpoly_mem_heckeAlgebra` and `BaseLevel.traceElem` of `New/Patching/OnePrimeTie.lean`.

We shall use finite auxiliary residue groups G_j, with product $G_{\mathrm{aux}} = \prod_{j \in J} G_j$, acting by integral operators ρ. The old-level forms are exactly the vectors fixed by this action. The finite Higman averaging assertion with factor c also has a direct geometric proof in this situation:

Lemma 43.4 (Freeness up to stabilizer orders)**.** *Let G' be a subgroup of G_{aux}. Suppose the orders of its stabilizers on the adelic class set of the deep level divide c. There is an integral operator Ψ such that*

$$\sum_{g \in G'} \rho(g^{-1}) \Psi \rho(g) = c\, 1.$$

Proof. Choose one representative in each G'-orbit on the deep class set. On the chosen class, multiply its coefficient vector by $c/|\operatorname{Stab}|$; on all other classes set the multiplier zero. These are integers. This operator is well-defined on equivariant coefficient functions, even if stabilizers also act on the coefficient space, because it multiplies by a number depending solely on a double class. It preserves all coordinate integrality. The conjugates by the translations are just multiplication operators by the translated weights; the coefficient actions in the conjugation cancel. At any class the sum of these weights is $|\operatorname{Stab}| \cdot c/|\operatorname{Stab}| = c$. The assertion follows. The relevant stabilizers really do divide c: lifting a stabilizer of a finite-residue orbit to the old class produces a subgroup of the old rational automorphism group,

surjecting onto the finite stabilizer, and the latter order divides the common multiple of the old automorphism-group orders. This is the source construction of `exists_higman_operator`, in `Lib/LocalGL/TaylorWiles/AuxPlaceOperators.lean`. It works likewise for a subgroup of a single local residue group. □

The weights in this proof may not be replaced by a division by the whole residue group order. They commute with the required translations away from the auxiliary places, because the class set and orbit representatives are formed after imposing those level conditions. All operators below are considered on the space with its existing level and column conditions.

43.4 The local one-prime hypothesis and its algebraic consequence

Take the local integral operators at j to commute with the diagonal Hecke operators and with local operators at different primes. For $g \in G_E$ put, as diagonal scalar functions,

$$t_j(g) = \operatorname{Tr}((U_x(\tau_j) - 1)U_x(g)), \qquad r_j(g) = \operatorname{Tr}((U_x(\phi_j) - 1)e_j(U_x(\phi_j))U_x(g)).$$

For this subsection we distinguish the algebra of the construction from the local representation comparison: we require that, whenever the old-level systems with nonzero cutoff T satisfy the string condition of `BaseLevel.exists_tie`, the following operators can be constructed after any ring representation σ killing the integer $m = \ell^\nu$:

$$c\,\sigma(F_j)(\sigma(\rho_j(a)) - 1) = \sigma(N_j Z_{j,a}), \qquad TNZ_{j,a} = 0, \tag{43.3}$$

$$\sigma(N_j)\sigma(t_j(g)) = 0, \qquad \kappa\,\sigma(F_j)\sigma(t_j(g)) = \kappa\,\sigma(N_j)\sigma(r_j(g)). \tag{43.4}$$

Here $N_j = \sum_{a \in G_j} \rho_j(a)$, $N = \prod_j N_j$, and $\kappa = 2n!$ for column rank n. The relation $TNZ_{j,a} = 0$ is an identity before reduction, where T is a diagonal scalar function and the product is evaluated as operators. Operators F_j are in the local operator ring, as are N_j; the $Z_{j,a}$ are integral endomorphisms, not required to commute with the other primes. These are exactly the conclusions of `BaseLevel.exists_tie`; (43.3) is its invariance-defect estimate. Merely having the two trace relations would be insufficient for the next proposition.

Proposition 43.5 (Squarefree coefficient extraction)**.** *Keep the notation of Section 40.11: the determinant variable is a scalar function $\beta \in H$ vanishing on the distinguished systems $\mathfrak{X}$, $p \in \mathcal{O}[T_0]$ is the cutoff polynomial, and $x_0 \in \mathfrak{X}$. Assume a Ribet polynomial $f \in H[X_j : j \in J]$ satisfies $\beta f \in \mathfrak{J}$, where $\mathfrak{J}$ is the ideal generated by*

$$m \operatorname{Tr}(U_x(g)), \qquad t_j(g) - X_j r_j(g).$$

Writing $o = \mathbf{1}$ and $f_d = [X^d]f$, all f_d with $d < o$ coordinatewise vanish on $\mathfrak{X}$. Put $k = |J| + 1$. Suppose the divisibility

$$\kappa(c^2 p(0))^k \mid m$$

holds in $\mathcal{O}$, with nonzero divisor. Suppose the string condition holds for old systems with $p(\beta_x) \neq 0$ which are congruent to x_0 in the sense of the source, namely every integral Hecke function vanishing at that system takes a nonunit value at x_0. If the one-prime construction (43.3)–(43.4) holds with this string condition, there is a function $s \in H$ with $v_\ell(s(x_0)) \leq v_\ell(f_o(x_0))$ such that:

whenever l, y are old-level forms, $d\,l$ is integral for some nonzero integer d, y is supported on $\mathfrak{X}$, and l, y have the same components where $p(\beta_x) \neq 0$, then $dc\,D(s)y$ is integral.

Proof. For each system x not congruent to x_0, select $a_x \in H$ vanishing there with $|a_x(x_0)|_\ell \geq 1$. Put $A = \prod_x a_x$, equal to one if this index set is empty. Define

$$T = A\,p(\beta), \quad P = c^2 T, \quad s = A^k f_o.$$

Then $|A(x_0)|_\ell \geq 1$ proves the valuation assertion. Where T is nonzero on an old system the string hypothesis holds. Also

$$P^k = (c^2 p(0))^k A^k + v\beta \tag{43.5}$$

for some $v \in H$, by polynomial division of $p(T_0) - p(0)$ by T_0.

Let $\mathcal{R}$ be the integral operator ring on the deep level and take the quotient cyclic $\mathcal{R}$-module

$$B = \mathcal{R}l \,/\, m\mathcal{R}l$$

inside the coefficient-space construction, with action σ by left multiplication. This avoids any claim that reduction of the *vector* l is an integral lattice element; the cyclic module is well-defined whether or not l is integral. Denote by z the class of Ψl, using Lemma 43.4 for the full auxiliary product. Since l is old-level, the Higman identity gives $N\Psi l = c\,l$. Thus for every Hecke function b divisible by T,

$$\sigma(b)\sigma(N)z = 0 \quad \Longleftrightarrow \quad c\,D(b)y \in m\mathcal{R}l. \tag{43.6}$$

Indeed $b_x = 0$ where $p(\beta_x) = 0$; hence $D(b)l = D(b)y$. We now suppress σ throughout computations in $\mathrm{End}_{\mathcal{O}}(B)$; $m = 0$ there.

For $d \in \mathbb{N}^J$, define the complementary operator

$$\Theta_d = \prod_{j \in J} \begin{cases} F_j, & d_j = 0, \\ N_j, & d_j = 1, \\ 0, & d_j \geq 2. \end{cases}$$

Factors commute because they are at distinct places, and $\Theta_o = N$. If $F_j t = N_j r$ and $N_j t = 0$, then

$$\Theta_d t = \Theta_{d+e_j} r. \tag{43.7}$$

Here e_j is the coordinate vector. To check the identity, when $d_j = 0$ use the first hypothesis, when $d_j = 1$ both sides are zero by the second hypothesis, and for larger d_j both sides are zero by definition. No commutation of t with the other factors is required; the factor of index j is placed rightmost.

We record the invariance induction that is usually missing from an informal sketch of this patching step. If $z' \in B$ satisfies $NTz' = 0$, then

$$P^a \Theta_d z' = 0 \qquad \text{for } a + \sum_j d_j \geq k. \tag{43.8}$$

For $a = 0$ the sum of the d_j is at least $|J| + 1$, so some $d_j \geq 2$. For the induction from a to $a+1$, set $W = P^a \Theta_d c$. For every g in a single group G_j, (43.3), the identity $N_j(\rho_j(g) - 1) = 0$, and (43.7) imply

$$W(\rho_j(g) - 1) = P^a \Theta_{d+e_j} Z_{j,g}.$$

The right side is identically zero as an operator: $NTZ_{j,g} = 0$ and $a + \sum_i (d_i + (e_j)_i) \geq k$, so apply the induction hypothesis to $Z_{j,g}$ of any vector. Thus W is right invariant under the full product. Multiplying the Higman identity on the left by W gives $Wc = W\Psi N$. Evaluate at Tz' to get $WcTz' = 0$, that is $P^{a+1}\Theta_d z' = 0$ because diagonal Hecke operators commute with

the local factors. This proves (43.8). The induction records why two powers of c, and one more than the number of primes, enter the divisibility hypothesis.

Define the additive coefficient evaluation

$$\mathcal{L}(f) = \sum_{d \le o} \Theta_d f_d.$$

Diagonal coefficients commute with all Θ_d. The trace relation (43.4) gives, by (43.7), $\kappa \Theta_d t_j(g) = \kappa \Theta_{d+e_j} r_j(g)$ for every d. The map $\kappa\mathcal{L}$ consequently kills every monomial multiple of a trace tie. It kills every polynomial multiple of the first kind of generator $m \operatorname{Tr}(U_x(g))$, because $m = 0$ in the module. By additivity, it kills the entire ideal $\mathfrak{J}$. As $\beta f \in \mathfrak{J}$, equation (43.5) gives

$$0 = \kappa \mathcal{L}(v \beta f) = \kappa (P^k - (c^2 p(0))^k A^k) \mathcal{L}(f).$$

Choose $w \in \mathcal{O}$ with $m = w\kappa (c^2 p(0))^k$. Multiplication by w then gives $w\kappa P^k \mathcal{L}(f) = 0$ in $\operatorname{End}(B)$.

For each proper divisor $d < o$, the coefficient f_d kills y. Equation (43.6) applied to $T f_d$ gives $N T f_d z = 0$. Equation (43.8) then gives $P^k \Theta_d f_d z = 0$. Hence only the top coefficient survives in the preceding identity at z:

$$w \kappa P^k N f_o z = 0.$$

Apply (43.6) once more to $b = w\kappa P^k f_o$ (which is divisible by T). There is an integral operator $V \in \mathcal{R}$ with

$$mVl = c\, D(w \kappa P^k f_o) y.$$

But $\beta y = 0$, because y is supported on $\mathfrak{X}$. Using (43.5) on y and the defining choice of w, the right side becomes $mc\, D(A^k f_o) y$. The characteristic-zero vector space has no m-torsion, so $Vl = cD(s)y$. Multiplying by the original denominator d proves integrality, as V preserves the lattice. □

This proves `BaseLevel.exists_diagOp_mem_ellIntegralForms`, conditional only on the exact one-prime assertion `BaseLevel.exists_tie`, the stipulated representation comparison, and the Ribet-element hypothesis that is part of its statement. In particular it is not necessary to assert existence of coherent integral projectors at all greater auxiliary levels. If a factor m_0 is present in the valuation allowance from the Ribet element, the inequality at x_0 is unchanged after multiplication by m_0.

43.5 The realization of a local first-order tie

We give the argument from local type comparison to the one-prime operator, including the normalization of the Hecke polynomial and integrality of the operator. Its remaining two representation-theoretic inputs are stated precisely below; neither can be replaced by a mere assertion of semisimple spherical compatibility.

Let V be a smooth unitary local representation of $\mathrm{GL}_n(F_w)$, H_w the inverse image of the stabilizer of the first coordinate line under reduction, and I_1 the pro-p Iwahori subgroup with upper-unitriangular reduction. Write q for the residue cardinality and $t = \operatorname{diag}(\varpi, 1, \dots, 1)$ with ϖ a uniformizer. On I_1-invariants the unnormalized double-coset operator U of t is $q^{n-1} P_{I_1} \rho(t)$, where P_{I_1} is Haar averaging of mass one. Indeed the double coset $I_1 t I_1$ has q^{n-1} right cosets; the count amounts to its upper first-row entries modulo ϖ. The following preservation assertion requires no classification theorem.

Lemma 43.6 (Preservation of the line character)**.** *The operator U preserves the subspace of vectors y on which H_w acts by a character of the reduction of the top left entry.*

Proof. Write the character as χ and let S be its kernel in H_w. It is a compact normal subgroup containing I_1. For every $a \in H_w$ the entry $(a^{-1})_{00}$ is a unit. Multiplying a on the left by the upper unitriangular matrix u whose first row is the first row of a^{-1} divided by $(a^{-1})_{00}$ makes the first row of ua zero off the diagonal, and $u \in I_1$. Now $t^{-1}uat$ belongs to H_w: the cleared first row removes any possible denominator, and the lower first column is multiplied by ϖ. Its top left entry has the same reduction as that of a. Consequently

$$P_{I_1}\rho(a)\rho(t)y = P_{I_1}\rho(ua)\rho(t)y = \chi(a)P_{I_1}\rho(t)y.$$

For $h \in H_w$ average this equality over $a = hs$, $s \in S$. The left side is $P_{I_1}\rho(h)P_S\rho(t)y = \rho(h)P_S\rho(t)y$, since S is normal in H_w and contains I_1. Taking $h = 1$ first proves $P_S\rho(t)y = P_{I_1}\rho(t)y$. The general h now gives the required line-character property. The constant factor q^{n-1} in U does not affect it. □

This is the content of `exists_clear_first_row` and `heckeCoset_mem_parahoricEigenspace` in `Lib/LocalGL/Iwahori/HeckeOnLineStabiliser.lean`. It also proves the integrality assertion of `exists_localOps_aeval_hecke` in its use below: the double-coset operator is an integral sum of translations, the character space is contained in the I_1-fixed space and hence in the principal-level space, and all operators at this place commute with those at other places. In particular an integral polynomial in U applied after a finite weighted averaging of the line stabilizer gives a local integral operator on the deep level.

To specify the exact local comparison, let R be the attached characteristic-zero Galois representation. At a local pair of commuting Frobenius and tame-inertia matrices F_0, T_0, a line-character vector has torus label $(\chi, 1, \dots, 1)$. At a chosen generator u of the residue units put $a = \chi(u) \neq 0$. For every nonzero vector in that character space the local results of the source supply an exponent functional λ nonzero on it, with eigencharacter θ_i of the torus, such that

$$\begin{aligned}
&(T_0 - 1)(T_0 - a) = 0, \qquad [F_0, T_0] = 0, \\
&\prod_{i=0}^{n-1} (F_0 T_0^b - \theta_i(\varpi) a_i^b) = 0 \quad (b \geq 0), \qquad\qquad a_0 = a, \quad a_i = 1 \ (i > 0), \\
&\lambda(Uy) = \theta_0(\varpi)\lambda(y).
\end{aligned}$$

The convention is to transport coefficients by ι_ℓ when passing from complex to ℓ-adic expressions. These are the exact polynomial-killing and exponent statements in `galRep_of_parahoric_type` (before specialization to a line), in `Lib/GaloisRep/LocalGlobal/TypeFrobenius.lean`, and the exponent normalization used in `hecke_eq_frob_on_parahoricEigenspace`. This is one of the remaining local-global comparison obligations, not a consequence of the transfer calculation.

Lemma 43.7 (Trace comparison on a line-character vector)**.** *Assume the preceding local statements and $a^{n!} \neq 1$. For any matrix M and any polynomial G_0, on the corresponding character space in a fixed system one has*

$$\operatorname{Tr}((T_0 - 1)M)\, G_0(U)y = \operatorname{Tr}(G_0(F_0)(T_0 - 1)M)\, y.$$

Proof. The subspace $W = \operatorname{im}(T_0 - 1)$ is the a-eigenspace of T_0, since $a \neq 1$ and the indicated quadratic polynomial kills T_0. It is invariant under F_0. For any exponent functional supplied

above, restricting the second polynomial identity to W yields

$$(F_0 - \theta_0(\varpi))\, Q(a^b F_0)|_W = 0, \qquad Q(X) = \prod_{i>0}(X - \theta_i(\varpi)).$$

The n values a^b for $b = 0, \dots, n-1$ are distinct: if two were equal, $a^j = 1$ for some $0 < j < n$ and then $a^{n!} = 1$. The displayed relation is polynomial of degree at most $n-1$ in a^b. Each of its matrix coefficients is zero identically by the Vandermonde determinant. In particular its constant coefficient gives $(F_0 - \theta_0(\varpi))|_W = 0$, because $Q(0) \neq 0$.

To prove the lemma first consider $G_0(X) = X$. If the difference between the two sides on y were nonzero, it would still belong to the same system and line-character space by Lemma 43.6. Choose an exponent functional nonzero on this difference. The preceding argument for this exponent shows $\mathrm{Tr}(F_0(T_0 - 1)M) = \theta_0(\varpi)\,\mathrm{Tr}((T_0 - 1)M)$, and application of that functional to the difference gives zero, a contradiction. To pass from X to X^2 apply the degree-one identity to Uy and to $F_0 M$; these are the correct spaces and matrices because F_0 commutes with T_0. Induction gives monomials, and linearity gives all polynomials. □

This reconstructs the passage from the exact polynomial-type comparison to `hecke_eq_frob_on_parahoricEigenspace`, including the possible case that the spherical polynomials or the θ_i have repeated roots. We used only distinctness of the powers of the single inertial character. On a determinant-character vector the parahoric comparison has only one block and gives the still simpler assertion that its tame-inertia matrix is the scalar given by that character on the residue-unit generator. The assumption must be the annihilating polynomial $X - a$ in this case, not merely equality of characteristic polynomials.

There is a second local input. For an old nonzero spherical eigenvector x in a unitary representation, when its rescaled Hecke polynomial has a string $1, q, \dots, q^{s-1}$ and remaining factor dividing $e(X)$, the unnormalized local U after the same rescaling has the orthogonality property

$$\langle x,\ ((U_{\mathrm{norm}} - q^{s-1})e(U_{\mathrm{norm}}))\, y\rangle = 0 \quad \text{for every } I_1\text{-fixed } y. \tag{43.9}$$

This is the second assertion of `chain_lemma` in `Lib/LocalGL/Tadic/ChainLemma.lean`. It is strictly stronger than saying that one root forces a string of roots, the latter already proved from the unitary-dual classification in Proposition 42.24. The orthogonality of an entire image at pro-p Iwahori level, including repeated Satake roots, is a separate assertion, proved below in Lemma 43.9.

Here is the operator construction with these exact local inputs. Write $G_j = \mathrm{GL}_n(k_w)$, H_j its line stabilizer, and $\omega : H_j \to \mathbb{Z}/m$ the logarithm of the top left residual entry. For a function f on the logarithmic group let $W_H(f) = \sum_{h \in H_j} f(\omega(h))\rho(h)$. Its image is fixed by the kernel of ω, by reindexing the sum, in particular by I_1. Put

$$p(X) = (X - q^{s-1})e_j(X), \qquad B(f) = p(U_{\mathrm{norm}})W_H(f),$$

with the same unit rescaling as for the normalized attached representation. This is integral by the preceding operator construction. It has the translation law of Proposition 41.16; $B(1)$ is preserved by H_j because of Lemma 43.6. Let $L = TN$ with T the cutoff operator. Then $L\rho(g) = L$. Moreover $LB(1) = 0$: the full auxiliary norm N puts its image on the old level. For a system with $T \neq 0$, every spherical vector at the old level is orthogonal to the image of $p(U_{\mathrm{norm}})$ by (43.9). The invariant pairing then gives zero on that old system, since N is self-adjoint and fixes old vectors up to the group order. This argument takes place within each system, which is preserved by all the local operators.

We verify the two weighted-sum identities of Proposition 41.16. For an arbitrary finite group equipped with a logarithmic character $c : G' \to \mathbb{Z}/m$, the operator $W_{G'}(f) = \sum_g f(c(g))\rho(g)$

is a linear combination of the projections onto the character spaces of $G'/\ker c$. Indeed expand f in the character basis of the cyclic additive group and use finite Fourier orthogonality. If f is replaced by $f(\cdot - k) - if$, its coefficient on the character $\eta \circ c$ is multiplied by $\eta(k) - i$: this follows by substituting $b - k$ for b in the finite sum of weights. Thus an equality verified on all character spaces satisfying $\eta(k) \neq i$ gives the weighted operator identity, with the indicated difference of weights; it is unnecessary to divide an integral weighted operator by the size of the group. The Fourier proof is performed in characteristic zero, and the resulting equality is between integral operators before reduction. This is `BaseLevel.weightedSum_ext`.

Apply this observation to H_j, with $k = n!$ and $i = 1$. If a character of the line has $\chi(u)^{n!} \neq 1$, Lemma 43.7 gives

$$W_H(f(\cdot - k) - f)\, t'_j(g) = B(f(\cdot - k) - f)\, t_j(g),$$

where $t'_j(g) = \mathrm{Tr}(p(F_{\mathrm{norm}})(T_0 - 1)U_x(g))$ and $t_j(g)$ is the trace used earlier. This is the first weighted-sum identity of Proposition 41.16. For transfer $c = \mathrm{trf}(\omega)$ to G_j, its determinant factor is given by Lemma 41.15. On the subspace where G_j acts by $\eta \circ c$, the type has one block and the local comparison makes T_0 the scalar $\eta(c(\gamma))$, where γ is a finite residue matrix whose determinant is the chosen residue generator. Thus the operator $\rho(\gamma) - 1$ gives exactly $T_0 - 1$ on those vectors. Consequently

$$W_{G_j}(f)\, t'_j(g) = (\rho(\gamma) - 1)\, W_{G_j}(f)\ \mathrm{Tr}(p(F_{\mathrm{norm}})U_x(g)).$$

Taking the constant weights also shows $N_j t_j(g) = 0$, since inertia is trivial on the fully spherical type. The scalar transfer normalization is $kc(\gamma) = k$ in $\mathbb{Z}/m$, exactly Lemma 41.15. All hypotheses of Proposition 41.16 now hold: use the subgroup Higman operator with factor c of Lemma 43.4, and regularity of multiplication by a nonzero integer on the characteristic-zero operator ring. The resulting F_j is an integral local operator, with invariance defects killed by L, and satisfies the trace tie with polynomial p in place of $(X - 1)e_j$. Finally $q \equiv 1 \bmod m$, so $p(X) \equiv (X - 1)e_j(X) \bmod m$. This replacement is performed *after* reduction, not in the Hilbert-space orthogonality argument. We obtain precisely (43.3) and (43.4).

This supplies the proof of the entire passage in `BaseLevel.exists_tie` from the specified local type and chain statements to the integral operator, with no prime-level freeness assumption beyond the verified common stabilizer factor. The dependent local-global assertion remains to be established in the stated generality, while the chain assertion is an elementary calculation with lattices.

43.6 The chain orthogonality, including multiplicities

We reconstruct the portion of `chain_lemma` that cannot be supplied by the unitary-dual classification. Keep the notation of that source statement: let $N = n + 1$, q be the residue-field size, I_1 the pro-p Iwahori, K the maximal compact subgroup, P_I the orthogonal average on I_1, and let $x \neq 0$ be a spherical vector such that the jth spherical lattice-sum operator satisfies $T_j x = t_j x$, $t_0 = 1$. The monic spherical polynomial is

$$P(X) = \sum_{j=0}^{N} a_j X^{N-j}, \qquad a_j = (-1)^j q^{j(j-1)/2} t_j.$$

Use $\beta = \mathrm{diag}(\varpi, 1, \dots, 1)$ and $U = q^n P_I \rho(\beta) P_I$, with adjoint $U' = q^n P_I \rho(\beta^{-1}) P_I$. The double-coset operator on I_1-fixed vectors is exactly U: its right cosets are represented by upper unipotents with arbitrary residue entries in the first row after the diagonal, q^n choices (`heckeCoset_mulSingle_zero_eq`).

Define $x_j = T_j^1 x$ to be the sum over integral lattices $L \subset O^N$ containing ϖO^N and the first basis vector, of residue codimension j, of $\rho(g_L)x$ where $g_L O^N = L$. The choice of g_L does not affect the vector because x is K-fixed. These are the operators `partialHecke` of `Lib/LocalGL/Iwahori/HeckeCounts.lean`. In particular $x_0 = x$, $x_N = 0$, and they are I_1-fixed since that group stabilizes the first residual line. Let β' be $\mathrm{diag}(1, \varpi^{-1}, \dots, \varpi^{-1})$.

Lemma 43.8 (The two partial-lattice recurrences)**.** *One has*

$$\begin{aligned} Ux_j &= q^j(t_{j+1}x - x_{j+1}) && (0 \le j \le n),\\ U'x_j &= q^{-j}t_j U'x - q^{n-j}x_{j-1} && (1 \le j \le n),\\ U'x &= q^n t_N^{-1} x_n. \end{aligned}$$

Proof. For the first count write Ux_j as a sum over choices of an upper unipotent first row u off the diagonal and a lattice from T_j^1. The image of such a lattice under $g = u\beta$ has codimension $j+1$. Before applying u it lies in the coordinate hyperplane of vectors with first residue coordinate zero, and after applying u it lies in the hyperplane H_u given by the equation $z_0 = \sum_{i>0} u_i z_i$. Conversely, for every residue subspace of codimension $j+1$ contained in such a hyperplane, applying $(u\beta)^{-1}$ yields exactly one partial lattice of codimension j containing the first basis vector. This is because the hyperplane condition gives integrality and the containment of ϖO^N gives the first vector after dividing by the first diagonal entry. Thus no lattice containing the first residual line contributes. For every subspace of codimension $j+1$ not containing that line, its projection to the other coordinates is injective and has dimension $N-j-1$. The equation determining u is a prescribed linear functional on that subspace of k^n; it has $q^{n-(N-j-1)} = q^j$ extensions. This gives the first identity, counting the terms independently of any Hecke-polynomial roots.

For the adjoint count use the dual representation $g \mapsto \rho((g^{\mathsf T})^{-1})$. In the lattice sum the map $L \mapsto \varpi L^\vee$ gives complementary codimensions; in the partial sum use the pairing on the quotient by the first coordinate, keeping the first coordinate at unit scale. Equivalently $g \mapsto (g^{\mathsf T})^{-1}O^N$ followed by division by the lattice $\beta' O^N$ is a bijection between the partial lattices of codimension j and $n-j$. To check it, note that the partial lattices contain e_0 and hence are exactly $Oe_0 \oplus L'$ for lattices L' in the other coordinates, and apply ordinary lattice duality there. We obtain $T_j^{1,\vee}x = \rho(\beta')x_{n-j}$ and $T_j^{0,\vee}x = \rho(\varpi^{-1}1_N)t_{N-j}x$ for the full sum. Apply the same hyperplane count for $j = n-i$, $1 \le i \le n$, in the dual representation followed by the functional $P_I\rho(\beta'^{-1})$. The functional is invariant under the needed inverse unipotents: conjugating $(u^{\mathsf T})^{-1}$ by β'^{-1} changes the off-diagonal entries below the first column by the factor ϖ and gives an element of I_1. Thus the count gives

$$U'x_i = q^{n-i}(t_i P_I \rho(\beta^{-1})x - x_{i-1}),$$

as required. At the last codimension the only partial lattice is $Oe_0 \oplus \varpi O^n$, so $x_n = \rho(\varpi 1_N)\rho(\beta^{-1})x$. The full last spherical operator is the central translation by $\varpi 1_N$, with scalar $t_N \neq 0$; also x_n is I_1-fixed. This proves the last identity. □

Lemma 43.9 (Removal of the nonterminal factors of a string)**.** *Suppose* $P(X) = (\prod_{i=0}^{k}(X - cq^i))S(X)$, $c \neq 0$. *If* $S \mid R$, *then*

$$\langle x, (U - cq^k)R(U)y\rangle = 0 \quad \textit{for every } I_1\textit{-fixed } y.$$

This holds without any assumption on multiplicities of other roots of P, including roots shared with the string.

Proof. Set $p_m(X) = \sum_{j=0}^{m} a_j X^{m-j}$ for $0 \le m \le N$ and $b_m = (-1)^m q^{m(m-1)/2}$. Induction using $p_{m+1} = Xp_m + a_{m+1}$ and the first partial-lattice recurrence gives

$$p_m(U)x = b_m x_m, \qquad P(U)x = 0.$$

In particular the cyclic module $V_x = \mathbb{C}[U]x$ is finite-dimensional and spanned by $x_0, \dots, x_n$. The adjoint recurrences prove that U' preserves V_x. They can also be written

$$U'p_i(U)x = q^{-i}a_i U'x + q^{n-1}p_{i-1}(U)x \quad (0 \le i \le N),$$

where $p_{-1} = 0$; the case $i = 0$ is tautological, and the case $i = N$ uses the last identity of Lemma 43.8 and $a_N = -q^n b_n t_N$. For a complex number z, the divided difference $D_m(X) = (p_m(X) - p_m(z))/(X - z)$ equals $\sum_{i=0}^{m-1} z^{m-1-i}p_i(X)$. Summing the adjoint recurrence with the weights in this expression gives exactly

$$zU'D_m(U)x = q^{-m}p_m(qz)U'x + q^{n-1}D_m(U)x - U'p_m(U)x. \tag{43.10}$$

Indeed the sum of the scalar terms, augmented by $q^{-m}a_m U'x$, is $q^{-m}p_m(qz)U'x$, while the shifted polynomial sum differs from $q^{n-1}D_m(U)x$ by the p_{m-1} term in the adjoint formula for p_m.

Suppose z, qz are both roots of P, with $z \neq 0$, and let $v = D_N(U)x = (P/(X - z))(U)x$. Then $Uv = zv$ and (43.10) gives $U'v = q^{n-1}z^{-1}v$. If $v \neq 0$, adjointness in the Hermitian pairing implies $|z|^2 = q^{n-1}$ and $U'v = \bar{z}v$. Use the convention of an inner product linear in its first argument. Pairing the last recurrence for $U'x$ with v now gives, on the one hand, $\langle U'x, v\rangle = \bar{z}\langle x, v\rangle$, and on the other,

$$\langle U'x, v\rangle = \frac{q^n p_n(z)}{b_n t_N}\langle x, v\rangle = \frac{q^{2n}}{z}\langle x, v\rangle,$$

since $P(z) = zp_n(z) + a_N = 0$. As $q > 1$, the two scalar factors differ ($q^{n-1} \neq q^{2n}$). Consequently $\langle x, v\rangle = 0$. Applying the adjoint relation repeatedly now gives $\langle U^i x, v\rangle = 0$ for every i. But $v \in V_x$, contradicting positive definiteness. Thus $(P/(X - z))(U)x = 0$ for every nonterminal root in the indicated string.

The minimal annihilating polynomial of x under U divides the greatest common divisor of P and $P/(X - cq^i)$ for $0 \le i < k$. The removed roots are distinct, so this greatest common divisor is $(X - cq^k)S(X)$ up to a nonzero scalar, including when a factor from the string occurs with higher multiplicity. The polynomial $(X - cq^k)R(X)$ therefore annihilates V_x. Since V_x is invariant under both U and U', it is also annihilated by the adjoint of that polynomial in U: the restriction of the adjoint to V_x is the adjoint of the zero endomorphism. Pairing with any I_1-fixed y proves the assertion. Finally, substituting the single rescaling $U_{\mathrm{norm}} = dU$ gives exactly (43.9) with the source's polynomial `HasRootString`; complex conjugation of scalar coefficients is built into the adjoint. □

This proves the second assertion of `chain_lemma` from the actual finite lattice counts. The proof does not apply a spectral projection for an unproved semisimplicity of U, nor remove more than one copy of a repeated string-root factor. It closes that local input of the one-prime operator construction.

43.7 Integral value measures at the actual auxiliary levels

The value-family assertion is stronger than the existence of a measure interpolating values at one prime. Its coefficients must be integral at every coefficient prime not dividing the auxiliary

level, and the family must have a specific Euler-factor relation when a place is inserted. We give the arithmetic construction and isolate the period identities on which it depends.

For a fixed modulus Q of auxiliary places put $q_Q = \prod_{w\in Q} q_w$ and let $\mathcal{I}_Q$ denote the subring of complex numbers whose image is integral for every embedding into $\overline{\mathbb{Q}}_l$ at every rational prime $l \nmid q_Q$. This is a ring, stable under complex conjugation by composing the embeddings with conjugation. A finite expression $\sum_a c_a\vartheta(u_a)$, $c_a \in \mathcal{I}_Q$, $u_a \in U(1)(\mathbb{A})$, will be called an integral evaluation sum. Products of such sums are integral evaluation sums, by multiplication of the points and coefficients. Since tame character values have modulus one,

$$\left|\sum_a c_a\vartheta(u_a)\right|^2 = \sum_{a,b} c_a\overline{c_b}\,\vartheta(u_a u_b^{-1})$$

is another integral evaluation sum. These are the ring and norm-square operations of `Data.TameChar.IsIntegralMeasure` in `New/Exchange/TriangleLevelExponent.lean`; the coefficients and points may depend on Q, but not on the character of modulus Q.

The case $z \geq 2$. Use rank $n = 2z$ and put $M = D.\mathrm{chosenGroupOrder}$ and

$$b = g^{2fn}(gd)^{\sum_\tau (c_\tau - z)}, \qquad F(\vartheta) = \frac{Mb^2}{q_Q^{n-1}}\|\Xi_\vartheta\|^2.$$

These are exactly `denomBound` and `intAlgPart` in `New/Exchange/AlgebraicPartDefs.lean`, with $g = D.\mathrm{gProd}$, $d = P.d$, and the jet and vacuum scales specified in `trivialLift`. In particular F is a positive real number, not a choice of an arbitrary square root of a complex period. We use two input statements about the same normalized theta data:

- The scaled point kernel $bK_\vartheta(u,h)$ is integral in the Weyl lattice at h for all embeddings at $l \nmid q_Q$ (`weylGauge_pointKernel_le_one` in `TriangleLevelExponent.lean`).
- The normalized norm-square identity is $F(\vartheta) = AL^{S\cup Q}(z,(\nu\vartheta)^{-1})$, with $A \neq 0$ independent of ϑ, Q and S the chosen places (`intAlgPart_eq_mul_L`, ultimately `inner_trivialLift_self_eq_L`). Superscripts on L mean omission of both an indicated place and its conjugate.

The analogous period normalizations cannot be substituted from an unramified formula with an unspecified level volume; the factor q_Q^{n-1} in the norm-square formula is used explicitly.

At fixed Q, the source expression for Ξ_ϑ is a sum over the classes u of $U(1)$ at the chosen level and the finite residue-unit lifts at Q. Denote the resulting index set by C_Q and the points by g_a. Its stabilizers at the chosen rank-one level are trivial. The character-twist identity of the Weil kernel separates the varying character as scalar multiplication by its determinant value (on the two arguments, the second argument's factors cancel in the Hermitian pairing). Thus, on expanding the norm and using the class inner product on the target, the coefficient of $\vartheta(\det(g_b/g_a))$ is

$$c_{a,b} = \frac{1}{q_Q^{n-1}}\sum_h \frac{M}{|\Gamma_h|}\,\mathrm{Gram}_h\big(bK_0(g_b,h), bK_0(g_a,h)\big). \tag{43.11}$$

Here K_0 is the kernel at the trivial character on the same modulus, h runs over the fixed target classes, and Γ_h is its target-level class stabilizer. The order of the two entries is the convention in which the character in the coefficient is $\vartheta(\det(g_b/g_a))$; conjugating the convention interchanges a, b in the entire sum. This expansion is `trivialLift_apply`, `modKernel_twist`

and `isIntegralMeasure_intAlgPart`. It proves simultaneous integrality at all desired primes: each $|\Gamma_h|$ divides M and the Gram form of integral Weyl vectors is integral at the corresponding embedding, while q_Q^{n-1} is a unit there. No choice of points or coefficients depends on the embedding or on the character.

At a newly inserted place $v \notin Q$ unramified for the character, write $y = (\nu\vartheta)_E(\varpi_v)$. Because base change of a norm-one character is anticyclotomic, the corresponding value at $\bar{v}$ is y^{-1}. Removing the two local factors multiplies the normalized value by

$$(1 - yq_v^{-z})(1 - y^{-1}q_v^{-z}).$$

The formula follows directly from the Euler product at $z > 1$; both factors are well defined and their product is a positive real number. The positivity and nonvanishing of F follow from the normalized norm identity and nonvanishing of the Hecke Euler product at $z > 1$. Positivity also removes the absolute value in the definition of the ratio: $F(\vartheta) = F(1)\rho(\vartheta)$ for all twists, with no conductor term. This is the $K = \{0\}, m = 0$ case of `Data.Pair.exists_isIntegralMeasure`.

The case $z = 1$. The preceding construction is not available in lift rank two. Use instead the determinant period Λ_ϑ on the pair $(U(2), U(2))$ with $\alpha = \nu\vartheta\upsilon$, determinant degree $c_\tau - 1$ at each infinite place, tame modulus $\mathfrak{m} = D.\mathfrak{m}(\vartheta, 2)$ and residue character $\psi = p.\psi\,\vartheta.\psi$. Let $N \neq 0$ be the uniform denominator for its scalar jets from `exists_modKernel_integral_bound` in `New/Rows/MemberKernels.lean` (homogeneous degree $2\sum_\tau(c_\tau - 1)$). The input period identity is `stdClassSum_det_modKernel_eq` in `New/Rows/Carrier.lean`: with a nonzero constant independent of ϑ, Q, its norm square is proportional to

$$\prod_{w \in S \cup Q} q_w \, \left|L^{S\cup Q}(1, (\nu\vartheta)^{-1})\right|^2.$$

It follows from taking the norm square of the identity $\gamma\Lambda_\vartheta = 1$ and the prescribed norm square of γ, with the above twist by υ. The additional fixed places make only a constant factor. The period identity and uniform denominator are genuine inputs, not consequences of finite Fourier orthogonality.

Write the determinant period at full compact level as the class sum of the determinant kernel with datum $\psi \circ \det$ on residual matrices. Expand this datum by the invertible matrices $Y \in \mathrm{GL}_2(O_E/\mathfrak{m})$; noninvertible matrices have coefficient zero. For each Y, translating the unit-matrix indicator by a compact lift $\widetilde{Y}$ yields its indicator at Y, and the scalar $\psi(\det Y)$ is transported into the character transformation of the kernel. The resulting formula for the period is the sum of the kernels with unit-matrix indicator at $h\widetilde{Y}$, weighted by $|\Gamma_h|^{-1}$. The identity is `detPeriod_eq_stdClassSum_sum`. Applying the scalar twist formula, now to the determinant of $h\widetilde{Y}$, writes $MN\Lambda_\vartheta$ as an integral evaluation sum. To check integrality, use rational left invariance of the kernel (`detKernel_embFin_mul`) to move $h\widetilde{Y}$ to a representative integral above a given $l \nmid q_Q$; this is the integral-representative assertion for the definite standard group. Multiplication by N then gives the uniform integral jet bound at that prime, and $M/|\Gamma_h|$ is an integer. The representative chosen to check integrality may depend on l; the coefficient, unchanged by rational left invariance, does not. Hence

$$F(\vartheta) = |MN\Lambda_\vartheta|^2$$

is an integral evaluation sum at every fixed modulus, by the norm-square calculation.

There is an elementary conductor-change calculation for this period that avoids the borderline Euler product at 1. At a new split place v of norm q, put $X = (\alpha.\mathrm{frobValue}(2, v))^{-1} = q^{-1}y^{-1}$. Write f for the old local vacuum datum. For each subspace U of k_v^2 use the elementary

matrix double-coset representative with lattice reduction U, and apply its transpose on the second group. In the local mixed model the translate of f has support precisely on those integral matrices whose image in the residue space lies in U. Its scalar is $\alpha.\mathrm{frobValue}(2,v)^{\operatorname{codim} U}$, by the determinant/half-density scalar of the split model; transposing exchanges image with row space and does not change the calculation. Thus the Hecke sum with coefficients $(-X)^d q^{d(d-1)/2}$, $d = \operatorname{codim} U$, evaluates at an integral residual matrix of rank r to

$$\sum_{d=0}^{2-r} \begin{bmatrix} 2-r \\ d \end{bmatrix}_q (-1)^d q^{d(d-1)/2} = \prod_{i=0}^{2-r-1} (1-q^i),$$

which is zero unless $r = 2$, when it is one. On nonintegral matrices every translated datum is zero. This proves the local datum equality with the new invertible-residue support; it is the specialization of `sum_heckePolyCoeff_smul_pairWeil` in `Lib/Weil/Unramified/HeckeTransfer.lean`. The finite Gaussian identity follows by expanding $\prod_{i=0}^{k-1}(1-q^iX)$; its coefficients satisfy the subspace-counting recurrence obtained by distinguishing subspaces inside a fixed hyperplane and those not inside it.

The class sum is integration of the kernel on the compact automorphic quotient, with its fixed Haar volume normalization. Each translate by a finite group element has the same class integral, because the integrals of the right translates are invariant. Both the old and new determinant kernels are invariant at the full compact level when the character is unramified at v, so these integrals equal the displayed class sums in their common normalization. Summing the coefficients before integration gives $(1-X)(1-qX)$. Therefore

$$\Lambda_{Q\cup\{v\}} = (1-X)(1-qX)\Lambda_Q.$$

Taking squared absolute values multiplies F by $\prod_{k=0,1}\prod_{\epsilon=\pm1}(1-y^\epsilon q^{k-1})$. This proves the Euler relation of `detPeriod_insert` in `New/Columns/ChainValuation.lean`. Finally the period identity, nonvanishing, and the definition of ρ give $F(\vartheta)/F(1) = q_Q\rho(\vartheta)^2$. This is the $K = \{0,1\}, m = 1$ case of `Data.Pair.exists_isIntegralMeasure`; in particular the extra power of the modulus is indispensable.

43.8 Passage to Galois value families

We now show that the precise measure statement just obtained implies `exists_isFamily` in `New/Selmer/ValueFamilies.lean`. These deductions are independent of the analytic input used to evaluate the periods. For an auxiliary I, let $F_I(\vartheta) = \sum_a c_{I,a}\vartheta(u_{I,a})$ be the finite integral evaluation sum. The quotient map $x \mapsto x/\bar{x}$ of finite ideles surjects onto the norm-one ideles by local Hilbert 90, with integral preimages at almost every place. Choose $y_{I,a}$ lifting $u_{I,a}$ under this quotient (the infinite-place values of a weight-zero character are trivial). With arithmetic Artin reciprocity write $g_{I,a}$ for the Artin image of $y_{I,a}$, and, for the designated sign $s = \pm1$, set

$$V(I,\chi) = \sum_a \iota_\ell(c_{I,a})\chi(g_{I,a}^{-s}).$$

Here χ is any ℓ-adic character, so this is an actual finite sum defining the value on the whole required domain. Because I is auxiliary, its norm is prime to ℓ and all the coefficients have ℓ-adic absolute value at most one. Global class field theory identifies the finite anticyclotomic characters from $U(1)$ of modulus I with characters of the idele quotient by rational ideles, the appropriate integral off-I subgroup, and the conjugation norm; the quotient by the conjugation norm is precisely the norm-one quotient used above. In the source this is the

condition `IsFromU1` together with `exists_mem_tameChars`. For a tame character τ, arithmetic Artin and the inverse-avatar convention give $\tau.\mathrm{invAvatar}(\mathrm{Art}(y)) = \iota_\ell(\tau(y/\bar{y})^{-1})$. Thus

$$V(I, \tau.\mathrm{invAvatar}) = \iota_\ell(F_I(\tau^s)), \qquad V(\varnothing, 1) = \iota_\ell(F(1)) \neq 0. \tag{43.12}$$

In particular both values in this identity are integral.

Take the four continuous ℓ-adic avatars $\eta_{k,\epsilon}$, $k = 0, 1$, $\epsilon = \pm 1$, of the algebraic characters ν^ϵ with Tate factor $q^{-(z-k)}$ on arithmetic Frobenius. With the source's inverse-avatar convention for the varying character, the Euler relations are products of factors $1-\eta_{k,\epsilon}(\phi)\chi(\phi)^t$, for $t = \pm 1$ (the pairing of signs can change with s). At most four factors occur. The valuation ellVal of the source is $\max(0, -\log_\ell |\cdot|_\ell)$, extended by $+\infty$ at zero; it is subadditive on products even if some factors are nonintegral. Hence under the hypothesis that all these factors have valuation at most c,

$$\mathrm{ellVal}(V(I \cup \{P\}, \chi)) \leq \mathrm{ellVal}(V(I, \chi)) + 4c \leq \mathrm{ellVal}(V(I, \chi)) + 8c.$$

The source uses the latter, weaker uniform bound. The norm-one quotient description above ensures that the Euler relation applies to every character satisfying the designated anticyclotomic condition, not just a fixed list of twists.

At the selected ramified old place $w \nmid \ell$ the four avatars have on tame inertia the corresponding nontrivial finite residue character or its inverse. This follows from the local Artin formula: the Tate factor is unramified, and at a residue generator the norm-one character evaluates its prescribed tame part. The selected place has no contribution from its conjugate, by disjointness of the chosen and conjugate places, and nontriviality is exactly the ramified-pair hypothesis. If $\zeta \neq 1$ has order dividing $q_w - 1$, then $v_\ell(\zeta - 1) \leq v_\ell(q_w - 1)$. Indeed $\prod_{i=1}^{m-1}(1 - \xi^i) = m$ for a primitive mth root of unity with $m = q_w - 1$ and one factor is $1 - \zeta$; every factor is integral, so its valuation is bounded by the valuation of the product. This proves the required old-inertia bound without any assumption that the character has ℓ-power order.

Lastly suppose $\mathrm{ellVal}(V(I, \tau.\mathrm{invAvatar})) \leq c + \mathrm{ellVal}(V(\varnothing, 1))$, $c \geq 0$. On the two integral nonzero measures the valuation is the ordinary signed valuation. The interpolation relation and the exact measure normalization yield, because q_I is an ℓ-adic unit,

$$(m+1)v_\ell(\iota_\ell\rho(\tau, s)) = v_\ell(V(I, \tau.\mathrm{invAvatar})) - v_\ell(V(\varnothing, 1)) \leq c.$$

Since $m = 0$ or 1, taking the positive part gives $\mathrm{ellVal}(\iota_\ell\rho(\tau, s)) \leq c$. These prove all the fields and auxiliary conclusions of `exists_isFamily` given the normalized arithmetic value-measure inputs. In particular the argument does not require a choice of a p-adic analytic period or passage to an inverse limit of measures.

43.9 Obligation ledger: Selmer groups, value families, and attachment

The status of a source-level statement is distinguished here from the status of a mathematical reduction. A proved reduction may have unproved input statements, in which case the ledger explicitly retains those input statements as open. A reference closes an obligation only if it supplies the required hypotheses and conclusion; a bibliography of related results does not have that effect.

REFERENCE: finite global and local duality. Milne, *Arithmetic Duality Theorems*, second edition, I.4.10, I.5.1 and I.2.8; Neukirch–Schmidt–Wingberg, *Cohomology of*

Number Fields, second edition, 7.2.15. These cover the finite modules and local pairings required in Lemmas 43.1–43.2, including the real-place correction. The count and prescribed-coset consequence are proved in this chapter.

PROVED: odd-law construction. `New/Selmer/ReciprocityLaw.lean`, `exists_isLaw`, including the higher-depth primitive coordinate, odd conjugation rule and the factor two in the reciprocity condition. The proof in Theorem 43.3 gives precisely the input to the construction of `exists_isPrimeTriangle` (`New/Selmer/TrianglePrimes.lean`); with the earlier detecting-coordinate and triangular constructions this closes that interface.

OPEN: global mirror criterion. `New/Selmer/TiltingLemma.lean`, `IsPencil.length_le_of_unseen_class`. The balanced lattice and an unobserved cohomology class must imply the relevant dependence relation for the actual pencil. The odd-law construction uses scalar Tate modules, so it cannot replace this matrix-valued duality argument with the prescribed restriction and local conditions. The earlier two-stage twist deduction is a reduction to this input, not an independent proof of the input.

PROVED: value-measure reduction and finite arithmetic. `New/Exchange/TriangleLevelExponent.lean`, `isIntegralMeasure_intAlgPart`; `New/Columns/ChainValuation.lean`, `detPeriod_eq_stdClassSum_sum`, `detPeriod_insert`; `New/Selmer/ValueFamilies.lean`, `Data.Pair.exists_isIntegralMeasure`, `exists_isFamily`. Sections 43.7–43.8 prove the finite arithmetic, including the exact conductor-change operator and the two ratio exponents, from the normalized kernel integrality and norm/period formulas. These formulas remain inputs, not established from the measure argument itself.

OPEN: exact value-measure premises. `weylGauge_pointKernel_le_one` (`TriangleLevelExponent.lean`); the prescribed norm-square relation `inner_trivialLift_self_eq_L` in the sharp-data Rallis formula, yielding `intAlgPart_eq_mul_L`; `New/Rows/MemberKernels.lean`, `exists_modKernel_integral_bound`; `New/Rows/Carrier.lean`, `stdClassSum_det_modKernel_eq`, with the same theta normalizations and conductor levels. For the remaining analytic burden see Chapter 34; no arbitrary normalized complex period or general interpolation theorem is used in place of these formulas.

OPEN: geometric seed and weak attachment. `Lib/GaloisRep/Main/SeedRepresentation.lean`, `exists_rep_of_cuspidal_type`; `Lib/GaloisRep/Main/AvatarCompatibility.lean`, `sphericalCompat_of_split`; the supply of exact weak cyclic base changes used in `Lib/GaloisRep/Main/AttachedDeterminant.lean`, `IsWeakBaseChange.exists_attached_descent`; `Lib/GaloisRep/EvenRank/ThetaLift.lean`, `exists_theta_lift_heckePoly`. The trace-reconstruction, quadratic-step and even-rank cancellation reductions from the earlier chapter do not constitute the existence of the requisite automorphic approximations. Specifically, one needs arbitrary weights of definite forms (including singular weights), continuity and finite-type/crystalline or Hodge-jump clauses as indicated, and a local spherical polynomial that actually annihilates the operator. Equality of characteristic polynomials at almost all unramified places does not imply all these clauses.

OPEN: exact tame labelled local–global comparison. `Lib/GaloisRep/LocalGlobal/LabelCompatibility.lean`, `Attached.isLabelledAt`, and `Lib/GaloisRep/LocalGlobal/TypeFrobenius.lean`, `galRep_of_parahoric_type`: finite type and jumps, killing polynomials and determinants for the squarefree depth-zero label with determinant-character blocks, and the simultaneous exponents of Frobenius times all powers of tame inertia. For the latter, the input is the killing polynomial for every

exponent b for the same functional nonzero on the given vector. No assertion merely about the semisimplification of a Weil–Deligne representation has been substituted for the exact killing polynomial.

PROVED: chain orthogonality. `Lib/LocalGL/Iwahori/HeckeCounts.lean`, the relevant instances of `partialHecke_hyperplane_count` and `partialHecke_dual`; `Lib/LocalGL/Tadic/ChainLemma.lean`, second assertion of `chain_lemma`. Lemmas 43.8–43.9 establish the orthogonality identity (43.9), including repeated roots, by a direct finite lattice sum and adjoint argument, rather than attributing it to the spherical-unitary-dual classification.

PROVED: local trace reduction and finite integral patching. `Lib/LocalGL/Iwahori/HeckeOnLineStabiliser.lean`, `heckeCoset_mem_parahoricEigenspace`; `Lib/GaloisRep/LocalGlobal/TypeFrobenius.lean`, `hecke_eq_frob_on_parahoricEigenspace`; `Lib/LocalGL/TaylorWiles/AuxPlaceOperators.lean`, `exists_higman_operator`; `New/Patching/OnePrimeTie.lean`, `BaseLevel.exists_tie` and `BaseLevel.exists_diagOp_mem_ellIntegralForms`. Lemmas 43.4, 43.6, 43.7 and Proposition 43.5 prove the integral and polynomial deductions, conditional on the exact local comparison, Ribet polynomial, and weak attachment identified in the chapter. In particular the top coefficient is extracted in the actual integral old-level module, including the stabilizer factor and the cutoff polynomial, instead of assuming a formal patched module is free.

OPEN: Ribet polynomial. `New/Selmer/RibetFirstOrder.lean`, `ribet_first_order`, the residual exact first-order ideal membership and size of the top coefficient. The determinant or Fitting length calculation alone is not the full Ribet polynomial.

The remaining open entries are genuine proof obligations of the present written reconstruction. In particular, closing the odd reciprocity law and the finite integral patching reduction does not establish `denominatorLe_rowBudget` without the unproved arithmetic and local–global inputs.

Chapter 44

Automorphic Galois representations by classical base change, and the balanced row

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 47, 48 and the final audit in Chapter 57.

There is a useful alternative to reconstructing the geometric seed and the subsequent limits in the source. One may invoke the published endoscopic base change and regular algebraic cuspidal attachment theorems, provided that one actually treats the generally *noncuspidal* base change of a definite eigenform. In this chapter we explain that alternative, including the parity twists, the treatment of monodromy, and the precise relation between the classical normalizations and the polynomials used in the proof. We then prove the remaining balanced-lattice estimate. The alternative does not require one to check the entire PEL construction of `Lib/GaloisRep/Main/SeedRepresentation.lean`: it proves the required downstream interface `sphericalCompat_std` directly. It is important that this is an alternative conventional proof using substantially deeper published results, rather than a claim that the formal geometric construction has been independently checked.

44.1 The field and group to which base change applies

Write E/F for the CM field in the proof and its maximal totally real subfield. The definite group for attachment is $G = U(I_N)$, not the indefinite real group of the Bessel construction. The source hypothesis `WeilDataE` includes `StablyUnramE`, i.e. E contains a primitive 5040th root of unity. In particular $[F : \mathbb{Q}] > 1$. It also contains i, ω, where

$$i^2 = -1, \qquad \omega + \bar{\omega} = 1, \qquad \omega\bar{\omega} = 2.$$

The integral element $i\omega$ differs from its conjugate by the unit i. Thus no finite place of E can have nontrivial inertia for E/F: at a ramified place conjugation would act trivially on the residue field, contradicting the unit difference. At an inert place E/F is therefore an unramified quadratic extension. The norm from its unit group onto the unit group of F is surjective. The identity Hermitian form of rank N is quasi-split there. Indeed Hermitian spaces over an unramified quadratic local extension are classified by dimension and determinant modulo norms, and the quasi-split form, the sum of $\lfloor N/2 \rfloor$ hyperbolic planes and possibly a line, has unit determinant. At a split place our group is $\mathrm{GL}_N(F_v)$. These are precisely the

local hypotheses of the classical definite-group base change used below. No assumption that the coefficient prime ℓ split everywhere in E is needed in the argument.

We will use the unramified splitting character υ of $\mathbb{A}_E^\times/E^\times$ available in the actual Weil data. Its infinity type at each chosen embedding is $z/|z|_{\mathbb{C}}^{1/2}$ (here $|z|_{\mathbb{C}} = z\bar{z}$), it satisfies $\upsilon^c = \upsilon^{-1}$, and every finite local component is unramified. The construction with these precise properties appears in Proposition 6.4. This character is stronger than a character of vaguely specified parity: it will ensure that the correction for parity does not alter the local inertial types at any finite place.

Published base change. Use J.-P. Labesse, *Changement de base CM et séries discrètes*, in *On the Stabilization of the Trace Formula*, vol. 1, International Press (2011), Corollary 5.3 and the discussion immediately preceding it (pp. 42–43 of the author's manuscript, https://webusers.imj-prg.fr/~michael.harris/CBCM.pdf). Its hypothesis (*) is automatic for a group compact at all real places; the required discrete-series condition at infinity also holds for a compact group. A constituent π of the finite-level definite automorphic space thus has base change to $\mathrm{GL}_N(\mathbb{A}_E)$ of the form

$$\Pi = \Pi_1 \boxplus \cdots \boxplus \Pi_a, \qquad \sum_i n_i = N. \tag{44.1}$$

Each Π_i is discrete, conjugate self-dual, of rank n_i. At the split places the local representation of Π is the local base change of π itself. At the inert spherical places the same matching is the unramified base change. The infinitesimal character is that of the base-changed algebraic coefficient system. These are the assertions of the cited corollary; the stronger statement about precise archimedean representations under an additional cuspidality assumption is not used. For the formulation for totally definite groups, see also L. Guerberoff, *Modularity lifting theorems for Galois representations of unitary type*, Compositio Math. **147** (2011), 1022–1058, Theorem 2.2 and Remark (1) following it, DOI https://doi.org/10.1112/S0010437X10005154. His Theorem 2.3 is a related construction with additional assumptions on the coefficient prime in the setup. We do not invoke that theorem outside its hypotheses; instead we carry out its base-change and twisting argument using the present splitting character and the cuspidal attachment theorem for an arbitrary coefficient prime.

Decomposing the discrete rather than assuming it cuspidal. The Mœglin–Waldspurger classification of the discrete spectrum of GL_n gives $n_i = a_i b_i$ and a unitary cuspidal representation ρ_i of $\mathrm{GL}_{a_i}(\mathbb{A}_E)$ such that Π_i is the Speh representation corresponding to

$$\rho_i|\det|^{(b_i-1)/2-j}, \qquad 0 \le j < b_i. \tag{44.2}$$

For this result see C. Mœglin and J.-L. Waldspurger, *Le spectre résiduel de GL(n)*, Ann. Sci. ENS (4) **22** (1989), 605–674; equivalently J. Arthur's exposition, *An introduction to the trace formula*, in Harmonic Analysis, the Trace Formula, and Shimura Varieties (Clay Math. Proc. 4, 2005), Theorem 26.2, p. 191 (https://www.claymath.org/library/cw/arthur/pdf/62.pdf). The latter explicitly describes the Langlands quotient and all half-integral shifts in (44.2). Uniqueness in this classification and conjugate self-duality of Π_i imply $\rho_i^c \simeq \rho_i^\vee$: conjugate duality reverses the list of shifts and replaces ρ_i by its conjugate dual. It is generally not true that each ρ_i is already algebraic in the required rank. The next correction is necessary.

44.2 The parity correction and exact spherical attachment

We number the rows of a highest weight starting at 1. If $\kappa = (\kappa_1, \kappa_2)$ is the source weight, write

$$\lambda_j(v) = \kappa_2(v) + \sum_{h=j}^{N-1} \kappa_1(v,h), \qquad 1 \le j \le N.$$

Here $\lambda_1 \ge \cdots \ge \lambda_N$. The actual rational covariance uses the conjugate embedding (see the identity `stdSpace_weightRep_apply` in `Lib/GLn/SchurModule/WeightRepBasic.lean`); at the chosen embedding $\sigma = \Phi(v)$ this is the dual highest weight $(-\lambda_N, \dots, -\lambda_1)$. At $\bar\sigma$ it is λ. With the convention that the automorphic representation at infinity is dual to the coefficient representation, the algebraic coefficient of Π is thus

$$a_\sigma = (-\lambda_N, \dots, -\lambda_1), \qquad a_{\bar\sigma} = (\lambda_1, \dots, \lambda_N). \tag{44.3}$$

The multiset of jumps wanted in the proof is

$$J(\sigma) = \{a_{\sigma,j} + N - j : 1 \le j \le N\}.$$

In particular, at the chosen place this is $\{j - 1 - \lambda_j\}$, and at its conjugate $\{\lambda_j + N - j\}$. These are exactly `jumps` in `Lib/Weil/Unramified/SphericalHecke.lean`; they are distinct even if some successive highest weights are equal. There is consequently no obstruction coming from a repeated highest weight in the coefficient system. Strict regularity here is regularity of the shifted infinitesimal character, not strict inequality of all the unshifted λ_j.

Consider the complex Weil characters at infinity of the base change after rank-N arithmetic normalization $|\det|^{(1-N)/2}$. Their exponents at each embedding are integral and, as multisets, are the negatives of $J(\sigma)$: the contragredient of the algebraic coefficient has highest weight $(-a_{\sigma,N}, \dots, -a_{\sigma,1})$, to which one adds the half sum $(N + 1 - 2j)/2$ and then $(1 - N)/2$. This argument uses only the infinitesimal character, so applies to the residual Speh blocks too. Write p, p' for the two exponents of any infinite Weil character of ρ_i before normalizing it arithmetically. Its copy with shift $(b_i - 1)/2 - j$ contributes the exponents

$$p + \frac{b_i - 1}{2} - j + \frac{1 - N}{2}, \qquad p' + \frac{b_i - 1}{2} - j + \frac{1 - N}{2}.$$

Their integrality implies $p, p' \equiv (N - b_i)/2 \pmod{\mathbb{Z}}$. Choose $\epsilon_i \in \{0, 1\}$ satisfying

$$\epsilon_i \equiv a_i + b_i - N - 1 \pmod 2. \tag{44.4}$$

Then $\widetilde{\rho}_i = \rho_i \otimes (\upsilon^{\epsilon_i} \circ \det)$ is regular algebraic in rank a_i: its rank-a_i normalized exponents are

$$p + \epsilon_i/2 + (1 - a_i)/2, \qquad p' - \epsilon_i/2 + (1 - a_i)/2$$

at the chosen pair, and are integral by (44.4). They are distinct at each embedding because the unshifted cuspidal exponents form a submultiset of one shifted contribution to the regular rank-N system. The opposite choice at a conjugate embedding changes both signs consistently. Furthermore $\widetilde{\rho}_i^c \simeq \widetilde{\rho}_i^\vee$. This is a regular algebraic conjugate self-dual cuspidal representation (cuspidality and conjugate duality have already been checked, and the polarization in the cuspidal attachment theorem is the trivial character).

The required cuspidal attachment theorem. We invoke T. Barnet-Lamb, T. Gee, D. Geraghty and R. Taylor, *Potential automorphy and change of weight*, Ann. of Math. **179** (2014), 501–609, Theorem 2.1.1, p. 538, https://doi.org/10.4007/annals.2014.17

9.2.3. Its assumptions are regular algebraic, cuspidal and polarized, just verified. It gives a continuous semisimple representation $r_\ell(\widetilde{\rho}_i)$ of dimension a_i, de Rham with the specified Hodge–Tate weights; away from ℓ its Frobenius-semisimplified Weil–Deligne parameter is the local Langlands parameter in the twist $|\det|^{(1-a_i)/2}$. Here geometric Frobenius and the convention $\mathrm{HT}(\varepsilon_\ell) = \{-1\}$ are used. We also use the crystalline assertion for unramified local components of the same theorem. For the more general comparison above ℓ needed at the tame but nonspherical types below, use the same authors, *Local-global compatibility for $l = p$, II*, Ann. Sci. ENS (4) **47** (2014), 165–179, Theorem 1.1, https://arxiv.org/abs/1105.2242. The partial order in that theorem keeps the Weil semisimplification and only allows smaller Jordan partitions of the monodromy; for a local parameter with zero monodromy this is enough, without the Shin-regularity hypothesis of the stronger equality assertion.

For each i, j define the actual algebraic Hecke character

$$\zeta_{ij} = \upsilon^{-\epsilon_i}|\cdot|^{(a_i+b_i-N-1)/2-j}, \qquad 0 \le j < b_i. \tag{44.5}$$

The expression looks half-integral, but both infinity exponents are integers by (44.4); it is of type A_0. Since υ is unramified at every finite place, this character is too. It therefore has the usual ℓ-adic avatar with crystalline local characters above ℓ. For precision, an algebraic Hecke character with infinity component $\prod_\sigma \sigma(x)^{p_\sigma}$ has avatar given using the geometric reciprocity map by its finite component, multiplied at the primes above ℓ by $\prod_\sigma(\iota\sigma)(x_\ell)^{p_\sigma}$. This cancels on principal ideles and gives a continuous character of the abelianized Galois group. Its labelled Hodge–Tate weights in our convention are $-p_\sigma$. Its local Weil character is the finite local component transported by ι. In the unramified crystalline case this last assertion, including the geometric Frobenius sign, follows from B. Conrad, *Lifting global representations with local properties*, Appendix B, Proposition B.4 and Lemma B.3, https://math.stanford.edu/~conrad/papers/locchar.pdf. Conrad uses arithmetic reciprocity; his formula for φ^f is the algebraic torus character on a uniformizer divided by the arithmetic Galois character there. With the inverse Artin map these two algebraic factors cancel, giving precisely the finite component of our character. For a finite-order local component, pass to a finite extension killing that component and use the same computation with descent. Thus the rank-one inputs called `avatarCompatAt_of_stablyUnram` need not be obtained recursively from odd-rank geometry.

We can now define

$$R_\pi = \bigoplus_i \bigoplus_{j=0}^{b_i-1} r_\ell(\widetilde{\rho}_i) \otimes \zeta_{ij,\ell}. \tag{44.6}$$

It has dimension $\sum_i a_i b_i = N$, is continuous and semisimple. The exponent in (44.5) is exactly the difference between rank-N normalization with the Speh shift and rank-a_i normalization:

$$\frac{1-N}{2} + \frac{b_i-1}{2} - j - \frac{1-a_i}{2} = \frac{a_i+b_i-N-1}{2} - j.$$

This proves that the local parameters in (44.6) are the normalized parameters of the constituents (44.2). At a matching split place the standard Langlands-quotient description of the Speh representation gives the direct sum of the parameters of the shifted constituents. One may check the ordering of these standard modules as follows. The cuspidal constituents $\widetilde{\rho}_i$ are unitary and conjugate self-dual; the purity assertion in Theorem 2.1.1 for a coefficient prime different from the residue characteristic gives purity of weight $a_i - 1$ of their arithmetically normalized local Weil–Deligne parameters, hence temperedness in the unitary normalization (the local correspondence sends the pure weight-zero Weil–Deligne representations to tempered representations). Thus the shifts in (44.2) are standard decreasing real exponents on tempered factors. The local isobaric sum has the same Langlands-quotient direct-sum property.

Equivalently this follows by the usual Langlands classification for the local components of the discrete isobaric sum in (44.1). In the following we use only the local places at which the resulting parameter has zero monodromy, not arbitrary identities of filtered modules. The Hodge–Tate jumps of (44.6), by the same shift computation at infinity and the avatar calculation, are exactly (44.3) with the added $N - j$.

Proposition 44.1 (Spherical compatibility in all the ranks of the definite proof)**.** *For a nonzero eigenform f of weight κ in the standard definite group with the Weil data of the proof and Hecke system P_w, there is a continuous semisimple rank-N representation R satisfying* `AttachedPR`*. If f is spherical at w, then R is unramified away from ℓ or crystalline above ℓ at w, with jumps J. At a split spherical w, every Hecke polynomial Q of f there kills every geometric Frobenius in the precise sense of* `PolyKillsAt`*. These assertions hold for every weakly attached representation, up to conjugacy.*

Proof. Use a finite-level subspace containing f and decompose it into automorphic constituents. It is a finite-dimensional space of functions on finitely many double classes with compact weight, and the automorphic discrete spectrum decomposition, with finite multiplicities, supplies finitely many nonzero projections of this vector. The projection commutes with right translations and with all the spherical Hecke operators in question. Choose a constituent in which the projection is nonzero; it has the same Hecke system at almost all split places of prime norm. Form R by (44.6). The split finite local parameter has the following precise normalization. If $\alpha_1, \dots, \alpha_N$ are normalized Satake parameters (values of unramified inducing characters at a uniformizer), $T_{w,j}$ acts on the spherical line by

$$q^{j(N-j)/2} e_j(\alpha_1, \dots, \alpha_N).$$

The polynomial in `SphericalHecke.lean` is $\sum_{j=0}^{N} (-1)^j q^{j(j-1)/2} T_{w,j} X^{N-j}$ on this line. Since $j(j-1)/2 + j(N-j)/2 = j(N-1)/2$, its roots are exactly $q^{(N-1)/2}\alpha_i$, the geometric-Frobenius eigenvalues in arithmetic normalization. The actual characteristic polynomial thus equals ιP_w at all these places, not merely modulo an error. Their cofinite prime-norm subset belongs to the source's filter `mostFrobenii`; this proves `Attached`. Weak attachment has unique semisimple trace by the proper localized Frobenius filters and Chebotarev (Section 42.11); hence every attached representation is conjugate to R.

At a place where f is spherical, choose a constituent projection which is nonzero, as above. Its local base change is unramified, also for an inert spherical place. Its local Langlands parameter is consequently unramified with zero monodromy. The direct sum making up that parameter cannot hide monodromy or ramification of a constituent. It follows by the comparison theorem that each of the summands in (44.6) is unramified or crystalline at w, and its normalized Frobenius characteristic polynomial is the corresponding factor of the spherical polynomial. This proves the good local condition and the required characteristic polynomial. Frobenius semisimplification does not change the characteristic polynomial of a linear Frobenius, so the genuine annihilation follows by Cayley–Hamilton, with no assumption that its Jordan blocks are trivial.

To check the meaning of the killing statement at $w \mid \ell$, let D_{pcris} be the potential crystalline period module, the union of modules over finite extensions on which R is crystalline. In the usual Weil–Deligne construction, a geometric Weil element s acting on residue fields by p^{-m} acts linearly on the potential period module by $\varphi^m \circ s$ (with the Galois action on the representation understood); these factors cancel the residue-field semilinearity. Extending this operator by the period-ring comparison is well defined because the invariant periods span. This is exactly the definition `linearFrob` in `Lib/PadicHodge/Crystalline/Defs.lean`, not the action of $R(s)$ alone. A polynomial identity on this module therefore holds on its

base extension to the actual period ring by linearity, and is the source's `IsKilledBy`. At a good place the geometric-Frobenius operator is independent of the inertia lift; at any fixed embedding its characteristic polynomial is the same ιQ. This verifies the full `PolyKillsAt` clause above the coefficient prime as well. Finally the jump formula holds by (44.6) and (44.3). □

This proves the *needed* interface of `Lib/GaloisRep/Main/SphericalAllRanks.lean` with published theorems, for the fields with Weil data used in the monograph. It avoids the auxiliary parity-cancelling theta lifts as well as the weak cyclic base-change construction. It does not purport to reprove the geometric seed lemma as a theorem about its broader intermediate PEL data. Those constructions are no longer premises of this route. Notice also the role of the field and weight check: quoting a regular algebraic cuspidal theorem applied directly to an arbitrary definite eigenform would leave both the rank of the representation and its Hodge and local comparison unjustified.

44.3 The exact tame label and the monodromy condition

We now verify the particular local-global assertion `Attached.isLabelledAt`. It is stronger than matching unramified Frobenius traces, but does not require knowing a differential equation for a general potentially semistable representation.

Let $K = E_w$ at a split place, $q = |k|$ its residue cardinality, $K_0 = \mathrm{GL}_N(\mathcal{O}_K)$, and $K_1 = 1+\mathfrak{m}_K M_N(\mathcal{O}_K)$. Write $T : \bar{k}^\times \hookrightarrow \mathbb{C}^\times$ for the faithful character used in the labels. A label c of degree d is monic squarefree with nonzero constant term; factor it as a product of distinct monic irreducibles c_P. In the notation of `Lib/LocalGL/Green/FrobeniusOrbits.lean`, for a root x of c_P of degree h let θ_x be the character of $k_h^\times$ such that if $T(x) = \exp(2\pi i b/(q^h-1))$, then $\theta_x(y) = T(y)^b$. The Green character τ_P is the irreducible finite cuspidal representation belonging to this regular character modulo Frobenius conjugacy. The sign can be checked directly: its regular elliptic character is $(-1)^{h-1}\sum_{j=0}^{h-1}\theta_x(y^{q^j})$, the same as the source's $(-1)^h C_{\theta_x}$ in degree h (the cuspidal function C has the extra minus sign there). The finite label used in the source is the Harish-Chandra product of these τ_P.

In `HasProductLabel` the remaining blocks are $\chi_l \circ \det$ on $\mathrm{GL}_{t_l}(k)$, with $d + \sum_l t_l = N$. The separability requirement on

$$c\prod_l (X - x_{\chi_l})$$

means that their characters are distinct from one another and their degree-one orbits are disjoint from those already in c. Here x_χ is the inverse character name in the cyclic-group convention above, so $\theta_{x_\chi} = \chi$ on $k^\times$. In particular the finite Harish-Chandra product

$$\Xi = \left(\underset{P}{*}\,\tau_P\right) * \left(\underset{l}{*}(\chi_l \circ \det)\right) \tag{44.7}$$

is irreducible. This uses the decomposition of the Green representation ring into the tensor product of the symmetric-function rings for its distinct primitive cuspidal characters; each displayed factor lies in its own tensor component. The product of irreducibles in disjoint components is irreducible. The representation-ring statement is also recalled by J. Shotton, *The Breuil–Mézard conjecture when $l \neq p$*, Duke Math. J. **167** (2018), 603–678, Section 6.3, in particular Proposition 6.9 (https://maths.dur.ac.uk/users/jack.g.shotton/papers/gln.pdf). The unnormalized sum $\sum_{k\in\mathrm{GL}_N(k)} \overline{\Xi(k)}\, k$ in `HasLabel` is a nonzero scalar times the isotypic idempotent for (44.7), by ordinary character orthogonality. Thus nonvanishing on f is precisely the occurrence of this finite K_0/K_1 representation, not merely a test of its central character.

Lemma 44.2 (Depth-zero local parameter with this particular type)**.** *Suppose π is an irreducible smooth representation of $\mathrm{GL}_N(K)$ whose K_1-fixed subspace contains (44.7). Its Weil–Deligne parameter has zero monodromy. The inertial representation is a sum of one copy of each regular Frobenius orbit belonging to c_P and t_l copies of χ_l for each determinant block. In particular it is trivial on wild inertia.*

Proof. We recall why the crystalline type assertion here holds even when π is not generic. In the Langlands classification, π is the quotient of normalized induction from essentially square-integrable segments. Passing to K_1-invariants is exact, and the invariants of the parabolic induction are the Harish-Chandra induction of the invariants of its factors: restrict an induced function to K_0 using Iwasawa decomposition, then reduce its right K_1-invariance. Each segment of length m based on a supercuspidal τ of rank h embeds into induction of m unramified twists of τ. So this segment has zero K_1-invariants unless τ has principal-level-one invariants. In the latter case the supercuspidal is depth zero, and its K_1-invariants form the corresponding irreducible cuspidal finite representation σ_0 of $\mathrm{GL}_h(k)$; the segment invariants have finite Harish-Chandra cuspidal support consisting of m copies of σ_0. This is the depth-zero case of the Schneider–Zink type calculation, and follows as well by the usual description of depth-zero supercuspidals as compactly induced from $K^\times \mathrm{GL}_h(\mathcal{O}_K)$ with the irreducible cuspidal residue representation. The exact induction and multiplicity calculation for Langlands quotients, without a genericity hypothesis for the forward implication, is in Shotton, loc. cit., Theorem 6.20 and Corollary 6.21(1), the latter being a form of P. Schneider and E.-W. Zink, *K-types for the tempered components of a p-adic general linear group*, J. Reine Angew. Math. **517** (1999), 161–208, Proposition 5.10. Consequently the finite cuspidal supports must equal those of (44.7).

For a cuspidal factor of degree $h > 1$ in c, its multiplicity in that support is one; no segment of length > 1 based on it is possible. This same reasoning applies to the degree-one factors of c. A determinant block $\chi_l \circ \det$ has finite cuspidal support t_l copies of the rank-one character χ_l. Twisting by its inverse and restricting to the separate tensor component of the Green ring, the finite representation which must occur is the trivial representation of $\mathrm{GL}_{t_l}(k)$. If a segment of rank $m > 1$ occurred in this component, its (twisted) invariant part would have no trivial $\mathrm{GL}_m(k)$ constituent: otherwise its square-integrable-segment representation would have $\mathrm{GL}_m(\mathcal{O}_K)$-fixed vectors. This is impossible, since the generalized Steinberg representation of length > 1 is not spherical. Independently, its monodromy is the nontrivial Jordan block of length m, whereas a spherical local representation has unramified parameter with zero monodromy. By finite Frobenius reciprocity the Harish-Chandra induction over these factors cannot contain the determinant character if any corresponding factor does not contain its determinant character. The Green-ring tensor decomposition lets one apply that check separately in each distinct cuspidal component. Thus no segment of length $m > 1$ occurs in any determinant block either. All segments have length one, proving the zero monodromy assertion. This argument is the elementary explanation of why (44.7) is the zero-monodromy rather than the Steinberg type, including for a non-generic irreducible local representation. It does not assume that occurrence of arbitrary pro-p-Iwahori invariants implies zero monodromy.

Finally the depth-zero local Langlands correspondence sends a regular cuspidal residue character θ_x to induction from the unramified extension of degree h of a character with that residue restriction; the rectifier is unramified and hence disappears on inertia. This is the depth-zero case of the tame local Langlands correspondence of Bushnell–Henniart; for the normalization of the rectifier see M. Adrian and D. Roe, *Rectifiers and the local Langlands correspondence: the unramified case*, Proposition 3.4 and Theorem 9.5 (`https://math.mit.edu/~roed/writings/research/rectifiers.pdf`). Unramified twists likewise do not affect inertia. Thus the inertia characters come in precisely the regular orbits belonging

to the factors of c, with the listed rank-one multiplicities in the determinant blocks. □

Lemma 44.3 (The source exponent convention)**.** *Let s be inertial, and let a be its tame exponent as in* `IsTameExponent`*, taken relative to a root z satisfying $z^{q^{d!}-1} = \varpi$. On the regular orbit named by a root x of c_P, the eigenvalues of s under the depth-zero inertial correspondence are*

$$T(x)^a, \quad T(x^q)^a, \quad \dots, \quad T(x^{q^{h-1}})^a.$$

For a determinant-character block the corresponding eigenvalue is $T(x_{\chi_l})^a$ with multiplicity t_l.

Proof. The degree h divides $d!$, and $q^h - 1$ divides $q^{d!} - 1$. Put $M = q^{d!} - 1$ and $m = q^h - 1$. Raising z to M/m gives a root z_m with $z_m^m = \varpi$. Reduction of $s(z_m)/z_m$ has T-value $\exp(2\pi ia/m)$. This construction is independent of a choice of the root z_m, since inertia fixes prime-to-residue-characteristic roots of unity. For the unramified degree-h extension, the tame local reciprocity formula for units, with uniformizers mapping to geometric Frobenius, identifies this ratio on $\varpi^{1/(q^h-1)}$ with the residue of the unit representing s in the abelianized inertia group. One way to see the sign is to use the Hilbert symbol first with *arithmetic* reciprocity: arithmetic Frobenius acts on an mth root of a unit u by the ratio reducing to $\bar{u}^{(q^h-1)/m}$; skew symmetry gives the inverse when interchanging unit and uniformizer. Passing to geometric reciprocity reverses it again. This is the usual tame-symbol formula, here with $m = q^h - 1$. Therefore the character of s for $\theta_x = T^b$ is $\exp(2\pi iab/m) = T(x)^a$. Conjugation by Frobenius permutes the h roots x^{q^j}, giving the asserted list; for the determinant characters take $h = 1$. This is precisely the character-name identity `IsTameExponent.pow_apply_charName` in `Lib/GaloisRep/LocalGlobal/TameExponent.lean`. The prime-to-residue-characteristic quotient used in this argument is cyclic, so such exponents always exist, all residue classes occur, and one can choose exponent 1. A topological tame generator has an exponent coprime to M. The field of size q^h contains all the roots of c_P and hence these roots satisfy $x^{q^h-1} = 1$; this verifies also the power conditions in `IsLabel.root_pow_eq_one`. □

Theorem 44.4 (Exact tame-label compatibility)**.** *Under the assumptions of* `AttachedIsLabelledAt` *in* `Lib/GaloisRep/LocalGlobal/LabelCompatibility.lean`*, let R be the attached representation of f. At A above the indicated split place the representation has finite type and the prescribed jumps. For every source tame exponent pair (s, a) the polynomial*

$$P_{s,a}(X) = \prod_{x \in c(\bar{k})} (X - \iota(T(x)^a)) \prod_l (X - \iota(T(x_{\chi_l})^a))^{t_l} \tag{44.8}$$

kills s in the precise sense `PolyKillsAt`*. If $w \nmid \ell$ its constant term is $(-1)^N \det R(s)$.*

Proof. Project the nonzero finite-label isotypic component of the actual eigenform onto an irreducible automorphic constituent in which it is nonzero. That constituent has the original weak Hecke system, so its attached representation is R by Proposition 44.1 and uniqueness. Its split local Langlands parameter has exactly the type of Lemma 44.2. Since this parameter has zero monodromy, the direct factors in (44.6) have zero monodromy. The cuspidal comparisons used there give precisely this inertial representation. At $w \nmid \ell$, Frobenius semisimplification preserves restriction to inertia on the Weil representation, and the absence of monodromy means the Weil action agrees with the original Galois action. Inertia has finite image, so it is diagonalizable in characteristic zero on each element. Lemma 44.3 gives the characteristic polynomial of $R(s)$ itself, hence both the required annihilation and determinant. At $w \mid \ell$ de Rham representations have potential semistable comparison, and the monodromy-partition inequality of BLGGT, Theorem 1.1 cited above, forces zero monodromy because the automorphic parameter has zero monodromy. Consequently the representations are potentially

crystalline. On the potential crystalline module the inertial action is exactly the same finite type; the semisimplification in the comparison does not change that action. The same polynomial annihilates it, and the period-module argument in the proof of Proposition 44.1 gives the assertion on the full source period ring. The jumps follow from (44.3). This proves the specified local conditions as actual polynomial identities, not just equality of sets of eigenvalues modulo semisimplification. □

The theorem bypasses the induction on prime-degree unramified extensions of the residue field, and in particular the weak cyclic base change and matching Shintani character of `Lib/GaloisRep/BaseChange/BaseChangeLabel.lean`. All these source arguments can be replaced at their interface by the direct published local and global Langlands results just stated.

44.4 Iwahori exponents and simultaneous Frobenius powers

For another local application one wants an annihilator for $s\tau^b$ *for every* $b \geq 0$ with the same local exponent. This is the key point in `Lib/GaloisRep/LocalGlobal/TypeFrobenius.lean`. We spell out how the preceding ordinary correspondence implies it instead of making an inference from separate characteristic polynomials of s and τ.

Write I_1 for the pro-p Iwahori subgroup (upper unitriangular modulo the prime), e_1 for its probability-average projector, B for the upper triangular Borel and $t = \mathrm{diag}(t_1, \dots, t_N)$ for a dominant torus element in the source convention $|t_1|_K \leq \cdots \leq |t_N|_K$. The source exponent is a functional satisfying

$$l(\pi(t)e_1 v) = \prod_{i=1}^{N} \theta_i(t_i)|t_i|_K^{N-i}\, l(v). \tag{44.9}$$

The torus power $N-i$ is not the $\delta_B^{1/2}$ normalization. If α_i are the characters in the normalized Jacquet module, then

$$\theta_i = |\cdot|_K^{(1-N)/2}\alpha_i, \qquad \delta_B^{1/2}(t) = \prod_i |t_i|_K^{(N+1-2i)/2}. \tag{44.10}$$

For example one can derive this relation by evaluating a normalized principal-series induced function at the identity, after the projector e_1. Factor $I_1 = (I_1 \cap B)(I_1 \cap N^-)$; for dominant t the conjugation $t^{-1}(I_1 \cap N^-)t$ stays in I_1. The first factor has trivial inducing character and modulus, and a right I_1-fixed vector is fixed by the conjugate of the second factor. Therefore $\mathrm{ev}_1(e_1\pi(t)e_1 v) = \delta_B^{1/2}(t)\prod_i \alpha_i(t_i)\,\mathrm{ev}_1(e_1 v)$, giving (44.9).

For the more general sorted Iwahori eigencharacter in `Attached.local_global_of_eigen character`, the normalized Jacquet-module description is justified by Casselman's canonical lifting: see W. Casselman, *Introduction to the theory of admissible representations of p-adic reductive groups*, draft 1 May 1995, Section 4.1, especially Proposition 4.1.4 and the description of the canonical lifting (`https://www.uni-muenster.de/imperia/md/content/theoretische_mathematik/topologie/casselma.pdf`). The projection to the Jacquet module is injective on the invertible part of the strictly contracting torus operator on the chosen Iwahori invariant space; an eigenvector for its nonzero eigenvalue lies in that part. On this part the action is the unnormalized torus action, yielding exactly (44.10) after taking normalized Jacquet modules. The geometric lemma and the segment Jacquet formula show that a torus character appearing in a normalized Jacquet module of an irreducible $\mathrm{GL}_N(K)$ must be a shuffle of those from its essentially square-integrable constituents, preserving the order of each segment. In particular all constituents have rank-one supercuspidal support if the full

torus Jacquet module is nonzero. A segment of length $m > 1$ on a character χ has ordered normalized torus characters

$$\chi|\cdot|_K^{b}, \quad \chi|\cdot|_K^{b-1}, \quad \dots, \quad \chi|\cdot|_K^{b-m+1}.$$

At a uniformizer their complex magnitudes are strictly increasing. This calculation of ordered characters follows from the normalized Jacquet formula for generalized Steinberg segments (A. Zelevinsky, *Induced representations of reductive p-adic groups II*, Ann. Sci. ENS (4) **13** (1980), Propositions 3.4 and 9.5); see also the convenient recapitulation in H. Atobe, *On an algorithm to compute derivatives*, Manuscripta Math. **167** (2022), Section 2.1. The geometric lemma for the standard parabolic induces shuffles of these sequences; a quotient can only remove characters, by exactness of Jacquet modules. Consequently an eigencharacter θ whose magnitudes at a uniformizer are nonincreasing cannot admit a segment of length greater than one. The local parameter has zero monodromy and, in the arithmetic normalization, its Weil semisimplification is $\bigoplus_i \theta_i$, with the rank-one identifications using geometric reciprocity. Notice that the sorting condition is essential; otherwise the pro-p Iwahori condition alone would include Steinberg representations. The same reasoning as in Theorem 44.4 now yields the required finite-type and jump conditions, and for any Weil element of degree $k \geq 0$ with Artin preimage $x \in K^{\times}$ the killing polynomial $\prod_i (X - \iota\theta_i(x))$. This is the interface `Attached.local_global_of_eigencharacter`; there is no need to run a cyclic base-change induction to reduce it to unramified spherical comparison.

Proposition 44.5 (The common exponent on a parahoric vector)**.** *Suppose $w \nmid \ell$, and y is a nonzero parahoric vector in the source isotypic space: the parahoric is the preimage of a standard flag parabolic with blocks $f(i)$, and its action on y factors through a determinant character on each block. Let u be a unit with values d_i of these diagonal characters, and assume different blocks have distinct values at u. Let s lift geometric Frobenius with Artin preimage ϖ, and τ be inertial with Artin preimage u. Put $F = R(s)$ and $T' = R(\tau)$. Then F and T' commute, and every polynomial vanishing at ιd_i annihilates T'. There exist characters θ_i and one functional l satisfying (44.9), with $l(y) \neq 0$, such that for every $b \geq 0$ one has*

$$\prod_{i=1}^{N} (F(T')^b - \iota(\theta_i(\varpi) d_i^b) 1) = 0.$$

Proof. Choose one irreducible constituent with nonzero projection of y. All the constituents in its global eigensystem have the same R by uniqueness. Finite Frobenius reciprocity gives the determinant-block type in (44.7), with $c = 1$, so the local parameter has zero monodromy and consists of characters. Its inertia characters are the specified distinct block characters, repeated according to their block sizes. We must arrange the exponents so as to have a functional nonzero on the particular vector, not just a character in the support of the representation.

The local Langlands classification with segments of length one embeds this representation into normalized induction from a list of characters α_i in a suitable order: take the dual of the Langlands quotient for the contragredient. Characters with different restrictions to units can be interchanged by invertible standard intertwiners, since the rank-two reducibility condition $\alpha_i/\alpha_j = |\cdot|^{\pm 1}$ is then excluded. Thus the list in the embedding can be ordered by the blocks of y, without changing the order used within each group of characters having that same restriction to units. This uses the ordinary rank-two intertwiner (or the Bernstein–Zelevinsky irreducibility criterion for two segments of length one). Write h for the induced function representing the projection of y. By Iwasawa decomposition it is determined by restriction to K_0, and, since the vector is K_1-fixed, on K_0 it is a finite function on $\mathrm{GL}_N(k)$. It transforms on the right by the indicated characters of the flag parabolic. Finite Bruhat decomposition writes

its possible nonzero double cosets as $B(k)wP(k)$. A torus element on the left, transported by w to the right, implies that $h(w) = 0$ unless w matches the unit characters on each index. As the characters of distinct blocks are distinct and the ordered α_i have the matching order, such a w must preserve every block, hence lies in $P(k)$. It follows that $h(1) \neq 0$. This applies also when the original vector has coefficients in several copies of the same automorphic constituent, by using its nonzero equivariant projection.

Take l to be evaluation at the identity of this embedding after e_1 and the equivariant projection. By the earlier computation it satisfies (44.9), with the θ_i of (44.10), and $l(y) \neq 0$. Its unit values are $\theta_i(u) = d_i$. The actual local Galois representation has finite inertia, and its Frobenius-semisimplified Weil parameter is $\bigoplus_i \theta_i$. Because the finite inertial characters factor through residue-field characters, the qth power fixes all of them. Conjugation by Frobenius consequently preserves every inertia eigenspace and acts identically on the action of every inertial element; thus F commutes with T' on the unsemisimplified representation. On these spaces T' is diagonalizable, with listed eigenvalues, and the characteristic polynomial of $F(T')^b$ is the product of the linear factors in the statement, by the Weil comparison and geometric reciprocity. This remains true if F has Jordan blocks: replacing F by its semisimple part does not change this characteristic polynomial, since the nilpotent part acts on the same inertia summands. Cayley–Hamilton finishes the proof for every b simultaneously. The chosen functional and characters were fixed before b was introduced. □

This is exactly the input called `galRep_of_parahoric_type` in `TypeFrobenius.lean`. Applying the polynomial factor comparison to this theorem gives the equality on the image of $T' - 1$ proved in Proposition 43.7, so the one-prime Hecke operator reduction no longer needs an independent assertion of the full Frobenius-polynomial law.

44.5 The primitive label and integral matrix span

The next obligation is not local Langlands: it is an integral consequence of the tame type which was used to control the Selmer rows. We give directly the arguments of `IsLabelledColumn.isRow` in `New/Selmer/LabelledColumns.lean` and `IsLatticeIn.span_eq_top_of_tame_pair` in `Lib/General/Groups/ResidualIrreducibility.lean`.

Lemma 44.6 (All Frobenius conjugates survive reduction)**.** *Let $q \equiv 1 \pmod 9$, $n = 3^{e+1}$, and ξ have multiplicative order $q^n - 1$. For any prime ℓ and $u \geq 0$, the numbers $(\xi^{\ell^u})^{q^i}$, $0 \leq i < n$, are distinct. If ξ is interpreted as a root of unity in $\overline{\mathbb{Q}}_\ell$, its n Frobenius conjugates are pairwise distinct modulo the maximal ideal.*

Proof. If the q-orbit of ξ^{ℓ^u} has length $< n$, its length divides $n/3$ because n is a power of three. Then $q^n - 1 \mid \ell^u(q^{n/3} - 1)$, whence

$$q^{2n/3} + q^{n/3} + 1 \mid \ell^u.$$

This integer is strictly larger than three and congruent to three modulo nine; it is divisible by three but not by nine, and has a prime divisor other than three. It cannot divide a power of a single prime. For the reduction assertion, the prime-to-ℓ part of a finite group of roots of unity injects into the residue field of any coefficient extension containing it. A root of unity reduces to one exactly when its order is a power of ℓ. Apply the first assertion to the ratio of any two distinct conjugates, taking u larger than its ℓ-primary exponent. This gives the exact unit-difference assertion in `norm_pow_sub_pow_eq_one` of the source. Scaling ξ by a base-field scalar does not change any of the ratios between its Frobenius conjugates. □

Lemma 44.7 (The metacyclic lattice argument)**.** *Suppose R is a continuous representation of a profinite group on an n-dimensional $\overline{\mathbb{Q}}_\ell$-space. Let γ, ϕ satisfy $R(\phi^{-1}\gamma\phi) = R(\gamma)^q$ and $R(\gamma)^{q^n} = R(\gamma)$. Assume that some eigenvalue ζ of $R(\gamma)$ has n pairwise residually distinct conjugates ζ^{q^i}. For every sufficiently large coefficient ring O and every stable integral form u of R over O, the O-span of $u(G)$ equals $M_n(O)$ and $u(\phi^{-n})$ is scalar.*

Proof. Enlarge O so it contains ζ. Write $A = u(\gamma)$, $B = u(\phi)$. Bars denote reductions. The characteristic polynomial of $\bar{A}$ has root $\bar{\zeta}$, so choose a nonzero eigenvector v. Since $AB = BA^q$, the vectors $v_i = \bar{B}^i v$, $0 \le i < n$, are eigenvectors with distinct eigenvalues $\bar{\zeta}^{q^i}$ and form a basis. The spectral idempotents E_i in this basis are polynomials in $\bar{A}$ of degree $< n$ by Lagrange interpolation. As $\bar{B}$ permutes the eigenlines cyclically, the products $\bar{B}^j E_i$ for $0 \le i, j < n$ give all matrix units up to nonzero scalars. They belong to the residue span of the group image (products of its elements are again group elements). Reduction of the integral span is consequently surjective onto $M_n(O/\mathfrak{m}_O)$, and Nakayama's lemma applies to this submodule of a finite free module. Finally $A^{q^n} = A$ implies B^n commutes with A; it also commutes with B. Their powers already span the complete matrix algebra modulo the maximal ideal, hence generate the full algebra over O by the same argument. The commutant of a full matrix algebra in itself consists of scalars. Thus B^n and B^{-n} are scalar. The proof works for every stable integral model, not only one obtained by a particular choice of basis. □

Stable integral models exist over every sufficiently large finite Galois coefficient field, by the compactness argument in Section 42.11. At the chosen level prime $g_i \neq \ell$, the tame inertia relation in Lemma 44.7 is the usual conjugation of a tame generator by arithmetic Frobenius. With the displayed relation $\phi^{-1}\gamma\phi = \gamma^q$ on the tame quotient, ϕ^{-1} is arithmetic, and ϕ^{-n} has cyclotomic value g_i^n. If one originally named the arithmetic lift ϕ instead, the scalar element is instead ϕ^n. Lemma 44.6 shows it has full integral matrix span whenever the primitive label of rank $n > 1$ occurs; for the rank-one row the assertion is automatic. There are two distinct level primes so one can do this for every coefficient prime.

Let $M = M_\omega = (N_0 d)^{s+6}$ be the exact tame order of ω in the data, and use the common scaling of the label by the inverse of the tame part at the chosen place. The scaling lies in $k^\times$ and has order dividing M in the untwisted row situation (twists are trivial at the chosen places). If the primitive norm is y of order $q-1$, the corresponding norm after scaling has order with p-part at least p^{s+6} for every prime $p \mid N_0 d$. Indeed the product of the conjugate roots is y times an element killed by M, and $M^2 \mid q-1$, so raising this product to M leaves a root whose order is at least $(q-1)/M$. In rank one the label $X-1$ is scaled by ω^{-j}, $1 \le j \le m$, and if its ath power is one then $M \mid ja$. Thus the same lower bound for order holds after allowing the factor $j \le m$. For any ℓ one may choose a prime divisor $p \mid N_0 d$ with $p \neq \ell$ (there are at least two). The size inequalities in the data give $(4n+2)m < 2^{s+6} \le p^{s+6}$: here $2^s = m^{22}$, $r \ll x^5$, and m grows as $\sqrt{x}$, so for the admissible large parameters the inequality follows with room; equivalently this is the instance of the logarithmic size comparison `D.nat_lt_of_hasLogBound` in `IsLabelledColumn.isRow`. In particular the p-part cannot be removed by an ℓ-power. Taking the power $1 + q + \cdots + q^{n-1}$ of a tame generator on a primitive orbit makes the common root value its norm, and in rank one use that generator itself. By exact label comparison the matrix then acts by one scalar y_0 and

$$|(y_0^{\ell^a})^j - 1|_\ell = 1 \qquad (a \ge 0,\ 1 \le j \le 4n+2).$$

This is the strong scalar-inertia condition in `IsRow`, not just the assertion that some Galois element acts by a scalar.

For every chosen-level place $v \nmid \ell$, put $c_\ell = \mathrm{crudeUnit}(\ell, D) = v_\ell(M_{\mathrm{gr}}) + 2^{-\ell}$, where $M_{\mathrm{gr}} = D.\mathrm{chosenGroupOrder}$. The labels exclude the root 1 for a row. At a tame generator

all roots in the polynomial (44.8) therefore differ from one, and each has order dividing $g_i^h - 1$ for some $1 \le h \le r$. If λ is such a nontrivial root and m_0 kills it, the factorization

$$\frac{\lambda^{m_0} - 1}{\lambda - 1} = 0$$

can equivalently be evaluated on the geometric sum and compared to its value m_0 at one: the polynomial $\sum_{k=0}^{m_0-1} X^k - m_0$ is divisible by $X - 1$, with integral coefficients. Hence $|m_0|_\ell \le |\lambda - 1|_\ell$. Now $g_i^h - 1 \mid M_{\mathrm{gr}}$, by the finite general linear group order factor for rank $h \le rt$ (the exact assertion `g_pow_sub_one_dvd_chosenGroupOrder` in `New/Selmer/AptTuples.lean`). Summing over the n characteristic roots gives

$$v_\ell \det(R(\gamma_v) - 1) \le nc_\ell, \qquad v_\ell \det(R((\gamma_v^c)^{-1}) - 1) \le nc_\ell. \tag{44.11}$$

Here the second inequality follows either from the same label at the conjugate place or directly from conjugate self-duality (the cyclotomic character is trivial on inertia away from ℓ). The representation is trivial on sufficiently high prime-to-g_i powers of inertia, because the tame labels factor through the finite prime-to-g_i quotient. One may use the exponent $g_i^{r!} - 1$ throughout. These are all the group-theoretic extra assertions needed to turn the indicated untwisted labelled column of rank n with its prescribed row jumps into a row.

44.6 The balanced-lattice estimate in full

We can now prove the result `IsRow.eventually_isBalanced` of `New/Selmer/RowBalanced.lean`. The following elementary local length calculations explain why even a highly ramified stable lattice can be used here.

Let $L = O^n$ be any stable integral lattice for a row R, with O the integer ring of a sufficiently large finite Galois extension of $\mathbb{Q}_\ell$. Write $e_O = v_O(\ell)$, v_O normalized to be one on a uniformizer, and $\mathrm{nlen}_O(X) = \mathrm{length}_O(X)/e_O$ for the normalized length of a finite module. Thus $v_O(z)/e_O = v_\ell(z)$, including for elements of an extension. Let S be the union of the places above ℓ and the two chosen level primes, and S_g the places in S not above ℓ. The data give $|S_g| \le 4f$, where $f = [E^+ : \mathbb{Q}]$; above each of the two completely split rational primes there are at most $2f$ places. The scalar and matrix-span assertions just proved hold for L; all its integral local comparisons are intrinsic to its rational representation, since base extension and a change of basis transport the period modules and the Hodge–Tate decompositions. At $v \mid \ell$ the field of potential crystallinity is a finite extension independent of the lattice. If a polynomial annihilates one of its inertial elements on the potential crystalline module, any polynomial vanishing at all its roots also annihilates that element, because this linear inertial action has finite image. This supplies the root-polynomial clause of `inClass_of_isLatticeIn`; the source's quantified condition over the whole class of columns is stronger than its specialization to this one column and therefore implies it. The remaining conditions of that definition (jumps and potential crystallinity, or actual crystallinity when the place is off the two level primes) are preserved by the coefficient extension and conjugation. When a collection of at most N such roots is killed individually by $X^{\ell^h-1} - 1$ with $1 \le h \le r$, each belongs to the unramified ℓ-adic extension of degree h by Teichmüller lifting, so its monic minimal polynomial over $\mathbb{Z}_\ell$ has degree at most r. The product has degree at most Nr and annihilates all of them; the common exponent $(\mathrm{lcm}_{1\le h\le r}(\ell^h - 1))$ works as well. This proves the separate assertion `isTameType_of_roots` at the start of `RowBalanced.lean`.

Lemma 44.8 (Dual relations and local finite lengths)**.** *Let $a \in O$, $a \mid \ell^M$, and put $B = O/aO$. If the crystalline cocycle lattice $Z = Z_f(L|_{G_{E_v}})$ at $v \mid \ell$ is defined by the period relations of the source, then the functions into B^n satisfying all its O-linear matrix evaluation relations have*

length at most $\operatorname{rank}_O Z \operatorname{length}_O B$. *At* $v \in S_g$, *every locally constant cocycle for* L *with values in* B^n *lies in a module whose length is at most*

$$n \operatorname{length}_O B + v_O \det(u(\gamma_v) - 1). \tag{44.12}$$

Here γ_v *may be the tame generator selected by the row; the bound does not require any restriction on the other local relations.*

Proof. For the first assertion, take the free module on all evaluation-coordinate symbols (g, i), and map it to $Z^* = \operatorname{Hom}_O(Z, O)$ by evaluation. Let A be the image. The module Z is finite free over the DVR, by the local finite-condition rank theorem, and A is therefore finite free of rank at most $\operatorname{rank} Z$. A function on the same symbols with values in B annihilating all relations vanishing on Z factors through A. The matrix-valued evaluation relations in the source include every scalar-coordinate relation, by putting the latter in a single matrix row. Hence the functions under consideration inject into $\operatorname{Hom}_O(A, B)$. Its length gives the bound, even when the image A is a proper non-saturated submodule of Z^*. This proves `length_obeying_relationsOn_le` without claiming that reduction of the cocycle lattice itself is saturated.

For the second assertion let P_v be the wild inertia at v. It is pro-residue-characteristic, whereas its linear inertia action and the order of the finite tame type factor through a prime-to-residue-characteristic quotient. Hence it acts trivially on L. Every continuous 1-cocycle restricted to it is an additive homomorphism to the finite ℓ-primary group B^n, hence zero. The affine action $g : x \mapsto u(g)x + c(g)$ consequently kills P_v and its restriction to tame inertia factors through a finite cyclic quotient, generated by the image of the tame generator. If evaluation of the cocycle at that generator is zero, it is zero on all inertia. Such a cocycle is determined on the decomposition group by its value $c(\phi)$ at an arithmetic Frobenius lift, because inertia and ϕ topologically generate the group modulo wild inertia. The two cocycle evaluations for $\tau\phi$ and $\phi(\phi^{-1}\tau\phi)$, for inertial τ, show

$$(u(\tau) - 1)c(\phi) = 0.$$

In particular this holds for the indicated generator. The kernel of evaluation at the tame generator thus injects into the kernel of $u(\gamma_v) - 1$ on B^n. The image of that evaluation has length at most $n \operatorname{length} B$. If the invariant factors of $u(\gamma_v) - 1$ have valuations d_i, Smith form bounds the length of the kernel modulo a by $\sum_i \min(d_i, v_O(a)) \le \sum_i d_i$. This is (44.12). The same proof gives the version of `length_localCocycles_le` for any chosen inertial element, by using its kernel in the last step; the tame generator is still used for the first evaluation. The local restrictions of global cocycles are a submodule of these local cocycles, so the inequality applies to the exact local modules in `IsBalanced`. □

The local rank input, proved with the classical Bloch–Kato formula in the preceding crystalline chapters, is

$$\operatorname{rank}_O Z_f(L|_{G_{E_v}}) = n + \sum_{\sigma \mid v} \#\{u \in J(\sigma) : u < 0\}. \tag{44.13}$$

Here $\sigma \mid v$ refers to the embeddings under ι, and the formula counts all of them, not just the selected half of a CM type. This is `finrank_crysCocycles_eq_add_neg_jumps` in `New/Selmer/CrystallineRank.lean`; the extra n accounts for cocycles rather than cohomology classes. Since the row type has at most n negative jumps at a conjugate pair together, summing its right side over all $v \mid \ell$ is at most $n(|S \setminus S_g| + f)$.

Let g be the scalar element of a row with $C = \mathrm{cyc}_\ell(g) = g_i^n$ and $u(g) = \alpha 1$. Conjugate self-duality in the source is

$$\operatorname{tr} R(h^c) = \mathrm{cyc}_\ell(h)^{1-t} \operatorname{tr} R(h^{-1}). \tag{44.14}$$

The identity implies $R^c \simeq R^\vee \otimes \mathrm{cyc}_\ell^{1-t}$, by semisimplicity and equality of characters. Thus $u((g^c)^{-1}) = \alpha' 1$ with $\alpha' = C^{t-1}\alpha$ in the same lattice. This can also be seen purely integrally by applying (44.14) to hg^{-1} and using nondegeneracy of the trace form on the integral matrix span to conclude equality of the two scalar matrices. In either proof α is a unit because the matrix is invertible over O. Consequently the normalized mirror loss in `IsBalanced` is exactly

$$\frac{n}{e_O} v_O(\alpha - \mathrm{cyc}_\ell(g^c)\alpha') = n v_\ell(1 - g_i^{nt}) \le n c_\ell,$$

using $1 \le nt \le rt$ and the finite group order divisibility from above. The sign of the exponent is immaterial after multiplying by a unit; the formula as written uses the inverse-cyclotomic convention of `IsConjSelfDual`.

Theorem 44.9 (The exact balanced row budget)**.** *For a row R satisfying `IsRow`, and for a chosen level place $w \nmid \ell$, let $c_\ell = \mathrm{crudeUnit}(\ell, D)$. For every $M \ge 0$ and every sufficiently large O, there is a stable model u and γ, β satisfying `IsBalanced` for its actual crystalline relations and*

$$\frac{1}{e_O}\left(\beta + \sum_{b \in \{1,c\}} v_O \det(u(\mathrm{idOrConjInv}_b(\gamma_w)) - 1)\right) \le 10 f n c_\ell. \tag{44.15}$$

Proof. Take O sufficiently large for the stable model and matrix-span assertions. For scalar inertia choose the element γ just above (44.11): its action is the scalar s with $s-1$ and $s+1$ units. Since it is at $v \nmid \ell$, $\mathrm{cyc}_\ell(\gamma) = 1$ and (44.14) gives the same scalar for $(\gamma^c)^{-1}$. Its residue has order $> 4n+2$ by the unit-difference assertion, so the residue field has cardinality $> 4n+2$. Continuity of u gives local constancy modulo ℓ^M, unramifiedness outside S is part of the row local conditions, and the matrix-span clause was proved above. Every coboundary $h \mapsto u(h)x - x$ obeys the crystalline relations modulo a: lift x from B^n to O^n, use the integral crystalline coboundary, and reduce the relations. This verifies the earlier clauses in the definition of `IsBalanced` without appealing to a special choice of stable model.

For each $v \in S_g$ fix a row tame generator at a place over v, and let $d_v = v_O \det(u(\gamma_v) - 1)$. The two local length assertions, (44.13) and the mirror calculation show that one can take

$$\beta = \sum_{v \in S_g} d_v + n v_O(1 - g_i^{nt}).$$

Indeed for every $a \mid \ell^M$ the sum of the least local lengths plus the scalar difference in `IsBalanced.local_length_le` is at most

$$(|S| + f) n \operatorname{length}_O(O/aO) + \beta.$$

The infimum over the places extending v is bounded by the chosen value; the inequality holds uniformly in a, not only for $a = \ell^M$. By (44.11) each $d_v/e_O \le n c_\ell$; the normalized mirror loss is $\le n c_\ell$. Adding the two determinant losses at w in (44.15) gives normalized sum at most

$$(4f + 1 + 2) n c_\ell \le 7 f n c_\ell \le 10 f n c_\ell,$$

as $f \ge 1$. In particular the estimate holds with the source's stated constant 10; the small improvement in this isolated bound is not used elsewhere. This proves precisely the needed balanced-lattice theorem. □

44.7 Row traces for all twists and the rigidity reduction

We can isolate the remaining geometric dependence in `New/Selmer/RowsExist.lean`. Its row construction first requires a nonzero component of the actual good form eigensystem with a tuple of good labels. This is the elementary projection step using orthogonal idempotents: the good projector is a sum of tuple idempotents, so if its value on a vector in the joint Hecke eigenspace is nonzero, at least one summand is nonzero there. The tuple projections commute with the off-level Hecke algebra. Apply the algebraic-form realization to this vector and Proposition 44.1; Theorem 44.4 gives a labelled column with the stipulated local conditions. When normalized by the algebraic twisting avatar of the pair, its jumps become those of the row type, using the formula `isRowType_auxPair` of `New/Rows/Separation.lean`. This last formula is the multiset of $\lfloor t/2 \rfloor - \mathrm{sgn}(\sigma)(\varsigma_q + w_j)$ for $0 \le q < r$; the required negative-jump count follows immediately because the signs at σ and $\bar{\sigma}$ are opposite and $t > 0$. Its other window-count bounds are as in the row-parameter calculations of the earlier chapters. Lemma 44.7 and the scalar-inertia calculations prove the full row conditions for this labelled column.

The trace assertion is then a coefficient identity, but its geometric premise must not be forgotten. Use the unramified theta Hecke-polynomial identity in the source normalization, and divide by the actual normalizing character `chiOneInv`. For a normalized theta lift on the auxiliary pair of rank difference t, this gives, at the places of prime norm where the lift and its twist are spherical, the polynomial factorization with the row factor twisted by the scalar ϑ^{-1} and the string $\prod_{e=0}^{t-1}(X - \mathrm{cycInv}(\mathrm{Frob}_w)^e)$. Taking the coefficient of degree $N-1$ gives the sum of traces of the two factors. The avatar identities hold at almost every such Frobenius. As the rank-r row representation is fixed, taking the limit in the localized Frobenius filter near each $h \in G_E$ yields the single identity

$$\mathrm{Tr}(R_{\mathrm{lift}}^{\mathrm{norm}}(h)) = \vartheta_\ell^{-1}(h)\,\mathrm{Tr}(R_{\mathrm{row}}(h)) + \sum_{e=0}^{t-1} \mathrm{cycInv}(h)^e.$$

Thus no new simultaneous choice of Galois representations or lattices for an infinite family of twists is necessary. The row depends only on the untwisted Hecke system and the fixed normalization; every twist satisfies this identity individually once its actual theta-lift Hecke identity is proved. This is the content of the quantified conclusion of `exists_row_for_all_twists`. The remaining premise is the genuine theta-transfer identity and actual nonzero algebraic realization of the indicated forms, not a missing theorem on the existence of attached representations.

For the dimension-transfer argument one can likewise remove its independent Galois assumption. Apply Theorem 44.4 to each of the two auxiliary injected forms in `Data.occurringLabels_subsingleton` of `New/GaloisRep/LabelUniqueness.lean`. Both have the same off-level Hecke system and therefore the same R by Proposition 44.1; take the coefficient prime $\ell = 2$, which is different from the chosen odd rational primes. For a single chosen tame exponent $a = 1$, the annihilation and determinant conclusions of Theorem 44.4 give exactly the common test representation and its determinant. In fact that theorem gives equality of the characteristic polynomials in this situation, so all the roots with order greater than M must agree on both sides. The labels are scaled by the same nonzero scalar from the common determinant twist; their primitive roots have order $q^r - 1 > M$, since $M^2 \mid q - 1$. The additional determinant blocks have root order dividing M. Hence the two scaled primitive orbit polynomials agree and, scaling inversely, so do the two original labels. This is the finite rigidity argument of Lemma 18.5, now with its arithmetic local and global Galois premises discharged. The necessary actual auxiliary injection and its parahoric/Hecke-equivariance

remain geometric inputs, as stated in the dimension-transfer chapter, independently of Galois attachment.

For the comparison with the period-ring convention, the source's rational crystalline ring is obtained from its maximal integral ring rather than by naming the divided-power integral lattice. The ordinary divided powers $\xi^j/j!$ lie in that maximal ring, since $(\xi/p)^j(p^j/j!)$ lies there. The inclusion extends by completion and is compatible with the embedding into B_{dR} and with the cyclotomic period. Thus an invariant basis in the usual rational crystalline period ring extends to the ring used here; the fixed-scalar comparison in Chapter 52 gives the same finite unramified field of coefficients. This is the normalization bridge for the ordinary potentially crystalline statements quoted above. It does not identify the two integral lattices.

44.8 Obligation ledger for this replacement route

- **S13 — proved from explicit classical references for the actual Weil data.** Proposition 44.1 replaces the final weak spherical attachment interface in all ranks, with exact polynomial annihilation and Hodge jumps. It does not require executing or re-proving the source's geometric seed, weight approximation, or weak base-change constructions. The replacement uses the full definite-group base change and cuspidal attachment literature explicitly identified above; these are ordinary published theorem inputs, not arbitrary declarations carrying hidden singular-weight hypotheses. The parity and rank normalization and the local period operator have been checked. The broader intermediate `SeedRepresentation` theorem for fields without the actual splitting data is not claimed separately.

- **S14 — proved from explicit classical references and local arguments.** Theorem 44.4 gives finite-type and jumps at every indicated split labelled place, the exact tame-root killing polynomial on the source period module, and determinant off ℓ. The sorted Iwahori comparison and Proposition 44.5 also give the polynomial for the simultaneous Frobenius–inertia powers, including the functional nonzero on the given vector. The passage from Weil–Deligne equality or its weaker monodromy inequality to the needed genuine killing statements is part of the proof. No regularity of a general local Langlands parameter has been silently imposed.

- **S02 — written proof.** Lemmas 44.6–44.8 and Theorem 44.9 prove the balanced lattice interface under its exact row hypothesis, with the prescribed scalar and matrix-span properties and budget. The row hypothesis is the subject of the preceding construction; it is not a new assumption in the balanced theorem.

- **S01 — partial/conditional only.** The Galois attachment, tame residual orbit and scalar bounds, and the simultaneous trace deduction from the unramified normalized theta polynomial have been supplied. The actual good-vector algebraic realization, normalized theta Hecke transfer, and the precise row-type window verification remain the dependencies already identified in the geometric chapters where not proved there.

- **B02 — conditional on the actual injective lift and its exact local intertwining.** The arithmetic rigidity deduction is now proved under those geometric premises, as explained above. It would still be incorrect to infer the existence of that auxiliary injection from the Galois comparison alone.

Chapter 45

The first-order Ribet polynomial over the integral coefficient algebra

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 47, 48 and the final audit in Chapter 57.

We prove the abstract first-order theorem in `New/Selmer/RibetFirstOrder.lean`; the assertion is `ribet_first_order`. This closes a different obligation from existence of the geometric family and local Galois labels. It says that once the indicated family of matrices, trace ideal and cocycle costs are supplied, the required polynomial really belongs to the integral coefficient algebra, has the required first coefficient, and obeys the stated valuation bound. All products are over finite sets; the group itself need not be finite. No semisimplicity for the coefficient algebra, division by the dimension of a matrix, or inversion of a residue characteristic is used.

45.1 Fitting ideals and the cost of imposing linear forms

Fix a DVR O, normalized valuation v, and nonzero $p \in O$. Write $D = O/pO$; this letter p denotes an element, not necessarily a prime element. For any finite free O-module P and submodule $L \supset pP$, the annihilator correspondence in the pairing

$$P \times \mathrm{Hom}_O(P, D) \longrightarrow D$$

identifies L with the annihilator of its annihilator. Indeed Smith normal form reduces this assertion to the equalities

$$\mathrm{Ann}_D(\pi^i O) = (\pi^{v(p)-i})D, \qquad \mathrm{Ann}_O((\pi^{v(p)-i})D) = (\pi^i) \quad (0 \le i \le v(p)).$$

Here π is a uniformizer. In particular, the length of the annihilator is $\mathrm{length}_O(P/L)$. This works without choosing a perfect ordinary trace pairing.

Let $\Lambda = O[X_j : j \in J]$, $T = \Lambda^d$, $N \subset T$ a Λ-submodule, $\Phi : P \to T$ an O-linear map, and $z : G \to P$ any map. Assume $p\Phi(P) \subset N$. Put

$$Z_N = \{\, g \mapsto \lambda(z(g)) : \lambda \in \mathrm{Hom}_O(P, D),\ \lambda(\Phi^{-1}N) = 0 \,\}.$$

Thus Z_N is finite length, and corresponds to `formsMod`. For submodules Z, V of the module of functions $G \to D$ denote by $\lambda(Z; V)$ the length of $Z/(Z \cap V)$. The zeroth Fitting ideal $\mathrm{Fitt}(N)$ means the ideal of Λ generated by determinants of $d \times d$ matrices with rows in N.

If $rw \in N$ then

$$r\ \mathrm{Fitt}(N + \Lambda w) \subset \mathrm{Fitt}(N). \tag{45.1}$$

Expand a row determinant by writing each row as its N part plus a multiple of w. All terms involving two or more copies of w are zero. The terms with no w or one w are in $\mathrm{Fitt}(N)$ after multiplication by r. Also, for an ideal $I \subset \Lambda$,

$$\mathrm{Fitt}(N + IT) \subset \mathrm{Fitt}(N) + I. \tag{45.2}$$

Indeed reduce each row modulo I before taking its determinant.

Lemma 45.1 (Exact charge for enlarging a relation module)**.** *Given an O-submodule $V \subset D^G$, there are $K \supset N$ and $a \in O$ such that*

$$v(a) \le \lambda(Z_N; V), \qquad a\ \mathrm{Fitt}(K) \subset \mathrm{Fitt}(N),$$

and $\Phi(w) \in K$ whenever every $\lambda : P \to D$ with $\lambda \circ z \in V$ satisfies $\lambda(w) = 0$.

Proof. Start with $K = N$. If the asserted last property fails, choose such a w with $\Phi(w) \notin K$. Choose the smallest $k \ge 1$ such that $\pi^k \Phi(w) \in K$, which exists because $p\Phi(w) \in N$. Replace K by $K' = K + \Lambda \pi^{k-1} \Phi(w)$. By (45.1) this costs one factor of π in the Fitting ideal. Moreover the image of $Z_{K'}$ in D^G/V is strictly smaller than the image of Z_K. Otherwise every λ annihilating $\Phi^{-1}K$ could be matched by a λ' annihilating $\Phi^{-1}K'$ such that $(\lambda - \lambda') \circ z \in V$. The defining property of w implies

$$\lambda(\pi^{k-1} w) = \lambda'(\pi^{k-1} w) = 0.$$

By annihilator separation this would imply $\pi^{k-1}\Phi(w) \in K$, a contradiction. Thus after at most $\lambda(Z_N; V)$ operations there can be no failure of the required last property. Multiply the factors of π used in the process. All valuations are exact integers; this proves the lemma. □

This is the statement `fittingIdeal.exists_cost_le_relLength` in `Lib/General/Algebra/FittingDuality.lean`. The proof also supplies its duality premise `mem_of_forall_form_eq_zero`. The distinction between annihilating a module of forms and imposing an equation on a single choice of coordinates is essential to the cost bound.

45.2 The triangular block theorem

Let J be linearly ordered, $I \subset J$, and for each j let $\tau_{j,+}, \tau_{j,-}, \phi_{j,+}, \phi_{j,-} \in G$. Let $T = \Lambda^s$ where s here denotes a finite set of detecting points of G; write $d_s = |s|$. Use the notation of Lemma 45.1. Suppose

$$\begin{aligned} p\Phi(P) \subset N, \quad \Phi(z(g)) - e_g \in N \quad (g \in s), \\ \Phi(z(\tau_{j,\pm})) - X_j \Phi(z(\phi_{j,\pm})) \in N, \\ X_i T \subset N \quad (i \notin I,\ i < j \in I). \end{aligned} \tag{45.3}$$

Here e_g is the standard row vector in coordinate g. The exact condition `HasTriangleCost` on $Z = Z_N$ is as follows. There exist a subset $L \subset G$, an element $C \in O$ and a bound b with two properties:

1. If $c \in Z$ is zero on L and $c(\tau_{i,+}) = c(\tau_{i,-})$ for $i < j$, and in addition $c(\phi_{i,+}) = c(\phi_{i,-})$ for $i \in I$, $i < j$, then
$$C(c(\tau_{j,+}) - c(\tau_{j,-})) = 0 \qquad (j \in I).$$

2. For every submodule $Z' \subset Z$ zero on L and on all $\tau_{j,\pm}$, with equal ϕ-pairs for $j \in I$, one has
$$\lambda(Z; \{c : c|_L = 0\}) + \lambda(Z'; D^G[C]) \le b.$$

Proposition 45.2 (Triangular Fitting polynomial)**.** *Under these assumptions, a polynomial* $\Gamma \in \Lambda$ *satisfies*
$$\Gamma \prod_{j \in I} X_j \in \mathrm{Fitt}(N), \qquad v(\Gamma(0)) \le b + (|I| + d_s)v(C).$$

Proof. It suffices to consider finite right-hand side. Apply Lemma 45.1 with $V = \{c : c|_L = 0\}$ to obtain K and multiplier a. Then $\Phi(z(g)) \in K$ for $g \in L$. Every form for any relation module containing K lies in Z and is zero on L.

For each $j \in I$ in increasing order, put $w_j = \Phi(z(\phi_{j,+}) - z(\phi_{j,-}))$, and add w_j to the current module. Write $v_j = \Phi(z(\tau_{j,+}) - z(\tau_{j,-}))$. At that step v_i lies in the current module for all $i < j$: either w_i has been added, or $X_i T \subset N$. Earlier ϕ-pairs for $i \in I$ agree on every form for the current module. By the first triangle hypothesis all such forms vanish on Cv_j. Annihilator separation, applied to P and the current relation module, gives Cv_j in that module. Since $v_j - X_j w_j \in N$, the new generator w_j is added at Fitting cost CX_j by (45.1). If T_1 denotes the result, then
$$aC^{|I|} \prod_{j \in I} X_j \ \mathrm{Fitt}(T_1) \subset \mathrm{Fitt}(N).$$

Put $T_2 = T_1 + (X_j : j \in J)T$. The forms Z_{T_2} satisfy the second triangle hypothesis: all τ vanish by (45.3), and all stipulated ϕ-pairs agree. Apply Lemma 45.1 to T_2 and $V = D^G[C]$. It gives $K_2 \supset T_2$ and a_2 with $v(a) + v(a_2) \le b$. For any $g \in s$, every form with function in $D^G[C]$ annihilates $Cz(g)$, so $Ce_g \in K_2$ by (45.3). Hence
$$a_2 C^{d_s} \in \mathrm{Fitt}(T_2) \subset \mathrm{Fitt}(T_1) + (X_j : j \in J).$$

Choose $d_1 \in \mathrm{Fitt}(T_1)$ with $d_1(0) = a_2 C^{d_s}$. Take $\Gamma = aC^{|I|} d_1$. The claimed membership and valuation follow, without assuming that $\mathrm{Fitt}(N)$ is principal. □

This proves `exists_mem_fittingIdeal_block` with the same coefficient loss and the same ordered variables. Notice that the proof controls the coefficient of the product of *all* variables in a block, rather than a determinant specialized at just one point.

45.3 Selection of a distinguished block determinant

Let A be an O-algebra, not necessarily commutative, $\psi : A \to M_r(O)$ surjective, and $N \subset A^m$ a left submodule. Suppose
$$Z = \{(w_1, \dots, w_m) \in (O/p_0O)^{rm} : \sum_j \psi(n_j) w_j = 0 \text{ for every } n \in N\}$$
has length at most $b_0 < v(p_0)$, where $p_0 \neq 0$. There exists $B_0 \in M_m(A)$, all rows in N, with
$$v(\det \mathrm{Flat}(\psi B_0)) \le b_0. \tag{45.4}$$

To prove this, take the O-module L in O^{rm} generated by all individual scalar rows of the block rows $(\psi(n_j))_j$. By multiplying on the left by preimages of matrix units, one can insert a scalar row into any chosen matrix row, putting zero in all other rows; adding these gives an arbitrary specified r-tuple of rows of L as one block row in $\psi(N)$. Smith normal form of

L gives Z length $\sum_i \min(v(p_0), d_i)$ with the missing diagonal entries assigned $d_i = \infty$. Since this is less than $v(p_0)$, no diagonal entry can be missing or have valuation $\geq v(p_0)$. Thus L is full rank and its exact colength is at most b_0. Choose an O-basis of L and assemble it in consecutive groups of r rows by the matrix-unit argument. This proves (45.4) and in particular `exists_rows_addVal_det_le` in `Lib/General/Algebra/BlockFittingBound.lean`. It is not necessary to find an idempotent over the entire family to isolate this distinguished block.

45.4 An integral two-adjugate calculation

Let a finite family of group-algebra representations, of a common dimension n, act over a commutative ring L' containing the given coefficient ring. We regard the family as one representation $u : A \to M_n(B)$, where B is the product of the ambient rings, $A = \Lambda[G]$, and let $S \subset B$ contain the scalar polynomial coefficients and the characteristic-polynomial coefficients of every $u(g)$. Fix an S-submodule $\mathcal{J} \subset B$. Write

$$T_{\mathcal{J}} = \{a \in A : \operatorname{tr}(u(a)u(g)) \in \mathcal{J} \text{ for all } g \in G\}.$$

By linearity in g and cyclicity of trace this is a two-sided ideal; its extension by S is still in the trace annihilator with test elements of $S[G]$.

For an A-linear map of left modules $V : A^m \to A$ and row $w \in A^m$, put $b = Vw$, $\beta = \det u(b)$. The Amitsur result of Corollary 8.6 shows that $\beta \in S$, and that all coefficients of the characteristic polynomial of $\operatorname{Flat}(u(M))$ for $M \in M_m(A)$ are in S. Hence each of the two adjugates is the image of an ordinary polynomial with coefficients in S applied to the corresponding group-algebra matrix: this follows by the universal adjugate polynomial, with no determinant inverted. Suppose the rows of M satisfy

$$V(M_i) = t_i + a_i b, \qquad t_i \in T_{\mathcal{J}}, \quad a_i \in \mathfrak{a}$$

for a left ideal $\mathfrak{a} \subset A$. Let H over $S[G]$ be the polynomial adjugate of M after flattening, and let $D_M = \det \operatorname{Flat}(u(M))$. Using the column of V-coefficients and multiplying the displayed relations by the block row wH after applying u, we get

$$u((D_M - wHa)b) = u(wHt).$$

Set $a' = D_M - wHa$ in $S[G]$ (coefficients act centrally). For $h \in A$ the function $f = \operatorname{tr}(u(a'h))$ lies in S, again by integral Amitsur. Multiplying the last relation by the polynomial adjugate of $u(b)$ and using the trace-annihilator condition gives

$$f\beta \in \mathcal{J}. \tag{45.5}$$

At a member x of the family for which $u_x(a_i h) = 0$ for all i, we have

$$f_x = D_{M,x} \operatorname{tr}(u_x(h)). \tag{45.6}$$

This proves the full ideal-membership and specialization assertion `exists_ribet_element_cramer`, not just an equality after division by β. The proof takes place in the image in $M_n(B)$; there is no assertion that the individual matrix entries lie in S.

45.5 The abstract first-order theorem

We now specify the exact source situation. Let Y be a finite family of representations $u_x : G \to GL_{r+t}(O)$, and $\mathfrak{X} \subset Y$ a distinguished subset with distinguished point x_0. On $x \in \mathfrak{X}$, after a common reindexing, suppose

$$u_x(g) = \rho_x(g) \oplus \operatorname{diag}(\xi_e(g))_{e=0}^{t-1}, \qquad \xi_0 = 1, \quad t \geq 1.$$

The characters ξ_e are independent of x. Assume that $\rho_{x_0}(G)$ spans $M_r(O)$ as an O-module. For each distinguished x fix $\gamma_x \in G$ and $y_x \in O$ with

$$\rho_x(\gamma_x) = y_x 1, \quad y_x - 1 \in O^\times, \qquad \xi_e(\gamma_x) = 1 \quad (0 \le e < t).$$

Let $p, p_0 \neq 0$ in O, $a \in G$, $C \in O$ satisfying

$$\prod_{k \neq e} (\xi_e(a) - \xi_k(a)) \mid C \quad (0 \le e < t). \tag{45.7}$$

The normalizer is the finitely supported function on G

$$R = [\gamma_{x_0} a] - y_{x_0}[a] + p_0 C \sum_g n_g [g],$$

where n_g are nonnegative integers with finite support. Put

$$\beta_x = \det\Big(\sum_g R_g (u_x(g) - 1)\Big).$$

As in the Lean definition `IsTraceIdeal`, let S be an O-subalgebra of $\prod_{x \in Y} L[X_j : j \in J]$ containing the variables, and all family-valued characteristic coefficients of $u(g)$, where L is a commutative O-algebra. Let $\mathcal{J}$ be an S-submodule of this ambient ring. For a finite linearly ordered J and its two copies of each $\tau_{j,\pm}, \phi_{j,\pm} \in G$, assume for all $g \in G$

$$\begin{aligned} p \operatorname{tr} u(g) &\in \mathcal{J}, \\ \operatorname{tr}\left((u(\tau_{j,\pm}) - 1)u(g)\right) - X_j \operatorname{tr}\left((u(\phi_{j,\pm}) - 1)\varepsilon_j(u(\phi_{j,\pm}))u(g)\right) &\in \mathcal{J}, \\ p \mid \varepsilon_j(\xi_e(\phi_{j,\pm})) - 1, & \end{aligned} \tag{45.8}$$

where $\varepsilon_j \in O[T]$ and the last line holds for every e.

Let $\delta_v \in G$, $v \in W$, be a finite collection and $o_v(T) \in O[T]$, with

$$\xi_e(\delta_v) = 1, \qquad (u_x(\delta_v) - 1) o_v(u_x(\delta_v)) = 0 \quad (x \in Y). \tag{45.9}$$

For a representation ρ and nonzero $q \in O$, denote by $Z(\rho, q)$ the cocycles $c : G \to (O/qO)^{\dim \rho}$ which obey every relation among the vectors $u(g) - 1$ in the *whole family*; more explicitly $\sum_g \theta_g c(g) = 0$ whenever $\sum_g \theta_g (u_x(g) - 1) = 0$ for every $x \in Y$. This is `obeyingCocycles`. For scalar ξ this has scalar values. Assume

$$\operatorname{length}_O\{c \in Z(\rho_{x_0}, p_0) : c(\tau_{j,\pm}) = 0,\ c(\gamma_{x_0}) = 0\} \le b_0 < v(p_0). \tag{45.10}$$

Let $\mathrm{ty} : J \to \{0, \dots, t-1\}$ be nondecreasing. Fix a finite detecting set $s \subset G$ and bounds b_e. Our last assumption is that for any module $Z \subset Z(\xi_e, p)$ with the following two properties:

1. $c(\delta_v) = 0$ for every $c \in Z$;
2. for each coboundary $c(g) = (\xi_e(g) - 1)m$ in Z, one has $(\xi_e(a) - 1)c(g) = 0$ for every g,

the exact triangle-cost condition above holds with $I_e = \{j : \mathrm{ty}(j) = e\}$, C, b_e, and $c \in Z$ zero on s implies $c = 0$. This is the source hypothesis `hb`. It is a condition on every such module Z, not just on the whole space of unramified cocycles.

Theorem 45.3 (First-order integral Ribet polynomial)**.** *Under the stated hypotheses there are $f \in S$ and $c \in O$ with $\beta \in S$, $\beta f \in \mathcal{J}$, and $\beta_x = 0$ on $\mathfrak{X}$. The coefficient of every monomial properly dividing $X_* = \prod_{j \in J} X_j$ in f_x is zero on $\mathfrak{X}$, and the coefficient of X_* in f_{x_0} is the image of c. Moreover*

$$v(c) \le b_0 + \sum_e b_e + t\left(\sum_{v \in W} v(o_v(1)) + 2|s| v(C)\right) + (|J| + 1) v(C). \tag{45.11}$$

Proof. The interesting case is finite right-hand side. If the right side is infinite, the only required inequality is trivial with $c = f = 0$; the membership $\beta \in S$ and its vanishing on $\mathfrak{X}$ still follow directly from Amitsur and $\xi_0 = 1$. In the finite case C and $o_v(1)$ are nonzero whenever their valuations are charged. We give the construction in six steps.

Step 1: The left relation module. Let $\Lambda = O[X_j : j \in J]$ and $A = \Lambda[G]$; all representations and characters extend to this algebra. Choose a finite $s_0 \subset G$ such that the elements $u(g) - 1$ for $g \in s_0$ span all these elements in $\prod_Y M_{r+t}(O)$ over O. This is possible by noetherianity. Include the extra, disjointly labelled coordinates W (for δ_v) and s (for detecting points) and put $m = |W| + |s_0| + |s|$. Write $x_i \in G$ for their group elements. Choose $z(g) \in O^m$ with

$$\sum_i z_i(g)(u_x(x_i) - 1) = u_x(g) - 1 \quad (x \in Y).$$

For a constant row q we identify $q \in O^m$ with its row in A^m. Define

$$V(q) = \sum_i q_i([x_i] - 1) \quad (q \in A^m), \qquad \omega = \sum_g R_g z(g) \in O^m.$$

Let $\eta_e : A \to \Lambda$ be evaluation through ξ_e, $\mathfrak{a} = \ker \eta_0$, and put

$$N = V^{-1}(T_{\mathcal{J}} + \mathfrak{a}V(\omega)).$$

All rows whose V-image evaluates to zero at every u_x are in N. In particular the following rows belong to N:

$$\begin{aligned} z(gg') - z(g) - [g]z(g'), \qquad & z(x_i) - e_i, \\ \sum_g \theta_g z(g) \quad \text{if } \sum_g \theta_g(u_x(g) - 1) = 0 \quad & (x \in Y), \\ z(\tau_{j,\pm}) - X_j \varepsilon_j([\phi_{j,\pm}])z(\phi_{j,\pm}), \qquad & ([g] - 1)\omega, \qquad pA^m. \end{aligned} \tag{45.12}$$

The first line follows from $u(gg') - 1 = (u(g) - 1) + u(g)(u(g') - 1)$. The second is the choice of z. The third uses (45.8), the fact that $\eta_0([g] - 1) = 0$, and the trace-ideal generators $p \operatorname{tr} u(g)$. Also

$$e_v\, o_v([\delta_v]) \in N \quad (v \in W) \tag{45.13}$$

by (45.9) and the definition of V (left coefficient in the v slot). In particular these are genuine relations in the original family, not merely relations on its distinguished subfamily.

Step 2: The ρ-block. Apply $\psi = \rho_{x_0}$ at $X = 0$ to coefficients in A. It is surjective by the spanning assumption. Let $w \in (O/p_0O)^{rm}$ be killed by all rows of N under ψ. Set $c(g) = \sum_i z_i(g)w_i$. From (45.12) this is a cocycle obeying the relations among the family, vanishes on all $\tau_{j,\pm}$, and satisfies $c(x_i) = w_i$. In the action on this quotient, the normalizer row ω has value

$$c(\gamma_{x_0} a) - y_{x_0} c(a) + p_0 C \sum_g n_g c(g) = c(\gamma_{x_0}).$$

The row $([\gamma_{x_0}] - 1)\omega$ and the unit $y_{x_0} - 1$ therefore give $c(\gamma_{x_0}) = 0$. The map $w \mapsto c$ is injective, and (45.10) bounds the length of the joint row annihilator by b_0. Section 45.3 supplies B_0 with rows in N and

$$v(D_0(x_0, 0)) \le b_0, \qquad D_0(x, X) = \det \operatorname{Flat}(\rho_x B_0) \quad (x \in \mathfrak{X}).$$

Step 3: The scalar forms and simultaneous normalized rows. For each e let $E_e : A^m \to \Lambda^{s_0 \amalg s}$ evaluate through η_e and drop the W slots. Suppose an O-linear map $\Phi = \Upsilon E_e|_{O^m}$ is used, with Υ any Λ-linear map, and the relation module K contains $\Upsilon E_e(N)$. For a form $\lambda : O^m \to O/pO$ annihilating $\Phi^{-1}K$, the function $c = \lambda \circ z$ is in $Z(\xi_e, p)$ and vanishes at all δ_v by (45.12); moreover all such functions form a module. If $c(g) = (\xi_e(g) - 1)m_1$ then, writing $u_e = \xi_e(a) - 1$ and using $\xi_e(\gamma_{x_0}) = 1$, we have

$$\lambda(\omega) = (1 - y_{x_0})u_e m_1 + p_0 C \sum_g n_g(\xi_e(g) - 1)m_1.$$

When $u_e = 0$ the required annihilation is automatic. Otherwise $e \neq 0$ and $u_e \mid C$ by (45.7). Since $v(p_0) > 0$, the scalar in brackets after factoring out $u_e m_1$ is a unit:

$$1 - y_{x_0} + p_0(C/u_e) \sum_g n_g(\xi_e(g) - 1).$$

The row $([g] - 1)\omega$ in (45.12) then gives $u_e c(g) = 0$. Thus the hypotheses of hb apply to the module of these forms.

In particular, take Υ to be projection to s_0. Any such form is zero on the detecting set s by the coordinate relations, so $c = 0$ by hb; it follows that the form kills every standard basis vector, because $c(x_i) = \lambda(e_i)$. By annihilator separation and pO^m in the preimage, the projection of $E_e(N)$ to Λ^{s_0} contains every constant basis vector. It is a Λ-submodule, hence is the full module. Select rows $m_{e,i} \in N$ with those projected character evaluations equal to e_i. One obtains a collection $\nu_i \in N$ $(i \in s_0)$ for which

$$(\eta_e(\nu_i))|_{s_0} = e_i \qquad \text{for all } e. \tag{45.14}$$

Start with zero and process the characters successively. A correction using only its evaluation at a single character would not retain the earlier identities. Instead update the whole collection in the group algebra by the row formula

$$\nu^{\text{new}} = \nu + (I - \nu|_{s_0})m_e.$$

Here $\nu|_{s_0}$ is the matrix of *group-algebra* coefficients in those slots, not its evaluation at e. Left multiplication of rows by these entries preserves N; at earlier characters the multiplying matrix evaluates to zero, and at e the new projection evaluates to $\eta_e(\nu|_{s_0}) + (I - \eta_e(\nu|_{s_0}))I = I$. This gives (45.14) without dividing by the character differences.

Step 4: Character separation in the square-free quotient. For each e define a map

$$A_e(b_0, b_s) = b_s - b_0\,(\eta_e(\nu_i)|_s)_{i \in s_0}.$$

It subtracts the row combination having the prescribed s_0 coordinates. Set $N_e = A_e E_e(N) \subset \Lambda^s$. Under $\Phi = A_e E_e|_{O^m}$, the coordinate relations in (45.12) give $\Phi(z(g)) - e_g \in N_e$ for $g \in s$. The ties become

$$\Phi(z(\tau_{j,\pm})) - X_j \Phi(z(\phi_{j,\pm})) \in N_e.$$

Indeed $\eta_e(\varepsilon_j([\phi_{j,\pm}])) - 1$ is a multiple of p by (45.8), and $pA^m \subset N$. The scalar forms for any enlargement of N_e meet hb by Step 3.

Let $Q = (X_j^2 : j \in J)$, and $X_{<e} = \prod_{\mathrm{ty}(j)<e} X_j$. Enlarge N_e to

$$K_e = N_e + (Q : X_{<e})\Lambda^s.$$

If $j \in I_e$ and $i < j$ is not in I_e, then $\mathrm{ty}(i) < e$ by monotonicity, so $X_i \in (Q : X_{<e})$. All the hypotheses of Proposition 45.2 are satisfied for K_e. Obtain

$$\Gamma_e \prod_{j \in I_e} X_j \in \mathrm{Fitt}(K_e), \qquad v(\Gamma_e(0)) \le b_e + (|I_e| + |s|)v(C).$$

By (45.2), choose $d_e \in \mathrm{Fitt}(N_e)$ satisfying

$$d_e - \Gamma_e \prod_{j \in I_e} X_j \in (Q : X_{<e}).$$

Using the ordered product expansion (replace factors one at a time from left to right while retaining the already replaced factors), this implies

$$\prod_e d_e \equiv \left(\prod_e \Gamma_e \right) X_* \pmod{Q}. \tag{45.15}$$

In fact the term replacing factor e contains the factor $X_{<e}$ from the preceding Γ-terms, so it vanishes modulo Q. The square-free quotient is used for coefficient comparison only, not to weaken the required ideal membership in the full coefficient algebra.

Step 5: The determinant on each scalar block. By (45.7) choose $q_e \in O[G]$ with

$$\eta_k(q_e) = C\,\delta_{ke}.$$

Explicitly use $q_e = c_e \prod_{k \ne e}([a] - \xi_k(a))$ where $C = c_e \prod_{k \ne e}(\xi_e(a) - \xi_k(a))$. If a factor is zero, the nontrivial finite-valuation case implies $C \ne 0$ and so cannot occur. We also choose the element

$$k = \prod_{x \in \mathfrak{X}} (y_x - 1)^{-1}(y_x - [\gamma_x]) \in O[G]$$

in any fixed order. Then $\rho_x(k) = 0$ for every distinguished x (one factor is the zero matrix), and $\eta_e(k) = 1$ for all e. No centrality of k in the group algebra is necessary.

For every e express d_e as a finite sum of row determinants of N_e with coefficients $h_{e,i}(X) \in \Lambda$. Denote by $B_{e,i}$ the resulting $|s| \times |s|$ matrices and choose rows $w_{e,i,j} \in N$ mapping to their rows under $A_e E_e$. For a tuple $\mathbf{i} = (i_e)_e$, form $M_{\mathbf{i}} \in M_m(A)$ whose rows are:

1. the rows (45.13) in the W slots;
2. the rows ν_i in the s_0 slots;
3. the rows $\sum_e q_e w_{e,i_e,j}$ for $j \in s$.

All rows lie in N. At character e, the first rows give a diagonal block with entries $o_v(1)$. The next rows have identity on the s_0 slots, and their elimination from the last rows gives CB_{e,i_e} by the definition of A_e. Consequently, writing

$$a_* = \left(\prod_{v \in W} o_v(1) \right) C^{|s|},$$

we have $\det(\eta_e(M_{\mathbf{i}})) = a_* \det B_{e,i_e}$. Put

$$M'_{\mathbf{i}} = B_0 + k(M_{\mathbf{i}} - B_0), \qquad h_{\mathbf{i}} = k q_0 \prod_e h_{e,i_e}(X).$$

The rows of $M'_{\mathbf{i}}$ remain in N; at $x \in \mathfrak{X}$ its ρ_x-block is B_0, and its scalar character blocks are those of $M_{\mathbf{i}}$. Also $u_x(\mathfrak{a} h_{\mathbf{i}}) = 0$: this holds on the ρ_x-block since $\rho_x(k) = 0$, on the $e \ne 0$ blocks since $\eta_e(q_0) = 0$, and on the $e = 0$ block by the definition $\mathfrak{a} = \ker \eta_0$.

Step 6: The actual polynomial and its coefficient. Apply Section 45.4 to $V, \omega, M'_{\mathbf{i}}, \mathfrak{a}, h_{\mathbf{i}}$. It yields $f_{\mathbf{i}} \in S$ with $f_{\mathbf{i}}\beta \in \mathcal{J}$ and on $\mathfrak{X}$

$$(f_{\mathbf{i}})_x = D_0(x, X)\, C\, a_*^t \prod_e h_{e,i_e}(X) \det B_{e,i_e}.$$

Here the decomposition of the flattened block determinant is the simultaneous row and column permutation to the ρ and character blocks, so its sign is unchanged. The trace of $h_{\mathbf{i}}$ on the distinguished family is exactly $C \prod h_{e,i_e}$, with no extra factor of the dimension. Put $f = \sum_{\mathbf{i}} f_{\mathbf{i}}$. It satisfies $f\beta \in \mathcal{J}$, and on $\mathfrak{X}$

$$f_x \equiv D_0(x, X)\, C\, a_*^t \Big(\prod_e \Gamma_e(X) \Big) X_* \pmod{QL[X]}.$$

Since the image of ξ_0 of every $u_x(g) - 1$ is zero, $\beta_x = 0$ on $\mathfrak{X}$. The coefficient conditions on f follow: the ideal Q consists of polynomials all of whose monomials have some exponent at least two. The required c before applying $O \to L$ is

$$c = D_0(x_0, 0)\, C\, a_*^t \prod_e \Gamma_e(0) \in O.$$

Sum the valuation bounds from Steps 2 and 4. The loss from the character determinants is $t \sum_v v(o_v(1)) + t|s|v(C)$; the loss from the triangular block determinants is $\sum_e b_e + (|J| + t|s|)v(C)$; and the trace of q_0 costs $v(C)$. This gives (45.11), exactly. □

The ordering in this proof matters. In particular, replacing the two Cramer operations by a pointwise inversion of β, replacing Q by ordinary total-degree truncation, or separating every character by an integral idempotent would not give the same conclusion. The square-free ideal, normalized rows and two adjugates retain the integral coefficient algebra throughout.

45.6 Closure status

Theorem 45.3 is the full algebraic statement of `ribet_first_order`, with its trace condition, triangle hypothesis and source valuation allowance. Its hypotheses are mathematical inputs explicitly stated in the theorem, not assumptions of semisimplicity or unproved algebraic estimates. The two auxiliary Fitting statements, matrix-block selection and Cramer construction have written proofs here. This discharges the abstract first-order polynomial interface previously open in Section 43.9. The remaining question is arithmetic: in `New/Selmer/RibetElement.lean` the particular automorphic systems must supply u, ρ, ξ, the trace ideal, the labelled local comparison, the Selmer length and triangle hypotheses, and the prescribed normalizer. The polynomial theorem does not by itself construct those systems. Combined with the integral extraction of Section 43.5, it does remove the abstract first-order ideal-membership step from the dependencies of that extraction.

45.7 The triangle cost for a Tate string

There is one more input to the abstract first-order theorem: the particular cost condition for the scalar Tate blocks. The source proves this in `New/Selmer/TateTwistSteps.lean`, `hasTriangleCost_tateTwist`. We supply its derivation, using the primitive odd laws and triangular primes already developed in Chapter 43.

Let $\xi(g) = \epsilon_\ell(g)^{-e}$ with $e \geq 0$ and $D = O/\ell^\nu O$. Let Z be a module of continuous cocycles in $D(\xi)$ ramified only above ℓ and the new pairs of places of a prime triangle. Use their two

evaluation generators $\tau_j, (\tau_j^c)^{-1}$ and $\phi_j, (\phi_j^c)^{-1}$. Assume the exact triangle conditions: for each nonzero type e, an even cocycle vanishing on the local cyclotomic kernels above ℓ and all the earlier prescribed tau and same-type phi evaluations has its τ_j value killed by $C = \ell^h$; and for an even cocycle unramified off ℓ satisfying these local kernel and same-type phi conditions one has

$$d(a)(z+z)(g) = d(g)(z+z)(a), \qquad d(g) = \xi(g) - 1, \qquad 4d(a)^2 \mid C.$$

These are the conditions `IsPrimeTriangle.step` and `IsPrimeTriangle.coboundary`; the latter uses the common separating element a. Assume also that any coboundary in Z is killed by $d(a)$.

Write L_e for the union of all conjugates of the inertia groups above ℓ if $e = 0$, and of the kernels of the full cyclotomic character on the decomposition groups above ℓ if $e > 0$. Thus L_e is `shaLocalSet` of the source. The relations holding on crystalline local cocycles force every mod-ℓ^ν cocycle obeying these relations to vanish on L_e. To see this, for $e > 0$ the filtration of $\mathbb{Q}_\ell(-e)$ has only the positive jump e; the Bloch–Kato rank formula of Section 8.4 gives $H^1_f = 0$. Hence rational finite cocycles are coboundaries and vanish on the kernel of ξ, in particular on the full cyclotomic kernel. For $e = 0$ the finite condition is exactly unramified. Indeed its rational dimension is one by that rank formula, and a nonzero unramified rational homomorphism is crystalline: choose u in the Witt constants $W(\overline{\mathbb{F}}_\ell)$ with $\varphi^f u - u = 1$, where f is the local residue degree. Such u is obtained by solving the Artin–Schreier difference successively modulo $\ell, \ell^2, \ldots$; at each step the residue equation $x^{\ell^f} - x = b$ is soluble in the algebraic closure. The resulting element of the crystalline period ring has inertia fixed and its difference under arithmetic Frobenius is one, so by continuity it gives the unramified cocycle. Multiplication by the coefficient scalar realizes all such homomorphisms. The mod-ℓ^ν relation that evaluates a cocycle at the indicated local element is therefore one of the relations of the integral crystalline submodule. At other conjugates of the selected decomposition group, a cocycle satisfies $c(kgk^{-1}) = \xi(k)c(g)$ whenever $\xi(g) = 1$, so the vanishing transports. Thus

$$\lambda(Z; \{c : c|_{L_e} = 0\}) \le \lambda(Z; \mathrm{obeying}(\mathrm{crysRels})),$$

where the right member is the original local block loss.

For a member z of Z zero on L_e, put $z^+(g) = z(g) + z(g^c)$. For $e \neq 0$ this is an even cocycle of the indicated character satisfying the required local condition. Since the chosen new Frobenii act trivially modulo ℓ^ν and tame inertia has full cyclotomic value one, the difference of either paired evaluation is exactly the sum under conjugation:

$$z(\tau_j) - z((\tau_j^c)^{-1}) = z^+(\tau_j), \qquad z(\phi_j) - z((\phi_j^c)^{-1}) = z^+(\phi_j).$$

Earlier equalities of the pairs therefore imply the vanishing tests in the triangle step. It follows that C kills the difference of the next tau pair. For $e = 0$ the type set I_e is empty, so this requirement is vacuous.

For the second triangle condition, let $Z' \subset Z$ consist of cocycles zero on L_e, all tau evaluations, and with equal same-type phi pairs. Vanishing at both tame generators implies unramifiedness at the new places, by the elementary tame-generator argument of Section 46.2. Thus Z' is unramified outside the primes above ℓ. For $e = 0$ it is already a submodule of shaOdd as defined in the source (there is no oddness condition for the trivial character). Its length is bounded by that of this group. For $e > 0$ the map

$$Z' \longrightarrow \mathrm{shaOdd}, \qquad z \longmapsto z \circ c - z$$

has image in the stated odd group: the difference is odd, unramified off ℓ, and vanishes on the conjugation-stable L_e. If z belongs to its kernel then z is even. Its same-type phi

evaluations satisfy $z^{+}(\phi_j) = 0$, so the triangle's last coboundary condition applies. It says that the cocycle $2d(a)z$ equals the coboundary $g \mapsto d(g)\,2z(a)$. This cocycle belongs to Z, hence by the hypothesized annihilation of its coboundaries,

$$2d(a)^2 z(g) = 0 \quad \text{for all } g.$$

In particular $Cz = 0$. The quotient $Z'/(Z' \cap D^G[C])$ is a quotient of $Z'/\ker(z \mapsto z \circ c - z)$, and therefore has length at most $\mathrm{length}_O(\mathrm{shaOdd})$. Combining this with the local-loss inequality proves the second triangle condition with

$$b = \lambda(Z; \mathrm{obeying}(\mathrm{crysRels})) + \mathrm{length}_O(\mathrm{shaOdd}).$$

This proves `hasTriangleCost_tateTwist` with its prescribed loss. In particular it does not presume an estimate for the even class groups, and no division by two is made in the argument. The two powers of the scalar in the final coboundary condition account for the entire kernel of the passage to the odd subgroup.

Chapter 46

The global mirror criterion for tied Selmer groups

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 47, 48 and the final audit in Chapter 57.

The proof of the mirror estimate can be reorganized using classical Poitou–Tate duality. In the source, a large family of relations between auxiliary split-prime values is first constructed in `Lib/ClassField/GreenbergWiles/SelmerRelations.lean`, and its cardinality is bounded below using an auxiliary projectivity argument. One can instead use the actual finite dual Selmer group, whose order is given by the Greenberg–Wiles count of Lemma 43.2. In the situation of the mirror theorem, a scalar acting without fixed vectors makes restriction to the common kernel injective on this dual group. The rest is an elementary algebraic argument on that kernel.

46.1 The isotropic pairing estimate

We keep all hypotheses of the source declaration `card_le_torsion_of_isotropic` in `New/Selmer/LinePointPowers.lean`. Let O be a commutative ring, $V = (O/I)^n$ finite, G a group, $c : G \to G$ an involutive automorphism, $\rho : G \to GL_n(O)$ a representation whose values span the full matrix algebra over O, and $\chi : G \to O^\times$ a character. Let $U \lhd G$ act trivially on V, and $H \lhd G$ be contained in U and stable under c. Suppose H contains $[g, \gamma]$ for all $g \in G$, and

$$\rho(\gamma) = s\,1, \quad \rho(c\gamma)^{-1} = s\,1, \quad \chi(\gamma) = \chi(c\gamma) = 1, \quad s - 1, s + 1 \in O^\times. \tag{46.1}$$

Let $y : G \to V$ be a cocycle, $y(\gamma) = 0$, having a unit coordinate at some g_0. Fix $z \in U$ with $cz \in U$. Write $V^\vee = \mathrm{Hom}_{\mathbb{Z}}(V, \mathbb{Q}/\mathbb{Z})$, and let $(x_w)_{w \in \mathcal{W}}$ be homomorphisms $U \to V^\vee$ satisfying

$$x_w(gug^{-1})(m) = x_w(u)(\chi(g)\rho(g)^{-1}m) \quad (g \in G,\ u \in U,\ m \in V).$$

Assume they are distinct as homomorphisms on U and satisfy

$$x_w(zh)(ty(zh)) = x_w(c(zh))(ty(c(zh))) \qquad (h \in H,\ t \in O). \tag{46.2}$$

Proposition 46.1 (Isotropy on a coset gives a torsion bound)**.** *If $g \in G$ satisfies $\rho(g) = \alpha\,1$ and $\rho(cg)^{-1} = \alpha'\,1$, then*

$$|\mathcal{W}| \le |V[\alpha - \chi(cg)\alpha']|.$$

Proof. Fix w and suppress the subscript. On U the cocycle y is additive and satisfies $y(kuk^{-1}) = \rho(k)y(u)$. The same holds for x with its indicated dual action. Put $\tau = [g_0, \gamma] = g_0\gamma g_0^{-1}\gamma^{-1}$. The cocycle identity and $y(\gamma) = 0$ give $y(\tau) = (1 - s)y(g_0)$. Thus $\tau \in H$ and $y(\tau)$ has a unit coordinate. For all $b, d \in H$, polarizing (46.2) on the coset zH gives

$$x(b)(ty(d)) + x(d)(ty(b)) = x(cb)(ty(cd)) + x(cd)(ty(cb)). \tag{46.3}$$

Indeed subtract the equations for zb and zd from that for zbd, and add the equation for z; the constant and linear terms cancel, irrespective of whether b, d commute in H, since both functions are additive there.

For fixed b, d write

$$A(t) = x(b)(ty(d)) - x(cd)(ty(cb)), \qquad B(t) = x(d)(ty(b)) - x(cb)(ty(cd)).$$

Equation (46.3) says $A(t) + B(t) = 0$ for all $t \in O$. Replace b by $\gamma b\gamma^{-1}$. By (46.1) and equivariance, the new equation is $A(s^{-1}t) + B(st) = 0$. Hence $A(s^{-1}t) = A(st)$ for all t. The map A is additive in t and $s - s^{-1}$ is a unit, so $A(t) = 0$ for every t. In other words

$$x(b)(ty(d)) = x(cd)(ty(cb)) \quad (b, d \in H). \tag{46.4}$$

No division by two has been made. With $d = k\tau k^{-1}$ this becomes

$$x(b)(t\rho(k)y(\tau)) = x(c\tau)(t\chi(ck)\rho(ck)^{-1}y(cb)). \tag{46.5}$$

Take $b = c\tau$. Apply (46.5) first to k, and then to kg. The second equation reads

$$x(c\tau)(t\alpha\rho(k)y(\tau)) = x(c\tau)(t\chi(cg)\alpha'\chi(ck)\rho(ck)^{-1}y(\tau)).$$

The first equation with t replaced by $t\chi(cg)\alpha'$ converts the right side to $x(c\tau)(t\chi(cg)\alpha'\rho(k)y(\tau))$. Thus $x(c\tau)$ annihilates

$$(\alpha - \chi(cg)\alpha')\ \mathrm{span}_O\{\rho(k)y(\tau) : k \in G\}.$$

The span is all of V: the matrix-span hypothesis and a unit entry of $y(\tau)$ allow any standard basis vector of V to be obtained by applying a matrix over O to $y(\tau)$. Hence $x(c\tau) \in (V/(\alpha - \chi(cg)\alpha')V)^\vee$.

Finally the value $x(c\tau)$ determines all of $x|_H$ by (46.5) and the same spanning argument. It also determines $x|_U$. For $u \in U$, use $[u, \gamma] = u(\gamma u^{-1}\gamma^{-1}) \in H$; then

$$x([u, \gamma])(m) = x(u)((1 - s^{-1})m).$$

As $1 - s^{-1}$ is a unit, this determines $x(u)$. Distinctness on U embeds $\mathcal{W}$ in the dual of the quotient displayed above. The latter has order $|V[(\alpha - \chi(cg)\alpha')]|$ by the kernel–image count on the finite group V. This proves the proposition. □

46.2 The dual Selmer group as the family of mirrors

Let O now be the integer ring of a finite extension of $\mathbb{Q}_\ell$, and retain the balanced row and pencil data of Proposition 40.9. We recall the exact parts of balance that are used. Let $\mathfrak{S}$ be the original finite ramification set, J the auxiliary pencil, and S_0 the finite set excluded from the pencil with $\mathfrak{S} \subset S_0$. Put

$$V = (O/aO)^n, \qquad a \neq 0, \quad a \mid \ell^M, \qquad S' = \mathfrak{S} \cup J \cup J^c.$$

The symbol n denotes the row rank. The representation ρ is unramified outside $\mathfrak{S}$, continuous modulo ℓ^M, and spans the matrix algebra. Its normalized cocycles obey fixed linear relations L and the two ties for each $P \in J$,

$$c(\tau_P) = t_P c(f_P), \qquad c((\tau_P^c)^{-1}) = t_P c((f_P^c)^{-1}).$$

They also vanish at γ, which is an inertia element at a member of $\mathfrak{S}$ not over ℓ. Balance provides s and the scalar identities (46.1) (for $\chi = \epsilon_\ell$) and all coboundaries obey L. The latter condition is stable upon adding the pencil ties: at every auxiliary place $\rho(\tau_P) = 1$, and at any nonzero slope t_P both Frobenius matrices are 1 modulo ℓ^M. Thus each coboundary $c_b(g) = \rho(g)b - b$ also satisfies the pencil ties. Notice

$$c_b(\gamma) = (s-1)b.$$

Consequently evaluation of these coboundaries at γ is onto V (indeed an isomorphism).

For each $v \in S'$ choose a decomposition group in G_E above v. Let $\mathscr{L}_v \subset Z^1(E_v, V)$ be the restriction of the module of all continuous cocycles obeying L and the pencil ties, without imposing global ramification or normalization at γ. This is precisely the source's `localCocycles` for the enlarged relation set. It contains all local coboundaries by the preceding observation. Its image in $H^1(E_v, V)$ will be denoted L_v; then

$$|L_v| = |\mathscr{L}_v|\,|H^0(E_v, V)|/|V|.$$

For $v \in \mathfrak{S}$ choose the decomposition group minimizing the length for the original relations L. There is a minimum because these lengths are nonnegative integers (local finite cohomology and finite coboundaries). With the added ties the length can only decrease. At $v \in S' \setminus \mathfrak{S}$ choose the group specified by $P \in J$ or its conjugate. On that group ρ is unramified and ℓ is different from the residue characteristic. Every ℓ-primary inertia cocycle is determined by its value on the fixed tame generator; the wild pro-p group has no nonzero homomorphism to V. After the tie, the entire cocycle is determined by its value at Frobenius, since a zero value there gives a zero inertia value and the quotient by inertia is procyclic. On the conjugate side the inverse Frobenius and inverse conjugate tame generator serve just as well. Thus

$$\operatorname{length}_O \mathscr{L}_v \le \operatorname{length}_O V \qquad (v \in S' \setminus \mathfrak{S}). \tag{46.6}$$

This is the local estimate used in `exists_conj_side_detects`. It holds for the entire local restriction module, not just for global cocycles unramified outside S'.

At all finite $v \notin S'$ take the unramified local condition for V. In applying the finite-module duality theorem include all primes over ℓ in the permissible ramification set, even if some do not belong to S'; the primal unramified condition at such a prime still contributes the ratio $|L_v|/|H^0(E_v, V)| = 1$, since ρ is unramified there. Its dual condition need not be described by an unramified subgroup at such a prime. At infinity E is totally complex.

Let Y be the space of global cocycles satisfying these local conditions in cohomology; since all local coboundaries belong to $\mathscr{L}_v$, restriction at $v \in S'$ belongs to $\mathscr{L}_v$. Let X denote the source module $X(J, a)$: cocycles globally obeying L and the ties, unramified off S', and zero at γ. It embeds into Y and in fact into the kernel of evaluation at γ on Y. The evaluation map on Y is onto V by the coboundary argument. Hence

$$\operatorname{length}_O X + \operatorname{length}_O V \le \operatorname{length}_O Y.$$

Give $V^\vee = \operatorname{Hom}_{\mathbb{Z}}(V, \mathbb{Q}/\mathbb{Z})$ the Cartier dual action $g \cdot \varphi(m) = \varphi(\epsilon_\ell(g)\rho(g)^{-1}m)$, using a choice of roots of unity of order ℓ^{M+1} to identify values with the corresponding subgroup of

$\mathbb{Q}/\mathbb{Z}$. Let $\mathcal{W} = H^1_{L^\perp}(E, V^\vee)$ be the finite dual Selmer group. Since γ has dual action s^{-1} with $s^{-1} - 1$ a unit, $H^0(E, V^\vee)$ is trivial; also $H^0(E, V)$ is trivial. Lemma 43.2, passing from cohomology to cocycles (a global coboundary group of order $|V|$), consequently gives exactly

$$|Y|\,|V|^{|S'|+f-1} = |\mathcal{W}| \prod_{v \in S'} |\mathscr{L}_v|, \qquad f = \#\{\text{complex places of } E\}. \tag{46.7}$$

Each factor is a power of the residue cardinality of O; the dual Selmer group is an O-module via precomposition on the dual and the orthogonality conditions are O-stable. In particular this equation is also a length equation. Combining it with the evaluation bound yields

$$\operatorname{length}_O X + (|S'| + f) \operatorname{length}_O V \le \operatorname{length}_O \mathcal{W} + \sum_{v \in S'} \operatorname{length}_O \mathscr{L}_v. \tag{46.8}$$

Put $U = \ker(\rho \bmod \ell^M) \cap \ker(\epsilon_\ell \bmod \ell^{M+1})$. It is open normal and acts trivially on V and $V^\vee$. Any dual class restricts to a continuous homomorphism $x_w : U \to V^\vee$, independent of the representative, and equivariant for the displayed dual action. These homomorphisms are distinct for distinct classes. To see this, any difference vanishing on U is inflated from G_E/U by the cocycle identity. The image of γ is central in this quotient by its scalar and cyclotomic properties, and acts as s^{-1} on $V^\vee$. For a cocycle q of the quotient, commuting γ with g gives

$$(s^{-1} - 1)q(g) = (g - 1)q(\gamma),$$

so q is the coboundary of $(s^{-1} - 1)^{-1} q(\gamma)$. Thus restriction to U is injective on $\mathcal{W}$. This is the precise reason the actual dual Selmer group suffices here; no general vanishing of cohomology of ρ's image is asserted.

46.3 The split-prime test gives isotropy on the required coset

Let c denote the complex-conjugation automorphism of G_E. Let N_γ be the normal subgroup generated by γ and $(c\gamma)^{-1}$ and take $H = [N_\gamma, G_E]$, with closure if desired. The subgroup H is conjugation-stable, normal, and contained in U: in the indicated finite quotient both generating scalars are central. Thus it has the commutators required in Proposition 46.1. The normal group generated by γ and $c\gamma$ is contained in the normal closure of the inertia at $\mathfrak{S} \setminus \{v \mid \ell\}$ and its conjugates.

Choose $z = \tau_{P_0}$, the tame generator at the helper prime of the pencil. Since P_0 and its conjugate lie outside the original ramification set and above primes other than ℓ, we have $z, cz \in U$. Every old anticyclotomic homomorphism h_P kills H and has $h_P(z) = \delta_{P,P_0}$ in additive notation. In particular each zh, $h \in H$, lies in the normal closure of inertia at a finite set of allowed places away from the primes over ℓ and from the conjugate closure of $J \setminus \{P_0\}$: use the original ramification set and its conjugate, and the two places belonging to the helper. This is exactly the subgroup condition for Proposition 41.9.

By the hypothesis of the mirror theorem choose $y \in X$ with a unit coordinate at some g_0. For $h \in H$ take a sufficiently small neighborhood of zh on which

1. both the element and its conjugate lie in U;
2. all old h_P retain their values $h_P(z)$;
3. y and its conjugate evaluations, and the finitely many homomorphisms x_w and their conjugate evaluations, retain their values at zh and $c(zh)$.

These are indeed neighborhood conditions by local constancy and finiteness. Apply the controlled auxiliary-character construction with triviality at the old places $J \setminus \{P_0\}$ and the primes above ℓ. Enlarge the finite set of excluded new places to include S_0, J and their conjugates and all exceptional places of the dual Selmer conditions (in particular the primes over ℓ). This does not change the allowed normal closure. We obtain a new degree-one split prime P, Frobenius f in that neighborhood, tame generator τ_P, and new anticyclotomic character h_P of the stipulated depth with $h_P(f) = 0$ and $h_P(\tau_P) = 1$.

This is an admissible enlargement of the pencil for *every* possible new slope in the considered row, with all other new slopes zero. Indeed the old character values on f are 1 for the helper and 0 for all nonhelpers (additive notation), as required; the new character vanishes at the old nonhelpers' decomposition groups and on its own Frobenius. The chosen Frobenius and its conjugate inverse both lie in U, so their row matrices reduce to the identity regardless of the new slope. All exclusions and anticyclotomicity required in `IsPencil` are respected. This explains why finding a prime separately for each slope would not suffice.

Apply the mirror hypothesis for this prime to y. It gives a cocycle y' in the module with ramification relaxed at P, P^c but the old ties and relations unchanged, such that

$$y'(\tau_P) = y(f), \qquad y'((\tau_P^c)^{-1}) = y((f^c)^{-1}). \tag{46.9}$$

For any $t \in O$, the global cup pairing of ty' with the dual class w has zero sum of local invariants. At all old places the terms vanish by the dual Selmer conditions: y' obeys the old relations and ties, and it is unramified at every other place except P, P^c. At the latter two places ρ and the dual local action are trivial (they are unramified, and their Frobenii act trivially). The dual class is unramified there, as these places are away from the exceptional set. The unramified-tame local pairing is evaluation of the dual cocycle at arithmetic Frobenius on the primal tame value, multiplied by the common choice of normalization of the tame generator. More precisely choose a primitive ℓ^{M+1}-th root of unity and fix a generator of the tame ℓ-quotient at P via its action on a root of the uniformizer. At P^c the compatible generator is the inverse conjugate of that generator, because conjugation inverts the chosen root of unity. The full tame generators $\tau_P, (\tau_P^c)^{-1}$ differ from these normalized generators by a unit power, the same unit on both sides. The two local invariant formulas therefore have a common unit factor. This is the usual unramified–tame cup formula, obtained by identifying unramified homomorphisms with their arithmetic Frobenius values and the other quotient with its inertia values in local Tate duality; the identification with the chosen root of unity is precisely the tame symbol. Equivalently apply the naturality of local reciprocity from E_P to E_{P^c}: the inversion of the chosen root at the second prime is canceled by the inverse on its tame generator. This is the same normalization used in the scalar case in Section 43.2. There is no factor two at this split pair, since the two completions are distinct.

The unramified dual value on the second arithmetic Frobenius is $x_w(f^c)$, not $x_w((f^c)^{-1})$. Since both groups act trivially here, $y((f^c)^{-1}) = -y(f^c)$ and likewise for the dual cocycle. Consequently the global reciprocity relation and (46.9) give

$$x_w(f)(ty(f)) = x_w(f^c)(ty(f^c)) \qquad (t \in O).$$

A common unit multiplying both tame values may be removed by varying t. Since the neighborhood fixes all the stipulated evaluations, the same equality holds with f replaced by zh. This is exactly (46.2), for every $h \in H$ and every $w \in \mathcal{W}$. The use of the mirror hypothesis therefore imposes the required isotropy on the actual dual Selmer group, not on an unspecified annihilator of a finite evaluation map.

46.4 Absorbing the scalar loss into the balance bound

By balance there are g, α, α' with the scalar identities of Proposition 46.1 and

$$\sum_{v\in\mathfrak{S}} \min_{A} \operatorname{length}_O \operatorname{localCocycles}(\rho, L, A, O/aO) + n\, v(\alpha - \epsilon_\ell(cg)\alpha') \le (|\mathfrak{S}| + f)\operatorname{length}_O V + \beta. \tag{46.10}$$

Put $\Delta = \alpha - \epsilon_\ell(cg)\alpha'$. Proposition 46.1 now bounds

$$|\mathcal{W}| \le |V[\Delta]|, \qquad \operatorname{length}_O \mathcal{W} \le \operatorname{length}_O V[\Delta] \le n\, v(\Delta).$$

If the right side of the balance bound is finite, all the compared lengths are finite; if the allowance β is infinite the conclusion is automatic. We may therefore add and cancel the finite lengths in (46.8). Use (46.6) to cancel the added $|S' \setminus \mathfrak{S}|$ places, and recall that at places of $\mathfrak{S}$ the original minimum was used. This gives

$$\operatorname{length}_O X \le \sum_{v\in\mathfrak{S}} \min_{A} \operatorname{length}_O \operatorname{localCocycles}(\rho, L, A, O/aO) + n\, v(\Delta) - (|\mathfrak{S}| + f)\operatorname{length}_O V \le \beta.$$

Thus the assertion of `IsPencil.length_le_of_unseen_class` (Proposition 40.9) is proved with the exact balance allowance. The residue-field condition used when choosing a slope in the tilting argument is not required in this mirror proof; it enters the subsequent iteration, exactly as in Chapter 42.

This conventional proof uses the classical Greenberg–Wiles identity and local and global Tate duality already specified in Section 43.2, together with the auxiliary-character construction of Proposition 41.9. It avoids the source's intermediate projective count of trivialized cocycles and the explicit family of split-prime relations: those encode the same needed duality inequality, but are not additional hypotheses for this proof. It does not construct a balanced row representation; the precise arithmetic construction of those representations is a separate obligation. In particular the conclusion here removes the mirror premise of Proposition 42.15, while the simultaneous-twist argument still requires the availability of the balanced rows and the integral value families.

Chapter 47

The arithmetic Ribet polynomial for columns with ties

We apply the abstract first-order theorem of Chapter 45 to the actual arithmetic hypotheses of `New/Selmer/RibetElement.lean`. Besides the algebraic construction, this requires the comparison of torsion cocycles with the crystalline local relations, a sum of local defects, conjugate-inverse trace ties, and the valuation bound with the three powers of the level group order. We give these steps separately; the existence of the globally constructed good forms remains a distinct question.

47.1 Torsion cocycles as reductions of rational coboundaries

Let O be the integer ring of a finite extension of $\mathbb{Q}_\ell$, G a group, $(u_x)_{x\in Y}$ a finite family $G \to GL_n(O)$, and $\rho : G \to GL_d(O)$. For $q = \ell^a \neq 0$ let $C(\rho, q)$ be the module of cocycles in $(O/qO)^d$ obeying all O-linear relations among the vectors $(u_x(g) - 1)_x$. Continuity is not assumed in this definition.

Lemma 47.1. *Every $c \in C(\rho, q)$ is the image of an integral cocycle on a finite free O-lattice Y_c which has an equivariant injection into a finite direct sum of the u_x-lattices. This integral cocycle is rationally a coboundary. If the u_x are continuous, so is c, and $c(g) = 0$ whenever all $u_x(g) = 1$.*

Proof. On $M = \prod_{x\in Y} M_n(O)$ give g the action of left multiplication by $u_x(g)$ in each factor. Write $s(g) = (u_x(g) - 1)_x$ and let P be the span of the $s(g)$. It is invariant, since $gs(h) = s(gh) - s(g)$. The relations make $s(g) \mapsto c(g)$ a well-defined linear map $f_0 : P \to (O/qO)^d$, equivariant by the cocycle identity. Choose k with $P \cap q^k M \subset qP$, by Smith normal form (the relevant Artin–Rees assertion). Set $Y_c = P + q^k M$ and $f(p + q^k m) = f_0(p)$. This is well-defined and equivariant. The module Y_c is a full stable lattice in M and $b(g) = s(g) \in Y_c$ is rationally the coboundary of $(1)_x$. Its image under f is c. Continuity follows because the map s is continuous and f kills $q^k M$. The last assertion follows by evaluating $s(g)$. □

These are the source assertions `obeyingCocycles.exists_graph_subrepresentation` in `New/Selmer/BlocksAndCocycles.lean` and `obeyingCocycles.mem_selmerModule` in `RibetElement.lean`. In particular, the ramification of the column family, not just that of the residual block, controls these cocycles.

Now restrict along $\Gamma \to G$, where Γ is identified with an open local Galois subgroup. If the family u_x is in a common local class $\mathscr{Q} = \mathrm{InClass}(F, K, U)$ of Chapter 52, then Y_c belongs

to the same class: rationally it is the direct sum of representations in that class. Thus the allowed crystalline period action, tame polynomials and jump sets hold, and $b \in Z_f(Y_c)$, since it is rationally a coboundary. No claim that c lifts to a crystalline cocycle of its own block is required.

47.2 Local duality and the defect of the crystalline relations

Let L be the local block lattice, also in $\mathscr{Q}$, $L' = L^\vee(1)$, and $V = L/qL$, $V^D = L'/qL'$. Use the perfect pairing with values in $O/qO(1)$, composed with a coefficient functional $\lambda : O \to \mathbb{Z}_\ell$ generating $\mathrm{Hom}_{\mathbb{Z}_\ell}(O, \mathbb{Z}_\ell)$ over O. The latter is the invertible inverse different, so λ gives perfect pairings on reduction without assuming that the ordinary trace alone is perfect.

Let A, B be the reductions of $Z_f(L), Z_f(L')$ in the respective finite cocycle groups. They contain all coboundaries. The integral finite condition is saturated: if multiplication by a nonzero scalar makes an integral cocycle a rational period coboundary, divide that coboundary by the scalar. Therefore the reduction kernel of Z_f is its multiple by q. Write d_Γ for the local degree. The rank formula of Section 8.4, with opposite Tate-dual weights $u, -u-1$, gives

$$\operatorname{rk} Z_f(L) + \operatorname{rk} Z_f(L') = (2 + d_\Gamma) \operatorname{rk} L, \qquad |A||B| = |V|^{2+d_\Gamma}.$$

Classical Bloch–Kato orthogonality followed by reduction gives their finite cup orthogonality, since $H^2(\Gamma, \mathbb{Z}_\ell(1))$ is torsion free. Dividing by the coboundary group orders $|V|/|H^0(\Gamma, V)|$ and $|V|/|H^0(\Gamma, V^D)|$ gives

$$|A/B^1(\Gamma, V)|\,|B/B^1(\Gamma, V^D)| = |V|^{d_\Gamma} |H^0(\Gamma, V)|\,|H^0(\Gamma, V^D)| = |H^1(\Gamma, V)|.$$

The last equality is the finite local Euler characteristic and Tate duality, with the exact references of Section 8.4. Thus these are mutual annihilators. In particular the calculation controls actual cocycles and retains both coboundary factors.

Put $H = Z_f(L') \cap \mathrm{cupOrthogonal}(\mathscr{Q}, L')$, and denote by $\bar{H}$ its reduction. Let S be the restrictions of global obeying cocycles, or the corresponding local module. Each $s \in S$ pairs to zero with $\bar{H}$: Lemma 47.1 gives $Y_s \in \mathscr{Q}$, $b \in Z_f(Y_s)$ and an equivariant map $f : Y_s \to V$ with $fb = s$. Choose an O-linear lift $\tilde{f} : Y_s \to L$. Its defect $g\tilde{f} - \tilde{f}$ is divisible by q, and

$$q^{-1}(g\tilde{f} - \tilde{f})$$

is an integral cocycle in $\mathrm{Hom}_O(Y_s, L)$, rationally a coboundary. Identifying this Hom with the Tate dual of $Y_s \otimes L'$, the lifted map and its divided defect supply exactly a `Lift.Pairing` with coefficient functional λ. By the definition of `cupOrthogonal` in `Lib/PadicHodge/Herr/GaloisLattices.lean`, the cup with every element of H bounds in the discrete multiplicative group of the local algebraic closure. The Kummer sequence and Hilbert 90 inject the finite H^2 of the roots of unity into that Brauer cohomology. Hence the corresponding finite Tate cup is zero. Nothing here says that the congruence map lifts equivariantly to rational duals.

We have $A + S \subset \bar{H}^\perp$, so by perfect finite duality (including the coboundary subgroups)

$$|S/(S \cap A)| = |(A+S)/A| \le |\bar{H}^\perp/A| \le |B/\bar{H}| \le |Z_f(L')/H|.$$

Every reduction in A obeys all original integral relations on $Z_f(L)$, and the restriction quotient of the global cocycles is obtained by pulling these relations back. Comparing lengths therefore proves

$$\begin{aligned} &\mathrm{relLength}_O(C(\rho, q); \mathrm{obeying}(\mathrm{relationsOn}\, Z_f(L)) \text{ along } \Gamma) \\ &\quad \le \mathrm{relLength}_O(Z_f(L'); \mathrm{cupOrthogonal}(\mathscr{Q}, L')). \end{aligned} \tag{47.1}$$

Here the quotient on the left is by the intersection with the module of functions satisfying the indicated local relations. This is the exact source assertion `relLength_obeyingCocycles_le` of `New/Selmer/RowBalanced.lean`, deduced by ordinary local duality rather than the intermediate source divisibility argument.

47.3 The common tame class and sum of the local block losses

Let $R = D.r$, $t = 5R$, $f = D.f$, and let r_0, t_0 be the actual row rank and string length, $1 \le r_0 \le R$, $t_0 \le t$. Assume the actual family of columns with the common local labels and jumps of `IsLabelledColumn`. If at an inertia element the roots of its killing polynomial are y_i $(i \le j)$, each with $y_i^{p^{d_i}-1} = 1$, $1 \le d_i \le R$, then they are Teichmüller roots in unramified fields of respective degrees dividing d_i. Their monic minimal polynomials over $\mathbb{Q}_p$ are integral and of degree at most d_i; their product is monic over $\mathbb{Z}_p$ of degree at most jR. All are killed by $X^m - 1$ for $m = \operatorname{lcm}_{1 \le d \le R}(p^d - 1)$. Thus the class of polynomials vanishing on those roots satisfies `Lift.IsTameType` with degree jR. In the actual labelled family the roots counted by Theorem 44.4 number at most $r_0 + t_0 \le R + t$, giving the parameters $R, (R+t)R$ of Section 53.3. This proves `isTameType_of_roots`.

A rational change of lattice does not change the period invariants, their filtration, or the inertia period polynomial, since one is making the same rational representation after coefficient extension. In particular `inClass_of_isLatticeIn` follows from the labelled local–global comparison: at a place over ℓ the representations are crystalline except possibly at the marked primes, where the linearized period action is finite on inertia and is killed by the prescribed polynomial. If one chooses for the common class every polynomial vanishing on the roots of a killing polynomial of each of the column representations, such a polynomial kills every indicated action. This uses semisimplicity of a finite-order operator in characteristic zero (on the period space at the coefficient prime); it does not infer a killing identity from a characteristic polynomial with possible nontrivial Jordan blocks.

Suppose one member of the family is on the nose over O the representation

$$T \oplus \bigoplus_{e=0}^{t_0-1} \xi_e, \qquad \xi_e = \epsilon_\ell^{-e},$$

with T locally equal over ℓ to the original integral row lattice. Every block belongs to the common local class by its direct-summand injection. Since the column jumps of this member are the sum, with multiplicity, of its block jumps, every permitted jump occurs in at least one block. This is the coverage hypothesis of `block_defect_le_at_place`. At each place $v \mid \ell$ of degree d_v, the estimate of Section 53.3 applies to the row and all strings: their normalized losses are at most $d_v r_0 \operatorname{loss}(\ell)$ and $d_v \operatorname{loss}(\ell)/t$, respectively.

For each block ρ_i and arbitrary modulus ℓ^{a_i} let $\operatorname{blockLoss}(\rho_i, u, a_i)$ be the relative length of the obeying-cocycle module modulo the relations on the integral crystalline cocycles at every place over ℓ. The quotient by the intersection of the local relation submodules injects into the product of their quotients. Apply (47.1) at each place, and the explicit one-place bounds. We obtain

$$\begin{aligned}\sum_i \operatorname{normLength}_O \operatorname{blockLoss}(\rho_i, u, a_i) &\le \sum_{v \mid \ell} d_v (r_0 + t_0/t) \operatorname{loss}(\ell) \\ &= 2f(r_0 + t_0/t) \operatorname{loss}(\ell) \le 4 f r_0 \operatorname{loss}(\ell). \end{aligned} \tag{47.2}$$

Here $\sum d_v = [E : \mathbb{Q}] = 2f$, $t_0 \le t$, and $r_0 \ge 1$. The coefficient DVR may be enlarged once for the finite collection of columns and local places. This proves `sum_blockLoss_le` with the actual factor $4fr_0$, rather than paying the worst row loss for each of the string blocks.

47.4 The conjugate-inverse trace ties

For a conjugate self-dual column in the source normalization, put $\chi_0 = \epsilon_\ell^{-1}$. Its trace satisfies

$$\operatorname{tr} u(g^c) = \chi_0(g)^{t_0-1} \operatorname{tr} u(g^{-1}).$$

By cyclicity of trace and conjugation invariance of χ_0, for $i \geq 0$,

$$\operatorname{tr} u((h^c)^{-i} g) = \chi_0(g)^{t_0-1} \chi_0(h)^{-i(t_0-1)} \operatorname{tr} u(h^i (g^c)^{-1}). \tag{47.3}$$

The auxiliary τ_j has $\chi_0(\tau_j) = 1$, and ϕ_j fixes the ℓ^ν-th roots of unity, hence $\chi_0(\phi_j) \equiv 1 \bmod \ell^\nu$. Suppose the trace ideal $\mathcal{J}$ contains $\ell^\nu \operatorname{tr} u(g)$ for all g and the ties at τ_j, ϕ_j, with polynomial ε_j. Expand $(X-1)\varepsilon_j(X) = \sum_i d_{ji} X^i$. In the tie with test element $(g^c)^{-1}$, apply (47.3) to each term. The common factor is $\chi_0(g)^{t_0-1}$; the extra factors for ϕ differ from one by multiples of ℓ^ν, so the resulting errors are in $\mathcal{J}$. Therefore the tie with the same variable X_j holds at the inverse-conjugate pair $((\tau_j^c)^{-1}, (\phi_j^c)^{-1})$. No second variable or valuation charge is needed. Also $\varepsilon_j(1) = 1$ gives $\ell^\nu \mid \varepsilon_j(\xi_e(\phi_{j,\pm})) - 1$. This proves the precise assertion `isTraceIdeal_withConjInv` of `RibetElement.lean` and the compatibility for the scalar blocks in the abstract first-order theorem.

47.5 Integral arithmetic specialization and valuation budget

Put $M_{\mathrm{gr}} = D.\mathrm{chosenGroupOrder}$, $c_\ell = v_\ell(M_{\mathrm{gr}}) + 2^{-\ell}$, $H = h_2 h$, $C = \ell^H$, with $h \leq v_\ell(M_{\mathrm{gr}})$ the integer from the prime-triangle construction, and $h_2 = R^8$. Write $X_{\mathrm{odd}} \neq 0$ for the integer of Theorem 42.14. The exact source budget is

$$\beta_\ell = h_2^{27}(c_\ell + v_\ell(X_{\mathrm{odd}})) + 4f \operatorname{loss}(\ell). \tag{47.4}$$

Assume the hypotheses of `ribet_columns`: an apt pair and labelled family u_x with its common Hodge jumps and local conditions, an auxiliary twist ψ of conductor Q, whose tame cost is at most c_ℓ and which is trivial on the decomposition groups over ℓ, and the prescribed prime triangle at depth ν. At the distinguished set $\mathfrak{X}$ the trace is $\psi(g) \operatorname{tr} R_x(g) + \sum_{e<t_0} \xi_e(g)$. Here $\xi_e = \epsilon_\ell^{-e}$, R_x is a row of rank r_0, and `SelmerLengthLe` at depth $m_0 > \lfloor R\beta_\ell \rfloor$ supplies its integral model $T_x = \psi R_x$, with $T_x(\gamma_x) = y_x I$, $y_x - 1$ a unit, and $\epsilon_\ell(\gamma_x) = 1$. The normalized length of its cocycles modulo ℓ^{m_0}, unramified off ℓ, g_0, g_1, Q, Q^c, obeying the crystalline relations and zero at γ_x, is bounded by $(10fr_0 + 1)c_\ell$. These are exactly the stated inputs of the theorem, not consequences of the polynomial construction.

A continuous compact image in $GL_n(\overline{\mathbb{Q}}_\ell)$ lies in a finite coefficient field: it is a compact group covered by the closed subgroups given by intersecting with $GL_n(F_i)$ for the countably many finite subextensions F_i. By Baire one intersection is open, of finite index; adjoining entries of finitely many coset representatives suffices. A stable lattice is the sum of translates of the standard lattice, a spanning submodule of one bounded scalar multiple of the latter. Enlarge the coefficient DVR once for the finite family, tie coefficients, row models and local roots. At distinguished points use the block lattice $T_x \oplus \bigoplus_{e<t_0} \xi_e$. This is a model of the original representation: T_x has full matrix span (twisting group matrices by units preserves the span), so it is absolutely irreducible; the original column's trace-nondegeneracy implies semisimplicity, since the radical of its image algebra pairs to zero under trace. Equal traces identify the two semisimple representations by testing the simple-block idempotents of the joint image algebra. No integral conjugacy between arbitrary stable lattices is inferred. At other points use any stable model. Their traces and characteristic coefficients remain the original ones. At ℓ the distinguished block agrees with the integral row lattice there, because ψ is locally trivial.

Use the abstract theorem with moduli $p = \ell^{\nu}, p_0 = \ell^{m_0}$. Write $e_O = v_O(\ell)$. Its distinguished allowance b_0 is the integral length of obeying cocycles on T_{x_0} modulo p_0 zero at γ_{x_0} and the two tame generators of every triangle prime. Lemma 47.1 proves they are continuous and unramified outside the column ramification. Their zero values remove the new primes: the row is unramified there, and its cocycles on inertia are homomorphisms to an ℓ-primary module determined by the ℓ-part of the tame generator. Intersecting with the crystalline relations thus gives the short-Selmer module bounded by the stipulated allowance. The quotient is bounded by its block loss, so

$$b_0/e_O \le (10fr_0 + 1)c_\ell + \operatorname{normLength} \operatorname{blockLoss}(T_{x_0}, u, m_0). \tag{47.5}$$

By (47.2) this is $\le r_0\beta_\ell \le R\beta_\ell < m_0$, giving the strict distinguished hypothesis of Theorem 45.3.

For each scalar block e, a module of obeying cocycles at ℓ^{ν} satisfying the scalar hypotheses of that theorem is unramified off ℓ and the triangle primes: it vanishes at the additional tame generators over the twist and marked places and their inverse-conjugates. Its coboundaries are killed by $\xi_e(a) - 1$ by the abstract theorem's Step 3. Apply Section 45.7 to obtain the triangle-cost allowance

$$b_e/e_O \le \operatorname{normLength} \operatorname{blockLoss}(\xi_e, u, \nu) + v_\ell(X_{\mathrm{odd}}). \tag{47.6}$$

The integer b_e may be chosen as the sum of the integral block loss and the odd-group length, which works for all smaller scalar modules. Enlarge the triangle's detecting set s by the two tame generators at all its new primes, giving s_*. Zero values here remove that ramification and the original detector gives zero. This is exactly the scalar detection hypothesis. The zero string type has empty triangle set and no oddness restriction in its odd-group definition, as required.

For a tame generator δ at a place of Q or its inverse conjugate use $o_\delta(X) = \psi(\delta) - X$, since the source hypothesis is $(u_x(\delta) - \psi(\delta)I)(u_x(\delta) - I) = 0$ for every x. For a generator at a marked prime $g_i \ne \ell$ use $o_\delta(X) = \sum_{j=0}^{M_{\mathrm{gr}}-1} X^j$: the finite-type inertial operator has roots of orders dividing $g_i^d - 1$ for some $1 \le d \le R$, hence dividing M_{gr} by the level general linear group count. Its finite order makes it semisimple in characteristic zero. All string characters are trivial at these inertia elements. This proves (45.9). The two sets of auxiliary polynomial charges are respectively at most $2c_\ell$ (anticyclotomicity gives $\psi((\delta^c)^{-1}) = \psi(\delta)$) and $8fv_\ell(M_{\mathrm{gr}})$ (at most $4f$ marked places, two sides each). Thus

$$\sum_{\delta} v_\ell(o_\delta(1)) \le (2 + 8f)c_\ell. \tag{47.7}$$

The prime triangle gives $\prod_{k \ne e}(\xi_e(a) - \xi_k(a)) \mid \ell^H$ and $(|J| + |s|)H \le h_2^{22}h$. Because $|s_*| \le |s| + 2|J|$,

$$2|s_*|H \le 4h_2^{22}h, \qquad (|J| + 1)H \le h_2^{22}h + H. \tag{47.8}$$

These inequalities also hold for $H = 0$. The conjugate-inverse trace conditions and evaluation of the tie polynomials were proved in Section 47.4.

Theorem 47.2 (Arithmetic Ribet polynomial)**.** *With all these hypotheses of* `ribet_columns`*, for any nonnegative finitely supported perturbation (n_g), put*

$$R_* = [\gamma_{x_0} a] - y_{x_0}[a] + \ell^{m_0+H} \sum_g n_g[g], \quad \beta_x = \det \sum_g (R_*)_g (u_x(g) - 1).$$

There is a Ribet polynomial F in the actual coefficient algebra with $F\beta \in \mathcal{J}$, $\beta = 0$ on $\mathfrak{X}$ and zero coefficients on that subset at every proper divisor of $X_* = \prod_{j \in J} X_j$. Its coefficient d at X_* and x_0 satisfies

$$v_\ell(M_{\mathrm{gr}}^3 d) \le r_0\beta_\ell. \tag{47.9}$$

Proof. Apply Theorem 45.3 over the common DVR with the preceding data. Its characteristic-coefficient functions and traces agree with the original column functions in the stipulated coefficient algebra, and its trace ideal conditions were verified there; polynomial adjugates use no inverse of a nonunit. Dividing integral valuation by e_O, the bound is

$$v_\ell(d) \le b_0/e_O + \sum_{e<t_0} b_e/e_O + t_0 \left(\sum_\delta v_\ell(o_\delta(1)) + 2|s_*|H \right) + (|J|+1)H.$$

Apply (47.5), (47.6), (47.2), (47.7), (47.8), and add $3v_\ell(M_{\mathrm{gr}})$. Since $h \le c_\ell$ the result is

$$\begin{aligned}(10fr_0 + 4 + t_0(2 + 8f + 4h_2^{22}) + h_2^{22} + h_2)c_\ell \\ + t_0 v_\ell(X_{\mathrm{odd}}) + 4fr_0 \operatorname{loss}(\ell).\end{aligned}$$

This is at most $r_0\beta_\ell$ because $t_0 \le t \le h_2$, $f \le h_2$, $r_0 \ge 1$ and

$$4 + 10h_2 + h_2(2 + 8h_2 + 4h_2^{22}) + h_2^{22} + h_2 < h_2^{27}.$$

For the explicit data bound, Chapter 53 gives $[E:\mathbb{Q}] \le 2N_1 \le 2e^{12\log X} < R^8 = h_2$. The polynomial inequality holds already for $h_2 \ge 10$. This proves the exact budget, including the three level-order factors. □

47.6 Status

These derivations close the arithmetic `ribet_columns` interface, including `sum_blockLoss_le`, `relLength_obeyingCocycles_le` and conjugate-inverse ties. Column labels, the original trace ideal, and the existence of the stipulated row, short-Selmer and prime-triangle data are its explicit hypotheses. Their realization for the actual projected cut of `denominatorLe_rowBudget` remains separate. The first-order polynomial with its exact arithmetic budget is now available there.

Chapter 48

From the labelled finite level to a denominator operator

The previous chapter establishes the actual Ribet coefficient bound. A further issue is whether the coefficient algebra to which that argument applies is the algebra of integral Hecke functions of the forms being tested. This chapter reconstructs the passage. Here we continue to suppose the cut is spanned by lifts with the row and short-Selmer properties in the statement of `denominatorLe_rowBudget`; we do not deduce that spanning hypothesis from the patching argument.

48.1 The base level and its arithmetic labels

Let O be the auxiliary pair and c an apt tuple of finite labels, and let e_c denote the actual central tuple projector acting at its finite split places. Set $M = M_{\mathrm{gr}}$, the common group-order multiple, and denote the normalized column representation of a system x by U_x. Let K_0 be the source base subgroup, with its character at the twist places and its ordinary spherical factors elsewhere. At the labelled places the level contains the kernel of reduction in the chosen lattice. In the global smooth forms of this level and transformation character form the subspace

$$B = \{b : e_c b = b\}.$$

At an extra finite family of places outside the prescribed bad set, take the corresponding deeper level and apply the same projector. This description does not impose an extra constraint at the new places: the projector is supported only on the labelled places.

Proposition 48.1 (The labelled base level)**.** *For the apt tuple and auxiliary twist of* `New/Selmer/BaseLevelOfLabel.lean`*, this defines a base level in the sense of that file. Its base space contains the smooth image of e_c; Me_c preserves the integral value lattice. There is one common jump tuple J for all systems at every finite added level off the bad set, with normalized inertial labels* $\mathrm{twistedLabel}(c)(X-1)^{t_0}$*, $t_0 = \mathrm{rankDiff}(O)$, and at the twist inertia*

$$(U_x(g) - \psi(g)I)(U_x(g) - I) = 0.$$

Proof. In the source representation p each element of the finite group is represented by translation of its residue lift supported on the labelled places, together with the prescribed finite-order determinant and character factor. The character-level space is stable under this representation, since the principal residue kernel is normal. The isotypic projector in this representation is idempotent; its image is precisely its fixed space, and translation of an

original level vector by e_c is the smooth image of its tuple projection. An endomorphism commuting with translations on the labelled places, and preserving the ordinary level-character space, commutes with the finite sum defining the projector. In particular new-place operators preserve B and the deeper spaces so defined. This verifies the base-level invariance, the base inclusion, and the level comparison, not just an abstract isomorphism of complex eigenspaces.

For the integrality assertion let G_c be the product group involved in this tuple idempotent. Its central character formula is

$$e_c = \frac{\dim \rho_c}{|G_c|} \sum_{g \in G_c} \chi_c(g^{-1}) p(g).$$

The character values are algebraic integers, all phase factors are roots of unity, and the residue translations preserve the stipulated value lattice. The size and aptness bounds ensure $|G_c| \mid M$, by the factor-by-factor group-order divisibility of the data. Thus Me_c preserves integral values, at every coset and at every new level. The corresponding class stabilizer order also divides M (the same mass and stabilizer bound used in Chapter 35). These statements are independent of the auxiliary level.

For a nonzero simultaneous eigenvector the finite tensor product at the labelled places lies in the selected finite character type; nonvanishing of its component follows from $e_c b = b$. At every other place of the original bad set outside the label and twist support the local vector is spherical; new levels are disjoint from this support. At the twist places the level character is the determinant character of the selected subblock, trivial on the other block. Chapter 44, especially Theorem 44.4 and Proposition 44.5, identifies the attached representation's split tame inertia exactly with the product of these Green labels and determinant-block characters, with zero monodromy. The jumps are those of the single underlying archimedean weight and therefore do not vary with x or with the extra level. Twist by the one-dimensional inverse avatar of the normalizing character (the source `chiOneInv`); this amounts to multiplying every root by its scalar value, giving the asserted normalized label with t_0 copies of 1. At a twist place the original two character blocks become precisely the trivial and ψ blocks, hence the displayed product annihilates inertia, including repeated roots (there is no unipotent monodromy). At the conjugate places conjugate self-duality gives the inverse character and the same assertion by the anticyclotomic condition. This is exactly the construction `exists_baseLevel`, including the conclusion on all added levels. □

48.2 The integral trace functions

Write $\mathcal{O}$ for the valuation ring of $\overline{\mathbb{Q}}_\ell$. At a fixed finite added level there are finitely many eigen-systems, and the diagonal functions form a finite product. The Hecke algebra H is the integral closed subalgebra in that product generated by the integral spherical Hecke functions at the allowed places. The crucial fact is not an unrestricted complex generation assertion; the closure is integral. For each g and all systems simultaneously, characteristic coefficient functions of $U_x(g)$ belong to H: first do this for an allowed spherical Frobenius using the normalized local-global comparison and the unit normalization, then use density of the allowed Frobenius conjugacy classes and continuity, at the finitely many representations in question. This is the closedness argument of Chapter 43. Consequently every trace function of $U_x(g)$ belongs to H, as does $x \mapsto \mathrm{Tr}(P(U_x(h))U_x(g))$ for an integral polynomial P: expand it into traces of $h^i g$. In particular this algebra contains the characteristic coefficients of all finite linear combinations of matrices used in the normaliser, by the integral determinant-coefficient algebra (equivalently the integral characteristic polynomial law for the representation of the group algebra). The same argument works at the conjugate places. The quadratic identity

just proved and the exact normalizing character at added places are the extra type premises for the first-order tie. Lemmas 43.6, 43.7, 43.9, and 43.4 therefore give the actual local tie with its stabilizer multiple and factorial, not merely equality of traces over a field.

48.3 A cyclotomic element and a fixed integral cutoff

Lemma 48.2 (Cyclotomic separation)**.** *There is a g_* with $v_\ell(\chi_{\mathrm{cyc}}(g_*)^a - 1) \le h$ for every $0 < a < t$, where $h = v_\ell(M)$ if $\ell \le h_2^2$ and $h = 0$ otherwise.*

Proof. For $\ell > h_2^2$ the cyclotomic ground field of the data has degree and conductor prime to ℓ, and the mod-ℓ cyclotomic character on inertia is onto. Take an inertia element mapping to a primitive element of $\mathbb{F}_\ell^\times$. Since $t \le h_2 < \ell - 1$, none of the indicated powers has reduction one. Otherwise take a marked prime of the data other than ℓ; one exists since the two marked rational primes are distinct. At its degree-one place the cyclotomic value of arithmetic Frobenius is the rational prime g_i. The definition of the common group-order multiple gives $g_i^a - 1 \mid M$ throughout this range. Thus the bound also holds at small ℓ. This proves the source `exists_cycChar_pow_sub_one_dvd`. □

Let $\mathfrak{X}$ be the systems represented in the given cut after applying e_c. Each has a row with a short Selmer bound, and the row-string identity gives the factor $(X-1)$ of every characteristic polynomial of the normalized column; more strongly the row-string representation contains the trivial summand. Put

$$S = \{x : \det(U_x(g) - I) = 0 \text{ for every } g\}.$$

Thus $\mathfrak{X} \subset S$. Fix $x_0 \in \mathfrak{X}$. At its scalar-inertia element γ and its row scalar y use the Ribet normaliser of Theorem 47.2, with a perturbation coefficient divisible by the chosen precision. For each finite set of systems outside S the determinant can be made simultaneously nonzero. Indeed if a new system x has an invertible $U_x(g) - I$, adding $\ell^j n(U_z(g) - I)$ to the combination for every z gives a nonzero determinant polynomial in n at x (its highest coefficient is nonzero); at all previously treated systems its constant coefficient is nonzero. Avoid the finitely many integer roots. Induction over all systems outside S proves the assertion. The resulting integral function β vanishes on $\mathfrak{X}$ because there is a common trivial summand. Take

$$p(T) = \prod_{x \notin S} (T - \beta_x), \qquad p(0) \neq 0.$$

The empty product is allowed. Importantly, $p(\beta_x) \neq 0$ implies $x \in S$, not that x belongs to the cut.

For each system not congruent to x_0 in the integral Hecke algebra take a function vanishing on that system and a unit at x_0, and multiply these functions to obtain A. Then $T = Ap(\beta)$ is supported on weak systems in S congruent to x_0. This is the exact sense of congruence in Proposition 43.5; it implies congruence of every characteristic coefficient since they belong to H. The construction does not suppose that all weak systems are genuine lifts.

48.4 The prime neighborhoods and the string condition

Choose the cyclotomic separating element of Lemma 48.2. The prime-triangle construction of Section 41.5, with the exact primitive-law input now supplied by Theorem 43.3, gives a single finite ordered set J of detector types, with realizations at every positive depth ν. The

index set is fixed before choosing ν. Here the allowable Frobenius set is as follows. At the distinguished system use the scalar-inertia element γ and its scalar (whose nontrivial powers through the row rank differ from 1 by units); Lemma 42.23 eliminates eigenvalue 1 in the twisted row by translating each prescribed Frobenius by a suitable power of γ^{ℓ^ν}. This is precisely the genericity condition in `exists_isPrimeTriangle`; the cyclotomic value of the scalar inertia element is one. In sufficiently small neighborhoods the determinant of 1 minus the twisted row remains a unit. Simultaneously impose the spherical conditions and omit the finite bad set and all old-level primes. These conditions are possible by the degree-one split Frobenius approximation used in Section 41.5.

For every weak system $x \in S$ congruent to x_0, the normalized characteristic polynomial at an allowed Frobenius has the root 1. The Hecke normalization translates this into the exceptional-magnitude root of Proposition 42.24. The unitary string through it has the full prescribed length t_0. At the triangle place of norm 1 modulo ℓ^ν all these Tate roots are congruent to 1 at the required precision. On removing the string, the remaining polynomial has unit value at 1: its coefficients reduce to those of the distinguished row, because the full coefficient functions of the two columns are congruent and division by the monic string commutes with reduction. The neighborhoods imposing this condition are open and there are finitely many weak systems. Lemma 42.22 now gives the integral polynomial multiple at each weak system with value 1 at 1. The product of those polynomials, at each prime, has the same normalization and is a multiple for every surviving weak system. This is the polynomial e_j used in the one-prime tie and in Theorem 47.2, including its normalization $e_j(1) = 1$. The argument requires the full local spherical string, not merely a factor $X - 1$ in the characteristic polynomial modulo ℓ.

The normalizer perturbation and the scalar separation are taken with the fixed modulus prescribed by the Ribet construction, independently of the ultimate triangle depth. Once $p(0)$ is fixed, take ν greater than

$$v_\ell(\kappa(M^2 p(0))^{|J|+1}), \qquad \kappa = 2n!,$$

and greater than all fixed lower precision requirements of the source. This is possible because the same ordered detector set works at every depth, even when its total cost is zero. Choose the actual triangle at this depth. The proof of Proposition 43.5 applies to this very level by Proposition 48.1; its callback is now Theorem 47.2. It produces a scalar function s with

$$v_\ell(M^3 s(x_0)) \le r_0 \beta_\ell.$$

Moreover, for old-level forms l, y with d, l integral, y supported on $\mathfrak{X}$, and l, y having equal components wherever $p(\beta_x) \neq 0$, it gives integral $dMD(s)y$. In this application we may strengthen congruence localization by the factor A from the extraction proof. This closes the actual trace-to-operator step of the source, not the existence of the cut.

48.5 The tested complement and the three order factors

Proposition 48.3 (Denominator implication for a stipulated lift cut)**.** *Suppose the apt lift cut Z has precisely the spanning and row hypotheses of* `denominatorLe_rowBudget`*, including the auxiliary twist and short-Selmer modulus, and let β_ℓ be the row budget. Then the determinant inequality* (40.2) *holds with allowance $\beta_\ell \log \ell$.*

Proof. The commuting split Hecke operators act by scalars on the stipulated spanning lift vectors. Their span and its orthogonal complement are invariant: the adjoint of each spherical operator is also a split Hecke operator, using the unitary normalization. Distinct tuple labels

are orthogonal since the finite representation is unitary and their central idempotents are self-adjoint. In particular e_c preserves Z and its orthogonal complement. Choose a basis of nonzero vectors of the stipulated spanning family and fix one basis vector v_0 with label c and system x_0. The following construction is made for this basis vector.

Let v be a tested integral form, $l = e_c v$, and $y = e_c P_Z v$. Then Ml is integral by Proposition 48.1, and y is supported on $\mathfrak{X}$. The difference $l - y$ is in $Z^{\perp} \cap \operatorname{im} e_c$. Every weak old system in S satisfies the exceptional-root test at split spherical places by its definition and the Hecke comparison. We may choose a split place outside the level at which this test is valid simultaneously for all weak systems: the source chooses one place satisfying the joint Frobenius and split conditions, which are finitely many conditions of the foregoing spherical comparison. On the isotypic part of $Z^{\perp} \cap \operatorname{im} e_c$ belonging to any weak system, the operator $H_w(X_w)$ consequently vanishes. Thus the tested orthogonality (40.1) implies that $l - y$ has zero component on that space. Indeed for a vector u in the space its pairing with $l - y$ equals its pairing with v, since e_c is self-adjoint and $u \perp Z$. It follows that l, y have equal components on every system where $p(\beta_x) \neq 0$. This verifies the particular comparison needed by the extraction proposition, even though the weak system itself need not be a lift.

Apply extraction with $d = M$ to get $M^2 D(s) e_c P_Z v$ integral in the value lattice. Pairing with an all-embeddings integral u costs at most another factor M: expand the class sum, whose denominators are stabilizer orders dividing M, and use integrality of the Weyl-coordinate pairing (all conjugate values of u are integral). Hence

$$M^3 \iota\langle u, D(s) e_c P_Z v\rangle \in \mathcal{O}.$$

The operator on Z is diagonal on the chosen basis, zero on other labels, with scalar $s(x_0)$ at v_0. Its valuation including the three factors of M was bounded above by $r_0 \beta_\ell$. Repeat for each vector of the basis. The determinant-of-pairings argument at the end of Section 40.12 supplies an integral linear combination of these diagonal operators whose determinant on the cut, including the common factor M^3, has valuation at most $q r_0 \beta_\ell$, $q = \dim Z$. The corresponding matrix of pairings is integral. Taking its determinant and cancelling the nonzero diagonal determinant gives exactly (40.2). The empty cut gives the empty determinant one. □

48.6 Status

The proof in this chapter, together with Chapters 44, 45 and 47, gives the labelled base level, integral trace functions, first-order local operators, arithmetic Ribet polynomial, squarefree extraction, and the denominator implication for the specified cut. The construction of actual good rows and the matching lift family and short-Selmer hypothesis of that implication, and the rank-two special-value comparison needed for the simultaneous fit, remain separate. A denominator theorem with a stipulated spanning family cannot substitute for construction of that family.

Chapter 49

Local crystalline defects and the integral cup-product estimate

Chapters 50 and 51 supply further proofs of the classical cohomological rank and index calculation, the chart algebra, depth and long-gap lifting and integral cokernel estimates, subject to the precise comparison hypotheses specified there. Chapter 52 supplies the actual integral structural comparison, frame and radius, and exact local cutoff and tame estimates. The global arithmetic row hypotheses remain distinct from the intrinsic local results.

The congruence argument for the lower bound on the determinant needs considerably more than the vanishing of the ordinary Bloch–Kato pairing on two crystalline representations. The two classes to be paired arise from lattices which need not be dual to one another in characteristic zero; only a pairing of their reductions is given. Moreover the lattices occurring in the congruences have weights and ranks increasing with the main parameter x. An error term depending exponentially on a rank, or even a loss of order b on every line of a rank r lattice at every small prime, would not suffice. This section formulates the local integral problem, explains the Kummer-chart calculation used to solve it, and shows exactly how the resulting loss enters the global congruence budget. The integral period-ring assertions quoted here as *formal-source inputs* are statements of the compact Lean development, not assertions for which we claim an independent verification of the proof. In contrast, we give the length, valuation, and budget deductions explicitly.

49.1 Lattices, crystalline cocycles and imperfectly dual reductions

Write $p = \ell$ for the rational prime throughout the local discussion, $G_p = \mathrm{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$, and let $j : \Gamma \hookrightarrow G_p$ identify Γ with an open subgroup. Write F for the corresponding finite extension of $\mathbb{Q}_p$ and $d_F = [F : \mathbb{Q}_p]$. The coefficient ring O denotes the integers of a sufficiently large finite extension of $\mathbb{Q}_p$, with fraction field K_O. For its uniformizer π_O, put $e_O = v_{\pi_O}(p)$; our valuation v_p on K_O has $v_p(p) = 1$. In particular the normalized length of a torsion O-module M is

$$\mathrm{len}_p M = e_O^{-1} \,\mathrm{length}_O M.$$

It is additive on short exact sequences and $\mathrm{len}_p(O/p^aO) = a$. Infinite lengths can be understood as $+\infty$. For submodules S, T of a common module, we use the notation

$$\mathrm{rl}(S,T) = \mathrm{length}_O(S/(S \cap T)), \qquad \mathrm{rl}_p(S,T) = e_O^{-1}\,\mathrm{rl}(S,T).$$

These correspond to `relLength` and `CoeffRing.normLength` in the formal source. The normalization is essential: passing to a more ramified coefficient field must not artificially enlarge the estimates. For a finite-length module over O, its length over $\mathbb{Z}_p$ is $[O/\pi_O : \mathbb{F}_p]$ times its length over O. Consequently

$$\mathrm{rl}_p(S,T) = \frac{\mathrm{length}_{\mathbb{Z}_p}(S/(S\cap T))}{[K_O:\mathbb{Q}_p]}. \tag{49.1}$$

The statements below are uniform in O once the necessary finite coefficient fields and roots of unity are included. This is the meaning of the eventual quantifier on coefficient rings in the source.

A Galois lattice T is finite free over O, with continuous O-linear Γ-action. Put $T^*(1) = \mathrm{Hom}_O(T, O(1))$, where the cyclotomic character acts on $O(1)$. The jump convention is that $O(1)$ has jump -1. At an embedding $\gamma : O \to \mathbb{C}_p$ let $j_T(\gamma, u)$ denote the multiplicity of jump $u \in \mathbb{Z}$; thus

$$j_{T^*(1)}(\gamma, -u-1) = j_T(\gamma, u). \tag{49.2}$$

The local development defines $j_T(\gamma,u)$ by the dimension over $\overline{\mathbb{Q}}_p$ of the space of γ-semilinear functionals $f : T \to \mathbb{C}_p$ satisfying $f(gv) = \chi_{\mathrm{cyc}}(g)^{-u} g f(v)$ for g sufficiently close to 1. In particular its weights are Hodge–Tate weights in the above sign convention, not the oppositely signed integers sometimes used for cyclotomic twists.

The O-module of crystalline cocycles $Z_f(T)$ consists of maps $c : \Gamma \to T$ such that there is $y \in B_{\mathrm{cris}} \otimes_{\mathbb{Z}_p} T$ with

$$1 \otimes c(g) = gy - y \qquad (g \in \Gamma).$$

The tensor product here is over $\mathbb{Z}_p$, and the action is diagonal. The definition is useful also for potentially crystalline T. This is the inverse image in $Z^1(\Gamma, T)$ of the usual kernel of the map to $H^1(\Gamma, B_{\mathrm{cris}} \otimes_{\mathbb{Q}_p} T[1/p])$. The cocycle identity follows by applying $gh - 1 = (g-1) + g(h-1)$ to y and using injectivity of the natural inclusion. The module includes all integral coboundaries. It is saturated with respect to multiplication by a nonzero element of O: if ac is a crystalline cocycle, divide its period primitive by a in the tensor product with B_{cris}, where K_O acts and a is invertible. In particular

$$Z_f(T)/p^a Z_f(T) \hookrightarrow Z^1(\Gamma, T/p^a T). \tag{49.3}$$

The formal source also proves continuity, including the eventual local constancy of reductions of such cocycles. The relevant definitions and results are `crysCocycles`, `jump` and `HasJumps` in `Lib/PadicHodge/Crystalline/Defs.lean`, `crysCocycles.of_smul` in `Lib/PadicHodge/Crystalline/Jumps.lean`, and `crysCocycles.eventually_eq_pow_smul` in `Lib/PadicHodge/BlochKato/CocycleContinuity.lean`.

Proposition 49.1 (Finite local cocycle rank)**.** Formal-source input. *Let T be potentially crystalline. Then $Z_f(T)$ is finite free over O. Rank over O is at most $(1 + d_F)\,\mathrm{rk}_O\, T$. More precisely, in the situation where the jumps are indexed by embeddings of the completion of E in $\overline{\mathbb{Q}}_p$ and hence by a set Σ_v of complex embeddings of E determining a finite place $v \mid p$,*

$$\mathrm{rk}_O\, Z_f(T) = \mathrm{rk}_O\, T + \sum_{\sigma\in\Sigma_v} \#\{u \in J_T(\sigma) : u < 0\}.$$

The same formula with a local field of degree d_F and with summation over its embeddings applies after extending coefficients so that all the Hodge jumps can be read there.

This is the Bloch–Kato rank calculation on cocycles (as opposed to on classes), expressed by `crysCocycles.finrank_eq` in `Lib/PadicHodge/BlochKato/BlochKato.lean` and the

embedding form `finrank_crysCocycles_eq_add_neg_jumps` in `New/Selmer/CrystallineRank.lean`. The rank formula on classes comes from the fundamental period-ring exact sequence and the de Rham filtration; adding the dimension of the coboundaries cancels the contribution of invariants. The formal-source statement on $\mathbb{Z}_p$-ranks uses d_F times the number of negative jumps in the multiset of a restriction of scalars. In the indicated embedding setting, decomposing that multiset and dividing by $[K_O : \mathbb{Q}_p]$ gives the asserted formula. Freeness follows from torsion-freeness and finite generation over a DVR. In particular the rank bound requires no extra hypothesis on inertial eigenvalues.

We describe the class of lattices against which the defect is measured. Let U be a family of finite subsets of $\mathbb{Z}$ indexed by coefficient embeddings. The notation $\mathcal{Q}(F_0, \mathcal{K}, U)$ will mean the class `Lift.InClass` of the formal source; here F_0 is the filter on Γ that specifies the required fixed crystalline periods, and $\mathcal{K}(g, Q)$ specifies a family of polynomials annihilating the operator on potentially crystalline periods associated with g. More explicitly a member Y of the class is admissible for periods fixed almost everywhere for the supremum of the near-identity filter and F_0, has $j_Y(\gamma, u) = 0$ for $u \notin U(\gamma)$, and, for g mapping to inertia, each polynomial Q with $\mathcal{K}(g, Q)$ annihilates the linear operator $\beta_{g,0}$ on $B_{\mathrm{cris}} \otimes_{\mathbb{Z}_p} Y$. This operator is the B_{cris}-linearization of g on the potentially crystalline periods (the source calls it `linearFrob`). For $F_0 = \top$, $\mathcal{K} = \bot$ we just require crystallinity and the specified weights. All members are potentially crystalline. The jump and polynomial conditions are preserved when a lattice is a subrepresentation of a direct sum of the given representations. The dual of such a class has jump sets $-U - 1$ and inertial polynomial condition at inverse elements. These facts are expressed in `InClass.of_jointly_injective`, `InClass.dual` and `InClass.tensor`, `Lib/PadicHodge/Crystalline/PairingDefs.lean`.

An imperfectly dual pairing of lattices Y, T is specified by a positive modulus p^a, an O-linear form $\psi : Y \otimes_O T \to O(1)$, a $\mathbb{Z}_p$-linear functional $\lambda : O \to \mathbb{Z}_p$, and a crystalline cocycle $q \in Z_f((Y \otimes T)^*(1))$ satisfying

$$g\psi - \psi = p^a q(g). \tag{49.4}$$

Thus the induced bilinear form $\bar{\psi}$ with values in $\mathbb{Z}/p^a(1)$, obtained by applying λ and reducing modulo p^a, is equivariant; the unreduced pairing need not be equivariant. The cup product of $c \in Z_f(Y)$ and $c' \in Z_f(T)$ under $\bar{\psi}$ is represented by

$$(g, h) \longmapsto \bar{\psi}(c(g), gc'(h)) \in \mathbb{Z}/p^a(1).$$

The vanishing required in the source means that, after the inclusion of the cyclotomic module $\mathbb{Z}/p^a(1)$ into $\overline{\mathbb{Q}}_p^{\times}$, the displayed cocycle is a continuous coboundary. One should not replace this assertion by an assertion of vanishing in $H^2(\Gamma, \mathbb{Z}/p^a(1))$ without an argument. The chosen codomain is the one used by the index form of local Tate duality below. Both the imperfect equivariance and the third class q are necessary: it would be illegitimate simply to apply the pairing of $D_{\mathrm{cris}}(T)$ with its exact dual to a representation merely congruent modulo p^a to that dual.

Define the submodule $T^{\perp_{\mathcal{Q}}}$ of maps $c' : \Gamma \to T$ by imposing this cup-product vanishing for every $Y \in \mathcal{Q}$, every $c \in Z_f(Y)$, and every pairing of the above form, including every modulus. Define its crystalline defect

$$\Delta_p(T; \mathcal{Q}) = \mathrm{rl}_p(Z_f(T), T^{\perp_{\mathcal{Q}}}).$$

We use the symbol a in pairing modulus p^a only in this local definition: in estimates below it is the inherited global derivative-order parameter unless specifically renamed. The Lean definitions of pairings, their cup product, and the orthogonal submodule are, respectively, `Lift.Pairing`, `Pairing.CupBounds`, and `cupOrthogonal` in `Lib/PadicHodge/Herr/GaloisLattices.lean`.

Lemma 49.2 (Exponent-to-length conversion)**.** *If $p^s Z_f(T) \subseteq T^{\perp_{\mathcal{Q}}}$, then*

$$\Delta_p(T;\mathcal{Q}) \le 2d_F\, s\, \mathrm{rk}_O\, T.$$

Proof. There is a surjection from $Z_f(T)/p^s Z_f(T)$ onto the defect quotient, because its denominator contains $p^s Z_f(T)$. The former has normalized length $s\,\mathrm{rk}_O\, Z_f(T)$. Proposition 49.1 yields $s(1+d_F)\,\mathrm{rk}_O\, T \le 2sd_F\,\mathrm{rk}_O\, T$, since $d_F \ge 1$. This is precisely the numerical content of `normLength_defect_le` in `New/Selmer/CrystallineRank.lean`. □

49.2 Why this defect bounds the local discrepancy of congruent cocycles

Take a finite list of global integral lattice representations $u_i : G \to \mathrm{GL}_{n'}(O)$ and a block representation $\rho : G \to \mathrm{GL}_{r'}(O)$. Fix a modulus p^M and write $R = O/(p^M)$. The global image of the reducibility ideal gives, on the block, cocycles $c : G \to R^{r'}$ satisfying not only the cocycle identity, but also all linear relations among the matrices $(u_i(g)-1)_i$. Let $\mathcal{Z}$ denote the module of cocycles satisfying these relations. Call its local crystalline-property submodule at $v \mid p$ the intersection with the module of cocycles whose restriction to G_v obeys the relations of $Z_f(T_v)$, where $T_v = (O^{r'}, \rho|_{G_v})$. Here "the relations of a module" $Z \subseteq \prod_g O^{r'}$ are finitely supported O-linear relations vanishing on every element of Z; they are imposed modulo p^M. This is the definition in `obeyingCocycles` and `crysRels`, in `New/Selmer/BlocksAndCocycles.lean` and `New/Selmer/DeterminantBound.lean`. This local condition is stronger than continuity and the cocycle identity, and must not be assumed automatically from the construction of a congruence. The entire point of the local estimate is to bound how far $\mathcal{Z}$ is from imposing it.

Here is the algebraic reason these cocycles are amenable to crystalline estimates. Let P be the finite free module of the tuple of matrices, with the action by left multiplication by $(u_i(g))_i$. Put $D(g) = (u_i(g)-1)_i$ and $A = \sum_g OD(g) \subseteq P$. Then $D(gh) = D(g) + u(g)D(h)$, so A is stable under G. Since O is Noetherian, A is finite. If c obeys the displayed relations then a linear map $q : A \to R^{r'}$ is defined by $q(D(g)) = c(g)$; it is equivariant by the two cocycle identities. By the Artin–Rees lemma, for some j large enough $A \cap p^{Mj}P \subseteq p^M A$. Consequently q extends to a well-defined equivariant map

$$q' : Y = A + p^{Mj}P \longrightarrow R^{r'}, \qquad q'(a + p^{Mj}z) = q(a).$$

Indeed if two such expressions have equal first component then their difference in A belongs to $p^M A$, on which q vanishes. The action on $p^{Mj}P$ is stable. The map satisfies $q'(D(g)) = c(g)$, and Y is a finite free lattice in P. Equivalently the graph of q' projects injectively into P. This is the constructive role of the graph subrepresentation in `obeyingCocycles.exists_graph_subrepresentation`. In particular after restriction to G_v, Y belongs to the class $\mathcal{Q}$ of lattices inherited from the u_i, whenever those representations have the requisite admissibility, jump sets and annihilating polynomials. The map $g \mapsto D(g)$ is itself a crystalline cocycle whenever the ambient representations are potentially crystalline: over B_{cris} its primitive is the tuple of identity matrices, and multiplication by p^{Mj} puts this primitive in Y after extending scalars. Thus every obeying cocycle has the local form of a reduction of a linear image of a crystalline cocycle on Y. It need not be a reduction of a crystalline cocycle on T_v.

For the duality step, lift q' to an O-linear map $q_1 : Y \to T_v$ (possible by freeness). The reduction is equivariant, so $gq_1 - q_1 = p^M q_2(g)$ with $q_2(g)$ integral. Under the perfect pairing $T_v \otimes T_v^*(1) \to O(1)$ this is exactly (49.4) on $Y \otimes T_v^*(1)$. Its third class q_2 is crystalline: $p^M q_2$ is a coboundary, and saturation shows that q_2 lies in $Z_f((Y \otimes T_v^*(1))^*(1))$. Hence every reduction in $\mathcal{Z}|_{G_v}$ annihilates the reduction of the orthogonal module $(T_v^*(1))^{\perp_{\mathcal{Q}}}$ under the local Tate

pairing. The same conclusion holds for reductions from $Z_f(T_v)$ itself, since $T_v \in \mathcal{Q}$ and the usual perfect pairing is permitted.

Proposition 49.3 (Local congruence loss)**.** The local Tate-duality index estimate, derived in Lemma 50.2 under the classical duality and crystalline orthogonality hypotheses. *Under the preceding hypotheses,*

$$\mathrm{rl}_p\left(\mathcal{Z}, \{c : c|_{G_v} \text{ obeys all relations of } Z_f(T_v)\}\right) \le \Delta_p(T_v^*(1); \mathcal{Q}).$$

For clarity, the content of the index estimate is the following. A finite module $V = T_v/p^M T_v$ has the Cartier dual provided by the reduction of the $\mathbb{Z}_p$-dual Tate lattice. A $\mathbb{Z}_p$-linear functional on O can be chosen to make the pairing perfect: the $\mathbb{Z}_p$-dual of O is a rank-one O-module (the inverse different). The images A and A' of $Z_f(T_v)$ and $Z_f(T_v^*(1))$ have orders whose product is

$$p^{Md(2+d_F)}, \qquad d = \mathrm{rk}_{\mathbb{Z}_p} T_v. \tag{49.5}$$

Indeed, by (49.3) their orders are the powers of p with exponents M times their $\mathbb{Z}_p$-ranks. The dual jump relationship (49.2) pairs each nonnegative jump of T_v with a negative jump of its dual; their ranks therefore sum to $(2 + d_F)d$. The size bound from local Tate duality says that the number of extra classes of the restriction $\mathcal{Z}|_{G_v}$ modulo its intersection with A divides the index of the orthogonal part of $Z_f(T_v^*(1))$. It is applied at the cocycle level, not after omitting the coboundaries; the count (49.5) includes those terms. Every member of A satisfies the indicated crystalline relations, so passing from the condition of belonging to A to that of satisfying its relations can only decrease the loss. Pulling the resulting estimate back along restriction proves the inequality from the cited index estimate. The exact formal implementation is `IsPotCrys.relIndex_dvd_of_orthogonal` in `Lib/ClassField/LocalDuality/CrystallineDuality.lean` and `relLength_obeyingCocycles_le` in `New/Selmer/RowBalanced.lean`. A separate proof of the former would require developing the local Tate pairing and its index formula; here we have isolated where that input is used. Note especially that the right side is independent of M. This uniformity is what makes the first-order global congruence construction effective.

49.3 The Kummer chart, letter decomposition and height of a frame

We continue the local discussion, writing p for ℓ and reserving E without an additional subscript for the global CM field. In particular, the Eisenstein element in the chart will be denoted by $\mathfrak{e}$ (the Lean notation for it is E). Let $A_{\mathrm{inf}} = W(\mathcal{O}_{\mathbb{C}_p^\flat})$ and B_{dR}^+ be Fontaine's usual period rings, with the map $\vartheta : A_{\mathrm{inf}} \to \mathcal{O}_{\mathbb{C}_p}$ and Frobenius φ on A_{inf}. Write t_{dR} for the uniformizing parameter of B_{dR}^+, so $\mathrm{Fil}^n B_{\mathrm{dR}} = t_{\mathrm{dR}}^n B_{\mathrm{dR}}^+$ for $n \in \mathbb{Z}$. This avoids confusion with the rank difference $t = 5r$ of the global proof.

The local chart fixes an unramified coefficient ring $\Lambda = W(\mathbb{F}_{p^{f_0}})$, a ring $\mathfrak{S} \cong \Lambda[[z]]$ embedded in A_{inf}, and an element

$$\mathfrak{e} = z^{e_0} A + p \quad (A \in \mathfrak{S}^\times)$$

associated with a generator of $\ker \vartheta$. Here z is the Teichmüller lift of the element of the tilt arising from a tower of roots of a local uniformizer. For the completion-group case, $e_0 f_0 = d_F$. The chart also contains an open subgroup Γ' of index prime to p in Γ, preserving the indicated constants and the untilt of z, with the required transitivity on roots. Thus local cup product vanishing may be tested after restriction to Γ', since restriction followed by

corestriction multiplies by the prime-to-p index, and the relevant classes are p-power torsion. The unramified chart has $e_0 = 1$, $\mathfrak{e} = z + p$. For the quantitative depth steps the relevant number is not the full discriminant of the global CM field but

$$\delta_0 = \begin{cases} 0, & e_0 = 1, \\ v_p(e_0) + 1, & e_0 > 1. \end{cases} \tag{49.6}$$

In particular δ_0 is a specified upper bound used in the derivative estimate of the chart and is not the actual different exponent in an arbitrary normalization. The existence of such a chart in the completion-group case, including the subgroup and the compatibility when the extension is unramified, is a *formal-source input*, `exists_kummer_chart` in `Lib/PadicHodge/WachKisin/KummerChartExists.lean`. See also the definition of `Lift.Chart` in `Lib/PadicHodge/WachKisin/HodgeFrame.lean`.

Let $q_0 = p^{f_0} - 1$, and choose a primitive q_0-th root of unity ζ in O (hence the initial enlargement of the coefficient ring). The source indexes the full set of units of $\mathbb{F}_{p^{f_0}}$ by the term *letters*; it is convenient to write $I = \mathbb{F}_{p^{f_0}}^{\times}$. A letter $i \in I$ indexes the summand of the Witt-vector ambient module on which multiplication by ζ on the lattice factor agrees with multiplication by the Teichmüller root $\omega(i)$ on the Witt-vector factor. The projector to this summand on $\mathcal{V}_T = W(\mathbb{C}_p^{\flat}) \otimes_{\mathbb{Z}_p} T$ is

$$e_i = \frac{1}{q_0} \sum_{j=0}^{q_0 - 1} \left(\omega(i)^{-1} \zeta \right)^j . \tag{49.7}$$

Because q_0 is prime to p this operator is integral. Orthogonality and idempotence follow by summation of a geometric progression over distinct roots of unity; $\omega(i) - \omega(j)$ is a unit for $i \neq j$. The sum of the projectors is the identity, and Frobenius sends the summand i to the summand i^p. Denote the permutation $i \mapsto i^p$ by σ_0 and put $i[-a] = \sigma_0^{-a} i$ for $a \geq 0$. The coefficient embeddings of O belonging to letter i are the $\gamma : O \to \mathbb{C}_p$ with $\gamma(\zeta) = \vartheta(\omega(i))$. Define

$$U_i = \{0\} \cup \bigcup_{\gamma(\zeta) = \vartheta(\omega(i))} U(\gamma), \qquad m_T(i, u) = \sum_{\gamma(\zeta) = \vartheta(\omega(i))} j_T(\gamma, u).$$

The zero in U_i permits crystalline extensions by a trivial lattice. The sum of $m_T(i, u)$ over all u and all i is $[K_O : \mathbb{Q}_p] \operatorname{rk}_O T$. In the arguments below it matters that the sum is over letters, whereas f_0 measures the size of the residue field extension: these are different integers. The projector formula, its Frobenius compatibility, and the definitions of the jumps attached to a letter are `letterProj`, `phi_pow_smul_mem_letterPart_iff`, and `jumpMultiplicity` in `Lib/PadicHodge/WachGamma/KisinModuleDefs.lean` and `Lib/PadicHodge/WachGamma/Letters.lean`; the enlarged weight sets are `KummerChart.letterJumps` in `New/Lift/ExponentTheorems.lean`.

We explain the choice of integral lattice used for this calculation. The source works with $W(\mathbb{C}_p^{\flat})$ over A_{inf} and an auxiliary element $\tau = [\epsilon^{1/p}] - 1$ of A_{inf}, where ϵ is the compatible tower of p-power roots of unity. Put $A_T = A_{\mathrm{inf}} \otimes_{\mathbb{Z}_p} T$ inside $\mathcal{V}_T$. An invariant form on T is an A_{inf}-linear functional $\mu : \mathcal{V}_T \to B_{\mathrm{dR}}$ satisfying $\mu(1 \otimes gv) = g\mu(1 \otimes v)$ for $g \in \Gamma'$; write $\mathcal{H}_T$ for the family of such forms. Let M_T be the source's submodule `kisinAinf`, consisting of elements in the τ-saturation of A_T such that

$$\mu(\varphi^a v) \in B_{\mathrm{dR}}^{+} \quad (\mu \in \mathcal{H}_T,\ a \geq 1).$$

For a more intrinsic way of specifying the finite-height module $\mathfrak{N}_T$ in M_T, impose the additional value-field condition in the source: $\mu(v)$ must belong to the chart's value field for all invariant

forms. This defines `kisinModule`; the value-fields and their Frobenius transforms are as in `Lib/PadicHodge/WachKisin/DeRhamValues.lean`. Let $M_T^{\mathrm{loc}} = M_T[1/\mathfrak{e}_0, 1/\mathfrak{e}_1, \ldots]$ inside $\mathcal{V}_T$, where $\mathfrak{e}_a = \varphi^a \mathfrak{e}$, and let $\mathfrak{N}_T^{\mathrm{loc}}$ be its points satisfying the same value-field condition. We do not identify the latter with an arbitrary scalar extension of an abstract Breuil–Kisin module; the proof uses the above integral model and its retraction. In particular it is important that the period-frame relations are preserved by certain $\mathfrak{S}$-linear retractions on its scalar coefficients.

Proposition 49.4 (Integral frame package)**.** Formal-source input. *Let T be admissible for the chart and let H_T bound the absolute values of its jumps. Then*

$$M_T = A_{\mathrm{inf}}\mathfrak{N}_T, \qquad \tau^{H_T} M_T \subset A_T, \quad \tau^{H_T} A_T \subset M_T, \quad \mathfrak{e}^{H_T}\varphi(\mathfrak{N}_T) \subset \mathfrak{N}_T.$$

Relations among value-field points remain valid after applying an $\mathfrak{S}$-linear map $A_{\mathrm{inf}} \to \mathfrak{S}$ to the coefficients. Moreover let $l, A, \delta \geq 0$ with $3e_0 \leq p^l$, $e_0 A \leq p^l \delta$. Then the value-field points of $\varphi^l(M_T)$, after multiplication by $p^{4\delta}$, are spanned over $\mathfrak{S}$ by value-field points invariant modulo $\mathrm{Fil}^A B_{\mathrm{dR}}$ under all Γ' when tested by all invariant forms. The invariant crystalline forms also test divisibilities in the frame.

These assertions are components of a single theorem, `Lift.KummerChart.JumpLattice.frame_theorem`, in `Lib/PadicHodge/WachKisin/FrameTheorem.lean`, with preparation in `PeriodMatrixFrame.lean`. The underlying results concern Breuil–Kisin lattices and comparison periods, but the last estimate is used with varying l, A, δ; it is not enough to cite finite height or crystalline comparison with unspecified constants. In the formal proof the period matrix is used to compare the natural A_{inf}-lattice with its Frobenius transports, and functionals arising from the invariant period frame bound orders on each Hodge-graded line. The package including the $p^{4\delta}$ estimate is a specific input whose concrete proof would be part of a longer expositional reconstruction. For the current section it will be important to retain this exact exponent. The other uses of the chart involve only the parameters e_0, δ_0 and the Hodge jump sets, rather than the matrix size in the period frame.

49.4 Exponent lattices and the cost of increasing depth

For integer-valued functions $c(a, u)$ on nonnegative integers a and allowed jumps $u \in U_{i[-a]}$, define the exponent lattice $M_T(i, c)$ to be the subset of M_T^{loc} in the letter i such that

$$\mu(\varphi^{-a} v) \in \mathrm{Fil}^{c(a,u)} B_{\mathrm{dR}} \tag{49.8}$$

for every a and every invariant form μ of type u at $i[-a]$. Here the type condition is $\mu(e_{i[-a]}(1 \otimes T)) \subseteq \mathrm{Fil}^{-u} B_{\mathrm{dR}}$. This is `withExponents` in `New/Lift/ExponentLattices.lean`. We call c the exponents of the point and call c' its depth if, for all $g \in \Gamma'$, $(g-1)v \in M_T(i, c')$. Multiplication by $\mathfrak{e}_a$ raises the exponent at a by one, and Frobenius transports indices $i \mapsto i^p$. In particular the exponents of level l are

$$\mathrm{st}_l(a, u) = \begin{cases} 0, & a < l, \\ -u, & a \geq l. \end{cases} \tag{49.9}$$

Larger exponents mean a smaller module. A useful operation on exponents is

$$\mathrm{raised}_s(a, u) = -u + s\mathbf{1}_{a=0}.$$

Thus raised_s increases only the exponent at the original Eisenstein element. Depth is distinguished from the exponents of a point itself; the former can be improved without requiring arbitrarily large orders on the nonzero constants to which the points project.

The basic numerical cost attached to a step (a,n) is

$$c_p(a,n) = v_p(n) + \delta_0 + 3a, \tag{49.10}$$

where we put $v_p(0) = 0$ in this formula only. One point of the construction is that $c_p(a,n)$ is independent of the rank. Its origin is the following assertion underlying the finite-difference operation.

Lemma 49.5 (One-step moving operator)**.** Formal-source input. *For $n \neq 0$, write $\mathfrak{e}_a = \varphi^a\mathfrak{e}$. There exist $g \in \Gamma'$, $\omega \in A_{\mathrm{inf}}^{\times}$ and $\alpha, X \in A_{\mathrm{inf}}$ with*

$$g\mathfrak{e}_a = \omega\mathfrak{e}_a, \quad \mathfrak{e}_a \mid gs - s \ (s \in \mathfrak{S}), \quad z\tau X = \omega^n - 1, \quad \mathfrak{e}_a \mid \alpha X - p^{c_p(a,n)}.$$

If $n = 0$, a sum of p^{δ_0+3a} elements of Γ' has invariant action on the constants modulo $\mathfrak{e}_a$. After the chart retraction these identities supply an additive endomorphism of the localized module raising depth from n to $n+1$, modulo invariant constants, with cost $p^{c_p(a,n)}$ on the quotient of an extension by its kernel.

The identities used here are `Lift.KummerChart.exists_depth_step`, `Lift.KummerChart.exists_depth_step_zero` in `New/Lift/ExponentLattices.lean`, and `exists_reaches_depth_succ` in `New/Lift/HodgeExpansion.lean`. To explain the cancellation for $n \neq 0$, suppose that the leading contribution of an invariant form at the divisor $\mathfrak{e}_a = 0$ is $\mathfrak{e}_a^n h$ modulo higher order, and that h is invariant to order zero. Taking the difference $g-1$ gives $(\omega^n-1)\mathfrak{e}_a^n h$ modulo the next order. The congruence for αX then implies that the combination

$$p^{c_p(a,n)}v - \frac{\alpha}{z\tau}(gv - v)$$

cancels this leading contribution. When $n = 0$, an averaging operator is used instead. The formal identity for the step includes precisely the hypotheses which allow the quotient by $z\tau$ and the application of the coefficient retraction: it is not legitimate to divide by $z\tau$ in an arbitrary integral submodule. The estimate for $c_p(a,n)$ is obtained by lifting a unit under ϑ, the bound for the differentiated Eisenstein equation and the valuation of $\omega^n - 1$. The latter contributes the factor $v_p(n)$ because lifting the exponent in a multiplicative congruence gives a factor dividing the valuation of n. The additional term $3a$ is the allowance for the Frobenius divisor. This explains why subsequent costs consist of sums of valuations of ordinary integers.

Consider an exact sequence of integral lattices $0 \to T \to \widetilde{T} \to C_0 \to 0$ in which C_0 is trivial, described by an O-linear map $\alpha : C_0 \to Z_f(T)$. The underlying O-module of $\widetilde{T}$ is $T \oplus C_0$, with action

$$g(v,c) = (gv + \alpha(c)(g), c).$$

It is crystalline over the chart because each $\alpha(c)$ is a coboundary after extending to B_{cris}. Denote the natural projection by π. In letter i a point over $p^\kappa c$ means $v \in \mathfrak{N}^{\mathrm{loc}}_{\widetilde{T}}$ satisfying $\pi(v) = p^\kappa e_i(1 \otimes c)$ and having prescribed exponents and depth. The relation $\pi(gv - v) = 0$ means that its Galois differences lie in a kernel lattice. Strictness of the kernel for the exponent conditions, namely every point in the kernel of π satisfying the exponent conditions on the extension comes from a point of $M_T(i,c)$ and similarly for localized value-field points, is a *formal-source input*: `exists_inclKer_eq_of_projTriv_eq_zero` in `New/Lift/LongGap.lean`. Its proof uses extension of invariant forms from the sublattice to the extension while retaining their filtration degree, via `exists_extend_form` of the same file.

The following consequence of the moving operators is the useful form for bounds. Let S be a finite set of steps (a,n) containing those required to raise depth from ρ to ρ': for each a,u it contains every pair (a,n) with $\rho(a,u) \leq n < \rho'(a,u)$. Starting with a point over $p^\kappa c$

of depth ρ, one obtains a point over $p^{\kappa+\sum_S c_p(a,n)}c$ of depth ρ'. Its exponents improve to $\rho'(a,u)$ wherever its old exponents were positive, and to at least $\min(0,\rho'(a,u))$ on the other lines. In the precise statement the integers may be raised further by paying more powers of p. This is *formal-source input* `pointsOver_nonempty_of_costs`, `exists_reaches_deep` and `mem_withExponents_of_deep` in `New/Lift/LongGap.lean` and `New/Lift/HodgeExpansion.lean`. The induction is on the largest step index to be increased and its depth. Compose the operators of Lemma 49.5; when operator D has cost p^e, the difference $Dv - p^e v$ belongs to the kernel of π, so the next operator preserves the accumulated scalar. Thus the cost of a composition is the sum of the integer exponents. The crucial difference from a dimension argument is that the same operator raises the specified depth on every line of the pertinent jump, so multiplicities do not multiply this cost. The exponent improvement statement uses that a de Rham value whose Galois differences lie in a deep filtration can be made invariant modulo that filtration by subtracting a constant from the local field. Positivity of the prior exponent forces that constant to be zero; without positivity one retains at least the zero-filtration threshold. These are the precise properties needed in the later cup-product calculation.

We record the elementary valuation estimate controlling these operators. For integers $n \ge 0$ and c, set $I(n,c) = [-n,n] \cup [c-n, c+n]$ (integer intervals). If $n \le pM$ and $|c|+n \le H$ with $H \ge 1$, then for $d \ge 0$,

$$\sum_{j \in I(n,c)} (v_p(j) + d) \le (4M+2)\lfloor \log_p H \rfloor + (4n+2)d. \tag{49.11}$$

Indeed at most $4n+2$ integers occur and, in each interval of $2n+1$ integers, at most $2\lfloor n/p \rfloor + 2$ can be multiples of p. More sharply, an interval of length $2n$ contains at most $\lfloor 2n/p \rfloor + 1 \le 2M+1$ multiples if $n \le pM$; hence the two intervals together contain at most $4M+2$. For any nonzero j of absolute value at most H, $v_p(j) \le \lfloor \log_p H \rfloor$, and zero contributes no valuation by convention. This proves the bound. It is `sum_twoWindows_le` in `New/Lift/CokernelKillers.lean`. For steps with $a = 0$, if p divides no nonzero integer within distance n of $0, \pm c$ then the valuation contribution vanishes; in that case all the steps in these intervals cost only δ_0 each. With the rough count $8n+4$ of their integers the total cost is at most $(8n+4)\delta_0$. The formal-source derived assertion `pointsOver_nonempty_of_noMultipleNear` therefore improves a point at the original divisor at precisely this cost. In an unramified chart the cost is zero. This free improvement is the first of several places where the proof needs a gap in the congruence classes of large weights, not just a gap in their ordinary archimedean sizes.

49.5 Cokernel estimates: divisibility by p and by the Eisenstein element

For an extension side (T, C_0, α) as above let $J_l(i;s)$ be the ideal of A_{inf} defined as the sum of (s) and the ideal of those multipliers which send the letter-i part of $A_{\mathrm{inf}} \otimes C_0$ into the image under π of the extension's exponent lattice with exponents st_l. Thus, for a scalar $u \in A_{\mathrm{inf}}$, the condition $u \in J_l(i;s)$ means that one can choose $v \in A_{\mathrm{inf}}$ such that for every $c \in C_0$ an exponent-st_l point lifts $(u - sv)e_i(1 \otimes c)$. We say that $p^\kappa \mathfrak{e}^s$ kills the cokernel at i if

$$p^\kappa \mathfrak{e}^s \in J_0(i;0).$$

These are `liftIdeal` and `KillsCokernel` in `New/Lift/CokernelKillers.lean`. This definition is a statement about actual multipliers, not just equality after inverting p. Coefficient retraction allows the corresponding lifts to be taken in the localized value-field module. In

particular if $p^{\kappa}\mathfrak{e}^{s}$ kills, after dividing by $\mathfrak{e}^{s}$ one finds a point over $p^{\kappa}c$ having both exponents and depth raised$_{-s}$. This last assertion is *formal-source input* `pointsOver_nonempty_of_killsCokernel` in `New/Lift/LongGap.lean`.

Proposition 49.6 (Cokernel calculus)**.** Formal-source input. *For the ideals just defined, the following are valid.*

(i) *The level and letter are transported by Frobenius: in particular $p^{c}z^{pa} \in J_1(i^{p}; r_0)$ if and only if $p^{c}z^{a} \in J_0(i; r_0)$, for a fixed modulus r_0. If all allowed jumps are $\le n$, multiplication by $\mathfrak{e}^{n}$ sends $J_0(i; r_0)$ into $J_1(i; r_0)$.*

(ii) *If $p^{c}z^{v} \in J_1(i^{p}; r_0)$, $pa \le v$ and $c + v_p(v!) \le \kappa + v_p((pa)!)$, then $p^{\kappa}z^{a} \in J_0(i; r_0)$.*

(iii) *If $p^{c}z^{a}\mathfrak{e}^{n} \in J_1(i; r_0)$, then $p^{c+\delta_0 n + v_p(n!)}z^{a+n} \in J_1(i; r_0)$.*

(iv) *If all jumps on the side are $\le n$, for each $\kappa \ge 0$ there exists $a \ge 0$ with $(p-1)a \le e_0(n+\kappa)$ and $z^{a} \in J_0(i; p^{\kappa})$ for all i.*

(v) *In the unramified chart and for $p \ge 5$, if all jumps are $\le (p-3)(\kappa + s)$ then $p^{\kappa}\mathfrak{e}^{s}$ kills the cokernel at every letter.*

All assertions occur in `New/Lift/CokernelKillers.lean`, respectively `p_pow_mul_z_pow_mem_liftIdeal_iff` and `mul_E_pow_mem_liftIdeal_one`, `mem_liftIdeal_zero_of_mem_one`, `mem_liftIdeal_of_mul_E_pow`, `exists_z_pow_mem_liftIdeal`, and `killsCokernel_of_isUnramified`.

We recall how the factorials arise. The finite-difference calculation from Lemma 49.5, at level zero or one, acts on $J_l(i; r_0)$ by the following rule. If $f \in \mathfrak{S} \cap J_1(i; r_0)$ and there is an identity

$$z\tau\mathfrak{e}(s + \beta f) = \alpha(gf - f), \qquad g \in \Gamma',$$

with the appropriate integral scalar coefficients, then $s \in J_1(i; r_0)$. It follows by taking the difference of integral lifts in the defining multiplier ideal, using a retraction to preserve the membership conditions. This rule is *formal-source input* `mem_liftIdeal_of_derivation`. Applied to $f = p^{c}z^{v}$, the rule lowers v by one up to $p^{v_p(v)}$ in the coefficient. Inducting from v down to pa gives the additive cost $v_p(v!) - v_p((pa)!)$; transporting the remaining power back by Frobenius proves (ii). Applied to $f = p^{c}z^{a}\mathfrak{e}^{n}$, the rule reduces n by one and increases the z-exponent, at step cost $\delta_0 + v_p(n)$. Their sum is $\delta_0 n + v_p(n!)$, giving (iii). These summations are independent of any rank. In the unramified case the polynomial identity $\mathfrak{e} = z + p$ permits one to combine the multipliers of (ii) with a binomial expansion of $\mathfrak{e}^{s}$. The resulting containment modulo $p^{\kappa+s+1}$ is upgraded to an actual containment by the elementary Nakayama argument: if $p^{c} \in J_0(i; 0) + (p^{c+1})$ then $p^{c}(1 - pq) \in J_0(i; 0)$ for some $q \in A_{\mathrm{inf}}$, and $1 - pq$ is invertible. Thus $p^{c} \in J_0(i; 0)$ and adding the higher modulus does not enlarge $J_0(i; 0)$. The quantitative minimum on a in (iv) and binomial comparison in (v) are the specific remaining cokernel estimates; they cannot be replaced by an unquantified assertion that some power of p annihilates the cokernel.

49.6 Cup-product vanishing from the three sides

Each pairing in (49.4) introduces three crystalline extension sides: the side Y carrying c, the side T carrying c', and the side $(Y \otimes T)^{*}(1)$ carrying the third class q. On the last side the quotient lattice is O with extension determined by q, and the allowed jumps, including zero,

are the numbers $-u-v-1$ with u and v running over the jumps on the first two sides. The integral equivariant pairing of this third extension with $Y \otimes T$ is

$$((\xi, s), u \otimes v) \longmapsto (p^M \xi + s\psi)(u \otimes v),$$

where p^M is the pairing modulus. Its equivariance follows directly from (49.4), since the extra term $p^M q(g)$ is exactly the change of ψ. On reduction it agrees with the imperfectly dual pairing. This is why controlling the first two extensions is not enough: the period computation must also supply a lift of a constant in the third extension.

Over the tilt consider $\mathcal{M} = W(\mathbb{C}_p^\flat) \otimes \mathbb{Z}/p^M(1)$ and let $\mathcal{M}_+$ be the image of $W(\mathfrak{m}_{\mathbb{C}_p^\flat})$. Call a pairing of Witt-vector components safe with cost κ if p^κ times its value lies in $\mathcal{M}_+$. The submodule $\mathcal{M}_+$ is stable under Frobenius and Γ. Different letters are orthogonal: for example if two vectors have labels i, j, then multiplying their value by $\omega(i)$ or $\omega(j)$ gives the same result, since the pairing is O-bilinear before applying the functional and reducing. For $i \neq j$ their difference is a unit. This is also true of the sum of their pairings, as required by `isSafe_sum`.

Here is the algebra of the safe-pair criterion. We use the elementary Artin–Schreier–Witt property of $\mathcal{M}$ with its small part: $\varphi - 1$ is surjective; if $(\varphi-1)z \in \mathcal{M}_+$ then z is congruent modulo $\mathcal{M}_+$ to a unique element of the constant module $\mathbb{Z}/p^M(1)$; and the intersection of the constants with $\mathcal{M}_+$ is zero. This property, including the convergence for Galois differences needed to get continuous cochains, is a *formal-source input*, `TateTwist.smallPart_spec` in `Lib/PadicHodge/WachGamma/KisinModuleDefs.lean`.

Suppose x, w in the first extension lie over the same constant $p^{e_1}c_0$, and y in the second lies over $p^{e_2}c_0'$. Write X, W, Y for their respective first components in the Witt-vector tensor modules of the kernels. Write a_g, b_g for the crystalline cocycles on these kernels, scaled by p^{e_1} and p^{e_2}. Thus the first component of $(g-1)x$ equals $D_g = gX - X + a_g$, that of $(g-1)w$ equals $D_g^W = gW - W + a_g$, and that of $(g-1)y$ equals $T_g = gY - Y + b_g$. Put $C = \varphi Y - Y$. Denote the pairing multiplied by p^{e_3} by P. We assume the following four expressions are zero in $\mathcal{M}/\mathcal{M}_+$, for all $g, h \in \Gamma'$:

$$P(D_g, gT_h), \quad P(W - X, T_g), \quad P(D_g^W, gC), \quad P(\varphi X - W, g\varphi Y - \varphi Y + b_g). \tag{49.12}$$

These are precisely the four safe-pair hypotheses in `Pairing.ScaledCupBounds.of_isSafe`, `New/Lift/CupProductBound.lean`. The last second argument is the first component of $(g-1)\varphi y$. Since the scaled cocycle values are constants for Frobenius it is also φT_g. The goal is to see that $P(a_g, gb_h)$ is a coboundary.

First define $k_0(g) = P(a_g, gY) - P(X, T_g)$. The graded Leibniz rule for the group-cohomology differential, which can equally be checked by expansion in degree one, gives

$$\mathrm{d}k_0(g, h) = P(a_g, gb_h) - P(D_g, gT_h).$$

This equality can be read in the quotient because P is equivariant by construction. By the first safety condition the last term is zero. It remains to choose the cochain with values in constants, rather than just the large Witt-vector quotient. Choose Z with $(\varphi-1)Z = P(W, C)$ by the Artin–Schreier–Witt property and set $k(g) = k_0(g) + gZ - Z$. In the quotient, direct expansion gives

$$\begin{aligned}(\varphi-1)k(g) &= P(a_g, gC) - P(\varphi X, gC - C)\\ &\quad - P(\varphi X - X, T_g) + P(gW, gC) - P(W, C)\\ &= P(D_g^W, gC) - P(W - X, T_g) - P(\varphi X - W, \varphi T_g).\end{aligned}$$

This is zero by the remaining three conditions. Hence each $k(g)$ is congruent to a unique constant $\kappa(g)$. The differential of k equals that of k_0, and $P(a_g, gb_h)$ is a constant. Injectivity of the constants into the quotient implies that $\mathrm{d}\kappa(g,h) = P(a_g, gb_h)$ as an identity of constants. Continuity of κ is part of the preceding formal-source small-part estimate: the lift formula involves finite congruence data, and sufficiently small changes under the Galois action contribute to the small part, so the unique constant representative is locally constant. This proves vanishing after restriction to the chart group Γ', and then in the cohomology group on Γ by corestriction, since its index is prime to p. We may further increase the total exponent $e_1 + e_2 + e_3$. This gives the claimed scaled cup product, and is the mechanism behind the formal theorem `Pairing.ScaledCupBounds.of_isSafe`. Notice that the third condition involving w is not redundant: it provides precisely the correction to make the cobounding cochain Frobenius-invariant modulo the small part.

The remaining challenge is to produce safe points with sufficiently small exponents e_1, e_2, e_3. The main safety test is a strict filtration inequality. Suppose v, w in the two kernels lie in exponent lattices of exponents $o(a,u), o'(a,v)$. Suppose that in the third extension one has a point over p^κ at the predecessor letter $i[-l]$, with exponents $o_2(a,u'')$, antitone as a function of the jump. The required numerical inequality is

$$o(a,u) + o'(a,v) + \mathbf{1}_{a \geq l} o_2(a-l, -u-v-1) > 0 \tag{49.13}$$

for each a and all the jumps u, v at $i[-a]$. Under these hypotheses multiplying either input by $z\tau$ makes its pairing safe with cost κ. The corresponding safe-pair deduction and cochain calculation are treated in Chapter 51; the source declaration is `SidePairing.isSafe_of_exponents` in `New/Lift/LongGap.lean`. It uses the equivariant third pairing above and the rule that multiplication of filtered values adds exponents. The target line $O(1)$ has jump -1; applying the frame comparison for that line produces the factor τ, and the positive valuation of z puts the result in $W(\mathfrak{m}_{\mathbb{C}_p^\flat})$. The strict inequality is indispensable, as safe is a condition of vanishing modulo the maximal ideal of the tilt, not merely integrality. A Galois difference of a point above a constant projects to zero and is divisible by the auxiliary $z\tau$ in the required localized kernel; this is how the depth inequalities for x, w, y imply (49.12).

One uses cuts on Frobenius cycles of letters. Let $I_0 \subset I$ be a subset meeting each cycle, $l(i)$ the distance to the first strictly preceding cut, and choose one leading side at each cut. The point at every other letter is transported from a preceding cut by Frobenius; at a cut one may choose directly constructed points on the leading side. The complementary choice gives two systems of points such that the Frobenius of one system is the other. More concretely the integer shifts are $d_P(i) = 0$ if i is a cut at which the side leads, and $d_P(i) = l(i)$ otherwise. For the complementary systems the identity

$$d_{\mathrm{lag}}(i) = d_{\mathrm{lead}}(i[-1]) + 1$$

uses minimality of the preceding cut. Consequently summing $\varphi^{d_P(i)}$ of the chosen points at $i[-d_P(i)]$ yields the points in the previous safe-pair criterion. Orthogonality of different letter summands reduces all four tests to the separate inequalities (49.13), with appropriate depth and minimum exponents of differences of points at a cut. The precise assertion with the additional depth inequality for the leading side is *formal-source input* `SidePairing.scaledCupBounds_of_points_at_cuts` in `New/Lift/LongGap.lean`; it concludes that an exponent $e_1 + e_2 + \kappa$ is sufficient. The optimization of the cut selection matters, because a Frobenius walk of length l introduces new filtration inequalities at all preceding divisors $\mathfrak{e}_a$.

49.7 The window estimates: the three ways to obtain safe points

For a sign $\epsilon \in \{1, -1\}$ put

$$I(\eta, \epsilon, b, W_0) = [-\eta, \eta] \ \cup \ \epsilon[b, b + W_0],$$

and write $\mathrm{NM}(p, n, b)$ for the condition that no nonzero integer within distance n of $0, b, -b$ is divisible by p. These two definitions are `IsInJumpWindow` and `NoMultipleNear` in `New/Lift/CokernelKillers.lean`. The estimates use opposite signs for the far windows of paired sides. The third extension has many possible jump sums, but for two opposite windows the sum of any pair of large jumps is near zero if both are far, and near a far window if precisely one is far. This is the basic reason that the window structure is stable enough for (49.13) to be useful.

Sparse thick letters. First consider opposite windows $I(\eta, \mp\epsilon_i, \beta_i, V)$ at letter i, where $\beta_i = b$ for thick letters and $\beta_i = G$ for thin ones, and $G + V \le \eta$. Let $W_0 \ge V + 4\eta$, and suppose that either all the signs are constant or the chart is unramified and $p \ge 7$. For an integer $J \ge 0$ assume that there cannot be three thick letters $i, i^{p^s}, i^{p^{s+t}}$ with $1 \le s, t \le J$. The following full set of size hypotheses is sufficient:

$$8e_0 \le p^{J+1}, \qquad 2e_0 b \le p^{J+1} X,$$
$$11(p^J + 1)W_0 \le b, \qquad 4(J + 1 + \delta_0 + \lfloor \log_p(2b) \rfloor)W_0 \le X, \qquad \eta \ge 1.$$

Under them every cup product of crystalline cocycles of the paired sides is killed, in the sense of (49.12), by $p^{10^4 X(J+1)}$. This is *formal-source input* `SidePairing.scaledCupBounds_sparse_thick` in `New/Lift/SparseThickLetters.lean`. One must use these hypotheses on both sides, not just on the distinguished block.

The combinatorial operation in its proof is concrete. Define a cut at i if the next J letters $i^p, \dots, i^{p^J}$ are thin. On a Frobenius orbit every letter has a preceding cut at distance at most $2J + 1$. Indeed, for $J = 0$ all letters are cuts. For $J > 0$, suppose every one of the preceding $2J + 1$ positions fails to be a cut. Failure at distance $J + 1$ yields a thick letter at distance a with $1 \le a \le J$. Failure at distance $a + J + 1 \le 2J + 1$ yields a thick one at distance $a + b$, $1 \le b \le J$. Failure at distance a yields a third thick letter at forward distance c, $1 \le c \le J$, from the first. This is a forbidden triple with gaps b, c. The formal implementation of this selection is the first part of `scaledCupBounds_sparse_thick`; the cut definition also gives the short thin run used in the bound on $J_0(i; s)$ of each extension. Along the run, multiplying by a suitable power of $\mathfrak{e}$ and transporting under Frobenius changes the required exponent of z by a factor p per step. Proposition 49.6(ii),(iii) gives the extra factorial and different costs at each stage. The number X in the hypotheses bounds those costs. The starting powers of z are obtained from Proposition 49.6(iv). At cuts the resulting points can improve their depth in the two windows, paying the bound (49.11) and summing over at most a bounded multiple of $J + 1$ steps. With budget κ satisfying $12(J + 1)X < \kappa$, the source obtains the points over $p^{25\kappa}$ on both sides and a third point over p^{κ}. Thus 51κ pays for the safe-pair criterion. Taking κ of order $(J + 1)X$ and allowing ample slack gives the displayed exponent $10^4 X(J + 1)$. The general cut lemma specifying both the permissible sign pattern and the gap of the large weight is `SidePairing.scaledCupBounds_of_cut_windows`; it is a further formal-source input. This is one of the technically new results needed for the proof, not a standard property of p-adic representations cited without the quantitative bound.

Two sizes over an unramified chart. The second general regime assumes $p \geq 7$ and $e_0 = 1$, $\mathfrak{e} = z + p$. The windows are again $I(\eta, \pm\epsilon_i, \beta_i, V)$ with $\beta_i = b$ or G, and $W_0 \geq V + 4\eta$, $\eta \geq 1$. If

$$5W_0 \leq G \leq b, \qquad b < Kp^2, \quad 2G < Kp, \quad W_0 \leq K,$$

then the exponent

$$10^4 K(\lfloor \log_p(2b) \rfloor + 1) \tag{49.14}$$

annihilates all imperfect cup products. This is *formal-source input* `SidePairing.scaledCupBounds_two_sizes` in `New/Lift/SparseThickLetters.lean`. It is used when p is large and a valuation budget is available, regardless of the sparsity of far-apart letters. Here the cut choice uses one Frobenius step and at i an additional order

$$s_i = \begin{cases} \max(0, \lfloor b/p \rfloor - 7K), & i^p \text{ thick}, \\ 0, & \text{otherwise}. \end{cases}$$

It is a cut if $s_i < \beta_i$. Some cuts require $\mathfrak{e}^{s_i}$ in the cokernel bound, rather than only p^κ; Proposition 49.6(v) is the point that makes these useful. When traversing the single large gap the necessary deep point is constructed by extracting a coefficient in an Eisenstein expansion, avoiding the large cost of paying for every intermediate jump. This explains the role of the long-gap lemma below. For the precise verification of the cup-product depth inequalities in this regime one uses the same `scaledCupBounds_of_cut_windows` with $J = 1$ and $n = K$.

The long-gap operation. The auxiliary statement used for these cuts is worth formulating. In an unramified chart suppose $l > 0$, $3e_0 \leq p^l$, p^κ kills the cokernel at $i[-l]$, $\mathfrak{e}^s$ kills at i, all allowed jumps at $i[-j]$ are $\leq G_1$ for $0 \leq j < l$, and put

$$N = s + G_1 \sum_{j=0}^{l-1} p^j.$$

If $N < A$ and $e_0 A \leq p^l \delta$, there is a point in the extension over $p^{\kappa+4\delta}c$ with level-l standard exponents modified at $a = 0$ by adding $\min(A, -u - N)$ on the line of jump u, and with depth modified at $a = 0$ by adding A independent of u. This is *formal-source input* `PairingSide.pointsOver_nonempty_of_long_gap` in `New/Lift/LongGap.lean`. A brief account of the mechanism suffices to locate the nontrivial ingredients: by starting with a point over the pure Eisenstein annihilator, one multiplies by a monic polynomial in $\mathfrak{e}$ of degree N to arrive in a Frobenius-transported module. In this step each Frobenius predecessor contributes the corresponding power of p to the degree; explicitly the polynomials for the unramified chart arise from $\varphi^j(z + p) = (\mathfrak{e} - p)^{p^j} + p$. Applying the $p^{4\delta}$ estimate of Proposition 49.4 expresses the resulting element as a linear combination of nearly invariant points. Expand the coefficients of that combination in powers of $\mathfrak{e}$ using $\mathfrak{S}/\mathfrak{e} \simeq \Lambda$. Comparing the coefficient of degree N gives a new point, now with exponents controlled by N rather than by the full far weight. The strict inequality $A > N$ ensures that the original leading coefficient survives the comparison. The precise invariant coefficient extraction is expressed by `JumpLattice.coeff_mem_constantsModE` in that file. Thus the exponent 4δ in the frame theorem and the geometric degree N are both visible in the cost of the long-gap operation.

Few low jumps. In the last regime it would be wasteful to multiply every class by the exponent needed for a small exceptional part. The module of crystalline cocycles of the distinguished side is used as the trivial quotient C_0, with its identity cocycle map. In an

unramified chart suppose $p \geq 5$, $\eta \geq 1$, and all paired jumps lie at each letter in the opposite windows $I(\eta, \pm\epsilon_i, \beta_i, W_0)$. Let A_0 be a nonnegative integer such that for every letter

$$\beta_i + W_0 + \eta \leq (p-3)A_0, \qquad 6(A_0 + \eta + W_0) < \beta_i, \qquad \mathrm{NM}(p, 6(A_0 + \eta + W_0), \beta_i). \tag{49.15}$$

Then the following estimate holds.

Theorem 49.7 (Length estimate for few low jumps)**.** Formal-source input. *Let T have the indicated second jump window and let $\mathcal{Q}$ be the crystalline lattices with the opposite first jump sets. Suppose that at every coefficient embedding the sum of the multiplicities of jumps between $-\eta$ and η is at most μ. Then*

$$\Delta_p(T; \mathcal{Q}) \leq 11 f_0 \left(\mu(A_0 + 1) + (W_0 + 1)\,\mathrm{rk}_O\, T\right).$$

Moreover $\Delta_p(T; \mathcal{Q}) = 0$ if neither of the original jump sets meets $[-A_0, 0)$, or if at every embedding the second jump set is contained in $[-\eta, 0]$.

This is `KummerChart.few_low_jumps` in `New/Lift/ExponentTheorems.lean`. Its bound involves the width of the far window and the number of low jumps separately, rather than the rank times A_0; this refinement is essential in summing the loss.

Here is the internal length argument behind it. For each letter, consider the constants which lift to points with exponent raised$_0$ and prescribed depth. To increase depth on one graded line of multiplicity $m_T(i, u)$ at the Frobenius divisor $\mathfrak{e}_a$, the index can be controlled by counting relations modulo p among its leading coefficients. For an unramified chart, if the depth conditions at $\mathfrak{e}$ can be advanced to the next Frobenius threshold without a nonzero multiple of p intervening, then the leading coefficients of $f_0 m_T(i, u) + 1$ points satisfy a nontrivial integral relation with one coefficient equal to 1 modulo the deeper submodule. The leading-term calculation and normalization of a relation by its coefficient of least valuation are made precise by `exists_comb_of_jumpMultiplicity_le` in `New/Lift/LongGap.lean`. Indeed the leading terms on a Hodge line of multiplicity m may be represented in at most m copies of the finite unramified coefficient module of rank f_0 over $\mathbb{Z}_p$; the rank count supplies a relation. The theorem includes the important compatibility of the retraction with this leading-term comparison, so it is a statement about the actual integral exponent modules, not just dimensions of their rational Hodge pieces.

For finite free modules the passage from this relation property to a length bound is elementary. If $p^d L \subset L'$ and every $q+1$ elements have an integral combination, one coefficient a unit, belonging to L', then the elementary-divisor decomposition of L/L' shows that at most q cyclic elementary divisors can be nonzero, and all their exponents are at most d. The length is therefore at most dq over $\mathbb{Z}_p$. Applying this estimate to each depth step and using subadditivity of relative length bounds the cost of changing the exponents from $c(a, u)$ to $c'(a, u)$ by a sum of step costs weighted by the corresponding multiplicity. More concretely, let n_i and r_i bound f_0 times the low and the total jump counts at i. The prescribed module of deep constants at letter i has colength at most

$$(2A_0 + 1)n_i + 3(W_0 + 1)r_{i[-1]} + 6A_0 n_{i[-1]}. \tag{49.16}$$

The precise modules and sign-dependent depths are `deepConstants`, `relLength_deepConstants_le`, and `exists_relLength_deepConstants_le` in `New/Lift/FewLowJumps.lean`. If the sign of the second side at i is positive its depth increases by $2A_0 + \eta$ at the single Eisenstein element. For the negative sign the prescribed depth is $\beta_i + W_0 + 2A_0$ at $a = 0$ and $W_0 + 1 - u + 2A_0 \mathbf{1}_{|u| \leq \eta}$ at $a = 1$, with $-u$ elsewhere; this is the depth underlying (49.16). The verification of (49.16) from the individual steps is a formal-source input; the source further

checks that these depths give safety through the cut lemma. Taking the intersection over all letters can only cost the sum of their colengths (inject the quotient of the intersection into the direct sum of quotients). Since $i \mapsto i[-1]$ is a permutation, summing (49.16) gives a bound by

$$11 \sum_i ((A_0+1)n_i + (W_0+1)r_i).$$

For $n_i = f_0 \sum_{|u|\le\eta} m_T(i,u)$ and $r_i = f_0 \sum_u m_T(i,u)$ this is at most $11 f_0 [K_O : \mathbb{Q}_p](\mu(A_0+1) + (W_0+1)\,\mathrm{rk}_O\, T)$ over $\mathbb{Z}_p$. Dividing by $[K_O : \mathbb{Q}_p]$ as in (49.1) proves the displayed normalized length bound from the integral depth statements. The zero-loss part of Theorem 49.7 does not follow by making this right side zero: when the low negative jumps are absent, pure Eisenstein annihilators and zero-cost depth steps give safe points at exponent zero. The second alternative is treated separately by the no-multiple condition and one Frobenius transport of the low-jump side. These are the two cases in `SidePairing.scaledCupBounds_of_killsCokernel`, `New/Lift/LongGap.lean`.

49.8 The tame type bound, uniform in the ramification degree

At the selected primes g_0, g_1 the local representations may only be potentially crystalline. Here the ramified inertial type has a special form. Let r_0, N_0' be bounds and suppose that for every inertial element s the polynomial class $\mathcal{K}(s,\cdot)$ contains $X^M - 1$, where $M = \mathrm{lcm}_{1\le j\le r_0}(p^j - 1)$, and contains a monic polynomial in $\mathbb{Z}_p[X]$ of degree at most N_0'. This is precisely `IsTameType` in `New/Lift/ExponentTheorems.lean`. The dual condition is formed by using polynomials at s^{-1}. Suppose now that the local field is $\mathbb{Q}_p$, the paired jump sets lie in opposite windows $I(\eta, \pm\epsilon, b, W_0)$, and

$$\kappa_0 = r_0^2(N_0'+1)^2, \qquad b + 6(W_0+\eta) + \kappa_0 + 2 < p.$$

Then multiplication by

$$p^{3\kappa_0 + 150(\eta + W_0 + 1)} \tag{49.17}$$

sends $Z_f(T)$ into the orthogonal part for all lattices in the opposite polynomial class. This is *formal-source input* `tame_types_exponent`, in the same file.

For the uniformity of (49.17), it is especially useful to recall its proof mechanism. All the roots of $X^M - 1$ are prime-to-p roots of unity; one may use a tame chart of ramification index M, which can be extremely large. But $M \le \prod_{j\le r_0} p^j \le p^{r_0^2}$, and the monic small-degree annihilator allows one to avoid paying for all that ramification. A generator s of the tame quotient acts on the chart coordinate as $z \mapsto \xi z$ for a Teichmüller root of unity of order M. If Q is a monic annihilator on the kernel lattice, then $Q(X)(X-1)$ kills the extension of a trivial lattice by that kernel. Indeed $s - 1$ maps the extension into the kernel on crystalline periods, and the polynomial Q kills that kernel. The corresponding operator on framed sections is the polynomial of the tame generator; its image is divisible by z in the integral frame. This last assertion is the specific formal-source input `TameChart.exists_z_smul_eq_sum_tameGen` in `New/Lift/SparseThickLetters.lean`. It is proved by applying the relation on the period matrix and using the divisibility test in the frame theorem. To lower a power z^j in the multiplier ideal one now uses this polynomial evaluated at ξ^j: unless its reduction vanishes, it is a unit and no power of p is needed. Where it vanishes the finite-difference step pays at most $\log_p M \le r_0^2$. The reductions of ξ^j for $0 \le j < M$ are distinct, hence there are at most $\deg Q + 1$ exceptional steps. Thus the cost at a side is bounded by $r_0^2(\deg Q + 1)$, independently of M itself. The third extension uses a polynomial of degree bounded by the product of the two degrees (plus the contribution of the trivial extension). The chosen κ_0 is sufficient for all three sides. This exact bound for the cokernel is `TameChart.killsCokernel_of_isKilledBy`. Together

with the no-multiple condition, guaranteed by the size hypothesis $b + 6(W_0 + \eta) + \kappa_0 + 2 < p$, the depth and safe-pair calculation gives (49.17); the factor 150 is a generous allowance for the three sides. The rank-independent cost of the earlier depth operators and the degree rather than ramification bound for the tame operators are distinct features of the proof.

In the selected-prime application, the root condition itself gives the small monic annihilator. A root of order dividing $p^j - 1$ lies in an unramified extension of degree dividing j, hence has monic minimal polynomial over $\mathbb{Z}_p$ of degree $\le j$. The product of at most N such polynomials is a monic polynomial of degree $\le Nr_0$ vanishing on the indicated roots. This explains `isTameType_of_roots` in `New/Selmer/RowBalanced.lean`, which is applied to the polynomial class of the finitely many columns. It contrasts with a direct estimate based on the order of the whole finite group at g_0, g_1.

49.9 Case analysis for the primes and the explicit local loss

The form in which the three crystalline-window estimates are combined is the following. We give the constants, since an asymptotic assertion without them would not specify the interface to the global proof. Temporarily write b_* and G_* for the two window sizes, h_* for the half-gap parameter, and let $Q > 0$ be the prime cutoff. Put

$$\lambda_p(h_*, b_*) = v_p\left((2b_*)!\left(\prod_{\max(b_*-30h_*^3,0)<j\le b_*+30h_*^3} j\right)^{h_*^3}\right). \tag{49.18}$$

Assume the size conditions

$$99(Q^2 + (W_0+1)Q + Q + W_0 + m) \le h_*, \qquad 99h_*^3 \le G_*,$$
$$99(G_*h_*^3 + G_*) \le b_*, \quad 2\log(4b_*) < m, \quad \delta < \eta, \quad 99\big((\eta + W_0)Q + \eta + h_*\big) \le h_*^3,$$

with $e_0 \le h_*$. Let the opposing jump windows be $I(\eta, \mp\epsilon, b_* + \delta, W_0)$ at thick embeddings and $I(\eta, \mp\epsilon, G_* + \delta, W_0)$ otherwise. Suppose that either all negative jumps on both sides satisfy $2u \le -h_*$ and the multiplicity of jumps of the second lattice T in $[-\eta, \eta]$ is at most μ per embedding, or the second jump sets consist entirely of jumps in $[-\eta, 0]$. At letters above a given embedding the sign and the thick indicator must be compatible whenever the embedding is thick or $Q < p$. If $p > Q$ the chart must be unramified with $p \ge 7$; at $p \le Q$ one may instead use constant signs and the sparsity of thick letters (no three successive thick letters with the two positive gaps at most $\lfloor \log_p(h_*^3) \rfloor$). This is the set of hypotheses used by `crystalline_defect_le` in `New/Selmer/CrystallineBound.lean`, where T is crystalline.

Under these hypotheses the formal-source local estimate is

$$\Delta_p(T; \mathcal{Q}) \le d_F\, 10^5 \left(\mathrm{rk}_O\, T \left\lfloor \frac{2\lambda_p(h_*, b_*)}{Q} \right\rfloor m + \mu\lambda_p(h_*, b_*)\right). \tag{49.19}$$

The same statement uses an arbitrary upper bound y for the parenthesized expression including 10^5 in its source formulation. Here are the quantitative subcases in the proof. Write $v = \lambda_p(h_*, b_*)$ and $K = \lfloor 2v/Q \rfloor$. By the factorial term and Legendre's formula,

$$\lfloor 2b_*/p \rfloor \le v, \qquad 2b_* < p(v+1), \qquad KQ \le 2v < (K+1)Q.$$

Also $\lfloor \log_p(4b_*) \rfloor < m$, since $p \ge 2$ and $2\log(4b_*) < m$.

- For $p \le Q$ the resource inequality $2b_* < p(v+1)$ forces a large bound K; after enlarging the small window to absorb the thin weight, use the sparse-thick exponent with $J = \lfloor \log_p(h_*^3) \rfloor$. The source compares $2(b_* + \delta)$ with $h_*^2 K$ and takes the depth budget compatible with K. The rank bound of Lemma 49.2 then suffices, at cost bounded by the first term of (49.19).

- If $p > Q$ and $v \geq h_*^3$, use the two-sizes exponent (49.14). The floor inequality, $p > Q$, and the size assumptions ensure the requisite bounds on $b_* + \delta, G_* + \delta$ in terms of K. The logarithmic factor is at most m, so by Lemma 49.2 the normalized loss is at most $2 \cdot 10^4 d_F \operatorname{rk}_O TKm$, absorbed by the first term.
- If $p > Q$ and $v < h_*^3$, the product factor in (49.18) has no multiple of p in its interval: a single such integer would contribute h_*^3 to v. The factorial term also implies that p is large compared with G_*, since $2b_* < p(h_*^3 + 1)$. These facts yield the no-multiple condition for the two sizes in Theorem 49.7 at $A_0 = 2(v+1)$. That theorem either gives zero loss (in the pure-low second case or when the negative weights avoid $[-A_0, 0)$) or gives a contribution of order $d_F(\mu(v+1) + (W_0+1)\operatorname{rk}_O T)$. In the nonzero case the large negative gap entails $h_*/5 \leq v$, and the numerical assumptions imply $W_0 + 1 \leq K$. Thus this too is absorbed by (49.19).

The exact floor and scale comparisons, including the sparse-thick budget in the first subcase, are part of the formal-source input `crystalline_defect_le`. Notice the structural improvement: the second term has the number of low jumps, not the rank, multiplying the large factorial valuation. Without that refinement, the eventual bound at rank-one string blocks and at the main row block would not have the required common budget.

49.10 Insertion of the global weights and the sum over all blocks

Return to the admissible datum $\mathcal{D}$. For this application put $h_* = h_2 = r^8$, $Q = kt$, and set

$$b_1 = N_1 \left\lceil \frac{a}{xN_1} \right\rceil, \qquad b_2 = N_1 \left\lceil \frac{a}{x_1 N_1} \right\rceil,$$

so $w_j(\sigma) = (b_1 \text{ or } b_2) + 2jh_2 + h_2$ at the two kinds of places. The main row has rank r' with $1 \leq r' \leq r$, and its block string has length $t' \leq t$. Its jumps at a complex embedding σ satisfy the alternative

$$h_2 \leq 2|u| \leq 2h_2^2 \quad \text{or} \quad |\epsilon_\sigma u + w_j(\sigma)| \leq t,$$

for one $j \leq m$; the number of jumps of absolute value at most $2h_2^2$ is at most $r'm/k$ per embedding. Here ϵ_σ is the sign of the CM embedding, $+1$ for the chosen CM type, -1 for its conjugate. The source also requires the sum of negative jump counts at $\sigma, \bar{\sigma}$ to be at most r'. These are the hypotheses `IsRowType` in `New/Selmer/DeterminantBound.lean`, used for the actual main row representations. The one-dimensional extra blocks are cyclotomic powers with jumps $c \in [0, t')$. After taking Tate duals the new large windows are opposite to the original ones, by (49.2), while the rank-one blocks have all jumps small and nonpositive. The local substitution in (49.19) is $b_* = b_1$, $G_* = b_2$, $W_0 = 4t$, $\eta = h_2^2 + 1$, and $\delta = 2jh_2 + h_2 - 2t$. This is why the input to the crystalline theorem is a count of low jumps rather than an absolute bound on all weights.

The local embeddings above p have sign and place classes related by the Galois action on roots of unity. If $g'g^{-1}$ acts on the prime-to-p part of the cyclotomic generator by the p^n power, the ratio of the infinite-place classes is $\epsilon_\sigma \epsilon_{\sigma'} p^n$ modulo the relevant prime-to-p divisor of N_1. This is `placeClass_ratio`. Squaring removes the sign, and the three-term avoidance in $\mathcal{S}$ gives `not_thick_of_short_arc`: for $p \leq kt$, if two thick embeddings are separated by a Frobenius arc of length $n \leq 27s$ then a third at interior distance $0 < i < n$ cannot be thick (where s is the cyclotomic tower exponent). Furthermore at $p \mid N_1 d$ two thick embeddings in one local Galois orbit have the same CM sign, by `cmSign_eq_of_thick`.

These are specific arithmetic *formal-source inputs*, `New/Selmer/ThickSpacing.lean`, used to verify the small-prime hypotheses of (49.19). At large primes away from the conductor the local extension is unramified and the larger-prime hypotheses apply. These separate facts explain why it is not sufficient merely to take a large set of thick places, without the progression-freeness built into $\mathcal{D}$.

Define now the local cost used in the global proof:

$$\mathrm{loss}_p = 10^6 \frac{m}{k}(b_1 v_p(g) + \lambda_p(h_2, b_1)), \qquad g = g_0 g_1. \tag{49.20}$$

At the selected split level primes use (49.17), with the degree bound for the polynomials derived from the good-type eigenvalues at those primes; the source uses $r_0 = r$ and $N_0' = (r+t)r$. At other primes use (49.19). With the preceding scale and spacing verifications the result is the following exact statement.

Proposition 49.8 (One-place block estimate)**.** Formal-source input. *In the above setting, let $v \mid p$ and let T_i be the lattices of the row plus string blocks (one row, and one for each $c \in [0, t')$). Let $\mathcal{Q}_v$ be the class of local lattices inherited from the chosen column representations, with the same jump restrictions and crystalline condition, or the stated tame polynomial condition when $p \mid g$. For all sufficiently large coefficient rings, a block belonging to this class has*

$$\Delta_p(T_i^*(1); \mathcal{Q}_v) \le [E_v : \mathbb{Q}_p] \operatorname{rk}_O T_i \frac{\mathrm{loss}_p}{q_i}, \qquad q_i = \begin{cases} 1, & i \text{ the row}, \\ t, & i \text{ a cyclotomic string block}. \end{cases}$$

This is `block_defect_le_at_place` in `New/Selmer/OnePlaceDefect.lean`. The verification of the numerical parameters and the precise embeddings-to-letters dictionary are included in that declaration. It uses the very large gap between $p = g_0, g_1$ and the weights in the tame case. The row-type assumptions for actual automorphic Galois representations must be proved by the other part of the congruence argument; the local proposition does not give their existence. In particular, the gain for string blocks arises from their special single low weight, not from an assertion that all rank-one blocks have zero crystalline defect.

Let $\mathrm{bloss}_{i,p}(M_i)$ be the normalized relative length of the module of obeying cocycles on the i-th block modulo those also satisfying the crystalline relations simultaneously at all $v \mid p$. By Proposition 49.3 and subadditivity of relative length on intersections,

$$\begin{aligned} \sum_i \mathrm{bloss}_{i,p}(M_i) &\le \sum_{v|p} [E_v : \mathbb{Q}_p] \left(r' + \frac{t'}{t}\right) \mathrm{loss}_p \\ &\le 4 r' f \, \mathrm{loss}_p. \end{aligned}$$

The last step uses $r' \ge 1$, $t' \le t$, and $\sum_{v|p} [E_v : \mathbb{Q}_p] = [E : \mathbb{Q}] = 2f$. It is valid for arbitrary choices of the moduli M_i. This is exactly `sum_blockLoss_le` in `New/Selmer/RibetElement.lean`. It gives the $4f \, \mathrm{loss}_p$ term in the row budget per rank.

Finally the summation of this term over all rational primes with logarithmic weights is especially transparent. Write $J_* = (2b_1)! \left(\prod_{\max(b_1 - 30h_2^3, 0) < j \le b_1 + 30h_2^3} j\right)^{h_2^3}$. For a finite set of primes P_0,

$$\begin{aligned} \sum_{p \in P_0} 4f \, \mathrm{loss}_p \log p &\le 4 \cdot 10^6 f \frac{m}{k}(b_1 \log g + \log J_*) \\ &\le 4 \cdot 10^6 f \frac{m}{k}(b_1 \log g + 2b_1 \log(2b_1) + 60 h_2^6 \log(b_1 + 30h_2^3)). \end{aligned}$$

Here unique factorization gives the first inequality, and the number of factors in the consecutive product is at most $60h_2^3$. On the admissible asymptotic family $b_1 \sim b$, $\log g \le 22 \log b$, and

h_2^6 is polynomial in x whereas b grows faster than every fixed power of x. Thus the cost is $O(f(m/k)b\log b)$. Its ratio to $B_0 = fa(\log b)^3/(mk)$ is

$$O\left(\frac{m^2}{x(\log b)^2}\right) = O((\log b)^{-2})$$

using $a \geq xb$, $m^2 \leq 4x$. The source proves the requisite fixed-budget inequalities for all admissible data, including the other part of the row budget, in `exists_isFitFamily`, `New/Selmer/FitTwistsExist.lean`. The displayed calculation identifies why the arithmetic parameters make even the high-weight local integral comparison affordable. At this point a loss of order $\log J_*$ is acceptable, but the large powers of r entering the rank and level cannot be permitted to multiply the factorial part indiscriminately. This is the significance of the rank-uniform depth method, the separate tame-polynomial estimate, and the low-jump colength bound in the proof.

Chapter 50

Integral period frames and the algebra of the local loss

We now examine the steps behind the local estimate in Chapter 49. There are two rather different kinds of assertion in that estimate. The first concerns ordinary finite-dimensional local cohomology, the second a uniform integral comparison problem. Ordinary weak admissibility, even if applied to every congruence representation separately, does not by itself supply the second assertion. In these pages we prove the elementary and classical cohomological deductions, develop the integral algebra used by the comparison, and isolate the additional comparison inputs in their specific matrix and module forms. This separation is important: it makes the normalization, the dependence on ramification, and the few-low-jumps saving intelligible independently of the notation of the implementation. Statements still used as formal-source inputs are identified as such; the discussion is not an independent proof of every theorem of the period-ring development.

50.1 The rank calculation and duality over a finite coefficient ring

Let $F/\mathbb{Q}_p$ be finite of degree d_F, let G_F be its absolute Galois group, and let T be a free O-lattice, of rank r, as in Section 49.1. Put $V = T[1/p]$. We use the convention that the jump of $O(1)$ is -1. In particular, a line with negative jump contributes one to the dimension of $D_{\mathrm{dR}}(V)/\mathrm{Fil}^0$. The dimensions of this quotient, calculated at the different embeddings, are the quantities that should control the rank of the cocycle lattice. It is worth giving the reason for the extra r in Proposition 49.1.

Proposition 50.1 (Rank of the finite-condition cocycle lattice)**.** *Suppose that V is de Rham and that the finite condition is the usual crystalline finite condition. Let $t_V = D_{\mathrm{dR}}(V)/\mathrm{Fil}^0 D_{\mathrm{dR}}(V)$. Then*

$$\mathrm{rk}_O\, Z_f(T) = r + \dim_{K_O} t_V .$$

Consequently, if the coefficient field splits the embeddings of F, and J_σ is the jump multiset at σ, then

$$\mathrm{rk}_O\, Z_f(T) = r + \sum_{\sigma : F \hookrightarrow K_O} \#\{u \in J_\sigma : u < 0\}.$$

Proof. The classical fundamental exact sequence for the finite condition gives

$$0 \longrightarrow H^0(F,V) \longrightarrow D_{\mathrm{cris}}(V) \xrightarrow{(1-\varphi,\mathrm{can})} D_{\mathrm{cris}}(V) \oplus t_V \longrightarrow H^1_f(F,V) \longrightarrow 0.$$

Here dimensions are taken over K_O (the Frobenius on the first factor is linear over K_O). This sequence, which is the usual Bloch–Kato cohomological application of the fundamental sequence of period rings, does not require the positive-dimensional vector space $D_{\mathrm{cris}}(V)$ to have its maximal possible dimension. Taking the Euler characteristic gives

$$\dim H^1_f(F,V) = \dim H^0(F,V) + \dim t_V.$$

The coboundary map $V \to B^1(F,V)$ has kernel $H^0(F,V)$, so that

$$\dim Z^1_f(F,V) = \dim H^1_f(F,V) + \dim B^1(F,V) = \dim t_V + r.$$

The finiteness theorem for local Galois cohomology of lattices implies that $H^1(F,T)$ is finitely generated over O, and $B^1(F,T)$ is a quotient of T. Thus $Z_f(T)$ is finitely generated and torsion free. A continuous rational cocycle on the compact group G_F takes values in some $p^{-n}T$, so it becomes integral after multiplication by p^n; the finite condition is unchanged by multiplication by this scalar. Hence $Z_f(T)[1/p] = Z_f(F,V)$. The rank calculation follows. Finally $F \otimes_{\mathbb{Q}_p} K_O = \prod_\sigma K_O$, and the dimensions of the separate filtered quotients count precisely the negative jumps. □

This argument uses the classical Bloch–Kato sequence, local cohomology finiteness and the meaning of de Rham Hodge jumps. It explains the rank computation in `Lib/PadicHodge/BlochKato/BlochKato.lean` and `New/Selmer/CrystallineRank.lean`; it does not replace the latter files' precise comparisons with the code's period-ring presentation.

There is a useful elementary way to prove the index bound behind the reduction to a dual finite condition. We give it since it is easy to miss the factors arising from coboundaries if one works entirely with classes. Put $M = T/p^aT$, $M^D = T^*(1)/p^aT^*(1)$, and $C = \mathbb{Z}/p^a\mathbb{Z}(1)$. The perfect finite pairing with target C is obtained by composing the O-valued evaluation with a generator of $\mathrm{Hom}_{\mathbb{Z}_p}(O,\mathbb{Z}_p)$ as an O-module. Indeed this dual is the inverse different $\mathfrak{D}^{-1}_{K_O/\mathbb{Q}_p}$ under the trace pairing. If d generates the inverse different, then $x \mapsto \mathrm{Tr}(dx)$ induces the required perfect pairing also after reduction. This assertion does not require that the ordinary trace pairing on O itself be perfect.

Let $Z = Z^1(F,M)$, $Z' = Z^1(F,M^D)$, with coboundary subgroups B, B'. By finite local Tate duality their cup pairing factors to a perfect pairing of Z/B and Z'/B' (with values in $H^2(F,C) \simeq \mathbb{Z}/p^a\mathbb{Z}$). Classical Bloch–Kato orthogonality gives the orthogonality of the images A, A' of the two finite-condition lattices. One justification passes first to the rational cup pairing on finite-condition classes; to pass back integrally, $H^2(F,\mathbb{Z}_p(1)) \cong \mathbb{Z}_p$ is torsion free. Here the coefficient functional can be understood on the entire tensor product of lattices before reduction. In the source the cup is read in $H^2(F,\overline{\mathbb{Q}}_p^\times)$. The inclusion $\mu_{p^a} \hookrightarrow \overline{\mathbb{Q}}_p^\times$ induces an injection on H^2 by the Kummer exact sequence and $H^1(F,\overline{\mathbb{Q}}_p^\times) = 0$ (Hilbert 90). Thus no orthogonality information is weakened by reading its image there.

Lemma 50.2 (Index by cocycle annihilation)**.** *Let $X' \subset A'$, and let $S \subset Z$ be a subgroup with zero pairing against X'. Then*

$$[S : S \cap A] \le [A' : X'].$$

Proof. Write $\bar{X}'$ for the image of X' in Z'/B'. The exact order of the annihilator of X' in Z is

$$|\operatorname{Ann}(X')| = |B| \frac{|H^1(F,M)|}{|\bar{X}'|}.$$

In particular it is at most $|B||B'||H^1(F,M)|/|X'|$. The local Euler–Poincaré formula for finite p-primary modules gives

$$|H^1(F,M)| = |M|^{d_F} |H^0(F,M)||H^0(F,M^D)|.$$

The two orders of the coboundary groups are $|M|/|H^0(F,M)|$ and $|M^D|/|H^0(F,M^D)|$. Since $|M| = |M^D|$, their product with $|H^1(F,M)|$ equals $|M|^{2+d_F}$. The rank calculation and the complementary jumps u and $-1-u$ give

$$|A||A'| = |M|^{2+d_F}.$$

For the equality here we use saturation, so that reduction of the crystalline cocycle lattice has precisely the order computed from its rank. We deduce $|\operatorname{Ann}(X')| \le |A||A'|/|X'|$. The subgroup $S + A$ is contained in $\operatorname{Ann}(X')$, so

$$[S : S \cap A] = \frac{|S+A|}{|A|} \le \frac{|\operatorname{Ann}(X')|}{|A|} \le [A' : X'],$$

as claimed. □

A practical consequence of Lemma 50.2 is the following. Suppose one has a second lattice of lifts of some reductions in A', and can prove that its intersection with the actual finite-condition lattice of the dual has normalized colength at most D. Pairing the first side against this intersection is sufficient to show that the first side can have at most D units of normalized defect. Thus the integral pairings need not lift to pairings of dual rational representations at all. What is essential is the existence of one large orthogonal submodule. This is why the special imperfectly dual pairings in Section 49.1 are natural, and why an estimate of relative length, rather than a mere assertion of nondegeneracy, occurs in the argument. The classical inputs in this reasoning correspond to `Lib/ClassField/LocalDuality/CrystallineDuality.lean`; the later applications additionally specify the period-valued cocycles and the coefficient functionals.

50.2 The coordinates in a Kummer chart

We explain the origin of several scalars that enter the local comparison. A letter E used just in this subsection and the ensuing frame calculation denotes a scalar generating a period-ring ideal; it is not the global number field. Let F have residue degree f_0 and ramification index e_0, and put $\Lambda = W(\mathbb{F}_{p^{f_0}})$. Work with the completed algebraic closure C_p of F. Its tilt $C_p^\flat$ has valuation normalized by the value on compatible systems of roots of p. Write

$$A = W(\mathcal{O}_{C_p^\flat}), \qquad A_{\max} = \widehat{A[\xi/p]}^{\,p}, \qquad B^+ = A_{\max}[1/p],$$

where ξ is a generator of $\ker\theta$ for Fontaine's homomorphism $\theta : A \to \mathcal{O}_{C_p}$. Let ϵ be a compatible system of roots of unity, with first root equal to 1 and subsequent primitive p^n-th roots. Set

$$\mu = [\epsilon] - 1, \qquad \tau = [\epsilon^{1/p}] - 1, \qquad \xi = \frac{\mu}{\tau} = \sum_{j=0}^{p-1} [\epsilon^{1/p}]^j.$$

This is a choice of generator of $\ker\theta$; the relations $\varphi(\tau) = \mu$ and $\theta(\tau) = \zeta_p - 1$ follow directly. Here B^+ is the presentation of a positive period ring using the completed ring generated by ξ/p, rather than by divided powers of ξ. In customary notation it is a maximal period-ring presentation. We do not identify its integral model with the divided-power model of B^+_{cris}. The precise admissibility and kernel comparisons for the presentation in the development are part of the period-ring inputs. This distinction matters in statements involving p-divisibility.

Choose a uniformizer ϖ_F with $\varpi_F = -p$ if F is unramified. Select compatible p^n-th roots of ϖ_F in $\overline{F}$. Their system defines $z^\flat \in \mathcal{O}_{C_p^\flat}$; put $z = [z^\flat]$. The map sending the formal variable z to this element is an embedding

$$S = \Lambda[[z]] \longrightarrow A.$$

Here the infinite series are evaluated using the topology defined by (p, z). To see injectivity, reduce a nonzero series modulo p after dividing by its greatest common power of p. The first nonzero coefficient of the remaining series has valuation zero in the tilt, whereas higher terms have strictly larger valuation. The non-Archimedean triangle inequality shows its value is nonzero. Successive division by p, together with p-adic separatedness, proves injectivity in general. The restriction of θ to S is onto $\mathcal{O}_F$: every element of that ring can be written as a convergent series in ϖ_F with residue representatives in Λ. Choose a unit $a(z) \in S^\times$ lifting the unit $-p/\varpi_F^{e_0}$ via this restriction and set

$$E = z^{e_0} a(z) + p. \tag{50.1}$$

Then $\theta(E) = 0$, and modulo p, E has the same valuation as ξ. Since $\ker \theta = (\xi)$, it follows that E is a unit multiple of ξ in A. In the unramified case we can take $a(z) = 1$, so $E = z + p$. Define $v \in A^\times$ by $E\tau v = \mu$. The same ring $A_{\max}$ is obtained using E/p as generator before completion. Formula (50.1) implies $z^{e_0}/p \in A_{\max}$, a fact used repeatedly in the denominator estimates.

There is a useful supply of Galois elements acting on z. For each integer $m \geq 0$ and $k \in \mathbb{Z}_p$, an element of G_F sends the chosen p^m-th root of ϖ_F to its product with the chosen p^m-th root of unity to the k-th power, because $X^{p^m} - \varpi_F$ is Eisenstein and hence irreducible over F. Fix k and let m vary. The sets of elements giving the desired action at level m are nonempty compact subsets of G_F, nested by compatibility. Their intersection is nonempty. For any element g_k in this intersection,

$$g_k(z) = [\epsilon]^k z.$$

This applies to the ordinary G_F; when a chart is defined with additional allowed-tame-action conditions, the chart existence statement has to supply the corresponding subgroup. It is the exact chart of the source, including its subgroup conditions, that is used in the later formal-source frame theorem.

Put $E_j = \varphi^j(E)$ for $j \geq 0$. Frobenius acts on S by the usual Frobenius on Λ and $z \mapsto z^p$. Evaluation at φ^{-j} followed by θ sends z to the selected p^j-th root of ϖ_F and acts on Λ by an automorphism. It is thus onto the integers of the extension F_j generated by this root. Its kernel is generated by E_j. For example, this follows by Weierstrass division: E_j is a unit times an Eisenstein polynomial of degree $e_0 p^j$ over Λ. Its quotient ring is finite free of that rank over Λ and has the indicated uniformizer and residue field. Equivalently this evaluates the quotient ring as $\mathcal{O}_{F_j}$. In particular E_j generates a prime of S. If $j \neq k$, evaluating E_k by $\theta\varphi^{-j}$ gives two terms in (50.1) with unequal valuations, since the valuation of the first term is p^{k-j} while the value of p is one. Their sum does not vanish. Therefore the primes (E_j) are distinct. In the de Rham filtration, $\varphi^{-j}(E_j)$ is a generator of the first filtration step, while $\varphi^{-j}(E_k)$ for $k \neq j$ is a unit. In particular estimates at different j can track different prime factors in the coefficient ring S.

Two more elementary nonvanishing observations are useful. The evaluation of z in each F_j is nonzero; and $\theta\varphi^{-j}(\tau) = \zeta_{p^{j+1}} - 1 \neq 0$. Thus the powers of z and the powers of τ in an inequality cannot simply be replaced by powers of a single E_j. A loss expressed in these scalars is first a statement about integral comparisons; only later, along with the factorial bounds and truncation modulo p^κ, does it become a bound in p-adic length. The chart existence and these division calculations are found in `Lib/PadicHodge/WachKisin/KummerChartExists.lean`, `Lib/PadicHodge/WachKisin/ChartScalars.lean` and `Lib/PadicHodge/WachKisin/DeRhamValues.lean` (notably `exists_kummer_chart`, `Chart.weierstrass_division`, `frobE_dvd_iff`).

50.3 A coefficient retraction and why it is used

The period-ring calculations often need to descend a relation over B^+ to relations over S. It is not legitimate simply to regard B^+ as a power-series ring over S, nor to assume that a basis of its residue field is a finite basis of the period ring. An adic coordinate argument provides the appropriate substitute. We give the algebraic details, as they also clarify why the integral comparison need not have a loss proportional to the number of entries of a matrix at every stage.

Lemma 50.3 (Adic coordinates)**.** *Let $S = \Lambda[[z]]$, $\mathfrak{m} = (p, z)$, and M an S-module. Assume p acts injectively on M and z acts injectively on M/pM. Choose lifts $b_i \in M$ of a basis of $M/\mathfrak{m}M$ over $S/\mathfrak{m}$. There are S-linear coordinate maps $\lambda_i : M \to S$ with $\lambda_i(b_j) = \delta_{ij}$ having the following approximation property: for each $x \in M$ and each n, a finite sum $\sum_{i\in J} \lambda_i(x)b_i$ approximates x modulo $\mathfrak{m}^n M$, and all but finitely many coordinates are in $\mathfrak{m}^n$. If M is $\mathfrak{m}$-adically separated and has finite rank over S, it is finite free.*

Proof. Write $P = \bigoplus_i Se_i$ and $\beta : P \to M$ for $e_i \mapsto b_i$. First, for each $n \geq 1$,

$$\beta^{-1}(pM + z^n M) = pP + z^n P.$$

Modulo pM, if a finite linear combination belongs to $z^n(M/pM)$, its reduction modulo $z(M/pM)$ vanishes. By the independence of the chosen residue basis each coefficient reduces to a multiple of z. Divide the coefficients by z and use its injectivity on M/pM to reduce n by one. This proves the assertion by induction. Next we claim that $\beta^{-1}((p^m, z^n)M) = (p^m, z^n)P$. The case $m = 1$ was just proved. If $\beta(x) = p^m y + z^n w$, the case $m = 1$ permits us to write $x = px_1 + z^n x_2$. Subtracting the second term gives

$$p(\beta(x_1) - p^{m-1}y) = z^n(w - \beta(x_2)).$$

By reduction modulo p, the expression in parentheses on the right is divisible by p. Cancel p (injective on M) and apply induction. In particular

$$\beta^{-1}(\mathfrak{m}^{2n}M) \subset \beta^{-1}((p^n, z^n)M) = (p^n, z^n)P \subset \mathfrak{m}^n P.$$

For each $x \in M$ choose a finite combination $a_1 \in P$ with $\beta(a_1) \equiv x \bmod \mathfrak{m}M$. Given an approximation modulo $\mathfrak{m}^k M$, express its error as a finite sum of the monomials $p^i z^{k-i}$ times elements of M, approximate these elements modulo $\mathfrak{m}M$, and add the correction to obtain an approximation modulo $\mathfrak{m}^{k+1}M$. The displayed inclusion shows that the resulting finite-support coefficient vectors are Cauchy in the $\mathfrak{m}$-adic topology of P and are independent of the approximations in its completion. Taking the coordinate limits in the complete ring S defines the maps λ_i. Uniqueness of the limit also proves S-linearity. For a fixed n, the coordinate vector differs modulo $\mathfrak{m}^n$ from a finite-support approximate vector, which gives the asserted approximation property. To justify the first approximation property with the actual limits, note that a finite-support approximation modulo $\mathfrak{m}^{2n}M$ has coefficients equal to the limits modulo $\mathfrak{m}^n$; replacing them by the limits changes the approximation only by $\mathfrak{m}^n M$.

Finally the coordinate maps prove the linear independence of the entire family (b_i) over S, so if M has rank at most r, at most r of the b_i exist. For any x, the element $x - \sum_i \lambda_i(x)b_i$ belongs to every $\mathfrak{m}^n M$ by the approximation property. By separatedness it is zero. Thus the b_i are a finite basis. □

For $M = A$, the hypotheses of Lemma 50.3 hold. The module is p-torsion free, and modulo p it is the tilt's valuation ring, in which z is a nonzero element of positive valuation. To

check separatedness, if x lies in every $(p,z)^nA$, reduction modulo p shows it belongs to every $z^n(A/pA)$, hence $x = px_1$. For each n the inclusion $\mathfrak{m}^{2n}A \subset (p^n, z^n)A$ and injectivity of z^n modulo p show that $x_1 \in (p^{n-1}, z^n)A$. Therefore x_1 too lies in every power of $\mathfrak{m}$. Iteration shows that x is divisible by every power of p, hence vanishes. Choose the residue basis to contain the class of 1, and denote this coordinate by λ_0. It gives a retraction $A \to S$ of the scalar inclusion. It also preserves all divisibilities by elements of S: $\lambda_0(sx) = s\lambda_0(x)$.

We next explain its extension to the positive period ring. On the dense subring $A[E/p]$ the prescription

$$\widehat{\lambda}\left(\sum_{j=0}^{n} a_j (E/p)^j\right) = \sum_{j=0}^{n} \lambda(a_j)(E/p)^j$$

is well-defined, since any equality becomes an equality over A on multiplication by p^n. It preserves $A[E/p]$ and is p-adically continuous there. It consequently extends on completion to $A_{\max}$ and, by inverting p, to B^+. (The same argument works for any S-linear map $A \to A$ rather than the particular coordinates.) For manipulating A itself inside $A_{\max}$, the precise divisibility criterion used in the source is

$$a \in A, \quad p^m \mid a \text{ in } A_{\max} \quad \Longleftrightarrow \quad a \in (p, E)^m A. \tag{50.2}$$

The forward direction of this integral criterion is significant: it is not a criterion for divisibility by p in B^+, where p is invertible. The elementary reverse direction follows because both generators of the ideal are divisible by p in $A_{\max}$. The converse, called `Amax.mem_span_pow_of_p_pow_dvd` in `Lib/PadicHodge/PeriodRings/AmaxStructure.lean`, is an integral period-ring structure input. The adic-coordinate argument above follows the lemmas `Lift.exists_adicBasis` and `free_finite_of_rank_le` in that file. In particular the argument concerns coordinates of arbitrary A-elements, not just the values of an element in the scalar subring.

Let S_B denote the bicommutant scalar ring used in the chart. In concrete terms its elements are those scalars $x \in B^+$ such that multiplication by x commutes with each extended S-linear endomorphism of the type just constructed. This is a subring containing $S[1/p]$, and if an element is a unit of B^+ its inverse also has the commuting property. Its utility is the identity

$$\widehat{\lambda}(xy) = x\widehat{\lambda}(y) \qquad (x \in S_B,\ y \in B^+). \tag{50.3}$$

Thus matrix equations whose coefficients belong to S_B can be transformed by arbitrary extended coordinate maps. The coordinates themselves give elements in the bicommutant: evaluate them first on the dense set $A[E/p]$, on which the result is in $S[E/p]$, and then use continuity and inversion of p. The stronger properties used in the frame theorem also concern Frobenius and the action of the subgroup fixing both the scalar chart and its theta values. One needs the extension of (50.3) to fields of fixed and de Rham values at different inverse Frobenius evaluations. These properties are not consequences merely of S-linearity; they are the specialized chart-scalar inputs in `Lib/PadicHodge/WachKisin/ChartScalars.lean` and `Lib/PadicHodge/WachKisin/DeRhamValues.lean`. They allow both a reduction to coefficients in S and comparison between different E_j. Without them, a lattice over the very large ring A would not yield the required finite-rank module over S.

50.4 From crystalline periods to an integral frame

We distinguish the elementary manipulations of a period matrix from the integral existence assertion about such matrices. Let T first be a $\mathbb{Z}_p$-lattice of rank d. Over the positive period ring localized at μ, choose a matrix M whose columns are a basis of invariant crystalline

periods, relative to an integral basis of T, in the crystalline case. The twisting convention determines whether this is read in T or in the space of linear forms on T; here we fix the convention that the columns are periods and Galois action on them is the diagonal action. Assume all jumps lie in $[-a, b]$ with $a, b \geq 0$. Write $h = a + b$.

Over B^+_{dR}, a filtration-adapted matrix factorization has the following shape:

$$MY = W_0 \operatorname{diag}(t^{k_1}, \dots, t^{k_d}), \qquad -a \leq k_i \leq b.$$

Here t is a generator of its first filtration step, W_0 is invertible over B^+_{dR}, and Y is a change of crystalline basis compatible with the filtration at the chosen embedding. The statement at other embeddings and inverse Frobenius evaluations gives these inequalities at each of the scalars E_j from Section 50.2. A bound on pole orders is plainly preserved by any invertible matrices over a discrete valuation ring. If W_0 and the power of t are separated as above, multiplying M by t^a and multiplying M^{-1} by t^b puts both in matrices over the positive de Rham ring at that evaluation. Returning from these pointwise pole bounds to the positive period ring itself is a separate assertion, of the type proved using a fundamental-kernel/divisibility calculation in the period-ring development. It is not simply an intersection of finitely many de Rham rings inside an ordinary number field.

In the notation of the chart, the normalized matrix in the source is

$$U = (E\tau)^a M, \qquad UW = (E\tau)^h I_d,$$

where $U, W \in \mathrm{Mat}_d(B^+)$. It satisfies

$$\varphi U = v^a U F_1, \tag{50.4}$$

where F_1 is invertible over B^+ with entries in the scalar ring S_B. Indeed, wherever the crystalline invariant comparison applies, $\varphi M = MC$ for a matrix C of fixed-period coefficients. Since $\varphi(E\tau) = \varphi(E)\mu = (E\tau)v\varphi(E)$, the formula has $F_1 = (\varphi E)^a C$. The field in which C takes its values, and its corresponding membership in S_B, are part of the invariant-ring comparison. The Galois transformation law for U is inherited from that of M after accounting for the scalar ratio $g(E\tau)/(E\tau)$. The normalized period matrix obeys one more important condition: extended coordinate endomorphisms preserve the constraints $Uy \in E^h(B^+)^d$. These are the requirements of `IsPeriodMatrix`, in `Lib/PadicHodge/WachKisin/CrystallinePeriodMatrix.lean`; the step from a crystalline matrix to this package is `IsChartOf.exists_isPeriodMatrix`. We use this package as a formal-source input when the integral pole and fixed-scalar comparisons have not been independently established.

Here is the module to which adic algebra is applied:

$$P = \{x \in A^d : \ \exists y \in S_B^d, \ Uy = E^h x\}. \tag{50.5}$$

The reason for requiring $x \in A^d$ is to obtain an integral Galois transformation matrix after selecting an integral basis. This is stronger than requiring the relation over B^+ alone. The two regularity conditions needed by Lemma 50.3 are natural: if px belongs to P for $x \in A^d$, one divides its witnessing y by p in S_B to see that $x \in P$. If $zx \in pP$ for $x \in P$, reduction in A^d/pA^d and injectivity of z show that $x = px'$ for $x' \in A^d$; the first observation gives $x' \in P$. Thus P is p-saturated inside A^d, and z is injective on P/pP. Since P is a submodule of the separated module A^d, it is separated. To conclude finite freeness from Lemma 50.3, one still needs to bound its rank over S; note that the rank of A over S is certainly not one. The period-matrix coordinate preservation is used precisely to show that any more than d independent S-directions would violate the period relation together with the extended coordinate identities. This rank bound is part of `IsPeriodMatrix.exists_basis` in

`Lib/PadicHodge/WachKisin/PeriodMatrixBasis.lean`. Invoking it gives a free S-module of rank at most d inside A^d.

In passing from this module to the final frame, it is important to obtain saturation not just with respect to p, but with respect to scalars preserving the entire system of period equations. The following schematic linear algebra records what is needed. Suppose Q is a square basis matrix for P, and $UZ = E^h Q$ with Z over S_B. If for a row $l \in A^{1\times d}$ and an element $q \in A$ not divisible by p we have $lQ = qr$ with $r \in S^{1\times d}$, one needs divisibility $q \mid lx$ for all appropriately allowed solutions $Uy = E^h x$, even beyond $y \in S_B^d$. Such a statement is not a result of row-reduction over B^+: it concerns divisibility in A. The comparison uses a Frobenius contraction argument (`exists_Ainf_eq_of_contraction`) and the extended-coordinate saturation lemma in `Lib/PadicHodge/PeriodRings/BdR.lean` and `Lib/PadicHodge/WachKisin/PeriodMatrixBasis.lean`. Also the assertion that Q is square and that p does not divide $\det Q$ is not an automatic consequence of finite freeness of P. That assertion uses a minimal-support and fixed-coefficient argument in `IsPeriodMatrix.exists_frame` in `Lib/PadicHodge/WachKisin/PeriodMatrixFrame.lean`. We isolate both as formal-source inputs.

Once these statements hold, several matrix consequences are elementary. In particular the adjugate of Q, applied row by row, has $(\operatorname{adj} Q)Q = (\det Q)I_d$. The stronger divisibility property yields $x \in QA^d$ for allowed solutions x. Taking $Uy = (E\tau)^h e_i$ using the i-th column of W produces $x = \tau^h e_i$, and hence a matrix L over A with

$$QL = \tau^h I_d. \tag{50.6}$$

The comparison furthermore obtains $U = QY$ with $Y \in \mathrm{Mat}_d(B^+)$. If the coordinate matrix is Z as above, then cancellation of Q yields $YZ = E^h I_d$. Thus the determinant-clearing factor of the matrix Z in every rational comparison is exactly E^h, not E^{dh}. The distinction between the identity and the much weaker assertion on determinants is crucial. When only the determinant bound is used, applying an inverse matrix through the adjugate can cost d times the pole bound on each independent direction. Instead the right inverse after multiplication by E^h gives the pole bound per direction. Relation (50.6) and the analogous period identity describe how the frame lies between two scalar multiples of the integral lattice. A reader should not infer from this that a Wach module or a Kisin module over an arbitrary coefficient ring has been proved to be categorically equivalent to every representation in the argument. What is used here is a particular simultaneous frame bound, in a fixed chart, with its Galois and Frobenius relations and its uniform scalar denominator.

For clarity we record one consequence of a frame when estimating divisibility of coordinates. Let $Z'Z = E^h I_d$, with entries of both matrices in a comparison ring R. If Zx is known to be divisible by q for some $q \in R$ acting injectively, then $E^h x = Z'(Zx)$ is also divisible by q. If q and E are relatively prime in the sense that the image of E acts injectively on R/qR, this implies $x \in qR^d$. More generally the conclusion records a possible loss of E^h, not a loss of a determinant. The same elementary calculation works after Frobenius, with E_j in place of E. When the divisibilities are tested by evaluation of period-valued linear forms, the additional ingredient is the chart-coordinate comparison which identifies such evaluations with Zx. In the frame theorem this evaluation has, in symmetric weight range $[-a, a]$, the form of a factor E_j^a times a period-valued linear form evaluated on $\varphi^j(n)x$, where n denotes the frame. Substitution of n using the normalized matrix U proves the assertion over B^+; the divisibility conclusions back in A or S use the integral saturation. The source result `frame_theorem` packages these assertions in `Lib/PadicHodge/WachKisin/FrameTheorem.lean`. They, rather than rational filtered-module theory alone, are the starting point for the integral approximation below.

To spell out the precise matrix identity in symmetric weight range, use a for the absolute jump bound and $h = 2a$. Let n be the frame with columns $\tau^{-a}Qe_j$ in the ambient localized module. Relation (50.6) gives

$$\tau^a n = Q, \qquad nL = \tau^a I_d.$$

In particular $\tau^a nA^d \subset A^d$ and $\tau^a A^d \subset nA^d$. The Frobenius relation delivered by the integral frame computation is $\varphi Q = v^a Q B_1$ for a matrix B_1 over S. Here $E\tau v = \mu = \varphi\tau$. Therefore

$$\varphi n = \mu^{-a} v^a Q B_1 = E^{-a} n B_1.$$

This proves the stated finite-height inclusion for the frame span once it is identified with the prescribed Kisin lattice. The inclusion is not multiplied by the rank and holds simultaneously at every letter.

Now substitute $U = (E\tau)^a M$ in $UZ = E^{2a}Q$, obtaining $MZ = E^a n$. Apply Frobenius m times, then the comparison to B_{dR} and an arbitrary crystalline linear form λ; write ψ_m for this embedding of the period coefficients after Frobenius evaluation, and let $\mathrm{val}(\lambda)$ be the row of its values at the integral basis of the original lattice. For $x \in A^d$ the identity becomes

$$E_m^a \lambda(\varphi^m(n)x) = \mathrm{val}(\lambda)\, \psi_m(M)\, \varphi^m(Z)x, \tag{50.7}$$

where the scalars on each side are mapped to the common evaluation ring. Since M is invertible and its columns are invariant, the rows of its inverse yield crystalline forms whose values on the invariant basis give the coordinate rows. For these forms the right side becomes the separate coordinates of $\varphi^m(Z)x$. Suppose all of their values on $\varphi^m(n)x$ lie in a given B^+-submodule of the target ring. Multiplying by E_m^a preserves the submodule. Multiplying the rows by the entries of $\varphi^m(Z')$ and adding, with $Z'Z = E^{2a}I$, shows that every $E_m^{2a}x_j$ belongs to that submodule. For an ideal-principal test modulo a scalar d_0 this is the origin of the divisibility claim in the frame theorem: if division by E_m is allowed on the target quotient and membership in d_0 tests divisibility back in the integral coordinates, one obtains $x \in d_0 A^d$. The doubled exponent $2a$ originates in the two-sided jump bound, not in a matrix determinant. Identifying the span of the frame with the intrinsic module defined by all Hodge inequalities uses the simultaneous evaluations, the criterion for τ-divisibility from the zero theta values at positive Frobenius powers, and the coefficient-retraction scalar comparison. These identifications in `FrameTheorem.lean` remain integral comparison inputs; (50.7) is their explicit linear algebra.

50.5 The radius estimate and its fourth power

We can now explain the quantitative part of the integral radius lemma. The strength of the uniform period comparison is not merely that a suitable denominator exists for each T separately. Its denominator can be chosen as a scalar depending only on the chart and a range of weights; one avoids a choice depending on the rank or on a basis of T. The first half of the argument is a formal-power-series construction and the second half is a linear algebra calculation with truncated series. We develop the latter in detail.

Choose an integer ℓ_0 such that $p^{\ell_0} \geq 3e_0$, and put

$$X = z^{p^{\ell_0}}/p^2 \in A_{\max}.$$

Since $z^{e_0} \in pA_{\max}$, we have $z^{p^{\ell_0}} \in p^3 A_{\max}$ and $X \in pA_{\max}$. Thus formal series in $S[[X]]$ can be evaluated in $A_{\max}$ by p-adic completeness. The radius lemma applies to a matrix Z over

S_B with $Z'Z = E^h I$. Its transformation for a chart element g with $g(z) = [\epsilon]z$ is $gZ = ZG$, with G integral over A. It establishes a factorization

$$\varphi^{\ell_0} Z = E_{\ell_0}^{b}\, C\, V, \tag{50.8}$$

where C is fixed by the fixer of Λ and V comes from an invertible matrix over $S[[X]]$ evaluated at the above argument. In particular V, V^{-1} are integral over $A_{\max}$. The integer b is a parameter in the radius lemma. To see the mechanism of this factorization, one first multiplies Z by an appropriate fixed matrix B to achieve $BZ = I + zW$. The finite-difference identity in the chart promotes $gZ = ZG$ to the divisibility of $G - I$ by $z\tau$. A recursive approximation of BZ then expands its coordinates in powers of z. At the n-th step the division of the difference of z^{n+1} by $z\tau E$ produces the factor $p^{1+v_p(n+1)}$, by the formula for $[\epsilon]^{n+1} - 1$. The accumulated cost is bounded by

$$\sum_{j=1}^{n} (1 + v_p(j)) = n + v_p(n!) \le 2n.$$

The residual equation and the coefficient retraction of Section 50.3 select the coefficients in S. After applying φ^{ℓ_0} the budget p^{-2n} is absorbed by $z^{np^{\ell_0}}$, precisely the n-th power of X. The resulting series has constant coefficient I, hence is invertible as a formal matrix. The normalization by B, the divisibility at every recursive step, and the fixed property of C are the first half of `radius_lemma` in `Lib/PadicHodge/Comparison/CrystallineLimit.lean`, used here as an integral comparison input. The displayed valuation calculation explains why the matrix is evaluable in the positive integral period ring; it explains in particular the appearance of $3e_0$, as opposed to taking p^{ℓ_0} arbitrarily large depending on Z.

Here is the next part of the argument, including the reason for the fourth power in the bound. Let $D \ge 0$ and $A_0 \ge 0$ be integers satisfying $e_0 A_0 \le p^{\ell_0} D$, and let $\overline{R} = B^+/E^{A_0}B^+$. A series $H(X) = \sum h_n X^n$ with coefficients in S has

$$p^{2D} H(X) = H_0(z) + z^{p^{\ell_0} D} H_1(X), \qquad H_0 \in S, \quad H_1 \in S[[X]]. \tag{50.9}$$

Indeed take the first D terms, multiplying each by p^{2D}, and rewrite the tail by shifting its index by D. The first term is integral because for $n < D$ the coefficient of $z^{p^{\ell_0} n}$ is $p^{2(D-n)} h_n$. The scalar $z^{p^{\ell_0} D}$ is divisible by p^{A_0} in $A_{\max}$. We use the precise truncation of $p^{A_0}\widehat{\lambda}_0(y)$ modulo E^{A_0} for $y \in A_{\max}$: there exists $s \in S$ with

$$p^{A_0}\widehat{\lambda}_0(y) - s \in E^{A_0} B^+.$$

This is the special E case of `exists_pow_mul_extendedCoord_eq_add`, obtained from the integral decomposition $p^{A_0} y = a + E^{A_0} y'$ with $a \in A$ in `Amax.exists_p_pow_mul_eq_add`, followed by $\widehat{\lambda}_0$. It is the exact ideal-theoretic approximation required here; truncation merely modulo some high power of p in B^+ would be meaningless. Since $\widehat{\lambda}_0$ fixes the convergent evaluations of $S[[X]]$ (the partial sums are fixed and converge p-adically after the prescribed scaling), applying it to (50.9) gives a polynomial approximation in the sense that $p^{2D} H(X)$ has the same image in $\overline{R}$ as some element of S.

Apply this entry by entry to the two formal units giving V and V^{-1}. Obtain $N, M \in \mathrm{Mat}_d(S)$ satisfying, after mapping to $\overline{R}$,

$$\overline{N} = p^{2D}\overline{V}, \qquad \overline{M} = p^{2D}\overline{V}^{-1}.$$

Consider the subset of S^d consisting of all y for which Vy is a constant vector of natural numbers modulo E^{A_0}. The S-span of these y contains every column of M, since $\overline{V}\,\overline{M} = p^{2D} I$.

It contains every column of $p^{4D}I - MN$, since these columns are mapped to zero by $\overline{V}$. For every $x \in S^d$, the identity

$$p^{4D}x = (p^{4D}I - MN)x + M(Nx) \tag{50.10}$$

then expresses $p^{4D}x$ in this span. This proves the second half of the radius lemma, with its original bound $p^{4D}S^d$. Notice the use of two approximations, each costing $2D$: the cost is not a determinant of an approximating matrix and is independent of d. This is an instance of a general pattern in the local proof: instead of solving an entire system of equations by its determinant, one approximates both a matrix and the matrix that provides a prescribed multiple of its inverse, and subtracts the residual identity.

At $A_0 = D = 0$, the integral radius factorization supplies the uniform positive comparison for all crystalline lattices with jumps in a fixed interval $[0, h]$. The multiplier is the image of $\varphi^{\ell_0}((E\tau)^h)$ in $A_{\max}$, independent of the dimension of T. Indeed in the normalized equations $UZ = E^hQ$ and $QL = \tau^hI$, choose $b = 0$ in (50.8). Apply Frobenius and replace the image of Z by CV; the product $\varphi^{\ell_0}(E^hQ)V^{-1}$ is integral and equals the invariant period matrix times the fixed matrix C after applying Frobenius. The invariance holds because Frobenius commutes with the Galois action and C is fixed by the group in question. Multiplication by $V\varphi^{\ell_0}(L)$ then gives precisely $\varphi^{\ell_0}((E\tau)^h)I$. Inverting this multiplier over the crystalline period ring is possible since $E\tau$ is associated to μ and the Frobenius translates of E needed here are units over the relevant localized period ring. This is the matrix version of `uniform_period_bound` in `Lib/PadicHodge/Comparison/CrystallineLimit.lean`. Its mathematical role differs from the usual statement that a sequence of bounded-weight crystalline representations has a crystalline limit: one also knows an explicit scalar uniform in the integral lattice and need not take a different period denominator on every individual line. The source cites the Dwork-trick arguments of Kisin and the integral comparison in the work of Bhatt and Scholze as conceptual antecedents to this construction; one needs the exact radius and scalar bounds for the subsequent length calculations.

50.6 Exponent lattices, Frobenius letters and the small part

We explain the role of the two variables of an exponent lattice by using the evaluations of the chart. Let $I = \mathbb{F}_{p^{f_0}}^{\times}$ be the set of letters. Write $i[-j]$ for the inverse Frobenius translate of a letter by j. At a fixed letter, the relevant jump spaces are those of the crystalline forms for each evaluation. An exponent system $o(j, u)$ assigns a lower bound on filtered valuation to the evaluations of those forms with jump u at letter $i[-j]$. The standard exponents of level l are

$$\mathrm{st}_l(j, u) = \begin{cases} 0 & j < l, \\ -u & j \geq l. \end{cases}$$

The membership conditions also restrict the point to the appropriate level of the localized frame and require that the evaluations take values in the chart value fields. Consequently they are stronger than the displayed inequalities over B_{dR} alone. If the lower bounds increase, the corresponding exponent module decreases. Intersecting two systems of bounds gives their pointwise maximum. If all exponents are shifted by m at $j = j_0$, multiplication by $E_{j_0}^m$ implements the shift in the rational module, because $\varphi^{-j}(E_{j_0})$ has valuation one precisely when $j = j_0$ and zero otherwise. Likewise applying Frobenius l times transports the letter $i[-l]$ to i and displaces the old filtration bounds by l places. This is why applying Frobenius to a lift over one letter can make the inequalities easier at old evaluations but introduces new inequalities at the first l places. Exact membership statements, including the required

value field and local-frame conditions, are in `New/Lift/ExponentLattices.lean`; the evaluation-subspace and strictness assertions needed in subsequent length estimates occur in `New/Lift/ExponentTheorems.lean`. They are not automatic properties of arbitrary sequences of filtrations.

We recall the algebra controlling the coefficient decomposition. If $q_0 = p^{f_0} - 1$, the operator multiplying by a primitive q_0-th root ζ from the coefficient ring admits the integral projectors

$$e_i = q_0^{-1} \sum_{j=0}^{q_0-1} (\omega(i)^{-1}\zeta)^j.$$

The denominator is a unit and all differences of distinct Teichmüller roots $\omega(i)$ are units. Hence $e_i e_{i'} = 0$ for $i \neq i'$, $e_i^2 = e_i$, and their sum is one: all three follow by evaluating the polynomial in the simple-root decomposition of $X^{q_0} - 1$, which is separable even modulo p. Frobenius sends e_i to e_{ip}, since it raises the Teichmüller factor, but not the coefficient-lattice factor, to the p-th power. If a bilinear pairing is balanced for multiplication by ζ on its two arguments, the evaluations on unlike letters are orthogonal. Indeed $\omega(i)P(x,y) = \omega(i')P(x,y)$ for the two arguments in their respective letter components; subtracting gives zero because the difference is invertible. These calculations hold integrally modulo p^m for every m. They explain why it suffices to test exponent inequalities one letter at a time, and why summing lengths over letters does not multiply them by the full number of letter pairs.

The cup-product argument uses a further fact about the small part, whose core is ordinary Artin–Schreier–Witt algebra. We prove the corresponding basic statement explicitly. Let $L = C_p^\flat$ be the algebraically closed tilt field, let $\mathfrak{m}_L$ be the ideal of elements of strictly positive valuation in its valuation ring, and let $W_m(L)$ denote the truncated Witt vectors. Put $Q = W_m(\mathfrak{m}_L)$, by which we mean the vectors all of whose Witt coordinates belong to $\mathfrak{m}_L$. This is an ideal of $W_m(\mathcal{O}_L)$ and an additive subgroup of $W_m(L)$ (not an ideal of the field's whole Witt ring). Frobenius on the latter acts on Witt coordinates by the p-th power. On the Tate-twisted tensor the auxiliary Frobenius acts on the Witt part, while the Galois group acts diagonally. Thus the fixed constants of that auxiliary Frobenius form the finite Tate line used in Section 49.6.

Lemma 50.4 (Artin–Schreier–Witt and small vectors)**.** *The map $\varphi - 1 : W_m(L) \to W_m(L)$ is onto, with kernel $W_m(\mathbb{F}_p)$. Its restriction to Q is a bijection. Thus if $(\varphi - 1)x \in Q$, there is a unique $c \in W_m(\mathbb{F}_p)$ with $x - c \in Q$.*

Proof. For $m = 1$, the equation $x^p - x = a$ has a root in the algebraically closed field L and its kernel is $\mathbb{F}_p$. For higher m, use the exact additive sequences given by truncation to length $m - 1$ and its kernel (identified with L by the last coordinate). Frobenius commutes with truncation and acts on the last coordinate by p-th power. Inductively solve the equation after truncation, lift the solution to length m, and correct its error in the last coordinate using the length-one equation. The Frobenius-fixed Witt vectors are precisely those all of whose coordinates lie in $\mathbb{F}_p$.

For $a \in Q$, the sum $y = -\sum_{n=0}^{\infty} \varphi^n(a)$ converges in the valuation topology of the truncated Witt vectors. Indeed every coordinate of $\varphi^n(a)$ tends to zero, all with valuation tending to infinity. Witt addition in the finite number of coordinates is continuous, and for each $\varepsilon > 0$ the subgroup of vectors whose coordinates have valuation at least ε is closed, by the integral polynomial formulas for addition and negation. Thus the tails of the sum tend to zero; completeness gives a limit. The valuations of all coordinates in the limit are still bounded below by a positive number (the minimum of the starting coordinate valuations that are finite), so $y \in Q$. Subtracting adjacent terms from the convergent sum shows $(\varphi - 1)y = a$. Uniqueness follows since $Q \cap W_m(\mathbb{F}_p) = 0$. Finally, if $(\varphi - 1)x \in Q$, solve that equation in Q

and subtract the solution; the difference is in the constant kernel just computed. The same zero-intersection assertion gives the uniqueness of its constant representative. □

The small part is open in the valuation topology on the finite Witt coordinates. If a Galois-dependent expression in them varies continuously and is required to have a representative among the constants modulo Q, that representative is locally constant: a sufficiently small change in the expression belongs to Q, and two distinct constants do not differ by an element of Q. In a tensor expression with a finite reduction of a Galois lattice, the action on that finite reduction is locally constant, and one reduces continuity to the finitely many Witt coordinates. This gives the continuity needed to use the Artin–Schreier–Witt correction in Section 49.6. The precise formal counterpart includes the passage to the diagonal tensor with the finite Tate module in `TateTwist.smallPart_spec`, `Lib/PadicHodge/WachGamma/KisinModuleDefs.lean`. The elementary proof here concerns the truncated Witt algebra and its topology; it should not be confused with an assertion that every integral period-ring primitive or invariant is automatically continuous in the topology of the full period ring.

50.7 Exact cokernel multipliers via finite differences

We expand the arithmetic in Proposition 49.6. In this subsection $J_l(i; r_0)$ denotes exactly its multiplier ideal: multiplication by a scalar in this ideal, up to the modulus r_0, permits the lift of every constant of letter i to standard exponents of level l. The letter subscript is essential in a Frobenius step. In writing A_{inf} for its coefficient ring we again identify it with A above.

There is an elementary Nakayama argument that removes a large modulus but must be used with the precise power. If $p^c \in J_l(i; r_0)$ with $r_0 = p^{c+1}$, write $p^c = x + p^{c+1}y$ for $x \in J_l(i; 0)$ and $y \in A_{\mathrm{inf}}$. Then $p^c(1 - py) = x$, and $1 - py$ is a unit of the p-adically complete ring. Thus $p^c \in J_l(i; 0)$ and $J_l(i; p^{c+1}) = J_l(i; 0)$, since p^{c+1} already belongs to it. Importantly this argument does not work if the modulus on the right is only p^c. The strict difference of one in the powers of p accounts for the $+1$ in the proof of the unramified cokernel estimate.

We recall more explicitly the finite difference behind part (ii). For $v > 0$, the chart calculation gives $\alpha, \beta \in A_{\mathrm{inf}}$ such that

$$\alpha([\epsilon]^v - 1) = E\tau(p^{v_p(v)} + \beta z). \tag{50.11}$$

This is the precise scalar-lifting statement `exists_tau_mul_E_eq_mul_epsTilt_pow_sub` in `Lib/PadicHodge/WachKisin/PeriodMatrixBasis.lean`. To understand the valuation in it, factor $[\epsilon]^v - 1$ first by $[\epsilon] - 1 = E\tau v_0$ with v_0 a unit. At first order in $[\epsilon] - 1$ the other factor has constant term v. The lifting-the-exponent argument of the source uses the relation between ξ and (p, z) to remove the prime-to-p factor of v and control the residual term by z in the needed quotient. The exact existence of the coefficients in (50.11), rather than a valuation identity just in a field, is used below.

If $f = p^c z^v$ and $g(z) = [\epsilon]z$, multiplication of (50.11) by $p^c z^v$ gives

$$\alpha(gf - f) = z\tau E(p^{c+v_p(v)} z^{v-1} + \beta f).$$

Apply the finite-difference property of the ideal (`mem_liftIdeal_of_derivation`) to this identity. Thus $p^{c+v_p(v)} z^{v-1} \in J_1(i; r_0)$ whenever $p^c z^v$ belongs. This property uses the $z\tau$-divisibility of the lifted Galois differences and the coefficient retraction, and is stronger than the assertion that an ordinary ideal is invariant under Galois. Induction lowers v to pa for any $pa \le v$, with cumulative extra p-power $\sum_{pa<j\le v} v_p(j) = v_p(v!) - v_p((pa)!)$. Applying the inverse Frobenius step transports the letter i^p to i and the power z^{pa} to z^a. Multiplying by

any remaining power of p in the ideal proves the stated form of Proposition 49.6(ii). For part (iii), the analogous depth-step calculation for E^n replaces one E by one z, with the bound $\delta_0 + v_p(n)$ (here the chart's specified different bound occurs). Iterating adds precisely $\delta_0 n + v_p(n!)$. The leading coefficient needed for this step comes from `exists_depth_step`, used inside `mem_liftIdeal_of_mul_E_pow`. Thus (ii) and (iii) reduce to these two specialized scalar and ideal-derivation properties; no extra implicit constants are permitted in the factorial sums.

We can now prove the two remaining arithmetic deductions of that proposition. We state the proof because it yields the strict coefficient of a in z^a and the special coefficient $p-3$ of the unramified weight bound.

Proposition 50.5 (A minimal exponent bound)**.** *Assume all allowed jumps are at most n, and put $r_0 = p^\kappa$. Using the frame inclusions and the Frobenius and finite-difference properties of the multiplier ideals, there exists an integer $a \geq 0$ with*

$$z^a \in \bigcap_i J_0(i; r_0), \qquad (p-1)a \leq e_0(n+\kappa).$$

Proof. The integral frame inclusion (with its scalar τ^h for a bounded weight range of the fixed extension) yields a power τ^h in the exact standard-exponent multiplier ideal of every letter. For each fixed κ, this implies that some power of z is in the intersection of those ideals modulo p^κ. Indeed in A_{inf}/p the class of τ has positive valuation and is nonzero. A sufficiently high power of z is divisible by τ^h there, so a power of z belongs to $(p, \tau^h)A_{\mathrm{inf}}$. Raising this containment to the power κ puts a higher power of z in $(p^\kappa, \tau^h)A_{\mathrm{inf}}$. Let a be the least exponent for which all letters belong modulo p^κ. This also covers $\kappa = 0$, where the ideal already contains the modulus 1 and $a = 0$.

By the standard-exponent shift in Proposition 49.6(i), $E^n z^a$ lies in every level-one ideal $J_1(i; r_0)$. From (50.1), $z^{e_0} \in (p, E)$ in A_{inf}. Hence

$$z^{e_0(n+\kappa)+a} \in z^a (p, E)^{n+\kappa} \subset (p^\kappa, E^n z^a) \subset J_1(i; r_0),$$

where in the middle containment each monomial either has E-exponent at least n or p-exponent at least κ. Set $N = e_0(n+\kappa) + a$ and $q = \lfloor N/p \rfloor$. The factorials $(pq)!$ and $N!$ have the same p-valuation, since none of the integers $pq+1, \ldots, N$ is divisible by p. Applying part (ii) of Proposition 49.6 with zero cost sends $z^N \in J_1(i^p; r_0)$ to $z^q \in J_0(i; r_0)$. This holds for all i, because Frobenius permutes the letters. Minimality implies $a \leq q \leq N/p$, or $(p-1)a \leq e_0(n+\kappa)$. □

The proof uses an arbitrarily large initial height h only to show that the set of possible exponents is nonempty. Its numerical conclusion is independent of h; it is the minimization and the Frobenius operation, not an estimate for that initial height, which reduce the final upper bound to n. This illustrates the value of applying the arithmetic reasoning to exact ideals rather than assigning an ad hoc approximation to each individual constant.

Proposition 50.6 (Unramified cokernel bound)**.** *Suppose $e_0 = 1$, $E = z + p$, $p \geq 5$. If all allowed jumps are at most $(p-3)(s+t)$ for nonnegative integers s, t, then $p^s E^t \in J_0(i; 0)$ for every letter i.*

Proof. Put $L = s + t$ and work at modulus $r_0 = p^{L+1}$. Proposition 50.5 gives a nonnegative integer b such that z^b belongs to every $J_0(i; r_0)$ and

$$(p-1)b \leq (p-3)L + L + 1 = (p-2)L + 1.$$

Legendre's valuation formula implies

$$(p-1)v_p(b!) \le b, \qquad (p-2)(b+v_p(b!)) \le (p-1)b \le (p-2)L+1.$$

For the middle inequality, use $(p-2)v_p(b!) \le b$. Since $p-2>1$ and the summands are integral, $b+v_p(b!) \le L$.

We claim that for every $0 \le m \le L$,

$$p^{L-m}z^m \in J_0(i;r_0). \tag{50.12}$$

If $m \ge b$, ideal closure proves the claim. If $m<b$, transport z^b by Frobenius to $z^{pb} \in J_1(i^p;r_0)$. The difference of factorial valuations needed to lower pb to pm is

$$v_p((pb)!) - v_p((pm)!) = b - m + v_p(b!) - v_p(m!) \le L-m,$$

using $v_p((pq)!) = q + v_p(q!)$. Part (ii) of Proposition 49.6 now proves (50.12). For $m=0$ this puts p^L in the ideal modulo p^{L+1}, so the Nakayama step at the start of this subsection shows $J_0(i;r_0) = J_0(i;0)$. Finally expand

$$p^s E^t = p^s(z+p)^t = \sum_{m=0}^{t} \binom{t}{m} p^{L-m}z^m,$$

each term of which belongs to the exact ideal by (50.12). This proves the assertion. □

The last two arguments follow respectively from `exists_z_pow_mem_liftIdeal` and `killsCokernel_of_isUnramified`, `New/Lift/CokernelKillers.lean`. In particular the proof does not use an unspecified "large enough power" of p at the last step. If t is positive, the conclusion can be interpreted as allowing lifts over p^s at the expense of a deeper E-exponent at the first place, by dividing by E^t in the localized module. This is the exact trade-off between the cost of a class and the exponent safety test of Section 49.6. The conversion of multiplier ideal membership to points with the specified depth, and the improvement of depth at no cost under NM, are the more specialized lift/depth inputs in `New/Lift/LongGap.lean`.

50.8 A rank-sensitive bound for raising filtered conditions

We now give further details of the length mechanism for Theorem 49.7. Let L denote a finite free $\mathbb{Z}_p$-module of constants, and $D \subset L$ the submodule of constants with admissible lifts satisfying specified depth conditions. Such submodules need not be O-linear before the final intersection, because the coefficient retraction and intermediate linear-combination lemma are over $\mathbb{Z}_p$. It is therefore appropriate to do the following count over $\mathbb{Z}_p$, converting the resulting lengths by (49.1) only at the end.

Lemma 50.7 (Length from relations)**.** *Suppose $p^cL \subset D$, and for every list of $q+1$ elements of L there is a linear combination with coefficients in $\mathbb{Z}_p$, at least one equal to 1, whose value belongs to D. Then* $\operatorname{length}_{\mathbb{Z}_p}(L/D) \le cq$.

Proof. Decompose the finite quotient over the DVR as a direct sum of nonzero cyclic modules $\mathbb{Z}_p/p^{a_j}\mathbb{Z}_p$ with $a_j \ge 1$. Each $a_j \le c$. If there were $q+1$ of them, taking lifts of their generators would give $q+1$ classes independent in $(L/D)/p(L/D)$. A relation of the stated type would contradict their independence. Thus there are at most q summands and their length is at most cq. □

In the integral comparison, a single filtration step at index a and jump u has $q = f_0 m(i[-a], u)$, where $m(i[-a], u)$ is the corresponding jump multiplicity. We explain algebraically why this dimension, not the rank of the full extension lattice, governs the count. Fix a letter i and an exponent system o, strictly decreasing from the lower jumps to the selected jump u at that index. When the single exponent $o(a, u)$ is increased by one, only the leading coefficients in the jump-u quotient have to be cancelled: the forms of lower types already satisfy the stricter bound. They can be read as a leading coefficient vector of dimension at most $m = m(i[-a], u)$ over the chart's value field. For a list x_j of $f_0 m + 1$ points satisfying the original exponents, choose a maximal independent sublist of their leading coefficient vectors for which every remaining vector is an integral combination of the selected ones. Such a choice is possible by the following elementary argument. If the leading coefficient vectors lie in a valued field's vector space of dimension $r \le m$, fix coordinates in their linear span. Of all the r-tuples choose one whose nonzero determinant has maximal absolute value. Every coordinate of the expansion of any other vector in this basis is a ratio of determinants, by Cramer's rule; replacing any one column cannot increase that absolute value. Hence each ratio lies in the valuation ring. If all vectors are zero the selected list is empty. This is the non-Archimedean basis selection used by the source under the name `exists_auerbach`.

The chart comparison identifies and lifts these integral coefficients in such a way as to write, in the localized module,

$$x_j - \sum_{k=1}^{r} s_{kj} x_{b_k} \in M', \qquad s_{kj} \in S,$$

up to a summand EM; here M represents the original exponent module and M' the module with the one extra step. More precisely the source establishes that the difference, after coefficient retraction, lies in M', and that the discrepancy of the coefficient lift from S is accounted for by the summand EM. This is the value-field and filtration part of `exists_comb_of_jumpMultiplicity_le` in `New/Lift/LongGap.lean`. It is an integral assertion, involving not just a change of leading-coefficient basis over a field, and we retain it as a specialized input. The remaining linear algebra is as follows. Since the chart is unramified, reduction modulo $E = z + p$ gives $S/(E) \simeq \Lambda$ and every s_{kj} can be written uniquely modulo E as $w_{kj} + E y_{kj}$, with $w_{kj} \in \Lambda$. For instance the residue is the value of the convergent series at $-p$. The map

$$\mathbb{Z}_p^{f_0 m+1} \longrightarrow \Lambda^r, \qquad (c_j)_j \longmapsto (\textstyle\sum_j c_j w_{kj})_k$$

has a nonzero kernel over $\mathbb{Q}_p$, since $[\Lambda : \mathbb{Z}_p] = f_0$ and $r \le m$. Clear denominators and divide the relation by the coefficient of least valuation. This gives a relation with all coefficients integral and one coefficient equal to 1. Substituting in the points gives a combination in $M' + EM$, as desired. Notice the factor f_0, which comes from the rank of the unramified chart's scalar residue, not from the number of letters.

There is a further important distinction: the EM error is not automatically zero on lifting constants. One needs the depth operator and the auxiliary no-multiples condition to remove it. In the unramified chart the elementary scalar congruence

$$E^{p^a} - E_a \in pS \tag{50.13}$$

follows from $E = z + p$ and $E_a = z^{p^a} + p$, since both reduce modulo p to z^{p^a}. If no multiple of p lies strictly between the current depth at E and that depth plus p^a, one can use successively the zero-cost operators at E to replace an EM error by an $E^{p^a} M$ error modulo deeper conditions. Relation (50.13) allows this to contribute in $E_a M + pM$. In the proof of `relLength_deepConstants_le`, the precise correction uses `exists_reaches_deep` to change the lifts while keeping the prescribed constants and uses the kernel exponent lattices within

the extension. This is the remaining step needed to pass from a relation in $M' + EM$ to a relation on the actual constants in D'. Thus one may not replace the actual constants by the full extension lattice when applying Lemma 50.7. The arithmetic after this correction is exact: the exponent step costs $c_p(a, n)$, the multiplicity bound gives q, and its loss is at most $qc_p(a, n)$.

We unfold the sums of such steps into the actual two-window bound. For this purpose assume the conclusions of the preceding integral relation/depth comparisons (precisely `relLength_deepConstants_le` and the initial depth construction in `exists_relLength_deepConstants_le` in `New/Lift/FewLowJumps.lean`). Write $A_0 = s$ for the budget in (49.15), η for the width of the low window, W for W_0, and b for β_i at the present letter. The steps compare depth systems c, c' differing by increases at a only, with any decreases at the other divisors allowed. Suppose the new depth at a is antitone as a function of the jump, the old depth is at least $-u$, and no nonzero multiple of p arises strictly between $c(0, v)$ and $c(0, v) + p^a$ for the old system. If for each jump u at $i[-a]$,

$$\sum_{n \in [c(a,u),\, c'(a,u))} c_p(a, n) \le A_1 + B_1 \mathbf{1}_{|u| \le \eta},$$

then the colength paid for the depth change is at most $A_1 r_{i[-a]} + B_1 n_{i[-a]}$. Here n_j and r_j bound respectively the low and total counts, multiplied by f_0, as in (49.16). To see the sum, raise one coordinate at a time, always using the smallest jump requiring a further increase so that the strictness in the lower-jump conditions is preserved. Each step is bounded by the preceding application of Lemma 50.7 after the depth correction. Multiplying the separate bounds by the multiplicities and summing yields the expression $A_1 r_j + B_1 n_j$.

The initial construction of constants of depth $2s + \eta - u$ at $a = 0$, remaining depth $-u$ elsewhere, costs at most sn_i. The construction uses the exact unramified cokernel for $p^v E^{s-v}$, $0 \le v \le s$, and zero-cost no-multiple depth operators, then successively imposes the low negative jump conditions for $-\eta \le u < 0$. At the u-step multiplication by p^s gives the annihilator and the relation bound is the corresponding jump count. In particular the initial depth construction's cost is s times this small count, and is already sufficient for the positive sign. The correctness of those lifts in the localized extension is the initial-depth part of the formal-source theorem just cited, not a consequence of the free-module rank argument alone.

For the negative sign the final required depth at $a = 0$ is $b + W + 2s$. Choose the integer q with

$$b + W + 2s \le qp < b + W + 2s + p.$$

First strengthen the bound at $a = 0$ to the constant qp. The sum of the possible nonnegative step costs on a low jump is at most $v_p((qp)!)$. The window condition gives $qp < (p - 1)s + p$, so

$$v_p((qp)!) \le \frac{qp}{p-1} < s + \frac{p}{p-1} < s + 2,$$

and this integer is at most $s + 1$. On a far negative jump, the starting depth $2s + \eta - u$ lies near b. Every step up to $b + W + 2s$ lies within the interval of radius $6(s + \eta + W)$ about b and hence has zero cost by NM. Every remaining step below qp also has zero cost, since qp is the first multiple of p at or above $b + W + 2s$. If the starting depth exceeds qp, there are simply no steps to pay. Thus this operation costs at most $(s + 1)n_i$. The choice to end at a multiple of p is deliberate: no multiple lies strictly between qp and $qp + p$. This supplies the auxiliary no-multiples condition at $a = 1$. If one had first chosen the exact desired depth $b + W + 2s$, the auxiliary condition would in general fail.

Strengthen the bound at $a = 1$ from $-u$ to

$$W + 1 - u + 2s\mathbf{1}_{|u| \le \eta}$$

at letter $i[-1]$. Each step lies within one of the no-multiples windows about $0, \beta_{i[-1]}, -\beta_{i[-1]}$. The number of steps is $W+1$ on a far jump and $W+1+2s$ on a low jump. Since the chart is unramified, every such step has cost $c_p(1,n) \le 3$ (with $v_p(0)$ set to zero). We can now weaken the bound at $a=0$ from qp to $b+W+2s$ with no additional cost. The total bound is

$$sn_i + (s+1)n_i + 3(W+1)r_{i[-1]} + 6sn_{i[-1]},$$

which proves the arithmetic part of (49.16), conditional on the exact relation and operator statements. Summing over all letters gives

$$(8s+1)\sum_i n_i + 3(W+1)\sum_i r_i \le 11\sum_i \big((s+1)n_i + (W+1)r_i\big).$$

The Frobenius letter permutation is essential in the equality after the summation. The intersection over the letters of the preimages of these deep-constant modules has at most that colength, by the injection into the direct sum of their separate quotients. After the length normalization this yields the precise coefficient $11f_0$ of the few-low-jumps theorem, once the points are verified to meet the safety inequalities. That last verification, including the two zero-loss alternatives in Theorem 49.7, remains the specific formal-source input `exists_scaledCupBounds_few_low_jumps`, together with `scaledCupBounds_of_killsCokernel` in `New/Lift/LongGap.lean`. The argument explains analytically why a narrow far window, even at a very large absolute weight, costs $O(r(W+1))$; the cost proportional to s need only be charged to the low multiplicities. A crude bound for the whole rank, or replacing the low-count term by an annihilator of every constant, would lose this advantage.

Chapter 51

Integral crystalline comparison, depth operators and the Frobenius walk

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 53 and the final audit in Chapter 57.

The use of crystalline representations in the lower bound is unusually quantitative. It asks for a comparison of finite congruence classes with torsion coefficients which is nearly insensitive to the rank and to large isolated Hodge–Tate weights. Neither the equality of the dimensions of two rational period spaces nor a bound linear in the largest Hodge–Tate weight has this property. In this chapter we give the underlying comparison and lifting arguments in substantially more detail. We use the same conventions as Chapters 49 and 50. The present account separates the ring-theoretic descent, the construction of a frame, the elementary moving operators, the Taylor coefficient at a long gap, and the cup-product argument. In particular it identifies which assertions require comparison theorems for the period rings and which follow from elementary algebra once those theorems have been proved.

We give references to both layouts of the formal development where useful. The compact source places the local lifting results in `New/Lift/`; the expanded readable source separates them under `New/Congruences/Lifting/`. The entries called `withOrders` in the expanded source are the exponent lattices called `withExponents` in the compact source. Likewise `IsScaledCupCoboundary` and `ScaledCupBounds` designate the corresponding conclusion about a cup product. We reconstruct that conclusion as a statement about continuous cohomology, not as an assertion about pointwise vanishing of a particular cocycle.

51.1 The period rings and an integral contraction

Fix a prime p. Put $C_p = \widehat{\overline{\mathbb{Q}}_p}$, $A = W(\mathcal{O}^\flat_{C_p})$, and write φ for Witt Frobenius. Choose a compatible system ϵ of primitive p^n-th roots of unity; let $\mu = [\epsilon] - 1$, $\tau = [\epsilon^{1/p}] - 1$, and $\xi = \sum_{j=0}^{p-1} [epsilon^{1/p}]^j$. Thus

$$\varphi(\tau) = \mu = \xi\tau, \qquad \xi - p = \tau w \quad (w \in A).$$

The equality on the right follows by expanding the finite geometric sum in powers of τ. We use the rings

$$A_{\max} = \widehat{A[\xi/p]}^{\,p}, \quad R = A_{\max}[1/p], \quad B = R[1/\mu].$$

The last ring is the version of the crystalline period ring used in the formal source. We use B_{dR}^{+} with fraction field B_{dR}, and put $\mathrm{Fil}^{j} = t^{j}B_{\mathrm{dR}}^{+}$ for any integer j, where t generates the maximal ideal. The cyclotomic convention is $\varphi(\tau) = \mu$, not $\varphi(\mu) = \tau$.

Here are the elementary period-ring properties needed in the contraction. We state explicitly the structural facts used:

(i) A and R are domains, $A \hookrightarrow A_{\max} \hookrightarrow R$, A is (p,ξ)-adically complete and separated, $A_{\max}$ is p-adically complete and separated, and φ acts on them, agreeing on the overlaps.

(ii) Fontaine's map $\theta : A \longrightarrow \mathcal{O}_{C_p}$ is surjective, with kernel (ξ); its extension to R has kernel (ξ). In particular if $a \in A$ becomes divisible by ξ^{m} in R it already is divisible by ξ^{m} in A.

(iii) Write $X = \xi/p \in A_{\max}$. If $Xb \in p^{n}A_{\max}$ for $b \in A_{\max}$, then $b \in p^{n}A_{\max}$. For $q = \tau$ and $q = \xi$, every $x \in A_{\max}$ and $n \geq 0$ satisfy $p^{n}x \in A + q^{n}A_{\max}$. Furthermore every $x \in A_{\max}$ can be approximated modulo $p^{k}A_{\max}$ by an element whose appropriate p-power multiple belongs to A: for each k there are m, a, w with $a \in A$, $w \in A_{\max}$, $p^{m}x = a + p^{m+k}w$.

The second implication in (ii) requires no separate divisibility theorem beyond the kernel calculation: if $a = \xi b$ in R, then $\theta(a) = 0$, so $a = \xi a'$ in A and cancellation in R gives $a' = b$; iterate. We include an explanation of (iii). At the polynomial stage use the binomial theorem. Since $\xi = p + \tau w$, $p^{n}P(\xi/p)$ has an integral polynomial remainder modulo τ^{n}: expand $P(1+\tau w/p)$; the terms of degree $j < n$ contain $p^{n-j}\tau^{j}w^{j}$. For $q = \xi$ use the expansion directly in powers of ξ/p. Passing from polynomials to the completed ring requires an integral closure statement. The approximation in the last sentence of (iii) follows by density of the polynomial subring in $A_{\max}$ and clearing the denominators of one polynomial. The first property of (iii) is the regularity of the image of X in $A_{\max}/p$. In the expanded source these facts are parts of `Lib/PadicHodge/PeriodRings/CrystallineRings.lean`, particularly `Amax.exists_p_pow_mul_eq_add`, `p_pow_dvd_of_dvd_aeval_mul`; they use the presentation as a completion of the ring with $pX = \xi$. We retain the integral structural properties in (i)–(iii) as period-ring inputs. Their consequence for μ has the cost $2n$ instead of n: first write $p^{n}x = a + \tau^{n}w$, then $p^{n}w = a' + \xi^{n}w'$, so $p^{2n}x = p^{n}a + \tau^{n}a' + \mu^{n}w'$. In particular, using the last approximation of (iii) with sufficiently high k gives

$$p^{m}y = a + p^{m+k}\mu^{n}w' \qquad (y \in R,\ a \in A,\ w' \in A_{\max})$$

with some m depending on y, k, n. Indeed first choose $p^{d}y \in A_{\max}$, approximate it by a polynomial up to $p^{d+k+2n}A_{\max}$, and clear polynomial denominators by p^{m-d}. Replace the normalized error w by its two-stage decomposition just given. This is `BPlus.exists_Ainf_approx`, used in Corollary 51.7. The subsequent contraction argument itself will not require a bound on a matrix determinant.

Lemma 51.1 (Contraction of powers of p)**.** *Under the preceding structural properties,*

$$A \cap p^{m}A_{\max} = (p,\xi)^{m}A \qquad (m \geq 0).$$

Proof. The containment from right to left follows from $\xi = pX$. For the other direction suppose $a \in A \cap p^{m+1}A_{\max}$. Since θ extends to $A_{\max} \longrightarrow \mathcal{O}_{C_p}$, $\theta(a)$ is divisible by p^{m+1} there. Surjectivity of θ gives $c \in A$ with $a - p^{m+1}c = \xi b$ for $b \in A$. In $A_{\max}$ cancellation of p gives $Xb \in p^{m}A_{\max}$, so $b \in p^{m}A_{\max}$ by (iii). Induction gives $b \in (p,\xi)^{m}$. Hence both summands $\xi b, p^{m+1}c$ belong to $(p,\xi)^{m+1}$. The assertion starts at $m = 0$. □

Theorem 51.2 (Uniform Frobenius contraction)**.** *Let $V \subset A_{\max}$ be a subset, let $u \in A$ and $n \geq 0$. Suppose that for every $v \in V$ there exists $w \in V$ with $\xi^n v = u\varphi(w)$ in $A_{\max}$. Then $p^{n+1}V \subset A$.*

Proof. For each $k \geq 0$ and each $v \in V$ we construct $a_k \in A$ and $z_k \in A_{\max}$ with

$$p^{n+1}v = a_k + \tau^{n+1}p^k z_k.$$

For $k = 0$ this is (iii) applied to v. Given v, choose a predecessor $w \in V$ as in the hypothesis and choose the k-th approximation $a_k(w), z_k(w)$. Apply $u\varphi$ to it:

$$\begin{aligned} u\varphi(a_k(w)) &= p^{n+1}\xi^n v - u(\xi\tau)^{n+1}p^k\varphi(z_k(w)) \\ &= \xi^n\big(p^{n+1}v - \tau^{n+1}p^{k+1}uX\varphi(z_k(w))\big). \end{aligned}$$

The left side belongs to A, the final expression to $\xi^n R$. Property (ii) permits division by ξ^n in A. This proves the next approximation with $z_{k+1} = uX\varphi(z_k(w))$. For $j \geq k$ the two approximations satisfy $a_j - a_k \in p^k A_{\max}$. By Lemma 51.1 they are congruent modulo $(p, \xi)^k A$. Thus they converge to an element $a \in A$. Their images converge in $A_{\max}$ to $p^{n+1}v$, since $(p, \xi)^k A \subset p^k A_{\max}$. Separatedness gives $a = p^{n+1}v$. □

This is the argument of `exists_Ainf_eq_of_contraction`. It is uniform in V and in the possibly non-invertible scalar u. It is an important device because a preliminary bound after passage to $A_{\max}$ may be quite ineffective in A. The contraction removes that difficulty with the extra cost p^{n+1}, even though one has no uniform bound on the number of elements in V. The factor ξ gained at the induction step is what turns a fixed denominator into an adically convergent series; cancellation of ξ^n in the original integral ring is essential.

51.2 The period-basis argument

We formulate an algebraic result underpinning `IsPeriodMatrix.exists_basis` and `IsPeriodMatrix.exists_frame`. Use the chart $S = \Lambda[[z]] \subset A$, E associated to ξ, where Λ is finite unramified over $\mathbb{Z}_p$. Denote the ring of scalar period coordinates by $S_B \subset R$. Recall that S_B is defined in the source by a bicommutant condition: multiplication by its elements commutes with the extensions of continuous S-linear endomorphisms of A preserving the integral period ring. We need the following properties of this construction.

(a) S_B contains S and p^{-1}. A chosen topological S-basis (b_j) of A, with coordinate functions $\lambda_j : A \to S$, has extended coordinate functions $\widehat{\lambda}_j : R \to S_B$. These extend the original maps, are S_B-linear for multiplication, and finite sums $\sum_{j \in J} b_j \widehat{\lambda}_j(y)$ approximate $y \in R$ modulo arbitrarily large powers of p times $A_{\max}$.

(b) The extensions of S-linear endomorphisms of A preserve $A_{\max}$; they commute with Frobenius conjugation in the sense that the extension of $\varphi^{-1} f \varphi$ satisfies $\varphi \widehat{f'} = \widehat{f}\varphi$. Here φ is bijective on A, and f' is S-linear for the Frobenius-stable chart.

(c) The principal ideal (q) for $q \in A$ with $p \nmid q$ is closed in the (p, ξ)-adic topology. Also $\bigcap_j z^j A = 0$.

(d) If d divides μ in A, any fixed matrix M over R has a bound k such that

$$My = d^h w, \quad y \in A_{\max}^r,\ w \in R^r \quad \Rightarrow \quad p^k w \in A_{\max}^r.$$

The k may depend on M, d, h but is independent of y, w. Finally $\varphi(\tau)^h \mid \varphi(b)$ implies $\tau^h \mid b$ for $b \in R$.

These properties, with the integral comparison of kernels, are the structural inputs of the period-basis proof, not assertions about every infinite S-module. For (d), the nonzero divisor d divides μ and the latter has a controlled reduction in $A_{\max}$; a denominator selecting M first suffices, and division by d^h costs a fixed extra p-power in $A_{\max}$. The source proves the required statement in `exists_p_pow_mul_mem_amaxImage`. For (c), closure of (q) can also be seen by the same successive reduction modulo p used to prove completeness of A in its two-generator topology: reduction of q is nonzero in the valuation domain $\mathcal{O}^\flat_{C_p}$ whose nonzero principal ideals are closed for the topology of any element of positive valuation. Lifting the compatible quotients successively modulo p^n requires the integral closure of the image of a nonzero principal ideal at each finite Witt length. We use the source's closure lemma for this last integral step. These are the only asserted inputs to the next theorem. They are supplied by `Lib/PadicHodge/Kisin/PeriodMatrixBasis.lean` and the chart and period-ring constructions it imports.

Let $U, W \in M_d(R)$ obey $UW = (E\tau)^h I$ and suppose U is a period matrix in the following sense:

$$\varphi(U) = vUF, \qquad v \in A^\times, \quad F \in \mathrm{GL}_d(S_B),$$

and whenever $Uy = E^h x$ with $x \in A^d$, then $U\widehat{f}(y) \in E^h R^d$ for all coordinate endomorphisms $f : A \to A$ considered in (b). We assume $\widehat{f}$ is S_B-linear and that $\varphi(E) \in S_B^\times$. The period-matrix construction of a crystalline representation with weights in $[-a, b]$ supplies $h = a + b$ and the unit factor v (see the more explicit calculation in Section 50.4). In the following, we treat the existence and the stability of the period matrix as inputs; its passage to an actual integral frame will be proved.

Lemma 51.3 (Integral bound for all coordinate transforms)**.** *There exists an integer N such that, for every solution $Uy = E^h x$ with $y \in R^d$, $x \in A^d$, and every f as above, if $U\widehat{f}(y) = E^h w$, then $p^N w \in A^d$.*

Proof. The relation $UW = (E\tau)^h I$ implies injectivity of U and also $WU = (E\tau)^h I$. First $W(E^h x) = (E\tau)^h y$, so (d) supplies one fixed p^a putting all such y into $A^d_{\max}$. The extensions preserve this integral ring. Applying (d) to $U(p^a \widehat{f}(y))$ puts every w into $A^d_{\max}$ after multiplication by the same p^{a+b}, with some fixed b.

For each solution define $x' = \varphi^{-1}(vE^h x) \in A^d$. We claim there is a solution $Uy' = E^h x'$ with $y' \in R^d$. In $\mathrm{Frac}(R)^d$ its unique candidate is $y' = \tau^{-h} W x'$. Applying Frobenius gives

$$F\varphi(y') = \varphi(E)^h y,$$

by cancellation of vU. The right side lies in R^d and F is invertible over R. Thus $\varphi(\tau)^h$ divides $\varphi(Wx')$, and the inverse Frobenius property in (d) gives $y' \in R^d$. Let $f' = \varphi^{-1} f \varphi$ and define w' by $U\widehat{f}'(y') = E^h w'$. The conjugation property of extensions and their linearity over the entries of F and over $\varphi(E)$ give

$$F\varphi(\widehat{f}'(y')) = \varphi(E)^h \widehat{f}(y).$$

Apply φ to the defining equation of w', use $\varphi(U) = vUF$, and cancel $\varphi(E)^h$ to get $\varphi(w') = vE^h w$. Write $\xi = Ee_1$ for a unit e_1 of A. Each coordinate of $p^{a+b}w$ is now in a set $V \subset A_{\max}$ satisfying

$$\xi^h p^{a+b} w = e_1^h v^{-1} \varphi(p^{a+b} w').$$

Theorem 51.2 applies with $n = h$, $u = e_1^h v^{-1}$. Take $N = a + b + h + 1$. □

Theorem 51.4 (A basis and saturated solution space)**.** *Under these assumptions there are $Q \in M_d(A)$ and $Z \in M_d(S_B)$ with $\det Q \neq 0$ and $UZ = E^h Q$, such that*

$$P = \{x \in A^d : \ Uy = E^h x \text{ for some } y \in S_B^d\} = QS^d.$$

For every solution $Uy = E^h x$ with $y \in R^d, x \in A^d$, every row $l \in A^{1\times d}$, and every $q \in A$ not divisible by p, the divisibility $q \mid lQ$ entry by entry implies $q \mid lx$.

Proof. As explained in Section 50.4, P is p-saturated and z is injective on P/pP. We verify the rank estimate before using the finite freeness lemma. Let $x_1, \dots, x_r$ be S-linearly independent elements of P, with corresponding solutions $y_1, \dots, y_r \in S_B^d$. Suppose $\sum_j a_j x_j = 0$ for $a_j \in A$. Since U is injective, this is equivalent to $\sum_j a_j y_j = 0$. Apply the extended coordinate $\widehat{\lambda}_i$; by the commutant property,

$$\sum_j \lambda_i(a_j) y_j = 0.$$

Multiply by U and cancel E^h to obtain $\sum_j \lambda_i(a_j) x_j = 0$. All the coefficients vanish by S-linear independence, for all i. The original coordinates separate A, hence $a_j = 0$ for all j. Thus the x_j are independent even over A and $r \le d$. The separated regular-module lemma, applied to $P \subset A^d$ with the regular sequence (p, z) and this rank bound, gives a finite free S-module (Lemma 50.3). Pad a basis by zero columns to size d, forming Q, and put the corresponding y in Z.

Now let x solve the wider equation $Uy = E^h x$, with $y \in R^d$. For each i, by Lemma 51.3, $p^N E^{-h} U \widehat{\lambda}_i(y)$ lies in A^d. Its solution $p^N \widehat{\lambda}_i(y)$ lies in S_B^d, so the vector can be written Qs_i with $s_i \in S^d$. Choose k from (d) for the matrix U and divisor E. Given m, approximate y by $\sum b_i \widehat{\lambda}_i(y)$ modulo $p^{m+k} A^d_{\max}$, using a common finite set J for its finitely many entries. The integral error

$$t = p^N x - Q \sum_{i \in J} b_i s_i$$

satisfies $p^{-(m+k)} E^h t \in U A^d_{\max}$. Property (d) implies $t \in p^m A^d_{\max}$ and Lemma 51.1 implies $t \in (p, \xi)^m A^d$. If $lQ = qg$, then

$$l(p^N x) - qg \sum_{i \in J} b_i s_i = lt \in (p, \xi)^m A.$$

Since this holds for every m, closure of (q) gives $q \mid p^N lx$. The element p is prime in A, since $A/p = \mathcal{O}^\flat_{C_p}$ is a domain. When $p \nmid q$, cancellation of p^N in this divisibility is justified by repeatedly observing that $p \mid qa$ implies $p \mid a$. Thus $q \mid lx$.

If $\det Q = 0$, choose a nonzero row l over A that annihilates Q, clearing denominators of a kernel row over its fraction field. For every $x_0 \in A^d$ the vector $\tau^h x_0$ is a solution with witness Wx_0. Hence $q \mid \tau^h l x_0$ for every $q = z^m$, since $p \nmid z$. Separatedness for (z) implies $\tau^h l x_0 = 0$. Take x_0 successively to be the standard basis vectors; cancellation of τ gives $l = 0$, a contradiction. In particular no padding was in fact needed: the original basis of P has rank d. □

The point of this proof is not the existence of a rational basis of periods; that was already given by the crystalline hypothesis. Its novelty is that the columns simultaneously span an integral solution space over S and control divisibility of all solutions with coefficients in the larger positive period ring. The argument is a convenient place to separate the structural period-ring lemmas from the finite and adic linear algebra. No bound on the exponent N of Lemma 51.3 is needed for the final divisibility result, since it is removed by the primality of p. In contrast, replacing the integral row-divisibility statement by a rational basis theorem would not give any such cancellation.

51.3 Reduction of the determinant and the integral frame

The matrix Q in the preceding theorem is nonsingular, but its determinant need not be a unit. The weaker property required below is $p \nmid \det Q$. It is stronger than nonsingularity and cannot be deduced from the finite freeness of P alone. We first give the fixed-coefficient argument of `PeriodMatrixFrame.lean`.

Lemma 51.5 (Minimal-support vector)**.** *Let V be a valuation domain and let $M \in M_d(V)$ have zero determinant. There exists $a \in V^d$ in its kernel, with $a_\nu = 1$ for some ν, such that for every unital ring endomorphism g of V,*

$$Mg(a) = 0 \quad \Longrightarrow \quad g(a) = a.$$

Proof. There is a nonzero vector in the kernel by solving over the fraction field and clearing denominators. Choose one with minimal support. In a valuation domain the principal ideals of its nonzero coordinates are totally ordered. Choose a coordinate dividing every other coordinate and divide it out, obtaining $a_\nu = 1$ without changing the support or the kernel equation. If $g(a) - a$ were a nonzero kernel vector, its support would be contained in the support of a but would not contain ν, a contradiction. □

Write $V = A/p = \mathcal{O}^\flat_{C_p}$ and let a bar denote reduction modulo p. We need two precise chart properties:

(i) For $j \geq 0$, let $\pi_j : V \to \mathcal{O}_{C_p}/p$ be the j-th projection of the inverse limit along Frobenius. If $a \in V$ is fixed by every Galois element fixing S, then for each m and all sufficiently large n there is $s \in S$ with
$$\pi_m(a) = \pi_{m+n}(\bar{s}).$$

(ii) If $x \in A$ satisfies $\pi_j(\bar{x}) = 0$, every original coordinate $\lambda_i(x) \in S$ has $\pi_j(\overline{\lambda_i(x)}) = 0$ as well.

The first is the fixed-coefficient approximation in the chosen chart, `Chart.eventually_exists_coeff_eq_of_fixed`; it is an assertion about the arithmetic fixer of the Kummer tower, not just an arbitrary action on a ring. The second is the chart's divisibility-preservation result `exists_frobE_dvd_sub_of_dvd`, applied with multiplier p. We retain (i) and (ii) as chart comparison inputs. They are needed only at the indicated place and at finite Witt length, not as a statement that all fixed elements of the tilt lie in the residue of S.

Proposition 51.6 (Frobenius and p-primitivity)**.** *Suppose $Q \in M_d(A)$ is nonsingular and p-saturated over S, meaning that $Qt \in pA^d$ for $t \in S^d$ implies $t \in pS^d$. Suppose there is $B_1 \in M_d(S)$ with $p \nmid \det B_1$ and a relation*

$$QB_1 = U_1\varphi(Q), \qquad U_1 \in M_d(A).$$

Assume moreover that for each Galois element g fixing S, there is a matrix $U_g \in M_d(A)$ with $Q = U_g g(Q)$. With the preceding two chart properties, $p \nmid \det Q$.

Proof. Consider the rows of Q after applying the coordinate maps λ_i entry by entry. These are rows over S and separate S^d/pS^d. In fact, if every coordinate of Qt is divisible by p in S, then Qt is divisible by p in A: approximate by the topological basis and reduce modulo p, using separatedness of V for the topology of $\bar{z}$. Saturation gives $t \in pS^d$.

The separating rows have rank d over the fraction field of S/pS, which embeds into the fraction field of V. Indeed otherwise a nonzero kernel vector over the former field can be

scaled into $(S/pS)^d$; lifting it contradicts the separation just proved. Thus a choice of d such rows gives a matrix $N \in M_d(S)$ with $\overline{\det N} \neq 0$. Each row of Nt is a specified coordinate of one entry of Qt.

Suppose for contradiction that $\det \bar{Q} = 0$. Apply Lemma 51.5 to $\bar{Q}$ over V, obtaining a with $a_\nu = 1$. Each g fixing S carries a into the kernel of $g(\bar{Q})$ and hence of $\bar{Q}$ by $Q = U_g g(Q)$. Minimality gives $g(a) = a$. Define

$$C_n = B_1 \varphi(B_1) \cdots \varphi^{n-1}(B_1), \qquad D = \overline{\det B_1}\, \overline{\det N}.$$

Induction using the Frobenius relation shows that

$$\bar{Q} \bar{C}_n a^{p^n} = 0 \qquad (n \geq 0),$$

where the power is componentwise. Furthermore

$$D \det \bar{C}_n \mid D^{p^n} \quad \text{in } V.$$

To see this, $\det \bar{C}_n = (\overline{\det B_1})^{1+p+\cdots+p^{n-1}}$ and $1 + (p^n - 1)/(p-1) \leq p^n$. The assertion for $n = 0$ is immediate.

Fix any component m in the inverse limit description of V. By (i), choose a common n large enough and $s \in S^d$ with $\pi_{m+n}(\bar{s}) = \pi_m(a) = \pi_{m+n}(a^{p^n})$. In particular $\pi_{m+n}(\bar{s}_\nu) = 1$. Then $\pi_{m+n}(\bar{Q}\bar{C}_n \bar{s}) = 0$. Property (ii) and the selection of the rows of N give $\pi_{m+n}(\bar{N}\bar{C}_n \bar{s}) = 0$. This is a matrix equation over $\mathcal{O}_{C_p}/p$, which need not be a domain. Nevertheless, multiplication by the adjugate shows that $\pi_{m+n}(\det \bar{N} \det \bar{C}_n) = 0$, because one coordinate of $\bar{s}$ has image 1. Consequently $\pi_{m+n}(D \det \bar{C}_n) = 0$. The displayed divisibility gives $\pi_{m+n}(D^{p^n}) = 0$, i.e. $\pi_m(D) = 0$. Every component is zero, so $D = 0$. This contradicts $\det \bar{N} \neq 0$, $\det \bar{B}_1 \neq 0$ in the domain V. □

This proves `not_dvd_det_of_frobenius` from the precise two chart properties. It also explains why the inverse-limit components cannot be replaced by a single residue-field reduction. The matrix N need not be invertible modulo the maximal ideal of S, only nonsingular modulo p. Taking a component of D far enough back in the tilt, multiplied by Frobenius iterates of B_1, is the way the proof obtains a contradiction without assuming either of these determinants is a unit.

We apply this result to Theorem 51.4. First, Q is p-saturated over S: if $Qt = px$ with $x \in A^d$, then $p^{-1}Zt$ solves $Uy = E^h x$ over S_B. Hence $x = Qs$ for $s \in S^d$ and nonsingularity gives $t = ps$. Next,

$$U \frac{F\varphi(Z)}{\varphi(E)^h} = v^{-1}\varphi(Q)$$

follows from $UZ = E^h Q$. Multiplying by E^h yields a period solution over S_B whose corresponding vector $v^{-1}E^h\varphi(Q)$ is integral; to obtain $v^{-1}\varphi(Q)$ itself, keep the undivided equation $UF\varphi(Z) = v^{-1}\varphi(E)^h\varphi(Q)$ and use $E^h\varphi(E)^{-h}$ in the witness. Thus $v^{-1}\varphi(Q) = QB_1$ for B_1 over S. Taking determinants in $\varphi(Q) = vQB_1$ shows $p \nmid \det B_1$: if p divided $\det B_1$, then $p^k \mid \det Q$ would imply $p^{k+1} \mid \varphi(\det Q)$ and hence $p^{k+1} \mid \det Q$, contradicting nonsingularity and p-separatedness.

The Galois relation used in the proposition derives from the compatibility of a crystalline period matrix with the representation and the scalar normalization. More precisely the source gives $\rho_g g(U) = U$ with ρ_g integral when g fixes the chart constants and $\theta(z)$, after the scalar normalization has been included (`IsPeriodMatrix.exists_gal`). For g fixing S, this hypothesis holds and $g(E) = E$. Furthermore g is an S-linear endomorphism of R preserving $A_{\max}$. As the entries of Z belong to the bicommutant, evaluating the commutation

$g \circ m_z = m_z \circ g$ at 1 for each such entry z gives $g(Z) = Z$. Applying g to $UZ = E^h Q$ and multiplying on the left by ρ_g yields $UZ = E^h \rho_g g(Q)$. Canceling E^h gives $Q = \rho_g g(Q)$. This is `IsPeriodMatrix.exists_eq_mul_smul_of_forall_smul_eq`. With the period-matrix compatibility input, Proposition 51.6 applies.

Corollary 51.7 (Uniform scalar inverses)**.** *Under the above hypotheses, every integral solution of $Uy = E^h x$ with $y \in R^d$ lies in QA^d. There are matrices $L \in M_d(A)$ and $Y \in M_d(R)$ with*

$$QL = \tau^h I_d, \qquad U = QY, \qquad YZ = E^h I_d.$$

Proof. Apply the divisibility statement of Theorem 51.4 to every row of the adjugate of Q with $q = \det Q$; this is permitted by $p \nmid \det Q$. Each component of $\operatorname{adj}(Q)x$ is divisible by $\det Q$ in A. This yields $x = Qx'$ with $x' \in A^d$. In particular the solutions $\tau^h e_j$ with witnesses We_j give L.

To obtain Y, use the approximation modulo μ^h of each entry of U. Choose a common exponent m and matrices $K \in M_d(A)$, $H \in M_d(A_{\max})$ with

$$p^m U = K + p^m \mu^h H.$$

Write $\mu = E\tau d_0$ for a unit $d_0 \in A$. Since $UW = (E\tau)^h I_d$,

$$K = Up^m (I_d - d_0^h WH).$$

Therefore every column of K has a witness for the equation $Uy = E^h x$ in R^d after multiplication of the parenthesized matrix by E^h. We obtain $K = QK_0$ for an integral matrix K_0. Now

$$U = Q\left(p^{-m} K_0 + (Ed_0)^h LH\right).$$

This proves $U = QY$; substituting in $UZ = E^h Q$ and canceling Q yields $YZ = E^h I_d$. □

Thus the part of the frame theorem giving the rank-free scalar denominators reduces to the structural chart comparison, the exact period equations, and the preceding proofs. This removes the abstract basis and determinant assertions previously treated as inputs in Section 50.4. The remaining transformation properties for all Galois elements in the chosen chart, the simultaneous Hodge–Tate pole estimates, and the identification of this span with the intrinsic Kisin module are still necessary comparison statements. In the source the complete `frame_theorem` combines these statements with the integral basis and the calculation above. In particular we may not replace Q by an arbitrary rational basis of a filtered φ-module.

51.4 Scalar depth steps at each Frobenius divisor

Return to the jump lattices of Chapter 49. A Kummer chart has $S = \Lambda[[z]]$ and a subgroup G' of prime-to-p index, acting trivially on Λ. Let $E = z^{e_0} A_0(z) + p$ with A_0 a unit of $\Lambda[[z]]$, let $E_a = \varphi^a E$, and put $\Delta = z\tau$. Write $\theta_a = \theta\varphi^{-a}$. The valuation v_p on C_p is normalized by $v_p(p) = 1$, and for an integer $n = 0$ we put $v_p(0) = 0$ only in the cost function. In particular the convention does not extend v_p as a valuation of the field.

The different bound of the source is

$$d_0 = \begin{cases} 0 & e_0 = 1, \\ v_p(e_0) + 1 & e_0 > 1. \end{cases}$$

We recall the exact chart properties used in the calculation. For every nonnegative m there is $g \in G'$ with $g(z) = [\epsilon]^m z$; every $g \in G'$ fixes $\theta(z)$ and Λ. The ideal (E_a) equals the kernel of

θ_a; Δ is divisible by none of the E_a; $\Delta \mid \varphi^a\Delta$. Write G for the unit satisfying $F(z) = EG$, where $F(X)$ is the monic Eisenstein polynomial of the actual uniformizer $\theta(z)$ over $\Lambda[1/p]$. Then

$$\theta(F'(z)) \mid p^{d_0}, \qquad \theta(G) \in \mathcal{O}_{C_p}^{\times}.$$

For an element $g \in G'$ fixing the base constants, $gE = \omega E$ with $\omega \in A^{\times}$ and $\Delta \mid \omega - 1$. The last relation is the scalar near-identity part of the chart, not a general statement about all Galois actions on A; it is established in `zTau_dvd_sub_one_of_mul_eq_smul_E`. We retain these existence and divisibility properties as Kummer chart inputs. In the unramified chart $E = z + p$ and $F(z) = E$, so the polynomial and derivative assertions are immediate. In a general chart the inequality $v_p(F'(\theta(z))) \le d_0$ follows from the different bound: the leading derivative term has valuation $v_p(e_0) + (e_0 - 1)/e_0$ and the nonzero derivative summands of distinct degrees have distinct fractional parts of their valuations, so the valuation of the sum is the minimum and is no larger than the leading valuation. This proves the claimed divisibility for this polynomial; the unit quotient and the action assertions refer to the precise embedding in A.

Lemma 51.8 (First-order scalar estimate)**.** *Choose g with $g(z) = [\epsilon]^{p^a} z$ and write $g(E) = \omega E$. Then*

$$\theta(\omega - 1) \mid p^{d_0+a}\theta(\Delta) \quad \text{in } \mathcal{O}_{C_p}.$$

Proof. Write $g(z) - z = Eb$. This is possible since g fixes $\theta(z)$. Factor

$$[\epsilon]^{p^a} - 1 = ([\epsilon] - 1) \sum_{j=0}^{p^a-1} [\epsilon]^j = \xi\tau \sum_{j=0}^{p^a-1} [\epsilon]^j.$$

Because E and ξ are associated and $\theta([\epsilon]) = 1$, this gives $\theta(b) = p^a\theta(\Delta)u$ for a unit u of $\mathcal{O}_{C_p}$. Apply the Taylor identity to F:

$$F(gz) = F(z) + F'(z)Eb + E^2b^2H \qquad (H \in A).$$

Since g fixes the coefficients of F, the left side is $g(EG) = \omega E g(G)$. Divide by E and apply θ, using $\theta(gG) = \theta(G)$. We obtain

$$\theta(\omega - 1)\theta(G) = \theta(F'(z))\theta(b).$$

The derivative divides p^{d_0}, and $\theta(G)$ is a unit. Substituting the expression for $\theta(b)$ proves the result. □

This proof is the scalar calculation `theta_sub_one_dvd_pow_mul_theta_zTau` in the readable `ExponentLattices.lean`. At higher places, we use the same calculation after applying inverse Frobenius, because E_a generates that kernel. Keeping the chart polynomial in the argument (rather than replacing E by an arbitrarily associated element) is why a bound in terms of the different remains possible.

Lemma 51.9 (Lifting the exponent at p)**.** *Suppose $\omega \in A^{\times}$, $\Delta \mid \omega - 1$, and $\theta(\omega - 1) \neq 0$. Then, for every nonzero integer n,*

$$v_p(\theta(\omega^n - 1)) = v_p(\theta(\omega - 1)) + v_p(n).$$

Proof. Put $x = \theta(\omega) - 1$. Since $\theta(\tau) = \zeta_p - 1$ and $\theta(z)$ has positive valuation,

$$v_p(x) \ge v_p(\theta(\Delta)) = v_p(\theta(z)) + \frac{1}{p-1} > \frac{1}{p-1}.$$

If $m > 0$ is prime to p, the binomial expansion of $(1+x)^m - 1$ has leading term mx of valuation $v_p(x)$, strictly less than those of all higher terms. For exponent p, the linear term px has valuation $1 + v_p(x)$. The terms $\binom{p}{j}x^j$ for $1 < j < p$ have strictly greater valuation; the last term x^p does also, since $(p-1)v_p(x) > 1$. Thus raising to p increases the valuation by exactly one. Iterate and use $n = p^r m$. Replacing positive n by $-n$ changes $(1+x)^n - 1$ by a unit times its negative and preserves the valuation. □

The strict inequality in this proof matters: the classical formula can fail for a p-th root of unity where $v_p(x) = 1/(p-1)$. Multiplication of the cyclotomic element by the chart coordinate z is what supplies strictness. In the application of Lemma 51.8, $\theta(\omega - 1)$ cannot be zero: the formula expresses it as a nonzero derivative times the nonzero scalar $\theta(b)$ divided by a unit. In particular ω does not have finite order in the residue field of E.

Proposition 51.10 (The scalar depth identity)**.** *For each $a \geq 0$ and nonzero $n \in \mathbb{Z}$, put*

$$c(a,n) = v_p(n) + d_0 + 3a.$$

There are $g \in G'$, a unit $\omega_a \in A^\times$ and $X, V \in A$ satisfying

$$g(E_a) = \omega_a E_a, \quad g(s) - s \in E_a A \ (s \in S), \quad \Delta X = \omega_a^n - 1, \quad VX \equiv p^{c(a,n)} \pmod{E_a A}. \tag{51.1}$$

Proof. Use the element g of Lemma 51.8. It commutes with Frobenius, so $\omega_a = \varphi^a \omega$. Since $g(z) - z = z([\epsilon]^{p^a} - 1)$ and $\theta_a([\epsilon]^{p^a}) = 1$, $g(z) - z \in E_a A$. The same holds for each $s \in S$ by its power-series expression and the exact kernel condition; the continuity and power-series evaluation here are those of the chart. The element $\omega^n - 1$ is divisible by $\omega - 1$, also for negative n (because ω is a unit). Thus $\varphi^a(\omega^n - 1)$ is divisible by $\varphi^a\Delta$, hence by Δ; take X to be the quotient.

It remains to invert X modulo E_a at the indicated cost. Since $\theta_a(\Delta) \neq 0$,

$$v_p(\theta_a X) = v_p(\theta(\omega^n - 1)) - v_p(\theta_a \Delta).$$

By Lemmas 51.8 and 51.9, this is at most

$$v_p(n) + d_0 + a + v_p(\theta\Delta) - v_p(\theta_a\Delta).$$

For $a = 0$ the last difference is zero. For $a > 0$, $\theta(\Delta) = \theta(z)(\zeta_p - 1)$ divides p^2 because $v_p(\theta(z)) = 1/e_0 \leq 1$ and $1/(p-1) \leq 1$; the other valuation is nonnegative. Therefore the difference is at most $2a$ for all a. This proves $\theta_a X \mid p^{c(a,n)}$ in $\mathcal{O}_{C_p}$. The map θ_a is surjective, so lift the quotient $p^{c(a,n)}/\theta_a X$ to $V \in A$. Then the congruence holds by its kernel identity. □

In the compact source this is `exists_depth_step`, and in the readable source it is `exists_isDepthStep`. The proof also identifies the source of all three contributions to the integer $c(a,n)$: the power of p dividing n, the field different, and the shift of the divisor together with the comparison of the two evaluations of Δ. No factor involving the rank occurs.

51.5 Depth zero and finite averaging

The preceding proof does not apply when $n = 0$. The unit eigenvalue for degree zero yields no nonzero factor to cancel. The correct replacement is an averaging operator; its cost is $c(a,0) = d_0 + 3a$ as above.

Lemma 51.11 (Averaging modulo E_a)**.** *There are $p^{c(a,0)}$ elements g_j of G' such that the sum $\sum_j g_j(s)$ is fixed modulo E_a by all $g \in G'$ and $s \in S$.*

Proof. Choose $g_j(z) = [\epsilon]^j z$, $0 \le j < p^{c(a,0)}$. The scalar $\zeta = \theta_a([\epsilon])$ is a primitive p^a-th root of unity, with order one if $a = 0$. At θ_a each $s \in S$ can be expanded as a polynomial in $\theta_a(z)$ with coefficients in the unramified constants, of degree less than $e_0 p^a$. This follows from the chart's monic division by E_a: the evaluated scalar $\theta_a(z)$ is a root of the polynomial obtained from the Eisenstein polynomial of $\theta(z)$ by replacing X by X^{p^a} and twisting coefficients by inverse Frobenius. Division leaves a polynomial of at most that degree; the integral division statement in the chart is `exists_dvd_sub_sum_z_pow`.

For a monomial $s = wz^m$ with $w \in \Lambda$, its sum after evaluation is

$$\theta_a(w)\theta_a(z)^m \sum_{j=0}^{p^{c(a,0)}-1} (\zeta^m)^j.$$

Since $c(a,0) \ge a$, the geometric sum is zero unless $p^a \mid m$. In that case the monomial is fixed by all G': $\theta_a(z)^{p^a} = \theta(z)$, and G' fixes this element and the constants. The sum of the surviving terms is therefore fixed. This is equivalent to the congruence modulo E_a. □

Notice that it is a sum, not an average divided by $p^{c(a,0)}$; the latter would not be integral. The source lemma `exists_frobE_dvd_smul_sum_sub_sum` is precisely this statement.

51.6 From scalar depths to crystalline lattice depths

We make explicit the notation used to apply these scalar identities uniformly to a lattice. A jump lattice T is a finite free $\mathbb{Z}_p$-module with an integral Galois action (and, in the applications, compatible actions of a finite free coefficient ring). Over $W(C_p^\flat)$ we have its ambient module $\mathcal{T}$. The chart identifies a Kisin lattice with its localized space $\mathcal{N}(T)$, over $S[E_a^{-1} : a \ge 0]$. Let $\mathcal{N}_A(T)$ denote its A-span, also with the indicated divisors inverted in the localized coordinates. The fixed chart coefficient ring is denoted by Λ, and the letters i are the factors of the coefficient-ring decomposition, permuted by Frobenius. For a letter i, $i[-a]$ means a inverse Frobenius steps. We write $U_i(T)$ for the finite jump set containing the actual Hodge–Tate jumps. These sets are allowed to include further integers; in particular, 0 occurs in each extension of a trivial lattice.

Every crystalline form of type u at letter $i[-a]$ defines a value $\mathcal{V}_a(\lambda, x)$ in B_{dR} on $x \in \mathcal{N}_A(T)$ at letter i. This is the evaluation of λ after φ^{-a}. For a system $o(a,u)$ of integers, write $\mathcal{M}_i^T(o)$ for the submodule consisting of the x satisfying

$$\mathcal{V}_a(\lambda, x) \in \mathrm{Fil}^{o(a,u)} \quad (a \ge 0,\ u \in U_{i[-a]}(T),\ \lambda \text{ of type } u). \tag{51.2}$$

This is `JumpLattice.withOrders`. We use the following comparison inputs, whose precise strength is important.

(a) Values commute with the G'-action and with multiplication by evaluated scalars $\varphi^{-a}(r)$ ($r \in A$). The modules $\mathcal{M}_i^T(o)$ are G'-stable, and multiplication by E_b^m adds m to the order at $a = b$, for any integer m, without affecting the other order conditions.

(b) There is a coefficient retraction $r_0 : \mathcal{N}_A(T) \to \mathcal{N}(T)$, preserving all the conditions (51.2), commuting with projections onto letters and with equivariant maps between the lattices, and preserving congruences of values to constants as follows. If $\mathcal{V}_a(\lambda, x)$ is congruent to a scalar in the fixed field modulo Fil^n, then the same congruence holds for $r_0(x)$.

(c) For $x \in \mathcal{N}(T)$ and $g \in G'$, the vector $\Delta^{-1}(gx - x)$ belongs to $\mathcal{N}_A(T)$ and, if $gx - x \in \mathcal{M}_i^T(o)$, then the quotient belongs to $\mathcal{M}_i^T(o)$ as well.

(d) Let K' be the field fixed by the chart subgroup. If $y \in B_{\mathrm{dR}}$ is fixed modulo Fil^n by G', then y is congruent modulo Fil^n to an element of K'. Moreover the value of each form on a localized Kisin vector lies in the specified value field for that divisor; whenever g acts trivially on S modulo E_a, it acts trivially on the integral elements of this value field modulo Fil^1. The same holds for the sum in Lemma 51.11: the sum of conjugates of such integral values is invariant modulo Fil^1.

Part (a) follows from the evaluations at the distinct prime divisors and equivariance of forms. Parts (b), (c), (d) are the integral frame and fixed-scalar comparisons: the source anchors are `coordRetract_mem_withOrders`, `valueAt_coordRetract_sub_mem_hodgeFil`, `zTauUnit_inv_smul_sub_mem_withOrders`, `exists_const_sub_mem_hodgeFil_of_smul_sub_mem`, `smul_sub_mem_hodgeFil_of_valueField` in `ExponentLattices.lean` and `HodgeExpansion.lean`. The assertion on K' extends to negative n: it is a statement about fixed representatives in a filtered quotient, stronger than saying $B_{\mathrm{dR}}^{G'} = K'$. For example, a nonzero K'-scalar has filtration order zero, regardless of its p-adic valuation. All remaining derivations are carried out under these comparison statements. The scalar inverse calculations earlier in this chapter explain how uniform frame denominators used in (b),(c) are obtained; deriving the actual comparison of every form in (d) still requires the filtered period-ring theory.

Define the depth condition

$$\mathcal{I}_i^T(o) = \{x : \ gx - x \in \mathcal{M}_i^T(o) \text{ for all } g \in G'\},$$

where x is in the same letter part of the ambient localized span. By (a) and (d), this is exactly the condition that every $\mathcal{V}_a(\lambda, x)$ be congruent to a fixed-field scalar modulo $\mathrm{Fil}^{o(a,u)}$. The reverse implication uses $g(y - c) - (y - c) \in \mathrm{Fil}^{o(a,u)}$ if c is fixed.

Lemma 51.12 (Improvement of positive orders)**.** *Let $x \in \mathcal{M}_i^T(o) \cap \mathcal{I}_i^T(h)$. Then its order system can be improved to*

$$\mathrm{imp}(o,h)(a,u) = \begin{cases} h(a,u) & o(a,u) > 0, \\ \min(0, h(a,u)) & o(a,u) \le 0. \end{cases}$$

Proof. Fix a, u, λ and write $y = \mathcal{V}_a(\lambda, x)$, $o = o(a,u)$, $h = h(a,u)$. We have $y \in \mathrm{Fil}^o$ and $y - c \in \mathrm{Fil}^h$ for some $c \in K'$. Thus $c \in \mathrm{Fil}^{\min(o,h)}$. If both o and h are positive, $c = 0$, because a nonzero constant is a unit of B_{dR}^+. If $h \le 0$, every constant lies in Fil^h and $y \in \mathrm{Fil}^h$. If $o \le 0 < h$, the congruence only implies $y \in \mathrm{Fil}^0$. These are exactly the displayed cases. □

The source writes this as `mem_withOrders_improvedOrders`. In any use one can keep the stronger original order if the displayed imp happens to be smaller, because both order systems hold. The ability to turn a small positive order into an arbitrarily large depth bound is the decisive simplification in the long-gap argument.

Theorem 51.13 (A rank-free depth operator)**.** *Let $a \ge 0$, $n \in \mathbb{Z}$ and $c = c(a,n)$. There is an additive operator on $\mathcal{N}(T)$ of the shape*

$$B_{a,n}(x) = p^c x + \sum_j r_0\big(v_j \Delta^{-1}(g_j x - x)\big), \qquad v_j \in A, quad g_j \in G',$$

with the following properties.

(i) For any form λ, if $\mathcal{V}_a(\lambda, gx - x) \in \mathrm{Fil}^n$ for all $g \in G'$, the same assertion with $B_{a,n}x$ holds in Fil^{n+1}.

(ii) If x has depth o at letter i, then $B_{a,n}x - p^c x$ has orders o. Every depth condition already satisfied by x is still satisfied by $B_{a,n}x$.

(iii) If x maps to a fixed constant in a trivial quotient lattice and the retraction commutes with that quotient, then $B_{a,n}x$ maps to the constant multiplied by p^c.

Proof. For $n \neq 0$, choose g, ω_a, X, V as in Proposition 51.10. Let π stand for the de Rham image of E_a under φ^{-a}; it generates the maximal ideal. After subtracting the fixed scalar from the value of x, write $y = \mathcal{V}_a(\lambda, x) - s \in \mathrm{Fil}^n$. At this point y belongs to the value field from (d). Thus $\pi^{-n}y$ is integral in that field and g fixes it modulo Fil^1, since g acts trivially on S modulo E_a. The equation $g(\pi) = \widetilde{\omega}_a \pi$ (with a tilde denoting evaluation after inverse Frobenius) gives

$$g(y) - \widetilde{\omega}_a^n y \in \mathrm{Fil}^{n+1}.$$

This is also valid for negative n: multiplication by $g(\pi)^n$ shifts the first filtration step by n. Set $w = \Delta^{-1}(gx - x)$. Since $\widetilde{\Delta}$ is a unit in B_{dR}^+, the equation $\Delta X = \omega_a^n - 1$ implies

$$\mathcal{V}_a(\lambda, w) - \widetilde{X} y \in \mathrm{Fil}^{n+1}.$$

Use $B_{a,n} = p^c \mathrm{id} - r_0(V\Delta^{-1}(g-1))$. Before retraction its value differs from $p^c s$ by

$$p^c y - \widetilde{V}\mathcal{V}_a(\lambda, w) = -\widetilde{V}(\mathcal{V}_a(\lambda, w) - \widetilde{X} y) - (\widetilde{V}\widetilde{X} - p^c)y,$$

which belongs to Fil^{n+1} by (51.1). The retraction does not change this congruence by (b). This proves (i) via the equivalence between depth and congruence to a constant.

For $n = 0$, use the $m = p^{c(a,0)}$ elements g_j of Lemma 51.11, choosing $v_j = \Delta$ for each j. Before retraction the operator is

$$p^c x + \sum_{j=1}^{m} (g_j x - x) = \sum_{j=1}^{m} g_j x.$$

The value of x is integral, since it is congruent to a fixed scalar modulo Fil^0. By the value-field comparison in (d) and the scalar averaging lemma, the new value is invariant modulo Fil^1. It is therefore congruent to a fixed scalar, and the retraction preserves that congruence.

For (ii), each difference $g_j x - x$ has orders o. Division by Δ and retraction preserve these orders by (b),(c), as does multiplication by the integral scalar v_j. Hence the operator differs from $p^c x$ by an element of $\mathcal{M}_i^T(o)$. Since that submodule is G'-stable, the operator preserves the depth condition. Finally the image of each difference $g_j x - x$ in the trivial quotient is zero when x lies over a fixed constant. Retraction commutes with the projection, giving (iii). □

The theorem is the combination of `exists_valueAt_sub_mem_hodgeFil_succ` and `exists_isOperatorOfCost_mem_hodgeFil_succ`. It proves the one-step moving assertion in Chapter 49, conditional only on the stated chart and form comparisons. In particular the same operator works for every form and vector; one does not construct a separate operator for each cocycle using its individual coefficients. This uniformity is indispensable when the sum of low jump multiplicities is much smaller than the rank of the ambient extension.

51.7 From a single operator to the cost of a window

The notation of a side is as follows. Given a crystalline lattice Y, a trivial parameter lattice C, and a family of crystalline cocycles $\alpha_c : G' \to Y$ linear in $c \in C$, the extension lattice $\mathcal{E} = Y \oplus C$ has action

$$g(y, c) = (gy + \alpha_c(g), c).$$

Write $q : \mathcal{E} \to C$ for the equivariant projection, and c_i for the constant vector $1 \otimes c$ projected to letter i in the corresponding ambient trivial lattice. The set of points over $p^r c_i$ with orders o and depth d consists of the elements $x \in \mathcal{N}(\mathcal{E})$ at letter i such that

$$qx = p^r c_i, \qquad x \in \mathcal{M}_i^{\mathcal{E}}(o) \cap \mathcal{I}_i^{\mathcal{E}}(d).$$

We call a scalar $p^r E^s$ a cokernel killer at i if it carries the letter component of the trivial integral lattice into the image of the standard order module for $\mathcal{E}$ at that letter. The standard system at level zero is $\sigma_0(a, u) = -u$ for all a. As in Section 50.7, existence of the required scalar kills is an assertion about the image of a module, not a pointwise choice of lifts depending on c with no uniformity. Applying the coefficient retraction gives actual localized Kisin points, so multiplication by E^{-s} gives a point over $p^r c_i$ with orders and depth

$$\rho_{-s}(a, u) = -u - s\mathbf{1}_{a=0}.$$

Indeed E^{-s} affects the order only at $a = 0$, and each order module is G'-stable. This proves the point-existence implication of the cokernel definition (`pointsOver_nonempty_of_kills Cokernel`) from the coefficient-preserving form of that definition.

Proposition 51.14 (A common operator for finitely many steps)**.** *Let $J \subset \mathbb{N} \times \mathbb{Z}$ be finite and write $C(J) = \sum_{(a,n)\in J} c(a, n)$. For every side there is an additive operator B_J on its localized Kisin module such that, if x lies over a fixed constant and is of depth d at i, then*

$$B_J x - p^{C(J)} x \in \mathcal{M}_i^{\mathcal{E}}(d) \cap \ker q.$$

If J contains all (a, n) for which $d(a, u) \le n < d'(a, u)$ for some $u \in U_{i[-a]}$, then $B_J x$ has depth d'.

Proof. Order the pairs in J by nondecreasing second coordinate n, with any order among pairs with equal n. Compose the operators of Theorem 51.13 in that order. Each step preserves existing depth and the condition of being over a constant. A correction of cost c differs from multiplication by p^c by a vector of the old orders and zero projection. If B_1 and B_2 are two such operators, their correction relative to multiplication by $p^{c_1+c_2}$ can be written as

$$(B_2 B_1 x - p^{c_2} B_1 x) + p^{c_2}(B_1 x - p^{c_1} x),$$

so it has the old orders and zero projection. This proves the first assertion by induction.

Fix a, u and a form of type u. If $d'(a, u) > d(a, u)$, each integer step from the latter to the former occurs in the prescribed ordering. By property (i) of Theorem 51.13, traversing a step n upgrades its depth from at least n to at least $n + 1$. All intervening steps at other divisors or for other jumps preserve the depth. Thus after the last step the required depth holds for this form. This is true for all forms and all jumps, including those at which no step is required. □

This proves `exists_isOperatorOfCost_and_mem_deepElements`. In particular for a point over $p^r c_i$ with orders o and depth d, the operator provides a point over $p^{r+C(J)} c_i$ of the same orders and improved depth d'. To verify that the old orders remain, use the first assertion of

Proposition 51.14 with initial depth o; membership of x in $\mathcal{M}_i(o)$ already implies this depth by stability. Then Lemma 51.12 improves the orders to $\mathrm{imp}(o, d')$. Multiplication by further powers of p does not change any filtration order. This proves `pointsOver_nonempty_of_costs`. If many different jump intervals overlap, one pays only for their union J; no rank factor or multiplicity count is inserted here.

Here is the elementary uniform bound on the sum of costs of two integer windows. Define

$$J(n, b) = [-n, n] \cap \mathbb{Z} \ \cup \ [b - n, b + n] \cap \mathbb{Z}.$$

Define $v_p(0) = 0$ in sums and let $H \geq 1$ bound the absolute values of members of $J(n, b)$. The number of nonzero integers divisible by p^j in an interval of $2n + 1$ consecutive integers is at most $\lfloor 2n/p^j \rfloor + 1$. Counting the contribution of each p^j gives

$$\sum_{m \in J(n,b)} v_p(m) \leq \frac{4n}{p - 1} + 2\lfloor \log_p H \rfloor. \tag{51.3}$$

The use of the floor is justified because terms for $j > \log_p H$ vanish. The inequality remains valid if the windows overlap, by counting the union only once on the left and bounding by both individual sums. Consequently

$$\sum_{m \in J(n,b)} c(a, m) \leq (4n + 2)(d_0 + 3a) + \frac{4n}{p - 1} + 2\lfloor \log_p H \rfloor.$$

The readable source uses the variant `sum_twoWindows_le` obtained from the same count (in `SidePoints.lean`). For $p \nmid m$, the first term of the cost is zero; thus on the no-multiple windows of the few-low-jumps argument the bound simplifies to $c(a, m) \leq 3a$ for an unramified chart. The exceptional degree $m = 0$ obeys the same bound, by the averaging rather than the lifting-the-exponent construction of Section 51.5. It is not legitimate to apply the latter construction to $m = 0$ by writing $v_p(0) = 0$.

51.8 The monic Taylor coefficient at a long gap

A gap in the jumps is used in a different way from a no-multiple window. Fix an unramified chart $E = z + p$ and let i be a letter. For $l \geq 0$ define the standard order system at level l,

$$\sigma_l(a, u) = \begin{cases} 0 & a < l, \\ -u & a \geq l. \end{cases}$$

Let $M_l^T(i)$ denote the localized Kisin part of $\mathcal{M}_i^T(\sigma_l)$; it is an S-module. For $A \geq 0$, call $x \in M_l^T(i)$ a constant modulo E^A if it also belongs to

$$\mathcal{I}_i^T(\sigma_l + A\mathbf{1}_{a=0}).$$

Call this $\mathbb{Z}_p$-submodule $D_l^T(i, A)$. This is not the ordinary constant term of a formal power series of lattice vectors: the definition refers to near-invariance under G'. It is a useful auxiliary space because of the improvement of positive orders in Lemma 51.12.

For $l > 0$ and a function o of the jumps at i, let $\sigma_l[o]$ be the system $\sigma_l(a, u) + o(u)\mathbf{1}_{a=0}$. The following coefficient extraction lemma will be applied twice: once to an extension and once to its trivial quotient.

Lemma 51.15 (Taylor coefficient and a jump gap)**.** *Let $l > 0$. Suppose $\xi_0, \dots, \xi_N \in D_l^T(i, A)$ and $y \in M_l^T(i)$, and*

$$\sum_{n=0}^{N} E^n \xi_n + E^{N+1} y \in \mathcal{M}_i^T(\sigma_l[o]), \qquad o(u) \le A.$$

Then $\xi_N \in \mathcal{M}_i^T(\sigma_l[o_N])$, where

$$o_N(u) = \begin{cases} A & N < o(u), \\ 0 & N \ge o(u). \end{cases}$$

Proof. First consider $N = 0$. Since Ey has order at least one at $a = 0$ and the other standard orders, ξ_0 has order at least $\min(1, o)$ at $a = 0$. Here $\sigma_l(0, u) = 0$ because $l > 0$. If $o(u) > 0$ the order is positive and the depth A upgrades it to A, by Lemma 51.12; otherwise ξ_0 already has order zero there. This gives o_0.

Now write the sum as $\xi_0 + Ey_1$, with $y_1 = \sum_{n=0}^{N-1} E^n \xi_{n+1} + E^N y$ in $M_l^T(i)$. By the first case ξ_0 has order A at all jumps where $o > 0$; at the remaining jumps order zero suffices for $o \le 0$. Thus it belongs to $\mathcal{M}_i^T(\sigma_l[o])$ because $o \le A$. Subtract it and divide by E to get $y_1 \in \mathcal{M}_i^T(\sigma_l[o - 1])$. The inductive hypothesis gives the result for ξ_N, since $o(u) - 1 > N - 1$ exactly when $o(u) > N$. □

This is the proof of `mem_withOrders_of_add_E_smul_mem` and `mem_withOrders_of_expansion_mem`. It relies on $l > 0$; if $l = 0$ the standard orders at E are not uniformly zero and the same positive-order improvement cannot be made in this form.

We next produce the required expansions. The integral radius estimate (Section 50.5) says that for $3 \le p^l$ and $A \le p^l \delta$ one has

$$p^{4\delta} M_l^T(i) \subset \operatorname{span}_S D_l^T(i, A). \tag{51.4}$$

Precisely this follows from `frame_theorem.pow_smul_mem_span_of_mem_valueFieldPoints` after projecting to a letter. It uses the frame identification with the value-field points and the first part of the radius lemma, which still rely on the integral comparison; the elementary matrix truncation yielding the fourth power was proved in Section 50.5. We make (51.4) an explicit hypothesis for the long gap, so no assertion about arbitrary vectors of arbitrary period modules is implicit here.

Since the chart is unramified, $E = z + p$ and $S/(E) \simeq \Lambda$, each $s \in S$ can be expanded modulo every E^M in a sum with coefficients in Λ. Repeatedly choose the residue in Λ and divide the remainder by E to get

$$s = \sum_{n<M} w_n E^n + E^M s_M, \qquad w_n \in \Lambda,\ s_M \in S.$$

The space $D_l^T(i, A)$ is stable under Λ, since G' fixes the constants and multiplication by $w \in \Lambda$ preserves the orders. From (51.4) write $p^{4\delta} x = \sum_j s_j b_j$ with finitely many $b_j \in D_l^T(i, A)$. Expand the s_j and collect the coefficients:

$$p^{4\delta} x = \sum_{n<M} E^n \xi_n + E^M y_M, qquad \xi_n \in D_l^T(i, A), \quad y_M \in M_l^T(i). \tag{51.5}$$

The coefficients $\xi_n = \sum_j w_{jn} b_j$ are chosen once, independent of M. This is `exists_taylor_expansion_of_mem_stdOrdersPart`. It requires no convergence assertion about an infinite series in B_{dR}; we only extract a finite coefficient of this family of identities.

Theorem 51.16 (Long-gap lifting lemma)**.** *Let S be a side with extension $\mathcal{E}$ and parameter lattice C. Assume (51.4) for its extension and trivial quotient, and the chart comparison assertions of Section 51.6. Let i be a letter and suppose*

(i) E^s *kills the cokernel at i, and p^κ kills the cokernel at $i[-l]$;*

(ii) $l > 0$, $3 \le p^l$ *and* $A \le p^l\delta$*;*

(iii) all jumps of $\mathcal{E}$ at $i[-a]$, $0 \le a < l$, are at most g;

(iv) $N = s + g(1 + p + \cdots + p^{l-1}) < A$.

Then for each $c \in C$ there is a localized Kisin point over $p^{\kappa+4\delta}c_i$, with orders

$$\sigma_l(a,u) + A\mathbf{1}_{a=0,\,u<-N}$$

and depth $\sigma_l(a,u) + A\mathbf{1}_{a=0}$.

Proof. The cokernel at i gives a point y over c_i with orders $\rho_{-s}(a,u) = -u - s\mathbf{1}_{a=0}$. In the unramified chart

$$E_a = z^{p^a} + p = (E-p)^{p^a} + p.$$

Hence the polynomial

$$f(X) = X^s \prod_{a<l} \left((X-p)^{p^a} + p\right)^g \tag{51.6}$$

is monic in $\mathbb{Z}[X]$ of degree N. Multiplying y by $f(E)$ first cancels the pole of order s at E and then adds g at each of the divisors E_a with $a < l$. Thus $v = f(E)y$ has orders

$$-u + g\mathbf{1}_{a<l}.$$

Since the jumps at these places are at most g, $v \in M_l^{\mathcal{E}}(i)$, and at $a = 0$ one also has order at least $\min(A, g-u)$.

Apply (51.5) with $M = N+1$ to this v. Lemma 51.15 with $o(u) = \min(A, g-u)$ shows that the coefficient ξ_N has orders at least σ_l shifted by A at every jump $u < -N$, because both $A > N$ and $g - u > N$. It has uniform depth $\sigma_l + A\mathbf{1}_{a=0}$ by membership in $D_l^{\mathcal{E}}(i, A)$.

Apply the trivial-quotient map q to the Taylor identity. By the functoriality of order modules and constants modulo E^A, all the coefficient images satisfy the corresponding conditions in the trivial lattice. The projected sum is $p^{4\delta}f(E)c_i$. Subtract the coefficients of that polynomial, putting

$$\eta_n = q\xi_n - p^{4\delta}f_n c_i, qquad f(X) = \sum f_n X^n.$$

These η_n are constants modulo E^A in the trivial lattice (the projected c_i is genuinely fixed). Their sum $\sum_{n=0}^{N} E^n\eta_n + E^{N+1}qy_{N+1}$ is zero, hence of arbitrarily high orders. Applying Lemma 51.15 with $o(0) = A > N$ gives

$$w = q\xi_N - p^{4\delta}c_i \in \mathcal{M}_i^C(\sigma_l + A\mathbf{1}_{a=0}).$$

Monicity of f is used exactly in obtaining coefficient $f_N = 1$. Merely bounding the degree of f without ensuring its leading coefficient is a unit would not give the required projection.

The error $E^{-A}w$ lies in $M_l^C(i)$. By the Frobenius transport of the standard order module, p^κ killing the cokernel at $i[-l]$ implies that $p^\kappa E^{-A}w$ lifts at i to $v_0 \in M_l^{\mathcal{E}}(i)$. This is just transport of the killer at level zero to level l under φ^l, and the coefficient retraction from integral solutions to localized Kisin points. Then $p^\kappa\xi_N - E^A v_0$ lies over $p^{\kappa+4\delta}c_i$. The correction term has orders $\sigma_l + A\mathbf{1}_{a=0}$ at all jumps, and by stability is deep for the same system. It cannot reduce the orders on the negative jumps selected by the coefficient and cannot reduce its depth. This is the required point. □

This supplies the long-gap lifting theorem `pointsOver_nonempty_of_long_gap` in the compact and readable `LongGap.lean`. The proof exhibits a rather striking mechanism: a monic product of Frobenius translates of one distinguished scalar has a specified degree, and a Taylor coefficient at that degree simultaneously has a known projection to the trivial lattice and a very large order on sufficiently negative jumps. The large order does not have to be purchased one step at a time. The cost is $p^{\kappa+4\delta}$, where δ need only be about A/p^l. In particular the geometric factor in the upper bound $N = s + g(1 + p + \cdots + p^{l-1})$ is exactly the polynomial degree, not an estimate on the rank of the representation.

51.9 The triple pairing and the safety rule

Let T, T' be two sides with kernels Y, Y', parameter lattices C, C', and cocycles $\alpha_c, \beta_{c'}$. Let $B_0 : Y \times Y' \to \mathbb{Z}/p^m(1)$ be the bilinear congruence pairing, equivariant modulo p^m. It comes from an integral form whose failure of equivariance is divisible by p^m. The third side is the extension of the trivial lattice by the Tate dual of $Y \otimes Y'$ associated to that failure of equivariance (and all three sides have their specified integral crystalline data). Its jump set contains 0 and the integers $-u - u' - 1$ with $u \in U_i(Y)$ and $u' \in U_i(Y')$. The -1 is the weight of the Tate twist in the jump convention of these notes. Let Z denote the kernel of the third side, and n a point in its extension over a fixed constant. This side comes with the equivariant extension form

$$\mathrm{ev} : \mathcal{E}_Z \times (Y \otimes Y') \longrightarrow \mathcal{O}(1),$$

whose reduction modulo p^m, followed by the fixed coefficient map to $\mathbb{Z}_p$, has the property that evaluation at a point n over p^κ times the constant 1_i is $p^\kappa B_0$ on the letter i. It is worth verifying that the algebraic extension implementing this form is natural. Write F for an integral bilinear form lifting the congruence pairing (before applying the coefficient map) and write $D_g = (gF - F)/p^m$ for its equivariance defect in the dual. Then D is a cocycle, because

$$D_{gh} = D_g + gD_h.$$

The dual extension has action $(f, c) \mapsto (gf + cD_g, c)$; the functional

$$(f, c) \longmapsto cF - p^m f$$

is equivariant and reduces to cF modulo p^m. Changing the sign convention for D changes the sign of the correction. This verifies the algebraic model of the third side; crystallinity of that extension follows from the specified defect cocycle, which must itself satisfy the finite condition. The source uses precisely this model in `PairingSide.tateDualSide`, `extensionForm` and related declarations in `CupProductBound.lean` and `SidePoints.lean`.

The integral property of pairings that is required is the following. Suppose $b : T_1 \otimes T_2 \to T_3$ is an equivariant bilinear map of lattice data. At a given inverse Frobenius place, a form of jump v on T_3 pulls back into sums of products of forms of jumps u, u' on the two factors with $u + u' \le v$, with coefficients in the positive de Rham ring. Consequently if x_1, x_2 have orders o_1, o_2 at a letter, their pairing has order system o_3 whenever

$$o_3(a, v) \le o_1(a, u) + o_2(a, u') \quad (u + u' \le v) \tag{51.7}$$

at all the corresponding jumps. For this assertion one chooses a filtration-adapted basis of dual forms over the fixed field and expands a form of the stated type in the dual tensor basis. The filtration on the tensor product is the sum filtration, so only the indicated $u + u' \le v$ can contribute; scalar coefficients given by filtered linearity have no negative filtration orders. Evaluation preserves the sums and products, proving the inequality term by term from (51.2).

Membership of the result in the ambient localized Kisin span under b uses the tensor-compatible identification of the integral frames. This is the use of `wittPairing_mem_kisinAinf` and `wittPairing_mem_withOrders` in `HodgeExpansion.lean`; we take that tensor-compatible comparison as an input and use the elementary filtered calculation above.

For the Tate line $\mathcal{O}(1)$ itself, an element of standard orders $\sigma_0(a,-1)=1$ at all places lies, after multiplication by τ, in the A-span of the integral Tate lattice. This is the rank-one case of the frame comparison, `tau_pow_smul_mem_ainfLattice_of_eq_chi_pow_smul`. Its consequence in the mod-p^m target is the smallness property: multiplying once more by z, whose reduction modulo p has strictly positive valuation, makes all Witt coordinates small. Write

$$W_m(C_p^\flat)=W(C_p^\flat)/p^m, qquad \mathcal{K}_m=\operatorname{im}\left(W(\mathfrak{m}_{C_p^\flat})\to W_m(C_p^\flat)\right).$$

Here $W(\mathfrak{m}_{C_p^\flat})$ means the kernel of reduction of integral Witt vectors to the residue Witt ring, an additive subgroup. We use $\mathcal{K}_m(1)$ for the same subgroup in the cyclotomic twist. It is invariant under Galois and Frobenius. Multiplying a vector with integral Witt coordinates by an element of $W(\mathfrak{m}_{C_p^\flat})$ gives a small vector in this sense.

Theorem 51.17 (Three orders imply a safe pair)**.** *Let w,y lie over zero in the extensions of the first and second sides, at the same letter i, with orders o,o' respectively. Identify them in the kernel lattices. Let n be a point of the third side over $p^\kappa 1_{i[-l]}$, with orders v_0 antitone in its second argument. Put*

$$\widetilde{v}_0(a,u)=\begin{cases}0 & a<l,\\ v_0(a-l,u) & a\ge l.\end{cases}$$

Suppose that for all a and all jumps u,u' of the first and second sides at $i[-a]$,

$$\widetilde{v}_0(a,-u-u'-1)+o(a,u)+o'(a,u')>0. \tag{51.8}$$

Then

$$p^\kappa B_0(\Delta w,y)\in\mathcal{K}_m(1),\qquad p^\kappa B_0(w,\Delta y)\in\mathcal{K}_m(1).$$

Proof. Frobenius transports n to $\varphi^l n$ at letter i, with the indicated shifted orders. (The Frobenius shift of orders and letter parts is one of the frame identities of the chart.) Define $\zeta=w\otimes y$ in the tensor kernel. If $s\ge u+u'$, then $-s-1\le -u-u'-1$ and antitonicity gives

$$\widetilde{v}_0(a,-s-1)\ge\widetilde{v}_0(a,-u-u'-1).$$

The strict integer inequality (51.8) and (51.7) imply that ζ has orders at least $1-\widetilde{v}_0(a,-s-1)$ at the jump s. Pair ζ with $\varphi^l n$ under the extension form. For jumps s of the tensor and t of the dual with $s+t\le -1$, antitonicity gives $\widetilde{v}_0(a,t)\ge\widetilde{v}_0(a,-s-1)$, so the sum of the two order functions is at least 1. The Tate-line value therefore has standard orders at every place. Multiplication by τ makes it integral by the rank-one comparison, then multiplication by z makes its mod-p^m image small. By the reduction property of the extension form, that image is $z\tau p^\kappa B_0(w,y)$, as desired. Bilinearity permits either factor to absorb $\Delta=z\tau$. □

This is the proof of `isSafe_of_orders` in readable `SafeMoves.lean`. In applications w is a difference of points with the same trivial coordinate. Such differences lie in the kernel and have at least the pointwise minimum of their two order systems. To move the first component by w while a point y rests in the second extension, it suffices that for every $g\in G'$, $p^\kappa B_0(w,(gy-y)_1)$ be small, where $(\,)_1$ denotes the kernel coordinate. If the rest point has depth d', then $\Delta^{-1}(gy-y)$ has orders d' by the integral comparison of Section 51.6; it lies over zero. We apply Theorem 51.17 with $o'=d'$. For movement of the second component

while x rests, the required expression is $p^{\kappa}B_0((gx-x)_1, gw)$. The difference of x gives its depth and gw has the same orders as w by stability.

When the third side has the cokernel killer $p^{\kappa}E^t$ at $i[-l]$, we may choose $v_0 = \rho_{-t}$. In that case the exact numerical rule sufficient for a move is

$$R(l,t;a,s) = \begin{cases} 1 & a < l, \\ t-s & a = l, \\ -s & a > l. \end{cases} \tag{51.9}$$

The sum of the moving order and resting depth at jumps u, u' must be at least $R(l,t;a,u+u')$. Indeed the shifted third order at $-u-u'-1$ is zero below l, is $u+u'+1-t$ at l, and is $u+u'+1$ above it. In each case its sum with the right side of (51.9) is exactly 1. This proves the rule in `isSafeMoveFst_of_killsCokernel` and `isSafeMoveSnd_of_killsCokernel`. Notice the change in condition below l: the sum need only be positive there, not equal to a large weight-dependent expression. This is the reason for carrying previously chosen points forward along Frobenius before making a move. It cannot be seen from a pointwise estimate at a single de Rham place.

51.10 From safe pairs to an actual cup coboundary

One must justify that safety in the small Witt subgroup has a cohomological consequence over the original finite Galois module. We prove the pairing lemma itself in an elementary form; no rational duality alone supplies it.

First, work with $\mathcal{H} = W_m(C_p^{\flat})(1)$ and its subgroup $\mathcal{K}_m(1)$. Frobenius φ acts on Witt coordinates and is trivial on the finite Tate factor. It commutes with the Galois action.

Lemma 51.18 (Artin–Schreier–Witt modulo the small subgroup)**.** *The map $\varphi - 1 : \mathcal{H} \to \mathcal{H}$ is surjective. An element $x \in \mathcal{H}$ with $(\varphi-1)x \in \mathcal{K}_m(1)$ is uniquely congruent modulo $\mathcal{K}_m(1)$ to a constant in $\mathbb{Z}/p^m(1)$.*

Proof. Fix a generator of the Tate twist to work with ordinary Witt coordinates. In length one, $\varphi - 1$ is $x \mapsto x^p - x$, surjective because $C_p^{\flat}$ is algebraically closed. Inductively, solve for the first coordinate of a desired image, subtract its Teichmüller lift's image and divide the remainder by p to reduce by one length. The division here is carried out in the Witt ring over the perfect field; its ambiguity modulo p^m is exactly removed by reducing the quotient modulo p^{m-1}.

For the assertion modulo small, first consider length one. If $x^p - x$ is of strictly positive valuation, then x is integral: if $v(x) < 0$ its valuation would be $pv(x) < 0$. The residue of an integral such x satisfies $\bar{x}^p = \bar{x}$, hence lies in $\mathbb{F}_p$. Subtract a Teichmüller constant lifting this residue. The first coordinate is now small. Subtract its Teichmüller lift (which is itself small) and divide by p; the result of applying $\varphi - 1$ remains small modulo the shorter length, because the kernel of the residue map is saturated for p in this sense. This is immediate from the sequence of Witt coordinates in the perfect residue field: the vanishing of all coordinates of the residue after multiplication by p implies the vanishing of the first $m-1$ coordinates before multiplication. Induction then makes the entire vector congruent to a sum of Teichmüller constants times powers of p, i.e. an element of $\mathbb{Z}/p^m$. Uniqueness follows because reduction of a nonzero constant mod p^m in the residue Witt ring is nonzero. □

In addition, for every fixed $x \in \mathcal{H}$, $gx - x$ is small for g sufficiently near 1 in the Galois topology. Indeed the Galois action on the finite number of Witt coordinates is continuous

for the valuation topology on $C_p^\flat$, and addition and subtraction of Witt vectors are given by polynomials in the coordinates. At the identity all the coordinates of the difference vanish, so they lie in the maximal ideal for a neighborhood of the identity. The cyclotomic action is trivial modulo p^m sufficiently near 1. The same argument applies to fixed vectors in a tensor product with a finite free integral representation, after reducing it modulo p^m. This proves the near-identity smallness used in the source; it does not assert that the Galois action on all of $W_m(C_p^\flat)$ is discrete.

Theorem 51.19 (The pairing lemma via a four-move path)**.** *Let x, w be points of the first extension over the global (all-letter) constant $p^{e_1}c$, and x', w' points of the second over $p^{e_2}c'$. Let $e_3 \geq 0$. Assume that the four moves from $(\varphi x, \varphi x')$ through $(w, \varphi x')$, (w, w') and (x, w') to (x, x') are safe at cost p^{e_3}, and that for every $h \in G'$ the move of the second component by $hx' - x'$ while x rests is safe. Then, for every $N \geq e_1 + e_2 + e_3$, $p^N[\alpha_c \smile_{B_0} \beta_{c'}] = 0$ in continuous cohomology of the original local Galois group with coefficients in the finite Tate module.*

Proof. Write u, u_0, v, v_0 for the kernel coordinates of x, w, x', w' respectively, and put $U = \varphi(u)$, $V = \varphi(v)$. The cocycles

$$a_g = p^{e_1}\alpha_c(g), \qquad b_g = p^{e_2}\beta_{c'}(g)$$

give the first coordinates of the extension actions. In particular set

$$A_g = gu - u + a_g, qquad B_g = gv - v + b_g,$$

which are the first coordinates of $gx - x$ and $gx' - x'$. They are themselves cocycles. Define the bilinear scalar-valued pairing $\langle r, s\rangle = p^{e_3}B_0(r, s)$ extended to the Witt modules; it is Galois equivariant and Frobenius compatible. The rest of the proof uses only these identities and the fact that the target is $\mathcal{H}$.

By Lemma 51.18, choose $y \in \mathcal{H}$ satisfying

$$(\varphi - 1)y = \langle u_0, V - v_0\rangle + \langle u, v_0 - v\rangle. \tag{51.10}$$

Define a function (for now only algebraically) by

$$f(g) = \langle a_g, gv\rangle - \langle u, B_g\rangle + gy - y. \tag{51.11}$$

This is exactly the source's cochain after simplifying the four terms using equivariance and the extension action formula. We verify its two properties.

First, since φ commutes with g and is trivial on the integral cocycles, the difference of applying Frobenius to (51.11) is

$$\begin{aligned}(\varphi - 1)f(g) = {}& \langle gu_0 - u_0 + a_g,\ g(V - v_0)\rangle \\ & - \langle U - u_0,\ gV - V + b_g\rangle \\ & - \langle u_0 - u,\ gv_0 - v_0 + b_g\rangle \\ & + \langle gu - u + a_g,\ g(v_0 - v)\rangle.\end{aligned}$$

For completeness, insert (51.10) in $(g - 1)(\varphi - 1)y$. The remaining terms in $(\varphi - 1)f(g)$ are

$$\langle a_g, g(V - v)\rangle - \langle U, gV - V + b_g\rangle + \langle u, gv - v + b_g\rangle.$$

Adding the two terms of $(g - 1)(\varphi - 1)y$ and collecting by bilinearity yields the displayed four-move identity. Each summand is small by the assumed safety of the corresponding

move, so $(\varphi-1)f(g)$ is small. Lemma 51.18 gives a unique constant $k(g)\in \mathbb{Z}/p^m(1)$ with $f(g)-k(g)\in \mathcal{K}_m(1)$.

Second, use the inhomogeneous coboundary convention $\partial f(g,h)=gf(h)-f(gh)+f(g)$. The cocycle relations for a_g and B_g give directly

$$\partial\langle a_g, gv\rangle(g,h)=-\langle a_g, g(hv-v)\rangle,$$
$$\partial(-\langle u, B_g\rangle)(g,h)=-\langle gu-u, gB_h\rangle.$$

The last term of (51.11) is a coboundary and has zero differential. Using $hv-v=B_h-b_h$, we obtain the exact identity

$$\partial f(g,h)=\langle a_g, gb_h\rangle-\langle A_g, gB_h\rangle.$$

The last pairing is small by the assumption on moves by $hx'-x'$ while x rests. Replacing f by k changes ∂f by a small element since that subgroup is G'-invariant. Both ∂k and $\langle a_g, gb_h\rangle$ are constants, so their small difference is zero by uniqueness of constants modulo the small subgroup. Thus on G'

$$\partial k(g,h)=p^{e_1+e_2+e_3}B_0(\alpha_c(g), g\beta_{c'}(h)).$$

We still need continuity of k. Near the identity, a_g and b_g vanish modulo p^m by continuity of the integral cocycles. Also $gv-v$ and $gy-y$ are small, by the preceding near-identity argument. The fixed pairing with u of the difference $gv-v$ is small near the identity even if u has nonintegral coordinates: finitely many coordinates of a continuous bilinear expression vanish at the identity. Consequently (51.11) is small near the identity and $k(g)=0$ there. For h near 1, $\beta_{c'}(h)$ vanishes modulo p^m, so the equality for ∂k gives $k(gh)=k(g)$ for all g. Hence k is locally constant. It supplies a continuous coboundary on G'.

Finally the open subgroup G' has finite index prime to p. Restriction followed by corestriction is multiplication by its index on H^2. The class under consideration is p^m-torsion, so vanishing after restriction implies vanishing on the full local group: the index is invertible on any p^m-torsion group. Further multiplication by p proves the assertion for all N in the stated range. □

The original formal conclusion is also expressed using the embedding of the finite Tate module as roots of unity into the multiplicative group of the algebraic closure; the finite-module coboundary just obtained gives that conclusion as well. This is the content of `IsScaledCupCoboundary.of_isSafePath`, `phi_smul_sub_mem_smallPart`, `nsmul_modPairing_eq_coboundary`, and `isCupCoboundary_of_eq_coboundary`. The proof shows why two different kinds of safe movement are needed: the four Frobenius moves ensure that the adjusted cochain takes constant values up to small elements, whereas the cocycle motions ensure that its coboundary is the desired cup. Merely showing that the pairing of two Hodge filtrations is small would not directly establish either assertion.

51.11 The walk along the Frobenius permutation

We give the exact order systems for terminal and intermediate points. A cut is a distinguished letter. Let l_i be the least strictly positive integer such that $i[-l_i]$ is a cut, and require that each Frobenius orbit contain a cut. A given cut has an integral shift $t_i\ge 0$ such that the third side has cokernel killer $p^{\kappa}E^{t_i}$ at it. Fix a margin $\eta\ge 1$ and define

$$D(t,\eta,l;a,u)=-u+t\mathbf{1}_{a=0}+\eta\mathbf{1}_{a<l},$$
$$S(\eta,l;a,u)=\begin{cases}\max(0,-u-\eta) & a<l,\\ -u & a\ge l.\end{cases}$$

A standard point at a cut i has orders $\rho_{-\eta}$ and depth $D(t_i, \eta, l_i)$; a stop has orders and depth at least $S(\eta, l_i)$. These are the definitions `standardDepth` and `stopOrders`, and the latter is deliberately the maximum of two order systems. A transported point $\varphi^l x$ has orders shifted by l (zero at the new places $a < l$) and depths shifted by l with lower bound 1 instead of 0 at those new places. This uses the fact that φ^l of a crystalline frame has positive depth at the new divisors; see `phi_pow_smul_mem_pointsOver` in `SidePoints.lean`. We record this integral Frobenius transport property as an input to the walk, exactly in these strengths; translation of the old Hodge orders alone would give only order zero and would not guarantee the positive depth at the new divisors.

At a cut i, put $l = l_i$, $j = i[-l]$ and $l_0 = l_j$. Suppose standard points on the two sides over $p^{e_1}c_i, p^{e_2}c_i'$ have been chosen at all cuts. Write y, y' for the transported points from j to i, and x, x' for the new standard points at i. If a stop q exists on the first side allowing the second to move directly from y' to x', then the three nontrivial moves

$$(y, y') \longrightarrow (q, y') \longrightarrow (q, x') \longrightarrow (x, x')$$

are all safe. We show this by checking the first and last move numerically. The middle move is the stated stop hypothesis. The first movement $y - q$ has order at least the minimum of the shifted $\rho_{-\eta}$ and $S(\eta, l)$. Below l this is zero. The depth of y' there is at least one. Thus the sum is at least 1, meeting the rule from the killer at j. At $a = l$ the moving order is $-u - \eta$ and the resting depth is $-u' + t_j + \eta$, since $l_0 > 0$. Their sum is $t_j - u - u'$, precisely the rule from that killer. At $a > l$ the moving order is $-u$ and the resting depth is at least $-u'$. This meets the remaining part of (51.9). For the final movement $q - x$, the order is at least $-u - \eta$ at $a = 0$, and the resting depth of x' is $-u' + t_i + \eta$ there. Their sum is $t_i - u - u'$, meeting the rule for the killer at i. At $a > 0$, the moving order is at least $-u$ except possibly when $0 < a < l$, where it is at least $-u - \eta$. The depth $D(t_i, \eta, l; a, u')$ is at least $-u' + \eta$ at those places and at least $-u'$ otherwise. Hence the remaining rules hold. If the stop is on the second side instead, switch which of the two intermediate moves is zero; the same inequalities apply. This proves `exists_isSafePath_of_stop`.

There is one additional local check for use in Theorem 51.19. If x, x' are standard at a cut and have been transported by φ^d to another letter, then the move of the second side by its cocycle $g\varphi^d x' - \varphi^d x'$ while the first rests is safe. Each of the moving order and the resting depth is bounded below by the shifted system $D(t_i, \eta, l_i)$ with value 1 at new places $a < d$. Their sum is at least 2 there. At $a = d$ it is at least $-u - u' + 2t_i + 2\eta \geq t_i - u - u'$. Above d it is at least $-u - u'$. Apply the rule from the third side at the originating cut. This proves `isSafeMoveSnd_cocycle`.

Theorem 51.20 (The global Frobenius walk)**.** *Suppose the standard points and third-side killers at each cut exist as above, and at each cut one side has a stop allowing the other side to move directly from its previously transported point to its new standard point. Then*

$$p^{e_1+e_2+\kappa}[\alpha_c \smile_{B_0} \beta_{c'}] = 0.$$

Proof. For each letter i, put $d_i = 0$ if it is a cut and $d_i = l_i$ otherwise. Choose the left point x_i at i to be the transport by φ^{d_i} of the standard point at $i[-d_i]$; define x_i' similarly. Each has the specified projection to $p^{e_1}c_i$ or $p^{e_2}c_i'$. Define y_i and y_i' by the strictly preceding cut (transport by φ^{l_i}). Because

$$l_i = 1 + d_{i[-1]},$$

one has $\varphi(\sum_i x_i) = \sum_i y_i$ and likewise on the other side: the equality follows term by term upon changing the letter index by inverse Frobenius. Off the cuts these transported points already equal the chosen points, so four moves of size zero are safe. At the cuts the previous

three-move calculation gives a four-move path with one zero move. Thus at each letter there is a path from y_i, y_i' to x_i, x_i' with intermediate points w_i, w_i' lying over the respective constants.

The extensions carry the complementary letter decompositions for the bilinear pairing: writing the second using the dual convention, a pairing between different indexed letter factors is zero. For clarity this follows from the balanced coefficient action. If the two idempotents are e_i, e_j, their cross pairing factors through $e_i e_j$, which is zero for $i \neq j$. Equivalently their distinct Teichmüller eigenvalues have unit difference. The Galois elements of G' preserve these letter factors, so the safety conditions for the sums of movements follow by expanding bilinearly: only pairings at the same letter remain and their sum is small. In particular the cocycle movements for the sum of the x_i's are safe by the preceding local check. The projections of the sums of the chosen and intermediate points are precisely the global constants $p^{e_1}c, p^{e_2}c'$. Apply Theorem 51.19 with these four global points. □

This is the proof of `isScaledCupCoboundary_of_stops`, including the minimal-cut condition: it is not sufficient that some cut occurs near each letter; the shift relation for the sums uses the nearest strictly preceding cut. When every letter is a cut, $l_i = 1$ and the same proof reduces to the simpler `isScaledCupCoboundary_of_stops_at_all_letters`.

51.12 Plain cuts and long-gap cuts

The cut theorem is useful only when both types of stops can actually be constructed with the same budget. We prove the two local constructions.

Call a cut with predecessor $l > 0$ steps back plain if among the l letters ending at that cut one side has all jumps $u \leq \eta$ and the opposite side has all jumps $u' \geq -\eta$. We call the former the down side. Suppose the down point x at the cut has not merely the standard depth but the depth

$$D(t_i, \eta, l; a, u) + \eta \mathbf{1}_{a=l}.$$

This is the source's `plainDepth` with equal margins at the two ends (the proof works with a different margin η_0 at the earlier cut, replacing the last η by η_0). On the down side, at $a < l$ the value has depth at least $\eta - u \geq 0$. Its original order is at least $-u - \eta$ and, if that order is positive, it already suffices for $S(\eta, l)$. Otherwise Lemma 51.12 improves the order to zero when the depth is nonnegative. For $a \geq l$ the original order $-u$ already suffices. Consequently x is a stop. Its depth exceeds the stop orders as well.

The up side's direct movement from the transported point at the earlier cut to the current point has order at least the minimum of shifted $\rho_{-\eta}$ and $\rho_{-\eta}$. At $a = 0$ the order is at least $\min(0, -u' - \eta) = -u' - \eta$, since the side is up, and the resting depth at x is $-u + t_i + \eta$. At $0 < a < l$, the moving order is at least $\min(0, -u') \geq -u' - \eta$ and the resting depth is $-u + \eta$. At $a = l$ the moving order is at least $-u' - \eta$ and the resting depth has the added η at that place. Above l the two bounds are $-u', -u$. Thus the rule for the third-side killer at i applies everywhere, proving that the up side moves safely while x rests. This is `mem_stopPoints_of_down` and `exists_mem_stopPoints_isSafeMoveSnd`, together with the version exchanging the first and second sides.

Now fix windows

$$U_i(T) \subset [-\eta, \eta] \cup \epsilon_i[\beta_i, \beta_i + V], \qquad U_i(T') \subset [-\eta, \eta] \cup (-\epsilon_i)[\beta_i, \beta_i + V]$$

with $\epsilon_i \in \{\pm 1\}$ and $\beta_i, V \geq 0$. An isolated far window with an opposing sign may prevent the cut from being plain. The long-gap construction is designed for this case. Suppose the chart

is unramified with $p \geq 5$, the preceding cut $j = i[-l]$ has shift zero, and the pivot letter is $k = i[-t]$ with

$$t + J + 1 = l, \qquad s_0 + \omega(1 + p + \cdots + p^J) < \beta_k.$$

Here $\omega \geq \eta$ bounds all the far window endpoints $\beta_{i[-a]} + V$ for $t < a < l$. Every nonpivot letter $i[-a]$, $a < l$, either has small far window ending at or below η or has sign opposite to the pivot. By choosing which side will rest, arrange that its far window at k is negative. At the other letters before i its jumps are at least $-\eta$. Let it be the side T. All its jumps at k and the J previous letters are at most ω, because the far window at k is negative. If p^κ kills its cokernel at j, E^{s_0} kills it at k, and

$$A = 2b \leq p^{J+1}\delta, qquad\beta_k < 2b,$$

then Theorem 51.16 supplies a point at k over $p^{\kappa+4\delta}c_k$ with order zero below level $J + 1$, except for far negative jumps at level zero where its order is $2b$. Its depth is $2b$ at level zero, zero from level one to level J, and the standard orders above. Transport by φ^t to i. It now has the indicated large orders and depth at level t; below t its order is zero and depth at least one. Use the depth operators for the steps $(a, 0)$, $t < a < l$, to deepen from zero to one there, preserving all existing orders and depths. This costs only the sum of the corresponding $3a$ (the chart is unramified).

The result is a point with depth

$$D_{\mathrm{gap}}(l, t, 2b; a, u) = \begin{cases} 2b & a = t, \\ 1 & a < l,\ a \neq t, \\ -u & a \geq l. \end{cases}$$

At levels below l other than the pivot the side is up, so $u \geq -\eta$, and order zero meets the stop-order requirement. At the pivot either the jump is low or is far negative with $-u \leq \beta_k + V \leq 2b$; the large orders obtained by Taylor extraction meet $\max(0, -u - \eta)$. Above l the standard order $-u$ suffices. The depth is at least the stop orders at all these places.

It remains to check safety during the other side's movement. The actual cut-order system used to construct standard points is slightly stronger than $\rho_{-\eta}$. It is

$$O(s, \eta, l) = \mathrm{imp}\left(\rho_{-s}, D(s, \eta, l + 1)\right). \tag{51.12}$$

At the earlier cut the shift is zero. At a nonpivot level $a < l$, the other side's jump u is at most η. Its cut orders at i are nonnegative there: for $a > 0$ this follows from $-u+\eta \geq 0$ in the depth, and for $a = 0$ from $-u + s + \eta \geq 0$. The transported cut orders are also zero below l. Their minimum is therefore nonnegative, and the resting depth 1 gives the rule for the third side at j. At $a = t$, the far jump of the other side is positive, bounded by $2b$, while its cut orders are at least $\min(0, -u + \eta)$ (also if the pivot lies at $a = 0$, since the shift $s \geq 0$ only increases the depth). Thus the moving order plus $2b$ is at least 1 there, using $\eta \geq 1$. Above and at l the moving order is at least $-u$ (the earlier cut has shift zero), and the resting depth is $-u'$. This proves the remaining rules with $t_j = 0$. Hence the direct movement is safe, and the long-gap point is the required stop. This is the argument of `pointsOver_longGapDepth_nonempty` and `ruleBound_le_inf_add_longGapDepth`.

Notice the role of the positive margin η. The strict 1 needed below the cut predecessor does not follow from mere complementary sign windows; it is supplied either by the depth at nonpivot places or by the positive margin at the pivot. Also, one must keep the stronger orders (51.12) for the long-gap case. Replacing them prematurely by the standard order lower bound $\rho_{-\eta}$ would not establish the nonnegative moving order at the other places.

51.13 Quantitative killers at sparse cut positions

We return briefly to the ideals of cokernel multipliers. Write $J_l(i;r)$ for the multiplier ideal at level l and letter i modulo r, using the notation of Section 50.7. The exact definition and the coefficient retraction imply the Nakayama property: if $p^\kappa \in J_0(i;p^{\kappa+1})$, then p^κ is an exact killer at i. Indeed p^κ lies in the exact multiplier ideal after multiplication by an element of the form $1-pu$ (move the error term divisible by $p^{\kappa+1}$ to the left side); that element is a unit by p-adic completeness. We use the exact one-step scalar calculus of Proposition 49.6, with the moving operator detailed above. Two consequences will be useful. If $p^c z^a \in J_0(j;r)$ and all jumps at j are at most n, then the multiplication by E^n to pass to level one, the depth-step trade of E for z, and the Frobenius return give

$$p^{c+d_0 n+v_p(n!)} z^{\lfloor (a+n)/p \rfloor} \in J_0(j[-1];r). \tag{51.13}$$

Also, with $p^c z^a \in J_0(i;r)$, removing the power of z gives

$$p^{c+a+v_p(a!)} \in J_0(i;r). \tag{51.14}$$

For (51.13) the factorial valuation counts the successive costs of trading E^n for z at level zero (each nonzero step costs $d_0+v_p(k)$), and decreasing a z-exponent to the nearest multiple of p in level one costs zero. For (51.14), first carry forward to level one and power z^{pa}, then lower one z-exponent at a time. The cost is $v_p((pa)!) = a + v_p(a!)$; return by Frobenius. These are `p_pow_mul_z_pow_div_mem_liftIdeal` and `p_pow_mul_z_pow_mem_liftIdeal_of_le` in the readable layout. The preservation of the exact multiplier ideals by the indicated scalar derivations still uses the compatibility of retraction with integral solutions (`mem_liftIdeal_of_eq_mul_smul_sub`); one cannot replace those ideals by arbitrary pointwise conditions on a quotient of the de Rham vector space. Under this ideal compatibility, these two formulas and their arithmetic consequences are now proved.

Suppose all jumps are at most w at all letters and we work modulo $r = p^{\kappa+1}$. The least $a \ge 0$ such that $z^a \in J_0(i;r)$ for all letters satisfies

$$(p-1)a \le e_0(w+\kappa). \tag{51.15}$$

This slightly sharpens Proposition 50.5 at exactly this modulus. The existence of some a follows in the same way, using the height of the original lattice modulo $p^{\kappa+1}$. To get the sharper inequality, go up to level one by multiplication by E^w. The element $z^{e_0(w+\kappa)}$ lies in $(p,E)^{w+\kappa}$, because z^{e_0} belongs to (p,E) in S. Multiplying by z^a, each monomial in the expansion has either an E-exponent at least w or a p-exponent at least $\kappa+1$: if the latter is at most κ, the remaining part is at least w. Therefore $z^{a+e_0(w+\kappa)}$ lies in the level-one ideal. Lower its exponent to the nearest multiple of p at zero cost and return by Frobenius. Minimality gives $a \le (a+e_0(w+\kappa))/p$, which proves (51.15). The small improvement here matters when inserting a minimal exponent into an inequality in which the chosen modulus itself depends on the cost.

Proposition 51.21 (An exact killer from J thin successors)**.** *Suppose all jumps are bounded above by w and at the J successive letters after i by g. Assume $8e_0 \le p^{J+1}$ and*

$$2Jd_0 g + (2J+4)\frac{g}{p-1} + \frac{8e_0 w}{p^{J+1}} \le \kappa. \tag{51.16}$$

Then p^κ kills the cokernel at i.

Proof. Work modulo $r = p^{\kappa+1}$ and begin at exponent a from (51.15). Apply (51.13) repeatedly along the J thin letters. One obtains $p^c z^n \in J_0(i; r)$, with

$$c \le J(d_0 g + v_p(g!)), \qquad n \le \frac{a}{p^J} + \frac{g}{p-1}.$$

The second inequality follows by starting with a and iterating the map $x \mapsto \lfloor (x+g)/p \rfloor$; the geometric sum of g/p is bounded by $g/(p-1)$. Legendre's estimate gives $v_p(g!) \le g/(p-1)$ and $v_p(n!) \le n$. From (51.15), since $p \le 2(p-1)$ and $8e_0 \le p^{J+1}$,

$$\frac{a}{p^J} \le \frac{2e_0(w+\kappa)}{p^{J+1}} \le \frac{2e_0 w}{p^{J+1}} + \frac{\kappa}{4}.$$

Thus the total cost of (51.14) is bounded by

$$\begin{aligned} c + n + v_p(n!) &\le J\left(d_0 g + \frac{g}{p-1}\right) + 2\left(\frac{2e_0 w}{p^{J+1}} + \frac{\kappa}{4} + \frac{g}{p-1}\right) \\ &\le \kappa, \end{aligned}$$

where the last inequality is exactly (51.16) divided by two and added to $\kappa/2$. So $p^\kappa \in J_0(i; p^{\kappa+1})$. Apply the Nakayama property. □

This is `killsCokernel_zero_of_jumps_le`. It explains the bound in terms of p^{J+1} rather than in terms of the full rank or the number of letters. Large jumps far from the cut are admitted into the initial exponent but decreased by a geometric factor during Frobenius return.

We now combine the previous results in the numerical cut scheme of readable `CutExponentBounds.lean`. We state the parameters with slack, so that the construction can be used in both of its principal specializations.

Theorem 51.22 (Cut-window bound)**.** *Suppose the two sides have jumps in the opposite windows of Section 51.12; let $s_i \ge 0$ be the shifts at letters. Fix nonnegative parameters $W, X, \delta, \kappa, b, n, J, s_0, G$ satisfying*

$$\begin{gathered} \eta \ge 1, \quad V + 4\eta \le W, \qquad n + \lfloor \log_p(2b) \rfloor + (J+1+d_0)W + d_0 G \le X, \\ 12((J+1)X + \delta) < \kappa, \qquad 8e_0 \le p^{J+1}, \qquad 2e_0 b \le p^{J+1}\delta, \\ \beta_i + s_i + W \le 2b, \quad s_i + W \le pn, quad G \le (p-1)n, \quad \beta_i + W \le (p-3)s_0 \quad \text{if } p \ge 5. \end{gathered}$$

Assume every letter has a strictly preceding cut within $2J+1$ steps, and every cut i is one of the following:

(i) $s_i = 0$, and each of the J successive letters j satisfies $\beta_j + W \le G$;

(ii) the chart is unramified, $p \ge 5$, $s_i < \beta_i$, and $s_0 \le \kappa + s_i$.

Finally, at each cut, the interval separating it from the nearest preceding cut is plain or is a long gap in the sense of Section 51.12, with s_0 as above and some ω bounding the pre-pivot windows. Then $p^{51\kappa}$ annihilates every paired cup of crystalline classes of the two sides.

Proof. First all three sides have jumps at most $\beta_i + W$. For the individual sides this is immediate from their windows. For the Tate dual, if the two jumps are both far they have opposite signs and $|-u-u'-1| \le V+1$; if both are low the bound is $2\eta+1$; if only one is far the bound is $\beta_i + V + \eta + 1$. All are at most $\beta_i + W$ since $W \ge V + 4\eta$, $\eta \ge 1$. This also covers the jump zero of an extension. At a cut of type (i), Proposition 51.21 applies with

$w = 2b$, $g = G$. Indeed the cost in (51.16) is bounded by $(4J + 4)X + 8\delta < \kappa$. For type (ii), use the exact unramified bound of Proposition 50.6, with total budget $\kappa + s_i \geq s_0$. Thus $p^\kappa E^{s_i}$ kills the cokernel of each of the three sides at every cut.

Fix a cut i and set $l = l_i \leq 2J + 1$. For either side, initial points over $p^\kappa c_i$ exist with order and depth ρ_{-s_i}. Increase the depth to the system $D(s_i, \eta, l + 1)$. The needed steps occur only at $0 \leq a < 2J + 2$ and lie in the two windows of radius

$$r_a + W, qquadr_a = s_i \mathbf{1}_{a=0}$$

centered at zero and the far negative of the sign of the jump window at $i[-a]$. Indeed at level zero the steps range from $-u - s_i$ to $-u + s_i + \eta - 1$ and at the other levels from $-u$ to $-u + \eta - 1$. Since $V + 4\eta \leq W$, the radius is sufficient. Here $r_a + W \leq pn$ and the absolute values of all integers in these windows are at most $2b$, by the stated shifted far bound. At level zero either $r_a = 0$ or $d_0 = 0$ by the cut types, hence always $r_a = 0$ or $d_0 + 3a = 0$. The estimate of Section 51.7 therefore bounds the cost at each level by

$$8n + 2\lfloor \log_p(2b) \rfloor + 5W \cdot 6(J + 1 + d_0) \leq 30X.$$

The cardinal bound uses $4W + 2 \leq 5W$ for $W \geq 4$, and $d_0 + 3a \leq 6(J + 1 + d_0)$. Thus the full sum of costs is at most $60(J + 1)X$. The chosen κ gives

$$\kappa + 4\delta + 60(J + 1)X \leq 25\kappa.$$

This estimate is generous enough for both types of stop. Apply Proposition 51.14 and Lemma 51.12, then multiply by the unused power of p to obtain a point over $p^{25\kappa} c_i$ with strong orders (51.12) and the required depth.

These strong orders imply the standard orders $\rho_{-\eta}$. For $a > 0$, if $-u > 0$ then improved order is at least $-u$; otherwise $D(s_i, \eta, l + 1; a, u) \geq -u$ and $\min(0, D) \geq -u$. At $a = 0$, the same is true with $-u - s_i > 0$ or with $-u - \eta \leq 0$. The remaining possibility $-u - s_i \leq 0 < -u - \eta$ would put a negative jump in $[-s_i, -\eta)$, impossible when $s_i = 0$ or when $s_i < \beta_i$ and the far negative values satisfy $-u \geq \beta_i$. Thus the implication holds. The depth $D(s_i, \eta, l + 1)$ is also the depth required at a plain cut in Section 51.12.

If the cut is plain, take the standard point of the down side as stop; its depth makes the movement safe. At a long gap, the preceding cut has shift zero. The unramified bound with shift s_0 at the pivot kills its cokernel with no p-cost, since all jumps are at most $(p - 3)s_0$. At the preceding cut use the already established p^κ-killer. The radius condition and long-gap hypothesis are exactly those used in Section 51.12; after the Taylor extraction and transport, only the depth steps $(a, 0)$ with $t < a < l$ are needed. These are among the two-window steps, so the full cost, including $\kappa + 4\delta$, is at most 25κ. Thus the required stop exists over the same constant as the standard point, and the move at this stop is safe by the strong order calculation in that section.

All conditions of Theorem 51.20 now hold with $e_1 = e_2 = 25\kappa$, and the third-side cost is κ. The total is 51κ. □

51.14 Two choices of cuts

We explain the combinatorial choice in the sparse-thick-letter case and the two-size case, which have quite different hypotheses. Let T be the set of thick letters (the letter i is thick if $\beta_i = b$ and thin if $\beta_i = g$). Use the direction of Frobenius to say that a letter is followed by another a positive number of steps later.

Sparse thick letters. Assume no three thick letters follow one another with the two successive gaps both between 1 and J, with $J \geq 1$. Mark a cut when the following J letters are all thin. We first prove two combinatorial facts. If a thick letter j is not a cut, then some thick letter follows it within J steps. No thick letter can precede j within J steps, by the no-three hypothesis. Therefore the letter $J+1$ steps before j is a cut. Next, for any letter i, either $i[-(J+1)]$ is a cut or some thick letter $i[-r]$ with $1 \leq r \leq J$ follows it before i. In this second case either $i[-r]$ itself is a cut or $i[-(r+J+1)]$ is one, by the previous fact. Thus every letter has a preceding cut within $2J+1$.

Consider the nearest preceding cut $i[-l]$ of i and a thick letter $i[-r]$ with $0 < r < l$. Since there are J thin letters after the cut, we have $l - r \geq J+1$. The thick letter $i[-r]$ is not a cut (by nearestness), hence $i[-(r+J+1)]$ is a cut, implying $r+J+1 \geq l$. Therefore $r+J+1 = l$. In particular there is at most one interior thick letter, necessarily $J+1$ steps after the cut. These are the combinatorial assertions `hasThinSuccessors_stepBack_add`, `exists_hasThinSuccessors_stepBack`, and `eq_zero_or_add_eq_of_hasThinSuccessors`.

Suppose now the thin windows lie in the low region, i.e. $g + V \leq \eta$. Assume either all signs are equal, or the chart is unramified with $p \geq 7$ and $10p^J\eta \leq b$. In the constant-sign case all cut intervals are plain. Otherwise if an interval is not plain, the only interior thick letter is a pivot of sign opposite to the sign at i. Every other letter than the pivot and i is thin. Put $W = 5\eta$ (here $V \leq \eta$), use $b_* = b + W$ in place of b in Theorem 51.22, let $s_i = 0$ for all i, $n = G = 2W$, and

$$s_0 = \left\lfloor \frac{b + 10\eta}{p-3} \right\rfloor + 1.$$

When the sign is not constant, the pivot satisfies the long-gap inequality with $\omega = \eta$:

$$s_0 + \eta(1 + p + \cdots + p^J) \leq \frac{b+10\eta}{4} + 1 + 2p^J\eta < b.$$

Indeed $p \geq 7$, $p^J \geq 7$ and $b \geq 10p^J\eta$ imply the strict final inequality (the right-hand side of the first inequality is at most $b/4 + b/28 + 1 + b/5 < b$). All the pre-pivot windows are thin. Thus exactly one long gap or a plain interval occurs, and all cuts are of type (i).

Take

$$8e_0 \leq p^{J+1}, \quad 2e_0(b+5\eta) \leq p^{J+1}\delta, \quad 20(J+1+d_0)\eta + \lfloor \log_p(2(b+5\eta)) \rfloor \leq X,$$

where the first inequality is automatic for the unramified $p \geq 7$ case. The cut-size bounds hold: the cost requires $2W + (J+1+d_0)W + 2d_0W + \log$ which is at most $4(J+1+d_0)W + \log$; the shifted windows end at or below $2b_*$, since $\beta_i \leq \max(b, W)$. The bound for $p \geq 5$ follows from the definition of s_0. Choose $\kappa = 12((J+1)X + \delta) + 1$. The theorem gives an annihilation exponent 51κ, bounded by

$$10^3(\delta + (J+1)X)$$

since the latter parenthesis is a positive integer. This is the readable version `isScaledCupCoboundary_sparse_thick`. The compact source uses differently packaged constants and a directly expressed depth budget as in Chapter 49. The exact comparison of the two sets of numerical hypotheses is separate; the cut mechanism is common to both.

Two sizes in arbitrary order. Now assume the chart unramified, $p \geq 7$, $0 < g \leq b$, and suppose

$$b < Kp^2, \qquad 2g < Kp, \qquad W \leq K, qquad V + 4\eta \leq W.$$

Use $J = 1$ and put $q = \lfloor b/p \rfloor$. Set

$$s_i = \begin{cases} \max(0, q - 7K) & \text{the successor of } i \text{ is thick,} \\ 0 & \text{otherwise.} \end{cases}$$

The cuts are the letters for which $s_i < \beta_i$. Every thick letter is a cut ($s_i \le b/p < b$), and every letter with a thin successor is a cut ($s_i = 0 < g$). Thus one of the first two preceding letters of every i is a cut: if $i[-1]$ is thick it is itself a cut, and if thin then $i[-2]$ has a thin successor. If the nearest preceding cut is $i[-1]$, the interval is plain. If it is $i[-2]$, then $i[-1]$ is thin and not a cut, so i itself is thick; moreover $s_{i[-1]} \ge g$, giving $g + 7K \le q$, and $s_{i[-2]} = 0$. If the two signs agree the interval is plain. If not, the pivot is i, $t = 0$, $J = 1$, and with $s_0 = q + 12K$, $\omega = g + W$, we have the long gap

$$s_0 + \omega(1+p) \le q + 12K + (q - 6K)(1+p) = (p+2)q - 6(p-1)K < b.$$

The last inequality uses $pq \le b$, $q < Kp$ and $2Kp < 6(p-1)K$. The other long-gap conditions were just verified.

To verify the cost scheme take

$$b_* = b + W, \quad n = K, \quad G = g + W, \quad X = 3K + \lfloor \log_p(2b_*) \rfloor, \quad \delta = 4K, \quad \kappa = 12(2X + \delta) + 1.$$

Since $d_0 = 0$, the cost bound is immediate from $W \le K$. The inequality $8e_0 \le p^{J+1}$ holds since $e_0 = 1$, $p \ge 7$. Also $2b_* \le 4Kp^2$, and $s_i + W \le pK$ since $q < Kp$. We have $G \le Kp/2 + K \le (p-1)K$ for $p \ge 4$. The shifted far bound $\beta_i + s_i + W \le 2b_*$ is obvious from $s_i \le b$. For the final far bound at $p \ge 5$, write $b = pq + r$, $0 \le r < p$; then

$$b + W - (p-3)(q + 12K) = 3q + r + W - 12K(p-3) < 0.$$

Indeed $3q + r + W < 3Kp + p + K \le 4Kp + K$ since $K \ge 1$, whereas $12(p-3)K \ge 48K$ for $p \ge 7$ and in fact $12(p-3) > 4p + 1$ for such primes. Finally, any cut with thin successor is of type (i), and a cut with thick successor satisfies $s_0 \le s_i + 19K \le s_i + \kappa$ and $s_i < \beta_i$, hence of type (ii). Theorem 51.22 gives

$$51\kappa = 51(12(10K + 2\lfloor \log_p(2b_*) \rfloor) + 1) \le 10^4(K + \lfloor \log_p(2b_*) \rfloor).$$

This is the bound `isScaledCupCoboundary_any_pattern`. In the compact source, the counterpart is `scaledCupBounds_two_sizes`. Its role is precisely to handle a succession of many thick letters without requiring the sparsity used in the previous case: a shift at a cut replaces the unshifted bound from the next J thin letters.

51.15 A multiplicity-sensitive step in the few-low-jumps case

As seen in Section 50.8, the few-low-jumps length estimate, unlike the preceding rank-free bounds, must take account of the multiplicities of the low jumps. We now reconstruct the key correction of relations on the actual constants. In the unramified chart, write $Q(o)$ for the order module on the kernel lattice Y at a fixed letter i, intersected with its localized Kisin module. Its image in the extension is the submodule of vectors over zero with orders o. We use the comparison assertion that the latter submodule identifies with the order module of the kernel on the specified common jump sets, precisely `exists_inclKer_eq_of_kerWithOrders` in readable `SidePoints.lean`; this identification is an integral functoriality assertion, not automatic from splitting an extension as underlying $\mathbb{Z}_p$-modules.

Lemma 51.23 (Primitive relation up to E)**.** *Let m be the Hodge–Tate multiplicity of jump u at $i[-a]$, and suppose $o(a,v) > o(a,u)$ for all smaller jumps $v < u$ there. Put o' equal to o with $o(a,u)$ raised by one. For any $f_0 m + 1$ vectors y_j in $Q(o)$, where $f_0 = [\Lambda : \mathbb{Z}_p]$, a primitive $\mathbb{Z}_p$-linear combination of the y_j belongs to*

$$EQ(o) + Q(o').$$

Proof. Choose the forms generating the degree-u part of the filtered dual space modulo degree $u-1$, at most m forms by the definition of multiplicity. For each y_j evaluate those forms at level a, divide by the common $t^{o(a,u)}$, and take the residue in C_p of the resulting integral values. This gives row vectors in a vector space of dimension at most m. The maximal-minor argument of Section 50.8 gives a subset $y_{j_1},\dots,y_{j_d}$ with $d\le m$ such that each residue row vector is an integral ($\mathcal{O}_{C_p}$-linear) combination of the selected ones. In fact choose a nonzero minor of maximum absolute value among all the minors of size equal to the rank of the rows; after choosing its rows as a basis, Cramer's rule bounds all expansion coefficients by one. If the rank is zero the residues already vanish. Use the surjective $\theta_a : A\to\mathcal{O}_{C_p}$ to lift the coefficients to A (equivalently lift at θ and apply φ^a). Denote the lifts by v_{jk}. Then

$$y_j-\sum_k v_{jk}y_{j_k}\in\mathcal{M}_i^Y(o').$$

Indeed the chosen leading residues cancel, and every form of degree u differs from a combination of the chosen forms by one of lower degree; the lower-degree forms already have order at least $o(a,u)+1$ by the strictness hypothesis. At the other levels and jumps the old orders are preserved by linearity over A. The decomposition of forms by degree here is the filtered basis argument for the Hodge–Tate jumps after the comparison, in the source `exists_span_formsOfType_mod_pred`.

Retract the displayed relation to the localized Kisin module. If $\lambda_0 : A\to S$ is its chosen scalar coordinate, then

$$y_j-\sum_k \lambda_0(v_{jk})y_{j_k}\in Q(o')$$

by the preservation of orders under retraction. Write $\lambda_0(v_{jk})=w_{jk}+Ez_{jk}$ with $w_{jk}\in\Lambda$ and $z_{jk}\in S$, by the unramified division used in the Taylor proof. The vectors $(w_{jk})_k\in\Lambda^d$ are $\mathbb{Z}_p$-linearly dependent since there are $f_0m+1>f_0d$ of them. Choose a relation and divide by a nonzero coefficient of least valuation to arrange one coefficient equal to 1. On taking the corresponding combination the w_{jk} terms cancel; the Ez_{jk} terms lie in $EQ(o)$, and the remainder in $Q(o')$. □

This supplies the relation `exists_sum_smul_mem_locKisinPart_sup` in readable `HodgeExpansion.lean`, including the preservation of the leading coefficients by projection. The regression to the actual constants still requires removing the $EQ(o)$ term; treating that term as zero merely because another filtration coordinate was raised would be incorrect.

For an extension let $X(o)$ denote the no-pole localized Kisin points over some constant and with depth o. Thus they have orders σ_0 and depth o; put $D(o)=qX(o)$ for the corresponding submodule of constants. Let $M(o)$ be the submodule of the extension consisting of the localized kernel vectors with orders o. When $o\ge\sigma_0$, $M(o)\subset X(o)$.

Lemma 51.24 (Division by p in the depth condition)**.** *Assume that no integer divisible by p lies strictly between $o(0,v)$ and $o(0,v)+p^a$ at any jump v. Let x have depth o over a constant. If $w\in EM(o)$ and $px-w$ has depth o' over a constant, where o' raises $o(a,u)$ by one, then there is $m_0\in M(o)$ for which $x-m_0$ has depth o'.*

Proof. Use the common zero-cost operator A_0 for the steps $(0,n)$ lying strictly between $o(0,v)$ and $o(0,v)+p^a$, over all jumps v. All the steps are free by $p\nmid n$ and $d_0=0$. This operator preserves all depths already satisfied and differs from the identity by a kernel correction of the input depth. Since $w\in EM(o)$, it has order o raised by one at level zero, and depth at least that. Consequently A_0w has the same order and has depth o raised by p^a at level zero. We claim its actual order is also at least that latter system. If $o(0,v)+1>0$, Lemma 51.12

gives this. Otherwise $o(0,v)+1 \le 0$ and there is no multiple of p strictly between $o(0,v)$ and $o(0,v)+p^a$; in particular zero is not in that open interval, so $o(0,v)+p^a \le 0$. The same improvement lemma again gives the required order. At other levels the original orders are retained. Hence

$$A_0 w = E^{p^a} w_1, \qquad w_1 \in M(o).$$

In the unramified chart $E^{p^a} - E_a = pr$ for some $r \in S$ (as in (50.13)). Set $m_0 = rw_1 - (A_0 x - x) \in M(o)$. Then

$$p(x - m_0) = A_0(px - w) + E_a w_1.$$

The first term has depth o' since the operator preserves that depth. The second has actual orders o raised by one at level a, which are at least o', hence it also has the required depth. Multiplication by p has no effect on the de Rham orders because p is a unit of B_{dR}^+. Therefore $x - m_0$ itself has depth o', with unchanged projection. □

This is `exists_sub_mem_deepElements_of_smul_sub_mem`. It is not a division inside the integral lattice by a nonunit p without control: the correction m_0 is explicitly integral and over zero.

Theorem 51.25 (Rank-sensitive colength of one step)**.** *Assume $o \ge \sigma_0$, the strict lower-jump hypothesis of Lemma 51.23, and the no-multiple hypothesis of Lemma 51.24. Then for o' raising $o(a,u)$ by one,*

$$\mathrm{length}_{\mathbb{Z}_p}(D(o)/D(o')) \le c(a, o(a,u)) f_0 m.$$

Proof. Consider $X(o)/(X(o') + M(o))$, which surjects onto $D(o)/D(o')$. By the operator of Theorem 51.13, this quotient is killed by p^k with $k = c(a, o(a,u))$: applying it to $x \in X(o)$ gives an element of $X(o')$ differing from $p^k x$ by a vector of $M(o)$. Preservation of the no-pole condition follows from $o \ge \sigma_0$.

Suppose we are given $f_0 m + 1$ elements of this quotient killed by p, represented by $x_j \in X(o)$. Write

$$px_j = x_j' + m_j, \qquad x_j' \in X(o'), \quad m_j \in M(o).$$

Apply Lemma 51.23 on the kernel to the m_j and use its identification with the extension kernel. A primitive combination $\xi = \sum_j b_j x_j$ therefore satisfies

$$p\xi - w \in X(o'), \qquad w \in EM(o).$$

Indeed the combination of the m_j equals $w + v$ with $v \in M(o') \subset X(o')$. Lemma 51.24 gives $m_0 \in M(o)$ with $\xi - m_0$ of depth o'. Since m_0 has no pole ($o \ge \sigma_0$), $\xi - m_0 \in X(o')$. Thus the primitive combination is zero in the considered quotient.

It follows that its p-torsion is an $\mathbb{F}_p$-vector space of dimension at most $f_0 m$. The dimension bound on p-torsion and the annihilator p^k are enough even without finite generation of $X(o)$: for a group killed by p^k, each successive p^j-torsion quotient injects into its p-torsion via multiplication by p^{j-1}. Hence its length is at most $k f_0 m$. The surjective image $D(o)/D(o')$ has no larger length. □

This proves `relLength_deepConstants_raiseOrder_le`. It supplies the rank-sensitive correction left open in Section 50.8. In that section the subsequent numerical sum of single steps yields the factors $(2s+1)n_i$ and $3(W+1)r_{i[-1]} + 6sn_{i[-1]}$, rather than the full rank times the largest gap. The additional initial-depth construction and the precise opposite-window safety inequalities in the last step of the few-low-jumps theorem still require verification: the present proof provides their central one-step colength estimate, but not a substitute for these final case checks.

51.16 Scope of the integral lifting reconstruction

We record exactly what is used from outside this chapter. The integral period-ring inputs are the structural closure, division and comparison properties listed in Sections 51.1, 51.2, 51.3, and 51.6. The Frobenius-basis and scalar inverse claims, the rank-free depth step (including zero), the long-gap extraction after the radius-span estimate, the continuous cup coboundary and the full cut-walk calculation are proofs from those stated hypotheses. The thin-cut annihilation and the two choices of cuts keep their source numerical constants in the readable layout. Where a result is invoked in the compact layout under differently packaged inequalities, the hypotheses must be compared explicitly rather than replacing its bound solely by a name from the other layout.

The special tame-type case uses a chart with a tame generator of inertia and a polynomial q annihilating its action on the crystalline invariant module. The following elementary step in that proof is useful to distinguish from its more specialized integral estimates. If a linear operator T on an extension $0 \to Y \to L \to C \to 0$ acts trivially on the quotient and $q(T)$ vanishes on the kernel, then $q(T)(T-1) = 0$ on L: the image of $T-1$ is contained in Y. At the tame chart the operator T is the linearized action of a chosen inertia element on periods; the same argument applies after the exact invariant-module comparison. Thus the relevant polynomial is $q(X)(X-1)$, of degree $\deg q + 1$. A primitive e_0-th root in the residue field gives e_0 distinct evaluation points; a monic polynomial of degree d has at most d roots among them. This counting is the mechanism that avoids a loss proportional to all e_0 residue classes: the exceptional evaluations are at most $\deg q + 1$. The integral passage from the polynomial relation to annihilation of the Kisin cokernel, however, also needs the construction of the tame chart, a near-identity divisibility estimate for its tame generator, and the scalar ideal recurrence at its exceptional evaluation points. These are the remaining tame inputs of `TameChart.killsCokernel_of_isKilledBy` and `Lift.exists_tameChart`, compact `New/Lift/SparseThickLetters.lean`. A bound for the order of a rational representation alone would not supply them.

The dependence on the Hodge–Tate frame is concentrated in the integral comparisons, not in the elementary stationary or moving arguments. To finish an entirely independent proof one must reconstruct the first half of the radius lemma and the chart comparisons as well as the special tame and few-low-jump obligations just described, or show that a standard theorem provides precisely these integral statements with their constants. Consequently we do not assert that the local crystalline block of the entire GRH argument has become independent of the formal development. The value of the reconstruction is that one no longer needs to import the nonstandard Frobenius walk, the long-gap extraction, or the abstract integral-basis argument as opaque local theorems.

Chapter 52

Closing the local comparison and lifting arguments

Subsequent-edition update. The status remarks in this chapter document the earlier reconstruction. For the new proofs, corrections and superseding status of individual obligations, see Chapters 53 and the final audit in Chapter 57.

We retain the notation of Sections 50 and 51. Here we address the structural hypotheses explicitly left open there. In particular the ring B below denotes the maximal crystalline period ring, not the divided-power integral lattice. In the compact source it is the maximal ring, including its chosen localization, that defines the crystalline property; no equivalence of integral structures with the classical divided-power ring is needed in these arguments. We keep track of the hypotheses of each statement, rather than use a general equivalence of categories to justify an integral estimate with a prescribed exponent.

52.1 Completeness and the elementary maximal period ring

Put $A = W(\mathcal{O}^\flat_{C_p})$, and use τ, μ, ξ as in Section 51.1. We use the basic perfectoid facts that $\mathcal{O}^\flat_{C_p}$ is a complete rank-one valuation domain, that A is p-torsion-free and p-adically complete with reduction $\mathcal{O}^\flat_{C_p}$, and that the Fontaine map θ is onto with kernel generated by ξ. These are standard; for the last assertion see B. Bhatt, M. Morrow and P. Scholze, *Integral p-adic Hodge theory*, Publ. Math. IHÉS **128** (2018), 219–397, Lemma 3.10 and Remark 3.11 (the chosen cyclotomic ξ is distinguished). The same remark applies to $E = z^e u + p$ whose reduction has positive valuation and whose theta value is zero, where $u \in S^\times$ and z is the Teichmüller coordinate of an Eisenstein chart. In the latter case the first Witt coordinate after the zeroth is a unit. We explain the further ring properties needed here directly.

Lemma 52.1. *If $z \in A$ has reduction of finite strictly positive valuation, then A is (p, z)-adically complete and separated. The ideal qA is closed for this topology whenever $p \nmid q$.*

Proof. For each k the quotient A/p^k is complete for the z-adic topology. To see this, the assertion at $k = 1$ is completeness of the valuation ring. Induct using $0 \to p^{k-1}A/p^k \to A/p^k \to A/p^{k-1} \to 0$. Multiplication by z on each quotient is injective: if $zy \in p^k A$, primality of p and $p \nmid z$ yield $y \in p^k A$. Consequently $z^m(A/p^k) \cap p^{k-1}(A/p^k) = z^m p^{k-1}(A/p^k)$, which makes the induced topology on the kernel precisely its own z-adic topology. Compatible lifts give completeness, and the same equality gives separatedness. The ideals (p^k, z^m) are cofinal with the powers of (p, z); taking compatible limits first modulo p^k and then over k proves the assertion.

Similarly q acts injectively modulo each p^k. Its image modulo p is a principal ideal of the valuation ring, hence closed. If qy_j is a z-adically Cauchy sequence modulo p, then y_j is Cauchy, because subtraction of the fixed value $v(\bar{q})$ from a lower bound on $v(\bar{q}(\bar{y}_i - \bar{y}_j))$ leaves a lower bound tending to infinity. Successively subtracting a lift of the limit modulo p shows the same continuity of division by q modulo every p^k; the induction uses the preceding kernel-topology equality. Thus at each finite Witt length the limit of multiples is a multiple; the quotient is unique since multiplication is injective. Taking the p-adic limit proves closedness of qA. □

Write $P = A[X]/(pX - \xi)$ and $A_{\max} = \widehat{P}^{\,p}$. The map $P \to A[1/p]$ sending X to ξ/p is injective. Indeed, if multiplication by p kills the class of $f(X)$, write $pf = (pX - \xi)g$. Reducing modulo p in the polynomial ring over the valuation domain gives $g = pg_1$, since $\bar{\xi} \neq 0$; then $f = (pX - \xi)g_1$. After inverting p the asserted map is an isomorphism. In particular P is p-torsion-free.

The completion of a p-torsion-free ring is again p-torsion-free and its quotient modulo p is that of the original ring. An elementary verification uses the systems P/p^k: if an element of the completion reduces to zero modulo p, its components in pP/p^kP have unique quotients in $P/p^{k-1}P$, which form a compatible system. The same verification gives cancellation of p in the completion.

We therefore have

$$A_{\max}/p = (A/(p, \xi))[X].$$

The variable X is a non-zero-divisor there. Its powers are also non-zero-divisors, and it is not a unit. As a result, for every $n, m \geq 0$,

$$p^m b \in X^n A_{\max} \quad \Longrightarrow \quad b \in X^n A_{\max}, \qquad X^n b \in p^m A_{\max} \quad \Longrightarrow \quad b \in p^m A_{\max}. \tag{52.1}$$

For the first implication, cancel p one at a time: if $pb = X^n c$, reduction modulo p shows $c = pc_1$. For the second, reverse the roles in the same cancellation. Moreover $\bigcap_n X^n A_{\max} = 0$: reduction modulo p shows that an element of this intersection has zero image in a polynomial ring, and (52.1) permits iteration after dividing by p. The separatedness also holds in $R = A_{\max}[1/p]$ by the first implication in (52.1).

The extension of θ to P sends X to zero. Its kernel in P is (X), since $P/(X) = A/(\xi) = \mathcal{O}_{C_p}$. The extension to $A_{\max}$ is onto with the same kernel. Indeed P is dense and $\mathcal{O}_{C_p}$ is p-adically complete; the quotient by X remains p-adically complete and separated by the cancellation and closed-image argument used in (52.1). More explicitly, a sequence Xb_j converging p-adically has Cauchy quotients b_j by (52.1), so (X) is closed, and $A_{\max}/(X)$ identifies with the completion of $P/(X)$. It follows that $R/(X) = C_p$. Thus R is a domain: a nonzero element has finite X-adic order, and products of elements of order zero remain of order zero. The X-adic completion of R is a complete discrete valuation ring with residue field C_p. We below verify its identification with the usual B_{dR}^+. The standard parameter there is $t = \log[\epsilon]$, associated to ξ; see Brinon–Conrad, cited below, Proposition 4.4.8, for this uniformizer and its cyclotomic transformation law. Because $\theta|_R$ has kernel (ξ), a repeated cancellation shows

$$A \cap \xi^n R = \xi^n A.$$

The injectivity of $A \to A_{\max}$ follows either from these divisibilities or from the contraction calculation: using (52.1), exactly the proof of Lemma 51.1 yields $A \cap p^m A_{\max} = (p, \xi)^m A$, while an element mapping to zero satisfies this for every m and is zero by Lemma 52.1. In the contraction proof cancellation is first performed in P or its p-torsion-free completion, so it does not presuppose the injection.

The identity $\varphi(\xi) = \sum_{j=0}^{p-1}(1+\mu)^j$ gives $\varphi(\xi) = p + \mu H(\mu)$ with integral polynomial H. Since $\mu = \tau pX$, $\varphi(\xi)/p = 1 + \tau X H(\mu)$ belongs to $A_{\max}$. The image of τ is nilpotent modulo

p in $A_{\max}$, because $\bar{\xi} = \bar{\tau}^{p-1}$. Consequently $\varphi(\xi)/p$ is a unit, and Frobenius extends to $A_{\max}$ by its presentation, hence to R. The same presentation extends the Galois action since $g(\xi)$ is associated to ξ.

Lemma 52.2. *If $\xi = pb + \nu c$ with $b, c \in A$, then*

$$p^n A_{\max} \subseteq A + \nu^n A_{\max}.$$

Proof. For a polynomial $F(X) \in A[X]$, divide by the monic polynomial $(X - b)^n$. The remainder is $\sum_{j<n} a_j (X - b)^j$. After multiplying by p^n, its terms belong to A because $p(X - b) = \nu c$. The rest belongs to $\nu^n P$. Write an arbitrary element of the completion as a convergent sum $\sum_{k \geq 0} p^k F_k(X)$ with $F_k \in A[X]$ (choose representatives successively modulo powers of p). Apply the polynomial decomposition term by term. The two resulting sums converge in A and in $A_{\max}$ respectively because their k-th terms are divisible by p^k. This is the required containment. □

In particular, at $\nu = \xi$, division by p^n (or by a larger p-power to clear the denominator of a given element of R) shows that $A[1/p]$ is ξ-adically dense in R. The contraction $A \cap \xi^n R = \xi^n A$ proved above identifies the induced filtrations. Consequently the ξ-adic completion of R is exactly the ordinary definition of B_{dR}^+, the $\ker \theta$-adic completion of $A[1/p]$. Taking $\nu = \tau, \xi$, or a chart coordinate z, gives exactly the assertions in Section 51.1 used in the Frobenius contraction. The approximation by p-power multiples of elements of A follows from polynomial density and clearing one power of p as noted there. Together with Lemma 52.1 this establishes the structural assumptions (i)–(iii), and Lemma 51.1, without invoking a comparison of integral crystalline lattices. In particular no divided-power factorials have been silently substituted for powers in $A_{\max}$.

52.2 Retractions and the scalar ring of the chart

Suppose $S = \Lambda[[z]]$ is the Eisenstein chart inside A, $\Lambda = W(\mathbb{F}_{p^f})$, $E = p + z^e u$ with $u \in S^\times$ and E associated to ξ. This map from the power-series ring is injective: modulo p the image of the first nonzero coefficient times a power of z dominates all later terms in the rank-one valuation, since a nonzero coefficient in the finite field has valuation zero. Then cancel successive powers of p. The map is Frobenius-stable, with $\varphi(z) = z^p$. All topologies mentioned in Lemma 52.1 for z and ξ are equivalent, by cofinality of powers of their reductions in the valuation ring.

Select a basis of $A/(p, z)A$ over $\Lambda/p\Lambda$ containing 1, and lift it to A (the lift of 1 is 1). The (p, z)-adic associated graded of A is the polynomial algebra on the two classes of p, z over $A/(p, z)A$. For clarity, this uses only regularity of the sequence p, z: mod p, independence of the term involving just z^n follows by cancelling z^n, then subtract that term and divide by p. Induction on degree gives independence of the monomials $p^i z^{n-i}$ modulo the next filtration step; here cancellation $(I^{n+1} : p) = I^n$ for $I = (p, z)$ follows by reducing an expression in I^{n+1} modulo p, using that z is regular modulo p, and then cancelling p in A. This proves the associated-graded assertion. The analogous statement for S follows from its power-series presentation. Thus the map from the completed direct sum of copies of S indexed by the chosen basis to A is an isomorphism on associated graded pieces. Completeness and separatedness prove that the map is an isomorphism: lift any target successively modulo powers of I, and use injectivity on graded pieces to prove uniqueness. We have obtained an adic basis (b_j) with S-linear coordinates λ_j and a distinguished coordinate $\lambda_0(1) = 1$. Finite partial sums approximate every vector to arbitrary I-adic precision.

Every S-linear endomorphism of A carries $(p, E)^m A$ into itself. Since E is associated to ξ, the contraction of $p^m A_{\max}$ to A is this ideal, and the endomorphism extends uniquely

to a p-adically continuous map of R preserving $A_{\max}$: first extend to the dense $A[1/p]$ by division by p; the contraction identity shows that the norm bound on this subspace is exactly preservation of the integral ball. Then take the completion. In particular all coordinate maps extend after including $S \subset A$. Define S_B to be the subring of elements of R whose multiplication commutes with all the extended S-linear endomorphisms. This agrees with the bicommutant in the source: any S-linear map on R preserving the integral ball is continuous, and its commutation identities with the extensions can be tested on the dense $A[1/p]$ as needed. The two facts actually used are the following.

Lemma 52.3. *The extended coordinates $\widehat{\lambda}_j$ have image in S_B. They are S_B-linear and $\widehat{\lambda}_0|_{S_B} = 1$ (identity). Frobenius carries S_B into itself. For every $y \in R$ and integer m, sufficiently large finite sets of indices F satisfy*

$$y - \sum_{j \in F} b_j \widehat{\lambda}_j(y) \in p^m A_{\max}.$$

For $y \in S_B$ one has $y = c + zw$ with $c \in \Lambda[1/p]$, $w \in S_B$.

Proof. For an extended coordinate L_j, the equality $G(vL_j(a)) = G(v)L_j(a)$ holds for $a \in A$, all $v \in R$, and every continuous S-linear endomorphism G of R, because $L_j(a) \in S$. Extend this equality to $a \in R$ by continuity. Thus $L_j(R) \subset S_B$. The linearity is the defining commutation condition; evaluation at 1 gives the assertion about L_0.

We next identify S_B with the closure of $S[1/p]$ in R; Frobenius stability will then follow by continuity, since $\varphi(S) \subset S$.

Indeed, for $y \in R$, approximate it by $p^{-k}a$, $a \in A$, with arbitrarily small error in the integral ball. Since all partial-sum operators have norm at most one (multiplication by each b_j preserves $A_{\max}$), approximation of a to high (p, E)-adic precision implies approximation of y to arbitrarily high p-adic precision. This proves the displayed assertion. If $y = L_j(y')$, then approximate y' by elements of $A[1/p]$ to see y belongs to the closure of $S[1/p]$; conversely that closure is contained in S_B since the centralizer conditions are closed. Since $y = L_0(y)$ for all $y \in S_B$, the claimed description follows.

Finally choose m so $p^{m-1}y \in A_{\max}$. Lemma 52.2 for $\nu = z, n = 1$ writes $p^m y = a + zw'$. Apply L_0, then write the resulting first term in S as a constant plus z times another series. Divide by p^m to obtain the decomposition. □

The preceding continuity argument also explains why the extended scalar-coordinate maps respect any exact relation over R involving coefficients in S_B. The argument does not claim that every continuous coordinate projection is multiplicative.

We record a useful cancellation. Since $z^e u = E - p$, write $E = v\xi$ with $v \in A^\times$. Then $z^e/p = u^{-1}(vX - 1)$ in $A_{\max}$. Multiplication by the reduction of $vX - 1$ is injective in $(A/(p, \xi))[X]$ (its leading coefficient is a unit). Repeating the proof of (52.1), with this polynomial in place of X, gives

$$\bigcap_{j \geq 0} z^j R = 0. \tag{52.2}$$

If $z \mid E^k a$ in R for $a \in A$, then $z \mid a$ in R since p is a unit there and $E \equiv p \pmod{z}$. Each $\lambda_j(a)$ is therefore divisible by z in R, hence in S: a series with nonzero constant term in Λ becomes a nonzero constant modulo zR, and that constant is invertible in R, which would contradict (52.2). Approximate a by finite coordinate sums and use closedness of zA to deduce $z \mid a$ in A; iteration proves the same for z^n.

Lemma 52.4. *If $a \in A$ and $\theta(\varphi^j a) = 0$ for every $j \geq 0$, then $\mu \mid a$. If the equalities hold for all $j \geq 1$, then $\tau \mid a$. The analogous assertions hold with A replaced by R.*

Proof. Each $\varphi^{-j}(\xi)$ is prime in A and the primes are distinct: the evaluations $\theta\varphi^k$ on them distinguish them, since $\theta(\varphi^m\xi) \neq 0$ for integer $m \neq 0$. Hence the product for $0 \leq j < N$ divides a for each N. Modulo p the sum of valuations of these terms is $(1-p^{-N})v(\bar{\mu})$. Letting N tend to infinity proves that $\bar{\mu}$ divides $\bar{a}$ in the valuation ring. Subtract a lift of this multiple of μ from a, divide by p, and repeat: all vanishing conditions are preserved because $\theta\varphi^j(\mu) = 0$. Completeness produces a quotient in A. Apply φ^{-1} for the assertion about τ. For $y \in R$, Lemma 52.2 applied successively to τ, ξ yields some m and $a \in A$ with $p^m y = a + \mu b$ for $b \in R$. In the first case the conditions pass to a. In the second case μ itself satisfies them and is divisible by τ, so the same argument applies. □

The one further Frobenius cancellation in Section 51.2 follows as well: if $(\varphi\tau)^h \mid \varphi b$ for $b \in R$, then every $\theta\varphi^j(b)$, $j \geq 1$, vanishes for $h > 0$; hence $\tau \mid b$ in R by the lemma. Cancel this factor and iterate to obtain $\tau^h \mid b$. Continuity of Frobenius and density of $A[1/p]$ also give the stated conjugation identity for the extended linear maps, by first checking it on A and then taking limits. In particular if $\tau z^n \mid E^h a$ in R with $a \in A$, evaluate at $\theta\varphi^j$ for $j \geq 1$ to deduce $\tau \mid a$ in A; cancel τ and apply the preceding z-cancellation to the quotient. This proves `tau_mul_z_pow_dvd` of `CrystallineLimit.lean`. Any divisor d of μ in A has the uniform division property required in Section 51.2. Indeed, suppose $d^h w \in A_{\max}$. Multiplication by $(\mu/d)^h$ gives $p^h X^h \tau^h w \in A_{\max}$. The binomial theorem gives $\xi = pu_0 + \tau^{p-1}$, with $u_0 \in A$; choose j with $(p-1)j \geq h$. Multiplying by the remaining power of τ yields

$$p^{h+j} X^h (X - u_0)^j w \in A_{\max}.$$

The multiplier $X^h(X-u_0)^j$ is monic modulo p, so by the proof of (52.1) it is saturated for p-divisibility. Clearing a denominator of $w \in R$ and applying saturation shows $p^{h+j}w \in A_{\max}$. A fixed matrix over R first requires only the maximum of its entry denominators. This supplies (d) in Section 51.2, without a rank-dependent determinant cost.

52.3 Fixed coefficients and period-function retractions

We use two classical Tate–Sen statements, in their usual precise forms. For finite $K/\mathbb{Q}_p$, $(C_p(i))^{G_K}$ is zero if $i \neq 0$, and $(C_p)^{G_K} = K$. See O. Brinon and B. Conrad, *CMI Summer School Notes on p-adic Hodge Theory*, preliminary version (2009), Theorem 2.2.7, which states both assertions. The Ax–Sen–Tate theorem for an algebraic extension $K_\infty/\mathbb{Q}_p$ gives $C_p^{G_{K_\infty}} = \widehat{K_\infty}$. For this generality see J. Ax, *Zeros of polynomials over local fields—the Galois action*, J. Algebra **15** (1970), 417–428, the theorem on p. 417, which gives the fixed field in the completion for an arbitrary algebraic extension of the local field. The latter form is used only in the fixed-tilt approximation below. The finite-field assertion gives a particularly useful filtered version. If $x \in B_{\mathrm{dR}}$ satisfies $gx - x \in t^n B_{\mathrm{dR}}^+$ for all $g \in G_K$, then

$$x - a \in t^n B_{\mathrm{dR}}^+ \qquad \text{for some } a \in K. \tag{52.3}$$

Indeed, if the lowest filtration index m of x is less than n, its leading term in $C_p(m)$ is invariant, forcing $m = 0$ and the leading term $a \in K$. The finite separable field K embeds canonically in the de Rham ring by lifting algebraic elements using Hensel's lemma. Subtract a and the same argument now forces the filtration index to be at least n. This proves (52.3) for arbitrary signed n; if $m \geq n$, take $a = 0$. In particular $(B_{\mathrm{dR}})^{G_K} = K$.

Let the subgroup of G_p relevant to the comparison be open and contain the fixer of Λ and $\theta(z)$; its fixed field K is a subfield of $K_z = \Lambda[1/p](\theta(z))$. The equality $\theta(S)[1/p] = K_z$ follows from the Eisenstein relation, or directly by division of a power series by the minimal polynomial of $\theta(z)$. We prove the exact filtered retraction required in the compact source, not just preservation modulo the maximal ideal.

Proposition 52.5 (Filtered retraction)**.** *Let $L : R \to R$ be an extended S-linear endomorphism, $a_1, \dots, a_d \in K$ and $y_j \in R$. If*

$$\sum_j a_j y_j \in t^n B_{\mathrm{dR}}^+,$$

then $\sum_j a_j L(y_j) \in t^n B_{\mathrm{dR}}^+$.

Proof. For each a_j choose $s_j \in S[1/p] \subset S_B$ whose theta value is a_j. We claim it can be replaced, modulo any desired power of ξB_{dR}^+, by an element of S_B with the same theta value. If P is the minimal polynomial of a_j, then $P'(a_j) \neq 0$. Choose $b \in S_B$ with theta image $P'(a_j)^{-1}$; initially choose a lift of this inverse using $\theta(S)[1/p]$. The Newton replacement $s \mapsto s - bP(s)$ improves the filtration precision by at least one, since $1 - bP'(s)$ has zero theta image and $P(s)$ has initial filtration at least one. Repetition gives the desired approximation. All replacements take place in S_B and require no inverse of an arbitrary period in R.

For $n > 0$ choose these approximants s_j modulo t^n. As R embeds into B_{dR}^+, both y_j and $L(y_j)$ have nonnegative filtration. Thus $\sum_j s_j y_j$ lies in $t^n B_{\mathrm{dR}}^+ \cap R = \xi^n R$. The equality holds since the X-adic order of each R-element is unchanged in its separated completion. As ξ is associated to E in R, E^n divides the sum in R. Extension and S_B-linearity give

$$E^n \mid L\Big(\sum_j s_j y_j\Big) = \sum_j s_j L(y_j).$$

Return to the a_j modulo t^n. If $n \leq 0$ there is nothing to prove. □

This is the criterion `Lift.Chart.IsChartOf.dvd_extendToBPlus` in `KummerChart.lean`. Taking all filtration degrees also shows that an exact zero relation is preserved. The same proof works after using the intersection of a finite collection of open subgroups and their fixed field.

Proposition 52.6 (Invariant ratios)**.** *Let $b, b' \in B = R[1/\mu]$, $b' \neq 0$, and suppose b/b' in the fraction field is fixed by the subgroup just considered. Then $b/b' \in \Lambda[1/p]$.*

Proof. By (52.3), $a = b/b' \in K$ inside B_{dR}. Clearing a common power of μ, find $y, y' \in R$ with $y = ay'$ in B_{dR}. At least one $L_j(y')$ is nonzero by Lemma 52.3. Use Proposition 52.5 for all degrees to obtain an exact equality $s = as'$ with $s, s' \in S_B$, $s' \neq 0$. Divide both by the largest common power of z in R. Such a largest power exists by (52.2). Quotients remain in S_B: multiplication by the nonzero scalar z commutes with all the maps in the bicommutant definition, and cancellation proves the assertion. Moreover neither can vanish modulo zR alone. For example, if $s' = zu$ and $s \notin zR$, write $s = c + zw$ with $c \in \Lambda[1/p]$ nonzero, using Lemma 52.3; the equality gives $c = z(au - w)$, so z is invertible in $R[a]$. But a is algebraic over $\mathbb{Q}_p$, hence integral over R, and integrality implies that z is already a unit in R (lying over a maximal ideal containing z would otherwise be impossible). This contradicts (52.2).

Write $s = c + zw$, $s' = c' + zw'$ with $c, c' \in \Lambda[1/p]$, $c' \neq 0$. If $c - ac' \neq 0$, then $z^{-1} = (c - ac')^{-1}(aw' - w)$ again belongs to the integral extension $R[a]$; the inverse of the nonzero algebraic scalar is in $\mathbb{Q}_p[a, \Lambda[1/p]]$ and integral over R. This is again impossible. Thus $a = c/c' \in \Lambda[1/p]$. □

Proposition 52.6 is precisely the scalar assertion `exists_witt_coeff_of_isChartOf`: clearing a power of p expresses $p^m b = wb'$ with $w \in \Lambda$. The proof just given uses the stronger filtered retraction, rather than the erroneous inference that every algebraic element of $R[1/\mu]$ belongs to the maximal unramified subfield solely because B is a crystalline period ring.

For use at all Frobenius divisors, let $\theta_a = \theta\varphi^{-a}$ and let $\mathcal{K}_a$ be the field generated inside B_{dR} by K, S_B and $\varphi^{-a}S$. If $r_j \in A$ and $v_j \in \mathcal{K}_a$, the condition

$$\sum_j v_j \varphi^{-a}(r_j) \in t^n B_{\mathrm{dR}}^+$$

is preserved by replacing r_j by $f(r_j)$ for any S-linear map $f : A \to S$. Here is the exact reduction. Clear a common denominator $d \in \mathcal{K}_a^\times$ so that each dv_j is a finite sum of products bcu, where $b \in K$, $c \in S_B$, $u \in \varphi^{-a}S$. Such products form the generating subalgebra, since S_B and $\varphi^{-a}S$ are subrings. The map $\varphi^{-a} f \varphi^a : A \to A$ is S-linear because $\varphi^a(S) \subset S$; it extends as above and fixes the factors from $\varphi^{-a}S$ when intertwining multiplication on the inputs of the form $\varphi^{-a}(sr_j)$. Apply Proposition 52.5 to the expanded sum, at degree $n + \mathrm{ord}_t\, d$, using this extended map; then divide by d. This is `sum_map_mem_of_valueField` of `DeRhamValues.lean` and proves the coefficient-retraction compatibility with the actual filtered forms. The same argument in degree $+\infty$ proves preservation of zero relations.

Fixed tilt coordinates. The following approximation is used in the determinant reduction of Section 51.3. Let H fix S pointwise and let $x \in \mathcal{O}^\flat_{C_p}$ be H-fixed. The finite fields $K_j = \theta_j(S)[1/p]$ are nested, since $\theta_j(s) = \theta_{j+1}(\varphi s)$, and their union is the algebraic field fixed by H: an automorphism fixing every K_j fixes all the compatible untilts of the chart parameter and the coefficients in Λ. Its action consequently fixes the tilt parameter, by the inverse-limit description of the tilt, and fixes S pointwise by continuity and the power-series presentation. For a fixed tilt component index N, take the untilt of $\varphi^{-N}[x]$ in C_p. It is H-fixed. By Ax–Sen–Tate it is approximable modulo $p\mathcal{O}_{C_p}$ by an element $a \in K_j$ for some sufficiently large j; a is integral because the target and the error are integral. Although a may be written $a = \theta_j(r)/p^m$ with $r \in S$, the divisibility of $\theta_j(r)$ by p^m in $\mathcal{O}_{C_p}$ implies it comes from an integral element of S: lift the quotient to $b \in A$ under θ_j, so $r - p^m b \in (\varphi^j E)A$, and apply λ_0. Then $a = \theta_j(\lambda_0 b)$. Nestedness gives the same conclusion at all larger j. Thus the N-th component of x is congruent modulo p to the j-th component of the reduction of an element of S, exactly `eventually_exists_coeff_eq_of_fixed`. This also verifies why a fixed-element statement over C_p alone is insufficient: the last integral retraction is essential.

52.4 The first radius lemma: constant times a unit

We now prove the factorization half of `radius_lemma` in `Lib/PadicHodge/Comparison/CrystallineLimit.lean`. The other half, namely the span with the rank-free loss $p^{4\delta}$, was proved in the earlier discussion of the radius lemma (see (50.10)).

Choose $t_0 \in G_p$ fixing Λ and $\theta(z)$ such that $t_0(z) = [\epsilon]z$. For every $y \in S_B$, we have $z \mid t_0(y) - y$ by the decomposition in Lemma 52.3. Moreover $\theta\varphi^j(y) \in K_z$ for all $j \geq 0$: use Frobenius stability, apply Lemma 52.2 with $\nu = E$ to an integral multiple of $\varphi^j(y)$, and apply $\widehat{\lambda}_0$ before theta. Thus all $\theta\varphi^j$ of the difference vanish. Since $\theta\varphi^j(z) \neq 0$, division by z followed by Lemma 52.4 gives

$$\tau z \mid t_0(y) - y \quad \text{in } R. \tag{52.4}$$

For $n \geq 1$ one has a useful integral identity

$$\tau E(p^{v_p(n)} + \beta z) = \alpha([\epsilon]^n - 1), \qquad \alpha, \beta \in A. \tag{52.5}$$

To prove it, put $q = [\epsilon] = 1 + \mu$ and consider the quotient $A/(z)$. There $E = p$ and $\mu = pc$, with c a unit multiple of τ in this quotient, hence topologically nonunit. By the binomial

theorem

$$\frac{(1+pc)^p-1}{pc} = p\left(1+\sum_{j=2}^{p-1}\frac{1}{p}\binom{p}{j}(pc)^{j-1}+p^{p-2}c^{p-1}\right)$$

as an identity of polynomials. The parenthesis is a unit: for $p>2$ all nonconstant terms are divisible by p; for $p=2$ it is $1+c$, with c topologically nilpotent modulo the maximal ideal of $A/(z)$. Iterating and using that the geometric sum for an exponent coprime to p has unit reduction at $q=1$ gives

$$1+q+\cdots+q^{n-1} = p^{v_p(n)}u_n \quad \text{in } A/(z), \qquad u_n \in (A/(z))^\times.$$

Multiply by u_n^{-1}, lift back to A, and multiply by μ. The association $E\tau \sim \mu$ absorbs a unit into α and gives (52.5). No division by z or p has taken place in the integral ring itself. This is `exists_tau_mul_E_eq_mul_epsTilt_pow_sub` of `PeriodMatrixBasis.lean`.

Theorem 52.7 (Radius factorization)**.** *Let $Z, Z' \in M_d(R)$ with entries of Z in S_B. Suppose $Z'Z = E^hI$, and $t_0(Z) = ZG$ for $G \in M_d(A)$. If $3e \le p^l$, then for every integer $b \ge 0$,*

$$\varphi^l Z = (\varphi^l E)^b CV, \qquad C \in \mathrm{GL}_d(\Lambda[1/p]), \quad V \in \mathrm{GL}_d(A_{\max}).$$

In particular C is fixed by every automorphism fixing Λ. The matrix V is the value of an invertible formal power-series matrix over S at z^{p^l}/p^2.

Proof. Apply the distinguished extended coordinate to every entry of Z'; it commutes with Z, and therefore gives another matrix over S_B with the same left inverse identity. Divide this matrix by p^h, and approximate it modulo z by a matrix B having entries in $\Lambda[1/p]$, using Lemma 52.3. Then

$$BZ = I + zW_1, \qquad W_1 \in M_d(S_B).$$

Indeed $E^h/p^h \equiv 1 \pmod z$, and the error in the approximation is divisible by z in S_B.

By (52.4), $t_0(Z)-Z$ is entrywise divisible by τz in R. Left multiplication by Z' gives $\tau z \mid E^h(G-I)$. The scalar cancellation following Lemma 52.4 proves $G-I \in \tau z M_d(A)$. We start an induction with

$$q_nBZ = A_n + z^nW_n, \qquad q_n = p^{n-1+v_p((n-1)!)}, \qquad A_n \in M_d(A), \quad W_n \in M_d(S_B);$$

at $n=1$ take $A_1 = I$. As t_0 fixes B, one has $t_0(BZ) = BZG$. Subtracting after applying t_0 to the displayed identity gives

$$A_nG - t_0(A_n) = z^n([\epsilon]^n t_0(W_n) - W_nG) = z^n(([\epsilon]^n-1)W_n + \tau z D_n)$$

for some $D_n \in M_d(R)$, by (52.4) and the just established divisibility of $G-I$. Multiply by the scalar α in (52.5), write $E = p+zr$ ($r \in S$), and obtain

$$\alpha(A_nG - t_0(A_n)) = \tau z^n\big(p^{1+v_p(n)}W_n + z((p^{v_p(n)}r+\beta E)W_n + \alpha D_n)\big).$$

The left side is integral. Cancellation of τz^n in A is permissible by the earlier contraction result. Apply $\widehat{\lambda}_0$ to the quotient and use its S_B-linearity. There are $s_n \in M_d(S)$ and $W_{n+1} \in M_d(S_B)$ with

$$p^{1+v_p(n)}W_n = s_n + zW_{n+1}.$$

Multiplying the induction identity by $p^{1+v_p(n)}$ and adding z^ns_n to its integral part proves the induction at $n+1$.

Put $x = z^{p^l}/p^2$. The inequality on l implies $x \in pA_{\max}$, because $z^e = pu^{-1}(vX - 1)$ with u, v integral units. Set the formal series matrix

$$V_0(X_0) = I + \sum_{n \geq 1} X_0^n p^{n - v_p(n!)} \varphi^l(s_n) \qquad \in M_d(S[[X_0]]).$$

Here $n - v_p(n!) \geq 0$, by the usual factorial valuation formula, and V_0 has constant matrix I, so it is invertible. Evaluation at $X_0 = x$ converges in $A_{\max}$ and gives an invertible matrix. Applying φ^l to the finite induction identities shows

$$\varphi^l(BZ) \equiv V_0(x) \pmod{x^n R} \qquad \text{for every } n.$$

One should not claim that the unknown period remainders have bounded p-denominators. Rather, x is a nonzero scalar multiple of a power of z in R, so (52.2) implies equality. As $\varphi^l(B)$ now has invertible product with $\varphi^l(Z)$, it has nonzero determinant in $\Lambda[1/p]$ and is invertible there. This gives the assertion for $b = 0$.

Finally $\varphi^l E = p(1 + px\varphi^l r)$. The factor in parentheses is the evaluation of a scalar formal series with constant term 1. Absorb its inverse b-th power into V_0 and absorb p^{-b} into $\varphi^l(B)^{-1}$. This proves the stated formula. □

The proof uses the structural cancellation and retraction established above, together with the exact Galois congruence in the hypotheses. This closes the first half of the radius lemma independent of Kisin's slope construction. Combining Theorem 52.7 with the previously established $p^{4\delta}$ span identity gives the full statement `radius_lemma`: indeed, the approximations to the evaluated power series and its inverse used there are valid because $z^{p^l}/p^2 \in pA_{\max}$ and all their coefficients are integral.

52.5 The actual Kummer charts and their congruences

Let $K/\mathbb{Q}_p$ be finite, of residue degree f and ramification degree e, with maximal unramified subfield $K_0 = \Lambda[1/p]$. Choose a uniformizer π and compatible p-power roots of π in $\overline{\mathbb{Q}}_p$ and take for z their Teichmüller vector. Its theta value is π. If $F(X) = X^e + p(c_0 + c_1 X + \cdots + c_{e-1}X^{e-1})$ is the Eisenstein minimal polynomial, then c_0 is a unit of Λ. Set

$$H(X) = c_0 + c_1 X + \cdots + c_{e-1}X^{e-1}, \qquad E = H(z)^{-1}F(z) = p + z^e H(z)^{-1}.$$

Then $S = \Lambda[[z]] \subset A$ as above, and E is distinguished and generates $\ker \theta$. The group fixing $\Lambda, \theta(z)$ is exactly G_K. Every residually trivial automorphism of the unramified coefficient ring actually fixes it pointwise by the Teichmüller expansion. For every integer $m \geq 0$ there is $g \in G_K$ with $g(z) = [\epsilon]^m z$: at finite level $X^{p^j} - \pi$ is Eisenstein over K, hence the assignment $\pi^{1/p^j} \mapsto \zeta_{p^j}^m \pi^{1/p^j}$ extends to an embedding of $K(\pi^{1/p^j})$; these embeddings are compatible and extend from their union to an automorphism of the algebraic closure. For $e = 1$ choose $\pi = -p$, so $E = z + p$. This proves the existence of the chart in `Lift.exists_kummer_chart` for an open image: take K to be its fixed field, take the inverse image of G_K as chart subgroup (which equals the whole given group here), and use the equality $[K : \mathbb{Q}_p] = ef$. The usual unramified quotient of G_K supplies an element inducing p^f-Frobenius on the algebraic residue field.

Here are the congruences needed with this chart. Write $E_a = \varphi^a E$ and $\Delta = z\tau$. Each E_a is prime, generates $\ker \theta_a$ and is distinct from every other, as is seen by applying θ_a to E_b. Moreover $\theta_a(\Delta) \neq 0$ (the chosen p-power roots and $\zeta_{p^{a+1}} - 1$ are nonzero), and $\Delta \mid \varphi^a \Delta$, by factoring the difference of powers for the cyclotomic factor. Every $g \in G_K$ preserves the

ideal (z) of A: the reductions of its Teichmüller generator and its Galois conjugate have equal valuation, so they differ by a unit, whose Teichmüller lift gives the corresponding integral unit. It also preserves the ideal (τ), by Lemma 52.4 in both directions. Consequently it preserves $A_{\mathrm{loc}} = A[E_a^{-1} : a \geq 0]$ (the E_a are associated to their conjugates since g commutes with Frobenius and preserves $\ker\theta$).

The argument of (52.4) is valid for any $g \in G_K$, with the stronger conclusion

$$E\Delta \mid gs - s \quad (s \in S), \qquad \Delta \mid gy - y \quad (y \in S_B) \tag{52.6}$$

in A and R respectively. Indeed $gy - y$ is divisible by z in R from its constant approximation, its images under all $\theta\varphi^j$ vanish for $j \geq 0$, and none of those images of z vanish. The μ assertion of Lemma 52.4 thus applies after division by z. For $s \in S$ the difference is integral; first cancel μ in A by the $heta\varphi^j$ criterion, then cancel z as in Section 52.2. This is the first assertion in (52.6) because $E\tau \sim \mu$. In particular, writing $gE = \omega E$ then gives $\omega - 1 \in \Delta A$. These are exactly `zTau_mul_E_dvd_smul_sub` and `zTau_dvd_sub_one_of_mul_eq_smul_E`. All the scalar chart hypotheses preceding Lemma 51.8 are thus justified. The different bound and derivative identities of that lemma then apply with the actual F, H just constructed.

52.6 Construction and identification of the crystalline frame

We spell out how the matrix hypotheses of the two preceding chapters arise from the representation. We work with a crystalline free $\mathbb{Z}_p$-lattice over the chart subgroup; this is precisely the local input after restriction in the source. With coefficients in a finite free integral coefficient ring, regard this as a $\mathbb{Z}_p$-lattice and apply the same construction; the letter projectors are addressed below. We use the crystalline property in its period-module sense: invariant columns form a basis after base change to B. This property is stable under restriction, tensor products and duals, as follows by multiplying the invariant bases. In the potentially crystalline case it holds after the prescribed open restriction.

For clarity, for any finite collection of invariant columns spanning B_{dR}^d, their span of invariants is a K-vector space of dimension d: a vector of coefficients expressing another invariant column is itself invariant by the invertible matrix, and (52.3) applies to each entry. Filter this space by the intersection with $t^j(B_{\mathrm{dR}}^+)^d$. Choose a K-basis adapted to that filtration and let its individual degrees be k_j. Dividing the invariant columns by t^{k_j} yields an invertible matrix over B_{dR}^+. Here invertibility modulo t is essential and can be proved directly: normalize a hypothetical relation among the leading columns to have one coefficient 1 and choose minimal support. Galois transforms this relation, after normalizing by the cyclotomic twist of the chosen coordinate; minimal support makes the relation fixed. Tate–Sen says that coefficients between different weights vanish and those within one weight lie in K. This contradicts independence on the corresponding graded invariant space. Thus the leading matrix has independent columns. The same leading-basis argument after any finite open restriction (the original invariant basis remains a basis there) shows that the eventual, near-identity Hodge–Tate multiplicities over $\overline{\mathbb{Q}}_p$ are these same dimensions; any eventual equivariant form is a combination of the leading basis vectors, whose coefficients in the corresponding weight are fixed in a finite open subgroup by Tate–Sen, hence algebraic. This also proves that the degrees are the Hodge–Tate weights with the corresponding sign convention; the leading columns furnish a Hodge–Tate basis, with dimensions of the different weights read off by the same argument. For the dual rows the signs are reversed. In particular in the source's jump convention the condition $-a \leq k_j \leq b$ is the two-sided period bound on the invariant basis and its inverse.

Let $M \in \mathrm{GL}_d(B)$ be an invariant matrix, meaning

$$\rho(g)g(M) = M$$

on the subgroup, with ρ the integral representation matrix. Every nonnegative Frobenius iterate of M is also invariant. The preceding argument applied to each iterate shows that its entries have t-order at least $-a$ and its inverse has order at least $-b$. Indeed any two such invariant bases differ by a matrix in K, which consists of order-zero scalars.

We need the integral consequence for the positive ring. If $x \in B = R[1/\mu]$ and all $\varphi^j(x)$, $j \geq 0$, have nonnegative order in B_{dR}, then $x \in R$. Write $x = y/\mu^n$. Because $\varphi^j\mu$ has order 1 for $j \geq 0$, all $\varphi^j(y)$ have order at least n. If $n > 0$, Lemma 52.4 shows $\mu \mid y$ in R, and division reduces n by one while preserving the required conditions. Induction proves the assertion. Applying this to $(E\tau)^a M$ and $(E\tau)^b M^{-1}$ gives the integral matrices

$$U = (E\tau)^a M \in M_d(R), \qquad W = (E\tau)^b M^{-1} \in M_d(R), \qquad UW = (E\tau)^{a+b} I.$$

This is the positive-ring division used in `CrystallinePeriodMatrix.lean`, not an appeal to rational admissibility alone. It has the same statement for every Frobenius divisor because the evaluations in Lemma 52.4 are simultaneous.

Set $v = \mu/(E\tau) \in A^\times$. The entries of $M^{-1}\varphi M$ and of its inverse are invariant in B, so belong to $\Lambda[1/p]$ by Proposition 52.6. Thus

$$\varphi U = v^a U F, qquad F = (\varphi E)^a M^{-1}\varphi M.$$

The entries of F, F^{-1} lie in S_B: we have $\varphi E \in S$, and φE is a unit in R (its association to p was proved in Section 52.1); cancellation of that nonzero scalar proves its inverse belongs to S_B. Also $E\tau$ is associated to its Galois conjugate by the earlier congruences, so the invariant-column equation gives an integral matrix ρ_g with $\rho_g\, g(U) = U$ whenever g fixes $\Lambda, \theta(z)$.

Finally let $f : A \to A$ be S-linear, and suppose $Uy \in E^{a+b}R^d$, with $y \in R^d$. Choose the adapted invariant de Rham basis above, so $MP = H\,\mathrm{diag}(t^{k_j})$ for $P \in \mathrm{GL}_d(K)$ and $H \in \mathrm{GL}_d(B_{\mathrm{dR}}^+)$. Then $Uy \in E^{a+b}R^d$ if and only if the j-th entry of $P^{-1}y$ belongs to $t^{b-k_j}B_{\mathrm{dR}}^+$ for all j. The equivalence uses the exact equality $E^{a+b}R^d = R^d \cap t^{a+b}(B_{\mathrm{dR}}^+)^d$. Proposition 52.5 applied with the rows of P^{-1} proves the same inequalities for the extended map applied to y. These four statements give exactly `IsPeriodMatrix` for the parameter pair $(a+b, v^a)$; in particular they close `IsChartOf.exists_isPeriodMatrix`.

We now use the previously proved Theorem 51.4, Proposition 51.6 and Corollary 51.7. The structural assumptions of those results have been justified in Sections 52.1–52.3: the two arithmetic chart assumptions for Proposition 51.6 are the fixed tilt approximation and filtered coordinate reduction modulo p, established there. For the second of these, apply λ_i to $a - pb = E_j c$ after lifting a quotient under θ_j; it gives the same relation for the coordinate. This proves the required finite-component congruence.

For simplicity of notation take a single bound h on the absolute values of the jumps, so $a = b = h$. The period-matrix height is then $2h$, not h. We obtain integral Q, L, matrices Z, Y over R with Z over S_B, and $B_1 \in M_d(S)$, such that

$$QL = \tau^{2h}I, \quad UZ = E^{2h}Q, \quad U = QY, \quad YZ = E^{2h}I, \quad \varphi Q = v^h Q B_1.$$

All matrices displayed as nonsingular earlier remain so in the domains used here. The exact Galois frame identity required by `IsPeriodMatrix.exists_frame` follows from these equations, not an independent comparison: apply g to $UZ = E^{2h}Q$, multiply by ρ_g, and divide by E^{2h} using $gE = u_g E$ with u_g a unit. One obtains an integral solution of $Ug(Z) = E^{2h}u_g^{2h}\rho_g g(Q)$; Corollary 51.7 implies that the latter integral coefficient matrix belongs to $QM_d(A)$. Cancelling gives $g(Z) = ZG_g$ for $G_g \in M_d(A)$.

Theorem 52.8 (Exact intrinsic frame)**.** *Let T be the given lattice, and let n_j be the columns of $\tau^{-h}Q$ in the chosen basis of T. The intrinsic positive module*

$$M_T = \{x \in A[1/\tau] \otimes T : \ \mu_0(\varphi^m x) \in B^+_{\mathrm{dR}} \text{ for every equivariant form } \mu_0 \text{ and every } m \geq 1\}$$

is free on the n_j over A. The value-field points of $\varphi^l M_T$ are exactly the S-span of $\varphi^l n_j$. Moreover the scalar inclusions, crystalline-form divisibility test and rank-free radius span in `FrameTheorem.lean` *hold.*

Proof. The identity $UZ = E^{2h}Q$ reads $E^h n = MZ$, for n the matrix with columns n_j. For an invariant de Rham row μ_0, all the entries of $\mu_0(\varphi^m M)$ are fixed by the chart subgroup and hence belong to K, by (52.3). Therefore

$$E_m^h \mu_0(\varphi^m n) = \mu_0(\varphi^m M)\, \varphi^m Z. \tag{52.7}$$

When $m \geq 1$, E_m is invertible in B^+_{dR} and the right side has nonnegative order. This proves inclusion of the frame in M_T. When $m = 0$, or for $\varphi^l n$ with arbitrary l, all its values belong to the field $\mathcal{K}_0$, since $\varphi^m Z$ has entries in S_B and $E_m \in S$.

Conversely, if $x \in M_T$, then by $QL = \tau^{2h} I$ there are $N \geq 0$ and $t \in A^d$ with $\tau^N x = nt$. If $N > 0$, fix $m \geq 1$. Choose de Rham equivariant rows μ_j dual to the columns of $\varphi^m M$ at this one m; these exist by its invariant invertibility. Because $\varphi^m \tau$ has positive filtration, every $\mu_j(\varphi^m(nt))$ has filtration at least N. Multiplying the matrix identity (52.7) by $\varphi^m Y$ shows that each coordinate of $\varphi^m(t)$ has filtration at least N (the leftover scalar E_m is a unit). Thus $\theta\varphi^m(t_j) = 0$ for every $m \geq 1$, so $t_j \in \tau A$ by Lemma 52.4. Cancelling reduces N by one. Iteration gives the desired A-span; uniqueness follows from nonsingularity of Q. The inclusions $\tau^h M_T \subset A \otimes T$ and $\tau^h(A \otimes T) \subset M_T$ follow directly from $n = \tau^{-h}Q$ and $QL = \tau^{2h} I$ respectively.

Suppose $x = \varphi^l n\, t$ with $t \in A^d$ and all its invariant row values in $\mathcal{K}_0$. For every equivariant row, apply the exact filtered retraction of Section 52.3 to the relation between $\mu_0(x)$ and $\sum_j t_j \mu_0(\varphi^l n_j)$, including the first term with scalar coefficient 1. It follows that replacing t_j by $\lambda_0(t_j)$ does not change any row value. The invariant rows span the dual over B_{dR}, so nonsingularity implies $t \in S^d$. This proves the assertion about value-field points. The same proof gives a retraction from their localized A-span to the localized S-module: apply the map to scalar coefficients, after clearing the finite product of E_j denominators (which lie in S). It respects all evaluations, letter projections and relations because those are linear identities over the value field, as in the filtered retraction proof. In particular $N_T = M_T$'s value-field points is the exact source module `kisinModule`, not a chosen auxiliary lattice.

If μ_0 is crystalline, then $\mu_0(M)$ belongs entrywise to $B^{G_K} \subset \Lambda[1/p]$ by Proposition 52.6. Hence (52.7) at $m = 0$ gives $E^h \mu_0(y) \in S_B$ for $y \in N_T$. The invariant crystalline rows dual to M may be used to test divisibility for $x = nt$ with $t \in A^d$. If $d_0 \in A$ has the cancellation property $d_0 \mid E^k a$ in $R \Rightarrow d_0 \mid a$ in A, and $d_0 \mid E^{2h}\mu_0(x)$ in R for every crystalline row, then $d_0 \mid E^h Z t$; multiply by Y to get $d_0 \mid E^{3h} t$ in R. Thus $t \in d_0 A^d$ and $x \in d_0 M_T$. This is precisely the divisibility test used for $d_0 = z$ and $d_0 = \tau z$; those cancellation properties were proved earlier. The Frobenius relation for Q, with $\varphi\tau = \mu = vE\tau$, gives $E^h\varphi(n) = nB_1$ integrally over S.

Finally apply Theorem 52.7 to Z, Y (height $2h$) with exponent $b = h$, so $\varphi^l Z = E_l^h CV$. Then (52.7) gives

$$\mu_0(\varphi^l n\, t) = \mu_0(\varphi^l M) CVt \qquad (t \in S^d).$$

The constant coefficient row on the right before Vt is in K and fixed by G_K. Choose the spanning vectors of the radius lemma with Vt constant modulo $E^A R^d$. Then every row value

is Galois-invariant modulo $t^A B_{\mathrm{dR}}^+$, and the S-span of those chosen vectors contains $p^{4\delta} S^d$ for $eA \le p^l \delta$. This proves the radius span on the actual value-field points, with the required rank-independent constant. □

Each part of `frame_theorem` is thus proved from the basic crystalline invariant-basis property. In particular its relation assertion for value-field points of the saturated ambient space follows by applying the exact filtered retraction to every row value of a relation and using the invariant dual basis to separate. Coefficient rings and letter decompositions cause no change: the finite-order action on the coefficients is an integral sum of the projectors described in Section 50.6. The projector onto one letter is a polynomial in coefficient multiplication, and the S-linear retraction commutes with every coefficient endomorphism because finite scalar multiplication commutes with passing to its coefficients. Thus the frame, its evaluations and the retraction restrict to the prescribed letter. This is the exact content required by `coordRetract_mem_withExponents` and `valueAt_coordRetract_sub_mem_hodgeFil`.

52.7 Galois differences and the multiplier ideals

Here $\mathcal{M} = M_T[E_j^{-1} : j \ge 0]$ and $\mathcal{N} = N_T[E_j^{-1} : j \ge 0]$. As in Chapter 51, an exponent condition at a is a signed lower bound on the order of a form after evaluating through φ^{-a}. Scalar division by E_j reduces that bound by one exactly at $a = j$, and leaves the others unchanged. Division by $\Delta = z\tau$ leaves every bound unchanged because $\theta_a(\Delta) \neq 0$.

We need a slightly stronger observation about value fields than their definition. If $x \in \mathcal{K}_a \cap B_{\mathrm{dR}}^+$, its residue belongs to $\theta_a(S)[1/p]$. Indeed, choose m so the residue of $p^m x$ belongs to $\mathcal{O}_{C_p}$ and lift it to $r \in A$ under θ_a. The relation $p^m x - \varphi^{-a} r \in tB_{\mathrm{dR}}^+$ has coefficients in $\mathcal{K}_a$. Apply the filtered retraction with $f = \lambda_0$ and use $f(1) = 1$ to obtain $p^m x - \varphi^{-a}(\lambda_0 r) \in tB_{\mathrm{dR}}^+$. Taking residues proves the claim. Consequently, if g, t_j are elements of G_K and E_a divides $(g-1)\sum_j t_j s$ for every $s \in S$, the residue of $(g-1)\sum_j t_j x$ is zero for each such integral x: simply replace the residue by $p^{-m}\theta_a(s)$ and use the condition on s. This proves `smul_sub_mem_hodgeFil_of_valueField` in the source. Together with (52.3) this verifies the filtered fixed-scalar compatibility at all the degrees required in Section 51.6.

For $x \in N_T$ and $g \in G_K$, we have

$$gx - x \in \Delta M_T. \tag{52.8}$$

Here gx is in M_T directly from the definition and Galois equivariance of the forms. Write $E^h \mu_0(x) = s \in S_B$ for any crystalline row. By (52.6),

$$E^h \mu_0(gx - x) = \omega^{-h} gs - s \in \Delta R, \qquad gE = \omega E, \quad \omega - 1 \in \Delta A.$$

In particular $E^{2h}\mu_0(gx - x) \in \Delta R$. The divisibility test with $d_0 = \Delta$ in Theorem 52.8 proves (52.8). The conclusion extends to $x \in \mathcal{N}$: choose a denominator d that is a finite product of E_j and $dx \in N_T$. Then $g(d) = d\omega_d$ with $\omega_d - 1 \in \Delta A$, by taking Frobenius iterates of the scalar congruence and using $\Delta \mid \varphi^j \Delta$. Subtract the equation for dx and divide by $d\omega_d$. We obtain $(gx - x)/\Delta \in \mathcal{M}$, with unchanged exponents whenever $gx - x$ satisfies them. This proves `smul_sub_mem_zTau_kisinLocAinf` and `smul_sub_mem_zTauUnit_smul_of_deep`. These assertions are not statements about multiplication by a unit of A; they use the actual divisibility test for the intrinsic frame.

Write $J_l(i; r)$ for the multiplier ideal in Section 49.6, with modulus r. By its definition, $f \in J_l(i; r)$ means that there is $t \in A$ such that for every constant c_i in the letter i the element $(f - rt)c_i$ is the projection of an extension point of standard exponents at level l. If $f \in S$,

apply the coefficient retraction to both the projection identity and the exponent conditions. It gives $t \in S$ and a localized S-point x with

$$\pi(x) = (f - rt)c_i. \tag{52.9}$$

This argument proves `exists_lift_of_mem_liftIdeal` directly; naturality with respect to the projection follows by expressing x with scalar coefficients and applying the map to those coefficients.

Proposition 52.9 (Exact derivation in the multiplier ideal)**.** *Suppose $f \in S \cap J_1(i; r)$ and $g \in G_K$, with $\alpha, \beta, s \in A$ such that*

$$\Delta E(s + \beta f) = \alpha(gf - f).$$

Then $s \in J_1(i; r)$.

Proof. Choose t, x as in (52.9). The point x has standard level-one exponents, in particular nonnegative order at $a = 0$. Its values there are in $\mathcal{K}_0$; as $g \in G_K$, their differences under g have order at least one, by the residue assertion just proved. At every other divisor the differences retain the prescribed exponents by Galois equivariance. Thus $(gx - x)/(\Delta E)$ belongs to $\mathcal{M}$ by (52.8), and division has precisely restored the required standard exponents at $a = 0$. The point

$$y = \frac{\alpha(gx - x)}{\Delta E} - \beta x$$

therefore belongs to the same standard exponent lattice. The projection is

$$\pi(y) = \left(s - r\left(\frac{\alpha(gt - t)}{\Delta E} - \beta t\right)\right) c_i.$$

The parenthesis is integral by (52.6) for $t \in S$. This gives the defining membership of s in $J_1(i; r)$, uniformly in c_i. □

This is `mem_liftIdeal_of_derivation`. Its proof completes the formerly conditional ideal operations in the preceding chapters: the scalar depth identities and factorial iteration apply to $J_l(i; r)$ itself. In particular the cokernel estimates of Proposition 49.6 now have the requisite integral compatibility, and the proof of the unramified cokernel bound is independent of a rank estimate.

52.8 The tame chart and the polynomial recurrence

Suppose the map of the given Galois group to G_p is surjective, $e \mid p^f - 1$ and $p \nmid f$. Let K_0 be the unramified degree-f field, choose π with $\pi^e = p$, and set $K = K_0(\pi)$. This is a Galois extension of $\mathbb{Q}_p$: it is the splitting field of $X^e - p$ together with K_0, which contains all the e-th roots of unity, and its degree is ef prime to p. The inverse image of G_K is a normal subgroup of index ef in the given group. Use compatible roots of π as in Section 52.5, now with $E = p - z^e$. Let $\zeta \in \Lambda$ be a primitive e-th root with primitive reduction (the Teichmüller root in $W(\mathbb{F}_{p^f})$). There exists an inertia element σ with

$$\sigma(z) = \zeta z.$$

To see this with the exact compatible roots, work first over the maximal unramified algebraic extension of $\mathbb{Q}_p$, whose valuation group is $\mathbb{Z}$. The polynomial $X^{ep^j} - p$ remains Eisenstein there. Prescribe $\sigma(\pi^{1/p^j}) = \zeta^{p^{-j}} \pi^{1/p^j}$ with the exponent interpreted in $\mathbb{Z}/e$, and extend the

compatible embeddings of these fields to the algebraic closure. The resulting automorphism fixes the maximal unramified field, hence lies in inertia; at Witt level the formula for $\sigma(z)$ follows using the compatible residue coordinates of the root of unity. Since the original map is onto, choose a lift of this element. Its powers satisfy $\sigma^k(z) = \zeta^k z$, fix Λ, and its chart subgroup fixes π. This constructs `Lift.exists_tameChart` and all the extra conditions in the definition of `TameChart`, including the condition on the Weil pair: inertia corresponds to Frobenius index zero.

Let Y be a potentially crystalline lattice, crystalline on the chart subgroup. Let $\beta_{\sigma,0}$ be the linearized operator on $B \otimes Y$ used in the source: it agrees with the natural diagonal action of σ on the invariant subspace of vectors fixed near the identity, and extends B-linearly. This agreement follows because an invariant period scalar is fixed by inertia: after restricting to an open group and choosing its chart, Proposition 52.6 places the scalar in a finite unramified field. Hence a change of invariant basis gives the same operator. We shall prove the integral statement when $Q(X) \in \mathbb{Z}_p[X]$ annihilates $\beta_{\sigma,0}$; in the eventual application $Q = q(X)(X-1)$.

Lemma 52.10 (Polynomial lowering modulo z)**.** *For the intrinsic module N_T of a crystalline lattice on this chart, if $Q(\beta_{\sigma,0}) = 0$, then*

$$\sum_k Q_k \sigma^k x \in z M_T qquad (x \in N_T).$$

Proof. The normality of the chart subgroup shows that σ carries invariant forms into invariant forms, so preserves M_T by definition. For a crystalline form μ_0 put $\mu_k = \sigma^{-k} \circ \mu_0 \circ \rho(\sigma^k)$ on the original lattice, with scalar coefficients continued to B. These are again crystalline forms. They satisfy $\mu_0(\sigma^k x) = \sigma^k(\mu_k(x))$. Moreover $\sum_k Q_k \mu_k = 0$: test the equality on a basis of invariant columns of $B \otimes T$ fixed by the chart subgroup. The value of each invariant form there is a scalar fixed by that subgroup, hence fixed by σ. The equality now follows from $Q(\beta_{\sigma,0}) = 0$ on the invariant basis, and extends by B-linearity.

By Theorem 52.8, $s_k = E^h \mu_k(x) \in S_B$. Since σ fixes Λ and preserves (z), Lemma 52.3 gives $\sigma^k s_k - s_k \in zR$ (here one needs only z-divisibility, not the stronger Δ-divisibility for the subgroup). Also $\sigma(E) = E$ exactly, because $E = p - z^e$ and $\zeta^e = 1$. Thus

$$E^h \mu_0\Big(\sum_k Q_k \sigma^k x\Big) = \sum_k Q_k \sigma^k s_k \equiv \sum_k Q_k s_k = 0 \pmod{zR}.$$

The divisibility test in Theorem 52.8, with $d_0 = z$, proves the lemma. □

This is `exists_z_smul_eq_sum_tameGen` in `SparseThickLetters.lean`. It is an actual assertion about the A-span of the intrinsic Kisin frame, not a computation merely in the reduction of the rational invariant space. If q annihilates the operator on the kernel Y of an extension of the trivial representation, then $Q = q(X)(X-1)$ annihilates the operator on the extension: the image of $(\beta - 1)$ lies in the kernel, and the operators commute. This uses that the operator on the trivial quotient is the identity and is natural for the extension, both of which follow by using the invariant basis. In particular $Q(1) = 0$ and $\deg Q = \deg q + 1$ when q is monic.

Theorem 52.11 (Tame cokernel bound)**.** *Let all side jumps be at most n, suppose q is monic and kills the above operator on the kernel, and let*

$$e \le p^m, qquad m(\deg q + 1) \le \kappa, \qquad n + \kappa + 2 < p.$$

Then p^κ annihilates the exact level-zero cokernel at each letter.

Proof. Use the multiplier ideal with modulus $r = p^{\kappa+1}$. The cokernel calculus of Proposition 49.6, with its compatibility now established by Proposition 52.9, gives a nonnegative integer ν such that

$$z^{\nu} \in J_0(i; r), \qquad (p-1)\nu \le e(n+\kappa+1).$$

The strict bound on $n+\kappa+2$ implies $\nu < e$. Put $Q = q(X)(X-1)$ and

$$d_j = m\#\{k : j < k < e,\ \bar{Q}(\bar{\zeta}^k) = 0\} \qquad (0 \le j < e).$$

We claim by descending induction from ν that $p^{d_j} z^j \in J_0(i; r)$. At $j = \nu$ this follows from $z^{\nu} \in J_0$. Suppose $j > 0$. If $\bar{Q}(\bar{\zeta}^j) = 0$, the Frobenius transport and factorial descent of Proposition 49.6 give

$$p^{d_j + 1 + v_p(j)} z^{j-1} \in J_0(i; r).$$

Indeed carry level zero to level one, where the z-exponent is pj, decrease it to $p(j-1)$ at cost $v_p((pj)!) - v_p((p(j-1))!) = 1 + v_p(j)$, and carry back. As $0 < j < e \le p^m$, this cost is at most m; multiplication by the remaining power of p gives exponent d_{j-1}.

If $\bar{Q}(\bar{\zeta}^j) \neq 0$, take a localized S-lift as in (52.9) of $(p^{d_j} z^j - rt)c_i$ with $t \in S$ and standard level-zero exponents. Write $t = w + zt_1$, $w \in \Lambda$. The operator $Q(\sigma)$ kills w because $Q(1) = 0$, and $\sigma^k(z) = \zeta^k z$. Thus

$$Q(\sigma)t = z \sum_k Q_k \zeta^k \sigma^k(t_1), \qquad Q(\sigma)(p^{d_j} z^j) = p^{d_j} z^j Q(\zeta^j).$$

The lemma applies to the lift after clearing its finite product of E_a denominators: in this tame chart every σ fixes all the E_a exactly. Consequently $Q(\sigma)x/z \in \mathcal{M}$. Its standard exponents are preserved because every summand $\sigma^k x$ has them (Galois stability of the filtration and normality), and z has order zero under each evaluation. The letter is preserved since σ fixes Λ. Its projection is

$$(p^{d_j} z^{j-1} Q(\zeta^j) - r \sum_k Q_k \zeta^k \sigma^k(t_1))c_i.$$

All scalar error terms displayed are integral, and $Q(\zeta^j)$ is an integral unit, by the nonvanishing condition on its reduction. Divide by this unit. This gives membership with exponent $d_j = d_{j-1}$ and completes the descent.

By primitivity the elements $\bar{\zeta}^k$ for $0 \le k < e$ are distinct, and the reduction of the monic Q has degree $\deg Q$. Thus $d_0 \le m \deg Q \le \kappa$. We conclude $p^{\kappa} \in J_0(i; p^{\kappa+1})$. The explicit Nakayama argument following Proposition 49.6 passes to the zero-modulus ideal, proving exact annihilation. □

This proves the precise statement `TameChart.killsCokernel_of_isKilledBy`; the exponent depends on the degree of the annihilating polynomial, not on the number of residue classes of the tame inertia group. The proof distinguishes two different mechanisms in the descent: factorial cost at a root of the polynomial, and integral division by z at every other exponent.

52.9 Initial depth and final safety for few low jumps

We now prove the remaining case calculations of `FewLowJumps.lean`. To avoid confusion with the Frobenius index, write s for the parameter denoted a in that file, and a for a Frobenius

index as in the preceding chapters. The chart is unramified, $p \geq 5$, and at every letter i there are a sign ε_i and numbers b_i, n_i, r_i with the following properties:

$$\begin{gathered} b_i + W + \eta \leq (p-3)s, \qquad 6(s+\eta+W) < b_i, \\ U_i \subset [-\eta, \eta] \cup \varepsilon_i[b_i, b_i + W], \\ p \nmid n \quad \text{for } 0 \neq n \text{ within } 6(s+\eta+W) \text{ of } 0, \pm b_i. \end{gathered}$$

Here n_i, r_i bound f times the low and all jump multiplicities respectively. We use the sign on the side whose low weights are few (denoted S); the paired side T has opposite sign. Put $\rho_t(a,u) = t\mathbf{1}_{a=0} - u$. The cost at level zero is $v_p(n)$ with zero cost at $n = 0$, since the different exponent of this chart is zero.

The initial submodule. For $-\eta \leq t \leq 0$, let $D_i(t)$ consist of the actual constants lifting to localized points with exponent system ρ_t and depth $\rho_{2s+\eta}$ in the extension on the side S. We claim

$$\operatorname{length}(C_i/D_i(t)) \leq s \sum_{u \in [-\eta, t)} f \operatorname{mult}_i(u). \tag{52.10}$$

Here length may equally be computed for the inverse image in the original constants under the injective letter-constant identification used in `constantAt`; this cannot increase the bound.

First, for each $0 \leq k \leq s$ the unramified cokernel theorem gives a lift over $p^k c_i$ with exponent and depth both ρ_{k-s}, because $p^k E^{s-k}$ is a cokernel killer. Improve its depth at $a = 0$ to $\rho_{2s+\eta}$; every intervening integer on the allowed jumps lies in the three no-multiple windows, so the cost is zero by Theorem 51.13. Its improved exponents at $a = 0$ are

$$\begin{cases} 2s + \eta - u & k - s - u > 0, \\ \min(2s + \eta - u, 0) & k - s - u \leq 0. \end{cases} \tag{52.11}$$

At $a > 0$ they remain $-u$. This is the improved-exponent rule of Section 51.6, now with the exact comparison from Sections 52.3 and 52.7.

For $k = 0$, (52.11) is at least $-\eta - u$: on the low and far-positive windows the latter is nonpositive, while a far-negative jump satisfies $-s - u > 0$, since $b_i > 6s$. Thus $D_i(-\eta)$ contains every constant. For $k = s$ and any $t \in [-\eta, -1]$, it is at least $t + 1 - u$: if $u < 0$ the first branch applies, and otherwise $t + 1 - u \leq 0$. Hence $p^s C_i \subset D_i(t+1)$.

We prove the step of (52.10). Choose $f \operatorname{mult}_i(t) + 1$ constants of $D_i(t)$ with lifts of the indicated exponent and depth. Apply the primitive-relation lemma of Section 51.15 at the level-zero jump $u = t$. The strict lower-jump condition holds for ρ_t, since $t - v > 0$ when $v < t$. The lemma gives a primitive combination of lifts lying in the sum of the exponent lattice with ρ_{t+1} and that with ρ_t raised by one at $(0, t)$. The former lattice is contained in the latter, so the combination has the single strengthened exponent and keeps the depth $\rho_{2s+\eta}$ by linearity. Apply the improved-exponent rule again: at $u < t$ the old exponent is strictly positive; at $u = t$ it is now 1; and at $u > t$ the desired new exponent $t + 1 - u$ is nonpositive. Thus the combination has all the exponents ρ_{t+1}, proving its projected constant belongs to $D_i(t+1)$. The extension has the same multiplicity as the kernel at $t < 0$ (only the trivial quotient weight 0 is added). This gives the rank bound $f \operatorname{mult}_i(t)$ on the p-torsion of the successive quotient. It is killed by p^s, as just proved, and therefore has length at most $sf \operatorname{mult}_i(t)$, by the finite-length argument of Theorem 51.25. Sum the steps to prove (52.10) and obtain cost at most sn_i at $t = 0$.

The two possible final depths. For $\varepsilon_i = +1$ no further change to the $t = 0$ module is needed. For $\varepsilon_i = -1$ let qp be the least multiple of p which is at least $b_i + W + 2s$. Raise the depth at $a = 0$ to the constant qp on every jump. The one-step colength bound of Theorem 51.25 can be iterated by always choosing the smallest jump that still needs an increase. Its strict lower-jump condition then holds because the completed target depth is antitone; for $a = 0$ the additional open interval of length $p^a = 1$ contains no integer. For a low jump the sum of nonzero costs is at most $v_p((qp)!) \le s + 1$, since

$$qp < b_i + W + 2s + p \le (p-1)s + p$$

and $p \ge 5$. For a far-negative jump start at $2s + \eta - u$. The steps up to $b_i + W + 2s$ lie in the stipulated no-multiple window near b_i (if the start is larger there are no such steps); the remaining steps below qp contain no multiple of p by its choice. Thus every far step costs zero. The total additional colength is at most $(s + 1)n_i$.

Replace the depth at $a = 0$ by exactly qp (weakening any larger depths). The open interval $(qp, qp + p)$ now contains no multiple of p, as required for the $a = 1$ one-step colength estimate. At that level raise the depths from $-u$ to

$$W + 1 - u + 2s\,\mathbf{1}_{|u|\le\eta}.$$

The target is antitone on the allowed jumps: the gap $b_i > 6(s + \eta + W)$ is larger than the possible extra increment $2s$ when crossing a gap. Each step occurs in one of the no-multiple windows near $0, \pm b_{i[-1]}$, so its cost is at most 3 including the Frobenius cost. Summing with multiplicities gives at most $3(W + 1)r_{i[-1]} + 6sn_{i[-1]}$. Finally weaken the order at $a = 0$ to $b_i + W + 2s$. We have proved the exact estimate

$$\operatorname{codim}_{\text{length}} D_i \le (2s + 1)n_i + 3(W + 1)r_{i[-1]} + 6sn_{i[-1]} \tag{52.12}$$

for standard exponent lifts with the desired sign-dependent depths:

$$\begin{aligned}
&D_S(a, v) = \rho_{2s+\eta}(a, v) && (\varepsilon_i = +1),\\
&D_S(0, v) = b_i + W + 2s,\\
&D_S(1, v) = W + 1 - v + 2s\mathbf{1}_{|v|\le\eta},\\
&D_S(a, v) = -v \ (a \ge 2) && (\varepsilon_i = -1).
\end{aligned}$$

These are precisely the asserted depths in `exists_relLength_deepConstants_le`. Intersecting the inverse images of these submodules in the parameter lattice and summing (52.12) gives at most

$$11 \sum_i \big(n_i(s + 1) + r_i(W + 1)\big),$$

using that $i \mapsto i[-1]$ is a permutation.

The safety verification. For completeness we check every inequality in `HasDeepPoint` and hence in `scaledCupBounds_of_points_at_cuts`, taking every letter to be a cut, so the nearest preceding cut is one step away. Fix a constant in the just constructed intersection on the side S and an arbitrary constant on the opposite side T. Write u for a jump on T, v for a jump on S, and $w = -u - v - 1$ for the third-side jump. The bounds imply $w \le b_i + W + \eta \le (p - 3)s$, and the possible w are within $2\eta + 1 + W$ of $0, \pm b_i$. Hence the unramified killer E^s is available for T and the third side, and the depth-raising intervals used below all lie in the no-multiple windows: their endpoints differ from the relevant centre by at most $2s + 3\eta + W + 1 \le 6(s + \eta + W)$ (here $s, \eta \ge 1$). Thus all these lifts have exponent cost zero.

Choose a T-point of depth $D_T = \rho_{2s+\eta}$ obtained from E^s as above. Its improved exponent O_T is $-u$ at $a > 0$, and at $a = 0$ it is

$$O_T(0,u) = \begin{cases} 2s+\eta-u & u < -s, \\ \min(2s+\eta-u, 0) & u \geq -s. \end{cases}$$

In particular $\min(-u, O_T(0,u)) \geq -u - s$, and the possible loss from $-u$ occurs only on the low negative window, where it is also at most η. Let O_S be the improved exponent of the chosen S-point after the above depth construction. Since it started with standard exponents, $O_S(a,v) \geq -v$ everywhere. If $\varepsilon_i = -1$, then at level zero $O_S(0,v) = D_S(0,v)$ for $v < 0$ and is zero otherwise, so is nonnegative. If $\varepsilon_i = +1$, then T has negative far window and every low $u \leq \eta$; hence $O_T(0,u) \geq 0$.

For the third side take exponent system Z from an E^s-lift of depth ρ_s:

$$Z(a,w) = -w \ (a > 0), \qquad Z(0,w) = \begin{cases} s-w & w < -s, \\ \min(s-w, 0) & w \geq -s. \end{cases}$$

The function in w is antitone. With $U = u + v$, note the useful alternatives

$$Z(0,-U-1) \in \{s+U+1, 0\}, \qquad Z(0,-U-1) = 0 \implies -s-1 \leq U \leq s-1. \tag{52.13}$$

Choose T as leading side for $\varepsilon_i = +1$, with third shift $r = 0$; choose S as leading side for $\varepsilon_i = -1$, with third shift $r = 1$. This is exactly the choice $Q(i) = (\varepsilon_i = 1)$ in the compact source. Both leading exponent systems dominate st_1, as just checked at level zero.

First, for either side at unshifted third exponent with resting order $\min(-v, O_S)$ or $\min(-u, O_T)$, the sums at every $a > 0$ are at least 1, because $Z = u + v + 1$, the resting order is at least the standard one, and both depths there dominate the standard orders. At $a = 0$ the T-sum is $Z + 2s + \eta - U$, which by (52.13) is at least $s + \eta + 1 > 0$. If $\varepsilon_i = +1$, the S-sum differs from this by a loss at most η, hence is positive. If $\varepsilon_i = -1$, its depth is $b_i + W + 2s$. When $Z \geq 0$ the resting order is at least $-(b_i + W)$, so the sum is positive. When $Z < 0$, the resting order is at least $-u - s$; the sum is at least $b_i + W + 2s + s + U + 1 - u - s \geq 2s + 1$, since $v \geq -(b_i + W)$. This verifies the first inequality of `HasDeepPoint` on both sides.

Now verify the second inequality for the leading side, with the resting exponent the minimum of its current and one-step transported exponents. When T leads, the third shift is 0. At level zero the resting exponent can be weakened to $\min(-v, 0)$, which loses at most η from $-v$, because S has positive far window. The earlier lower bound $s + \eta + 1$ remains positive. At positive indices the minimum is at least the standard order; the sum is at least 1.

When S leads, the third shift is one. At $a = 0$ its exponent is zero, and the resting order is at least $-(b_i + W)$ (the opposite side T has positive far window). Thus the sum is at least $2s$. At $a = 1$ work at letter $i[-1]$; put $U = u + v$ there. The shifted third order is $Z(0, -U - 1)$ and the resting exponent is at least $-u - s$, with loss possible only if u is low negative. Write this loss as $d \in [0, s]$. The sum is at least

$$Z(0,-U-1) + W + 1 - U + 2s\mathbf{1}_{|v| \leq \eta} - d.$$

If $Z = s + U + 1$, this is at least $W + 2$. If $Z = 0$, then $U \leq s - 1$. For low v the extra $2s$ again makes it at least $W + 2$. If v is far and u is low, the opposite sign windows and $b_{i[-1]} > 6(s + \eta + W)$ imply $|U| > s + 1$, contradicting (52.13). If both are far, the opposite windows imply $U \leq W$, and $d = 0$, so the sum is at least 1. At $a \geq 2$ all orders in the sum are standard, including the shifted third order, so the sum is 1. This finishes the last safety inequalities.

The third-side points have antitone exponents and the least positive preceding cut is 1 at every letter. We give the direct walk, using the exact safety criterion of Theorem 51.17. Write x_i, x_i' for the new points at letter i, and y_i, y_i' for the points transported by one Frobenius from $i[-1]$. Move the leading side first, while the transported other side rests. At $a = 0$ the moving order is at least zero (the leading standard order), the transported resting depth is at least one, and the shifted third order is zero. At $a \geq 1$, the moving order is at least the minimum of the earlier-cut order and the standard order; apply the first inequality previously proved, at $i[-1]$ and $a - 1$, with its third point transported by one. The remaining move, by the nonleading side while the new leading point rests, uses exactly the second inequality proved above with shift r. Implement these local two-move paths in four global moves by choosing the first intermediate component at each letter to be either the old or new one according to which side leads, and the second intermediate component to be the new one. A cocycle move of either side while the other rests is safe by the first inequality (the other-side depth dominates the standard order). Distinct letter parts pair to zero, as in the proof of Theorem 51.20, and $\varphi(\sum_i x_i) = \sum_i y_i$. Thus Theorem 51.19 applies with $e = e' = \kappa = 0$ and proves that every constant in the intersection just constructed pairs to a coboundary with an arbitrary constant of any paired side with the opposite windows. Together with the colength bound this is `exists_scaledCupBounds_few_low_jumps`. In particular the initial-depth and safety obligations that were left open in Section 51.15 have been verified with the compact source's exact hypotheses.

52.10 Exact compact-source numerical hypotheses for the cut bounds

The constants in the more readable layout of the source were packaged differently. Here we compare the hypotheses directly, so the compact source's two principal large-jump estimates need not be taken as additional black boxes. Keep the terminology of Theorem 51.22. In this paragraph only, b, G, X denote the parameters of the compact source, while we put stars on the corresponding chosen parameters of that theorem when they differ.

Sparse thick letters. The compact statement assumes $V + 4\eta \leq W$, $G + V \leq \eta$, $11(p^J + 1)W \leq b$, and

$$4(J + 1 + d_0 + L)W \leq X, \qquad 2e_0 b \leq p^{J+1}X, \qquad 8e_0 \leq p^{J+1}, \qquad L = \lfloor \log_p(2b) \rfloor.$$

It assumes the same no-three condition and either constant signs or an unramified chart with $p \geq 7$. Let $b_* = b + W$ and take in Theorem 51.22

$$s_i = 0, \quad n_* = G_* = 2W, \quad X_* = X, \quad \delta_* = 2X, \quad \kappa = 12((J + 1)X + 2X) + 1.$$

When $p \geq 5$ set $s_0 = \lceil (b + W)/(p - 3) \rceil$; for smaller primes no bound involving s_0 is needed, so take it zero. Mark precisely the cuts whose next J letters are thin. The combinatorial facts preceding Section 51.14's choice of constants follow from the no-three condition, regardless of the value of W. If $J = 0$, all letters are cuts, nearest predecessor distance is one and every interval is plain. If $J > 0$, an interval which is not plain must have a thick interior letter with a sign different from the terminal letter, and this unique interior pivot satisfies $t + J + 1 = l$. Every pre-pivot window has endpoint at most η. For $p \geq 7$, using $\eta \leq W/4$ and $b \geq 11(p^J + 1)W$ gives

$$s_0 + \eta \sum_{a=0}^{J} p^a \leq \frac{b + W}{4} + 1 + 2\eta p^J \leq \frac{b}{4} + \frac{b}{44} + 1 + \frac{b}{22} < b.$$

Thus the long-gap condition holds with $\omega = \eta$. All cuts are of type (i), since their thin successors have $\beta_i + W \le \eta + W \le 2W$. The far bound at $p \ge 5$ holds by our definition of s_0. The other window constraints are immediate. In particular, $\beta_i + W \le b_*$, $W \le pn_*$, $G_* \le (p-1)n_*$. As $b + W \le 2b$, the logarithmic term $L_* = \lfloor \log_p(2b_*) \rfloor$ is at most $L + 1$. The cost inequality of the readable theorem follows from

$$2W + L_* + (J + 1 + d_0)W + 2d_0W \le (J + 3 + 3d_0)W + L + 1 \le 4(J + 1 + d_0 + L)W \le X.$$

The penultimate inequality uses $W \ge 4$. Moreover $2e_0b_* \le 2 \cdot 2e_0b \le p^{J+1}\delta_*$. The theorem applies and gives

$$51\kappa = 612(J + 3)X + 51 \le 10^4(J + 1)X,$$

since $J + 3 \le 3(J + 1)$ and $X \ge 1$. This proves exactly `scaledCupBounds_sparse_thick` in the compact source, including both its size and sparsity assumptions.

Two sizes. The compact statement has an unramified chart, $p \ge 7$, $5W \le G \le b$, $b < Kp^2$, $2G < Kp$ and $W \le K$. Use the choice made in Section 51.14, with $g = G$, $b_* = b + W$, $n_* = K$, $G_* = G + W$, $J = 1$, $X_* = 3K + L_*$, $\delta_* = 4K$ and $\kappa = 12(2X_* + 4K) + 1$. There all the numerical hypotheses and the two-step predecessor choice were verified using only the weaker $0 < g \le b$, so they apply here unchanged. The expanded radius bound $2b_* \le 4Kp^2$ holds, since $W \le K$ and $b < Kp^2$. In particular this specialization does not require inferring an extra jump hypothesis from $5W \le G$. The resulting exponent is

$$51\kappa = 6120K + 1224L_* + 51 \le 6120K + 1224(L + 1) + 51 \le 10^4K(L + 1).$$

Indeed $K \ge 1$ and $L \ge 0$ and $L_* \le L + 1$. This is the compact source's stated numerical bound `scaledCupBounds_two_sizes`. Multiplication by additional powers of p preserves the coboundary conclusion. These two specializations therefore use the proved readable cut scheme with explicit constants, rather than requiring an identification of differently parameterized lemmas by name.

52.11 Functorial frame operations used in the lifting proof

For use in the cup calculations we give the remaining elementary compatibility details. Let T_1, T_2, T_3 be crystalline lattices on the chart, with the intrinsic positive modules and localized modules as in Theorem 52.8; take crystalline forms over the invariant field. A map of lattices carries the intrinsic positive modules into one another: it preserves $A[1/\tau]$-coefficients, and composing an equivariant form on the target with the map gives an equivariant form on the source. In localized coordinates all the same statements hold after multiplying by a common product of the E_a. When both source and target are value-field points, the map preserves such points, again by composing forms. Consequently its matrix with respect to the chosen integral frames has entries in S (apply the map to each basis vector). The coefficient retractions therefore commute with this map, including after localization. This proves the functorial retraction statement for projections, inclusions, and letter projectors used in Sections 51.6 and 51.15; an abstract splitting of an extension as representations is neither asserted nor needed.

For an exact sequence with trivial quotient, choose an integral splitting of the underlying lattices just to identify the kernel inside $A[1/\tau] \otimes T$. If a localized point over zero is given, it corresponds to a unique element of the kernel in these ambient coordinates. Every equivariant de Rham form on the kernel extends to an equivariant form on the extension; choose a basis of invariant columns for the kernel, extend those to a basis for the extension over the fixed field,

and extend the dual basis. The same construction is compatible with the Hodge filtration: the invariant spaces in all three terms have adapted bases as in Section 52.6; the induced filtration on the kernel is its intersection with that of the extension, directly from their embedding after extending scalars to B_{dR} and using the integral underlying splitting. For the dual forms, restriction of a form of type u preserves membership in the filtration, and a form on the kernel of type u has an invariant extension of that type. To see the latter assertion, dualize a basis adapted to the filtration and choose the basis for the kernel filtration as part of that of the extension. The adapted bases have leading independent columns after extension of scalars, proved in Section 52.6; an invariant form written in their dual basis therefore has no coefficients in the degrees forbidden by its type. Extend the coefficients by zero on the other basis vectors. Evaluation then shows that the kernel of the localized projection intersected with the prescribed exponent conditions is exactly the kernel's localized module intersected with those conditions (on the common sets of jumps, as in the source). This supplies the identification used in `exists_inclKer_eq_of_kerWithOrders`. It also gives the assertion `exists_span_formsOfType_mod_pred`: the difference between filtration degrees u and $u-1$ is spanned by the degree-u dual basis vectors, precisely the Hodge multiplicity. The calculation uses only the actual forms and hence applies to a jump set enlarged by integers of zero multiplicity.

For a bilinear equivariant lattice map, the tensor of invariant de Rham bases gives an invariant basis on the tensor product. Their filtration is the sum filtration, by the leading-matrix argument of Section 52.6 applied to the tensor of the two invertible leading matrices. Thus the pullback of a target form is the sum of the products of the dual forms with the stated filtration inequalities. The map preserves the intrinsic positive modules by this calculation at every positive Frobenius iterate, and preserves value-field points by the same calculation in the value fields. This justifies the tensor-compatible use of the localized frames and the order addition in (51.7). In particular the form on the third extension causes no change of integral lattice when used to test safe pairs in Theorem 51.17.

Finally consider Frobenius transport in `phi_pow_smul_mem_pointsOver`. The basis at positive level l is $\varphi^{l}n$ by its construction, and $E^{h}\varphi(n) = nB_1$ over S. Thus transport is a map of the actual localized spaces; the projections and letters transform as prescribed. At a newly appearing divisor $a < l$, the evaluation of the transported point lies in the positive ring by the defining intrinsic positive-module property at $l - a \geq 1$. Its Galois difference has positive filtration: the integral value has residue in the fixed field by the value-field assertion at level zero (applied to the positive iterate), so the residue of its difference is zero. At all other places the evaluations are the old ones with shifted index. This proves precisely the order-zero and depth-one new-place bounds used in the walk. The multiplication by τ in the Tate-line version follows directly from its one-dimensional intrinsic module: an element of standard orders on the Tate line has coefficients in $A[1/\tau]$ whose evaluations, after multiplying by τ, have nonnegative order; cancelling the powers of τ by Lemma 52.4 puts the multiplied element into the integral span. In this way the comparison statements required before Theorem 51.17 reduce to the integral intrinsic-frame theorem, not to an external tensor-compatibility assertion.

52.12 Both alternatives of the no-multiple bound

The compact `scaledCupBounds_of_killsCokernel` is a second all-letter walk, used also outside the few-low-jumps specialization. We verify its two alternatives separately. Suppose each of the three sides has cokernel killer $p^{\kappa}E^{s}$ at every letter, the first two sides have the opposite windows of radius η near zero and width W near $\pm b_i$, and the no-multiple condition

holds within radius $6(s+\eta+W)$ of each centre $0, \pm b_i$. Put

$$K = \kappa + (48(s+\eta+W)+4)d_0, \qquad D(a,u) = \rho_{2s+\eta}(a,u), \qquad Z(a,w) = \rho_{-s}(a,w).$$

From an initial point of exponent and depth ρ_{-s}, raise the depth to D. The only steps occur at $a=0$, and they belong to the union of the three windows: their distance from the relevant centre is at most $2s+2\eta+W$. Each has cost d_0 (the integer is zero or prime to p). The union is contained in the union of the two two-window sets, whose total cardinal is at most $8 \cdot 6(s+\eta+W)+4$. Thus the new point is over $p^K c_i$ after multiplying by any unused power of p, and it has improved exponent

$$O(a,u) = -u \quad (a>0), \qquad O(0,u) = \begin{cases} 2s+\eta-u & u < -s, \\ \min(2s+\eta-u, 0) & u \geq -s. \end{cases}$$

In particular $O(0,u) \geq -u-s$ and $O(0,u) \geq 0$ whenever $u \leq \eta$. Take the third-side point from the original cokernel killer and multiply it by the extra power of p to put it over p^K; its antitone exponents are Z.

In the first alternative neither side has a jump in $[-s, 0)$. Consequently $O \geq -u$ everywhere, and the first safety inequality at a cut (with both sides of depth D) is

$$Z(0, -u-v-1) + D(0,u) + \min(-v, O(0,v)) \geq s+\eta+1 > 0.$$

At $a>0$ it is at least 1. Let the side with negative far window lead, so its jumps at the current letter satisfy $u \leq \eta$ and its order at $a=0$ is nonnegative. Its order system dominates the standard one at level 1. The other side has $v \geq -\eta$; the minimum of its order at $a=0$ and transported order zero is at least $-v-\eta$. Thus the second safety inequality with third shift $r=0$ at $a=0$ is at least $s+1$; at $a>0$ it is at least 1 because both resting orders dominate the standard order. The direct all-cut walk proved in Section 52.9 applies.

For the second alternative the second side S' has all jumps in $[-\eta, 0]$, the chart is unramified, $p \geq 5$ and $\eta \geq 1$. The zero-shift and zero-power killer is available for S' by the unramified cokernel theorem, because its jumps are nonpositive. Starting from this killer, raise the depth to D as before (zero cost), transport from $i[-1]$ to i by one Frobenius, and deepen at the new divisor $a=0$ to $2s+\eta-v$ with the same zero-cost operators. The new orders and depth on S' can be taken as

	O'	D'
$a=0$	0	$2s+\eta-v$
$a=1$	$\geq -v$	$2s+\eta-v$
$a \geq 2$	$-v$	$-v$.

The other side S has the point (O, D) above. All points may be put over p^K times their respective constants. For the first safety inequality at level zero, with S resting, the sum is at least $s+\eta+v+1 \geq s+1$; with S' resting it is at least $\eta+1$, since the possible loss of O from the standard order $-u$ is at most s. At positive levels the sums are at least 1. Always let S' lead. If the first side has positive far window, use third shift $r=0$: at level zero its moving order (minimum of the present and shifted earlier order) is at least $-u-\eta$ because $u \geq -\eta$, and the sum is at least $s+1$; at level one its moving order is at least $-u-s$, and the sum is at least $s+\eta+1$. If the first side has negative far window use third shift $r=1$: its order at zero is nonnegative, since $u \leq \eta$, so the moving order is zero, the shifted third order is zero, and the sum is at least $\eta > 0$. At level one the third term is $-s+u+v+1$, the moving order is at least $-u-s$, the leading depth is $2s+\eta-v$ and the sum is at least $\eta+1$. At levels $a \geq 2$ in both cases all terms meet the standard orders and give sum at least 1. This verifies the second inequality with exactly the shift prescribed by the compact theorem. The

all-cut path and Theorem 51.19 apply in both alternatives with $e_1 = e_2 = e_3 = K$ and give the asserted bound for every $H \geq 3K$. In particular, in the unramified chart, if $\kappa = 0$ the claimed loss is exactly zero, not a bound depending on the rank or the size of the far jump.

Standard exponents and the intrinsic lattice. For clarity, the intrinsic module identification also shows `mem_withExponents_stdExponent_iff`, used implicitly in the lifting recurrence. Write x in the localized A-span of the integral frame, so denominators of its plain-lattice coordinates are finite powers of τ and E_a. At a letter, normalize a basis of invariant dual forms by its individual filtration degrees, and use the invertibility of its leading matrix from Section 52.6. At a place $a \geq l$, the standard order conditions $-u$ then say exactly that the plain-lattice coordinates of $\varphi^{-a}x$ are integral in B_{dR}^{+}. Since E_a is the sole possible denominator vanishing there, the full power of that denominator cancels in A (use primality and theta, successively). At places $a < l$ the remaining possible powers of E_a are absorbed by a power of $\varphi^l\tau$: it is divisible by every E_a with $a < l$, by Lemma 52.4 or the factorization of $\varphi^l(\tau)$. Hence $\varphi^{-l}x \in A[1/\tau] \otimes T$. Its positive-iterate evaluations are nonnegative: for the new places this is the order-zero part of the standard conditions, and for the remaining positive iterates it follows by the defining intrinsic property, since localization by any E_a has no pole at such an iterate. Thus $x \in \varphi^l M_T$. Conversely a vector in $\varphi^l M_T$ has the order-zero conditions at $a < l$ by the same definition, and at $a \geq l$ its plain-lattice coordinates have nonnegative filtration (the denominator is a power of $\varphi^{l-a}\tau$, which has nonzero theta value). Applying any form of type u now gives order at least $-u$. This proves the equivalence on the prescribed letter. For the cyclotomic power χ^c the unique weight has absolute value c; the inclusion $\tau^c M_T \subset A \otimes T$ in Theorem 52.8 proves `tau_pow_smul_mem_ainfLattice_of_chi`, including $c = 0$ and any multiplicity.

52.13 From side pairings to the local defect theorem

We record the two final implications from `New/Lift/ExponentTheorems.lean`, so that the coefficient ring and the tame restriction do not remain implicit hypotheses.

Let O be the ring of integers of a finite extension of $\mathbb{Q}_p$, with residue degree f_O, ramification degree e_O and degree $d_O = e_O f_O$. The module $Z_f(T)$ of finite-condition cocycles is a finite free $\mathbb{Z}_p$-lattice: the exact sequence of coboundaries and finite-condition classes embeds the quotient of the former into the finitely generated local integral cohomology, and the coboundary module is itself a quotient of the finite free lattice. The submodule of actual cocycles is torsion-free. Finiteness of the local integral cohomology and the finite-class interpretation are the standard local cohomology facts used in Chapter 50. Give $C = Z_f(T)$ the trivial action, and form the extension of C by T with identity cocycle map $C \to Z_f(T)$. It has the crystalline condition after the same restriction as T: for a basis of C, every cocycle becomes a coboundary in the crystalline period module, so choose period primitives and subtract them from the trivial lifts. Together with a crystalline invariant basis of T, these corrected lifts form an invariant basis of the extension. Its jump multiset is that of the kernel with the zero weights of the trivial quotient added, by the filtration-adapted basis and exact sequence argument of Section 52.11. No bound in the exponent proofs charges a multiplicity of the trivial quotient at a nonzero weight. In the few-low-jumps estimate the length calculation is on the kernel multiplicities (as in Section 51.15), not on the multiplicity of the trivial quotient.

This is `KummerChart.universalSide`. The third side for a congruence pairing modulo p^m is the extension by its defect cocycle, as constructed in Section 51.17. It is rationally split, since its defect is the coboundary of the chosen lifting form divided by p^m, and therefore crystalline on the subgroup on which the tensor dual is crystalline. Our side

pairing statements apply to each first lattice, congruence pairing, and cocycle; thus their annihilation exponents (or the common submodule of constants in the few-low estimate) place the resulting second-side cocycles in the intersection of the cup annihilators. This is precisely `mem_cupOrthogonal_of_scaledCupBounds`: the original pairing between integral reductions is the reduction of the third-extension form, not a pairing between dual rational Galois representations.

For the few-low bound let μ bound the sum of the low multiplicities at each coefficient embedding, and let $r = \mathrm{rk}_O T$. The letters partition the d_O embeddings of the coefficient field according to their value on the specified root of unity. Their total multiplicity counts are the sums over the corresponding coefficients, by the Hodge–Tate basis after scalar extension; the distinct Teichmüller roots have unit differences. Thus the bounds n_i, r_i of Section 52.9 can be chosen with

$$\sum_i n_i \le f d_O \mu, \qquad \sum_i r_i \le f d_O r.$$

In particular the colength estimate (52.12), summed over the letters and applied to the universal cocycle extension, gives a common submodule of $Z_f(T)$, contained in the cup orthogonal for the whole class of paired lattices, with $\mathbb{Z}_p$-length of the quotient at most

$$11 f d_O\big((s+1)\mu + (W+1)r\big).$$

For any finite-length O-module, its $\mathbb{Z}_p$-length is f_O times its O-length, since every simple O-factor has cardinality p^{f_O}. The cup orthogonal intersection is an O-submodule; its quotient length is no larger than that of the quotient by the constructed $\mathbb{Z}_p$-submodule. Divide by f_O and use $\mathrm{length}_O(O/pO) = e_O$ to obtain precisely

$$\mathrm{length}_O Z_f(T)/(Z_f(T) \cap \mathrm{cupOrthogonal}) \le 11 f\big((s+1)\mu + (W+1)r\big)\,\mathrm{length}_O(O/pO).$$

The unramified killer is available for the third side since its jumps $-u-v-1$ are bounded above by $b_i + W + \eta$; the same bound holds for the first two sides. Therefore Section 52.12, with $\kappa = 0$, proves both zero-loss alternatives. These are the two conclusions of `KummerChart.few_low_jumps`, or equivalently Theorem 49.7 after the stated length normalization.

The tame type. Here the Galois group is identified with G_p by the assumed open embedding and surjection. Put

$$M = \mathrm{lcm}_{1\le j\le r}(p^j - 1), \quad m = r^2, \quad \kappa = r^2(N+1)^2.$$

The empty least common multiple is 1. The size hypothesis $b + 6(W+\eta) + \kappa + 2 < p$ implies $p \nmid r!$. Each $p^j - 1$ for $1 \le j \le r$ divides $p^{r!} - 1$, so M divides it, and $M \le \prod_{1\le j\le r}(p^j - 1) \le p^{r^2}$. Construct the tame chart of Section 52.8 with $f = r!$, $e = M$.

We justify why every potentially crystalline lattice of the specified tame type is crystalline on that chart subgroup. Its space of near-identity invariant periods spans over B. For each near-identity invariant scalar, the invariant ratio proposition applied to an open subgroup and its chart puts it in a finite unramified extension of $\mathbb{Q}_p$. The union of these fields has trivial inertia action. Consequently on near-identity invariant vectors the linearized inertia action agrees with the ordinary action and is multiplicative: changing the invariant basis introduces only such unramified scalars, fixed by inertia. It is trivial on an open subgroup of inertia, since finitely many invariant basis vectors suffice. Its finite image has exponent dividing M by the annihilator $X^M - 1$. Wild inertia is pro-p, hence maps trivially into that image; tame inertia is procyclic with prime-to-p finite quotients. The kernel therefore contains the subgroup of tame inertia of index M, namely the inertia fixing the chosen M-th root of p.

Here we use the classical description of tame inertia; see J.-P. Serre, *Local Fields*, Springer GTM 67 (1979), Chapter IV, §2, pp. 67–68, Corollaries 1, 3 and 4 (the tame quotient and the pro-p wild subgroup), together with the Eisenstein construction of the root extension in Section 52.8. The chart inertia thus fixes every near-identity invariant period.

For completeness, the passage from inertia-fixed periods to crystalline periods on the chart is finite unramified descent. Choose a finite Galois extension L/K on which an invariant basis exists (K is the chart field). Its invariant coefficient ring after inverting p is precisely L_0, the maximal unramified subfield: Proposition 52.6 gives inclusion and the Teichmüller coefficients give the reverse inclusion. The G_K/G_L action on its invariant space is semilinear over this field and trivial on the inertia quotient, as shown above. The remaining action is that of $\mathrm{Gal}(L_0/K_0)$; ordinary finite Galois descent (equivalently matrix Hilbert 90) gives a K_0-basis of fixed vectors spanning after extension to L_0. Thus these vectors also span after base extension to B, giving the crystalline property on G_K. This proves the restriction step used before the tame cokernel theorem, even when the original representation is not crystalline on G_p.

We also need the polynomial for the third side. For linear operators killed by monic integral polynomials q_1, q_2 of degrees d_1, d_2, the tensor operator is killed by a monic polynomial of degree at most $d_1 d_2$: form the finite free $\mathbb{Z}_p$-algebra $\mathbb{Z}_p[X,Y]/(q_1(X), q_2(Y))$, of rank $d_1 d_2$, and use the characteristic polynomial of multiplication by XY. The element XY satisfies that polynomial by Cayley–Hamilton, and the identity passes to every module on which X, Y act via the two commuting tensor factors. For duals one uses the given polynomials at the inverse inertia element, then transposes; the Tate twist has trivial linearized inertia operator, its invariant basis being the cyclotomic period times the dual lattice generator. The degree for the tensor dual is therefore at most N^2. Tensor compatibility and the rational splitting of the third extension are as above. Thus the monic kernel polynomials on the two sides and the third side of a pairing have degrees at most N, N, N^2. The hypotheses of Theorem 52.11 hold for each, since

$$m(N^2+1) \le \kappa, \qquad n = b + 6(W+\eta) \text{ bounds their upper jumps}, \qquad n + \kappa + 2 < p.$$

They each give the exact killer p^κ. This time $s = 0$; the no-multiple condition in Section 52.12 follows from the strict size bound, since the absolute value of every integer in the three windows is smaller than p. Its first alternative is automatic for the empty interval $[-s, 0)$. The chart uses the coarse parameter $d_0 = 0$ for $e = 1$ and $d_0 = 1 + v_p(e)$ otherwise. In the tame chart $d_0 \le 1$. Thus Section 52.12 gives a scaled cup bound with exponent at most $3\kappa + 144(\eta + W) + 12$, which is at most the compact statement's exponent

$$3r^2(N+1)^2 + 150(\eta + W + 1).$$

Applying the universal cocycle construction and intersecting all pairing annihilators gives precisely `tame_types_exponent`. The proof explains separately the three uses of tame type: field restriction to a crystalline chart, a small degree bound rather than a rank bound, and an exact cokernel killer prior to the cup operation.

52.14 Obligation-by-obligation closure record

PROVED below means a mathematical derivation in this volume, using the elementary classical inputs explicitly identified. **REFERENCE** means an external precise standard theorem whose hypotheses match the use. **OPEN** identifies a residual obligation of the translation, not an assertion that the relevant Lean statement is false.

1. **REFERENCE:** construction of the perfect tilt and Witt ring, and the surjective Fontaine map with distinguished principal kernel, as in Bhatt–Morrow–Scholze, *Integral p-adic Hodge theory*, Lemma 3.10 and Remark 3.11. De Rham uniformizer and its cyclotomic transformation law: Brinon–Conrad, *CMI Summer School Notes on p-adic Hodge Theory*, Proposition 4.4.8 and the preceding construction. The maximal-ring integral lattice used here is not identified with the divided-power integral one.

2. **PROVED:** `Amax.mem_span_pow_of_p_pow_dvd`, `Amax.exists_p_pow_mul_eq_add`, `BPlus.exists_Ainf_approx`, closure of principal ideals, regularity of ξ/p modulo p, and separatedness and contraction from the actual maximal period ring. Sections 52.1–52.2, together with Lemma 51.1.

3. **PROVED:** adic coordinates, bicommutant approximation, bounded denominator removal for divisors of μ, Frobenius conjugation, both cyclotomic cancellation tests and z-separation. Sections 52.2–52.4. These justify the structural assumptions of the earlier period-basis and determinant reduction.

4. **REFERENCE:** Tate–Sen finite-field invariants and vanishing of the nontrivial cyclotomic twists, Brinon–Conrad Theorem 2.2.7. Arbitrary algebraic-extension invariants of the completed algebraic closure: Ax, J. Algebra **15** (1970), 417–428, theorem on p. 417.

5. **PROVED:** filtered invariant scalar representatives, `dvd_extendToBPlus`, `exists_witt_coeff_of_isChartOf`, `sum_map_mem_of_valueField`, `eventually_exists_coeff_eq_of_fixed`, and `smul_sub_mem_hodgeFil_of_valueField`. Sections 52.3 and 52.7 give the required exact filtrations, not just rational field invariants.

6. **PROVED:** first half of `radius_lemma`, including the rank-free integral formal-series unit factorization, Theorem 52.7; combine it with (50.10) for the required $p^{4\delta}$ span.

7. **PROVED:** actual Kummer chart and scalar congruences, Sections 52.5 and 52.8; `IsChartOf.exists_isPeriodMatrix`, `IsPeriodMatrix.exists_frame`, `frame_theorem`, and the scalar linear algebra needed to identify the intrinsic frame, Section 52.6 and earlier Theorem 51.4. The hypotheses here are exactly the explicitly defined crystalline invariant-basis condition on the chosen chart, not an assertion of crystallinity for arbitrary lattices.

8. **PROVED:** functoriality and value-field/letter retraction of the actual modules, filtration-adapted form expansion and kernel comparison `exists_inclKer_eq_of_kerWithOrders`, standard-exponent characterization, Frobenius transport including new-place depth one, and tensor-compatible addition of orders. Sections 52.11 and 52.12.

9. **PROVED:** integral derivation compatibility of the multiplier ideals `mem_liftIdeal_of_derivation`. Section 52.7. This supplies the remaining premise of the factorial cokernel recurrence and the rank-free depth operators in the preceding chapter.

10. **PROVED:** tame chart construction and near-identity integral polynomial lowering, `exists_z_smul_eq_sum_tameGen`, the monic polynomial $q(X)(X-1)$ on the extension, and the exact recurrence `TameChart.killsCokernel_of_isKilledBy`. Section 52.8. The exceptional polynomial roots and the zero-cost nonroot lowering are distinguished explicitly.

11. **REFERENCE:** the wild/tame inertia decomposition, Serre, *Local Fields*, Chapter IV, §2, corollaries on pp. 67–68. **PROVED:** the tame type restriction by unramified descent, degree-N^2 kernel polynomial on the tensor dual, and `tame_types_exponent`, Section 52.13.

12. **PROVED:** exact compact-source `scaledCupBounds_sparse_thick` and `scaledCupBounds_two_sizes`. Section 52.10 checks their published-in-source numerical hypotheses against the already proved cut theorem, including the two output exponents.

13. **OPEN (auxiliary generality only):** a proof of the standalone compact `scaledCupBounds_of_cut_windows` for every one of its abstract parameter choices, with its own optimal 51κ constant, rather than the slightly slack readable theorem. The preceding item proves both of its actual uses within `SparseThickLetters.lean` with the exact advertised final constants, so the additional generality of this helper is not invoked to deduce those two corollaries.

14. **PROVED:** initial depth construction, one-step multiplicity estimate, sign-dependent depth arithmetic, final all-letter safety and both zero-loss alternatives, Sections 52.9, 52.12 and 52.13, together with Theorem 51.25. In particular `exists_relLength_deepConstants_le`, `exists_scaledCupBounds_few_low_jumps`, `scaledCupBounds_of_killsCokernel` and `KummerChart.few_low_jumps` receive a mathematical derivation on the actual constant lattices.

These are the specific structural and local crystalline lifting obligations listed in Section 51.16 and in the earlier audit. Their local conclusions still require the stated admissibility and Hodge-jump hypotheses for the particular global congruence representations; establishing those hypotheses for the arithmetic family, including its local–global type comparisons and existence, is a separate problem, not a consequence of the intrinsic-frame or cup-product arguments. No claim of a completed independent global proof is made on the basis of this chapter alone.

Chapter 53

The quantitative crystalline loss for the actual weight windows

We now apply the integral cup-product results of Chapter 52 to the exact windows and constants in the global construction. The point of this step is to retain the small loss per string block and not merely a uniform estimate depending on the largest Hodge–Tate weight. We first give the general numerical reduction in `New/Selmer/CrystallineBound.lean`, and then verify the arithmetic spacing and the parameters of `block_defect_le_at_place`.

53.1 A numerical consequence of the integral cutoff theorems

Let p be a prime and use a Kummer chart over a local field of ramification index e_0, residue degree f_0 and absolute degree $d = e_0 f_0$. Write $d_0 = 0$ if $e_0 = 1$, and $d_0 = 1 + v_p(e_0)$ otherwise, as in the source. Let Q, m, h, μ be nonnegative integers with $Q > 0$. Let G, b, δ, η, W have the source meanings: the two high-window centres are $G + \delta, b + \delta$, their width is W, and the low window is $[-\eta, \eta]$. Assume exactly the inequalities of `crystalline_defect_le`:

$$\begin{aligned} 99(Q^2 + (W+1)Q + Q + W + m) \le h, \quad 99h^3 \le G, \\ 99(Gh^3 + G) \le b, \quad 2\log(4b) < m, \quad \delta < \eta, \\ 99((\eta + W)Q + \eta + h) \le h^3, \qquad e_0 \le h. \end{aligned} \tag{53.1}$$

Define

$$\lambda = v_p\left((2b)!\left(\prod_{b-30h^3<j\le b+30h^3,\ j\ge 1} j\right)^{h^3}\right), \qquad K = \lfloor 2\lambda/Q \rfloor. \tag{53.2}$$

Here v_p on a positive integer is the usual normalized integral valuation. The product is nonzero. In particular

$$\lambda \ge \lfloor 2b/p \rfloor, \qquad 2b < p(\lambda + 1). \tag{53.3}$$

Also, if $1 \le n \le 4b$, then $\lfloor \log_p n \rfloor < m$, since $\log 2 > 1/2$.

The chart has letters labelled by the images of a primitive Teichmüller root of order $p^{f_0} - 1$ under the coefficient embeddings. Let T be a subset of these letters and $s(z) \in \{\pm 1\}$ a sign for each letter. For an embedding g, assume the two opposite jump windows of the source, of signs $-\varepsilon(g)$ and $\varepsilon(g)$ and centre $\beta_g \in \{b + \delta, G + \delta\}$. If g is thick (centre $b + \delta$), or $Q < p$, assume that its thickness and sign agree with T and s at its letter. Assume either that the chart is unramified and $p \ge 7$, or that $p \le Q$ and all signs on the letters are equal. When

$p \le Q$, suppose no three thick letters occur successively with either positive Frobenius gap at most

$$J = \lfloor \log_p(h^3) \rfloor.$$

As in the source, the jump sets U, U' of the paired classes are permitted sets, not necessarily multisets of a single representation. For a crystalline lattice L with jumps in U', of rank r_L, suppose either

1. all negative permitted jumps in U and U' are $\le -h/2$ and the number of jumps of L in $[-\eta, \eta]$ at every embedding is at most μ (counting multiplicities), or
2. U' is contained in $[-\eta, 0]$ at every embedding.

We prove the precise bound

$$\mathrm{normLength}_O\big(Z_f(L)/(Z_f(L) \cap \mathrm{cupOrthogonal}(\mathrm{InClass}(\top, \bot, U), L))\big) \le 10^5 d\,(r_L K m + \mu\lambda). \tag{53.4}$$

The normalized length here is length divided by the ramification index of the coefficient DVR. The symbol $\mathrm{InClass}(\top, \bot, U)$ means the crystalline class with the prescribed permitted jumps and no extra tame polynomial conditions, and the cup orthogonal is precisely the simultaneous annihilator used in the earlier local chapter.

Passage to letters. Adjoin zero to the permitted jump sets on each letter. When $p > Q$, every embedding on that letter has the same centre and sign by hypothesis; the letter jump sets therefore obey the corresponding original opposite windows. When $p \le Q$, enlarge the low radius to $2G$: the entire thin high window now lies in that radius, because

$$G + \delta + W \le G + \eta + W < 2G$$

by (53.1). A thick embedding already has the stated letter centre and sign. Thus every letter jump set in the small-prime case lies in the two corresponding opposite windows with low radius $2G$ and high width W. The centre assigned to the thick letters is $b + \delta$, to the thin letters $G + \delta$. This justifies the use of the union of embedding jump sets even when a letter contains both a thick and thin embedding.

The case $p \le Q$. Apply the exact sparse-thick bound in Section 52.10 with

$$b_* = b + \delta, \quad G_* = G + \delta, \quad \eta_* = 2G, \quad V_* = W, \quad W_* = 9G, \quad X_* = K.$$

Here V_* denotes the width of the high window in that theorem and W_* its accounting parameter. We verify its hypotheses explicitly. The inequalities (53.1) give

$$W + 8G \le 9G, \qquad G + \delta + W \le 2G, \qquad 11(p^J + 1)W_* \le 99(h^3 + 1)G \le b_*.$$

Also $p^{J+1} > h^3 \ge 8h \ge 8e_0$. Since $b \ge Q^2$ and $p \le Q$, (53.3) gives $\lambda \ge b/p$ (because $b \ge p$), hence

$$K \ge b/Q^2.$$

For example, $\lfloor 2\lambda/Q \rfloor \ge 2b/Q^2 - 1 \ge b/Q^2$. Since $h \ge 99Q^2$ and $\delta < b$, we obtain $2b_* < h^2 K$. Therefore $2e_0 b_* < p^{J+1} K$. Next,

$$K \ge b/Q^2 \ge 99Gh^3/Q^2 \ge 99^2 G h^2.$$

We have $J+1 \le m$, $d_0 \le m$ (use $e_0 \le h \le 4b$), and $\lfloor \log_p(2b_*) \rfloor < m$. In particular

$$4(J+1+d_0+\lfloor \log_p(2b_*) \rfloor)W_* \le 108mG \le (108/99)hG \le K.$$

The sign or unramified-chart hypothesis is precisely the one in the sparse-thick theorem, and the stipulated no-three condition is its letter-gap hypothesis. Thus the exponent is at most $10^4K(J+1) \le 10^4Km$ on every paired side.

For a potentially crystalline lattice of rank r_L, the rank of its crystalline cocycles over O is at most $(1+d)r_L \le 2dr_L$ by the rank formula of Section 8.4. Hence if $p^k Z_f(L)$ is in the cup orthogonal, the normalized quotient length is at most $2dr_Lk$ (apply the elementary length of a free O-module modulo p^k). The universal-side and cup application of Section 52.13 now bounds the defect by $2 \cdot 10^4 dr_L Km$; this is no larger than (53.4).

The case $p > Q$ and $\lambda \ge h^3$. The chart is unramified and $p \ge 7$. Use the exact two-sizes bound of Section 52.10 with $b_* = b+\delta$, $G_* = G+\delta$, low radius η, width W, accounting width $W_* = W + 4\eta$, and K as in (53.2). We check

$$5W_* \le G_* \le b_*, \qquad b_* < Kp^2, \qquad 2G_* < Kp, \qquad W_* \le K.$$

First $W_* \le 4(W+\eta) \le 4h^3/(99Q)$, so $5W_* \le G_*$ and $W_* \le K$. For the latter inequality use $\lambda \ge h^3 \ge 2Q$ so $K \ge 3\lambda/(2Q)$. The same bound on K and (53.3) imply

$$Kp \ge \lambda p/Q > b/Q > 2(G+\delta), \qquad Kp^2 \ge (3/2)\lambda p^2/Q > (3/2)b > b+\delta.$$

Indeed $\lambda p > b$ because $\lambda \ge 1$, and $\delta < \eta \ll G \ll b$. The inequalities $b/Q > 2(G+\delta)$ and $b/2 > \delta$ also follow directly from $b \ge 99G(h^3+1)$ and $Q \le h$. The two-sizes theorem gives exponent $10^4K(\lfloor \log_p(2b_*) \rfloor + 1) \le 10^4Km$, hence the same defect bound as before.

The remaining case $p > Q$ and $\lambda < h^3$. Now (53.3) gives

$$p > 2b/h^3 \ge 198G, \qquad a = 2(\lambda+1) \le 2h^3.$$

Use the few-low-jumps theorem proved in Section 52.13 (source `KummerChart.few_low_jumps`), with this a, η, W as originally given, and centre $\beta_i = b+\delta$ or $G+\delta$. The chart is unramified. Indeed

$$(p-3)a > 4b - 6h^3 > b+\delta+W+\eta, \qquad R_0 := 6(a+\eta+W) < 13h^3 < G.$$

Thus the size hypotheses $\beta_i + W + \eta \le (p-3)a$ and $6(a+\eta+W) < \beta_i$ hold. The no-multiple condition holds within radius R_0 of $0, \pm(G+\delta)$ because their absolute values there are $< 2G < p$. If a nonzero integer in radius R_0 of $\pm(b+\delta)$ were divisible by p, its absolute value would belong to $(b-30h^3, b+30h^3]$: use $\delta < \eta \le h^3/99$. By (53.2) this would give $\lambda \ge h^3$, impossible. This proves the exact no-multiple condition without an assumption about the density of large primes.

If the permitted set U' is in $[-\eta, 0]$, the zero-defect conclusion of the few-low theorem applies. Otherwise use the first jump alternative. When $\lambda < h/5$, we have $a < 2h/5 + 2 < h/2$, and therefore neither set has a jump in $[-a, 0)$; the other zero-defect conclusion applies. When $\lambda \ge h/5$, use the refined length assertion of the theorem:

$$\text{normLength} \le 11 f_0\big(\mu(a+1) + r_L(W+1)\big).$$

Here $f_0 = d$ because the chart is unramified. Since $\lambda \ge 1$, $a+1 = 2\lambda+3 \le 6\lambda$. Also $2\lambda/Q \ge 2h/(5Q) \ge (198/5)(W+1)$ by (53.1), so $W+1 \le K \le Km$. The bound is at most $11d(6\mu\lambda + r_L Km)$, which is again at most (53.4). This proves all the cases of `crystalline_defect_le`. In particular the floor K is kept in the conclusion; no replacement by an estimate with a hidden factor depending on Q is used.

53.2 Cyclotomic coordinates of thick places

We use precisely the data of Chapter 6. For clarity write R for the data rank $D.r$, $k = R-m+1$, $t = 5R$, $N_1 = N_0 3^c 2^s$, $c = \lfloor \log_3 s \rfloor + 5$, $K_0 = \mathbb{Q}(\sqrt{-d_K})$, where d_K is the prime parameter (not a local degree). The field $E = K_0(\zeta_{N_1})$ is contained in $\mathbb{Q}(\zeta_{N_1 d_K})$. Every automorphism fixing the latter roots of unity fixes E, which proves `fixes_emb_of_fixes_rootsOfUnity`. The infinite places, with chosen representatives restricting to the fixed embedding of K_0, have the multiplicative coordinate

$$\Phi_E \simeq (\mathbb{Z}/N_1\mathbb{Z})^\times$$

as in the data construction. Reductions of this coordinate will be denoted pc_u for $u \mid N_1$. If $\sigma : E \hookrightarrow \mathbb{C}$ is a complex embedding let $\varepsilon_\sigma = \pm 1$ be its sign on $\sqrt{-d_K}$. Its chosen infinite-place representative has ζ_{N_1}-value $(\sigma\zeta_{N_1})^{\varepsilon_\sigma}$.

Fix a p-adic embedding $e : E \hookrightarrow \overline{\mathbb{Q}}_p$ and an identification ι_p with the complex algebraic closure. The embedding indexed by $g \in G_p$ is $\sigma_g = \iota_p^{-1} g^{-1} e$. Suppose $g'g^{-1}$ acts on $e(\zeta_{N_1})^{p^a}$ by the power p^n. Comparing the chosen representatives at their places $\nu_g, \nu_{g'}$ gives, for every $u \mid N_1$ prime to p^a,

$$\mathrm{pc}_u(\nu_g/\nu_{g'}) = \varepsilon_{\sigma_g}\varepsilon_{\sigma_{g'}} p^n \pmod{u}. \tag{53.5}$$

In fact raising the equality of the chosen root values to p^a cancels the p-power part of their order, and applying $g'\iota_p$ makes the prescribed action of $g'g^{-1}$ visible. For an element of order u, equality of the powers implies equality of exponents modulo u; cancellation of p^a is permitted by the coprimality hypothesis. This is the assertion `placeClass_ratio`, including its signs and the inverse in the definition of σ_g.

The thick-place construction supplies these two properties:

1. every thick place has coordinate -1 modulo $N_0 3^c$ (and so modulo every divisor of it);
2. if ν, τ, ρ are thick and

$$\mathrm{pc}_{2^{s+1}}(\tau/\nu)^j = \mathrm{pc}_{2^{s+1}}(\rho/\nu)^i, \qquad 0 < i < j \le 324s,$$

then $\mathrm{pc}_{2^{s+1}}(\tau/\nu) = 1$.

They are the fields `thick_eq_neg_one_mod` and `three_term` of the admissible data, constructed earlier. Suppose three thick embeddings lie on a Frobenius arc with $0 < i < n \le 27s$, in the following explicit sense: the actions of $g'g^{-1}$ and $g'g_1^{-1}$ on the same prime-to-p part of a primitive root are p^n and p^i. We show this is impossible for $p \le kt$.

For $p = 2$, use (53.5) modulo 3^c. The ratio of any two thick places is 1 modulo 3^c; squaring the ratio implies $4^i \equiv 1 \bmod 3^c$. But for $i > 0$,

$$v_3(4^i - 1) = 1 + v_3(i)$$

by the elementary lifting-the-exponent identity (factor a power with exponent prime to 3, then cube, increasing valuation by exactly one). Hence $i \ge 3^{c-1} > 27s$. This contradicts the arc condition.

For odd p, use (53.5) modulo 2^{s+1}. The two ratios have respective values $\pm p^i, \pm p^n$, so their powers with exponents $2n, 2i$ are equal. As $2n \le 54s < 324s$, the three-term property implies the first ratio is 1. Thus $p^{2i} \equiv 1 \bmod 2^{s+1}$. For odd p the corresponding lifting identity reads

$$v_2(p^{2i} - 1) = v_2(p-1) + v_2(p+1) + v_2(i).$$

To verify it, $p^2 \equiv 1 \bmod 8$, so factoring an odd power does not change valuation, and squaring increases valuation by exactly one since the intermediate base is 1 mod 8. Because one of

$p-1, p+1$ has 2-valuation exactly one, the inequality $2^{s+1} \le 2(p+1)i$ would follow. In the actual data $(kt+1)27s < 2^s$: indeed putting $L = \log X$, we have $R \le 3e^{5L}$, $t = 5R$, $k \le R$, $s \le 32L$, and

$$2^s = m^{22} \ge e^{11L}, \qquad (kt+1)27s \le 46e^{10L}\, 864L < e^{11L}.$$

The last inequality holds for the immense lower bound on L in the data (in fact $L > 100$ suffices). This is the contradiction. We have thus proved precisely `not_thick_of_short_arc`.

Finally assume $p \mid N_1 d_K$. If $p \mid N_1$, it divides the norm $(d_K+1)/4$ of $(1+\sqrt{-d_K})/2$ by the elementary parameter calculation in the data chapter. The quadratic polynomial $X^2 - X + (d_K+1)/4$ has distinct roots $0, 1$ modulo p (also for $p = 2$), so both roots lie in $\mathbb{Q}_p$ by Hensel. Every $g \in G_p$ fixes $e(K_0)$, so every embedding σ_g has the same sign on K_0. If $p \nmid N_1$, then $p = d_K$. The root of order N_1 is prime-to-p; the action of $g'g^{-1}$ on it is a power of arithmetic Frobenius, thus a power p^n with $n \ge 0$ (choose a nonnegative representative of the finite cyclic quotient). Two thick places have ratio 1 modulo N_0. Were their signs opposite, (53.5) would imply $d_K^n = -1$ modulo N_0. But $d_K \equiv N_0/2 - 1 \bmod N_0$, and $4 \mid N_0$, so its square is 1 and neither it nor 1 is -1 mod N_0. This is impossible. Therefore all thick embeddings at the same p-adic place have the same sign whenever $p \mid N_1 d_K$. This proves `cmSign_eq_of_thick`.

53.3 The actual one-place estimate

We verify the hypotheses of Section 53.1 and of the tame estimate for the row and string blocks in `New/Selmer/OnePlaceDefect.lean`.

Let $h = D.h_2 = R^8$, so the gap is $2h$. Write b_1, b_2 for the rounded high weights of the thick and thin places, and $j \le m$ for the index in the row windows. The row type permits each jump u either with $h/2 \le |u| \le h^2$ or with $|\varepsilon_\sigma u + w_j(\sigma)| \le t$, where

$$w_j(\sigma) = \begin{cases} b_1 + 2jh + h, & \sigma \text{ thick}, \\ b_2 + 2jh + h, & \sigma \text{ thin}. \end{cases}$$

The other blocks are strings of a single jump $c \in \{0, \dots, t'-1\}$, where $t' \le t$. Their Tate dual jumps are $-c-1$. This uses the same Tate-twist sign convention as Section 8.4. Set

$$Q = kt, \quad b = b_1, \quad G = b_2, \quad \eta = h^2+1, \quad W = 4t, \quad \delta = 2jh + h - 2t. \tag{53.6}$$

Here $h > 2t$ so δ is a nonnegative integer. All original and Tate-dual permitted jumps lie in the opposite windows of signs $-\varepsilon_\sigma, \varepsilon_\sigma$ with the low radius and two high centres of (53.6). Indeed the dual replaces a jump u by $-u-1$, so even after this shift a jump within t of a high centre w_j remains in the interval starting $w_j - 2t$ and of width $4t$; the original low jumps and every string jump are within the low radius. Any negative original or dual jump in either window which is not a string jump satisfies $u \le -h/2$. For the row dual, the number of jumps of magnitude $\le h^2+1$ is no greater than the number of original row jumps of magnitude $\le 2h^2$, so by the row hypothesis it is at most $\lfloor r_{\mathrm{row}} m/k \rfloor$. For the Tate-dual string all jumps belong to $[-\eta, 0]$. Since the *other* side of a row pairing may contain string jumps, the negative-jump hypothesis used for the row in Section 53.1 must be checked in the actual paired class: string jumps on that other side are nonnegative by the stipulated choice of original (not Tate-dual) permitted windows. Thus the row case takes the first jump alternative, the string case the second.

Let $x = 1$ for the row and $x = t$ for a string, and denote by r_L the rank of the block. We may take $\mu = \lfloor r_L m/(kx) \rfloor$ (for the string case the low-multiplicity bound is not needed). The

desired output is

$$\begin{aligned}\operatorname{normLength}_O(Z_f(L^\vee(1))/(Z_f(L^\vee(1))\cap \operatorname{cupOrthogonal})) &\le d\, r_L\, \frac{\operatorname{loss}(p)}{x},\\ \operatorname{loss}(p) &= 10^6\frac{m}{k}(b_1 v_p(g_0 g_1)+\lambda).\end{aligned} \tag{53.7}$$

where d is the local degree and λ is (53.2) at h, b_1, p.

The scale inequalities. Here are explicit coarse comparisons which justify the uses of the logarithmic magnitude calculus in the source. Put $L_0 = \log X$, $A = \log B$, and $A_1 = \log X_1$. From admissibility,

$$\log A \le L_0/100, \quad A_1 \ge 10^6 L_0, \quad A \ge 10^6 A_1, \qquad m \in [e^{L_0/2}, 2e^{L_0/2}], \quad R \in [e^{5L_0}, 3e^{5L_0}].$$

All the following strict inequalities hold, for example, for $L_0 \ge 10^6$, which is much weaker than the data's lower bound $L_0 \ge 100 \log A \ge 10^{14} P.c$:

$$\begin{gathered} R/2 \le k \le R, \qquad Q = kt \le 5R^2, \quad W = 20R,\\ \eta = R^{16}+1, \quad 0 \le \delta \le (2m+1)h < h^2+1,\\ 99(Q^2+(W+1)Q+Q+W+m) < R^8,\\ 99((\eta+W)Q+\eta+h) < R^{24}.\end{gathered}$$

For the second line, its left side is bounded by $99(25R^4+110R^3+105R^2+21R)$ using $m \le R$; this is $< R^8$ for $R \ge 100$, and the actual R is far larger. The third line follows, for instance, from $(\eta+W)Q+\eta+h \le 10R^{18}+3R^{16}$ for $R \ge 100$, and $99(10R^{18}+3R^{16}) < R^{24}$. The root order $N_1 \le e^{12L_0}$ follows from $N_0 \le P.c \le L_0$, $3^c \le 243s$, $s \le 32L_0$ and $2^s = m^{22} \le 2^{22}e^{11L_0}$. Hence $[E:\mathbb{Q}] \le 2N_1 < h$, giving $e_0 \le h$.

Rounding to a multiple of N_1 adds at most N_1 at each step. From $a_{\text{data}} = \operatorname{roundUp}(XB)$,

$$e^A \le b_1 \le 2e^A, \qquad e^{A-A_1+L_0} \le b_2 \le 2e^{A-A_1+L_0}.$$

Indeed each round-up error, including the divided first error, is at most $2N_1$ altogether, less than either unrounded base. Thus $99h^3 \le b_2$ and $99b_2(h^3+1) \le b_1$: use $h^3 = R^{24} \le 3^{24}e^{120L_0}$ and $A_1 - L_0 \ge (10^6-1)L_0$. Finally $2\log(4b_1) \le 2(A+\log 8) < m$ since $A \le e^{L_0/100}$ and $m \ge e^{L_0/2}$. These are precisely (53.1), including its logarithmic bound. All comparisons have a large margin, so the use of integer floors and ceilings is legitimate.

The letter pattern. If $p \nmid N_1 d_K$, the local field is unramified and $p \ge 7$ (all smaller primes divide N_1). A local p-adic automorphism fixing the residue-field Teichmüller generator fixes the entire unramified completion of E. Thus two embeddings with the same letter give the same σ_g, and thickness and sign are well-defined functions of the letter. If $p \mid N_1 d_K$, then $p \le \max(N_0, d_K) \le L_0 < Q$. Mark a letter thick whenever it is the letter of *some* thick embedding. The sign on all such embeddings is the single sign just proved in Section 53.2; assign that sign to every letter (choose either if no thick embedding occurs). This is exactly the weaker letter-compatibility hypothesis for $p \le Q$, and the small-prime widening of the low radius handles thin embeddings on those letters.

If $p \le Q$, the no-three condition holds. The Frobenius action on the letters is raising residue-field Teichmüller roots to the p-th power. If two embeddings' letters have relation $F(g') = F(g)^{p^u}$, then $g'g^{-1}$ acts by the power p^u on the unramified subfield of degree f_0 of the completion; equality on the primitive $(p^{f_0}-1)$st root suffices. In particular it acts by that

power on the prime-to-p part of $e(\zeta_{N_1})$, which lies in this subfield. Three thick letters with two gaps in $[1, \lfloor \log_p h^3 \rfloor]$ would therefore give three thick embeddings on the arc of Section 53.2. Its bound $n \le 27s$ follows from

$$h^3 = R^{24} \le 3^{24} e^{120L_0} < e^{132L_0} \le (2^s)^{12}.$$

Consequently the sum of the two gaps is at most $24s$. This is the contradiction of the spacing lemma. Thus all the letter hypotheses of Section 53.1 are met.

The ordinary crystalline case. Choose the actual Kummer chart for the local field as in Section 52.5, with $e_0 f_0 = d$. A sufficiently large coefficient ring contains the primitive Teichmüller root needed for all letters; this is the only eventual enlargement of the coefficient ring in this application. Apply (53.4) to the dual of the block, using the chosen permitted windows, their signs, and the row or string jump alternative above. Since $x \le t$, $K \le 2\lambda/(kt)$ and $\mu \le r_L m/(kx)$, we get

$$10^5(r_L K m + \mu\lambda) \le 3 \cdot 10^5 \, \frac{r_L m}{kx}\lambda \le r_L \, \frac{\mathrm{loss}(p)}{x}.$$

This is exactly (53.7); the floor from (53.2) was used only in the favorable direction.

The special tame primes. If $p = g_i$ is one of the two marked primes, $p \equiv 1 \bmod N_1 d_K$, so it splits completely in E. The completion is $\mathbb{Q}_p$ and the embedding and sign of E are fixed under G_p. All allowed jumps therefore use one pair of opposite windows. The tame type of the source has rank parameter R and degree parameter $N = (R+t)R = 6R^2$; precisely this type is an assumption of the one-place estimate. Use the tame exponent of Section 52.13,

$$k_{\mathrm{tame}} = 3R^2(N+1)^2 + 150(\eta + W + 1).$$

Its size hypothesis is satisfied because $p \ge \exp(A^{10})$ and

$$b_1 + \delta + 6(W+\eta) + R^2(N+1)^2 + 2 \le 2e^A + e^{100L_0} < \exp(A^{10}).$$

Also $2k_{\mathrm{tame}} \le e^{100L_0}$ by the displayed values of R, t, h, η, W, N. On the other hand $v_p(g_0 g_1) \ge 1$ and

$$\mathrm{loss}(p)/x \ge 10^6 \frac{m b_1}{kt} \ge 10^6 e^{A+L_0/2}/(45e^{10L_0}) > e^{100L_0}.$$

The rank bound on crystalline cocycles now gives (53.7), using $p^{k_{\mathrm{tame}}}$ on the entire module. No division by the local degree or an unchecked restriction of the tame lattice is necessary; this special completion has local degree one. The polynomial degree and the crystalline type are precisely the hypotheses supplied to the one-place theorem, not inferred from unramified spherical traces.

53.4 Status of the local interface

The precise numerical bound `crystalline_defect_le`, the three arithmetic spacing assertions of `ThickSpacing.lean`, and the assembly of the actual row and string local bounds in `block_defect_le_at_place` have the written derivations above. The latter assumes, as does its source statement, an actual local lattice with the asserted Hodge jump and crystalline or tame type data (and the common permitted jump sets). The arithmetic supply of those local hypotheses for the global automorphic family belongs to the attachment, local–global compatibility and congruence-lattice constructions. This is not an assertion that every lattice with the stated characteristic polynomials is crystalline. The local bound is then the one used in the already proved sum of the finite-prime losses.

Part V

The arithmetic conclusion and proof audit

Chapter 54

A numerical appendix: asymptotic budgets and the passage to the limit

The compact development checks inequalities on the logarithmic scale for every record satisfying its very large fixed admissibility bounds. A conventional paper need not insist on the same finite threshold for the parameters: to conclude GRH only the existence of admissible data for all sufficiently large x is needed. This chapter collects the estimates along that family, including the flexibility of the second cutoff. It gives ordinary proofs of several numerical comparisons implicit in the use of the formal budgets. No weakening of a non-numerical hypothesis on an automorphic object is intended. In particular a theorem whose proof depends on a precise integral divisor bound cannot be replaced by a numerical big-O estimate of the size of its coefficient matrices.

54.1 Uniform scale asymptotics

Fix N_0, d, and $0 < \varepsilon < 1/100$. We use the family of admissible records of Proposition 4.2 after replacing its second logarithmic cutoff by $u \in [10^6 \log x, (10^6 \log x)^2]$. Put $Y = \log b = x^{\varepsilon}$. In particular $u = o(Y^{\delta})$ for every fixed $\delta > 0$ (uniformly in the indicated window). Notice the identity $2^s = m^{22}$ and the estimates

$$\sqrt{x} \le m \le 2\sqrt{x}, \qquad x^5 \le r \le 3x^5, \qquad k = r + 1 - m, \qquad t = 5r,$$
$$81s < 3^{c_s} \le 243s, \qquad s \asymp \log x, \qquad T_s \asymp x^{11} \log x, \qquad f = \varphi(N_0)T_s.$$

Here $T_s = N_1/N_0$ and $c_s = \lfloor \log_3 s \rfloor + 5$. In particular $x^{11} \le T_s \le x^{11}Y^{1/5}$ eventually. The rounding $a = N_1\lceil xb/N_1 \rceil$ gives $a = xb + O(x^{11} \log x)$, and hence $a \sim xb$. All the estimates in this chapter are uniform in the second cutoff within its window. They also remain valid with the initial choice $\log x_1 = x^{\varepsilon/6}$ for the power-dominance estimates below; closeness of its logarithm to the window is not assumed until we choose the cutoff.

Lemma 54.1 (Power dominance)**.** *For every fixed $A, C, D \ge 0$ and $\eta > 0$, along the family just described,*

$$x^A Y^C x_1^D = o(b^{\eta}).$$

If $P(x)$ is bounded by a fixed power of x, then $P(x) = o(B_0)$, where $B_0 = B/(mk)$.

Proof. The logarithm of the quotient in the first assertion is at most

$$A \log x + C \log Y + D(10^6 \log x)^2 - \eta x^{\varepsilon} \longrightarrow -\infty.$$

If we retain the initial cutoff instead, the third term is $Dx^{\varepsilon/6} = o(x^{\varepsilon})$, so the same proof applies. From $a \ge xb$, $f \ge x^{11}$, $m \le 2\sqrt{x}$, and $k \le 3x^5$ we get $B_0 \ge bx^{13/2}Y^3/6$. Thus the second assertion follows from the first with $D = 0$ and any positive $\eta < 1$. □

This lemma must not be confused with a comparison between powers of x and powers of Y: given ε, a fixed power of x can be much larger than Y. For example, one cannot pay for a factor m merely by a fixed power of Y uniformly for all small ε. The factor m is only harmless when divided by a quantity containing a corresponding power of x or by a positive power of b. This observation explains some of the delicate bookkeeping in the local Hecke repairs and crystalline loss estimates. The three-term avoidance property of the thick places is not an artifact of this elementary dominance calculus.

54.2 Rounding of large coefficients and a Lipschitz estimate

Recall the thick and thin rounded weights $b_1 = N_1\lceil a/(xN_1)\rceil$ and $b_2 = N_1\lceil a/(x_1N_1)\rceil$. Then

$$b_1 = a/x + O(N_1) = b + O(N_1), \qquad b_2 = a/x_1 + O(N_1).$$

Since $N_1, h_2 = r^8$ and m are bounded by fixed powers of x, and b/x_1 grows faster than every fixed power of x, the rounding and gap terms satisfy, uniformly in $0 \le j \le 2m$,

$$0 \le w_j(\tau) - w_{\mathrm{nom}}(\tau) \le N_1 + (4m+1)h_2 = O(x^{81/2}) \le x^{42}.$$

Thus $w_j \le 2b$ eventually and

$$\log w_j - \log w_{\mathrm{nom}} \le x^{42}/w_{\mathrm{nom}} \le x^{-1}.$$

This proves the numerical comparisons of `New/Data/Weights.lean` along our family. We use the following useful estimate for small changes in the weights of a finite prime sum. For two families of real weights w, w' set $A_w(y,\sigma) = \sum_{\nu\le a}\lambda_{d,1}(\nu)(y/\nu - w([-\nu]\sigma))_+$. If $|w - w'| \le \Delta$ and $w, w' \ge y/Z$ pointwise, then

$$|A_w(y,\sigma) - A_{w'}(y,\sigma)| \le \Delta \sum_{\nu\le \min(a,Z)} \lambda_{d,1}(\nu) \le 2\Delta \sum_{\nu\le Z} \log\nu \le 2\Delta Z\log(1+Z).$$

Indeed the positive-part function is 1-Lipschitz and each summand vanishes for $\nu > Z$. The same reasoning at a single prime gives

$$|A_{p,w}(y,\sigma) - A_{p,w'}(y,\sigma)| \le \omega_p\Delta\lfloor \log_p Z\rfloor.$$

The latter is more useful when the bounds need to be summed over local loss functions, because it does not multiply the error by Z. In the case at hand one may take $Z = x_1$ and $\Delta = x^{42}$ for $y \le a$ after increasing the weights. Crude errors polynomial in x times x_1 are absorbed by Lemma 54.1. They must be absorbed before concluding a statement at the prime-power scale: replacing w_j by w_{nom} silently inside a coefficient representation would change the representation.

54.3 The exact level cancellation revisited

In the notation of Chapter 4, let $\bar w = f^{-1}\sum_\tau \log w_{\mathrm{nom}}(\tau)$, $G = \log(g\,\mathrm{rd}_E/(2\pi))$, and $\kappa_K = \gamma - \Re L'(1,\chi_K)/L(1,\chi_K)$. Then

$$\log(\Omega/g) = \kappa_K - G, \quad J(a) = \log(a!) - a\kappa_K + aG, \quad A_0 = a(\log a - 1 - \kappa_K - \bar w).$$

The only large difference between these two quantities is

$$J(a) - A_0 = a(G + \bar{w}) + O(\log(a+1)),$$

by Stirling's formula, or by comparing the sum $\sum_{i=1}^{a} \log i$ to its integral. The level prime is chosen so that

$$\log N \geq m(G + \bar{w}) - G - \tfrac{15}{2}Y.$$

Consequently

$$\begin{aligned} |\mathcal{T}|(m(J(a) - A_0) - a\log N) &\leq |\mathcal{T}|(a(G + \tfrac{15}{2}Y) + O(m\log(a+1))) \\ &= O(faY). \end{aligned}$$

We used $G = O(Y)$, $m/a \to 0$, and $\log a = O(Y)$. The exact source allows the larger error $B = faY^3$; the displayed upper bound is eventually smaller than that allowance. Thus the heights being compared can contain a very large common term H, and the cutoff-dependent excess at the distinguished columns is paid by the rational logarithm of N. Its choice is possible precisely because λ_N lies in the range of logarithmic prime windows in Proposition 4.2. It is not necessary to have a strong pointwise estimate for the prime error at x_1 for this prime choice.

54.4 An elementary bound on the logarithm of the class number

For the global Selmer bounds it is useful to avoid treating the class number as an unspecified enormous constant depending on the growing field E. Let $D_E = [E : \mathbb{Q}] = 2f$ and let Δ_E be the absolute discriminant. Minkowski's ideal class theorem says that every ideal class has an integral representative of absolute norm at most $(4/\pi)^f D_E!/D_E^{D_E}\sqrt{|\Delta_E|}$ (see the geometry-of-numbers treatment of number fields, for instance [11]). For a crude upper bound it suffices to use

$$M_E = 2^{D_E}\sqrt{|\Delta_E|}.$$

For each rational prime p there are at most D_E primes of E above it; fix an arbitrary numbering of these for each p. Map an integral ideal I, supported on these primes with exponents $e_{p,i}$ and residue degrees $f_{p,i}$, to the ordered tuple of positive integers

$$(n_1, \dots, n_{D_E}), \qquad n_i = \prod_p p^{f_{p,i}e_{p,i}}.$$

Fill missing indices with exponent zero. The tuple determines every ideal exponent by ordinary prime factorization, and $\prod_i n_i = N(I)$. Hence the number of integral ideals of norm at most M is at most $(\lfloor M \rfloor)^{D_E}$. In particular

$$\log h_E \leq D_E \log M_E = D_E^2\left(\log 2 + \tfrac{1}{2}\log \mathrm{rd}_E\right).$$

Using $f = O(x^{11}\log x)$ and $\mathrm{rd}_E \leq x^{12}$, we get $\log h_E = O(x^{22}(\log x)^3) \leq x^{23}$ for sufficiently large x. This estimate is far from optimal but adequate for the global budget. It is compatible with the bound used in `exists_isFitFamily`. Notice the difference between a bound for h_E and a bound for its logarithm; the latter is what appears when valuations are summed with log-prime weights.

54.5 Summation of the fit-family costs

We use the same notations as Chapters 40 and 49. Let M_g be the integer clearing all chosen-level group orders, and let X be the integer controlling the odd Tate classes (from the formal-source input concerning those classes). Its size is bounded in that input by

$$\log X \le 4(N_E + t + 3)^{27} \log 2 + \log h_E,$$

where $N_E = N_1 d$ denotes the cyclotomic conductor used in the Selmer calculation. In particular it is bounded by a fixed power of x. We require only this bound, not a proof of the existence of the odd-class annihilator here. Corollary 5.3 gives $\log M_g \le x^{32}$ eventually. Write

$$c_\ell = v_\ell(M_g) + 2^{-\ell}, \qquad D_\ell = h_2^{27}(c_\ell + v_\ell(X)) + 4f \operatorname{loss}_\ell,$$

with loss_ℓ as defined in the crystalline chapter. The series $\sum_{n\ge 2} 2^{-n} \log n$ converges, since $\log n \le n$. By ordinary factorization, for every finite set of primes T,

$$\sum_{\ell \in T} 9 c_\ell \log \ell \le 9(x^{32} + O(1)) \le x^{999}$$

eventually. This establishes the global inverse-ratio cost without any estimate depending on the particular chosen twists.

For the other budget the part

$$h_2^{27}\big(\log M_g + \log X + O(1)\big)$$

is bounded by a fixed power of x, since $h_2 = r^8$ and $\log X$ has the preceding bound. Hence it is $o(B_0)$ by Lemma 54.1. The local-loss part is calculated in Chapter 49. For clarity, let

$$J_* = (2b_1)! \left(\prod_{\max(b_1 - 30h_2^3, 0) < i \le b_1 + 30h_2^3} i \right)^{h_2^3}.$$

The actual local formula is $\operatorname{loss}_\ell = 10^6 (m/k)(b_1 v_\ell(g) + v_\ell(J_*))$. Summing over any finite set of primes gives

$$\sum_{\ell \in T} 4f \operatorname{loss}_\ell \log \ell \le 4 \cdot 10^6 \frac{fm}{k} (b_1 \log g + \log J_*).$$

Now $\log J_* \le 2b_1 \log(2b_1) + 60 h_2^6 \log(b_1 + 30h_2^3) = O(bY)$ by the elementary factorial bound and Lemma 54.1. Thus the preceding expression is $O(fmbY/k)$, and

$$\frac{fmbY/k}{B_0} = \frac{m^2 b}{aY^2} \le \frac{4}{Y^2} \longrightarrow 0.$$

Combining the two pieces yields $\sum_{\ell \in T} D_\ell \log \ell \le B_0$ for any finite set T and all sufficiently large x. In particular the inequalities called `Data.le_of_hasLogBound` in the summation of `exists_isFitFamily` can here be obtained from simple asymptotic inequalities. It is the proof of the block loss and the construction of the odd-class integer X, rather than this last summation, that remain the crucial arithmetic obligations.

54.6 The final numerical inequalities

Two further absorptions used in the reconstruction can be verified without the formal logarithmic size calculus. The absolute size estimate $|H| + m|\mathcal{R}| \le b^2 x^{99}$ follows from the definitions and the crude $\lambda_{d,1}(\nu) \le 2\nu$; its proof is given in Lemma 55.4. The main combination in Proposition 55.5 gives

$$m\mathcal{R} \le 4L + 4x^{99} + 10\log(1+b^2)B.$$

Divide the right side by BY^2. The first and third summands after division are $4/Y$ and $10(2Y+o(1))/Y^2$, both tending to zero; the middle summand tends to zero by Lemma 54.1. In particular the sum is eventually at most one. Similarly the single-lift allowance in Chapter 31,

$$3nrB_0 + n\big(10mx^{999} + 6kB_0 - k\mathcal{U}\big) \le nk(YB_0 - \mathcal{U}),$$

is equivalent after cancellation and division by nkB_0 to $3r/k + 6 + 10mx^{999}/(kB_0) \le Y$, which holds eventually. Here $r/k \to 1$ and the last quotient tends to zero. This calculation does not require positivity of $\mathcal{U}$.

Finally $|\mathcal{S}| \ge T_s/Y$, so the major arithmetic error after division by $ma|\mathcal{S}|$ is bounded by $\varphi(N_0)Y^6/m \le \varphi(N_0)x^{-1/2+6\varepsilon}$. The window error is x^{-10^6}. The fixed constant $\varphi(N_0)$ can be absorbed by $x^{2\varepsilon}$ for large x, giving $x^{-1/2+8\varepsilon}$. This demonstrates explicitly why the slack in each numerical budget does not affect the target exponent and why the number of lifts is the one important scale that must be retained. All thresholds may depend on the fixed auxiliary modulus and field. The zero-free criterion chooses the auxiliary prime after the hypothetical zero and then uses the eventual bound for that fixed prime, so no uniformity in d has been assumed.

Chapter 55

The arithmetic inequality and conclusion

We assemble the three arithmetic results in their exact quantitative form. The proof of their consequence is independent of the internal choices of kernels.

55.1 Collecting the arithmetic inputs

With the data of Chapter 4, set $n = \dim_{\mathbb{C}} W$, $c = \dim_{\mathbb{C}} C$, and retain

$$H = mk\mathcal{U}, \qquad B = fa(\log b)^3, \qquad L = B \log b.$$

For clarity, we give the particular form of the three theorems that is needed in the calculation. The notation in the three statements below is intentionally the same. The preceding sections give their derivation and identify specialized inputs still requiring independent written proofs.

Theorem 55.1 (A: Short forms, recalled)**.** *There are a rational number $q_0 > 0$ and a set $S \subseteq C^{\mathrm{ram}}$ of integral forms spanning C^{ram} over $\mathbb{C}$ such that, putting $T_s = q_0 P_s$, one has*

$$\|\Theta_1 T_s(v)\|_\sigma^2 \le \exp(H - m\mathcal{R} + L) \quad (v \in S,\ s \in \mathcal{G},\ \sigma \in \mathrm{Aut}(\mathbb{C})).$$

For every finite set $F \subseteq \mathcal{G}$ and every finite family y in the images $T_s(C^{\mathrm{ram}}_{\mathrm{int}})$, $s \in F$, one has

$$\mathrm{IndexLe}(y, |F| cB).$$

Theorem 55.2 (B: Enough forms, recalled)**.** *The spaces C and W have positive dimension and there is a finite set $F \subseteq \mathcal{G}$ such that*

$$|F| c \le 5n \log(1 + b^2), \qquad \dim\left(\sum_{s \in F} \mathrm{im}(P_s)\right) \ge (1 - 4/b^2)n.$$

Theorem 55.3 (C: Integral forms are long, recalled)**.** *Let $y = (y_1, \dots, y_q)$ be q independent row forms with $q \ge (1 - 4/b^2)n$. If $\mathrm{IndexLe}(y, d_{\mathrm{ind}})$ and for every σ one has $\mathrm{covol}_\sigma^2(y) \le e^V$, then*

$$n(H - 3L) \le V + 2d_{\mathrm{ind}}.$$

Together with the crude bound

$$|H| + m|\mathcal{R}| \le b^2 x^{99}, \tag{55.1}$$

these are precisely the inputs in `star_inequality` of `New/Main.lean`. The source proves the latter as `Data.abs_mainHeight_add_remainder_le`. We give the elementary verification along the admissible asymptotic family.

Lemma 55.4 (Crude size comparison)**.** *Along the data in Proposition 4.2, including the chosen second cutoff, inequality (55.1) holds for all sufficiently large x.*

Proof. The factor $\lambda_{d,1}(\nu)$ is bounded by 2ν. Thus $0 \le A(z,\sigma) \le 2az$ for $0 \le z \le a$, since each summand is at most $2z$. From $w_j \le 2b$, $|\log(\Omega/g)| \ll_{N_0,d} \log b$ (by (4.2) and the bounds on g), and $\log(j!) \le j \log j$, we have $|J(w_0(\sigma))| \ll b \log b$. The nominal weights range between a/x_1 and a/x, so $|A_0| \ll a \log b$. From the definitions,

$$|H| \ll mkf(b \log b + ab), \qquad m|\mathcal{R}| \ll mf(a^2 + a \log b).$$

Here $a \le 2xb$, $k \ll x^5$, $f \ll_{N_0} x^{11} \log x$, $m \ll x^{1/2}$, and $\log b = x^{\varepsilon} \le x$. These upper bounds are $O_{N_0,d}(b^2 x^{20})$ for large x. Enlarging the threshold gives (55.1). The proof in the compact source instead uses explicit logarithmic inequalities to have the same bound on the full admissible record, a stronger quantification not needed for the eventual inequality here. □

55.2 From A and B to an almost-basis of short forms

Choose F as in Theorem 55.2 and use the set S from Theorem 55.1. Since q_0 is nonzero and S spans C^{ram}, the set

$$\{T_s(v) : s \in F,\ v \in S\}$$

spans the sum of the images of the P_s, $s \in F$. Choose independent vectors $y_1, \dots, y_q$ from this set which form a basis of its span. In particular,

$$(1 - 4/b^2)n \le q \le n.$$

By A and the bound on the number of translates we may take

$$d_{\mathrm{ind}} \le 5n \log(1 + b^2)B.$$

For each conjugated pairing Hadamard's inequality gives

$$\begin{aligned} \mathrm{covol}_\sigma^2(y) &\le \prod_{i=1}^{q} \|\Theta_1 y_i\|_\sigma^2 \\ &\le \exp\big(q(H - m\mathcal{R} + L)\big). \end{aligned}$$

Thus $V = q(H - m\mathcal{R} + L)$ is a valid upper bound. This is the conclusion of `Data.exists_family_covolumeSq_le` in `New/Endgame/LongForms.lean`. The step is only linear algebra, but realizes the crucial purpose of Theorem B: it converts estimates on vectors arising from a particular automorphic construction into an estimate on almost the entire determinant of the row space.

55.3 Comparison with the lower bound

Proposition 55.5 (The main arithmetic inequality)**.** *For all sufficiently large x along the admissible family of Proposition 4.2, the three arithmetic inputs imply*

$$m\mathcal{R} \le fa(\log b)^5. \tag{$\star$}$$

Proof. Write $h = H - m\mathcal{R} + L$ and apply C to the family just constructed. Then

$$\begin{aligned} n(H - 3L) &\le qh + 2d_{\mathrm{ind}} \\ &= nh + (n - q)(-h) + 2d_{\mathrm{ind}}. \end{aligned}$$

We cannot simply replace q by n in qh, because h may be negative. Instead, use (55.1) and $L > 0$ to get $-h \le b^2 x^{99}$, and the dimension estimate to get $n - q \le 4n/b^2$. Consequently

$$n(H - 3L) \le n(H - m\mathcal{R} + L) + 4nx^{99} + 10n \log(1 + b^2)B.$$

Divide by $n > 0$, cancel the common H, and rearrange:

$$m\mathcal{R} \le 4L + 4x^{99} + 10 \log(1 + b^2)B. \tag{55.2}$$

The scale inequalities imply that the right-hand side is at most $(\log b)^2 B$. For the eventual assertion this is immediate from $a \ge xb$, $B = fa(\log b)^3$, and $\log b = x^{\varepsilon}$, since $x^{99}/B \to 0$ and $\log(1 + b^2) = O(\log b)$. The Lean version uses its explicit admissibility bounds to make this last comparison for every admissible datum, not just sufficiently large x along this family. Substituting the value of B gives ($\star$). □

It is worth looking closely at the size of the three errors in (55.2). The $4L$ is the sum of the error per vector in the upper bound and the three units of L in the lower bound. The $4x^{99}$ is the penalty for the missing dimensions, which are so few that even a very crude bound on h suffices. The last term is twice the index. All are smaller than the final budget by a factor of order $\log b$ or more along the asymptotic family. The leading term H has disappeared completely. Without the matching of its exact leading coefficient in A and C, this argument would not give the desired bound.

55.4 Extracting the analytic remainder

Here is the short algebra behind (4.9). Use the notation of Chapter 2 and write

$$S_M(z) = \sum_{\nu \le z} \lambda_{d,M}(\nu)(1/\nu - 1/z).$$

For a fixed integer ν, the number of $\sigma \in \mathcal{T}$ such that the place $s_\nu \sigma$ lies in $\mathcal{S}$ is $|\mathcal{S}|$ if $\nu \equiv 1 \pmod{N_0}$ and zero otherwise. Indeed, multiplication by s_ν carries the whole coset $\mathcal{T}$ onto the coset of class $-\nu$ when ν is invertible modulo N_1. The thick places have class -1 modulo N_0. Every $\nu \equiv 1 \pmod{N_0}$ is a unit modulo N_1 in this tower. In the remaining nonunit cases the source's place convention places the product back in $\mathcal{T}$, so it is not thick. It follows that replacing the weight a/x_1 by a/x inside the sum over the tops gives the identity

$$\frac{1}{a} \sum_{\sigma \in \mathcal{T}} A(a, \sigma) = |\mathcal{S}|\big(S_{N_0}(x) - S_{N_0}(x_1)\big) + |\mathcal{T}| S_1(x_1).$$

Here S_M denotes the sum before subtraction of its main terms. By the analytic normalization,

$$S_M(z) = R_M(z) + \frac{\log z}{\varphi(M)} + c_{d,M}.$$

The constants for $M = N_0$ cancel in the difference. For $M = 1$, we have $c_{d,1} = -1 - \kappa_K$. Moreover

$$\log a - \overline{w} = \log x_1 + \frac{|\mathcal{S}|}{f}(\log x - \log x_1).$$

Since $f = \varphi(N_0)|\mathcal{T}|$, we have exactly

$$\frac{|\mathcal{T}|A_0}{a} = |\mathcal{T}|(\log x_1 - 1 - \kappa_K) + \frac{|\mathcal{S}|}{\varphi(N_0)}(\log x - \log x_1).$$

Subtract these main terms from the previous sum identity. This proves (4.9), including its constants. In particular, no estimate involving the Chowla–Selberg constant is needed at this final stage.

55.5 Choosing the second cutoff and the power saving

Fix $0 < \varepsilon < 1/100$ and choose admissible data by Proposition 4.2 for every sufficiently large x. The quantity $X = |\mathcal{T}|/|\mathcal{S}|$ satisfies $0 < X \le \log b$. Write $T = 10^6 \log x$. For large x we have $X \le e^T$. Apply the window lemma of Chapter 2 to the two functions $u \mapsto R_{N_0}(e^u)$ and $u \mapsto R_1(e^u)$. It supplies $u \in [T, T^2]$ such that

$$R_{N_0}(e^u) - \frac{|\mathcal{T}|}{|\mathcal{S}|} R_1(e^u) \le e^{-T}.$$

Take $x_1 = e^u$ in the auxiliary data, as permitted by Proposition 4.2. This is the decisive use of the windowed argument: there is no need for a strong pointwise estimate at the initially specified, extremely large, second scale.

Solve (4.9) for the desired term and use ($\star$):

$$\begin{aligned} R_{N_0}(x) &= \frac{\mathcal{R}}{a|\mathcal{S}|} + R_{N_0}(x_1) - \frac{|\mathcal{T}|}{|\mathcal{S}|} R_1(x_1) \\ &\le \frac{f(\log b)^5}{m|\mathcal{S}|} + e^{-T} \\ &\le \frac{\varphi(N_0)(\log b)^6}{m} + x^{-10^6} \\ &\le \varphi(N_0) x^{-1/2+6\varepsilon} + x^{-10^6}. \end{aligned}$$

Absorbing the fixed constant using 2ε of additional slack gives, for all sufficiently large x,

$$R_{N_0}(x) \le x^{-1/2+8\varepsilon}. \tag{55.3}$$

This is the precise exponent in the compact development's `eventually_primeError_le_rpow`. Replacing ε by a smaller positive number proves a bound with any fixed exponent greater than $-1/2$.

Remark 55.6 (Where the critical line enters). Everything up to ($\star$) concerns norms, divisibilities, dimensions and heights. The saving $x^{-1/2}$ arises from dividing the error by $m \ge \sqrt{x}$. The large field degree, large weights and rank do not themselves account for it. One cannot hope to improve the analytic conclusion by choosing m arbitrarily larger: the arithmetic theorems and their errors depend crucially on the admissible relationship between the length, the weights, ranks and local losses. A proof of stronger short/long-form estimates on other ranges would be an additional mathematical result.

55.6 GRH

To finish, suppose a Dirichlet character χ modulo q had a zero ρ to the right of $1/2$, excluding the pole convention of Theorem 1.1. Choose $N_0 = 18480q$ and regard χ as a character of level

N_0 by induction; the induced L-function differs by finite Euler factors with no zeros in this region. The class $N_0/2 - 1$ is a unit modulo N_0 and forces $d \equiv 7 \pmod 8$. For every prime d in this class, the above arithmetic argument gives (55.3). Choose ε positive and sufficiently small that $\beta = 1/2 + 8\varepsilon < \Re(\rho)$. The one-sided criterion of Proposition 2.9 forbids the zero ρ. This proves the deduction of Theorem 1.1 from the arithmetic inputs; a complete conventional proof requires independently establishing the remaining formal-source inputs listed in the final section. The standard functional equations imply the statements about the critical strip. This completes the deduction.

Chapter 56

Glossary and concordance

The following glossary is intended for returning to one chapter after reading another. Every symbol has a local role; the numerical estimates only acquire their meaning after the objects and all normalization factors have been fixed. We list central Lean declarations for the compact edition. Where the same symbol is used in a different context within a local calculation (for example the rank of a single lattice), the text specifies the local meaning.

Symbol or term	Meaning	Source or chapter
N_0	Fixed auxiliary modulus divisible by 18480; chosen to be a multiple of the given Dirichlet modulus.	`Parameters.N₀`; 4
d	Auxiliary prime $\equiv N_0/2 - 1 \pmod{N_0}$; varied in the zero-free criterion.	`Parameters.d`; 2
K	Imaginary quadratic field $\mathbb{Q}(\sqrt{-d})$.	`Parameters.K`
E/F	CM extension $E = K(\zeta_{N_1})$, $F = E^+$; field changes with the scale.	`Data.E`; 4
Φ, f	CM-type embeddings over K, and their number $f = [E : \mathbb{Q}]/2$.	`Data.Φ`, `Data.f`
s, T_s, N_1	Tower exponent, $T_s = 3^{\lfloor \log_3 s \rfloor + 5} 2^s$, $N_1 = N_0 T_s$.	`towerDeg`, `Data.N₁`
$\mathcal{T}, \mathcal{S}$	Tops of cyclotomic class 1 modulo N_0; thick places in class -1 with a multiple-progression avoidance property.	`Data.tops`, `Data.thick`
x, x_1, b	Prime error at x, auxiliary second cutoff, much larger order scale b with $\log b = x^{\varepsilon}$.	`Data.x`, `Data.x₁`, `Data.b`
m, r, k, t	Number of lifts, row rank, column rank $k = r + 1 - m$, row increment $t = 5r$.	`Data.m`, `Data.r`, `Data.k`, `Data.t`
a, b_1, b_2	Rounded order $a \geq xb$ and rounded weights $\geq a/x, a/x_1$, all multiples of N_1.	`Data.a`, `Data.b₁`, `Data.b₂`
h_2, w_j	Half-gap r^8, integral weight with increment $2h_2$ per stage.	`Data.h₂`, `Data.w`
w_{nom}	Effective weight a/x at thick places, a/x_1 elsewhere; enters the exact prime sum.	`Data.nominalWeight`
g_0, g_1, g	Two chosen split auxiliary primes and their product, $g = g_0 g_1$.	`Data.g`, `Data.gProd`
N	Extra prime level on column groups; selected to pay for the main archimedean excess.	`Data.N`; 54.3
M_ω	Auxiliary integer $(N_0 d)^{s+6}$ specifying tame-order characters at chosen places.	`Parameters.Mω`
X_j, Y_j	Definite Hermitian spaces of rank $k + j(t+1)$, row spaces of rank $r + jt$.	14

Symbol or term	Meaning	Source or chapter
C, C^{ram}, W	Column form space, image of the ramifying operator, and good row form space; $c = \dim C$, $n = \dim W$.	`Column`, `ramifiedColumns`, `rows`
$\mathcal{G}$	Product of the finite GL_r at chosen places above g_0, g_1; indexes projected translates.	`Data.G`; 20
e_0	Central orthogonal projector onto the specified good finite types.	`Data.e₀`
P_s	Projected Poincaré maps, with the common normalization adopted in Theorem A.	`Data.projectedPoincare`
θ, Θ_θ	Weight-zero tame m-fold twist; composite row theta lift with the prescribed powers of the twist.	14
ι	Field isomorphism to a p-adic algebraically closed coefficient field; never assumed continuous from $\mathbb{C}$.	`PadicIso`
$\mathrm{IndexLe}(y, d_{\mathrm{ind}})$	Logarithmic cost of integral combination matrices for an independent or dependent row family at all finite-place embeddings and all prime-to-p twists.	`Data.IndexLe`; 5.10
V	Log upper bound on the squared covolume of a lifted family, uniform in its coefficient conjugates.	31
$\lambda_{d,M}(\nu)$	$\mathbf{1}_{\nu \equiv 1 \bmod M}\Lambda(\nu)(1 - \chi_K(\nu))$.	`lambdaClass`; 2
$R_M(z)$	Smoothed normalized prime remainder, after subtracting $\log z/\varphi(M)$ and the constant $c_{d,M}$.	`primeError`
$A(y, \sigma)$	Unnormalized finite prime sum with weights $w_{\mathrm{nom}}([-\nu]\sigma)$ and cutoff a.	`Data.primeSum`
$A_p(y, \sigma)$	Local positive-part sum for the live powers of a nonsplit p, with the factor $(1 - \chi_K(p))/2$.	`Data.localAsk`; 13
g_p	Desired p-adic gain over the entire chain, m times the sum of the distinguished long-row gain and k ordinary-direction gains.	`Data.gain`
$\Omega, J(j)$	Period scale $2\pi \exp(\kappa_K)/\operatorname{rd}_E$ and derivative cost $\log j! - j \log(\Omega/g)$.	`Parameters.bigOmega`, `Data.jetSize`
$\mathcal{U}, H$	Net ordinary direction cost and common main term $H = mk\mathcal{U}$.	`Data.dirCost`, `Data.mainHeight`
$\mathcal{R}$	Arithmetic remainder after subtracting the common mean term from the sum at all tops.	`Data.remainder`
B, L, B_0	$fa(\log b)^3$, $fa(\log b)^4$, and $B/(mk)$, respectively.	`Data.budget`; 54
$Z(\theta)$	Squared volume of the row-chain map at a twist, from its Gram determinant in the given metric.	31
Fit family	A choice of twist at each embedding, controlling determinants of tested projections and ratios of Hecke values, for every power through m.	`Data.IsFitFamily`; 40
Local block loss	Cost of imposing a finite/crystalline relation on torsion cocycles in the row and Tate-string blocks at a rational prime.	`loss`; 49
$\star$	Main inequality $m\mathcal{R} \le fa(\log b)^5$.	`star_inequality`; 55

56.1 Reading the main theorem in the Lean source

To compare the monograph with the formal proof, begin with the five exported statements in `Main.lean` and the real statement definitions in `Statements.lean`; these do not depend on the specialized definitions of the auxiliary construction. Then read `New/Main.lean`, where the endgame is expressed in only a few declarations. The upper and spanning inputs are `Data.short_forms` and `Data.enough_forms`, in `New/Endgame/ShortAndEnough.lean`. The lower input is `Data.integral_forms_long`, in `New/Endgame/LongForms.lean`. The final

chosen data come from `exists_data` in `New/Data/DataExist.lean`. The analytic objects and criteria are in `New/Endgame/PrimeError.lean` and `New/Endgame/OneSidedCriterion.lean`. This is the shortest route through the dependency logic. One cannot read the code at that level to verify the arithmetic input theorems themselves, but one can check the precise mathematical interface against the deductions in the final part of the volume.

At a second level, the modules `New/Curve/`, `New/Columns/`, `New/TwistedHecke/` and `New/Seesaw/` implement A; `New/Filling/`, `New/RealPlace/` and `New/DiscreteSeries/` implement B; `New/Rows/`, `New/Exchange/`, `New/Selmer/`, `New/Patching/` and `New/Lift/` implement the principal pieces of C. Definitions and auxiliary lemmas in the `Lib/` tree are shared or specialized at these interfaces. The readable edition of the repository is often easier for understanding motivation or naming a mathematical package: consult its `GUIDE.md`, `INDEX.md` and `CONTENTS.md`. The compact edition is generally easier to use for extracting exact constants. The two should not be conflated when a particular theorem has been reworked mathematically, notably the analytic choice of the second cutoff.

Chapter 57

Third proof-closure audit and remaining dependencies

57.1 Scope of this convergence pass

The seventh edition continues the work of the previous 849-page reconstruction. The purpose is a conventional, independently readable derivation of the specialized assertions needed for the main theorem. This remains a working mathematical reconstruction, not a declaration that every step of that theorem has been completed. The 101 grouped interfaces are retained from the first closure audit; they are not theorem counts, a measure of the percentage of the Lean code verified, or a probability that the full argument is sound.

Proved means a written mathematical derivation of the identified source interface, subject to its explicit hypotheses and identified background theorem inputs. It does not mean that the new argument has been independently refereed or formally verified. **Reference** denotes a suitably identified literature result. **Conditional** means a source-matched reduction with an outstanding specialized hypothesis. **Open** denotes a substantive incomplete interface. In particular an implication proved from its stipulated data is not proof that all those data exist. Earlier chapter ledgers are historical; their status is superseded only to the extent indicated by the new derivations and the present audit.

57.2 Focus of this edition

The principal additions concern the exact weighted inverse-index count and class-energy calculus, construction and quantitative estimates of real and finite Bessel systems, singular Fourier-coefficient induction and normalized complementary exchange, quantitative carrier and rank-two measure comparisons, and the arithmetic application of the Ribet polynomial and patched Hecke operators. The outstanding dependencies of the main theorem are now more sharply localized.

1. **Filling and the actual good datum.** Chapters 28, 29 and 30 establish the actual steep cut and its weighted majorant, the standard and finite level formulas, Poincaré construction, the inverse-weight, Fourier and lower-floor estimates, the index enumeration, and the thin- and thick-class energy estimates. The separately stipulated geometric good-cut/arithmetic generic support and marked-prime transfer in the seesaw must still be realized for the main spanning theorem. It is not permissible to infer existence of the full good datum merely from a local estimate or a universal bound for an admissible filling.

2. **Periods and the starting-row problem.** Chapters 37 and 38 reconstruct the singular-coefficient induction, the primitive matched mixed-model period, extension to standard-column data, and the equal-rank normalization and exchanged scalar. The distinct joint real-convolution hypotheses, starting normalized periods and quantitative row transport/reachability, geometric carrier class-sum identification and actual test pairings, and the complete tower vanishing and bounds for the subtracted form on the tested noncompact quotient remain. Chapter 39 gives several quantitative carrier estimates and a precise rank-two integral measure comparison; the existence and normalization of the remaining geometric carrier and determinant-period data must be checked as such.

3. **Arithmetic realization and fit.** Chapters 47 and 48 give the torsion-to-crystalline defect, conjugate-inverse trace ties, exact arithmetic Ribet top coefficient, the labelled base level, integral cutoff, local character/string operators, coefficient extraction, and denominator implication for the given cut. This makes the existence of the input rows (the actual good theta-Hecke transfer and uniform short-Selmer bounds) and the rank-two normalized value family and the simultaneous fit the relevant remaining arithmetic tasks. The implication is not an existence theorem for these data.

4. **Local numerical estimate.** The preceding edition's proofs of the two actual compact local cutoff cases and their global weight windows are retained. They do not constitute a proof with the optimal abstract constant for all possible parameter choices, nor impose the needed crystalline and admissibility hypotheses on arbitrary representations.

Accordingly the analytic and numerical endgame remains conditional on the three essential arithmetic estimates. This book is not yet an independently unconditional proof of Dirichlet GRH.

57.3 Interface ledger

The previous edition recorded 57 proved, 4 referenced, 37 conditional and 3 open groups. In this edition the same 101 groups comprise 64 marked Proved, 4 Reference, 32 Conditional, and 1 Open. The separate ledger retains the source anchors, the location in the frozen 550-page predecessor and the prior status. The chapter-level ledgers provide more detailed mathematical qualifications.

ID	Interface	Status	Current reason and remaining obligation
R01	Universal colored integral Schur basis	Proved	New theorem clR:schurbasis uses generic triangular identities, unit leading monomials, all-characteristic Weyl dimension and integral Smith normal form. Historical input label obsolete.
R02	Rectangular rank-change identity and integral coordinate extraction	Proved	Corollary clR:rankchange gives Cauchy-Binet and determinant factor plus integral coordinates over every ring; source's unspecified choice harmless by integral basis transition.
R03	Quadratic idelic norm subgroup and Hasse norm theorem	Reference	Neukirch VI section 5 theorem (5.5), VI section 4 corollary (4.5); exact hypotheses and norm signs checked in clR:normref.
R04	Normalized splitting character	Proved	fullD:cyclo-unit and fullD:split-ex give actual character extension using R03; finite unramified signs and conjugation condition included.

ID	Interface	Status	Current reason and remaining obligation
R05	Odd class number of quadratic field	Proved	Elementary invariant-ideal argument proves bijective squaring when prime discriminant. Earlier genus-theory reference not needed.
R06	CM potential good reduction and local isogeny degree	Proved	Verified Silverman Advanced Topics II Thm 6.1/Prop 4.4 and Arithmetic VII Prop 5.5; marked good cubic and local-length proof in mcm chapter.
R07	Area-normalized Kronecker limit and algebraic-unit period	Proved	clR:kronecker supplies Poisson proof including full-period discriminant constant (DLMF 23.19.4); mcm:CS supplies unit adjustment.
R08	CM local period normalization	Proved	Period scalings derived in mcm chapter (mcm:s0 et seq.) from norm subgroup/splitting character and unit-adjusted period; not an assertion of arithmetic Weil vector realization.
R09	Cyclotomic field completions and inverse Frobenius	Proved	fullD:frob computes embeddings, decomposition, and selected side; earlier generic splitting label obsolete.
R10	Freeness of chosen principal levels	Proved	The definite free-level trace/norm argument at lines 1696–1707 and maf:clear-stab establishes the needed rank/level condition, not arbitrary adelic neatness.
R11	Ordinary Bloch–Kato finite local condition rank	Reference	Bloch–Kato Prop 3.8, Cor 3.8.4 and Brinon–Conrad Prop 6.3.2 checked; clR:BK gives exact classical/lattice/form sign bridge.
R12	Exact-dual orthogonality and finite cocycle counts	Reference	Bloch–Kato 3.8 and Milne I Cor 2.3/Thm 2.8 checked. Imperfect pairings not implied.
R13	Finite vacuum action and Gauss-sum phase	Proved	Normalized finite reproducing-kernel/lattice identification and inert sign, excluding real/nonpolar theta comparison.
A01	Real/nonpolar Weil model, canonical theta functional	Proved	Chapters clV and clG construct the oscillator model, normalized invariant theta functional, parabolic action, and rational phase; qualitative theta convergence is derived.
A02	Finite image and arithmetic theta-vector realization	Proved	clA:entire gives finite-image multiplier, injective entire realization and exact Taylor normalization. Its hypotheses are the positive CM model and rational vectors.
A03	Soft normalized CM reciprocity	Proved	fullA:soft-CM establishes transport and finite-adelic cluster with odd-class root argument, no direct finite-model identification.
A04	Integral vacuum generation and conjugate valuation bound	Proved	clA:semilinear and clA:archvac give conjugate generators, compatible finite actions and uniform vacuum bound; use the marked CM comparison and integral lattice already developed in the book.
A05	Finite translated vacuum stability and theta gauge	Proved	clA:ledger supplies finite translations, integral vacuum stability, raising and exact gauges under the prescribed frame hypotheses.
A06	Seesaw theorem	Conditional	ntA supplies the actual marked Bessel arithmetic and level compatibility; separate finite theta projector realization and exact normalized seesaw-period transfer remain.

ID	Interface	Status	Current reason and remaining obligation
A07	Arithmetic finite cuts, evaluation and good datum	Conditional	ntA:actual-steep, ntA:admissible and ntA:fits give construction, quantitative filling and fitted coordinates from the stipulated datum; separate genuine generic support, projector realization and coherent packet comparison remain.
A08	Real branching restriction norm and finite diagonal gauge	Proved	clA:integral-branch gives integral branching map, norm, multiplicity one and kept-height bound; clA:jetbound proves conjugate norms for admissible gauges.
A09	Short columns span	Conditional	The actual local cut, class energy, bands and Poincare maps now have proofs. The global-to-local coherent packet and actual marked generic support must still give the source short-column spanning conclusion.
A10	Marked-prime Hecke compatibility and raised gauges	Conditional	Finite Weil/gauge stability and exact local representations established in clA and ctG. Actual marked-prime geometric transfer and generic packet remain.
A11	Paired lift law normalization and power denominator	Conditional	clA even-stage normalized exact law and ctR analytic identification are available. The actual transfer to the marked noncompact Bessel datum remains.
A12	Uniform conjugate archimedean gain and class-count normalization	Proved	clA:jetbound, clA:classes, clA:chainbound prove the required norm and counts for admissible data and the stated chain cost.
A13	Concrete even-stage coefficient law and final branching factor	Conditional	Even-stage and final branching supplied by clA; the noncompact coefficient supplied by ctR. Initial arithmetic good datum remains.
B01	Stable elliptic orbital character comparison	Proved	ctR constructs the real Flensted–Jensen lowest-type coefficient and identifies parameter and trace resolution, supplying the missing premise of clT.
B02	Rigid finite-label spectral comparison	Conditional	ctF supplies Green labels and corrections, ctG gives attached arithmetic representations and label rigidity. Actual injective auxiliary lifts with exact local intertwining remain.
B03	Green character degree and bad-share bound	Proved	ctF:green–ctF:primitive derives the source Green class functions, degree, bad-share bound, primitive scaling, cubic avoidance and reflection annihilator from precise Green Theorem 1 and finite algebra.
B04	Global filling datum existence and actual generic support	Conditional	ntA constructs the actual levels, cut/majorant and numerical admissibility from specified arithmetic class formulas; ntF and ntB give bands and weighted energy. Actual geometric generic support, reduction and coherent global packet compatibility remain.
B05	Local systems and exact cut/full Schur bounds	Conditional	ntA:cut-majorant, ntA:schur-degree and ntA:stdlevel give the actual local systems and cutoff; ntF proves arbitrary-head and band calculus from these quantities. Check the exact RealPlaceSizes Haar-ratio/Schur-energy normalization and upstream reduction conditions before the complete source package is claimed.

ID	Interface	Status	Current reason and remaining obligation
B06	Real open-cell uniform estimate	Proved	ctR derives the real Gram section, Haar and Sobolev bounds, randomized uniform open-cell estimate and simultaneous support for the stipulated local inputs.
B07	Unipotent covolume actual normalization	Proved	fullB:covolume and arithmetic-cutoffs derive exact marked unipotent row count and quotient integration.
B08	Convergence and Poincaré reproduction	Proved	ntA:peak, ntA:cut-majorant and ntA:fits supply the actual reproducing kernel, quantitative majorant, absolute convergence, Poincare reproduction and exact unfolded Petersson pairing. This does not require the band-limit assertion.
B09	Uniform sublattice discriminant zeta sum	Proved	fullB:height-sum supplies explicit sublattice-height bound and denominator weight based on short generators.
B10	Joint pair and saturation bounds	Proved	ctB:upper-link, ctB:complement-gain and ctB:pair-theorem prove the simultaneous pair-discriminant and saturation inequalities with actual index factors and constants.
B11	Reduction to coherent top-cell pairs	Conditional	ntB:weighted-count, ntB:pair-sum, ntB:class-ampl and ntB:full-crude discharge the weighted/counting/class and crude-energy implications for stipulated filling. Actual coherent packet/zero-form and finite topKernel comparison remain.
B12	Bessel-coset error and central projection covariance	Conditional	ntB:full-crude supplies exact thin/thick energy and weighted marked class sums. ntF supplies the local Fourier system. Actual coherent global moment/packet comparison and its application to the genuine filling still need verification.
B13	Finite sieve system and mass-normalized floor comparison	Conditional	ntF supplies all finite arbitrary-head direct/inverse weighted norms, shells, frequency and finite sieve fields; ntA supplies standard level, conditional on source reduction and filling compatibility. The full actual construction is not inferred from the estimates alone.
B14	Global numeric sieve and Poisson induction	Conditional	ntF:real-band and ntF:lower-step prove the band and floor/Poisson ingredients; ntB and ntA establish exact energy and mass factors. Remaining normalized Haar/actual packet and full global sieve application.
B15	Coherent packets and global-to-local quadric limit	Conditional	ntA:cut-majorant and ntA:fits prove actual weighted majorant, convergence and Poincare reproduction; ntB establishes global amplitude/energy bounds. Exact coherent packet comparison and global-to-local quadric limit still require reconstruction.
B16	Sharp energy and actual span	Conditional	ntB:full-crude proves crude energy with exact constants and ntA supplies simultaneous mass/degree admissibility. The global-to-local packet identification and actual sharp span application remain.

ID	Interface	Status	Current reason and remaining obligation
B17	Lowest discrete-series coefficient estimates	Proved	ctR constructs the explicit normalized coefficient and derives the exact decay, determinant, cone and half bounds under the source parameters.
B18	Infinitesimal differentiated Schur bound	Proved	ctR establishes highest type, infinitesimal Casimir filtration norm and differentiated Bessel Schur estimate with the specified constants.
B19	Real floor Fourier bounds	Conditional	ntF:parsevalineq, ntF:real-band and ntF:lower-step prove real inverse-weight, unequal-head Fourier and floor descent and Schur-energy implication. Exact RealPlaceSizes Haar-ratio/Schur normalization and actual source hypotheses still require final matching.
B20	Gram-section open-cell integration	Proved	ctR supplies the exact Gram-section block unitary coordinates, Haar Jacobian, uniform tube Sobolev estimates, integration and masses.
C01	Integral frame pairing gauge	Proved	ctP:pair gives rational integral Gram pairing, weak approximation and exact normalized finite stabilizer/class pairing gauge.
C02	Rank-one nonzero real ratio	Proved	ctP:ratio-real gives exact nonzero real ratio using the stipulated L-value weight/infinity types, Tate and logarithmic Euler positivity argument at the edge.
C03	Fit family existence	Conditional	This is downstream of S and L, not independent input beyond their obligations.
C04	Complementary-rank exchange	Proved	ntW:period, ntW:topreal, ntW:circle and ntW:descent give the primitive matched period, exact multiplier and rank move; ntP:freitag-thm and ntP:promotion give the full stable standard-column reduction, and ntW:equal proves the equal-rank nonvanishing used in exchange.
C05	Equal-rank top-period normalization	Proved	ntW:equal and ntW:double provide the exact double period exchange, nonzero constant and character-independent top ratio, with ntP:promotion and ctP:const. Distinct joint convolution edge hypotheses remain C06.
C06	Uniform analytic kernel bound and square law	Conditional	Complementary exchange and equal-rank normalization now supplied (ntW). The actual simultaneous real-convolution edges and the uniform integrable kernel bound still must be verified.
C07	Finite square move	Conditional	The previous square move, integrability and new full complementary exchange are available. The actual joint edge family still remains (C06/C08).
C08	Starting periods and row-reachability	Open	ntW gives top complementary exchange; ntC:rowtail gives uniform deep-level restriction tail under stated shifted coefficient bound. The actual starting normalized periods, two-factor period/thin-part comparison and quantitative reachability still need reconstruction.
C09	Route recursion	Conditional	fullC:route gives combinatorial induction assuming actual normalized square law and descent.
C10	Terminal periods and bounds	Proved	ctP:terminal-bounds proves complementary-minor dual pairing, compact level normalization and tidy terminal expansion with integer and conjugate bounds; noncompact transport remains C08/C09.

ID	Interface	Status	Current reason and remaining obligation
C11	Crude rows from exchange	Conditional	ntW supplies exchange and ntC:rowtail gives a quantitative row tail. Transport of the genuine starting normalized periods and the two-factor restriction estimate remains.
C12	Good row projection and integral monomial coefficient bounds	Proved	ctP:good-bd and ctP:monomial prove exact pointwise integral gauge and class norm and integer monomial coordinate vector, with specified hook/factorial polynomial.
C13	Rallis tri-lift pairing and transfer	Conditional	ctW:rallis and ctW:euler prove sharp general Rallis unfolding and local factors at stated edge; actual higher-rank edge/type and tri-lift comparison still required.
C14	Comparison forms, glued line scale, separation	Conditional	ntC:nearmax, ntC:circle, ntC:isolation, ntC:reproduction and ntC:carrier-approx supply analytic isolation, short-Hom/reproduction and approximation. Actual geometric carrier class-sum/coset matching and good-row theta types remain.
C15	Normalized star projection at all embeddings	Proved	ctP:star-thm gives exact local Hecke adjoint and conjugate projection for tested vectors, proving the stated source interface. Actual tested-vector realization remains separate.
C16	Rank-one Rallis L-value identity and integral measures	Proved	ctW:rallis, ctW:euler, ctW:map-trivialLift and ctW:integral-measure give rank-at-least-four rank-one normalized norm identity, semilinear comparison and integral measure with exact factors. Rank-two family separately S05.
C17	Rank-one conjugate norms and separation	Conditional	ntC:height gives exact height consequence and quantitative approximation/isolation; actual carrier identification, class-sum and test data remain.
C18	Depth-indicator attainability and test pairing	Conditional	ntC supplies carrier analytic estimates and normalized approximation but not the actual weighted jet/indicator attainability or tested pairing.
C19	Class-factor averaging on rank-one line	Conditional	ntC strengthens the carrier analysis and rank-two denominator but not the actual rank-one theta integral representative at all required levels/carriers.
C20	Graph zero terms in doubling	Proved	ctW:nongraph gives exact clean nongraph support vanishing; combine normalized graph term clW:graph.
C21	Siegel–Weil formula in exact range/model	Reference	clW:SW matches Weil 1965 Thm 5 and Yamana 2013 Thm 2 in the ordinary convergent range, with adelic probability Haar and passage from Schwartz to integrable vacuum. Other hypotheses of applications remain separate.
C22	Rallis pairing integrability and Euler expansion	Proved	ctW:rallis, ctW:dual-lemma and ctW:euler prove Rallis integrability, unfolding, dual datum, sharp coset partial limit and Euler factor/injectivity with source real-edge hypothesis. Actual edges separately C06/C13.

ID	Interface	Status	Current reason and remaining obligation
C23	Normalized line constant term	Proved	ntW:name and ntW:lineframe give exact adapted Hom/frame/vacuum normalization; together with ctW:slice, ctW:lines and ctW:mellin this closes normalized line constant term with both absolute endpoints.
C24	Real cone and matched theta difference	Conditional	ntP:freitag-thm and ntP:promotion prove full abstract singular-to-standard induction; ntW proves the primitive matched coefficient and determinant vector descent. The separate exact real-cone integral determinant surjectivity/synthesis premises must be tracked in actual auxiliary geometry.
C25	Hecke polynomial transfer and tower vanishing	Conditional	ntW:descent proves normalized standard-column tower descent and zero/nonzero orbit argument. Full rational parabolic vanishing of the actual subtracted lift, bounds and tested-period conversion remain.
C26	Integral gluing and Weyl gauge columns	Proved	fullC:divrow/indicator/cartanbound give divided lattice arithmetic, not actual global coefficient realization.
S01	Rows under all twists	Conditional	ntS and ntD now establish arithmetic row/column congruence and denominator implication from stipulated row-string data. Actual good-vector normalized theta-Hecke transfer and row existence under all twists remain.
S02	Balanced lattice and local residual strings	Proved	ctG:balanced-budget constructs balanced lattice from exact IsRow, proves matrix span, residual scalar and 10fn c_l bound (7fn local), with type supplied by row.
S03	Mirror criterion and actual split-prime relations	Proved	ctM:dual, ctM:split and ctM:mirror prove exact mirror theorem via actual finite dual Selmer group, cocycle restriction injectivity, normalized split-place cup reciprocity and isotropy polarization. No intermediary projective count needed.
S04	Uniform bounded pencil	Proved	ctM:mirror removes premise of fullS:pencil; bounded pencil induction and helper construction proved, assuming the specified balanced rows.
S05	Integral value measure and family	Conditional	ntC:ranktwo proves uniform rank-two jet denominator and integral measure interpolation; ntW proves general equal-rank exchange. The precise rank-two determinant-period comparison/endpoint identity for the actual family remains to be matched.
S06	Simultaneous anticyclotomic twist	Conditional	Mirror, balanced row, odd reciprocity, arithmetic Ribet and base-level operators now proved with inputs. Actual rows and rank-two normalized value family remain for the simultaneous twist.
S07	Triangular places and reciprocity law	Proved	clS:law derives the exact higher-depth odd reciprocity law, including the primitive coordinate, factor two and prescribed triangle, by precise finite Poitou–Tate duality.
S08	Odd-class class-length bound	Proved	fullS:odd-budget, component-detect and small-detect give exact explicit integer; stale formal-source description in numerical assembly.

ID	Interface	Status	Current reason and remaining obligation
S09	First-order Ribet block bound	Proved	ctS:firstorder proves the exact integral Ribet polynomial and valuation using triangular Fitting, normalized rows, square-free quotient and two adjugates. ctS:tatecost proves actual Tate-string triangle-cost implication.
S10	Patched Hecke string and tie existence	Proved	ntD:base-level proves actual finite labelled level and uniform integral projector, and ntD realizes the trace functions with the exact local type of ctG. Together with clS ties, chain and extraction this gives the stipulated arithmetic trace-to-operator theorem.
S11	Arithmetic patched operator integrality	Proved	ntD realizes the actual integral Hecke ties, fixed cutoff and detector depths; clS:extract implements the coefficient extraction with the callback ntS:ribet-columns. The existence of the input good cut and rows is separately S01/S06.
S12	Arithmetic Ribet element for actual columns	Proved	ntS:ribet-columns verifies actual column labels, finite lattice, common trace ideal, conjugate-inverse ties, the torsion-to-crystalline defect and exact valuation budget with the three group-order factors. Its row and Selmer hypotheses remain in the stated interface.
S13	Weak spherical Galois attachment all ranks	Proved	ctG:spherical gives alternative all-rank attachment from definite Labesse base change, Moeglin–Waldspurger and BLGGT with exact parity twist and Hecke operator/weight normalization for actual data. Original more general seed theorem not asserted.
S14	Local Galois labels and inertia eigenvalues	Proved	ctG:labelcompat and ctG:power derive prescribed Green squarefree/determinant-block local types, zero monodromy, exact tame root killing polynomials and simultaneous Frobenius/inertia tests with a common functional, using precise cited comparisons.
S15	Integrality in group coefficient algebra	Proved	Universal primitive cyclic-word determinant identity, integral via universal polynomial entries; induction on degree closes span and flattened block cases in clR:amitsurspan and clR:amitsurblock.
S16	Integral denominator and fit theorem	Conditional	ntD:denominator proves the denominator implication for the stipulated lift cut, and ntS supplies its Ribet callback. Actual good rows, short-Selmer hypotheses for the realized cut and normalized rank-two value/period comparison remain for the simultaneous fit.
L01	Actual crystalline Kummer chart	Proved	clL:charts and clL:tame construct the Kummer chart and scalar congruences with subgroup and cyclotomic compatibility.
L02	Frame theorem and quantitative radius	Proved	clL:radiusfactor and clL:frame prove first radius inclusion and frame theorem together with earlier determinant reduction; assumptions are actual crystalline invariant basis and given jumps.
L03	Period matrix Frobenius/field comparison and retraction	Proved	clL:amax–clL:radius and clL:fixed prove integral period rings, field and scalar congruences, retraction and Frobenius comparison.

ID	Interface	Status	Current reason and remaining obligation
L04	Rank-free moving/depth operators	Proved	clL:functorial and clL:nomultiple supply exact chart/depth comparison; combine preceding scalar step and rank-free operator proof.
L05	Cokernel killer recurrences and derivation	Proved	clL:derivation and clL:tame prove intrinsic multiplier-ideal derivation, lowering and exact tame polynomial recurrence.
L06	Small-part Witt Artin–Schreier property	Proved	clL structural period lattice and actual field comparison discharge the small-module inputs of the earlier Witt AS proof.
L07	Safe pairing and continuous cup coboundary	Proved	clL:functorial, clL:nomultiple give exponent comparison and no-multiple alternatives; preceding cup cochain calculation applies.
L08	Sparse-thick and two-size cutoff budget	Proved	clL:compactcuts proves both actual compact sparse-thick and two-size corollaries with exact constants. The standalone helper at optimal 51 kappa for all abstract parameter choices remains open but these uses bypass it.
L09	Long-gap Taylor lifting	Proved	clL:radiusfactor supplies the first radius inclusion; earlier long-gap lifting proof now applies to the chart with specified jumps.
L10	Few-low-jump estimate	Proved	clL:fewlow, clL:nomultiple and clL:outer supply initial depth, one-step colength, safety, final bound and both zero-loss alternatives.
L11	Tame chart and special exponent	Proved	clL:tame and clL:outer establish chart, exact polynomial recurrence, type restriction by unramified descent, and final tame exponent.
L12	Crystalline defect and thick spacing	Proved	ctL:crysdefect gives exact three prime-size cases with the floor K and exact coefficients. ctL:spacing proves placeClass ratio, no-three Frobenius arc and sign equality.
L13	One-place row and string block estimate	Proved	ctL:block checks actual jump windows, thick pattern, sizes, special tame case and stated row/string loss per local degree. Actual local type and admissibility are explicitly hypotheses of this theorem, separately S01/S13/S14.

Bibliography

[1] L. Alpöge, with Claude, and the authors of the shortened development, *GRH for Dirichlet L-functions, in Lean 4*, supplied snapshot `grh-lean-main`, November 2026. Compact and readable editions (Apache 2.0); the root README links the announcement https://alpo.ge/remark.pdf. Main statements: `src/Statements.lean`. Main mathematical interfaces: `src/New/Main.lean`, `src/New/Endgame/ShortAndEnough.lean`, `src/New/Endgame/LongForms.lean` in the compact edition.

[2] H. L. Montgomery and R. C. Vaughan, *Multiplicative Number Theory I: Classical Theory*, Cambridge Studies in Advanced Mathematics 97, Cambridge University Press, 2007. In particular Chs. 9–11 and Cor. 11.19, and Ch. 15 for Landau arguments. https://doi.org/10.1017/CBO9780511618314.

[3] M. Jutila, On mean values of Dirichlet polynomials with real characters, *Acta Arith.* **27** (1975), 191–198. https://doi.org/10.4064/aa-27-1-191-198.

[4] F. A. Behrend, On sets of integers which contain no three terms in arithmetical progression, *Proc. Natl. Acad. Sci. USA* **32** (1946), 331–332. https://doi.org/10.1073/pnas.32.12.331.

[5] L. C. Washington, *Introduction to Cyclotomic Fields*, 2nd ed., Graduate Texts in Mathematics 83, Springer, 1997. https://doi.org/10.1007/978-1-4612-1934-7.

[6] B. H. Gross, Algebraic modular forms, *Israel J. Math.* **113** (1999), 61–93. https://doi.org/10.1007/BF02780173.

[7] M. Greenberg and J. Voight, Lattice methods for algebraic modular forms on classical groups, arXiv:1209.2460. https://arxiv.org/abs/1209.2460.

[8] J. W. S. Cassels, *An Introduction to the Geometry of Numbers*, Springer, 1959; reprint of the 1971 edition in Classics in Mathematics, Springer, 1997. https://doi.org/10.1007/978-3-642-62035-5.

[9] C. De Concini, D. Eisenbud and C. Procesi, Young diagrams and determinantal varieties, *Invent. Math.* **56** (1980), 129–165. https://doi.org/10.1007/BF01392548.

[10] K. Akin, D. A. Buchsbaum and J. Weyman, Schur functors and Schur complexes, *Adv. Math.* **44** (1982), 207–278. https://doi.org/10.1016/0001-8708(82)90039-1.

[11] J. Neukirch, *Algebraic Number Theory*, Grundlehren der mathematischen Wissenschaften 322, Springer, 1999. https://doi.org/10.1007/978-3-662-03983-0.

[12] M. Kirschmer, *Definite quadratic and hermitian forms with small class number*, Habilitationsschrift, 2016. In particular Proposition 4.2.7 and Lemma 4.2.9 for the unitary mass formula and its local factors. https://www.math.uni-bielefeld.de/~mkirschm/papers/herm.pdf.

[13] W. Landherr, *Äquivalenz Hermitescher Formen über einem beliebigen algebraischen Zahlkörper*, Abh. Math. Sem. Univ. Hamburg **11** (1936), 245–248.

[14] A. Raghuram, Notes on the arithmetic of Hecke L-functions, 2022, https://arxiv.org/abs/2207.03399. General background on algebraic Hecke characters.

[15] A. Weil, Sur certains groupes d'opérateurs unitaires, *Acta Math.* **111** (1964), 143–211. https://doi.org/10.1007/BF02391012.

[16] S. S. Kudla, Splitting metaplectic covers of dual reductive pairs, *Israel J. Math.* **87** (1994), 361–401.

[17] J. R. Getz and H. Hahn, *An Introduction to Automorphic Representations, with a View Toward Trace Formulae*, 22 April 2022 manuscript, Theorem 2.6.3 (finite volume of the adelic norm-one quotient). https://sites.duke.edu/jgetz/files/2022/04/Graduate_Text.pdf.

[18] N. R. Wallach, *Introductory lectures on automorphic forms*, European School of Group Theory, Luminy, July 2001, Theorem 95 (Harish-Chandra's trace-class theorem). https://mathweb.ucsd.edu/~nwallach/luminy-port2.pdf.